

KRAKÓW



ICNPR 2024

INTERNATIONAL
CONGRESS ON
NNATURAL
PRODUCTS
RESEARCH

Kraków, 2024, July 13-17

ABSTRACTS BOOK

SILVER SPONSORS



BRONZE SPONSORS



LOGISTICS PARTNER



PARTNERS



Content

Welcome Letter	23

Committee	24

Chairs of the Scientific Committee	24

Members of the Scientific Committee	24

Chair of the Organizing Committee	26

Members of the Organizing Committee	26

Workshop ECRW	28

The natural diversity of acyltransferases reveals versatility and specificity in the synthesis of gene-encoded lipopeptides	28

Anti-inflammatory biscebranoids from aquaculture-derived soft coral <i>Sarcophyton trocheliophorum</i>	29

Integration of <i>Caenorhabditis elegans</i> bioactivity assays for a sustainable identification of health promoting natural products	30

Novel natural product inhibitors targeting oncogenic MAPK/ERK and PI3K/AKT signaling in melanoma: from large library screening to target identification	31

Towards the engineering of the plant endoplasmic reticulum for sustainable production of specialized metabolites	32

PKC- α activation with new semi-synthetic 7 α -acetoxy-6 β -hydroxyroleanone derivatives for breast cancer therapeutics	33

Development of a high throughput cell-based assay to highlight potential bioactive antiviral natural products against the respiratory syncytial virus (RSV)	34

<i>In vitro</i> model for predicting oral bioavailability of isoflavone C-glycosides, O-glycosides, and their aglycons	35

<i>Dracaena cambodiana</i> biflavonoid CB2 counteracts ferroptosis by directly inhibiting lipid peroxidation and inducing the key ferroptosis suppressor FSP1	36

Dietary molecules from <i>Glycyrrhiza foetida</i> and their modulation on metabolic syndrome pathways	37

How to make "extreme" CPC greener: how to substitute alkanes in biphasic solvent systems using COSMO-RS predicting tools	38

Investigating the anxiolytic activity of selected essential oils in zebrafish larvae and identification of active constituents using biochemometrics	39

Animal Healthcare and Veterinary Phytotherapy: Science and Practice (VET)	41

Plant-derived compounds as a solution to robust health status of chickens and turkeys	41

Medicinal Plant based Treatments for Infectious Disease and Infertility in US Organic Dairy Cattle	42

Effects of Oregano Essential Oil and Purple Garlic Powder in weaning piglets as an alternative to Zinc Oxide	43

Effect of gastro-resistant pellets of thymol on its absorption rate, intestinal wall integrity and accumulation in the rabbit tissues	44

Use of botanical extracts as tranquilizer in fattening animals	45

The food-chain transfer of ecdysteroids and its potential significance in view of human and animal health	46

Behavioral and pharmacological investigation of putative self-medicative natural materials used by wild chimpanzees and mountain gorillas	47

Variation in chemical profile and antibacterial activity of different parts of <i>Balanites aegyptiaca</i> (L.) Delile, (Zygophyllaceae) among two contrasted zones in Benin	48

Opening Lecture	50

Carbohydrate-based vaccines, therapeutics and diagnostics and medicines produced in flow	50

Plenary Lectures	51

Innovative analytical methods for evaluation of efficacy, safety and quality of raw materials and final products in the phyto-area	51

Foundational principles can inspire bioactive natural products research	52

Unusual enzyme reactions in natural product biosynthesis	53

AFERP Association francophone pour l'enseignement et la recherche en pharmacognosie	54

Medicinal plants in the fight against malaria: investigating the <i>Strychnos</i> genus, <i>Artemisia</i> sp. and ellagic acid derivatives.	54

Pharmacognosy faces new challenges: proposals to combine chemometrics and annotations based on mass spectrometry and NMR for more accurate and faster identification of bioactive natural products in complex matrices	55

ASP American Society of Pharmacognosy

57

The evolution of botanical product quality testing

57

Discovery and engineering of natural product molecules for drug development

58

Session I: Marine organisms, fungi, microbes and plants as chemical factories

59

Topical Lecture

59

Efficient discovery of bioactive natural products by targeted metabologenomic approaches

59

Oral contributions

60

Unprecedented bicyclic and tricyclic non-canonical terpenoids via synthetic biology platforms: Isolation and structure elucidation of unique C16 and C17 terpenoids

60

Peptide discovery and development – known and dark matter

61

Co-culture exploration of some selected species of *Fusarium*, *Pestalotiopsis* and *Colletotrichum* in different culture media

62

Evaluation of biodiverse *Sorghum bicolor* and related genotypes for root exudation, sorgoleone production and potential carbon capture

63

‘Two neutrons walk into a bar’. The power of the $^{35}\text{Cl}/^{37}\text{Cl}$ isotope effect in structure elucidation of natural products by NMR and mass spectrometry

64

Charting the chemical content of toxic and lethal French fungi through a multiply annotated molecular networking-based approach

65

Discovery of novel biopesticides from an agricultural disease nursery – a proof of concept study

66

Deep ocean biodiscovery – diving into abyss

67

Specialized metabolites of the herpetile gut fungus, *Basidiobolus*

68

Session II: Natural products in the era of emerging infectious diseases

69

Topical Lecture

69

From microbes to medicine: leveraging nature's arsenal for innovative anti-infectives

69

Oral contributions

70

Observations and insights into the prolific and varied metabolites of the fascinating plant genus *Dalea* (Fabaceae)

70

Target identification of 3-(phenethylamino)demethyl(oxy)apamine (PDOA) as an anti-dormant Mycobacterial substance.

71

Revealing anti-SARS-CoV-2 natural products from traditional remedies using molecular networking

72

Identification and SAR studies of indole-functionalized polyamines as antibiotic enhancers

73

Hope in hop

74

Hops (*Humulus lupulus* L.) compounds against two tropical infectious diseases: *Chikungunya* and *Oropouche* virus (Bunyavirales)

75

The phototherapeutic potential of fungal pigments

76

Villosane A, an allene containing bisabolane sesquiterpene with potent antiplasmodial activity isolated from cultures of the marine derived basidiomycete *Halocyphina villosa*

77

Naphthylisoquinoline alkaloids streamlining mycetoma drug discover by circumventing differential diagnosis

78

Antibacterial marinopyrroles and pseudilins act as protonophores

79

Session III: Biotransformation and biosynthesis of natural products

80

Topical Lecture

80

Flavoenzyme-driven structural diversification of bacterial polyketide and tropone natural products

80

Oral contributions

81

Streamlined chemoenzymatic synthesis of cyclic peptides by non-ribosomal peptide cyclases

81

Enzymatic synthesis of human phase II metabolites of flavonoids and other (poly)phenols

82

Fungi – important biocatalysts in production of natural pigments and fragrances

83

First insights in the metabolic faith of paraconic acids from *Cetraria islandica*

84

Biotransformation of favelines from *Cnidocolus quercifolius* into a new bis-nor-diterpene with anti-melanoma potential

85

Enterolactone and enterodiol produced by fermentation of Valerian root extract with human fecal samples

86

Biosynthesis of bicyclo[3.3.1]nonane derivatives in *Hypericum sampsonii*

87

Unravelling the biosynthesis of phenylphenalenones: insights from metabolites and enzymes in *Musella lasiocarpa*

88

A synthetic biology platform for refactoring and combinatorial biosynthesis of anthracyclines

89

Discovery of new bacterial sesterterpene synthases and their products

90

JSP The Japanese Society of Pharmacognosy

91

Colletofragarone A2 from the fungus *Colletotrichum* sp. promoted the degradation and aggregation of mutant p53

91

Novel thiosugar sulfoniums, salacinol and neokotalanol, with antidiabetic activity obtained from plants of genus *Salacia*

92

3-epi-18β-Glycyrrhetic acid and its 30-O-glucuronide involves the development of licorice-induced pseudoaldosteronism in humans with individual differences

93

Amway-Nutrilit	94
Roses and buds, in the current state of botanical dietary supplements – a global perspective	94
Society Session: GA	95
Session I: Medicinal natural products: from bench to bedside	96
Topical Lecture	96
Will artificial intelligence become a game changer in natural product drug discovery?	96
Oral contributions	97
Exploring lipidomic changes: the impact of St. John's wort extract ze 117 and escitalopram in a rat model of depression induced by chronic stress	97
Proton spin network fingerprints for dereplication, pharmacophore identification and standardization of the botanical supplement <i>Centella asiatica</i>	98
Drug discovery pipeline reveals ellagic acid to reduce forgetting in <i>Caenorhabditis elegans</i> through specific Musashi inhibition	99
Anti-inflammatory and cytoprotective polypharmacology of BNO 2103 reveals targeting of IKK-NF-κB and p38-MK2-RIPK1 axes	100
Verification of the medicinal value of Patriniae Herba for colon cancer using preclinical models	101
<i>Symphytum</i> genus: novel insights into its phytochemistry and bioactivity	102
Insight into the mechanism of anti-migraine activity of butterbur (<i>Petasites hybridus</i> L.)	103
Study of the prebiotic potential of <i>Althaea officinalis</i> L. root extract using the high throughput <i>ex vivo</i> technology SIFR	104
Development of bio-based active materials for textiles and high-traffic objects	105
Macroalgae derived natural product scavengers of reactive aldehyde and carbonyl species and their implications in slowing the progression of prevalent chronic diseases	106
Mitochondrial modulators: the defenders	107
Bioassay-guided fractionation, identification and docking experiments of α-glucosidase inhibitors from <i>Geum kokanicum</i> root	108
Session II: Natural product research and the challenges of climate change – ecological and environmental solutions	109
Topical Lecture	109
Marine chemical ecology and natural products research in our changing ocean	109
Oral contributions	110
Search for anti-phytopathogenic compounds by studying biotic interactions between <i>Fusarium</i> sp. and the endophytic fungi of the palm tree model <i>Astrocoryum sciophilum</i>	110
Using natural, greener, biodegradable and recyclable solvents to enhance lignin extraction from Acacia wood	111
Development of biobased films from 'Annurca' apple cellulose dopped with phenolic compounds with novel characteristics	112
Potential and barriers of triterpenoid saponins as tomorrow's plant based bio-pesticides	113
Purification of derivatives of 2-amino-3H-phenoxazin-3-one using CPC	114
Leveraging phytochemical variation in <i>Piper scintillans</i> Trel. to discover novel flavanone meroterpenes	115
Invasive weeds: a treasure chest of medicinal chemicals from Palmer amaranth (<i>Amaranthus palmeri</i>)	116
Ecotoxicity studies of natural sesquiterpene lactones on <i>C. elegans</i> as soil model organism for potential bioherbicides	117
Session III: Regulation for the use of natural products in food and drugs	118
Topical Lecture	118
Three-pronged strategy of TCM quality control: deep research, holistic standard and precision testing	118
Oral contributions	119
Herbal Health Products in Europe – an overview on challenges and chances	119
Understanding psilocybin mushrooms and aiding in setting standards for a burgeoning industry in Jamaica	120
<i>Lavandula angustifolia</i> Mill. based soft gel capsules: authenticity and quality assessment	121
Session: Animal Health Care (VET)	122
Topical Lecture	122
Nature's Pharmacy" in the context of insects – is there anything to be learned?	122
Oral contributions	123
Gut-health histomorphometry in fattening lambs fed with hazelnut skin and linseed	123
<i>Malassezia pachydermatis</i> sensitivity to plant extracts from the Brazilian Amazon Rain Forest	124
Orangutan herbal remedies: lessons for human health	125

Willow (<i>Salix acmophylla</i>) leaf and twig extracts inhibit sporulation of coccidia (<i>Eimeria spp.</i>) in goats	126
Hot Pepper (<i>Capsicum annuum</i>) extract: not only a flavor enhancer, but also a useful fish feed additive	127

Session IV: A.I.M.S. Cluster: Prospecting aquatic and terrestrial natural biological resources for biologically active compounds 128

Oral contributions	128
Enhancing bio-based compound diversity through advanced biotechnological approaches: The SECRETed project	128
Marine biodiversity as sustainable resource of disease-suppressive microbes and bioprotectants for aquaculture and crop diseases "MARBLES"	129
Sustainable skincare ingredients from plant cells with scientifically proven efficacy	130

New trends in pharmacognosy and cosmeceuticals 131

Pharmacognosy in the digital era: exciting opportunities offered by metabolomics and accessible digitised data	131
Nanophytocosmetics – a new trend in cosmeceuticals	132
Nitrogenous flavonoid derivatives production by reacting with amino acids	133

SIF Italian Society of Phytochemistry and Sciences of Medicinal, Food and Perfume Plants 134

Investigating structural features and bioactivity of natural products by NMR and computational techniques	134
The epidemic of antibiotic-resistant: a call to research and development of new natural antimicrobial molecules	135
Metabolomic patterns in response to abiotic stimuli of <i>Cladonia</i> lichens from dry grasslands	136
Natural products as active substances and structure components of delivery systems: the case of nanovesicles	137

Session I: Cutting edge technologies in natural product drug discovery, formulation and development 138

Topical Lecture	138
Mass spectrometry in natural products from Brazil: from understanding chemical signaling to pre-clinical studies	138
Oral contributions	139
Ion channels: beyond native mass spectrometry	139
High-throughput cell phenotype screening coupled with ion mobility spectrometry of American <i>Aconitum</i>	140
Antibiotic discovery from a bacterial colony: isolation of new peptidic antibiotics using robotics, 3D printed bioassays, and MS-based bioinformatics	141
Bioactive triterpenic acid metabolism investigation through mass spectrometry approaches	142
Applications of ChemGPS-NP in natural product research – exploring the Outer Rim	143
Addressing the challenge of protein target identification of bioactive phytochemicals with a yeast-3-hybrid screening	144
Remediation of plant extracts using liquid-liquid chromatography: removal of pesticides and tetrahydrocannabinol from hemp extracts as a case study	145
nmrXiv enhances the utility of NMR spectroscopic data of natural products	146
Natural product-inspired chemical biology: development of highly sensitive labeling reagents for amino acids and scarce natural products	147
Role of Metabolomics in natural product discovery and chemical ecology	148

Session IIA: Total synthesis and biomimetic synthesis of natural products 149

Topical Lecture	149
<i>Olea europaea</i> : a compelling source of health promoting biomolecules and an inspiration for the semi-synthetic design of bioactive analogues	149
Oral contributions	150
Synthetic homoisoflavonoids to treat wet-stage macular degeneration	150
Minor Amaryllidaceae alkaloids as inspiration for development of highly selective butyrylcholinesterase inhibitors: design, synthesis, and biological evaluation	151
Naturally based prototypes for vector-control: synthesis and modification of carlina oxide scaffold	152
Design and anti-inflammatory evaluation of ellagitannin-derived postbiotic metabolite conjugated with serotonin and dopamine	153

Session IIB: Genetic engineering of food crops and medicinal plants 154

Topical Lecture	154
Unravelling the fitness programs that steer plant metabolism allows creating novel tools for the engineering of plant-based production of bioactive metabolites	154

Oral contributions	155
Towards the engineering of the plant endoplasmic reticulum for sustainable production of specialized metabolites	155
Session IIC: Medicinal natural products: from bench to bedside	156
A new cytotoxic diphenazine from a deep-sea fungus <i>Cystobasidium laryngis</i>	156
Safety and efficacy of topical Arnica tincture for the treatment of uncomplicated human cutaneous leishmaniasis	157
<i>Mesembryanthemum tortuosum</i> improves Parkinson's Disease symptoms in a zebrafish larvae model	158
Tailoring plants as photosynthetic production chassis for small molecules	159
Session III: Natural cosmetics, nutraceuticals, and food supplements	160
Topical Lecture	160
How to harness the potential of eutectic solvents for the formulation and enhancement of natural ingredients	160
Oral contributions	161
A multi-target approach to evaluate the efficacy of a capsiate rich fraction of <i>Capsicum annuum</i> fruit against typeII diabetes and obesity	161
<i>Cimicifuga racemosa</i> extract Ze 450 mediates resilience against oxidative stress, inflammatory responses and UV-induced aging through metabolic rewiring	162
Phenolic-rich <i>Cratogeomys cochinchinense</i> leaves extract: a promising natural cosmeceutical	163
Dual ¹³ C NMR and LC-MS ² analysis for chemical profiling of Polynesian plant extracts	164
Chemical composition and nutritional value of the Sonora gum	165
Cannabinoids and olive oil secoiridoids: a fascinating case of synergy	166
<i>Lignosus rhinoceros</i> extract ameliorates glutamate-induced neuronal toxicity	167
Novel health potential of isomers of 13 ² -hydroxypheophytin a from <i>Arthrospira platensis</i>	168
Green extraction via NaDES of waste and by-products of the Aglianico cv. vinery supply chain	169
Bioassay-guided isolation of potential cancer chemopreventive agents from the fruits of <i>Rhodomyrtus tomentosa</i>	170
PSE Phytochemical Society of Europe	171
Honeypot strategy for parasite weed control: An offer they cannot refuse	171
A (sub)cellular mapping of natural substances	172
Utilizing natural products for weight management and longevity promotion	173
KSP Korean Society of Pharmacognosy	174
Treatment strategies of pulmonary fibrosis using natural products	174
Coding metabolic profile of medicinal herbs using 2D-HPTLC	175
Status and prospect of natural deep eutectic solvents in current natural products research	176
Discovery and biosynthesis of specialized metabolites from human gut microbiome	177
Poster Session 1, Sunday, July 14, 2024	179
Marine organisms, fungi, microbes and plants as chemical factories	179
S1.P1 Anti-cancer effect of <i>Willughbeia edulis</i> Roxb. stem extract in hepatocellular carcinoma cells through triggering apoptosis and G2/M cell cycle arrest	179
S1.P2 Chemical constituents isolated from leaves of seashore plant <i>Aglaia rimosa</i>	180
S1.P3 Fragment-guided genome mining leads to unprecedented structural complexity in natural products	181
S1.P4 OSMAC approach, the tool to unlock the biosynthetic potential of deep-sea actinobacteria	182
S1.P5 Antiprotozoal activity of alkaloid fractions and isolation of four new steroidal alkaloids from <i>Buxus obtusifolia</i>	183
S1.P6 Genome mining-based discovery of new glycosylated terpenoids from <i>Micromonospora</i> sp. isolated from <i>Leontopodium nivale</i> subsp. <i>alpinum</i> rhizosphere	184
S1.P7 Changing paradigms in natural product discovery: A molecule to microbe approach	185
S1.P8 Bioactive scalarane sesterterpenoids and polybrominated diphenyl ethers from the two sponges <i>Hippospongia</i> sp. and <i>Dysidea</i> sp.	186
S1.P9 New triterpenoid saponins from the aerial parts of <i>Glinus oppositifolius</i> and their anti-adipogenic activity.	187
S1.P10 Chlorine-containing polyacetoxymethanes from the octocoral <i>Juncella fragilis</i>	188
S1.P11 Apoptotic agents from endolichenic fungi against chemoresistant cancers	189
S1.P12 <i>Aetokthonos hydrillicola</i> – an epiphytic cyanobacterium with surprising toxic potential	190
S1.P13 Cellulamides, a new family of marine-sourced peptides from a seaweed-associated <i>Cellulosimicrobium</i>	191
S1.P14 A high-performance thin layer chromatography (HPTLC) method to identify lichen compounds and highlight their antibacterial activity	192
S1.P15 Potent antiproliferative dimeric biaryl-cyclohexapeptides from <i>Lentzea flaviverrucosa</i>	193
S1.P16 A validated RP-HPLC-DAD method for analyzing flavonoids in Caatinga Brazilian green propolis from <i>Mimosa tenuiflora</i> (Willd.) Poir. produced by <i>Apis mellifera</i> L.	194
S1.P17 Functional oligosaccharides derived from the degradation of natural <i>Dendrobium</i> polysaccharides by human gut microbes regulate physiological factors associated with type-2 diabetes	195

S1.P18	Discovering original antiparasitic compounds through mass-guided exploration of the alkaloid diversity in the <i>Strychnos</i> genus	196
S1.P19	Characterization and antimicrobial activity of extracts from microalgae <i>Chlorococcum</i> sp. with green organic solvents	197
S1.P20	Exploration of new biological activity of <i>Mikania micrantha</i> , a widespread weed in Asia	198
S1.P21	Preclinical study on the anti-proliferative effects and underlying mechanisms of broussonetone F in colon cancer	199
S1.P22	Isolates of <i>Pavonia multiflora</i> as alternative control of the phytopathogen <i>Fusarium solani piperis</i>	200
S1.P23	Co-culturing <i>Streptomyces albidoflavus</i> , an epiphyte isolated from <i>Lavandula angustifolia</i> Mill., increased the anti- <i>Porphyromonas gingivalis</i> effect in exo-metabolome extracts	201
S1.P24	Tyrosinase and hyaluronidase inhibitors in <i>Eleutherococcus divaricatus</i>	202
S1.P25	Four sesquiterpene-acylated flavonol glycosides from the sprouts of <i>Astragalus membranaceus</i>	203
S1.P26	Effects of Astrafavonoids A and D, flavonol glycosides from the aerial parts of <i>Astragalus membranaceus</i> , on TNF- α -induced human dermal fibroblasts	204
S1.P27	Harnessing the microbial richness of <i>Ips perturbatus</i> for novel antimycotic agents	205
S1.P28	Insights into the mechanism of antibacterial action for prenylated (iso)flavonoids from Fabaceae: a systematic study on inhibition and permeabilization of Methicillin-resistant <i>Staphylococcus aureus</i>	206
S1.P29	<i>Porphyridium cruentum</i> as a chemical factory: valorization of high-value phycobiliproteins and lipids using an innovative solid-liquid-liquid method based entirely on eutectic solvents	207
S1.P30	Formicine ant venoms put to the acid test - novel antimicrobial peptide families revealed by proteotranscriptomics	208
S1.P31	Antibacterial activity of a novel callus culture line from <i>Cistus creticus</i>	209
S1.P32	Secondary metabolites produced by <i>Nigrospora sphaerica</i> associated with the invasive weed <i>Cenchrus ciliaris</i>	210
S1.P33	Anti-inflammatory activity of a methanol extract of <i>Adiantum capillus-veneris</i> L.	211
S1.P34	Fatty acid profile of lipids from microalgae grown on media with hydrolysed agro-food biowaste	212
S1.P35	Cytotoxic cyclopeptides from the fungus <i>Gymnopus fusipes</i> (Omphalotaceae)	213
S1.P36	Cytotoxic hederagenin saponins from the seeds of <i>Oxybasis rubra</i> (L.) S. Fuentes, Uotila & Borsch (Amaranthaceae)	214
S1.P37	Untargeted and targeted LC-MS metabolomic analyses of putative chemical defenses in <i>Taenaris</i> butterflies	215
S1.P38	Multimodal analysis of oak wood metabolites for a comprehensive understanding of the ageing process in French typical spirits barrels	216
S1.P39	Exometabolites from the Mediterranean sponge <i>Agelas oroides</i> : where is oroidin?	217
S1.P40	Anti-platelet aggregation and vasorelaxant activity of rapanone from <i>Ardisia crenata</i> Sims	218
S1.P41	Unlocking bioactivity of endophytic fungi from Mokrzyński Forest in Wrocław, Poland	219
S1.P42	HME-DDS-applied astaxanthin nanocapsule suppresses DNCB-induced atopic dermatitis-like inflammation in BALB/c mice	220
S1.P43	Spermidine alkaloids of the native Australian plant <i>Acacia auriculiformis</i>	221
S1.P44	New bioactive peptaibols from a soil fungus, <i>Trichoderma strigosum</i> (Hypocreaceae)	222
S1.P45	Evaluation of bioisosteric 6,5-fused heterocyclic salvinorin A analogues	223
S1.P46	Two novel saturated alicyclic carboxylic acids in <i>Ginkgo biloba</i> (Ginkgoaceae) leaf extract EGb 761	224
S1.P47	Microalgae: versatile microbial factories for natural product synthesis	225
S1.P48	Chemical and biological characterization of species of the genus <i>Weinmannia</i> (Cunoniaceae) endemic to Reunion Island for valorization purposes	226
S1.P49	Enhancing caffeoylquinic acids production in <i>Limbarda crithmoides</i> (L.) Dumort. through yeast extract elicitation in liquid root cultures	227
S1.P50	Daphnane diterpenoids from the fruits of <i>Daphne pseudomezereum</i>	228
S1.P51	Nephroprotective <i>Ganoderma</i> sp. extracts towards cisplatin <i>in vitro</i> : chemical composition, metabolomics study and potential impact on cancer cells	229
S1.P52	Phytochemical differentiation of wild thyme from thyme	230
S1.P53	Chemical ecology in marine fungi: diversity and dynamics of pyran-2-ones in a mussel- derived <i>Penicillium restrictum</i>	231
S1.P54	Development and optimization of different eco-extraction and analytical methods on <i>Cannabis sativa</i> L.	232
S1.P55	Impact of biomass hydration on the valorization of non-polar metabolites from <i>Spirulina</i> using eutectic solvents	233
S1.P56	Chemical investigation of the beetle <i>Agrilus cyanescens</i> (Buprestidae)	234
S1.P57	New lipopeptides from the marine bacteria <i>Pseudovibrio denitrificans</i>	235
S1.P58	The Arabian Peninsula: A largely untapped resource for biotechnology	236
S1.P59	Valorisation of north atlantic macroalgae: antimicrobial activity of phlorotannins	237
S1.P60	Assessment of the antioxidant potential of leaf extracts from twelve <i>Sambucus nigra</i> varieties and their phytochemical characteristic	238
S1.P61	Unlocking the potential of <i>Geissospermum</i> trees: exploring bioactive alkaloids for neurodegenerative diseases	239
S1.P62	Bioguided-isolation of antimicrobial compounds against phytopathogens and human pathogens from lichen-associated bacteria	240
S1.P63	Cardiometabolic effects on obese mice of the Mediterranean seagrass <i>Cymodocea nodosa</i> hydroalcoholic extract	241
S1.P64	Obtaining natural and sustainable blue dyes from the fungus <i>Terana coerulea</i> (Lam.) for textile application – The TEXTBLUESTYLE project	242
S1.P65	Using low levels of antimicrobials and antiseptics as elicitors for environmental isolates to activate or re-activate antimicrobial producing biosynthetic gene clusters	243
S1.P66	Biotransformation of the secondary metabolites from <i>Picralima nitida</i> and their antibacterial activities	244
S1.P67	Natural products isolation from subterranean fungi of Wind Cave National Park (US)	245
S1.P68	Chemical and biological study of <i>Glutamicibacter</i> sp. NCA-315 isolated from the sinkhole Tza- Itzá in the Yucatan Peninsula	246
S1.P69	Comparative analysis of chemical variations across plant parts, seasons, and species in three <i>Osmanthus</i> plants	247
S1.P70	Siderophores-mediated antagonism of specialized metabolites in fungal-bacterial interactions	248
S1.P71	Absolute configuration of unisolated epimers of pseurotin D from the marine-derived fungus <i>Penicillium janczewskii</i>	249
S1.P72	A solution from the past: exploring the untapped potential of museum specimens to preserve secondary metabolites	250
S1.P73	Discovery of a brominated Ahp-containing cyclic depsipeptide from the marine cyanobacterium <i>Moorena</i> sp. collected in Panama	251
S1.P74	The utilization of LED lamps to enhance metabolite production in plants – fagopyrins from buckwheat	252
S1.P75	From bugs to drugs: genomic characterisation and bioinformatic analysis of <i>Streptomyces</i> strains from the bark beetle microbiome	253
S1.P76	A microscale cultivation platform integrating OSMAC to explore metabolomic changes and antimicrobial activity of a fungal strain collection	254
S1.P77	Enhanced production of hexylaconitic anhydride compounds from endolichenic fungus <i>Xylaria</i> sp. by co-culture with <i>Chaetomium</i> sp.	255
S1.P78	Bioprospection of bioactive compounds from fungi in Cameroon: current status and outlook	256
S1.P79	Natural product biosynthesis regulation in fungi. What can system biology tells us?	257
S1.P80	Exploration of the clusoid clade metabolome through molecular networking: In the search for new pharmacophoric scaffolds of fungal UPR inhibitors	258

S1.P81	Differences in chemical compounds of selected strains of <i>Hericium erinaceus</i>	259
S1.P82	Genome mining paired with stable isotope labeling allows for discovery of new cyanopeptolins in cyanobacterium <i>Nostoc</i> sp. UIC10890	260
S1.P83	Diversity of <i>Origanum onites</i> essential oils in the ecological regions of Eastern Mediterranean	261
S1.P84	Comparative GC/MS analysis of essential oils from two varieties of <i>Vitex negundo</i> L.	262
S1.P85	Molecular interactions in the ocean microbiome: tracking fungal metabolites modifying <i>Prorocentrum lima</i> physiology and toxin production	263
S1.P86	Chemical mediation of algal microbiome across temperate and subtropical regions	264
S1.P87	Marine actinobacterium as a source of natural products against multiple myeloma	265
S1.P88	Parishin derivatives from the aerial parts of <i>Sedirea japonica</i>	266
S1.P89	Nitrobenzoyl sesquiterpenoids from a Hawaiian marine-derived fungus FM451	267
S1.P90	Steroid derivatives from cannabis-associated endophytic fungus <i>Hypoxylon perforatum</i>	268
S1.P91	Sesquiterpene-phthalide hybrids from cannabis-associated endophytic fungus <i>Zopfiella</i> sp.	269
S1.P92	A new cyclic peptide and two new ellagic acid glycoside derivatives from <i>Jatropha podagrica</i> leaves	270
S1.P93	Discovery of new sesqui-lignans from <i>Salix gracilistyla</i> leaves and their neuroprotective effects	271
S1.P94	Lipopeptides from the bioluminescent bacterium <i>Photorhabdus luminescens</i>	272
S1.P95	A new phenolic glycoside from the leaves of <i>Salix chaenomeloides</i> and identification of anti- skin active compounds	273
S1.P96	Transcriptome mining, immunochemistry, and MSMS <i>de novo</i> sequencing – phosphorylated peptides from marine sponge <i>Stryphnus fortis</i>	274
S1.P97	Distribution of cyanochelins, cyanobacterial beta-hydroxy aspartate lipopeptides capable of iron chelation	275
S1.P98	Syringin and phillygenin - natural compounds with a potential role in preventing of foam cells formation in human atherosclerotic plaque	276
S1.P99	Deciphering monoterpene indole alkaloid metabolism in <i>Heliconius</i> butterfly: a MS-based computational chemistry to describe plant-host interactions	277
S1.P100	Aporphine alkaloids from <i>Phaeanthus lucidus</i> Oliv. and their α -glucosidase inhibitory activity	278
S1.P101	Tychonemazole, a linear depsipeptide, from the cyanobacterium <i>Tychonema bourrellyi</i>	279
S1.P102	Isocoumarin derivatives from cannabis-associated endophytic fungus <i>Phialemonium</i> sp. SUNP1021	280
S1.P103	Novel polyketide metabolites, skeletocutins M–Q, from <i>Tyromyces chioneus</i> and their antibacterial activity against <i>Staphylococcus aureus</i>	281
S1.P104	praca wycofana	282
S1.P105	Usnic acid enantiomers in the lichen-forming fungal family Parmeliaceae	282
S1.P106	The potential of Amazonian microorganisms in drug discovery	283
S1.P107	From kitchen to lab: unveiling methyl 5-vinylnicotinate and other compounds formation during cooking with olive oil	284
S1.P108	C15 acetogenins and sesquiterpenes with anti-inflammatory activity from red algae of the genus <i>Laurencia</i> collected in the Red Sea	285
S1.P109	Diterpenes featuring the spatane, seco-spatane and prenylcubebene carbocycles from the invasive brown alga <i>Rugulopteryx okamurae</i>	286
S1.P110	Metabolic and biosynthetic potential of microbes isolated from North American snake venom	287
S1.P111	Functionalized diterpenes featuring the cembrane skeleton from soft corals of the genus <i>Sarcophyton</i> collected in the Red Sea	288
S1.P112	Neuroprotective activity of <i>Swertia iberica</i> (Gentianaceae) on BV-2 microglia cells	289
S1.P113	Tyrosinase inhibitory and cytotoxic activities of two <i>Inula</i> species (Asteraceae)	290
S1.P114	Linking soil chemistry to Parkinson's disease pathogenesis: novel neurodegenerative compounds from Actinomycetes	291
S1.P115	Isolation and bioactivity evaluation of sesquiterpenes from soft corals of the genus <i>Lemnalia</i> collected in the Red Sea	292
S1.P116	Production of natural products by using plant cell cultures as chemical factories	293
S1.P117	Synergistic approaches in natural products discovery: metabolomic, genomic, and bioactivity- guided isolation of new compounds from the cyanobacterium <i>Phormidium</i> sp. LEGE 15488	294
S1.P118	Peptide toxins from ribbon worms (Nemertea)	295
S1.P119	Exploring the pharmaceutical potential of the Antarctic fungus <i>Epicoccum nigrum</i> isolated from <i>Kallymenia antarctica</i> seaweed	296
S1.P120	Antiparasitic activity of the antarctic fungus <i>Aspergillus unguis</i> metabolites against <i>Trypanosoma brucei</i>	297
S1.P121	Unlocking nature's pharmacy : <i>Iris pseudacarus</i> from Ireland as potential source for antimicrobial and anti-inflammatory compounds	298
S1.P122	Antitrichomonal activity of secondary metabolites of <i>Alternaria</i> sp., a fungal endophyte of alpine Edelweiss	299
S1.P123	Isolation and structural elucidation of bioactive cycloartane glycosides from <i>Astragalus glycyphyllos</i>	300
S1.P124	Screening marine bacterial compounds for adhesion blockers of enteropathogenic <i>Escherichia coli</i>	301
S1.P125	NMR-based metabolomics approach and anti-proliferative activity of <i>Olea europaea</i> products	302
S1.P126	Potential nematocidal agents from a bioactive fraction of the fungus <i>Trichoderma parceramosum</i>	303
S1.P127	Antimicrobial secondary metabolites of <i>Trochila</i> sp., a new endophytic fungus of <i>Lilium carnolicum</i>	304
S1.P128	Seasonal variation of dimeric flavonoids in ginkgo (<i>Ginkgo biloba</i> L.) leaves	305
S1.P129	Efficient production of some bioactive depsides by <i>Aronia</i> $\times$ <i>prunifolia</i> (Rosaceae) microshoot agitated <i>in vitro</i> cultures as the effect of precursor feeding strategy	306
S1.P130	Acridone alkaloids and lignans from the leaves of <i>Luvunga scandens</i> (Rutaceae)	307
S1.P131	<i>Schisandra henryi</i> bioreactor grown <i>in vitro</i> culture as a valuable raw material – biotechnological, phytochemical and biological studies	308
S1.P132	The phenolic profile and antioxidant potential of different types of <i>Schisandra henryi</i> <i>in vitro</i> cultures	309
S1.P133	Isolation and characterisation of stigmasterol and β -sitosterol from <i>Starchytarpheta jamaicensis</i> (L) Vahl (Verbennaceae) leaves of Eastern Nigeria	310
S1.P134	UHPLC-HRMS profiling of sulphated constituents in different plant parts of <i>Salvadora persica</i>	311
S1.P135	Koninginins DS A-C, novel polyketides from <i>Trichoderma koningii</i> , a soil fungus collected from medicinal herbal garden	312
S1.P136	Anti-parkinson pyridone alkaloids from an endolichenic fungus, <i>Thielavia</i> sp. CNC14	313
S1.P137	The anticancer activity and DNA synthesis inhibition potential of <i>Teucrium sandrasicum</i> and <i>Teucrium divaricatum</i> subsp. <i>graceum</i>	314
S1.P138	Kronopolitides A-D, a new family of 36-membered polyol macrolides with antiparasitic activity	315
S1.P139	A correlative metabologenomics approach for the elucidation of the biosynthetic pathway of 3-(3-Furyl)alanine-containing endolides produced in <i>Stachylidium bicolor</i>	316
S1.P140	Discovery and structure elucidation of novel antagonistic cyclic lipopeptides for the control of fungal agricultural pathogens	317
S1.P141	Hydrogen sulfide is a key mediator in the antioxidant effects of osthole on mouse corpus cavernosum and liver	318
S1.P142	Isoimperatorin induces relaxation in mice corpus cavernosum via increasing hydrogen sulfide (H ₂ S) synthesis	319
S1.P143	Purple willow (<i>Salix purpurea</i> L.) varieties as a source of valuable herbal raw material with a high content of salicylic glycosides	320
S1.P144	Exploring myxobacteria for bioactive compounds against malaria and neglected tropical diseases	321
S1.P145	Dereplication of hidden saponins in bioactive extracts	322
S1.P146	Understanding the chemistry of the genus <i>Amylostereum</i> in its multilateral symbioses	323

S1.P147	<i>In vitro</i> evaluation of proliferative activity of Turkish endemic <i>Prangos hulussii</i> S.G. Şenol, H. Yıldırım & Ö. Seçmen and its major compounds	324
S1.P148	Spectral data mining of museum-sourced soft corals collected in the 19 th century for natural product discovery	325
S1.P149	Influence of flavonoid 3',8'' dimerization on antioxidant, neuroprotective, anti- hyperpigmentation and anti-diabetic activities: comparative <i>in vitro</i> analysis of acacetin and isoginkgetin	326
S1.P150	Development of quality criteria for <i>Vitex pinnata</i> Leaf as an herbal medicine	327
S1.P151	Monographic characterization of <i>Vitex peduncularis</i> Leaf as an Herbal Medicine	328
S1.P152	Identification of ergot alkaloids and lysergic acid derivatives in Kykeon, a traditional concoction of the Eleusinian Mysteries	329
S1.P153	Molecular networking-based dereplication and structural elucidation of compounds derived from liquid culture supernatants of marine-sourced <i>Streptomyces</i> strains	330
S1.P154	Src tyrosine kinase inhibitory activity of selected flavonoids	331
S1.P155	Green extraction of bioactive volatile terpenes from Croatian <i>Salvia</i> species	332
S1.P156	Anti- <i>Fusarium</i> activities of microemulsions of essential oils of <i>Artemisia annua</i> L. (Asteraceae) and <i>Salvia fruticosa</i> L. (Lamiaceae) to be used as pre-harvest treatment of fruits	333
S1.P157	Antibacterial and cytotoxic activities of bark and branches extracts from <i>Lafoensia replicate</i> Pohl (Lythraceae)	334
S1.P158	Phytochemical composition and biological activities of <i>Pleroma granulosum</i> (Melastomataceae) leaf extract	335
S1.P159	Combining a 'multi-bioassay screening' approach with metabolomics for in depth exploration of four Andean-Patagonian fungi	336
S1.P160	Diketopiperazines from the marine sediment-derived bacterium <i>Streptomyces hoynatensis</i>	337
S1.P161	The phenolic acids production in stationary and agitated shoot cultures of <i>Aronia melanocarpa</i> , <i>A. arbutifolia</i> and <i>A. × prunifolia</i>	338
S1.P162	Meroterpenoids from <i>Centrapalus pauciflorus</i> leaves with antiproliferative activity	339
S1.P163	Volatiles derived from Philippine plant species exhibited antimicrobial activity, antioxidative and cytotoxicity using a newly developed microplate-based assay	340
S1.P164	Metabolomics-driven discovery of new ophiobolin-type sesterterpenes from marine-derived fungus <i>Pleosporales</i> sp. and their antimicrobial activities	341
S1.P165	Molecular networking of actinobacteria sourced from thai environments reveals the presence of metabolites with antimicrobial properties	342
S1.P166	MSMS-based molecular networking and fragmentation patterns reveal new insights into the astins chemodiversity from the fungal endophyte <i>Cyanodermella asteris</i>	343
S1.P167	Phytochemicals and Enzyme inhibition potential of <i>Rydingia</i> and <i>Moluccella</i> Species in Iran	344
S1.P168	Phenanthrenes from the aerial parts of <i>Juncus tenuis</i>	345
S1.P169	Discovery of anti-inflammatory cyanobacterial secondary metabolites targeting the Keap1/Nrf2 pathway	346
S1.P170	A terpenoids-enriched fraction of <i>Inula viscosa</i> : its chemical composition and anti-cancerous effects against human lung cancer cells	347
S1.P171	Molecular identification and metabolite profiling of endophytes isolated from <i>Bambusa vulgaris</i> var. <i>striata</i> shoot	348
S1.P172	<i>Annulohypoxyton stygium</i> , an endophyte from the marine red alga <i>Asparagopsis taxiformis</i> is a source of antiviral binaphthalenes	349
S1.P173	Antibacterial anthraquinone produced by the fungal endophyte <i>Botryosphaeria rhodina</i> , isolated from the mangrove <i>Xylocarpus granatum</i> J. Koenig	350
S1.P174	Elucidating phytochemical profiles and antibacterial potential of <i>Helminthostachys zeylanica</i> (L.) through untargeted LC-HRMS metabolomics analysis	351
S1.P175	Sesquiterpenoids and hexanorcucurbitacin from <i>Aquilaria malaccensis</i> agarwood with anti- inflammatory effects by inhibiting STAT1/AKT/MAPK/NLRP3 pathway	352
S1.P176	Isolation of <i>Salvia apiana</i> Jeps. (Lamiaceae) essential oil—the influence of distillation technique and process parameters on the yield and composition of volatile fraction	353
S1.P177	Biomasses of <i>Phyllanthus</i> spp. (Euphorbiaceae) as a source of sulfonic derivatives of flavonoids	354
S1.P178	Antibiotic potentiating effects of <i>Salix</i> twig extracts against <i>Bacillus cereus</i>	355
S1.P179	HPLC-MS and 2D-TLC analysis of secondary metabolites from the leaves of selected species and hybrids from the genus <i>Populus</i> with the estimation of their antioxidant properties	356
S1.P180	Antibacterial and phytochemical study of Bornean Myristicaceae and Clusiaceae species	357
S1.P181	Screening for inhibition of β-haematin formation in <i>Rhaphiostylis beninensis</i> (Hook. f. ex Planch) Planch ex Benth. (Icacinaeae)	358
S1.P182	New bryostatin derivatives from the North Atlantic Nudibranch <i>Janolus cristatus</i>	359
S1.P183	Study of the metabolic diversity of Antarctic endophytic fungus <i>Aspergillus unguis</i> in relation to different macroalgal hosts and environmental stressors	360
S1.P184	Oxidation studies on selected betacyanins from the fruits of <i>Hylocereus polyrhizus</i> under influence of ABTS cation radicals	361
S1.P185	Identification of novel betaxanthin pigments in jewel-like fruits of <i>Talinum paniculatum</i> (Jact.) Gaertn	362
S1.P187	Evaluation of a standardized <i>Melissa officinalis</i> extract anti-obesity effects in high fat high sugar diet-induced obesity in rat	363
S1.P188	Biotransformation and antimicrobial studies of flavonoids with bromine and chlorine atoms	364
S1.P189	<i>In situ</i> approaches to marine and terrestrial metabolomics	365
S1.P190	Correlation between saponin content and biological activities of <i>Beta vulgaris</i> L.	366

Natural products in the era of emerging infectious diseases

367

S1.P191	A metabolomic approach in assessing the anti-biofilm potentials of <i>F. proliferatum</i> -a fungal endophyte isolated from an indigenous Nigerian medicinal plant	367
S1.P192	Flavonoids content, anti-inflammatory effect and antioxidant activities methanol leaf extract of <i>Funtumia africana</i> (Benth)	368
S1.P193	Demystifying Bangladeshi ethnomedicines with modern approaches: <i>Congea tomentosa</i> as a source of potential anti-microbial and anti-cancer agents	369
S1.P194	In silico testing of flavonoids of the genus <i>Mentha</i> (Lamiaceae) as potential inhibitors of NS3- protease domains of dengue viruses	370
S1.P195	Application of feature based molecular network analysis for the investigation of the Covid-19 remedy Hanshiyi	371
S1.P196	Pathophysiological assessment of trypanocidal activities of ethanolic leaf extract and fractions of <i>Azadiractha indica</i> on infected experimental model	372
S1.P197	Antioxidant activity and Structural-Activity Relationship (SAR) study of stilbenoids isolated from the stem bark of <i>Anisoptera laevis</i> (Dipterocarpaceae)	373
S1.P198	Resveratrol dimer and tetramers from the isolation of <i>Shorea pauciflora</i> King. (Dipterocarpaceae)	374
S1.P199	In-silico and <i>in-vitro</i> evaluation of natural product scaffolds to overcome the scourge of drug resistant tuberculosis by virtual based drug design approach	375
S1.P200	Exploring Malaysian biodiversity in the fight against antibiotic resistance	376
S1.P201	Effect of seasonal and geographical location on the secondary metabolic contents of <i>Artemisia afra</i> and <i>Artemisia annua</i> . Anti-plasmodial properties	377

S1.P202	Ecology, metabolome and occurrence of <i>Pseudallescheria</i> sp. associated with termites	378
S1.P203	<i>Tetracera alnifolia</i> (Wild) Drake popular in Guinean traditional medicines possesses some pharmacological activities	379
S1.P204	Specialized metabolites with antiviral properties from a Vietnamese plant: isolation and synthesis	380
S1.P205	From signaling molecules to secondary metabolites: harnessing c-di-GMP for enhanced natural product biosynthesis in <i>Streptomyces</i>	381
S1.P206	A novel natural product from the leaves of a <i>Solanum spec.</i> (Solanaceae) shows antimicrobial activity against multiple multi-resistant pathogens	382
S1.P207	Prenylated phenolics from Fabaceae as inhibitors of the multidrug efflux pump NorA	383
S1.P208	Improving the activity of pleuromutilin antibiotics	384
S1.P209	Bioactivity-guided investigations on biofilm inhibition with <i>Rubus chamaemorus</i> leaf extracts	385
S1.P210	Insights into the nature of the microalgal toxins from the <i>Chrysochromulina leadbeateri</i> blooms in Northern Norwegian fjords	386
S1.P211	Revisiting F.P. Porcher's Resources of Southern Fields and Forests (pub. 1863): Insights into the Forgotten Medicinal Botany of the Southeastern United States	387
S1.P212	Bioassay-guided interpretation of antimicrobial compounds from <i>Limonia acidissima</i>	388
S1.P213	Targeted isolation of potent antiviral cyclopeptides from tropical plant extracts	389
S1.P214	Anti- <i>Helicobacter pylori</i> compounds isolated from <i>Sambucus williamsii</i>	390
S1.P215	Lignans from autochthonous Argentinians <i>Larrea</i> species: potential use in the control of microorganisms in agriculture	391
S1.P216	Traditional Middle Eastern spice blends (Baharat): Antimicrobial activity, metabolomic profile, and trace element analysis	392
S1.P217	Ethnopharmacological survey of medicinal plants traditionally used in Tanzania for the management of sickle cell disease	393
S1.P218	Bio-elicitation of isoquinoline alkaloids in <i>Chelidonium majus</i> in vivo & in vitro	394
S1.P219	In vitro anti- <i>Helicobacter pylori</i> activity of extracts and essential oil of <i>Thymus pannonicus</i> All. (Lamiaceae)	395
S1.P220	Assessing the structural interaction profile differences between natural products and synthetics, a data-driven approach	396
S1.P221	Gutierrezones-targeted dereplication integrating LC-HRMS/MS and ¹³ C-NMR data: A comparative metabolomic study of <i>Symphonia globulifera</i> latex	397
S1.P222	QSAR-derived opportunities of prenylated phenolic compounds from Fabaceae as antimicrobial phytochemicals	398
S1.P223	Phytochemistry and biological activities of <i>Ostericum palustre</i> (<i>Angelica palustris</i>) (Apiaceae)	399
S1.P224	Discovery of two new bicyclic macrolides from <i>Kitasatospora acidiphila</i>	400
S1.P225	The potential of curry leaf extract in controlling SARS-CoV-2 infectious syndromes	401
S1.P226	Antiviral effect of <i>Hoveniae Semen Seu Fructus</i> against Influenza A viral infection	402
S1.P227	<i>Thuja orientalis</i> Folium contains antiviral effect against Influenza A viral infection via modulating hemagglutinin and neuraminidase	403
S1.P228	Exploring the phytochemistry, antibacterial and anti-inflammatory activities of three Ericaceous species native to Ireland	404
S1.P229	Funicone-like compounds: potential antiviral agents towards canine coronavirus	405
S1.P230	Efficacy of food waste as a potential source of antibacterial agents and antibiotic adjuvants against MRSA and MSSA isolates	406
S1.P231	Coumarins of <i>Angelica archangelica</i> roots inhibit efflux pumps in <i>Escherichia coli</i>	407
S1.P232	Thymoquinone potentiates activity of antibiotics against clinical isolates of <i>Mycobacterium tuberculosis</i>	408
S1.P233	Exploring olivomycin A: a novel lipopeptide antibiotic - insights into its mechanism of action	409
S1.P234	GC-MS analysis of essential sandalwood oils with in vitro activity against <i>Madurella mycetomatis</i>	410
S1.P235	Bio-guided approach as a valuable tool in the phytochemical investigation of three underexplored Myrtaceae plants	411
S1.P236	From poison to potential: Nb-methyl usambarensine reveals antiviral activity against SARS-CoV-2	412
S1.P237	Semi-synthetic ecdysteroids as promising new antichagasic agents	413
S1.P238	Exploring the potential biopesticide application of nemertide alpha-1 on parasitic and disease vector mosquitoes	414
S1.P239	Chemical constituents of <i>Sambucus nigra</i> fruit (European Elderberry)	415
S1.P240	Invasive plants as promising solutions against biofilms	416
S1.P241	Isolation and characterization of antimicrobial compounds from <i>Picralima nitida</i> (Apocynaceae)	417
S1.P242	Identification of a Potent Antibiotic Compound from <i>Geodermatophilus</i> Bacteria	418
S1.P243	Ethnobotany as a tool for targeted discovery of bioactive natural products for emerging infectious diseases	419
S1.P244	Amaryllidaceae alkaloids as promising anti-dengue phytochemicals	420
S1.P245	Multi-target action of volatile compounds from essential oils in the treatment of periodontitis	421
S1.P246	Metabolomic-guided phytochemical investigation and antimicrobial activity of <i>Morus alba</i> L. twigs extract	422
S1.P247	A new approach to study the mechanisms of action of antimicrobial plant-derived molecules in prokaryotic cells	423
S1.P248	Harnessing the constrained macrocyclic peptide frameworks for the development of antimicrobial peptides to treat infections	424
S1.P249	A steroid-rich fraction from parasitic climber <i>Cocculus hirsutus</i> (L.) W.Theob inhibits coronavirus disease and neutrophilic inflammation	425
S1.P250	<i>Sargassum fusiforme</i> and its components inhibits Respiratory Syncytial Virus replication in vitro and in vivo	426
S1.P251	Inhibition of Respiratory Syncytial Virus in vitro and in vivo by <i>Apostichopus japonicus</i> (Selenka) extract	427
S1.P252	<i>Aster tataricus</i> and its components display a broad spectrum of antiviral effects in vitro and in vivo	428
S1.P253	Extracts of <i>Costus speciosus</i> suppresses influenza A H1N1 activity in vitro and in vivo	429
S1.P254	Exploring faba bean pods as a potential nutritional supplement for piglet gut health	430
S1.P255	Anti-inflammatory properties of twelve Norwegian medicinal plants and their protective properties against impairments in the intestinal epithelial barrier	431
S1.P256	Could Bistort rhizome serve as a potential alternative to antibiotics in preventing diarrhea in weaned pigs? In vitro study	432
S1.P257	Detection and insight into structural requirements of SARS-CoV2 antibodies by peptide- oligonucleotides derived from sunflower trypsin inhibitor 1 (SFTI-1) peptide	433
S1.P258	Chemical analyses and in vitro biological properties of <i>Lythrum salicaria</i> L. for potential use in piglet nutrition	434
S1.P259	Therapeutic potential of stilbene/ β -diketone/gingerol allied naturally occurring non- flavonoids in triple-positive breast cancer: tracing the path from extraction to efficacy	435
S1.P260	Drug discovery based on essential oils concurrently active against eumycetoma and actinomycetoma with <i>Pimpinella anisum</i> emerged as a forerunner	436
S1.P261	Synthesis of autoinducer-2 prodrugs and analogues as a new antimicrobial strategy	437
S1.P262	Bioactive compounds, antibacterial and antioxidant activities of methanol extract of <i>Axonopus compressus</i> (Sw.) P. Beauv	438
S1.P263	The anti-inflammatory and pro-resolving role of selected South African medicinal plants used traditionally as immunomodulatory remedies	439
S1.P264	Phytochemical analysis and antiplasmodial activity of <i>Pseudolachnostylis maprouneifolia</i> fruits extracts	440
S1.P265	Reconstruction of an ancient treatment of infections by incense	441
S1.P266	Chelation of metal ions by tannins: from protolytic equilibrium to antibacterial activity	442

S1.P267	Effect of pH on UV/Vis and CD spectrum of cohumulone and colupulone	443
S1.P268	Inhibitory activity of <i>Vitex negundo</i> (Lagundi) leaf extracts against biofilm formation and quorum sensing of <i>Vibrio cholerae</i>	444
S1.P269	Exploring the role of tannin-metal ion complex formation in antimicrobial activity	445
S1.P270	Isolation of flavonoids from the stem bark of <i>Cynometra malaccensis</i>	446
S1.P271	Chemical constituents from the leaves of <i>Dipterocarpus costulatus</i> (Dipterocarpaceae)	447
S1.P272	16-Angeloyloxy tiglane diterpenoid esters from <i>Euphorbia cooperi</i> with anti-HIV activity	448
S1.P273	Anti-inflammatory and anti-oxidative effects of STW 42 and root extract of <i>Althaea officinalis</i> L. on endothelial cells, fibroblasts and macrophages <i>in vitro</i>	449
S1.P274	Ancient remedies for modern challenges: novel anti-infectives from Egyptian mummy embalming	450
S1.P275	Identification of natural compounds targeting the <i>Plasmodium falciparum</i> pyruvate kinases	451
S1.P276	The relationship between bioactivity and taxonomy of <i>Euphorbia</i> species in South Africa	452
S1.P277	Structure activity studies of anti-parasitic norditerpenes from subterranean fungi	453
S1.P278	Bioactive terpenoids from members of the Hericiaceae	454
S1.P279	Isolation, structural characterization and antiprotazoal activity of steroidal alkaloids from <i>Pachysandra terminalis</i>	455
S1.P280	Antibacterial Activity of different extracts of <i>Plicosepalus acacia</i>	456
S1.P281	<i>Epimedium koreanum</i> attenuates Influenza A viral infection via modulation of hemagglutinin and neuraminidase	457
S1.P282	Anti-biofilm and anti-quorum sensing activity of galloylquinic acids from <i>Copaifera lucens</i> leaves against clinical isolates of multidrug-resistant <i>Pseudomonas aeruginosa</i> in open wound infection: <i>in vitro</i> and <i>in vivo</i> study	458
S1.P283	Effects of raw and boiled ginger (<i>Zingiber officinale</i> Roscoe) on hyperlipidemia in high cholesterol diet-induced obese rats – A comparative therapeutic study	459
S1.P284	Triterpenoids from the roots of <i>Anthocleista djalonensis</i> and their antiplasmodial, antimicrobial, and antitrypanosomal properties	460
S1.P285	Enzymatic Synthesis of Tryptophan-Linked DKP Dimers: Expanding Biosynthetic Routes for Antifungal Drug Discovery	461
S1.P286	An investigation of <i>Origanum onites</i> and <i>Thymus fallax</i> in terms of the European Pharmacopoeia criteria	462
S1.P287	Transformation of food by-products into high value added extracts with antimicrobial properties	463
S1.P288	Secondary metabolites as chemotaxonomic markers in <i>Salix myrsinifolia</i> and <i>S. starkeana</i> (Salicaceae) twigs extracts	464
S1.P289	Carvacrol solution and copper nanoparticles as an effective and non-phytotoxic treatments of carrot seeds against the pathogenic bacterium <i>Xanthomonas hortorum</i> pv. <i>carotae</i>	465
S1.P290	Characterization of polyphenolic compounds from an antibacterial twig methanol extract of <i>Salix aurita</i>	466
S1.P291	Antimicrobial and anti-biofilm properties of geranylated flavonoids from <i>Paulownia tomentosa</i> Steud. fruit	467
S1.P292	Navigating Experimental Metabolomics Knowledge Graphs for Targeted Isolation of Anti-Infective Natural Products	468
Biotransformation and biosynthesis of natural products		469
S1.P293	Discovery of novel terpene hydrocarbon scaffolds and their associated tailoring enzymes from <i>Streptomyces</i> bacteria	469
S1.P294	Effects of difference LED light colors on isoflavonoid production in callus cultures of <i>Pueraria candollei</i> Wall. ex Benth. (Fabaceae)	470
S1.P295	Elicitation of licorice callus for improving flavonoids production	471
S1.P296	Chemopreventive properties of <i>Alchemilla vulgaris</i> L. extract biotransformed by the human gut microflora – preliminary study of a human cellular model of colon cancer	472
S1.P297	Chemical characterization, prebiotic and anti-diabetic effects of galactomannans from two traditional Chinese medicines	473
S1.P298	Phytochemical profiling of phenolic metabolites in flower buds of <i>Magnolia kobus</i> DC. and their antioxidant activity	474
S1.P299	A research of the optimization of recombinant protein expression based on plant system	475
S1.P300	Cyanobacterial natural products: bioactive cyclic and linear peptides	476
S1.P301	Investigation of the effect of solid state fermentation on the release of phenolic compounds from by-products of the agro-food industry	477
S1.P302	Biosynthesis of gramine in barley by a cryptic oxidative rearrangement	478
S1.P303	Urolithin production in <i>ex vivo</i> incubations of gut microbiota – survey on different substrates and antibiotics' influence	479
S1.P304	Identification of gene cluster for biosynthesis of antiviral polyketide limocrocin	480
S1.P305	Reconstitution of natural products biosynthetic pathways using cell-free synthetic biology	481
S1.P306	Biotransformation of phlorotannins from brown algae to enhance bioactivity	482
S1.P307	Enhancing polyphenol profiles and production: a novel approach through synergistic non symbiotic fungi and organic fertilization	483
S1.P308	The potential role of aqueous extract from fruits of <i>Chaenomeles japonica</i> and its post-digestive fractions in inflammation-related intercellular communication	484
S1.P309	<i>Phlegmariurus carinatus</i> <i>in vitro</i> culture and study on <i>Lycopodium</i> alkaloid biosynthesis	485
S1.P310	Biosynthesis of α -aminopyrones produced by the marine-derived fungus <i>Aspergillus</i> sp. DLM38	486
S1.P311	Plant tissue culture for production of bioactive metabolites in <i>Agrostemma githago</i> L.	487
S1.P312	Analysis of host and gut microbial biotransformation products of kratom (<i>Mitragyna speciosa</i>) by mass spectrometry	488
S1.P313	Interactions with gut microbiota and bioavailability of natural products contained in lavender flowers	489
S1.P314	Using banana peels in fermentation: upstream bioprocesses	490
S1.P315	Waste to wellness: hypocholesterolemic solutions from mushroom residues, boosting circular economy	491
S1.P316	Bilateral interactions between human gut microbiota and xanthohumol	492
S1.P317	Engineering potential of the homologation pathway of L-phenylalanine and L-tyrosine	493
S1.P318	Unraveling the biosynthetic pathway of aorimycin: insights from gene deletion experiments	494
S1.P319	Synthetic biology platform for production of novel anticancer agents	495
S1.P320	Symbiotic relation between <i>Ammi visnaga</i> and <i>Rhizobium rhizogenes</i> in bioreactor conditions	496
S1.P321	Application of 1,2,4-triazolium in structural modification of natural products and their biological activity studies	497
S1.P322	Biotransformation of bufadienolides from <i>Drimys maritima</i> and <i>in silico</i> biological studies	498
S1.P323	Diverse ketoreductases involved in tailoring of atypical angucyclines	499
S1.P324	CBD biotransformation potential of endophytic fungi from <i>Cannabis sativa</i>	500
S1.P325	Structure and biosynthesis of hyellamide, a glycosylated N-acyltyrosine derivative, from the cyanobacterium <i>Hyella patelloides</i> LEGE 07179	501
S1.P326	Transcriptome analysis of <i>Triphyophyllum peltatum</i> , a liana producing naphthylisoquinoline alkaloids	502
S1.P327	The roots of <i>Scorzonera kotschy</i> Boiss. a new source for unusual dihydroisocoumarin and saponin derivatives	503
S1.P328	Genome-wide identification and characterization of two potential antiviral plant protein families (Major Latex Protein and Glycine-Rich Protein) in <i>Chelidonium majus</i>	504

Medicinal natural products: from bench to bedside

505

S2.P1	Stellettin B inhibits VEGF-induced angiogenesis in human endothelial progenitor cells <i>in vitro</i> and <i>in vivo</i>	505
S2.P2	Antidiarrheal coumarins from <i>Psydrax schimperianus</i> (Rubiaceae)	506
S2.P3	Ethnopharmacological validation of antiproliferative and anti-inflammatory activities of four <i>Curcuma</i> species through chemical characterization and <i>in-vitro</i> studies	507
S2.P4	<i>In vitro</i> and <i>in silico</i> study of protein tyrosine phosphatase 1B (PTP1B) inhibitory compounds from the leaves of <i>Cleistocalyx operculatus</i> (Myrtaceae)	508
S2.P5	Revival of a traditional Austrian wound healing preparation	509
S2.P6	Enhancing product stability by microencapsulation of a nanoemulsion of plant extracts	510
S2.P7	Homoisoflavonoids from the Hyacinthaceae for use against ocular angiogenesis	511
S2.P8	<i>Pistacia lentiscus</i> : a plant with multiple virtues for use in human medicine as palliative in the treatment of inflammatory-related disorders	512
S2.P9	Evaluation of the effective dose of the aqueous root extract of <i>Sphenocentrum jollyanum</i> Pierre: Role as an antioxidant and phosphodiesterase inhibitor for erectile function	513
S2.P10	Antiepileptic effects of some medicinal plant hydrosols	514
S2.P11	Alisol B 23-acetate and Alisol A 24-acetate reverse MDR cancer via ATPase modulation and membrane fluidity enhancement	515
S2.P12	The constituents and anti-inflammatory activity from <i>Murraya paniculata</i> var. <i>omphalocarpa</i> Hayata	516
S2.P13	Exploring Chemical Composition and α -Glucosidase Inhibitory of <i>Nelumbo nucifera</i> Seedpods	517
S2.P14	<i>Fragaria x ananassa</i> Duch aerial by-product activity against <i>Candida albicans</i>	518
S2.P15	Integrated <i>in vitro</i> and <i>in vivo</i> antilithiatic efficacy of a synergistic herbal formulation, URO-5 and exploration of <i>in silico</i> guided mechanism of action	519
S2.P16	Ashwagandha improves cognition and attenuates depressive-like behavior in an Alzheimer's disease mouse model	520
S2.P17	Chemical structure and bioactivities of a sulfated polysaccharide from precious medicinal fungi	521
S2.P18	Unlocking the therapeutic potential: Comparative analysis of tormentic acid and its ester in combating inflammation and oxidative stress	522
S2.P19	Anti-diarrheal activity of <i>Psidium guajava</i> extract and partitions in <i>Drosophila melanogaster</i>	523
S2.P20	The influence of growth conditions on the amount of active substances in <i>Mentha x piperita</i> L.	524
S2.P21	Inhalation - A promising administration route for anti-infectives from <i>Morus alba</i> root bark?	525
S2.P22	Reporting of medical and health research - An assessment of studies on medicinal plant extracts and herbal medicines registered on PROSPERO	526
S2.P23	Anti-inflammatory and antioxidant effects of <i>Marrubium vulgare</i> L. herb extracts in human immune cells and blood plasma	527
S2.P24	Antioxidant, anti-diabetic and anti-inflammatory potential of <i>Sorbus commixta</i> (Rosaceae) leaves in cellular, cell-free and human plasma models	528
S2.P25	<i>In vitro</i> antioxidant capacity screening of <i>Piper auritum</i> extracts	529
S2.P26	Chemical constituents and antioxidant properties of <i>Verbesina montanoifolia</i> leaf extracts	530
S2.P27	Isolation and characterization of α -glucosidase inhibitors from <i>Ziziphora tenuior</i> using high- performance counter-current chromatography	531
S2.P28	Triterpenoids and flavonoids of Corncockle - a ('n almost) forgotten weed and the (future) medicinal herb	532
S2.P29	Therapeutic effects of Menthacarin and its individual components peppermint oil and caraway oil in an experimental colitis model	533
S2.P30	<i>Bryophyllum pinnatum</i> inhibits oxytocin and vasopressin signalling in myometrial cells	534
S2.P31	Didesmethylocaglamide Cytotoxic Activity in High Grade Serous Ovarian Cancer in Preclinical <i>In Vitro</i> and <i>In Vivo</i> Models	535
S2.P32	Isobavachalcone attenuates TNF- α -induced vascular inflammation in human umbilical vein endothelial cells	536
S2.P33	Synergistic therapeutic potential of quercetin, curcumin, and berberine combination in triple negative breast cancer: insights from <i>in vitro</i> models	537
S2.P34	Quercetin aims at the HuR-signaling axis to impede the migration of Triple Negative Breast Cancer (TNBC): Association with metabolic processes	538
S2.P35	Biological activities of the polar and nonpolar extracts of Philippine Bamboo species	539
S2.P36	Terpenoids and polyphenols from <i>Carpesium cernuum</i> (Asteraceae) of European origin	540
S2.P37	Diet-induced differences in gut microbiota determine the responder or non-responder of traditional herbal medicine	541
S2.P38	Antioxidant activity and cell viability assay on L6 myotubes of different parts of <i>Mitragyna speciosa</i>	542
S2.P39	pH-dependent membrane permeabilisation through saponins and how it can enhance endosomal escape	543
S2.P40	Cytotoxic potential of saponins and fruit extracts from two genotypes of <i>Sechium edule</i> (Cucurbitaceae)	544
S2.P41	Studies <i>in vitro</i> and <i>in vivo</i> of anti-tumor activity of extracts derived from <i>Plukenetia volubilis</i> and <i>Moringa oleifera</i> in gastric tumor	545
S2.P42	Differential modulatory effects of <i>Populus balsamifera</i> (Salicaceae) and <i>Pinus Spp.</i> (Pinaceae) hydroethanolic extracts on IL-6 and TNF production by activated human peripheral blood mononuclear cells	546
S2.P43	Optimization of extraction conditions for salidroside isolated from <i>Acer tegmentosum</i>	547
S2.P44	<i>Eleutherococcus senticosus</i> fruits intract - no toxicity in a balb/c mouse model	548
S2.P45	Potential molecular mechanisms of Silexan identified by RNA-Sequencing in a mouse model of chronic social stress-induced excessive aversion sensitivity	549
S2.P46	The <i>Urtica dioica</i> root extract component in WS 1541 diminishes parameters of epithelial mesenchymal transition in human benign hyperplastic prostate epithelial cells	550
S2.P47	Antitumor activity of evodiamine analogs by targeting PKC β I in gemcitabine-resistant pancreatic cancer cells	551
S2.P48	Perceived changes in anxiety symptom burden during treatment with <i>Bryophyllum pinnatum</i> : a prospective, non-randomised, observational study	552
S2.P49	Effects of orally applied sanggenons C and D in BALB/c mice	553
S2.P50	Novel natural product inhibitors targeting oncogenic MAPK/ERK and PI3K/AKT signaling in melanoma: From large library screening to target identification	554
S2.P51	Screening for natural products inhibiting PI3K/AKT pathways in melanoma: investigation of <i>Mammea americana</i> EtOAc extract	555
S2.P52	Exploring the anti-inflammatory potential of an herbal mixture preparation and individual herbal extracts on inflammatory bowel disease	556
S2.P54	Extracts of <i>Sisymbrium officinalis</i> and <i>Plantago lanceolata</i> reduced inflammation induced by lipoteichoic acid from <i>Streptococcus pyogenes</i> in human tonsil epithelial cells	557

S2.P55	Protein kinase activity of chromatographic fractions of <i>Acacia auriculiformis</i> stem bark as potentials for anti-cancer drug discovery	558
S2.P56	Comparative protein kinase inhibitory activity of three <i>Acacia</i> species growing in Nigeria	559
S2.P57	Diversification of kratom alkaloids using biocatalysis for opioid use disorders	560
S2.P58	Study of <i>Carpesium abrotanoides</i> in dextran sulfate sodium-induced IBD and Caco-2 cells	561
S2.P59	<i>In vitro</i> co-culture analysis of Caribbean medicinal plant extracts for bacterial vaginosis	562
S2.P60	Pharmacokinetic dynamics of the hybrid <i>Sechium</i> H387 07 and phytochemical characterization of its fruit	563
S2.P61	Neolignans and diarylnonanoid derivatives from the seed of <i>Myristica fragrans</i> Houtt. and their cytotoxic activities against human ovarian cancer cell lines	564
S2.P62	The ameliorative effects of <i>Moschus</i> and its major compound L-muscone on traumatic brain injury mice model focusing on the mechanisms of action	565
S2.P63	Cycloartane-type triterpenoids from <i>Combretum quadrangulare</i> (Combretaceae) and their potential for the treatment of hypercholesterolemia	566
S2.P64	Bitter-tasting plant extracts stimulate mechanisms of gastric acid secretion of immortalized human parietal cells via TAS2Rs depending on their polyphenol content	567
S2.P65	Anti-inflammatory testing of some isolated sesquiterpene lactones and methoxylated flavonoids from <i>Achillea millefolium</i> L. in an ICAM-1 <i>in vitro</i> assay	568
S2.P66	Evaluation of aromatase enzyme inhibition by extracts of <i>Bergenia crassifolia</i> (Saxifragaceae)	569
S2.P67	Assessment of the anticonvulsant activity of <i>Myrica gale</i> L., a traditional Irish plant in ethnoveterinary medicine, using a 4-AP induced seizure, electrophysiological model, <i>in vitro</i>	570
S2.P68	Centrifugal partition chromatography as a tool for the isolation of polyphenols from ginger rhizomes	571
S2.P69	Comparative assessment of the antioxidant capacity of Domat olive leaf and Sari Yaprak olive leaf	572
S2.P70	Syringin accelerated wound healing and reduced inflammation in human skin fibroblastst and keratinocytes <i>in vitro</i>	573
S2.P71	Exploring the Impact of Different Heat Extraction Processes on the phytochemical profile of <i>Cichorium intybus</i> Extracts	574
S2.P72	Modulation of Gastric Motility by a Traditional Herbal Preparation in Functional Dyspepsia: Implications of Muscarinic Receptor Inhibition	575
S2.P73	<i>In vitro</i> liver metabolism of four biflavonoids and their corresponding monomers	576
S2.P74	Screening of flavonoids and phenolic compounds in <i>Betula litwinowii</i> Doluch. by LC-Q- TOF/MS and determination of their antioxidant potential	577
S2.P75	The extracts of <i>Capparis cartilaginea</i> Decne attenuate inflammasome activation in LPS- stimulated THP-1 macrophages	578
S2.P76	<i>Pistacia chinensis</i> Bunge subsp. <i>falcata</i> (Becc. ex Martelli) aqueous extract possesses acetylcholinesterase and monoamine oxidase inhibitory activities	579
S2.P77	Cyclotides from Bolivian plants	580
S2.P78	Cytotoxicity assessment of <i>Rosa pisiformis</i> (Christ) D. Sosn. (Rosaceae) endemic to Turkiye	581
S2.P79	Confirming (-)-epicatechin's biological signature; universal modulation of NOS in chronic disease	582
S2.P80	Analytical development on <i>Pulsatilla</i> species: comparison of two species	583
S2.P81	Wound healing effects of <i>Arnica montana</i> formulations administered by different routes in mice	584
S2.P82	Effect of <i>Tormentillae tinctura</i> and its postbiotic metabolites on human intestinal epithelium cellular model	585
S2.P83	Effects of the Korean Herbal Prescription, Bojangggunbi-tang, on indomethacin--induced Small Bowel Injury focusing on mechanisms of action	586
S2.P84	Interaction of an herbal multi-component medicine for treatment of digestive disorders with the growth of selected intestinal lactic acid bacteria	587
S2.P85	<i>Vangueria infausta</i> Burch (wild medlar) extract improves semen quality by regulating testosterone levels and oxidative stress in testicular Leydig cells	588
S2.P86	Phytochemical, antioxidant, and androgenic effects of multi-solvent extracts from <i>Strychnos spinosa</i> Lam on mouse testicular TM3 Leydig cells	589
S2.P87	Unlocking longevity through the use of plant adaptogens: lifespan extension by Maral root (<i>Rhaponticum carthamoides</i> (Willd.) Iljin) extract in <i>Caenorhabditis elegans</i>	590
S2.P88	Revealing the veil of barberry alkaloids: high-throughput non-targeted UHPLC-DIA-HRMS profiling enhanced by solid phase extraction based on mixed-mode sorbent	591
S2.P89	<i>Viscum album</i> extracts in the perioperative oncological setting: evidence from bench to bedside	592
S2.P90	Potential benefits of <i>Malva sylvestris</i> in treating dry-eye pathology <i>in vitro</i>	593
S2.P91	The relationship among essential oil composition, morphological and climatic diversification - the case of <i>Teucrium montanum</i> from the Balkan Peninsula	594
S2.P92	Salicylic alcohol content in the leaves of different <i>Salix</i> species and individuals: impact on pharmaceutical potential	595
S2.P93	Plant metabolites as glucose-induced insulin secretion modulators	596
S2.P94	Antioxidative capacity of European tree bark extracts	597
S2.P95	Exploring the potential of ADAPT-232: plant adaptogens for healthy aging, and longevity in <i>Caenorhabditis elegans</i>	598
S2.P96	<i>Harpagophytum procumbens</i> DC. ex Meisn. hairy roots extract alleviates inflammatory response in psoriasis dermatitis	599
S2.P97	Caloric restriction mimicking through NRF-2-mediated mechanisms by plant-derived bioactive leads	600
S2.P98	<i>SKN1/NRF2</i> -dependent modulation of nutrient sensing network in <i>Caenorhabditis elegans</i> upon curcumin treatment	601
S2.P99	The phenylethanoid glycoside forsythoside B as a promising candidate for photoprotective dermatological products	602
S2.P100	Exploring the therapeutic potential of <i>Punica granatum</i> leaves extract in enhancing healthspan and lifespan in <i>Caenorhabditis elegans</i>	603
S2.P101	Baicalein, a major component of the root of <i>Oroxylum indicum</i> (Bignoniaceae) vasodilates isolated pulmonary arteries through endothelium-dependent and -independent mechanisms	604
S2.P102	Chemical profile and cytotoxic activity of methanol extracts of eight <i>Geranium</i> L. species	605
S2.P103	The composition of the essential oil of <i>Pimpinella serbica</i> and its relationship to other <i>Pimpinella</i> species from the central part of Balkan Peninsula	606
S2.P104	Different extracts of three <i>Veronica</i> species: the study of their antioxidant activity	607
S2.P105	Anti-inflammatory potential of chamomile flowers (<i>Matricariae flos</i>) extract in topical applications	608
S2.P106	A comparative study on biological activities of different extracts of <i>Scrophularia xanthoglossa</i> Boiss	609
S2.P107	Andrographolide as a potential anti-inflammatory molecule in prostate cancer	610
S2.P108	The impact on proinflammatory cytokines secretion by human neutrophils and keratinocytes by extracts from <i>Arctii lappae fol</i> , <i>Serpylli hba</i> and <i>Millefolii hba</i> - raw materials traditionally used in Poland in skin disorders	611
S2.P109	Enhancing the anticancer activity of oleander from <i>Nerium oleander</i> L. (Apocynaceae) through novel formulations	612
S2.P110	Oral acute toxicity and preliminary phytochemical characterization of <i>Hyptis albida</i>	613

S2.P111	<i>Ex vivo</i> assessment of potential <i>Eucalyptus cinerea</i> antihypertensive	614
S2.P112	The anti-asthmatic effect of artichoke in BALB/c mouse model	615
S2.P113	Antimicrobial activity and chemical composition of <i>Taraxacum aleppicum</i> Dahlst. (Asteraceae)	616
S2.P114	The hydroethanolic extract from <i>Calendula officinalis</i> L. herba demonstrated potent anti- inflammatory properties <i>in vitro</i>	617
S2.P115	A study on the antiproliferative activity of the triterpenoid constituents in <i>Erica erigena</i>	618
S2.P116	Cyclic peptides derived from sunflower trypsin inhibitor 1 (SFTI-1) for autoantibodies neutralization in rheumatoid arthritis	619
S2.P117	Detailed LC-MS analysis and evaluation of cytotoxic effects of various extracts obtained from aerial parts of <i>Glaucium acutidentatum</i> Hausskn. & Bornm	620
S2.P118	<i>In vivo</i> evaluation of the anti-inflammatory activity of the marine pigment echinochrome A on hairless mice skin	621
S2.P119	Anti-inflammatory effect of <i>Glaucium flavum</i> extracts	622
S2.P120	Alpha-glucosidase inhibitory profile of sesquiterpenes lactones from <i>Centaurea</i> species	623
S2.P121	Phytochemical profiling and <i>in vitro</i> biological activities on <i>Myrtus communis</i> L.	624
S2.P122	Dietary molecules from <i>Glycyrrhiza foetida</i> and their modulation on metabolic syndrome pathways	625
S2.P123	Smoking hot – the story of the “pepper bark tree” <i>Warburgia salutaris</i>	626
S2.P124	The effect of cannabidiol, cannabidiol and cannabigerol on the motoric activity of the rat proximal colon – <i>ex vivo</i> study	627
S2.P125	Coumarin polypharmacology and elucidation of an underlying molecular mechanism of action	628
S2.P126	Deciphering the targets of trypanocidal chalcones in <i>Trypanosoma cruzi</i>	629
S2.P127	Innovative methods for analysing flavonolignan complex from <i>Silybum marianum</i> L.	630
S2.P128	Phytochemical analysis of <i>Hyssopus officinalis</i> L. cultivated in western Turkey	631
S2.P129	Enhancing wound healing: sericin and <i>Chelidonium majus</i> L. as potential dressings	632
S2.P130	Evaluating the efficacy of topical application of <i>Genista tridentata</i> (L.) (Fabaceae) leaf extract on wound healing in experimental diabetes	633
S2.P131	<i>Lepidium meyenii</i> – isolation of imidazole alkaloids by centrifugal partition chromatography	634
S2.P132	Connecting targets of the standardized herbal combination STW 5-II to gastrointestinal disorders	635
S2.P133	The anti-inflammatory and skin barrier function recovery effects of <i>Schisandra chinensis</i> in mice with atopic dermatitis	636
S2.P134	Comparison of amount of tannins in Ericaceae plants using different water extraction methods	637
S2.P135	Biosynthetic relationship and self-assembly nanoparticles of royleanone isolated from <i>Plectranthus</i> species	638
S2.P136	Yamogenin – a steroidal plant metabolite triggers extrinsic and intrinsic pathway of apoptosis in gastric cancer cells	639
S2.P137	Phytochemical analysis and neuroprotective activity of extracts from <i>Paliurus spina-christi</i> (Rhamnaceae)	640
S2.P138	Bioactive molecules separation from <i>Acmella oleracea</i> (Jambu) from Amazon Florest	641
S2.P139	The protective impact of <i>Aronia melanocarpa</i> L. berries extract against cadmium-induced changes in the concentrations of calcitropic hormones in the serum	642
S2.P140	LC-MS/MS analyses of triterpenes and polyphenols from less-polar fractions of <i>Rosa rugosa</i> leaf extract and their biological activity	643
S2.P141	Volatiles, triterpenes, phytosterols and anti-inflammatory activity of <i>Rhododendron luteum</i> Sweet (Ericaceae) supercritical CO2 extracts	644
S2.P142	Antioxidant and anti-inflammatory activity of the <i>Berberis vulgaris</i> L. bark extract	645
S2.P143	The use of herbal medicines for gynecological ailments: patient-reported outcomes	646
S2.P144	An analysis of real-world data regarding therapeutic efficacy and adverse effects of herbal medicines in mental health issues	647
S2.P145	Antiproliferative activity of homolupane triterpenoids	648
S2.P146	Anticancer and antioxidant properties of extracts from <i>Rubus caesius</i> L. (Rosaceae)	649
S2.P147	Broad antiplatelet activity of the flavones zapotin, chrysin, and tropoflavin in human platelet - rich plasma	650
S2.P148	Anti-aging, anti-neurodegenerative and healthspan-promoting compounds from <i>Dendrobium officinale</i> Kamura et Migo (Orchidaceae)	651
S2.P149	Effects of STW 3-VI (St. John's wort) on the paracrine signaling between neurons and microglia under pro-inflammatory conditions <i>in vitro</i>	652
S2.P150	Gut microbiota - a new aspect for understanding the antidepressant effects of <i>Hypericum perforatum</i> ?	653
S2.P151	The cytotoxic effect of chrysosplenetin, a natural product from <i>Artemisia annua</i> , investigated with different human cancer cell lines, as well as <i>in vitro</i> CYP450 enzymatic activity in human liver microsomes	654
S2.P152	Bioprospecting for plant extracts with protective effects on endothelial function	655
S2.P153	A purified fraction from 70% hydroethanol extract of whole herb of <i>Platostoma africanum</i> P. Beauv. (Lamiaceae) inhibited formation of β -haematin formation	656
S2.P154	Studies of St. John's Wort (<i>Hypericum perforatum</i> L.) dry extracts: no hints on mutagenicity	657
S2.P155	Shortcutting iridals to irones transformation in orris root: unleashing the potential of microwave drying for rapid irone production	658
S2.P156	Didesmethylocaglamide cytotoxic activity in high grade serous ovarian cancer in preclinical <i>in vitro</i> and <i>in vivo</i> models	659
S2.P157	The herbal preparation, STW 5, protects against changes in the microbiome in experimental functional dyspepsia	660
S2.P158	Modulatory effects of <i>Glycyrrhiza glabra</i> L. on endocrinological shift in ovariectomy-induced menopausal mice: Network pharmacology guided <i>in-vivo</i> study	661
S2.P159	<i>Ziziphus nummularia</i> crude and fractionated extracts attenuate the malignant phenotype of human triple negative breast cancer cells	662
S2.P160	Anti-inflammatory properties of jaboticaba peel bioactive compounds during <i>in vitro</i> gastrointestinal and colonic digestion	663
S2.P161	Improving sleep quality and daytime function with a synergistic combination of tryptophan, magnesium, <i>Lactuca</i> , and <i>Melissa</i> : A prospective pilot study in adults with sleep disturbances	664
S2.P162	Harnessing gut microbiota through immunomodulation by LM-1 attenuates HDM-induced asthma	665
S2.P163	Ethnopharmacological study of the bark traditionally used for diabetes of <i>Croton guatemalensis</i> Lotsy: from the field to the action mechanisms	666
S2.P164	<i>Eryngium cymosum</i> F. Delaroché, a Mexican species traditionally used in the treatment of diabetes - ethnopharmacological insights	667
S2.P165	Onychocolone A produced by the fungus <i>Onychocola</i> sp. targets cancer stem cells and stops pancreatic cancer progression by inhibiting MEK2-dependent cell signaling	668
S2.P166	Human prolyl endopeptidase inhibitory activity of the peptide enriched extracts of <i>Momordica charantia</i> (Cucurbitaceae) fruits	669
S2.P167	Impact of phytocannabinoids THC, CBD and their acidic counterparts THCA and CBDA on TGF- β 1-induced epithelial-mesenchymal transition in human conjunctival epithelial cells	670
S2.P168	Development of the immunopharmacognostic interventions on potentiating tumor recognition against prostate cancer	671
S2.P169	Investigating the anti-inflammatory effects of a TCM formulation used for the prevention of rheumatoid arthritis	672
S2.P170	Biological potential of endophytes from liverwort <i>Marchantia polymorpha</i> L.	673
Natural product research and the challenges of climate change – ecological and environmental solutions		674
S2.P171	Small molecule metallophores from bacteria, adaptation and isolation strategies	674

S2.P172	Exploring Tuscany onion (<i>Allium cepa</i> L.) heritage: unveiling ancient varieties through molecular network analysis and bioinformatic tools	675
S2.P173	Extraction of <i>Cannabis sativa</i> : subcritical water vs. supercritical CO ₂ and organic solvents	676
S2.P174	Response surface methodology guided RP-HPLC-PDA method optimization for the quantification of multi-analytes (six) in Indian kudzu and its greenness assessment	677
S2.P175	Development and validation of RP-HPLC method for estimation of anti-oomycete diterpenoids in <i>Larix decidua</i> extracts	678
S2.P176	Eutectic solvents as a sustainable solution for the recovery of high value free fatty acids from microalgal co-products	679
S2.P177	EKO-YAM - investigating <i>Dioscorea</i> for a sustainable future	680
S2.P178	Wound healing and antimicrobial activity of European tree bark extracts	681
S2.P179	An ethnobotanical survey investigating traditional remedies to treat infectious diseases in French Polynesia	682
S2.P180	Water as a green solvent to extract polar and non-polar plant pigments	683
S2.P181	Long term experiments on applying a pulsed electric field (PEF) to living plant roots in order to extract flavonoids in <i>S. baicalensis</i> Georgi model	684
S2.P182	Enhancing fire ant bait selectivity: Synthesis and evaluation of feeding deterrent activity of fire ant venom alkaloids and their analogs against invasive fire ants and native ants	685
S2.P183	Sucrosephenylpropanoid esters and isoflavonoids from <i>Belamcanda chinensis</i> (Iridaceae) roots and their potential anti-osteoclastogenic activity	686
S2.P184	PTP1B inhibitory activity of serratene-type triterpenoids from <i>Lycopodium serratum</i> (Lycopodiaceae)	687
S2.P185	Introducing NaDES-based biphasic solvent families inspired by the Arizona family: a new approach for the sustainable use of centrifugal partition chromatography	688
S2.P186	Beyond the tuber: crafting high-value molecules from potato residues	689
S2.P187	Mulberry cell cultures: a promising source for health care	690
S2.P188	Spatio temporal dynamics of favourable habitats for <i>Terminalia leiocarpa</i> , a traditional plant used in ethnoveterinary medicine in Benin (West Africa)	691
S2.P189	From the production of targeted metabolites to the complete valorization of <i>Coleus forskohlii</i> Briq. using aeroponic cultivation	692
S2.P190	ViscY NMR experiments combined with chiral discrimination for complex mixture analysis	693
S2.P191	A green approach for preparing alkaloid-rich extracts and fractions	694
S2.P192	Optimization of a green method for quantifying harmine in <i>Banisteriopsis caapi</i> stems used in the preparation of ayahuasca	695
S2.P193	Nutrient density and mineral content of radishes grown in black soldier fly frass-enriched compost	696
S2.P194	Myristoleic acid ameliorates the inhibition of rat dermal papilla cell growth by fine particulate matter via the activation of autophagy and Wnt/ β -catenin pathways	697
S2.P195	Identification of phenolic compounds in three <i>Veronica</i> species: <i>Veronica anagallis-aquatica</i> L., <i>Veronica persica</i> L. and <i>Veronica polita</i> L.	698
S2.P196	Turning the Tide: Harnessing the Invasive algae <i>Rugulopterix okamurae</i> for Commercial Applications	699
S2.P197	Scale-up production of g-mangostin using Centrifugal Partition Chromatography	700
S2.P198	<i>In vitro</i> callus culture of <i>Mespilus germanica</i> to improve the production of bioactive compounds from a forgotten fruit	701
S2.P199	The phytochemical diversity of wild hops from northern France for potential brewing application	702
S2.P200	Effect of the aquaponic system on productivity and quality of <i>Acmella oleracea</i> R.K. Jansen	703
S2.P201	Comparison of free volatile compounds of wild and cultivated species <i>Veronica catenata</i> Pannell	704
S2.P202	Urban parks biowaste as a sustainable source of new antidiabetics	705
S2.P203	Unravelling the secrets of fungicides through actinobacteria-phytopathogen interactions	706
S2.P204	The involvement of cyclotides in plant response to heavy metals in <i>Viola</i> spp.	707
S2.P205	Applying the global nature products social molecular networking (GNPS) platform to identify natural products involved in tolerance to heavy metals in <i>Viola</i> spp.	708
S2.P206	Screening plant extract for biological activity. A work spanning 60° Latitude South	709
S2.P207	Polyphenol and anthocyanin content in early producing maqui hybrids (<i>Aristotelia chilensis</i>)	710
S2.P208	Effect of different ethylene levels on other phytohormones and <i>Arabidopsis</i> resilience	711
S2.P209	Copper-induced siderophore production and copper complex formation in vitamin-producing bacteria at a Vermont copper mine	712
S2.P210	Disentangling metabolite dynamics in extreme environments: life detection strategies for extraterrestrial worlds	713
S2.P211	Effect of green-based extraction approaches on antioxidant activity and phenolic compounds of Oregano (<i>Origanum vulgare</i> ssp. <i>hirtum</i> (Link) Ietswaart)	714
S2.P212	Enhancing polyphenol production in <i>Cistus albidus</i> L. through strategic mycorrhization and irrigation management: An off-season approach	715
S2.P213	The potential of chitosan from <i>Hermetia illucens</i> as a sustainable biopesticide	716
S2.P214	Screening of Nettle Clones with the Ph Eur HPTLC method for identification testing	717
S2.P215	Creation of a medicinal plants collection and research consortium to catalyze drug discovery in San Diego and beyond: An update from 2022-2024	718

Regulation for the use of natural products in food and drugs

S2.P216	The cannabis/marijuana (<i>Cannabis sativa</i> L.) landscape in Africa: an overview of its cultivation and legal aspects	719
S2.P217	A comparative toxicity study depending on type of clinical use of the roots of <i>Asarum heterotropoides</i> var. <i>mandshuricum</i> (Maxim.) Kitag	720
S2.P218	Determination of triterpenoids of the native <i>Centella asiatica</i> L. from the Jeju Island, South Korea	721
S2.P219	Streoselective inhibitory effects of ginsenosides on six uridine 5'-diphosphoglucuronosyl transferases in human liver microsomes	722
S2.P220	Discrimination of herbal medicines originated from soy beans (<i>Glycine max</i> Merrill.) using high-performance thin-layer chromatography	723
S2.P221	Analytical Quality by Design (AQbD) methodology approach for simultaneous quantification of Hesperidin and Magnolol in Daikumeum-ja using HPLC-PDA	724
S2.P222	Discrimination between similar herbal medicines of Campanulaceae by high-performance thin layer chromatography (HPLC)	725
S2.P223	Quantification of saponin-rich plants mixture in complete feed with UHPLC-MS/MS using a "one-point" calibration	726
S2.P224	Interactions of ivy (<i>Hedera helix</i>) with other herbs or medicines: a systematic review	727
S2.P225	Brazilian regulation for food supplements containing bioactive substances from plant extracts	728
S2.P226	Chemical profile of natural musk (<i>Moschus moschiferus</i>) by GC-MS/MS	729
S2.P227	Advances in the Brazilian Regulation for the use of plant extracts in medicines	730
S2.P228	Comparing the volatilome of Spanish-style and natural cv. Chalkidiki green table olives at industrial scale	731
S2.P229	Addressing regulatory challenges and ensuring safety in the e-commerce of herbal remedies in Mexico	732

S2.P230	Questions for consumer reporting on risks and benefits from herbal medicine usage: A preliminary international Delphi study	733
S2.P231	Analysis of the chromatographic profile and <i>in vitro</i> antiurolytic activity of species of the genus <i>Phyllanthus</i> (Phyllanthaceae), endemic and non-endemic of the flora of Santa Catarina	734
S2.P232	Hot chips - is it a real 'challenge'?	735
S2.P233	Botanical adventure in pharmacy education: implementing an escape game for quality control training	736
S2.P234	Call for a contextual reform of the regulations on traditional herbal medicine in Burkina Faso	737
S2.P235	The <i>in vitro</i> modulatory effects of <i>Artemisia sieberi</i> on phase-I cytochrome P-450 isoenzyme gene expression using human hepatoma cells (HepG2)	738
S2.P236	Comparison of qNMR and HPLC-UV techniques for quantitation of secoiridoid phenolics in olive oil	739
S2.P237	Anticancer efficacy of Korean medicine and sorafenib (concomitant drug) for hepatic tumors	740
S2.P238	<i>Erythrina</i> spp. as natural anti-inflammatory ingredients for cosmeceuticals: An <i>in-silico</i> study	741

Medicinal Plants and Natural Products in Animal Healthcare and Veterinary Medicine

742

S2.P239	Chemical composition of resin extract from <i>Corymbia</i> spp. an indigenous Australian native plant	742
S2.P240	<i>Melissa officinalis</i> (Lemon balm) extract to promote calming effect before stressful events	743
S2.P241	Equine Stromal Cells: effects of non-ribosomal peptides on proliferation and viability	744
S2.P242	Extracts from the roots of <i>Scutellaria baicalensis</i> reduce honeybee death caused by nosemosis	745
S2.P243	Larch (<i>Larix decidua</i> L.) fibre waste: characterization of a functional feed ingredient	746
S2.P244	Identification of novel caffeoyl- and feruloyl-derivatives in <i>Solanum glaucophyllum</i>	747
S2.P245	Phytochemical screening and study of the hypotensive effect of the aqueous extract of <i>Rubus ulmifolius</i> Schott leaves	748
S2.P246	Evaluating the antibacterial properties of macroalgae against swine pathogens	749
S2.P247	A survey-based study on knowledge, attitude, and perception of pet owners towards the use of plant extracts from bio-refinery in pet food formulations	750
S2.P248	Anti-inflammatory effects of Cinnamon essential oil assessed on diverse cell models	751
S2.P249	Induction of apoptosis in pancreatic cancer cells by <i>Albizia julibrissin</i> extract through inhibition of an oncogenic signaling pathway	752
S2.P250	Antibacterial, antibiofilm and quorum sensing inhibitory activities of fractions of the ethanol extract of <i>Searsia lancea</i> against pathogens causing bovine mastitis	753
S2.P251	Potential use of <i>Paulownia</i> leaves as feed additive to enhance animal health condition	754
S2.P252	Exploring ethnoveterinary medicine in dogs in Ireland, Norway and Poland	755
S2.P253	Polish perspective on phytotherapeutic approaches to pet anxiety: an analysis of popular preparations	756
S2.P254	Anti-tumor effects of <i>Auricularia polytricha</i> extracts on human breast cancer cell lines	757
S2.P255	Analysis of veterinary historical German-language textbooks regarding medicinal plants used for the treatment of gynecological diseases and fertility disorders	758
S2.P256	Phyto-synergy: a holistic approach to tackle <i>paratuberculosis</i>	759
S2.P257	Preliminary study of the neuroprotective capacity of <i>Corema album</i> (L.) D.Don ex Steud (Ericaceae) juice and its butanolic fraction	760
S2.P258	Anti-inflammatory and antioxidant ability of <i>Vitis vinifera</i> L. leaves	761
S2.P259	Pre- and post- treatment effect of water-soluble phenolics from date palm (<i>Phoenix dactylifera</i> L.) fruits in cisplatin-induced hepatotoxicity in rats	762
S2.P260	Determination of anticancer properties of <i>Moringa oleifera</i> seed bioactive fraction and the interaction effect of its combination with the classical cytotoxic drugs	763
S2.P261	Fertility parameters of dairy cows after intrauterine treatment of clinical endometritis with an antibiotic or an herbal veterinary medicinal product	764
S2.P262	Medicinal plants for animal health care: translating tradition into modern veterinary medicine - the COST Action MedPlants4Vet	765
S2.P263	Effectiveness of replacing antibiotics with botanicals in the treatment of bovine mastitis	766
S2.P264	Pumpkin seed cake waste as a sustainable functional feed ingredient	767
S2.P265	Herbal bioenhancers: an example of inhibition of ABCB1 using <i>Brugmansia</i> tincture	768
S2.P266	Antibacterial activity of essential oils against streptococci and staphylococci causing bovine mastitis	769
S2.P267	Absorption, distribution, accumulation and excretion of thymol after its sustained administration in rabbits	770
S2.P268	<i>In-vitro</i> investigation of the gastroprotective effects of a natural extract blend: a promising avenue for canine gastritis prevention	771
S2.P269	Acaricidal activity of essential oil of <i>Satureja montana</i> L. against <i>Dermanyssus gallinae</i> <i>in vitro</i> conditions	772
S2.P270	A novel phytogenic supplement that prevents coccidiosis in chickens	773

Poster Session 3, Tuesday, July 16, 2024

774

Cutting edge technologies in natural product drug discovery, formulation and development

774

S3.P1	<i>In silico</i> investigation of <i>Pterocarpus marsupium</i> against type-2 diabetes mellitus through network pharmacology approach	774
S3.P2	Integration of High-Throughput Screening, AI-ML Strategies, and ECD/DP4+ for the Discovery and Elucidation of Novel Natural Products	775
S3.P3	On-tissue derivatization and MALDI-MS Imaging reveal hemlock alkaloid distribution in <i>Conium maculatum</i>	776
S3.P4	New world and old world <i>Salvia</i> species (sages): searching of diterpenoids with cytotoxic activity	777
S3.P5	Use of <i>Galleria</i> moth larvae model to assess the pathogenesis of AIEC and the therapeutic potential of Geraniol as an antibacterial agent	778
S3.P6	Inhibition of transforming growth factor β signaling by prodigiosin through disruption of receptor recycling and subcellular translocation in epithelial cells	779
S3.P7	Assessing intra- and inter-variability of Macedonian wild-types and commercial <i>Cannabis</i> strains using short tandem repeat markers	780
S3.P8	Preclinical development of natural products into agents for pediatric brain cancers	781
S3.P9	Exploring the anti-mycobacterial potential of French Guiana's flora through a metabolomic study	782
S3.P10	Arnica of Provence: an alternative to <i>Arnica montana</i> L.? Investigations about polyphenolic composition and antioxidant potential	783
S3.P11	Biosynthesis of anti-inflammatory compounds from <i>Calendula officinalis</i>	784
S3.P12	Development and validation of an analytical method for the analysis of standardized red propolis extract and its chemopreventive effect against colon carcinogenesis	785
S3.P13	Extract rich in polyprenylated benzophenones from Brazilian red propolis: obtaining, analytical characterization and <i>in vitro</i> evaluation against breast cancer lines	786
S3.P14	Computational NMR enhances metabolomic analysis and sustainability	787

S3.P15	Exploring on the networking of chemical compositions and their cytotoxicity toward KBCC <i>Nepenthes</i> plants	788
S3.P16	Chemical composition and bioactivity of <i>Uvaria micrantha</i> from Vietnam	789
S3.P17	Ontology based ethnobotany data to support natural product-based drug discovery – growth of data, large-scale insights and case studies	790
S3.P18	Optimization formulation of <i>Rhodiola rosea</i> L. nanoemulsion containing natural oil using response surface method (RSM): preparation, characterization, stability test and antioxidant activity	791
S3.P19	Oxidoreductase FSP1-mediated redox-regenerating natural product E18C6 potentially suppresses ferroptotic cell death	792
S3.P20	Chinese traditional medicine for preventing hair loss in androgenic alopecia mice	793
S3.P21	Qualitative and semi-quantitative screening of triterpenes in <i>Vaccinium uliginosum</i> L. fruits using supercritical fluid chromatography	794
S3.P22	Matrix free laser desorption ion mobility mass spectrometry in chemometrics: A new approach to the rapid identification of activity markers in complex mixtures of natural products	795
S3.P23	Metabolomic study of tea tree oil and <i>Melaleuca</i> spp. using HPTLC, LC-MS and GC-MS	796
S3.P24	Preclinical evaluation of oral nanoformulation based on natural extract as a potential treatment for type 2 diabetes mellitus	797
S3.P25	Employing Plant-Bacterial Interactions to Access the Metabolic Capacity of Bacteroidetes	798
S3.P26	<i>Epilobii</i> herba: Quality assessment using chromatographic techniques	799
S3.P27	Phytochemical analysis and quality control study of <i>Dendrobium fimbriatum</i> Hook. flowers	800
S3.P28	Identification of chemical defenses of a tropical tree <i>Sextonia rubra</i> using molecular networks from GC-MS, LC-MS/MS and MALDI-MS/MS imaging data	801
S3.P29	An ethnobotanical, pharmacological and phytochemical review on flowers of Shi-hu (<i>Dendrobium</i> spp.)	802
S3.P30	Patients' use of medicinal plants in the urology service of Ibn-Sina hospital in Rabat, Morocco: integrating ethnobotanical and pharmacovigilance approaches	803
S3.P31	Unveiling the multifaceted efficacy of Gu Sui Bu in osteoporosis treatment: insights from a zebrafish model and advanced analytical approaches	804
S3.P32	Phytochemical profiling and the wound healing potential of five <i>Bulbine</i> species using <i>in vitro</i> and <i>in vivo</i> assays	805
S3.P33	Translating community-wide mass spectral libraries into actionable chemical knowledge: a proof of concept with monoterpene indole alkaloids	806
S3.P34	Non-targeted profiling analysis by GC/MS and LC/MS/MS of extracts derived from <i>Eucalyptus tereticornis</i> with antiobesogenic and antidiabetic activity	807
S3.P35	Design, synthesis and screening of novel prenylated chalcones as optimised anti-cancer agents	808
S3.P36	Does the combined effect of natural abietane diterpenes with temozolomide have therapeutic potential against glioblastoma?	809
S3.P37	Assay-directed method development for natural product purification	810
S3.P38	Comparative analysis of the anti-inflammatory properties of nanoparticles derived from the medicinal plants <i>Zingiber officinale</i> (Ginger) and <i>Caralluma edulis</i> (Chunga), in the treatment of Inflammatory Bowel Disease	811
S3.P39	Isolation, characterization, and bioactivity of compounds from the stem bark of <i>Prunus africana</i>	812
S3.P40	Lanostane triterpenoids of <i>Inonotus obliquus</i> are modulators of inflammation regulators ROR γ and GPBAR1	813
S3.P41	COSMO-RS as cutting-edge technology for design and pre-screening of Natural Deep Eutectic Solvents for selective extraction: Spirulina as case study	814
S3.P42	Tetrahydrofuran sesquiterpenes and the bronchodilator effect of <i>Artemisia judaica</i>	815
S3.P43	Development of non-targeted metabolomic approaches for the screening of bioactive molecules in <i>Clematis</i> species	816
S3.P44	Semisynthetic pyrazole ring containing hydrocurcumin derivatives as potential new antitumor agents	817
S3.P46	A decision theory-based aggregation tool for tandem mass spectrometry multi-annotation outputs	818
S3.P47	<i>Plectranthus</i> : an important source of bioactive diterpenoids for therapeutic applications	819
S3.P48	Characterization of volatile and non-volatile compounds in <i>Larix decidua</i> (Pinaceae) oleoresin and its nanoemulsion development	820
S3.P49	<i>In vitro</i> antihypertensive effects of <i>Viscum album</i> mother tinctures in kidney proximal tubule cells	821
S3.P50	^1H NMR-based metabolomics on 30 east asian traditional herbal medicines for quality control	822
S3.P51	Unveiling terpenoid diversity in <i>Sideritis</i> spp. through molecular networks	823
S3.P52	Targeted discovery of sesquiterpene indole alkaloids from Gabonese <i>Greenwayodendron suaveolens</i>	824
S3.P53	The benefit of classification machine learning in virtual drug discovery for beta-lactamase inhibitors identification based on an in-house library	825
S3.P54	A bioactive chlorophyll-derived yellow phyllobilin from senescent leaves of <i>Carica papaya</i> (Caricaceae) and its potential role in the ethnopharmacological use of withered leaf extracts	826
S3.P55	Evaluating the use of Raman spectroscopy for the characterization of volatile compounds by direct measurement on the oregano leaf glands	827
S3.P56	New benzothiazines with pain-reliving activity from marine bacterium <i>Croceibacter atlanticus</i>	828
S3.P57	3D printed polymeric scaffolds loaded with antibiotic and anti-inflammatory drugs for tissue engineering applications	829
S3.P58	ChromAnnot – a webserver for reproducible LC-HRMS/MS chromatogram annotation	830
S3.P59	Exploring the anticancer potential of <i>Polygonum barbatum</i> (L.) against colorectal cancer: <i>In vitro</i> and <i>in vivo</i> investigations	831
S3.P60	Physical interactions between <i>Streptomyces</i> (Streptomycetaceae) and competing microbes trigger the production of natural products	832
S3.P61	Energy Saving Efficacy and Optimized condition of pulse energy fields technology on <i>Centella asiatica</i> extraction	833
S3.P62	Identification of a leoligin-inspired synthetic lignan as a novel TGR5 agonist	834
S3.P63	Nano-phytosome delivery system loaded with <i>Crataegus laciniata</i> (Rosaceae) flowers extract: whitening properties	835
S3.P64	Leoligin derivatives as novel phosphodiesterase 4 inhibitors	836
S3.P65	The search for nontraditional synergy: pairing <i>in vitro</i> omics and <i>in silico</i> machine learning methods to discover novel combinations of antimicrobial agents from natural products	837
S3.P66	Electron diffraction-guided natural product discovery: identification and isolation of new scaffold class from <i>Podospora australis</i>	838
S3.P67	Metaboseek 2.0: Deconvoluting the complexity of MS/MS Big Data through interactive data processing	839
S3.P68	Cytotoxic activities of terpenoids from genera <i>Euphorbia rowlandii</i> and <i>Euphorbia grandocornis</i> (Euphorbiaceae) against the Triple-negative breast cancer (TNBC)	840
S3.P69	Simultaneous extraction and synthesis of new cannabinoid acid esters and their evaluation on TRPA1, cell proliferation and in MDA-MB-231 xenograft model of cancer	841
S3.P70	Exploring potential breast and melanoma cancer drug candidates from <i>Crocus sativus</i> : molecular docking insights	842
S3.P71	Metabolic fate of oleocanthal and oleacein after oral consumption: Novel biomarkers, human bioactivity insights and TRPA1 channel activation	843
S3.P72	An innovative methodology for the production of a new food supplement enriched in oleocanthal and oleacein with multi target protective properties	844

S3.P73	Online dried spot LC-Orbitrap analysis for the quality assessment of cannabis products	845
S3.P74	Anti-prostatic effects of <i>V. album</i> L extracts are influenced by season and the use of hydrogel delivery system	846
S3.P75	A “head-to-tail” approach towards foodomics advancement: LC-HRMS and NMR correlation in the case of olive products	847
S3.P76	Systematic cheminformatics analysis of herbal plants in malawi: uncovering chemical scaffolds with anti-candidal potential	848
S3.P77	From nature's palette to drug candidates: exploring the chemodiversity of <i>Piper fimbriatum</i> and expanding the bioactive potential for the development of novel drug leads	849
S3.P78	Inhibition of autotaxin by quercetin and its derivatives	850
S3.P79	Exploring vibrational spectroscopic approaches for quantifying main phytocannabinoids in medicinal <i>Cannabis sativa</i> inflorescence	851
S3.P80	Exploring the therapeutic synergy: comprehensive assays unveil the potential of sericin and <i>Chelidonium majus</i> L. in developing a possible wound dressing	852
S3.P81	Synergistic impact of natural stilbene-linked mono-substituted nitroarenes in hormone- dependent breast cancer: a greener approach	853
S3.P82	The natural diversity of acyltransferases reveals versatility and specificity in the synthesis of gene-encoded lipopeptides	854
S3.P83	Exploring new dimensions: analytical method development for enhanced quality control of natural products	855
S3.P84	Simultaneous quantification of terpenes and cannabinoids in <i>Cannabis sativa</i> by reversed phase LC-APCI-MS/MS	856
S3.P85	LumiNose: a novel instrument to apply the receptomics biosensor technology for identifying and quantifying medicinal bioactivities in plant extracts	857
S3.P86	pH-Dependent degradation of andrographolide; kinetics, product identification and anti- inflammatory and cytotoxic activity assessment	858
S3.P87	Immunohistochemical evaluation of the effect of <i>Kigelia africana</i> stem-bark extracts on benign prostatic hyperplasia induced by testosterone and estradiol in albino Wistar rats	859
S3.P88	Styryl lactone derivatives and aristolactam alkaloids from <i>Goniiothalamus tapis</i> Miq. and their α -glucosidase inhibitory activity	860
S3.P89	Exploring synergies between hemp cannabinoids and β -caryophyllene on triple-negative breast cancer cell line	861
S3.P90	Response Surface Methodology as a tool to obtain a potential satiating bitter enriched extract from <i>Gentiana lutea</i> L.	862
S3.P91	The Botanical Safety Consortium: collaborative effort to improve botanical safety methods	863
S3.P92	Natural digestion mimicking curcuminoid formulation enhances oral bioavailability of curcuminoids	864
S3.P93	Reveal the structure of plants by X-ray imaging and biomedical beamline at SSRF	865
S3.P94	A cyclodextrin based tertiary complex of curcuminoids to improve bioavailability -its basic characteristics and immunomodulatory properties	866
S3.P95	Phytochemical analysis of artichoke (<i>Cynara scolymus</i>) herb: Implications for <i>in vitro</i> neuroprotective and anti-obesity potential	867
S3.P96	A preliminary study investigating the wound healing potential of <i>Cedrus brevifolia</i> extracts in mice	868
S3.P97	Preformulation studies of lycopene-rich tomato extract	869
S3.P98	Investigating the mechanism of action of the anti-aging ability of geraniin nano-phytosome	870
S3.P99	QbD approach for the development of CBD-loaded polymeric micelles for oral delivery	871
S3.P100	Native metabolomics for the discovery of new small molecule gai protein modulators	872
S3.P101	NMR-HetCA uncovers bioactive secondary metabolites from propolis extracts prior to isolation	873
S3.P102	Discovery of new bioactive compounds: cloning, activation and engineering of specialized metabolites biosynthetic gene clusters	874
S3.P103	<i>Origanum dictamnus</i> infusion characterization by HPLC-PDA-MS and nanoformulation development	875
S3.P104	Extract from the seedlings of <i>Brassica oleracea</i> var. <i>italica</i> (Brassicaceae) reduces tissue damage and alters inflammatory markers over time: development of new medications	876
S3.P105	Identification of naturally-derived lipids as GPR84 agonists	877
S3.P106	Quantitative analysis of endogenous steroids by UPLC-MS/MS in various medicinal plant species	878
S3.P107	Enhancing drug discovery efficiency in plant extracts through polyphenol removal	879
S3.P108	Bioactive <i>Lycopodium</i> alkaloids from clubmoss cultivars and transformation to enhance anti- acetylcholinesterase activity	880
S3.P109	Depicting the chemical diversity of bioactive meroterpenoids produced by the largest organism on earth	881
S3.P110	Comparative <i>In Vitro</i> enzyme inhibitory activities of different plant parts of five <i>Albizia</i> species (Fabaceae)	882
S3.P111	Study and nutritional screening of some notanicals used as herbal remedies in managing eye and ear ailments in Ado-Odo Ota Local Government Areas, Ogun State	883
S3.P112	New data about chemical composition of <i>Cynanchum acutum</i> L.	884
S3.P113	A comparative study between CPC and flash chromatography for natural product purification	885
S3.P114	Molecular docking and dynamics simulations study of Cathecin from <i>Mentha cordifolia</i> as potential HIV-1 protease pnhibitor and Tat1-Cyclin competitor	886
S3.P115	Phytochemical profiling of <i>Acalypha australis</i> and screening for its most potent anti-inflammatory compounds with bio-affinity ultrafiltration LC-MS	887
S3.P116	The natural products magnetic resonance database (NP-MRD): comprehensive resource for natural products NMR data enabling discovery and understanding	888
S3.P117	Exploring the impact of moderate water stress on anthocyanin, flavonol glycosides and terpene dynamics across fruit development and ripening in <i>Pistacia lentiscus</i> L.	889
S3.P118	Isolation and Characterization of novel compound from <i>Anaphalis triplinervis</i> , its HPTLC quantification and antioxidant potential	890
S3.P119	Trapping and detecting electrophilic metabolites of botanical natural products as glutathione conjugates	891
S3.P120	Validation of artificial intelligence tools for expedited innovation and globalization of medicinal plants in the dietary supplement industry	892
S3.P121	Modified proanthocyanidin DESIGNER materials with dentin biomodification activity	893
S3.P122	Antimicrobial activity of 2'-hydroxychalcones with a chlorine atom and their glycosides	894
S3.P123	Novel ardisiphenol D derivatives 1279P1 and 1279P2: promising anti-colorectal cancer agents targeting the PI3K/Akt pathway and lipogenesis enzymes	895
S3.P124	Antifibrotic activity study of natural stilbenoids from <i>Dendrobium nobile</i> Lindl	896
S3.P125	Planar chromatography method development for the herbal metabolomic study on <i>Achillea millefolium</i> (Asteraceae)	897
S3.P126	Metabolomics and microbiomics analysis reveals the dynamic changes during <i>Citrus depressa</i> Hayata fermentation	898
S3.P127	LC-MS analysis-based molecular networking for identifying anthelmintic compounds in <i>Terminalia leiocarpa</i> , a traditional plant used in livestock farming in West Africa	899
S3.P128	Integration of Wnt-inhibitory activity and metabolomics structural novelty results to discovery novel bioactive natural products: new bicyclo[3.3.1]non-3-ene-2,9-diones from the leaves of <i>Hymenocardia punctata</i>	900
S3.P129	Enhancing prioritization strategies of natural extracts: integrating heterogeneous data from metabolomics datasets and biological screenings into knowledge graphs	901

S3.P130	The impact of various technological parameters and a place of origin of <i>Nigella sativa</i> . L seeds on the thymoquinone content in black cumin oil	902
S3.P131	Unravelling enzymatic hydrolysis inhibitors in lignocellulosic biomass samples: a combined LC-HRMS/MS and NMR multiblock modelling approach.	903
Total synthesis and biomimetic synthesis of natural products		904
S3.P132	Obtaining new [6]-gingerol derivatives with potential antiparasitary activity	904
S3.P133	Identification of the C2-sulfonamidyl chromone scaffold for developing anti-inflammatory agents targeting hyperactivation of human neutrophils	905
S3.P134	New xanthone analogues with potent intracellular reactive oxygen species inhibition effects	906
S3.P135	A new complementary process toward functionalized dihydropyridinones: application in total synthesis of phenanthroindolizidine alkaloids	907
S3.P136	Divergent synthesis of prenylated xanthenes as novel fungal IRE1 inhibitors	908
S3.P137	Cytotoxic effects of xanthone derivatives towards leukemia cells	909
S3.P138	Bimatoprost potentiates hair growth via Akt/Wnt/ β -catenin signaling by targeting the KATP channels	910
S3.P139	Large scale synthesis, anticonvulsant effect and toxicity assessment of cannabigerol and cannabinol	911
S3.P140	Thymol-based heterocyclic derivatives: synthesis and SARS-CoV-2 Mpro molecular docking studies	912
S3.P141	Expanding chemical space around structures from <i>Strychnos variabilis</i> : late-stage functionalization strategies using photochemistry	913
S3.P142	Effects of metabolic modulators on the accumulation of specialized metabolites in <i>Hypogymnia physodes</i> (L.) Nyl.	914
S3.P143	Synthesis of polymethoxyflavone - zapotin (5,6,2',6'-tetramethoxyflavone) and its new structural analogues	915
S3.P144	Anti-leishmanial activity of synthesized and characterized semisynthetic derivatives of betulin	916
S3.P145	Exploring novel amino acid synthesis for plant-derived active cyclopeptide acquisition	917
S3.P146	Structure modification of plant-derived milisanes compounds to explore their therapeutic potential as anticancer agents	918
Genetic engineering of food crops and medicinal plants		919
S3.P147	Seed propagation, tissue culture and <i>in-vitro</i> conservation studies in <i>Aristolochia tagala</i> Cham: an endangered medicinal plant of western ghats of India	919
S3.P148	Efficient genetic transformation of <i>Nigella damascena</i> callus	920
Natural cosmetics, nutraceuticals, and food supplements		921
S3.P149	Identifying antioxidant compounds within a mixture through dereplication and online DPPH strategies: A study on the CO ₂ byproduct of the Makwaen Pepper	921
S3.P150	Establishing human intestinal slice cultures as an ex vivo model of inflammatory bowel diseases for the pharmacological characterization of herbal extracts	922
S3.P151	Comparative investigation of frankincense nutraceuticals: pharmacokinetic/pharmacodynamic study	923
S3.P152	Bioactive potential of deglycosylated flavonoids extracted from immature oranges in skin health	924
S3.P153	Protective effect on H ₂ O ₂ -induced oxidative stress in L6 myotubes of <i>Mitragyna diversifolia</i> leaf extract	925
S3.P154	<i>Diospyros lotus</i> leaf extract and its main component myricitrin inhibit both histamine- dependent and histamine-independent itching	926
S3.P155	Beneficial effects of <i>Gryllus bimaculatus</i> extract on muscle atrophy in dexamethasone- stimulated C2C12 Cells	927
S3.P156	Anti-inflammatory effect of hyuganin from <i>Peucedanum japonicum</i> Thunberg leaves in lipopolysaccharide-stimulated RAW264.7 cells	928
S3.P157	Triterpene glycosides from the roots of <i>Deutzia x hybrida</i> "Strawberry Fields" (Hydrangeaceae) as sweet taste modulators	929
S3.P158	Simultaneous analysis of bergapten and schinifoline in <i>Zanthoxylum schinifolium</i> Siebold & Zucc. seeds using HPLC and UPLC-MS/MS systems	930
S3.P159	Unveiling the mystery of Australian <i>Apis mellifera</i> propolis	931
S3.P160	Chemical profile and antioxidant capacity of Chilean traditional fermented beverages of algarrobo and chañar	932
S3.P161	Anti-glycation activity of corn silk water extract and establishment of methods for evaluating its quality	933
S3.P162	Bioactivity-Based Molecular Networking for the comparison of <i>Calophyllum inophyllum</i> (tamanu) oil obtained via three ecofriendly extraction processes	934
S3.P163	Regenerative farming of medicinal crops as a source of robust, sustainable and novel supply chains for the cosmetic industry (BIOSURE)	935
S3.P164	Immunomodulatory activities of aqueous extracts of <i>Dendrobium officinale</i> and <i>Dendrobium nobile</i> in peripheral blood mononuclear cells of tongue cancer patients	936
S3.P165	An innovative bioactive ingredient preparation method without column chromatography and gel filtration: High-purity 3S,3'S-astaxanthin from bioresources	937
S3.P166	Bioguided isolation and feature-based identification of bioactive molecules from <i>Fagraea berteriana</i> targeting dermal papilla cells and the Wnt pathway	938
S3.P167	GABA prevents sarcopenia through regulation of protein turnover and inflammaging in 21- 25-month-old C57BL/6J mice	939
S3.P168	The influence of disturbed gravity on the content of bioactive phytochemicals in selected sprouts of edible plants from the Brassicaceae family	940
S3.P169	Equivalency of DNA sequencing, HPTLC chromatographic analysis, botanical microscopy, for botanical identity/ authentication: A statistical evaluation and quantitation of identification uncertainty	941
S3.P170	Bioassay-guided characterization of the Bergamot (<i>Citrus aurantium</i> var. <i>bergamia</i>) leaf- extract	942
S3.P171	From nature to hair: using lignin from <i>Acacia</i> wood to produce hair conditioners	943
S3.P172	Phytochemical exploration and biological evaluation of four <i>Thymus</i> L. species from Türkiye	944
S3.P173	Isoflavone-rich clover sprouts and their effect on thyroid cells in vitro	945
S3.P174	Anti-histaminic and anti-inflammatory activities of <i>Ribes nigrum</i> L. leaf extract CPSP-01	946
S3.P175	In vitro physiological effects and phytochemical exploration of four <i>Sideritis</i> L. species from Türkiye	947
S3.P176	Inhibitory effects of dietary Schisandra supplements on CYP2C19 and CYP2J2 in human liver microsomes	948
S3.P177	Inhibitory effects of dietary <i>Schisandra</i> supplements on cytochrome P450 3A in human liver microsomes	949
S3.P178	Direct extraction and formulation of <i>Bixa orellana</i> L. using a green biphasic solvent	950
S3.P179	Phenylpropanoids and lignans in hemp seeds (<i>Cannabis sativa</i> L. <i>semen</i>): identification and quantification	951
S3.P180	A refined extract of Rosehip modulates skin genes: New evidence for a nutraceutical	952
S3.P181	Complementary LC-MS/MS analyses of alkaloid-rich fractions of the seeds of <i>Plukenetia volubilis</i> L. by LTQ Orbitrap XL and ZenoTOF 7600	953
S3.P182	Compounds of edible fruits vs. bacterial nucleases – <i>in silico</i> study	954

S3.P183	Iridoids from <i>Stenaria nigricans</i> and their insect repellency assessment	955
S3.P184	Natural deep eutectic solvents: a sustainable alternative for the extraction and stabilization of carotenoids and chlorophylls from spinach leaves	956
S3.P185	Baking a difference: elevating bread with <i>Portulaca oleracea</i> L. seeds	957
S3.P186	Chemical and sensory profile of different oregano individuals originated from Vertisco mountain	958
S3.P187	Valuing the medicinal flora of C��a valley (Portugal) through the development of new plant- based cosmetics	959
S3.P188	Identification of anti-inflammatory compounds in <i>Parasenecio adenostyloides</i> using bioactivity-based molecular networking	960
S3.P189	Metabolic response of <i>Panax ginseng</i> to root-rot: a molecular networking approach	961
S3.P190	Collagenase and tyrosinase inhibitory compounds from fish gut bacteria <i>Ruegeria atlantica</i> and <i>Pseudoalteromonas neustonica</i>	962
S3.P191	Phytochemical study of five Polynesian plant extracts for skin care application	963
S3.P192	Eco-extraction and characterization by HPLC-HRMS/MS of <i>in vitro</i> culture of the carnivorous plant <i>Drosera</i>	964
S3.P193	Antioxidant potential and phytochemical analysis of an infusion obtained from the leaves of <i>Talisia esculenta</i>	965
S3.P194	Prospecting the phytochemical and antioxidant effects in vivo and in vitro of the ethanolic and aqueous extracts of the seeds of <i>Geoffroea decorticans</i> burkart (Fabaceae) on the animal model <i>Tenebrio molitor</i>	966
S3.P195	Standardisation of secondary metabolite extraction and characterisation of <i>Citrus aurantium var. bergamia</i> leaves using green extraction technology	967
S3.P196	Sustainable isolation and improved structural characterization of mycosporin-like amino acids: natural UV-filters from red algae	968
S3.P197	Cosmeceutical stability assessment of bergamot: behavioural studies on its potential as an active ingredient	969
S3.P198	<i>Erythrina</i> spp. as natural anti-inflammatory ingredients for cosmeceuticals: an <i>in-silico</i> study	970
S3.P199	Residual solvents in botanical extracts used in food supplements: a regulatory conundrum for scientific justification	971
S3.P200	<i>Grateloupia lanceolata</i> promotes hair growth by inhibiting 5��-reductase activity and opening KATP channels	972
S3.P201	Anti-inflammatory effect of <i>Lysimachia mauritiana</i> extract on particulate matter 10 Plus Diesel Exhaust Particles (PM10D)-induced respiratory disease mouse model and standardization using UPLC-qTof/MS analysis	973
S3.P202	Analysis of dietary supplements containing adaptogenic herb <i>Astragalus trimestris</i>	974
S3.P203	The GreenValue project: exploring salt tolerant plants as sources of innovative food additives	975
S3.P204	Seasonal variation of the indole alkaloid profile of <i>Vinca minor</i> determined by GC-MS	976
S3.P205	Review of honeybee-sourced, ingested natural products and their health uses	977
S3.P206	Evaluating the analytical profile of ten hydrophilic <i>Cucurbita pepo</i> seed extracts	978
S3.P207	Uncovering the potential of cyanobacterial pheophorbide derivatives for metabolic disorders	979
S3.P208	Phyllobilins derived from chlorophyll degradation are overlooked, bioactive phytochemicals in plant-based food	980
S3.P209	Development of a simple liquid chromatographic method for the determination of sitosterol in dietary supplements	981
S3.P210	Effects of Crude leaf extract of <i>Ficus</i> sp. on the proliferation and differentiation of C2C12 muscle cell line	982
S3.P211	<i>In vitro</i> analysis of antioxidant and antidiabetic activity of four plant extracts from the Algarve region	983
S3.P212	Innovative approach towards the qualitative analysis of herbal products using UHPLC-PDA- HRMS/MS: case study on species discrimination within the Lamiaceae family	984
S3.P213	Phenolic composition, antioxidant, and enzyme inhibitory activities of <i>Saxifraga paniculata</i> Mill.	985
S3.P214	What does the percentage of TBA% say about <i>Boswellia serrata</i> extract?	986
S3.P215	Pharmacokinetic study of cycloastragenol administered in a liposomal formulation to rats	987
S3.P216	An extract of <i>Boehmeria nivea</i> induces apoptosis by disrupting oncogenic signaling in human pancreatic cancer cells	988
S3.P217	Chemical screening and biological activity of various species of wood-decay fungi of the genus <i>Ganoderma</i>	989
S3.P218	Identification of a marine cyanobacterium <i>Rivularia</i> sp. LEGE 07159 with promising bioactivity on appetite regulation	990
S3.P219	Potent xanthine oxidase inhibitory activity of acacetin from <i>Agastache rugosa</i> (Fisch. & C.A.Mey.) Kuntze <i>in vitro</i> and molecular docking studies	991
S3.P220	Analysis and biological evaluation of Chios Mastic Gum towards its anti-ageing properties	992
S3.P221	Metabolomics of <i>Withania somnifera</i> L. extracts by a combined LC-MS and NMR approach	993
S3.P222	A food supplement derived from high-phenolic Kalamon table olives reduced cholesterol in patients with mild dyslipidemia	994
S3.P223	The rising phenomenon of cannabis ‘tea’ drinking: a safe and healthy alternative?	995
S3.P224	HPLC-MS analysis and AChE activity of aerial parts and root extracts of <i>Taraxacum mirabile</i> Wagenitz, an endemic species of Turkey	996
S3.P225	Dietary soft electrophiles upregulate lipid pro-resolving molecules in a <i>Drosophila</i> model of Parkinson’s disease	997
S3.P226	Comprehensive analysis of commercial kratom products: alkaloids, polyphenols, heavy metals and microbial contaminations	998
S3.P227	Diarylheptanoids from <i>Corylus avellana</i> cultivar ‘Tonda di Giffoni’: antioxidant effects through the activation of the Nrf2 pathway as assessed via molecular docking and biological evaluation	999
S3.P228	Valorization of Almond Okara as a valuable source of polyphenols and proteins	1000
S3.P229	Application of annular centrifugal extraction (ACE) in honey - isolation and identification of honey bioactive compounds	1001
S3.P230	Ornamental varieties of <i>Achillea millefolium</i> L. as a source of skin-lightening ingredients for natural cosmetics	1002
S3.P231	Impact of the drying process and incorporation of natural preservatives on the viability of lactic acid bacteria in yogurts	1003
S3.P232	NMR metabolomics and chemometrics of <i>Malus domestica</i> (Suckow) Borkh. varieties from Molise region of Italy	1004
S3.P233	Exploring <i>Genista tridentata</i> leaves extract: novel insights for functional food in diabetes management	1005
S3.P234	A formulation containing palmitoylethanolamide and phenolic compounds counteracts diabetes and related hepatic damage in mice: in vivo and in vitro evidence	1006
S3.P235	The <i>Vitis vinifera</i> cvs. callus <i>in vitro</i> culture extracts as a potential cosmetic antioxidant	1007
S3.P236	Phytochemical analysis of <i>Echinacea</i> preparations from the pharmacies and herbal markets in T��rkiye	1008
S3.P237	Untargeted metabolomics analyses to identify a new sweet compound released during post- fermentation maceration of wine	1009
S3.P238	Do you know what’s really hiding in popular herbal supplements? – evaluation of products containing <i>Cistus</i> leaves and extracts	1010
S3.P239	A smartphone-based methodology for analyte quantification and antioxidant activity determination directly in emulsions	1011
S3.P240	Phytochemical analysis of <i>Laurus nobilis</i> L. leaves essential oil: rationalization of antioxidant and anti-inflammatory activities by combined <i>in vitro</i> and <i>in silico</i> approach	1012
S3.P241	A study on the biological activities of loquat leaf extract using various extraction methods	1013
S3.P242	Study on modulation of inflammation expression by <i>Brassica oleracea</i> L. leaves extract	1014
S3.P243	Glucosinolates in <i>Brassica incana</i> subsp. <i>raimondoi</i> (Brassicaceae) leaf extract: composition, content, and anti-inflammatory activity	1015
S3.P244	<i>Ex vivo</i> and clinical evaluation of efficacy and mode of action of an <i>Anogeissus leiocarpus</i> and retinol-containing eye cream	1016
S3.P245	Targeted Isolation of natural products that modify sweet taste perception through metabolomics, molecular networking and sensory evaluation of <i>Symplocos</i> spp. (Symplocaceae)	1017
S3.P247	Investigation of the Greek microbial diversity for the discovery of novel antiaging products	1018

S3.P248	EthnoHERBS: Conservation of european biodiversity through exploitation of traditional herbal knowledge for the development of innovative products	1019
S3.P249	<i>Artocarpus thailandicus</i> heartwood extract extends lifespan and promotes healthy aging behaviour in <i>Caenorhabditis elegans</i>	1020
S3.P250	Utilization of ethnobotanical data to discover potent inhibitors of skin related enzymes through the application of in silico based methodology	1021
S3.P251	Chemical characterization of a <i>Schisandra sphenanthera</i> fruit extract, an adaptogen from a sustainable panda-friendly conservation program	1022
S3.P252	LC-MS/MS analysis of polyphenolic and triterpene components and antioxidant activity of cold-pressed rose oils	1023
S3.P253	Chemical insights into a bioactive polyphenolic extract from <i>Camellia japonica</i> L. flower	1024
S3.P254	Complementary LC-MS/MS analyses of alkaloid-rich fractions of the seeds of <i>Plukenetia volubilis</i> L. by LTQ Orbitrap XL and ZenoTOF 7600	1025
S3.P255	Effect of different extraction methods on determination of <i>Drosera capensis</i> active ingredients	1026
S3.P256	Valorization of <i>Olea europaea</i> L. cv. Caiazzana pruning waste: a step towards a sustainable approach for resource functional recovery	1027
S3.P257	Plants from Lublin region as potential sources of ingredients with cosmetic value	1028
S3.P258	Anti-neuroinflammatory effects of <i>Elsholtzia blanda</i> benth on lipopolysaccharide-stimulated BV-2 microglial cells	1029
S3.P259	Metabolite profiles and biological activities of <i>Capsicum chinense</i> cell cultures	1030
S3.P260	Liposome encapsulation of <i>Lavandula austroapennina</i> N.G. Passal., Tundis & Upson leaf and stem extracts for cosmeceutical innovation	1031
S3.P261	Effect of combining proteolytic enzymes papain and chymotrypsin with Alexandrite laser hair removal on the degree of facial hirsutism	1032
S3.P262	Simultaneous analysis of bergapten and schinifoline in <i>Zanthoxylum schinifolium</i> Siebold & Zucc. seeds using HPLC and UPLC-MS/MS systems	1033
S3.P263	Health-promoting effects of the hydroalcoholic extract from the <i>Allium sativum</i> L. ecotype “Sulmona red garlic” in the colon	1034
S3.P264	Anti-inflammatory effect of <i>Lysimachia mauritiana</i> extract on particulate matter 10 plus diesel exhaust particles (PM10D)-induced respiratory disease mouse model and standardization using UPLC-qTof/MS analysis	1035
S3.P265	Optimization, estimation, and improvement of lutein content from marigold flowers using <i>Lactobacillus</i> species by solid-state fermentation	1036
S3.P266	Metabolite Profiling and Molecular Network Shows Kinkeloids as promoting of collagen synthesis from <i>Combretum micranthum</i>	1037
S3.P267	Determination of the anti-inflammatory activity of fruits of some varieties of <i>Rubus idaeus</i> and <i>Rubus occidentalis</i> (Rosaceae)	1038
S3.P268	Unveiling Thailand's ethnopharmacological heritage to find new natural sweeteners	1039
S3.P269	Recovery effect of phloroglucinol 1-O-β-D-glucopyranoside from the brown alga <i>Agarum clathratum</i> subsp. <i>yakishiriense</i> on neomycin-induced otic hair cell damage in zebrafish	1040
S3.P270	Giuseppe Raddi and the plants: Taxonomy, Uses and Conservation in the 21st century	1041

Welcome Letter

Dear Participants, dear Colleagues,

on behalf of the Organizing Committee and the Scientific Committee it is our great pleasure to welcome you at the International Congress on Natural Products Research 2024 (ICNPR2024) which will be held in Kraków, Poland, July 13-17, 2024.

ICNPR2024 is co-organized by the Phytochemical Society of Europe (PSE), and our six sister societies representing the US, Europe and Asia, which include the Society for Medicinal Plant and Natural Products Research (GA), the Association Francophone pour l'Enseignement et la Recherche en Pharmacognosie (AFERP), the Italian Society of Phytochemistry and Sciences of Medicinal, Food and Perfume Plants (SIF), the American Society of Pharmacognosy (ASP), the Japanese Society of Pharmacognosy (JSP) and the Korean Society of Pharmacognosy (KSP). Every four years all of them meet together to share common interest in natural products research and exchange experience and simply unite together. It is always a unique possibility to experience this. ICNPR2024 is one of the largest gatherings of scientists involved in natural products and related fields from nearly every continent. The congress brings together experts and researchers from all over the world to share their latest findings and discuss the latest developments in the field of natural products research.

More than 1000 participants from 68 countries will deliver all together around 130 oral and almost 900 poster presentations. As a participant, you will have the opportunity to learn about cutting-edge research, network with other professionals in your field and getting involved in the extensive scientific program in many ways. Aside from that, you should also take the chance to explore the beautiful city of Kraków. It is one of the best recognizable cities in this part of Europe visited by about 10 million visitors annually. They are attracted by one-of-a-kind atmosphere and largest historical complex in Poland, unique on global scale. It is a city with a rich history and culture. The old town, with its beautiful architecture and narrow streets, is a UNESCO World Heritage site.

This congress is only possible with the collaboration of many people who were dedicated to do their best to make this event happen. First of all, we would like to thank the representatives of the seven organizing societies for their great support. Without the work of the members of the Scientific Committee it would not have been possible to handle more than 1000 abstracts and to create a scientific program of high quality, your efforts are very much appreciated. We also would like to thank plenary and keynote speakers for accepting our invitation to Krakow. We are grateful to the congress organizing secretariat, Symposium Cracoviense Sp. z o.o., for excellent cooperation in preparation of this event. Finally, we sincerely would like to acknowledge all financial support provided by the sponsors of this conference.

We hope that you will enjoy the time at ICNPR2024 and you feel not only the spirit of the city and place but also the touch of Polish culture and hospitality.



Krystyna Skalicka-Woźniak

Prof. Dr. Krystyna Skalicka-Woźniak

Chairman of the Scientific Committee



Fran Bucar

Prof. Dr. Franz Bucar

Chairman of the Scientific Committee

Committee

Chairs of the Scientific Committee

Prof. Krystyna Skalicka-Woźniak

Medical University of Lublin (Poland)

Prof. Franz Bucar

University of Graz (Austria)

Members of the Scientific Committee

Prof. Sayeed Ahmad

Centre of Excellence in Unani Medicine (India)

Prof. Shunji Aoki

Department of Pharmacy, School of Pharmacy,
Hyogo Medical University (Japan)

Ass. Prof. Dr. Maria Pia Argentieri

Department of Pharmacy, University of Bari (Italy)

Prof. Randolph Arroo

De Montfort University (United Kingdom)

Prof. Søren Bak

University of Copenhagen (Denmark)

Ass. Prof. Dr. Stefania Benvenuti

Department of Life Sciences, University
of Modena e Reggio (Italy)

Prof. Clara Bik-San Lau

The University of Hong Kong (Hong Kong SAR, China)

Prof. Dr. Anna Rita Bilia

Department of Chemistry „Ugo Schiff”,
University of Firenze (Italy)

Prof. Dr. Angela Bisio

Department of Pharmacy, University of Genova (Italy)

Dr. Sabrina Bouteftouchet

Université Paris-Cité (France)

Prof. Dr. Alessandra Braca

University of Pisa (Italy)

Prof. Ian Castro-Gamboa

State University of São Paulo (IQ-UNESP) (Brasil)

Prof. Lucie Cahliková

Charles University (Czech Republic)

Prof. Paula Castillo

University of Madeira (Portugal)

Prof. Pierre Champy

Université Paris-Saclay (France)

Dr. François Chassagne

IRD Pharmadev (France)

Dr. Antigoni Cheilari

National and Kapodistrian University of Athens (Greece)

Dr. Sanem Hosbas Coskun

National Institutes of Health (Bethesda, MD, USA)

Ass. Prof. Esperanza Carache De Blanco

Ohio State University (Columbus, OH, USA)

Prof. Dr. Nunziatina De Tommasi

Department of Pharmacy, University of Salerno (Italy)

Dr. Steven Dentali

Dentali Botanical Sciences (Redondo Beach, CA, USA)

Prof. Thomas Efferth

Johannes Gutenberg University (Germany)

Prof. Halina Ekiert

Jagiellonian University, Kraków (Poland)

Prof. Francesco Epifano

University „Gabriele D'Annunzio” of Chieti, Pescara (Italy)

Prof. Maria José Umbelino Ferreira

University of Lisbon (Portugal)

Ass. Prof. Nikola Fokialakis

University of Athens (Greece)

Prof. Dr. Robert Fürst

LMU Munich (Germany)

Dr. Stefan Gafner

American Botanical Council (United States)

Ass. Prof. Katarzyna Gaweł-Bęben

University of Information Technology and
Management in Rzeszów (Poland)

Prof. Jürg Gertsch

University of Bern (Switzerland)

Prof. Simon Gibbons	University of Nizwa (Sultanate of Oman)
Prof. Sebastian Granica	Warsaw Medical University (Poland)
Dr. Tanja Grkovic	National Cancer Institute, National Institutes of Health (Frederick, MD, USA)
Prof. Olivier Grovel	Nantes Université (France)
Prof. De-an Guo	Shanghai Institute of Materia Medica, Chinese Academy of Sciences (China)
Ass. Prof. Dr. Maria Halabalaki	National and Kapodistrian University of Athens (Greece)
Dr. Christophe Hano	University of Orleans (France)
Prof. Dr. Michael Heinrich	University College London (United Kingdom)
Dr. Mah Siau Hui	Taylor's University (Malaysia)
Prof. Takuya Ito	Department of Pharmacy, Osaka Ohtani University (Japan)
Prof. Sanjay Jachak	National Institute of Pharmaceutical Education and Research (India)
Ass. Prof. Kyo Bin Kang	Sookmyung Women's University (South Korea)
Prof. Dr. Oliver Kayser	Technische Universität Dortmund (Germany)
Dr. Olaf Kelber	Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt (Germany)
Dr. Isabelle Kerzaon	Université de Lyon (France)
Dr. Stefanos Kikionis	National and Kapodistrian University of Athens (Greece)
Dr. Chungsub Kim	Sungkyunkwan University (South Korea)
Prof. Ki Hyun Kim	Sungkyunkwan University (South Korea)
Prof. Dr. Luiz Carlos Klein Júnior	University of Vale do Itajaí (Brazil)
Prof. Ilina Krasteva	Medical University of Sofia (Bulgaria)
Prof. Wirginia Kukuła-Koch	Medical University of Lublin (Poland)
Dr. Marina Kritsanida	Université Paris-Cité (France)
Prof. Norberto Peoporine Lopes	Universidade de São Paulo (Brasil)
Dr. Simon Vlad Luca	Technical University of Munich (Germany)
Ass. Prof. Agnieszka Ludwiczuk	Medical University of Lublin (Poland)
Prof. Francisco A. Macias	University of Cadiz (Spain)
Prof. Toshiaki Makino	Graduate School of Pharmaceutical Sciences, Nagoya City University
Ass. Prof. Maria Carla Marcotullio	Department of Pharmaceutical Science, University of Perugia (Italy)
Prof. Marta Mendel	Warsaw University of Life Science (Poland)
Ass. Prof. Dr. Luigi Menghini	Department of Pharmacy, University G. d'Annunzio of Chieti, Pescara (Italy)
Ass. Prof. Tomasz Mroczek	Medical University of Lublin (Poland)
Ass. Prof. Dr. Luigi Milella	Department of Science, University of Basilicata (Italy)
Prof. Hiroyuki Morita	Institute of Natural Medicine, University of Toyama (Japan)
Prof. Brian Murphy	University of Illinois at Chicago (Chicago, IL, USA)
Prof. Dr. Lutfun Nahar	Liverpool John Moores University (United Kingdom)
Prof. Ilkay Erdogan Orhan	Gazi University (Turkey)
Prof. Giorgio Pintore	Department of Medicine, Surgery and Pharmacy, University of Sassari (Italy)

Prof. Jakub Piwowarski
Prof. Monica Tallarico Pupo
Ass. Prof. Dr. Federica Pollastro

Prof. Young Pyo Jang
Dr. Emerson Ferreira Queiroz
Prof. Ronald J Quinn
Prof. Pascal Richomme-Peniguel
Dr. Kenneth Ritchie
Prof. Céline Rivi re
Prof. Dr. Judith Maria Rollinger
Dr. Caroline Rouger
Prof. Catherine Roullier
Dr. Fanny Roussi
Dr. Gaia Scalabrino
Ass. Prof. Elwira Sieniawska
Prof. Caroline St vigny
Prof. Nazım Şekero lu
Ass. Prof. Kevin Tidgewell
Prof. Micha  Tomczyk
Dr. Adriana Trifan
Prof. Alvaro Viljoen
Dr. Cica Vissiennon
Dr. Michael Walkenhorst
Ass. Prof. Jaclyn M. Winter
Dr. Evelyn Wolfram
Prof. Jean-Luc Wolfender
Ass. Prof. Heejung Yang
Prof. Jianbo Xiao

Warsaw Medical University (Poland)
University of Sao Paulo (FCFRP-USP) (Brasil)
Department of Pharmaceutical Science, University of
Piemonte Orientale „Amedeo Avogadro” (Italy)
Kyunghee University (South Korea)
Universit  de Gen ve CMU (Switzerland)
Griffith University (Australia)
Universit  d'Angers (France)
Liverpool John Moores University (United Kingdom)
University of Lille (France)
University of Vienna (Austria)
Universit  de Bordeaux (France)
Nantes Universit  (France)
ICSN-CNRS (France)
The University of Dublin (Ireland)
Medical University of Lublin (Poland)
Universit  Libre de Bruxelles (Belgium)
Gaziantep University (Turkey)
University of Kentucky (Lexington, KY, USA)
Medical University of Białystok (Poland)
University of Medicine and Pharmacy (Romania)
Tshwane University of Technology, Pretoria (South Africa)
Leipzig University (Germany)
Research Institute of Organic Agriculture (Switzerland)
University of Utah (Salt Lake City, UT, USA)
Z rich University of Applied Sciences (Switzerland)
Universit  de Gen ve (Switzerland)
Kangwon National University (South Korea)
University of Vigo (Spain)

Chair of the Organizing Committee

Prof. Krystyna Skalicka-Wo niak Medical University of Lublin (Poland)

Members of the Organizing Committee

Prof. Franz Bucar University of Graz (Austria)
Prof. Sebastian Granica Warsaw Medical University (Poland)
Ass. Prof. Agnieszka Ludwiczuk Medical University of Lublin (Poland)
Prof. Jakub Piwowarski Warsaw Medical University (Poland)
Prof. Agnieszka Szopa Jagiellonian University, Krak w (Poland)
Prof. Micha  Tomczyk Medical University of Białystok (Poland)

Workshop ECRW

The natural diversity of acyltransferases reveals versatility and specificity in the synthesis of gene-encoded lipopeptides

Johannes A. Eckert^{a,*}, Nina M. Bösch^a, Florian Hubrich^a, Serina L. Robinson^b, Jörn Piel^a, Anna L. Vagstad^a

^a Eidgenössische Technische Hochschule (ETH) Zürich, Institute of Microbiology, Vladimir-Prelog-Weg 4, 8093 Zürich, Switzerland

^b Swiss Federal Institute of Aquatic Science and Technology (Eawag), Department of Environmental Microbiology, Überlandstrasse 133, 8600 Dübendorf, Switzerland

Lipopeptides comprise biochemical compounds that can be naturally produced or chemically synthesized. They exhibit diverse bioactive functions and are widely used in health care (Hamley, 2015). Despite their versatility, the production and diversification of lipopeptide compounds continue to be a challenge for state-of-the-art biological and chemical synthesis. In contrast, we offer a more streamlined and adaptable route to lipopeptide natural products via ribosomally synthesized and post-translationally modified peptides (RiPPs). Their primary structure is determined by a genetically encoded precursor protein, which is modified by maturases within the same biosynthetic gene cluster (Montalbán-López et al., 2021). The crucial lipidation reaction is catalyzed by a novel class of maturases of the GCN5-related *N*-acetyltransferase (GNAT) family, which introduce medium-chain fatty acids onto the sidechain amino group of lysines or arginine-derived ornithines (Hubrich et al., 2022) (Fig. 1).




Fig. 1. Transfer of fatty acids to the sidechain of ornithines and lysines by RiPP GNATs.

Here we present the natural diversity of this GNAT family and their corresponding gene-encoded lipopeptides by characterizing the substrate scope of selected GNATs with their cognate precursor peptides. We could show that a variety of fatty acids can be introduced including chain lengths from C10 to C18. We are currently exploring this enzyme class to enable custom peptide engineering efforts towards generating bioactive lipopeptide compounds. We envision this platform to be used to facilitate the production of crucial lipopeptide compounds for human health and to quickly screen for antimicrobial lipopeptide compounds and non-ribosomal lipopeptide mimics.

Keywords: GNAT, RiPPs, lipopeptides, synthetic biology

References

- Hamley, I. W., 2015. Lipopeptides: from self-assembly to bioactivity. *Chem. Comm.* 51 (41), 8574-8583.
- Montalbán-López, M., et al., 2021. New developments in RiPP discovery, enzymology and engineering. *Nat. Prod. Rep.* 38, 130-239.
- Hubrich, F. & Bösch, N. M., et al. 2022. Ribosomally derived lipopeptides containing distinct fatty acyl moieties. *PNAS* 119 (3), e2113120119.

* Presenting author. E-mail address: johannes.eckert@biol.ethz.ch (Johannes Eckert)

Anti-inflammatory biscembranoids from aquaculture-derived soft coral *Sarcophyton trocheliophorum*

Lo-Yun Chen^{a,*}, Bo-Rong Peng^b, Ngoc Bao An Nguyen^b, Tsong-Long Hwang^c, Kuei-Hung Lai^{a,b}, Jui-Hsin Su^{d,e}

^a PhD Program in Clinical Drug Development of Herbal Medicine, College of Pharmacy, Taipei Medical University, 11031, Taipei, Taiwan

^b Graduate Institute of Pharmacognosy, College of Pharmacy, Taipei Medical University, 11031, Taipei, Taiwan

^c Research Center for Chinese Herbal Medicine, College of Human Ecology, Chang Gung University of Science and Technology, Taoyuan, 33303, Taiwan

^d National Museum of Marine Biology & Aquarium, Pingtung, 94450, Taiwan

^e Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung, 80424, Taiwan

As the most abundant white blood cells in the human body, neutrophils are a key component of the immune system, playing a crucial role in defending against infections. However, excessive activation or improper regulation may lead to severe inflammatory diseases. Therefore, the development of drugs capable of inhibiting the activation of neutrophils holds the potential to effectively treat various inflammatory conditions. In our quest to discover bioactive biscembranoids sourced from cultured soft coral *Sarcophyton trocheliophorum*, we utilized molecular networking (MN) analysis. Our focus was on clusters within the *m/z* range of 650 to 750, as well as the clusters exhibiting a substantial inhibition rate (>70%) of superoxide anion generation. These clusters were prioritized for purification, leading to the isolation of ten biscembranoids, including four new compounds (sarcotrochelides A–D, **1**, **2**, **5**, and **6**) and six known analogues. Noteworthy, **5** and **6** exhibited an unprecedented 3-hydroperoxy-3-methylcyclohex-1-ene moiety. The structures of all isolated metabolites were elucidated through a series of spectral analyses, and their anti-inflammatory activities were assessed. Compound **8** demonstrated the highest inhibition against superoxide anion generation and elastase release in activated neutrophils, with IC₅₀ values of 1.98 and 2.76 μM, respectively. The observed variations in bioactivity underscored the significance of lactone rings and double bond geometry within the biscembranoid skeleton. These results confirmed the potential of biscembranoids derived from cultured *S. trocheliophorum* as promising candidates for future development of anti-inflammatory agents tailored to target neutrophils.

Keywords: neutrophil, *Sarcophyton trocheliophorum*, biscembranoid, molecular networking, anti-inflammation

References

- Hegazi, N.M., Mohamed, T. A., Saad, H.H., Al-Hammady, M.A., Hussien, T.A., Hegazy, M.E.F., Gross, H., 2022. Molecular network guided cataloging of the secondary metabolome of selected Egyptian red sea soft corals. *Marine Drugs*, 20(10).
- Nguyen, N.B.A., Chen, L.Y., Chen, P.J., El-Shazly, M., Hwang, T.L., Su, J.H., Lai, K.H., 2022. MS/MS Molecular networking unveils the chemical diversity of biscembranoid derivatives, neutrophilic inflammatory mediators from the cultured soft coral *Sarcophyton trocheliophorum*. *International Journal of Molecular Sciences*, 23(24).

* Presenting author. E-mail address: d339112001@tmu.edu.tw (Lo-Yun Chen).

Integration of *Caenorhabditis elegans* bioactivity assays for a sustainable identification of health promoting natural products

Martina Redl^{a,b,*}, Judith M. Rollinger^a

^a Department of Pharmaceutical Sciences, Division of Pharmacognosy, University of Vienna, Josef-Holaubek Platz 2, 1090 Vienna, Austria

^b Vienna Doctoral School of Pharmaceutical, Nutritional, and Sport Sciences, University of Vienna, Josef-Holaubek Platz 2, 1090 Vienna, Austria

The increasing life expectancy of the world's population has a profound impact on our society and healthcare system, resulting in an urgent need for innovative regimens to improve healthy aging. Additionally, great emphasis ought to be placed on sustainability of environmental and time constraints in early drug discovery studies. This project aims for the sustainable discovery of natural products (NPs) to decelerate the aging process and attenuate age-related diseases. The presented screening strategy is based on the 3Rs “robustness, reliability and resource conservation”, which are inspired by both the 3Rs in animal studies and environmental protection. The screening approach uses the small model organism *Caenorhabditis elegans* as a simple phenotypic tool in different readouts (Kirchweger et al, 2023) to select the most promising starting materials for in depth phytochemical and pharmacological investigations including LC-MS/MS based dereplication, *in silico* prediction tools and target-oriented assays. Starting from *C. elegans* based screenings of >70 extracts from herbal and fungal origin, we applied a semi-automatic IR-based wormtracker and optimized methodological parameters in various *C. elegans* assays focusing on lifespan, health span, metabolic and neurodegenerative disorders. To best comply with the 3Rs, a workflow has been established, enabling to start with time- and cost saving high content assays to continuously scale down the number of bioactive NPs to be probed on more specific, but elaborate assays with genetically modified nematode strains (Redl et al, 2024). This procedure allows a stepwise increase of information regarding involved pathways, molecular mechanism, and involved chemistries.

Funding: The project is funded by the “Schwabe Phyto Innovation Challenge” by Dr. Willmar Schwabe GmbH & Co.KG.

Keywords: healthy ageing, *Caenorhabditis elegans*, natural product screening, sustainability

References

- Kirchweger, B., Zwirchmayr, J., Grienke, U., Rollinger, J.M., 2023 The role of *Caenorhabditis elegans* in the discovery of natural products for healthy aging. Nat Prod Rep. 40, 1849-1873
- Redl, M., Shayegan, A., Rollinger, J.M., 2024 Application of 3Rs in *Caenorhabditis elegans* research for the identification of health promoting natural products. Planta Med. 90, ahead of print.

* Presenting author. E-mail address: martina.redl@univie.ac.at (Martina Redl).

Novel natural product inhibitors targeting oncogenic MAPK/ERK and PI3K/AKT signaling in melanoma: from large library screening to target identification

Maria Karpouchtsi^{a,*}, Lara Dürr^a, Maciej Dobrzynski^b, Silke Radetzki^c, Tanja Hell^a, Matthias Hamburger^a, von Jens-Peter von Kries^c, Olivier Pertz^b, Robin Teufel^a, Eliane Garo^a

^a Division of Pharmaceutical Biology, Department of Pharmaceutical Sciences, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

^b Institute of Cell Biology, University of Bern, Baltzerstrasse 4, 3012 Bern, Switzerland

^c Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Robert-Rössle-Straße 10, 13125 Berlin, Germany

Malignant melanoma is the deadliest type of skin cancer with unmatched mutation rates arising in both MAPK/ERK and PI3K/AKT signaling pathways. Although specific inhibitors of these critical pathways show spectacular initial results, most patients relapse within just a few months. Combination therapy can improve overall survival rates, but the currently available options are limited (Tanda et al., 2020). Therefore, novel inhibitors targeting oncogenic ERK and AKT signaling in melanoma are urgently needed. To tackle this issue, our in-house library of crude plant extracts was combined with an innovative high-content screen (HCS) that quantifies downstream inhibitory activity at ERK and AKT level. HPLC-based activity profiling of the active hits and subsequent targeted isolation of the bioactive constituents was performed (Dürr et al., 2022). To further explore the coverage of chemical space for such inhibitors, we also screened large pure compound libraries by accessing high-throughput screening pipelines through an EU-OPENSSCREEN program. To this end, we screened 2,576 plant extracts and additional 25,696 pure natural and synthetic compounds. A total of 46 active compounds were confirmed as downstream inhibitors of ERK and/or AKT with IC₅₀ values in the low micromolar range. The current challenge aims towards target identification of the most promising hits. Several key kinases of the ERK-AKT network were produced through different cloning techniques and heterologous expression systems. Our strategy further includes the assessment of physical binding as well as enzymatic inhibition. Ultimately, we envisage to develop these newly discovered inhibitors into lead compounds for future drug development.

Keywords: melanoma, high-content screening, EU-OPENSSCREEN, MAPK/ERK and PI3K/AKT signaling, target identification

References

- Tanda, E.T., Vanni, I., Boutros, A., Andreotti, V., Bruno, W., Ghiorzo, P., Spagnolo, F., 2020. Current State of Target Treatment in BRAF Mutated Melanoma. *Front. Mol. Biosci.* 7, 154.
- Dürr, L., Hell, T., Dobrzynski, M., Mattei, A., John, A., Augsburger, N., Bradanini, G., Reinhardt, J.K., Rossberg, F., Drobnjakovic, M., Gupta, M.P., Hamburger, M., Pertz, O., Garo, E., 2022. High-Content Screening Pipeline for Natural Products Targeting Oncogenic Signaling in Melanoma. *J. Nat. Prod.* 85, 4, 1006–1017.

* Presenting author. E-mail address: maria.karpouchtsi@unibas.ch (Maria Karpouchtsi).

Towards the engineering of the plant endoplasmic reticulum for sustainable production of specialized metabolites

Elke Clicque^{a,b,*}, Alain Goossens^{a,b}

^a Department of Plant Biotechnology and Bioinformatics, Ghent University, Technologiepark 71, B-9052 Ghent, Belgium

^b VIB Center for Plant Systems Biology, Technologiepark71, B-9052 Ghent, Belgium

Plants are among the most creative survivors on the planet. They have optimized their own defence mechanisms against attackers, typically by producing a species-specific arsenal of specialized metabolites. These metabolites not only play vital roles within the plants themselves but also offer a multitude of applications as pharmaceuticals, fragrances etc. However, industrial production of these valuable compounds faces many bottlenecks and often requires intensive adaptations. To facilitate sustainable large-scale production of plant metabolites, we aim to develop a universal tool to boost plant-based production of specialized metabolites. Our strategy is based on recent findings in yeast, in which engineering of the endoplasmic reticulum (ER), which harbours a main part of the enzymatic machinery for many metabolite biosynthesis pathways, results in an enlarged ER size and an increased metabolite accumulation (Arendt et al., 2017). By interfering with the phospholipid biosynthesis pathway, a higher production of building blocks for the ER-membrane and ultimately an enlarged ER were achieved. Our project aims to translate the findings from yeast to plants. Thus far, we have obtained an increased accumulation of phospholipid building blocks in tomato (*Solanum lycopersicum*) roots, by knocking out the *PHOSPHATIDIC ACID PHOSPHOHYDROLASE* (PAH) genes. Currently, we are investigating whether this correlates with an increased production of bioactive specialized metabolites. Additionally, we are also exploring the potential of ER engineering by overexpressing other key phospholipid biosynthetic enzymes. With our obtained *in planta* results, we believe that ER engineering holds great promise for a more efficient and sustainable production of plant specialized metabolites.

Keywords: *Solanum lycopersicum*, endoplasmic reticulum, phospholipids, engineering, roots

References

Arendt, P., Miettinen, K., Pollier, J., De Rycke, R., Callewaert, N., Goossens, A., 2017. An endoplasmic reticulum-engineered yeast platform for overproduction of triterpenoids. *Metab. Eng.* 40, 165-175.

* Presenting author. E-mail address: elcli@psb.vib-ugent.be (Elke Clicque).

PKC- α activation with new semi-synthetic 7 α -acetoxy-6 β -hydroxyroyleanone derivatives for breast cancer therapeutics

Vera M. S. Isca^{a,b,*}, Ana C. Matos^c, Joana Almeida^c, Lucília Saraiva^c, Carlos A.M Afonso^b, Patrícia Rijo^{a,b}

^a CBIOS – Universidade Lusófona's Research Center for Biosciences & Health Technologies, Lisbon, Portugal

^b Research Institute for Medicines (iMED.Ulisboa), Faculdade de Farmácia, Universidade de Lisboa, Portugal

^c LAQV/REQUIMTE, Laboratório de Microbiologia, Departamento de Ciências Biológicas, Faculdade de Farmácia, Universidade do Porto, Portugal

Breast cancer is the most prevalent cancer worldwide (WHO). PKC- α is an attractive anticancer target often associated with breast cancer (Tam et al., 2013). Species of the *Plectranthus* genus (Lamiaceae) are traditionally used in the treatment of several ailments, including cancer. The diterpenoid 7 α -acetoxy-6 β -hydroxyroyleanone (**1**) is the major compound of *P. grandidentatus* Gürke. and demonstrated cytotoxic effects in several cancer cell lines (Ladeiras et al., 2019). Accordingly, the main goal of this study was to synthesize new derivatives of **1** and evaluate them as new antitumoral agents focus on breast cancer therapy. In the context of the study, it was possible to extract and isolated 1.0 g the lead compound **1** and subsequently, synthesize thirty new derivatives (**2** to **31**) through hydrogenation, photochemical and esterification reactions. The cytotoxic effects of all compounds (**1** to **31**) were evaluated on breast cancer cell lines. Notably, the analogues **6**, **7**, **18** and **21** revealed promising outcomes, displaying selectivity towards cancer cells and low IC₅₀, mainly against the aggressive triple negative breast cancer cell lines. Specifically, analogues **6**, **7**, **18** and **21** displayed IC₅₀ values of 1.36, 1.92, 2.11, 6.90 μ M in MDBA-MB-468 cells and 1.70, 3.63, 3.52, 9.44 μ M in MDBA-MB-231 cells, respectively. Therefore, these derivatives were studied as PKC- α activators. Derivative **7** emerged as the most promising PKC- α activator, outperforming the control (PMA) and confirming its potential as antitumoral agent with application on breast cancer. These findings denote a significant progress in our attempts to develop new antitumor agents from natural sources.

Funding: This research was funded by Foundation for Science and Technology (FCT, Portugal) for financial support through the projects with reference DOI 10.54499/UIBP/04567/2020 and DOI 10.54499/UIDB/04567/2020 and the PhD grant (SFRH/BD/137671/2018).

Keywords: *Plectranthus*, royleanone, semi-synthetic derivatives, antitumoral activity, PKC- α

References

- World Health Organization (WHO), fact sheet on Cancer. <https://www.who.int/news-room/fact-sheets/detail/cancer>, accessed October 3, 2023.
- Tam, et al., 2013. Cancer Cell, 24, 347-3649.
- Ladeiras, D., et al., 2016. Current Pharmaceutical Design, 22, 1682–1714.

* Presenting author. E-mail address: vera.isca@ulusofona.pt (Vera M. S. Isca).

Development of a high throughput cell-based assay to highlight potential bioactive antiviral natural products against the respiratory syncytial virus (RSV)

Paola Haemmerli^{a,b,*}, Luis-Manuel Quiros-Guerrero^{a,b}, Laurence Marcourt^{a,b}, Bruno David^c, Antonio Grondin^c, Clément Fage^d, Sophie-Clément^d, Emerson Ferreira Queiroz^{a,b}, Caroline Tapparel^d, Jean-Luc Wolfender^{a,b}

^a Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, CMU, 1211 Geneva, Switzerland

^b School of Pharmaceutical Sciences, University of Geneva, CMU, 1211 Geneva, Switzerland

^c Green Mission Department, Herbal Products Laboratory, Pierre Fabre Research Institute, Toulouse, France

^d Department of Microbiology and Molecular Medicine University of Geneva Geneva 1206, Switzerland

Respiratory syncytial virus (RSV) is a leading cause of bronchiolitis in children resulting in three million hospitalizations (Mazur et al., 2023). To address the need for effective RSV treatments, we present the development of a high-throughput fluorescence-based screening to identify antiviral activity in natural extracts (NEs). Our assay, which is based on a 2D cell model run in a 384-well plate, uses a mCherry-carrying recombinant virus to identify hits based on viral replication inhibition. The method was validated using 192 chemodiverse NEs from a plant collection of Pierre-Fabre Laboratoires (PFL) (Allard et al., 2023). All NEs were analyzed by UHPLC-HRMS² to partially assess their chemical composition. Dereplication strategies identified betulinic acid among several active NEs. This compound is known for such activity (Amiri et al., 2020) and its bioactivity profile obtained after targeted isolation indicates that our assay provides data comparable to literature under different screening conditions. It further demonstrates the efficacy of our dereplication pipeline to discard NEs containing betulinic acid. Based on this proof of concept, we have extended our screening to 800 NEs from the PFL. Hits showing >70% inhibition of viral replication and >70% cell viability represented 21% of NEs. Further data mining is currently underway using both the newly developed Experimental NP Knowledge Graph (Burdet et al., 2024) and prior knowledge of NP antivirals with the Lotus initiative (Rutz et al., 2022). This approach aims to identify compounds in NEs with efficacy against RSV, thereby contributing to the discovery of new RSV lead compounds.

Keywords: natural products, respiratory syncytial virus, antiviral activity, screening, UHPLC-HRMS²

References

- Mazur NI, Terstappen J, Baral R, Bardají A, Beutels P, Buchholz UJ, et al. Respiratory syncytial virus prevention within reach: the vaccine and monoclonal antibody landscape. *Lancet Infect Dis.* 2023;23(1):e2-21.
- Allard PM, Gaudry A, Quirós-Guerrero LM, Rutz A, Dounoue-Kubo M, Walker TW, et al. Open and reusable annotated mass spectrometry dataset of a chemodiverse collection of 1,600 plant extracts. *GigaScience.* 2023;12:giac124.
- Amiri S, Dastghaib S, Ahmadi M, Mehrbod P, Khadem F, Behrouj H, et al. Betulin and its derivatives as novel compounds with different pharmacological effects. *Biotechnol Adv.* 2020;38:107409.
- Burdet F, Allard PM, Nothias LF, Kirchhoffer O, Gaudry A, Moretti S, et al. Application of the RDF framework to integrate heterogeneous experimental data of a large chemo- and biodiverse collection from a research collaborative project [Internet]. *ChemRxiv*; 2023.
- Rutz A, Sorokina M, Galgonek J, Mietchen D, Willighagen E, Gaudry A, et al. The LOTUS initiative for open knowledge management in natural products research. *Donoso DA, Akhmanova A, Tapley Hoyt C, eLife.* 2022;11:e70780.

* Presenting author. E-mail address: paola.haemmerli@unige.ch (Paola Haemmerli).

***In vitro* model for predicting oral bioavailability of isoflavone C-glycosides, O-glycosides, and their aglycons**

Sabrina Zölch*, Jörg Heilmann

Department for Pharmaceutical Biology, Faculty of Chemistry and Pharmacy, University of Regensburg, Universitätsstr. 31, 93053 Regensburg, Germany

Isoflavones are widely distributed in plants of the Fabaceae family mainly in legumes such as soy, clover and beans (Křížová et al. 2019). They recently received increased attention due to their putative health properties as a potential alternative therapy against cancer, cardiovascular diseases, and menopausal symptoms like osteoporosis (Gómez-Zorita et al. 2020). The potential biological activity of isoflavones in the human body highly depends on their absorption and metabolism in the small intestine and liver. Thus, in the scope of this project, an *in vitro* model for predicting oral bioavailability was developed. The approach consists of three stages (**Fig. 1**): the small-scale *in vitro* digestion model to study the gastrointestinal behaviour of the flavones, the Caco-2 cell model to analyse the absorption and metabolism in the small intestine, and the HepaRG™ cell model to investigate the hepatic metabolism.




Fig. 1 Overview of the *in vitro* model for predicting oral bioavailability.

The investigations identified various structural motifs of isoflavones influencing their gastrointestinal stability and intestinal absorption. Based on the absorption in the Caco-2 model, the permeability coefficients of the compounds were determined. Moreover, numerous intestinal and hepatic metabolites were detected, and hepatic clearance was calculated. By calibrating the experiments with well-investigated model substances, these results could be correlated to the *in vivo* oral bioavailability. Therefore, the amount of isoflavones available for tissue distribution and thus for exertion of potential physiological effects can be estimated.

Keywords: Isoflavones, glycosides, bioavailability, hepatic clearance, intestinal absorption

References

- Gómez-Zorita, S., González-Arceo, M., Fernández-Quintela, A., Eseberri, I., Trepiana, J., Portillo, M.P., 2020. Scientific evidence supporting the beneficial effects of isoflavones on human health. *Nutrients* 12, 3853.
- Křížová, L., Dadáková, K., Kašparovská, J., Kašparovský, T., 2019. Isoflavones. *Molecules* 24, 1076.

* Presenting author. E-mail address: sabrina.zoelch@ur.de (Sabrina Zölch).

***Dracaena cambodiana* biflavonoid CB2 counteracts ferroptosis by directly inhibiting lipid peroxidation and inducing the key ferroptosis suppressor FSP1**

Loc Le-Xuan^{a,b,*}, Minh Bui-Hoang^a, Lena Hofmann^a, Katharina Hille^a, Pawel Mlotek^a, Stefan Schwaiger^c, Hermann Stuppner^c, Van Anh Tran-Thi^d, Andreas Koeberle^a, Solveigh C. Koeberle^a

^a Michael Popp Institute and Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, Innsbruck 6020, Austria

^b School of Medicine, Vietnam National University Ho Chi Minh City, Ho Chi Minh City 70000, Vietnam

^c Pharmacognosy - Institute of Pharmacy and Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, 6020, Innsbruck, Austria

^d Faculty of Pharmacy, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City 70000, Vietnam

The pathophysiology of many degenerative diseases has recently been linked to ferroptosis, a form of programmed cell death dependent on iron-mediated phospholipid hydroperoxide formation (Dixon et al, 2012). Inhibition of ferroptosis therefore represents a promising therapeutic approach for degenerative diseases characterized by excessive cell death (Stockwell, 2022). Various ferroptosis inhibitors have been identified, but their clinical translation has been hampered (Koeberle et al, 2023). In our search for small molecules targeting ferroptosis, we screened an in-house library of over 200 structurally diverse natural products in human hepatocytes and identified the biflavonoid CB2 ($EC_{50} = 150$ nM) from *Dracaena cambodiana* as one of the most active biogenic ferroptosis inhibitors described so far. CB2 is efficient across different cell types (liver, brain, kidney and lung) and prevents ferroptosis irrespective of the triggering mechanism. CB2 efficiently scavenges radicals and suppresses lipid peroxidation in artificial membranes. At the cellular level, besides suppressing lipid peroxidation, it counteracts ferroptosis by inducing the expression of the key ferroptosis suppressor FSP1 without activating NRF2, the master regulator of antioxidant defense, or altering membrane unsaturation, which determines the susceptibility to ferroptosis. We conclude that CB2 prevents ferroptosis by inhibiting lipid peroxidation and promoting redox homeostasis by targeting a yet elusive factor in the control of FSP1.

Keywords: *Dracaena cambodiana*, ferroptosis, lipid peroxidation inhibition, FSP1

References

- Dixon, S.J., Lemberg, K.M., Lamprecht, M.R., Skouta, R., Zaitsev, E.M., Gleason, C.E., Patel, D.N., Bauer, A.J., Cantley, A.M., Yang, W.S. Morrison, B., 2012. Ferroptosis: an iron-dependent form of nonapoptotic cell death. *Cell* 149, 1060-1072.
- Stockwell, B.R., 2022. Ferroptosis turns 10: Emerging mechanisms, physiological functions, and therapeutic applications. *Cell* 185, 2401-2421.
- Koeberle, S.C., Kipp, A.P., Stuppner, H. Koeberle, A., 2023. Ferroptosis-modulating small molecules for targeting drug-resistant cancer: Challenges and opportunities in manipulating redox signaling. *Med. Res. Rev.* 43, 614-682.

* Presenting author. E-mail address: Loc.Le-Xuan@student.uibk.ac.at (Loc Le-Xuan).

Dietary molecules from *Glycyrrhiza foetida* and their modulation on metabolic syndrome pathways

Elena Serino^{a,*}, Hekmat B. Al-Hmadi^{b,c}, Fabio Arturo Iannotti^d, Arianna Pastore^a, Rosa Maria Vitale^d, Giuseppina Chianese^a, Saoussen Hammami^c, Mariano Stornaiuolo^a, Orazio Taglialatela-Scafati^a

^a Department of Pharmacy, School of Medicine and Surgery, University of Naples Federico II, Via D. Montesano 49, 80131 Napoli, Italy

^b Department of Chemistry, College of Medicine, AL-Muthanna University, Samawah, Iraq

^c Laboratory of Environmental Chemistry and Clean Processes (LR21ES04), Faculty of Sciences of Monastir, Monastir University, Monastir, 5000, Tunisia

^d Institute of Biomolecular Chemistry, National Research Council (ICB-CNR), Via Campi Flegrei 34, 80078, Pozzuoli (NA), Italy

Glycyrrhiza is a genus of edible plants, commonly known as licorice, with a long use in folk medicine for their adaptogenic, anti-inflammatory, antioxidant effects, estrogenic activity and hepatoprotective properties. Among the less investigated species of this genus, *G. foetida* seems to be a new promising tool to treat metabolic syndrome (MetS). In fact, this plant has been pointed out as a source of amorfrutins, acetate-deriving polyphenols that explicate a potent insulin-sensitizer effect as PPAR γ agonists that are quite rare in this genus (Weidner et al., 2012). The minor members of this class of compounds have not been in-depth characterized, thus, after a preliminary LC-UV-MS² dereplication step, we have carried out a NMR-based phytochemical screening of *G. foetida* aerial parts that led us to the characterization of 29 compounds, mainly polyphenols, including 11 previously undescribed amorfrutins and an unprecedented flavanone. The amorfrutins were tested on PPAR γ and PPAR α and, besides amorfrutin A, the most active compound on PPAR γ , we have identified the new amorfrutin H as a selective PPAR α agonist and the unprecedented amorfrutin E as a dual PPAR γ/α agonist (Serino et al., 2023). Moreover, all the compounds were tested on other two MetS dysfunctional pathways, highlighting amorfrutin M and decarboxyamorfrutin A as mitochondrial stimulators, while amorfrutin 2 as a glucose uptake promoter. Thus, *G. foetida* is rich source of many bioactive ingredients that, through to the concomitant modulation of different pathways, can ameliorate MetS, emerging as a new potential functional food.

Keywords: *Glycyrrhiza foetida*, metabolic syndrome, amorfrutins, mitochondrial activity, GLUT

References

- Weidner, C., de Groot, J. C., Prasad, A., Freiwald, A., Quedenau, C., Kliem, M., Sauer, S., 2012. Amorfrutins are potent antidiabetic dietary natural products. PNAS 109, 7257-7262.
- Serino, E., Iannotti, F. A., Al-Hmadi, H. B., Caprioglio, D., Moriello, C., Masi, F., Taglialatela-Scafati, O., 2023. PPAR α/γ -targeting amorfrutin phytocannabinoids from aerial parts of *Glycyrrhiza foetida*. J. Nat. Prod. 86, 2435-2447.

* Presenting author. E-mail address: elena.serino@unina.it (Elena Serino).

How to make “extreme” CPC greener: how to substitute alkanes in biphasic solvent systems using COSMO-RS predicting tools

Clément Nève^{a,*}, Nicolas Borie^a, Sébastien Chollet^d, Luc Marchal^c, Jean-Hugues Renault^a, Alexios-Leandros Skaltsounis^b

^a Université de Reims Champagne Ardenne, CNRS, ICMR UMR 7312, Reims, France

^b Division of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimioupoli Zografou, Athens, Greece

^c Nantes Université, CNRS, GEPEA UMR 6144, Saint-Nazaire, France

^d CPCEng, Capacités, Nantes, France

Centrifugal Partition Chromatography (CPC) is a support-free chromatographic technique involving the distribution and transfer of solutes between two immiscible liquid phases according to their partition coefficients, that avoids irreversible adsorption of the analytes, allows a total recovery of injected samples and enables separations of compounds within a large polarity range. To date, there are few alternatives to alkanes when it comes to purifying highly lipophilic compounds. However, most alkanes are classified as CMR or ecotoxic compounds. Replacing them with «greener» alternatives is an essential objective for the CPC development as an industrial-scale purification technique. To replace alkanes, a lot of alternatives are currently being studied (oils, ethers, esters, NaDES, ionic liquids, *etc.*), but testing all of them in different ternary or quaternary systems, at different compositions and on various extracts is an extremely time and money-consuming approach. COSMO-RS software is a based equilibrium thermodynamics method, placing compounds in a conductor-like environment and calculating a so-called « σ -profile », allowing to compare the similarity between two compounds (*i.e.*, their capacity as hydrogen donor or acceptor), and thus their miscibility. This software allowed the prediction of theoretical intermolecular properties and of ternary, quaternary and solubility diagrams of biphasic solvent systems containing “green” solvents such as CPME, MeTHF, CyreneTM, vegetable oils, *etc.* The predicted properties were compared with experimental data on hydrodynamic behavior with the visual-CPC tool, partitioning, selectivity, *etc.* As a proof of concept, the isolation of neutral triterpenes from *Pistacia lentiscus* has been achieved without using petroleum-based solvents.

Keywords: centrifugal partition chromatography, COSMO-RS, green chemistry, *Pistacius lentiscus*

References

- Claux, O., Santerre, C., Abert-Vian, M., Touboul, D., Vallet, N., Chemat, F., 2021. Alternative and sustainable solvents for green analytical chemistry. *Current Opinion in Green and Sustainable Chemistry*, 31, 100510.
- M. Vilas-Boas, S., W. Cordova, I., A. Kurnia, K., H.S. Almeida, H., S. Gaschi, P., A.P. Coutinho, J., P. Pinho, S., Ferreira, O., 2022, Comparison of two computational methods for solvent screening in countercurrent and centrifugal partition chromatography. *Journal of Chromatography A*, 1666, 462859.
- Sicaire, A.-G., Abert-Vian, M., Fine, F., Carré, P., Tostain, S., Chemat, F., 2015. Experimental approach versus COSMO-RS assisted solvent screening for predicting the solubility of rapeseed oil. *OCL*, 22(4), D404.
- W. Armstrong, D., 1988, Theory and Use of Centrifugal Partition Chromatography. *Journal of Liquid Chromatography*, 11(12), 2433-2446.

* Presenting author. E-mail address: clement.neve@univ-reims.fr (Clément Nève).

Investigating the anxiolytic activity of selected essential oils in zebrafish larvae and identification of active constituents using biochemometrics

Veronica Maphanga^{a,*}, Maxleene Sandasi^{a,b}, Guy Kamatou^a, Gill Enslin^a, Krystyna Skalicka-Woźniak^c, Alvaro Viljoen^{a,b}

^a Department of Pharmaceutical Sciences, Tshwane University of Technology, Private Bag X680, Pretoria, 0001, South Africa

^b SAMRC Herbal Drugs Research Unit, Faculty of Science, Tshwane University of Technology, Pretoria, 0001, South Africa

^c Department of Natural Products Chemistry, Medical University of Lublin, 1 Chodzki Street, 20-093, Lublin, Poland

Anxiety disorders have become prevalent in modern day society mainly due to socioeconomic pressures and pandemics such as COVID-19. Current pharmaceutical interventions which include benzodiazepines have become undesirable due to unpleasant side effects (Hosein Farzaei et al., 2016). Traditional medicines on the other hand, have become popular over the years as they are deemed safe, inexpensive and have fewer side effects. Essential oils (EOs) are herein investigated as natural alternatives for the management of anxiety as they have a long history of traditional use for various central nervous system disorders. Commercially available essential oils ($n = 63$) reported to exhibit possible anxiolytic activity were sourced from Pranarôm International® (Belgium) and Scatters Oils® (South Africa). The oils were chemically profiled on an Agilent gas chromatograph, coupled to a quadrupole mass spectrometer (MS) and a flame ionisation detector (FID) (GC-MS/FID). In order to assess anxiety-like behaviour in five days post fertilisation (5-dpf) zebrafish larvae, the light/dark transitions test was performed in a DanioVision observational chamber (Noldus) and locomotor activity was monitored. A biochemometrics approach was used to correlate essential oil profiles to anxiolytic activity data and subsequently identify bioactive compounds in the essential oils. Twenty nine out of the 63 EOs alleviated anxiety in the larvae to some extent, while *Boswellia carterii*, *Pogostemon cablin* and *Lavandula burnatii* essential oils displayed the best anxiolytic activity with significant reduction in locomotor activity ($p < 0.05$), as well as increased distance travelled, and time spent in the central arena during the dark phase (reverse-thigmotaxis behaviour). Biochemometric analysis identified α -pinene, *p*-cymene, camphene, camphor, eucalyptol, linalyl acetate, and *trans*-anethole as putative biomarkers, and experimental validation confirmed the anxiolytic activity of (+)- α -pinene in zebrafish larvae. The study provides scientific evidence to support the use of some essential oils to alleviate anxiety. Biochemometrics could successfully identify bioactive constituents in the essential oils.

Keywords: essential oils, anxiolytic activity, biochemometrics, zebrafish larvae

References

Hosein Farzaei, M., Bahramsoltani, R., Rahimi, R., Abbasabadi, F., Abdollahi, M., 2016. A systematic review of plant-derived natural compounds for anxiety disorders. *Curr. Top. Med. Chem.* 16, 1924-1942.

* Presenting author. E-mail address: MaphangaBV@tut.ac.za (Veronica Maphanga).

Animal Healthcare and Veterinary Phytotherapy: Science and Practice (VET)

Animal Healthcare and Veterinary Phytotherapy: Science and Practice (VET)

Plant-derived compounds as a solution to robust health status of chickens and turkeys

Paweł Konieczka^{a,b,*}

^a Department of Poultry Science and Apiculture, University of Warmia and Mazury, Oczapowskiego 5, 10-718 Olsztyn, Poland

^b Department of Animal Nutrition, The Kielanowski Institute of Animal Physiology and Nutrition, Polish Academy of Sciences, Instytutcka 3, 05-110 Jabłonna, Poland

Despite the introduction of restrictions on the use of antimicrobials in poultry production, their use remains at a high level, which creates the risk of increasing drug resistance among many pathogens. Poland is currently the EU leader in the production of chickens and turkeys, therefore the introduction of modern solutions in prevention and biosecurity is a key aspect.

This study presents the most up-to-date research conducted in our center on the use of alternative methods of preventing the most important stress factors, including coccidiosis and necrotic enteritis in chickens and turkeys, using bioactive substances of plant origin. Among other, our data indicated that phytogenic products from hops manifested effects that supported the proliferation, in the caeca, of bacteria that were beneficial to the host, which could contribute to the improvement of the litter quality during the last period of rearing. Application of *Yucca schidigera* was effective alternative for anticoccidial treatment in broilers whereas, the water extract of yarrow leaf increased the tissue antioxidant status and improved liver function and lipid metabolism in turkeys. Eventually, we evidence that *Cannabis*-originated bioactive compounds may have a great potential in maintaining gut health status in chickens and turkeys.

Our research results indicate that the use of bioactive substances of plant origin can effectively support the immune system of chickens and turkeys and may constitute an effective prevention of many diseases in these species.

Keywords: bioactive compounds, cannabidiol, gut health, chicken, turkey

* Presenting author. E-mail address: pawel.konieczka@uwm.edu.pl (Paweł Konieczka).

Medicinal Plant based Treatments for Infectious Disease and Infertility in US Organic Dairy Cattle

Hubert J. Karreman, VMD FACVBM

Veterinary Practitioner and co-owner of Reverence Farms, PO Box 129, Graham, NC 27253

The USDA National Organic Program (NOP) requires removal of any animal from organic production if antibiotics are used¹. Some veterinarians use botanicals, biologics and other alternative modalities to treat infection and reproduction issues. While botanical and biological medicines were widely used between 1900-1940², there are still botanicals incorporated into protocols for primary treatment of mastitis and systemic infection such as pneumonia, salmonella and toxic mastitis. Phytotherapy is also being utilized to help stimulate cows to display estrus. Veterinary protocols for successful use of antibiotic alternatives will be shared³. Data from publicly-funded university research will be shown for *in vitro* and *in vivo* clinical studies as well as analytic residue depletion rate studies regarding a specific natural product for udder health^{4,5,6,7,8,9,10,11}. Two studies, done 10 years apart, show differences and similarities between conventional and organic udder health and reproductive health.¹²

Keywords: USDA NOP, antibiotics, infection, mastitis, reproduction

References

USDA NOP 7CFR205.238(c)(7) 2Milks, Howard J. Practical Veterinary Pharmacology and Therapeutics (Balliere, Tindall and Cox), 1930 (among others) 3see www.vcpr.org Karreman protocols 4Mullen, et.al, Journal of Dairy Science Volume 97, Issue 6, June 2014, Pages 3509-3522 5Mullen, et.al, Journal of Dairy Science Volume 97, Issue 9, September 2014, Pages 5587-5591 6Mason, et.al, Pharmacol Clin Toxicol 3(1):1040 7Mullen, et.al, Vet Rec Open 2017 Aug 11;4(1):e000214. 8Mason, et.al, Food Addit Contam Part A Chem Anal Control Expo Risk Assess 2017May;2017May;34(5):740-749. 9McPhee, et.al, Dairy Sci 2011 Apr;94(4):1738-43. doi: 10.3168/jds.2010-3988. 11Pinedo, et.al, Can Vet J. 2013 May; 54(5): 479-484. 12Karreman, World Buiatric Association Proceedings, Dublin, 2016.

* Presenting author. E-mail address: Hubert.karreman@gmail.com (Hubert Karreman).

Effects of Oregano Essential Oil and Purple Garlic Powder in weaning piglets as an alternative to Zinc Oxide

Jorge Rivera-Gomis^{a,*}, Daniel Serrano-Jara^b, María José Cubero Pablo^c

^a Scotland's Rural College (SRUC), Centre for Epidemiology and Planetary Health, Inverness, Scotland IV2 5NA, UK

^b CIAGRO Institute, University of Elche Miguel Hernández, Ctra de Benial, Km 3,2, 03312 Orihuela, Spain

^c Animal Health Department, Veterinary Medicine Faculty, Regional Campus of International Excellence "Campus Mare Nostrum", University of Murcia, Espinardo, 30100 Murcia, Spain

This talk discusses the effects of including Oregano Essential Oil (OEO) and Purple Garlic Powder (PGP) in weaning piglet feed, comparing different doses and combinations with Zinc Oxide (ZnO). The studies were conducted under commercial farm conditions with crossbred piglets of 21 days of age. We studied the effect of these compounds on biomarkers of oxidative status, stress, and inflammation using saliva and serum samples, alongside average daily gain and feed conversion ratio (Rivera-Gomis et al., 2020). Additionally, impacts on intestinal health and growth performance were analysed over seven weeks post-weaning. Samples from the jejunum and ileum were used to quantify goblet cells, intraepithelial lymphocytes, and IgA-producing cells, while growth performance was monitored at the start and end of the trial (Serrano-Jara et al., 2023a). Degradation of the bioactive components of OEO and PGP during storage in silos, their effects on jejunum and ileum morphometry, and cecum microbiota as health indicators were examined, along with monitoring antimicrobial resistance in commensal *E. coli* (Serrano-Jara et al., 2023b).

High doses of OEO and PGP were found ineffective and potentially harmful regarding oxidative status, stress, and inflammation. Conversely, higher doses improved growth performance similarly or better than ZnO, enhancing piglet intestinal health without compromising growth. Supplementation with PGP 2%, OEO 1.2%, and PGP 2% + OEO 1.2% offered a natural alternative to ZnO without sacrificing efficacy. Overall, optimal dosages of these additives yielded results comparable or superior to ZnO, highlighting the importance of dosage when administering these products to piglets.

Keywords: Weaning piglets, oregano essential oil, purple garlic powder, zinc oxide, intestinal health

References

- Rivera-Gomis, J., Peres Rubio, C., Martínez Conesa, C., Otal Salaverri, J., Cerón, J. J., Tortosa, D. E., Cubero Pablo, M. J., 2020. Effects of dietary supplementation of garlic and oregano essential oil on biomarkers of oxidative status, stress and inflammation in postweaning piglets. *Animals*. 10(11), 2093.
- Serrano-Jara, D., Rivera-Gomis, J., Tornel, J. A., Bernabé, A., Martínez-Conesa, C., Navarro, J. A., Cánovas, R., Otal, J., Cubero, M. J., 2023a. Effects of dietary supplementation with purple garlic powder and oregano essential oil on intestinal health in post-weaning piglets from commercial farms. *Vet. Res. Commun.* 47(2), 901-909.
- Serrano-Jara, D., Rivera-Gomis, J., Tornel, J. A., Jordán, M. J., Martínez-Conesa, C., Pablo, M. J. C., 2023b. Oregano Essential Oil and Purple Garlic Powder Effects on Intestinal Health, Microbiota Indicators and Antimicrobial Resistance as Feed Additives in Weaning Piglets. *Animals*. 14(1), 111.

* Presenting author. E-mail address: Jorge.Gomis@sruc.ac.uk (Jorge Rivera-Gomis).

Effect of gastro-resistant pellets of thymol on its absorption rate, intestinal wall integrity and accumulation in the rabbit tissues

Radoslava Kristofova^{a,*}, Karin Zitterl-Eglseer^b, Maria Teresa Capucchio^{c,e}, Dorotea Ippolito^{c,d}, Francesco Gai^e, Fardad Firooznia^f, Ales Franc^g, Iveta Placha^a

^a Centre of Biosciences-Slovak Academy of Sciences-Institute of Animal Physiology, 04001 Kosice, Slovak Republic

^b Centre of Animal Nutrition and Welfare, University of Veterinary Medicine Vienna, 1210 Vienna, Austria

^c Department of Veterinary Sciences, University of Turin, 10095 Grugliasco, Italy

^d National Institute of Health, Department of Food Safety, Nutrition and Veterinary Public Health, 00161 Roma, Italy

^e National Research Council of Italy-Institute of Sciences of Food Production, 10095 Grugliasco, Italy

^f Department of Biology, City College of New York, NY 10031, New York, USA

^g Department of Pharmaceutical Technology, Faculty of Pharmacy, Masaryk University Brno, 61200 Brno, Czech Republic

Thymol, a major compound of thyme (*Thymus vulgaris* L.), exhibits strong antioxidant properties related to its phenolic structure. One of the crucial aspects of the beneficial effect of thymol is the amount present in the gut as a result of its release from feed, and consequent ability to pass through the intestinal wall in each part of the gastrointestinal tract.

To improve the stability of thymol against enzymatic degradation, ensure its penetration into the intestinal mucosa and achieve desired therapeutic levels in rabbit tissues, we incorporated coated self-emulsifying pellets of thymol into rabbit diets. Forty-eight rabbits were allocated to control and experimental group with thymol addition (250 mg/kg feed) for 21 days and withdrawal for 7 days. The level of thymol in different tissues was analysed by means of solid phase microextraction and GC/MS; while histomorphometry of different intestinal segments and metabolic active organs (liver and kidney) were investigated.

Intensive absorption of thymol in duodenal and jejunal wall was shown. It could be detected in colon, caecum, liver, kidney, spleen, fat and faeces at both sampling time points, but only in trace amounts in muscle after its withdrawal. Statistical analyses showed that the treatment did not impair the gut histomorphometry and did not influence the severity and distribution of the histological lesions observed in the metabolic active organs.

Keywords: thymol pellets, metabolic pathway, rabbit

Funding: VEGA 2/0009/20, 2/0005/24; CNR-SAS-2022-02, Open-Mob-2022-01, COST CA22109, Action Austria-Slovakia MPC-2023-01046

* Presenting author. E-mail address: kristofova@saske.sk (Radoslava Kristofova).

Use of botanical extracts as tranquilizer in fattening animals

Josep Casadellà-Xifra^a, Javier Álvarez-Rodríguez^a, Emma Fàbregab^b, Isabel Blanco-Penedo^{a,*}

^a Department of Animal Science, University of Lleida, 25198 Lleida, Spain

^b Animal Welfare Program, IRTA, Monells (Spain)

The stress response induces significant behavioral, biochemical, and immunological changes that directly affect animal welfare and farm productivity. This study aimed to evaluate the effectiveness of a tranquilizer based on botanical extracts (Quiet-farm®) in reducing stress levels in a commercial fattening cattle farm (Study 1) and in fattening pigs (Study 2). A total of 142 uncastrated Frisian males were used in Study 1 (145 ± 1 days; 179 ± 5 kg). In Study 2, a total of 135 Danbread x Pietrian pigs (females and uncastrated males) were used (11-12 weeks; 47 ± 5 kg). In both studies, animals were randomly assigned to a control group (fed without tranquilliser supplementation) and a treatment group (fed with tranquilliser supplementation). Several measures of animal behaviour and health proposed in the Welfare Quality® protocols (2009a, b) were used to assess product efficacy, as well as animal performance data. In Study 2, stress biomarkers (salivary and hair cortisol) were also assessed. In term of outcomes, in both studies there was a significantly lower frequency of agonistic and sexual behaviors, as well as a lower prevalence of integument alterations in the treatment group. In terms of productive outcomes, average daily gain was also higher in the treatment group in both studies. In study 2, the treatment group had significantly lower hair cortisol than the control group (no statistical differences were found for salivary cortisol). The present results suggest that the phytogenic tranquiliser could improve animal welfare and production parameters in these production systems.

Keywords: Botanical extracts, stress, fattening calves, fattening pigs

References

Welfare Quality® (2009a). Welfare Quality® assessment protocol for cattle. Welfare Quality® Consortium.

Welfare Quality® (2009b). Welfare Quality® assessment protocol for pigs (sows and piglets, growing and finishing pigs). Welfare Quality® Consortium.

* Presenting author. E-mail address: isabel.blancopenedo@udl.cat (Isabel Blanco-Penedo).

The food-chain transfer of ecdysteroids and its potential significance in view of human and animal health

Attila Hunyadi^{a,b,*}

^a *Institute of Pharmacognosy, University of Szeged, Eötvös str. 6, 6720 Szeged, Hungary*

^b *HUN-REN-SZTE Biologically Active Natural Products Research Group, Eötvös str. 6, H-6720 Szeged, Hungary*

Phytoecdysteroids are a highly diverse group of natural products. As of today, 577 of these compounds have been identified in plants and fungi. They are analogs of the insect molting hormone, 20-hydroxyecdysone (20E). When consumed, phytoecdysteroids are toxic to insects but in vertebrates they act as non-hormonal anabolic and cytoprotective agents. In mammals, 20E activates the protective arm of the renin-angiotensin system via the Mas (angiotensin(1–7)) receptor (Lafont et al, 2021); this mechanism is highly relevant in cardiovascular and pulmonary diseases.

We have discovered that pharmacologically relevant amounts of minor phytoecdysteroids may pass the food chain from plants through insects to insectivorous animals including songbirds and bats. As a resistance mechanism, insects can detoxify ecdysteroids through ester formation with long-chain fatty acids. Our work revealed that such lipophilic conjugates may accumulate in insects and act as ecdysteroid pro-drugs in insectivorous vertebrates. Our studies on a large insectivorous bird population revealed ecdysteroid blood levels at the several micromolar concentration range, which could be linked with a pathological moulting of the birds' ectoparasites such as ticks. This presentation will cover most recent developments in the study of this remarkable chemical interaction between the Kingdoms of Nature and provide an overview to its significance concerning human and animal health.

Keywords: phytoecdysteroid, zooecdysteroid, ester pro-drug, insectivorism, food chain

Acknowledgments: Financial support from the NRDIO, Hungary (K-146359 and TKP2021-EGA-32) is acknowledged.

References

- Lafont, R., Serova, M., Didry-Barca, B., Raynal, S., Guibout, L., Dinan, L., Veillet, S., Latil, M., Dioh, W., Dilda, P.J., 2021. 20-Hydroxyecdysone activates the protective arm of the RAAS via the MAS receptor. *J. Mol. Endocrinol.* 68, 77-87.
- Hornok, S., Kováts, D., Flaisz, B., Csörgő, T., Könczöl, Á., Balogh, G.T., Csorba, A., Hunyadi A., 2016. An unexpected advantage of insectivorism: insect moulting hormones ingested by song birds affect their ticks. *Sci. Rep.* 6, 23390.
- Hornok, S., Csorba, A., Kováts, D., Csörgő, T., Hunyadi A., 2019. Ecdysteroids are present in the blood of wild passerine birds. *Sci. Rep.* 9, 17002.
- Hornok, S., Berkecz, R., Sós, E., Sándor, D.A., Körmöczi, T., Solymosi, N., Kontschán, J., Hunyadi, A., 2022. Arthropod moulting hormones (ecdysteroids) are present in the blood of insectivorous bats. 52, 317-321.

* Presenting author. E-mail address: hunyadi.attila@szte.hu (Attila Hunyadi).

Behavioral and pharmacological investigation of putative self-medicative natural materials used by wild chimpanzees and mountain gorillas

Fabien Schultz^{a,b}, Inken Dworak-Schultz^a, Florian Wald^{a,x}, Lisa Grohs^a, Tabitha Iker^a, Godwin Anywar^c, Gladys Kalema-Zikusoka^d, Matthew McLennan^e, Catherine Hobaiter^f, Lewis Marquez^g, Sunmin Woo^g, Luc Pieters^h, Guy Caljonⁱ, Cassandra Quave^g, Elodie Freymann^j, Leif-Alexander Garbe^{a,j}

^a Ethnopharmacology & Zoopharmacognosy Research Group, Department of Agriculture and Food Sciences, Neubrandenburg University of Applied Sciences, Brodaer Str. 2, 17033 Neubrandenburg, Germany

^b Pharmacognosy and Phytotherapy, UCL School of Pharmacy, 29-39 Brunswick Square, London WC1N 1AX, London, UK

^c Department of Plant Sciences, Microbiology and Biotechnology, Makerere University, Uganda

^d CTPH - Conservation Through Public Health, Uganda

^e Bulindi Chimpanzee & Community Project, Bulindi, Hoima District, Uganda

^f School of Psychology and Neuroscience, University of St. Andrews, Scotland

^g School of Medicine, Emory University, Atlanta, GA, USA

^h Department of Pharmaceutical Sciences, University of Antwerp, Belgium

ⁱ Department of Biomedical Sciences, University of Antwerp, Belgium

^j Primate Models for Behavioural Evolution Lab, Institute of Human Sciences, Department of Anthropology and Museum Ethnography, University of Oxford, Oxford, UK

^k ZELT - Center for Nutrition and Food Technology gGmbH, Seestr. 7a, 17033 Neubrandenburg, Germany

This study examined the self-medicative behavior of the African Great Apes, e.g. ingestion or topical application of medicinal plants, mushrooms or insects. Presumed self-medication was first described in East African chimpanzees in the early 1960s. Today, most of the natural remedies used by great apes remains undiscovered, and only a few have been investigated for their medicinal efficacy potential. Awareness of the pharmacological properties of plants, mushrooms or insects must have originated from a long-running evolutionary connection between apes (and humans) and their search for remedies in nature. To gain insight into apes' potential medicinal use, we observed wild chimpanzees and mountain gorillas in their natural habitats in tropical Uganda in: 1) the remote montane Bwindi Impenetrable Forest; 2) the moist, semi-deciduous Budongo Forest and 3) a highly disturbed unforested area in Bulindi. Many species were reported for the first time. A total of 83 plant, mushroom and insect samples were collected from sites across Uganda, extracted and taken to the lab. Here, the "traditional use" of our closest animal relatives was evaluated through observational data and pharmacological *in vitro* screenings, including a) antimalarial assays against chloroquine-resistant *Plasmodium falciparum* K1; b) antiinflammatory assays via a COX-2/1 inhibition model, c) antibiotic assays against multiresistant ESKAPE strains and d) cytotoxicity evaluation against human MRC-5 lung fibroblasts. Results may reveal novel natural remedies and drug leads that are effective in humans, therefore addressing the worldwide need for new painkillers, antiparasitics, antimalarials and antibiotics.

Keywords: animal self-medication, chimpanzees, mountain gorillas, Bwindi, Bulindi, Budongo, Uganda, zoopharmacognosy, ESKAPE pathogens

* Presenting author. E-mail address: schultz@hs-nb.de (Fabien Schultz).

Variation in chemical profile and antibacterial activity of different parts of *Balanites aegyptiaca* (L.) Delile, (Zygophyllaceae) among two contrasted zones in Benin

Jéronime Ouachinou^a, Hospice Dassou^{a,*}, Yaya Koudoro^b, Jérrold Agbankpé^c, Aristide Adomou^a

^a Laboratoire de Botanique et Ecologie Végétale, Département de Biologie Végétale, Faculté des Sciences et Techniques, Université d'Abomey-Calavi, Benin

^b Laboratoire d'Etude et de Recherche en Chimie Appliquée, Ecole Polytechnique d'Abomey-Calavi, Université d'Abomey-Calavi, 01 BP 2009 Cotonou, Benin

^c Laboratoire de Recherche en Biologie Appliquée, Ecole Polytechnique d'Abomey-Calavi, Université d'Abomey-Calavi, 01 BP 2009 Cotonou, Benin

Balanites aegyptiaca (L.) Delile (Zygophyllaceae), native to Africa and Asia, is widely used for its ethnoveterinary value (Dassou et al. 2015). In Benin, the plant grows in two ecological (dry and humid) zones (Akoègninou et al. 2006) where it is strongly involved in the treatment of bovine diarrhea (Ouachinou et al. 2019). So far, influence of ecological region and plant parts on the bioavailability of chemical compounds and subsequent biological activities are rarely documented. In this study, we evaluated the chemical composition and the antibacterial effects of leaves, stem barks and roots from the both zones in Benin. Many secondary metabolites notably flavonoids, tannins, alkaloids, saponins were identified and their content varied with the plant part and the ecological zone. The total phenolic contents (TPC) of the different extracts varied from 0.99 mg GAE/g in the humid zone to 8.03 mg GAE/g in the dry zone. Considering plant parts, the TPC is more preponderant in leaves than any other part with the highest value in the humid zone (17.29 ± 0.01 mg GAE/g). The trend is the same for total flavonoids and tannins. The antibiotic power of different extracts depends on the ecological zone and plant parts. Root extracts showed the best antibacterial effects on isolates of clinical and reference strains with bactericidal power on *Salmonella*. *B. aegyptiaca* extracts possesses antidiarrheal properties justifying its traditional use for the treatment of bovine diarrhea. Our further research efforts will be focused on elucidating the molecular structures of the bioactive molecules and establishing their mechanism of action and probable toxicological properties of the species.

Keywords: *Balanites aegyptiaca*, aqueous extract, chemical compounds, antidiarrheal.

References

- Akoègninou, A., Van der Burg, W.J., van der Maesen, L.J.G., 2006. Flore Analytique du Bénin. Backhuys Publishers, Leiden, Pays-Bas, 1063 pp.
- Dassou, G.H., Adomou, A.C., Yédomonhan, H., Ogni, A.C., Tossou, G.M., Dougnon, J.T., Akoègninou, A., 2015. Flore médicinale utilisée dans le traitement des maladies et symptômes animaux au Bénin. *Journal of Animal and Plant Sciences* 26: 4036-4057.
- Ouachinou, J.M.-A.S., Dassou, G.H., Idohou, R., Adomou, A.C., Yédomonhan, H., 2019. National inventory and usage of plant-based medicine to treat gastrointestinal disorders with cattle in Benin (West Africa). *South African Journal of Botany* 122: 432-446.

* Presenting author. E-mail address : daspice2@gmail.com (Dassou Hospice)

Orals

Opening Lecture

Carbohydrate-based vaccines, therapeutics and diagnostics and medicines produced in flow

Peter H. Seeberger

Max-Planck Institute for Colloids and Interfaces Potsdam, Germany

Rapid preparation of polysaccharides is by automated glycan assembly (AGA) (Guberman et al., 2019) using a synthesizer (Hahm et al., 2017) provides access to diverse glycans as large as 151-mers (Joseph et al., 2020). I will describe a medicinal chemistry approach to the development of semi- and fully synthetic glycoconjugate vaccines against severe bacterial infections, including resistant hospital microorganisms (Seeberger et al., 2021). Synthetic glycans are key in combination with single molecule imaging, (Wu et al., 2020) to address fundamental questions of carbohydrate structure, folding and material science (Fittolani et al., 2023). Continuous flow systems are useful for automated reaction optimization (Chatterjee et al., 2018) and the modular assembly of active pharmaceutical ingredients (Ghislieri et al., 2015) have become increasingly interesting to practitioners of synthetic chemistry. The new “radial synthesis” concept enables the autonomous production of medications (Chatterjee et al., 2020). Production of the anti-malaria drug artemisinin and its use against different cancers will serve as an example for a drug produced in flow (Triemer et al., 2018).

Keywords: carbohydrate-based vaccines, bacterial infections

References

- Chatterjee, S., Guidi, M., Seeberger, P.H., Gilmore, K., 2020. Automated radial synthesis of organic molecules. *Nature* 579, 379-384.
- Chatterjee, S., Moon, S., Hentschel, F., Gilmore, K., Seeberger, P.H., 2018. An empirical understanding of the glycosylation reaction. *J. Am. Chem. Soc.* 140, 11942-11953.
- Fittolani, G., Tyrikos-Ergas T, Poveda A, Yu Y, Yadav N, et al., 2023. Synthesis of a glycan hairpin. *Nature Chem.* 15, 1461-1469.
- Ghislieri, D., Gilmore, K., Seeberger, P.H., 2015. Chemical assembly systems: layered control for divergent, continuous, multistep syntheses of active pharmaceutical ingredients. *Angew. Chem. Int. Ed. Engl.* 54, 678-682.
- Guberman, M., Seeberger, P.H., 2019. Automated glycan assembly: a perspective. *J. Am. Chem. Soc.* 141, 5581-5592.
- Hahm, H.S., Schlegel, M.K., Hurevich, M., Eller, S., Schuhmacher, F., et al., 2017. Automated glycan assembly using the Glyconeer 2.1 synthesizer. *Proc. Nat. Acad. Sci. USA* 114, E3385-E3389.
- Joseph, A., Pardo-Vargas, A., Seeberger, P.H., 2020. Total synthesis of polysaccharides by automated glycan assembly. *J. Am. Chem. Soc.* 142, 8561-8564.
- Seeberger, P.H., 2021. Discovery of semi- and fully-synthetic carbohydrate vaccines against bacterial infections using a medicinal chemistry approach. *Chem. Rev.* 121, 3598-3626.
- Triemer, S., Gilmore K., Vu G.T., Seeberger P.H., Seidel-Morgenstern A., 2018. Literally green chemical synthesis of artemisinin from plant extracts. *Angew. Chem. Int. Ed. Engl.* 57, 5525-5528.
- Wu, X., Delbianco, M., Anggara, K., Michnowicz T., Pardo-Vargas A., et al., 2020. Imaging single glycans. *Nature* 582, 375-378.

* Presenting author. E-mail address: peter.seeberger@mpikg.mpg.de (Peter H. Seeberger).

Plenary Lectures

Innovative analytical methods for evaluation of efficacy, safety and quality of raw materials and final products in the phyto-area

Guenther K. Bonn

ADSI - Austrian Drug Screening Institute, University of Innsbruck, Innrain 66a, 6020 Innsbruck, Austria

Analytical chemistry and cell biology have significantly advanced natural product research. Developments in chromatography have opened new avenues for the analysis and isolation of natural products, crucial due to the complexity of plant extracts, which contain hundreds to thousands of compounds at varying concentrations. Novel enrichment and purification methods (Huber et al., 2020), particularly advanced solid-phase extraction combined with high-resolution chromatographic separation and mass spectrometric detection (LC-MS), are essential for accurate phytochemical profiling and the quantification of specific metabolites (Kreidl et al., 2020). Combining separation science with spectroscopy, such as near- and mid-infrared (IR) spectroscopy, facilitates rapid, non-invasive qualitative and quantitative analyses of raw plants and extracts. The integration of modern drone technology with NIR spectroscopy expands applications in plant cultivation, enabling efficient determination of optimal harvest times. Traditional analytical methods focus on detecting low molecular weight compounds like phenolics, but the detection of DNA and proteins in plants is gaining importance (Mergner and Kuster, 2022). Plant proteins can serve as active ingredients or allergens, and their identification and quantification are critical for quality control, ensuring the effectiveness, quality, safety, and authenticity of plant products. The latest advancements in protein sequencing, specifically using Quantum-Si's Platinum® instrument for Next-generation protein sequencing™ (NGPS), allow for the direct sequencing of peptide barcodes with single-molecule resolution. This benchtop technology merges DNA-based method accessibility with the innovative capabilities of peptide-based barcoding, enabling the precise identification of unknown phyto-proteins and bioactive protein isoforms. These analytical methods not only enhance research and development but also improve quality control in phyto-analysis (Mair et al., 2023). Comprehensive biological studies complement these analytical approaches, identifying biological activities (Leitner et al., 2022) and conducting necessary safety assessments in phyto-pharmacy, phyto-cosmetics, and phyto-nutrition.

Keywords: natural compounds, analytical chemistry, phyto, NIR-spectroscopy, cell-biology

References

- Huber, S., Harder, M., Funck, K., Erharter, K., Popp, M., Bonn, G.K., Rainer, M., 2020. Novel room temperature ionic liquid for liquid-phase microextraction of cannabidiol from natural cosmetics. *Separations* 7, 45.
- Kreidl, M., Rainer, M., Jakschitz, T., Bonn G.K., 2020. Determination of phototoxic furanocoumarins in natural cosmetics using SPE with LC-MS. *Anal. Chim. Acta* 1101, 211-221.
- Leitner P. Jakschitz, T., Gstir, R., Stuppner, S., Perkams, S., et al., 2022. Anti-inflammatory extract from soil algae *Chromochloris zofingiensis* targeting TNFR/NF- κ B signaling at different levels. *Cells* 11, 1407.
- Mair, A., Fischnaller, M., Lutz, O.M.D., Corradini, D., Skaltsounis, A.L. et al., 2023. Quantification of health claim-relevant tyrosol and hydroxytyrosol after direct hydrolysis improves customer understanding and mitigates market distortion. *Separations* 10, 268.
- Mergner, J. Kuster B., 2022. Plant proteome dynamics. *Annu. Rev. Plant Biol.* 73, 67-92.

* Presenting author. E-mail address: guenther.bonn@uibk.ac.at (Guenther K. Bonn).

Foundational principles can inspire bioactive natural products research

Guido F. Pauli

Pharmacognosy Institute and Department of Pharmaceutical Sciences, University of Illinois Chicago College of Pharmacy, Chicago, IL 60612, USA

Pharmacognosy approaches to the discovery of bioactive natural products (NP) can progress by two main mechanisms: improving interdisciplinary integration and advancing the basic sciences of the disciplines involved. Thus, the challenges in our field arise equally from the interdisciplinarity and basic science perspectives. Geoff Cordell's recent work, which can be acronymized as "L.O.V.E. for cyberecoethnopharmacolomics" (Cordell, 2019; 2024), addresses important public health needs for and the potential pathways toward better scientific integration. This presentation will focus on the flipside of the advancement challenge coin and discuss how foundational biological, chemical, physical, and mathematical principles affect today's integrated NP research methodology. The aim is to gain insights into how the application of overarching fundamental knowledge is a prerequisite for subsequent integration and can play a crucial role in advancing biomedicine collectively. To demonstrate how both long-existing and recently advanced foundational knowledge and methodology can inspire progress in applied NP research, we will examine a range of questions along the biology to mathematics continuum:

- What processes drive the discovery of bioactives and who produces them?
- What steps should be considered to help develop screening hits to clinical translation?
- Can bioactive NPs be "too good to be true"?
- What is a NP "compound" or "isolate"? How many structures does it have?
- What physiochemistry stands behind the purification of bioactive NPs?
- Is methodological innovation the first objective or a last resort?
- What is the difference between peak overlap and a multiplet? How can they be resolved?

To answer these questions as concisely as possible, the discussion will include experimental evidence from our own recent work in the discovery of antibiotics, dental biomodifiers, bioactive principles from medicinal and dietary plants, and methodology for pharmaceutical integrity analysis. The broader discussion and outlook part of the presentation will seek to reflect on the contemporary role of philosophy in Ph.D.-level NP research and the nature of experimental evidence in our framework of scientific understanding.

Keywords: bioactive natural products, discovery, advancement of research methodology

References

Cordell, G.A., 2019. Cyberecoethnopharmacolomics. *J. Ethnopharmacol.* 244, 112134.

Cordell, G.A., 2024. The contemporary nexus of medicines security and bioprospecting: a future perspective for prioritizing the patient. *Nat. Prod. Bioprospect.* 14, 11-41.

* Presenting author. E-mail address: gfp@uic.edu (Guido F. Pauli).

Unusual enzyme reactions in natural product biosynthesis

Ikuro Abe

Graduate School of Pharmaceutical Sciences, The University of Tokyo, Tokyo, Japan

β -Nicotinamide adenine dinucleotide (β -NAD) is a pivotal metabolite for all living organisms and functions as a diffusible electron acceptor and carrier in the catabolic arms of metabolism. Furthermore, β -NAD is involved in diverse epigenetic, immunological, and stress-associated processes, where it is known to be sacrificially utilized as an ADP-ribosyl donor for protein and DNA modifications, or the generation of cell-signaling molecules. Here, we report the function of β -NAD in secondary metabolite biosynthetic pathways, in which the nicotinamide dinucleotide framework is heavily decorated and serves as a building block for the assembly of a novel class of natural products. The gatekeeping enzyme of the discovered pathway (SbzP) hereby catalyzes a sophisticated, pyridoxal phosphate (PLP)-dependent (3+2)-annulation reaction between β -NAD and S-adenosylmethionine (SAM), generating a 6-azatetrahydroindane scaffold. Members of this novel family of β -NAD-tailoring enzymes are widely distributed in the bacterial kingdom and encoded in diverse biosynthetic gene clusters. The findings of this work set the stage for the discovery and exploitation of β -NAD-derived natural products (Hu et al. 2019; Barra et al., 2021).

Keywords: β -NAD, S-adenosylmethionine, β -NAD-derived natural products, PLP-dependent enzyme

References

- Hu, Z., Awakawa, T., Ma, Z. Abe, I., 2019. Aminoacyl sulfonamide assembly in SB-203208. *Biosynthesis*. Nat. Commun. 10, 184, 10 pages.
- Barra, L., Awakawa, T., Shirai, K., Hu, Z., Bashiri, G., Abe, I., 2021. β -NAD as a building block in natural product biosynthesis. *Nature* 600, 754-767.

* Presenting author. E-mail address: abei@mol.f.u-tokyo.ac.jp (Ikuro Abe).

AFERP Association francophone pour l'enseignement et la recherche en pharmacognosie

Medicinal plants in the fight against malaria: investigating the *Strychnos* genus, *Artemisia* sp. and ellagic acid derivatives.

Michel Frédérick^{a,*}, Olivier Bonnet^a, Gilles Degotte^a, Lucia Mamede^{a,b}, Methodius Lahngong^a, Jacob Souopgui^c, Luc Angenot^a, Joëlle Quetin-Leclercq^b, Pierre Francotte^a, Allison Ledoux^a

^a CIRM – Center for Interdisciplinary Research on Medicines, Université de Liège, Avenue Hippocrate 15, B4000 Liège, Belgium

^b LDRI -Louvain Drug Research Institute, Université Catholique de Louvain, Avenue Mounier 72/B1.72.03, 1200 Woluwe-Saint-Lambert, Belgium

^c IBMM – Institute of Molecular Biology and Medicine, Université Libre de Bruxelles, Rue Prof. Jeener & Brachet, 12, 6041 Gosselies, Belgium

Malaria, a parasitic disease transmitted by *Anopheles* mosquitoes, remains a significant global health threat despite advancements in control efforts. In 2022, it accounted for 608,000 deaths and afflicted 249 million individuals worldwide, with 94% of fatalities occurring in Africa (WHO, 2023). The Pharmacognosy Laboratory in Liège, Belgium, has dedicated years to exploring medicinal plants for novel antimalarial compounds, particularly focusing on African flora. Over three decades, we have investigated more than a hundred plants, aiming to identify promising compounds or validate traditional remedies. Three projects of the laboratory will be shortly presented. The first project concerns bisindole alkaloids derived from *Strychnos* species, some showing potent antiplasmodial activity in both *in vitro* and animal studies, suggesting potential as lead compounds (Beaufays et al., 2018). Recently, molecular networking methods were applied to unveil new original antiplasmodial indole alkaloids (Bonnet et al., 2023). Another initiative began with screening Rwandan medicinal plants, leading to the discovery of ellagic acid as active compound (Muganga et al., 2014) and the development of semi-synthetic derivatives with antimalarial properties (Degotte et al., 2023). The third project is dedicated to *Artemisia annua* and *Artemisia afra*, well-known medicinal plants, focusing on standardization, metabolomics-based mode of action studies, and development of a clinical trial. These projects highlight the ongoing relevance of medicinal plants in malaria research. Despite modern pharmaceutical advancements, natural compounds continue to offer significant potential in the fight against this deadly disease.

Keywords: *Strychnos* sp., *Artemisia annua*, *Artemisia afra*, ellagic acid, malaria

References

- Beaufay, C., Ledoux, A., Jansen, O., Bordignon, A., Zhao, S., Teijaro, C.N., Andrade, R.B., Quetin-Leclercq, J., Frédérick, M., 2018. *In vivo* antimalarial and antitrypanosomal activity of strychnogucine B, a bisindole alkaloid from *Strychnos icaja*. *Planta Medica* 84, 881–885.
- Bonnet, O., Beniddir, M.A., Champy, P., Degotte, G., Mamede, L., Desdemoustier, P., Ledoux, A., Tchinda, A.T., Angenot, L., Frédérick, M., 2022. Unveiling antiplasmodial alkaloids from a cumulative collection of *Strychnos* extracts by multi-informative molecular networks. *Front. Mol. Biosci.* 26, 967012.
- Degotte, G., Pendeville, H., di Chio, C., Ettari, R., Pirotte, B., Frédérick, M., Francotte, P., 2023. Dimeric polyphenols to pave the way for new antimalarial drugs. *RSC Med. Chem.* 14, 715–733.
- Muganga, R., Angenot, L., Tits, M., Frédérick, M., 2014. *In vitro* and *in vivo* antiplasmodial activity of three Rwandan medicinal plants and identification of their active compounds. *Planta Med.* 80, 482–489.
- WHO, 2023, World Malaria Report. <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2023>, accessed 9th of April, 2023.

* Presenting author. E-mail address: M.Frederich@uliege.be (Michel Frédérick).

Pharmacognosy faces new challenges: proposals to combine chemometrics and annotations based on mass spectrometry and NMR for more accurate and faster identification of bioactive natural products in complex matrices

S  verine Derbr  ^{a,*}, Manon Meunier^a, Antoine Brugui  re^a, Dimitri Br  ard^a, S  verine Boisard^a, Patricia Blanchard^a, Marc Litaudon^{b,c}, Khalijah Awang^c, Fr  d  ric Saubion^d, Pascal Richomme^a, Andreas Schinkovitz^a

^a Univ Angers, SONAS, SFR QUASAV, F-49000 Angers, France

^b Institut de Chimie des Substances Naturelles, CNRS-ICSN, UPR 2301, Universit   Paris-Saclay, 91198 Gif-sur-Yvette, France

^c IFM NatProLab, University of Malaya, Faculty of Sciences, Department of Chemistry, Malaysia

^d Univ Angers, LERIA, SFR MathSTIC, F-49000 Angers, France

With global warming, reduced biodiversity and the emergence of new pathogens for humans and their resources, researchers including natural products (NPs) chemists need to make the most of new regulatory, modelling and data-sharing tools to make faster and better progress regarding NPs valorization. In that context, as NMR spectrometers now provide useful data sets in a reasonable time frame, we made the MixONat software freely available for dereplication in 2020 (Brugui  re et al. 2020, 2021, <https://sourceforge.net/projects/mixonat>). It allows the processing of ¹³C chemical shifts (δ_c) as well as DEPT 135 and 90 data for a critical carbon type (*i.e.* CH₃, CH₂, CH and C) filtering. It requires experimental or predicted δ_c databases (DB) and displays interactive results that can be refined based on the user's phytochemical knowledge. To go further, this presentation proposes an integrated strategy combining tandem mass spectrometry (MS²)-based molecular networking (MN), a partial least squares (PLS) chemometric model, as well as ¹³C NMR-based dereplication using MixONat software. In addition, an advanced glycation end products (AGEs) assay was used for activity evaluation. The approach was implemented on a *Garcinia parvifolia* bark extract, which contains a high content of prenylated xanthenes and had previously shown a remarkable inhibitory effect on AGE formation (Fig. 1) (Meunier et al. 2023).



Fig. 1. Workflow for an activity-based molecular network approach that integrates statistical data (VIP score and RC) and annotation based on both ¹³C NMR and MS²-based dereplication for a higher level of confidence.

As a main result the proposed strategy permitted the identification of potentially active NPs within complex mixtures and their annotation with a higher level of confidence by both MS² and NMR data. Overall, this comprehensive approach provides a powerful and efficient solution for the targeting and annotating of active compounds in complex NPs mixtures.

Keywords: annotation, bioactivity-based molecular networking, Clusiaceae, dereplication, *Garcinia parvifolia*, in silico databases, level of confidence, MixONat, xanthenes

References

- Brugui  re, A.; Derbr  , S.; Dietsch, J.; Leguy, J.; Rahier, V.; Pottier, Q.; Br  ard, D.; Suor-Cherer, S.; Viault, G.; Le Ray, A.-M.; Saubion, F.; Richomme, P., 2020. MixONat, a Software for the Dereplication of Mixtures Based on ¹³C NMR Spectroscopy. *Anal. Chem.* 92 (13), 8793–8801.
- Brugui  re, A.; Derbr  , S.; Br  ard, D.; Tomi, F.; Nuzillard, J.-M.; Richomme, P., 2021. ¹³C NMR Dereplication Using MixONat Software: A Practical Guide to Decipher Natural Products Mixtures. *Planta Med.* 87 (12/13), 1061–1068.
- Meunier, M.; Br  ard, D.; Awang, K.; Boisard, S.; Guilet, D.; Richomme, P.; Derbr  , S.; Schinkovitz, 2023. A. Matrix Free Laser Desorption Ionization Assisted by ¹³C NMR Dereplication: A Complementary Approach to LC-MS² Based Chemometrics. *Talanta*, 253, 123998.

* Presenting author. E-mail address: severine.derbre@univ-angers.fr (S  verine Derbr  ).

Empowering natural products research with machine learning and expert knowledge integration

Mehdi A. Beniddir^{a,*}

^a Équipe Chimie des Substances Naturelles, Université Paris-Saclay, CNRS, BioCIS, 17 Avenue des Sciences, 91400 Orsay, France

Landmark advances in artificial intelligence approaches and computational omics technologies, particularly in mass spectrometry (MS) and nuclear magnetic resonance (NMR), have recently enhanced the field of natural product (NP) research, putting today's practicing chemists in the enviable position of being able to efficiently speed up the NP discovery process (Mullowney et al., 2023). For some extent, the “art of NP isolation”, shifted from the traditional “grind and find” model to the streamlined hypothesis driven targeting of NPs. This presentation is intended to describe through two selected cornerstone examples the new thinking in natural product chemistry resulting from the emergence of this new generation of sophisticated tools. The first part will be dedicated to the structure validation, of the first trimeric monoterpene indole alkaloids isolated from *Catharanthus roseus*, using a combination of ML-JDP4 (Tsai et al., 2022) approach and empirical computation (Szwarc et al., 2024). The second example will showcase how information-rich multicomponent reactions could be unravelled to extract reactivity patterns in line with Baldwin and Whitehead's manzamine alkaloids biosynthesis scenario (Leblond et al., 2024).

Keywords: metabolomics, structure elucidation, biosynthesis, alkaloids

References

- Leblond, A., A. Nguyen, C. Alcover, K. Leblanc, J.-F. Gallard, D. Joseph, E. Poupon, E., Beniddir, M.A., 2024. Baldwin and Whitehead's manzamine alkaloids biosynthesis hypothesis involves a finely tuned reactivity of acrolein: automated extraction of reactivity patterns from LC-MS² data. *Org. Lett.* 26, 2163-2168.
- Mullowney, M. W., K. R. Duncan, S. S. Elsayed, N. Garg, J. J. van der Hooft, N. I. Martin, D. Meijer, B.R. Terlouw, F. Biermann, K. Blin, J. Durairaj, M. Gorostiola González, E. J. N. Helfrich, F. Huber, S. Leopold-Messer, K. Rajan, T. de Rond, J. A. van Santen, M. Sorokina, M. J. Balunas, M. A. Beniddir, D. A. van Bergeijk, L. M. Carroll, C. M. Clark, D.-A. Clevert, C. A. Dejong, C. Du, S. Ferrinho, F. Grisoni, A. Hofstetter, W. Jespers, O. V. Kalina, S. A. Kautsar, H. Kim, T. F. Leao, J. Masschelein, E. R. Rees, R. Reher, D. Reker, P. Schwaller, M. Segler, M. A. Skinnider, A. S. Walker, E. L. Willighagen, B. Zdrazil, N. Ziemert, R. J. M. Goss, P. Guyomard, A. Volkamer, W. H. Gerwick, H. U. Kim, R. Müller, G. P. van Wezel, G. J. P. van Westen, A. K. H. Hirsch, R. G. Linington, S. L. Robinson and M. H. Medema, 2023. Artificial intelligence for natural product drug discovery. *Nat. Rev. Drug Discov.* 22, 895-916.
- Szwarc, S., A. Jagora, S. Derbré, K. Leblanc, S. Rharrabti, C. Said-Hassane, C. El Kalamouni, J.-F. Gallard, P. Le Pogam and M. A. Beniddir, 2024. Combination of Machine Learning and Empirical Computation for the Structural Validation of Trirosaline, a Natural Trimeric Monoterpene Indole Alkaloid from *Catharanthus roseus*. *Org. Lett.* 26, 274-279.
- Tsai, Y.-H., M. Amichetti, M. M. Zanardi, R. Grimson, A. H. Daranas and A. M. Sarotti, 2022. ML-J-DP4: An Integrated Quantum Mechanics-Machine Learning Approach for Ultrafast NMR Structural Elucidation. *Org. Lett.* 24, 7487-7491.

* Presenting author. E-mail address: mehdi.beniddir@universite-paris-saclay.fr (Mehdi Beniddir).

The evolution of botanical product quality testing

Paula N. Brown*

Natural Health and Food Products Research Group, British Columbia Institute of Technology, Burnaby, British Columbia, Canada

Concerns about product quality and the means through which it is assessed has existed since the earliest development of commerce and trade. Both Pliny the Elder's (23–79 CE) *Naturalis Historia* and Dioscorides' (40–90 CE) *Materia Medica* described products based on their predominant characteristics and the methods through which adulteration could be detected. Despite scientific advances since the original publication of these books, our definition of quality has had to evolve to keep pace with an ever-changing environment. The development and appropriate use of analytical methods is dependent on many factors, including knowledge of analyte(s) and matrix, current state of the art, and nature of the analytical need. Some analytical challenges are more easily addressed, such as determination of target analytes in complex matrices, while reliable analytical solutions for qualitative analyses, such as authentication, remain more elusive. Given the complexity of botanical products and the lack of a direct link between traditional determinants of quality and efficacy, the concept of exploring plant chemistry by a more holistic means is attractive. Comprehensive characterizations can provide pathways to differentiating closely related species, detecting adulteration, and exploring intrinsic and extrinsic influences. Irrespective of the complexity of the analytical challenge or the innovation involved in application of new technologies, methods must be demonstrated to be fit for purpose, rugged, precise, and accurate. This award address will present an overview of experiences, ranging from method development on new analytical platforms to experimental approaches for demonstrating fitness for purpose for qualitative methods.

Keywords: Varro E. Tyler Award, botanical products, quality, authentication, analytical methods, validation, chemometrics

* Presenting author. E-mail address: paula_brown@bcit.ca (Paula N. Brown).

Discovery and engineering of natural product molecules for drug development

David H. Sherman*

University of Michigan, Life Sciences Institute, Departments of Medicinal Chemistry, Chemistry, Microbiology & Immunology, Ann Arbor, Michigan, USA

My talk will focus on various highlights from select research programs initiated and on-going since 1990. One project that has spanned this period relates to modular polyketide synthases, and our development of chemoenzymatic synthesis/cascade biocatalysis to generate new macrolide antibiotic systems. Our work has probed the mechanisms, and engineering of key catalytic domains, and we have linked our ability to generate new molecules with antibiotic activity and ribosome inhibition. A second area of long-standing interest has focused on fungal derived indole alkaloids and assembly of complex multi-ring systems through two classes of enzymes that catalyze intramolecular Diels-Alder reactions. Late-stage C-H oxidation and halogenation has also featured as an important aspect of macrolides and indole alkaloids as the precise modification of their core molecules ultimately enables potent biological activity. More recently, we have developed new semi-synthetic natural product molecules to advance an anti-HIV Nef series of lead compounds that provide a potential path to eradicate the virus. Pathway engineering has been essential for strain optimization to assure a sustainable supply of the two plecomacrolide source materials. I will also highlight studies to identify the endosymbiont producer of ET-743, and current programs to characterize cyanobacterial bloom-derived metabolites in the Great Lakes. Finally, our field collection programs have resulted in a growing marine/terrestrial-derived microbial strain collection and pure natural products/extract library. Centered at the Life Sciences Institute Natural Product Discovery Core, we have benefitted from an institutional Biosciences Initiative that has grown our capabilities significantly, and highlights will be described.

Keywords: natural products, polyketides, indole alkaloids, C-H functionalization, HIV Nef

* Presenting author. E-mail address: davidhs@umich.edu (David H. Sherman).

Session I: Marine organisms, fungi, microbes and plants as chemical factories

Topical Lecture

Efficient discovery of bioactive natural products by targeted metabologenomic approaches

Dong-Chan Oh

Natural Products Research Institute, College of Pharmacy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

Natural products, which were considered as natural gifts, are small molecule material produced by living organisms. In the human history, they have been playing a key role in drug discovery. However, bioactive small molecules with new carbon backbones are still urgently required because drug-resistant diseases have become more widespread and the side effects of clinically-used drugs are problematic. In the past, discovery of new natural products was primarily based on serendipity. However, in the cotemporary genomic era, natural products are recognized as genetically-encoded small molecules. For logical and selective discovery of bioactive natural products, targeted metabologenomic methods were developed utilizing genomic and spectroscopic signatures of specific structural motifs. The application of the metabologenomic methods allowed for the efficient discovery of new macrolactams, piperazic acid-containing peptides, and terminal oxazole-bearing natural products from bacteria. Even though the integration of these metabologenomic techniques in natural product discovery is still in its early stages, it would lead a new paradigm from serendipity to logical discovery and eventually contribute to development of new drug leads.

Keywords: metabologenomic approach, bacteria, natural product, genomics, spectroscopy

References

- Shin, Y.H., Im, J. H., Oh, D.C. 2023. Genomic and spectroscopic signature-based discovery of natural macrolactams. *J. Am. Chem. Soc.* 145, 1886-1896.
- Shin, D., Byun, W. S., Kang, S., Lee, S.K., Oh, D.C. 2023. Targeted and logical discovery of piperazic acid-bearing natural products based on genomic and spectroscopic signatures. *J. Am. Chem. Soc.* 145, 19676-19690.
- Park, J., Shin, Y.H., Hwang, S., Oh, D.C. 2024. Discovery of terminal oxazole-bearing natural products by a targeted metabologenomic approach. *Angew. Chem. Int. Ed.* 2024, accepted.

* Presenting author. E-mail address: dongchanoh@snu.ac.kr (Dong-Chan Oh).

Oral contributions

Unprecedented bicyclic and tricyclic non-canonical terpenoids via synthetic biology platforms: Isolation and structure elucidation of unique C₁₆ and C₁₇ terpenoids

Aikaterini Koutsaviti^a, Yao-Tao Duan^b, Mohamed A. Tammam^a, Maria Harizani^a, Codruta Ignea^b, Vassilios Roussis^a, Yong Zhao^b, Sotirios C. Kampranis^b, Efstathia Ioannou^{a,*}

^a Section of Pharmacognosy and Chemistry of Natural Products, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771 Athens, Greece

^b Biochemical Engineering Group, Plant Biochemistry Section, Department of Plant and Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark

Terpenoids are the most diverse group of specialized metabolites having essential ecological roles, exhibiting diverse biological activities, and finding numerous applications in various industrial sectors. Their biosynthesis is based on the C₅ isoprene building block; thus, almost all terpenoids isolated are based on backbones that contain multiples of five carbon atoms. Nonetheless, a few terpenoids with non-canonical carbon numbers in their backbones have been reported. To explore if these unexpected isoprenoids are limited to the few examples identified so far, gene mining was combined with synthetic biology and natural products chemistry to investigate the extent and diversity of their biosynthesis. After mining available bacterial genomic information for putative non-canonical isoprenoid biosynthetic gene clusters, representative clusters were chosen and studied for their gene organization to reveal complex architectures of methyltransferase and terpene synthase genes. In the next step, the yeast *Saccharomyces cerevisiae* was used as a platform to functionally characterize the selected methyltransferases and terpene synthases. This allowed for the detection of several different non-canonical terpenoids. Copious chromatographic separations enabled the isolation and structure elucidation of a number of C₁₆ (Ignea et al., 2022; Duan et al., 2023) and C₁₇ metabolites, revealing intricate highly strained bi- and tricyclic backbones that are either rare or unprecedented in natural products. Our findings reveal the existence of an extensive class of non-canonical terpenoids in bacteria and prompt further investigation into their biosynthesis and potential pharmacological activity or ecological role, relevant for industrial applications.

Keywords: non-canonical terpenoids, isolation, structure elucidation, synthetic biology

References

- Duan, Y.T., Koutsaviti, A., Harizani, M., Ignea, C., Roussis, V., Zhao, Y., Ioannou, E., Kampranis, S.C., 2023. Widespread biosynthesis of 16-carbon terpenoids in bacteria. *Nat. Chem. Biol.* s41589-023-01445-9.
- Ignea, C., Raadam, M.H., Koutsaviti, A., Zhao, Y., Duan, Y.T., Harizani, M., Miettinen, K., Georgantea, P., Rosenfeldt, M., Viejo-Ledesma, S.E., Petersen, M.A., Bredie, W.L.P., Staerk, D., Roussis, V., Ioannou, E., Kampranis, S.C., 2022. Expanding the terpene biosynthetic code with non-canonical 16 carbon atom building blocks. *Nat. Commun.* 13, 5188.

* Presenting author. E-mail address: eioannou@pharm.uoa.gr (Efstathia Ioannou).

Peptide discovery and development – known and dark matter

Ulf Göransson*, Taj Muhammad, Erik Jacobsson, Luke Robertson, Blazej Slazak, Sunithi Gunasekera

Pharmacognosy, Department of Pharmaceutical Biosciences, Faculty of Pharmacy, Uppsala University, Sweden

Interest in peptides for drug development is surging. This surge is propelled by the genomic revolution, enhanced comprehension of protein function, and advancements in peptide characterization techniques, driven by recent successful precedents. Furthermore, peptides give access to targets non tractable for small molecules, not the least in the modulation of protein-protein interactions. Importantly, techniques to study peptides are mature: genome sequencing to uncover ribosomally produced natural products; bioinformatic tools, e.g. AI, for structural determination and functional annotation; and synthetic and recombinant methods for their production. So, what contribution can pharmacognosy and natural product chemistry make? While bioinformatics and structural predictions are widely accessible, they inherently rely on existing knowledge. In this presentation, I will illustrate how natural product chemistry continue to contribute to the exploration of the unknown—the dark—peptide matter. We have been focusing on plant derived peptides, and our interest has long circled around the cyclotides (Burman et al., 2014). This family of peptides revealed the occurrence of naturally occurring head-to-tail cyclic peptides. I will show how we exploit this feature in the design of scavengers of antibodies in rheumatoid arthritis and Covid 19, and for development of antimicrobials against gram negative bacteria (Gunasekera et al., 2018; 2020; Eriksson et al., 2023). But the remaining question is: are there other families of peptides to be discovered? The answer is “Yes”, albeit if genomes are known or not, and I will demonstrate this with examples from both plants and animals (Jacobsson et al., 2018).

Keywords: peptide chemical biology, peptide discovery and design, antimicrobial, toxins

References

- Burman R, Gunasekera S, Strömstedt AA, Göransson U. 2014. Chemistry and Biology of Cyclotides: Circular Plant Peptides Outside the Box. *J Nat Prod.* 77, 724–736
- Eriksson C, Gunasekera S, Muhammad T, Zhang M, Laurén I, Mangsbo SM, Lord M, Göransson U., 2023. Epitopes displayed in a cyclic peptide scaffold bind SARS-CoV-2 antibodies. *ChemBioChem.* 24:e202300103.
- Gunasekera S, Fernandes-Cerqueira C, Wennmalm S, Wähämaa H, Sommarin Y, Catrina AI, Jakobsson PJ, and Göransson U., 2018. Stabilized cyclic peptides as scavengers of autoantibodies: neutralization of anti-citrullinated protein/peptide antibodies in rheumatoid arthritis. *ACS Chem. Biol.* 13(6):1525-1535.
- Gunasekera S, Muhammad T, Strömstedt AA, Rosengren KJ, Göransson U., 2020. Backbone Cyclization and Dimerization of LL-37-Derived Peptides Enhance Antimicrobial Activity and Proteolytic Stability. *Front Microbiol.* 11, 168.
- Jacobsson J, Andersson HS, Strand M, Peigneur S, Eriksson C, Lodén H, Shariatgorji M, Andrén PE, Lebbe EKM, Rosengren KJ, Tytgat J, Göransson U., 2018. Peptide ion channel toxins from the bootlace worm, the longest animal on earth. *Sci Rep.* 8, 4596.

*Presenting author. E-mail address: ulf.goransson@uu.se (Ulf Göransson).

Co-culture exploration of some selected species of *Fusarium*, *Pestalotiopsis* and *Colletotrichum* in different culture media

Mustapha Ngaski Abubakar, Vitor de Souza Mazucato, Paulo Cezar Vieira*

Faculdade de Ciências Farmacêuticas de Ribeirão Preto, USP - Avenida do Café s/n - 14040-903 - Ribeirão Preto, SP- Brazil

Exploration of the axenic and co-cultures of phytopathogenic species of *Fusarium guttiforme* **F1**, *Pestalotiopsis diospyri* **F2**, *Colletotrichum horii* **F4** and *C. gloeosporioides* **F6** were carried out. The roles of these phytopathogens have been implicated in the infestation faced by protease-rich edible fruits and other agricultural produce. This results in huge economic losses every year (Savary et al., 2012). However, harnessing the co-cultivation of these fungi could be an avenue to understanding their specific needs for the biosynthesis of some specialized metabolites. Hence, informing a novel approach that is capable of affording metabolites with desired/anticipated potentials. The trio co- culture permutations [**CoF126** and **CoF246**] were carried out in rice and PDB (Potato Dextrose Broth) media to ascertain fungi-substrate effects. Although, in PDB fewer metabolites were produced due to **F1** generally, the rice medium favors the fungal exploration in terms of yields and diversity of metabolites. The EtOAc/Acetone extracts were subjected to chromatographic separation/purification and spectroscopic methods for the structure elucidation/identification of compounds. Comparison between ¹H NMR profiles of the co-culture with axenic revealed a significant increase in peaks signal in the former suggesting that certain biosynthetic pathways leading to enhanced production of metabolites were activated. Except in **CoF126**, where fusaric acid from **F1** slightly showed co- expression with the α -pyrone derivatives from **F2**, generally, there was an overexpression/predominance of these α -pyrones followed by chaetiacandin-type polyketide and colletotric acid from **F6**. This implies that the trio-fungal species co-existence was non-mutualistic but of cause beneficial to the search for antagonistic specialized compounds with antimicrobial and protease inhibitory activity.

Keywords: co-culture, *Fusarium guttiforme*, *Pestalotiopsis diospyri*, *Colletotrichum horii*, *Colletotrichum gloeosporioides*, metabolites

References

Savary, S., Ficke, A., Aubertot, J.N., Hollier, C. 2012. Crop losses due to diseases and their implications for global food production losses and food security. Food Security, 4, 519-537.

* Presenting author. E-mail address: pcvieira@fcrp.usp.br (Paulo Cezar Vieira).

Evaluation of biodiverse *Sorghum bicolor* and related genotypes for root exudation, sorgoleone production and potential carbon capture

Leslie A. Weston*, Paul A. Weston, Jesmin Aktar, Nirodha Weeraratne

Gulbali Institute of Agriculture, Water and the Environment, National Life Sciences Hub, Charles Sturt University, Wagga Wagga NSW, Australia

Many sorghum species express allelopathic properties resulting from the biosynthesis and release of long chain hydroquinone metabolites impacting the growth of plants and certain microbes (Weston et al. 2013). Earliest studies on the hydrophobic root exudates from the roots of *Sorghum bicolor* demonstrated their phytotoxicity to broadleaf crop and weed species, suggesting that root exudates play an important role in the growth inhibition of broadleaf weeds. Most recently, multiple long-chain hydroquinones have been identified in exudates, but sorgoleone, 2-hydroxy-5-methoxy-3-[(Z,Z)- 80,110,140-pentadecatriene]-p-benzoquinone, is the major constituent. Interestingly, sorgoleone affects multiple molecular targets (Weston et al., 2013) and has proven to be a potent inhibitor of photosystem II and respiration (Czarnota et al., 2001). Sorgoleone also strongly inhibits the enzyme HPPD (p-hydroxyphenylpyruvate dioxygenase), which is involved in the formation of plastoquinone and also acts as a respiratory inhibitor. Its phytotoxic properties coupled with multiple target sites and the quantity of carbon released in the soil over time offer promise for sorgoleone as a natural product alternative to synthetic herbicides and also as a means to effectively capture carbon release into the surrounding soil environment. The biosynthesis of sorgoleone occurs in sorghum root hairs, and exudate is rapidly released from the root hair in the form of oily yellow droplets (Pan et al. 2016). Sorgoleone has been observed as osmiophilic globules in root hair cells, which are deposited between the plasmalemma and cell wall, and are associated with sorgoleone rhizosecretion (Czarnota et al., 2003). The biosynthetic pathway leading to the formation of sorgoleone was determined in recent years and was it later demonstrated that the biosynthesis of sorgoleone (Fig. 1) involves the production of an alkylresorcinolic intermediate (Weston et al. 2013). *Sorghum bicolor*, *S. Sudanese*, *S. halpense* and *S. leiocladum* have all been found to produce bioactive root exudates containing long chain hydroquinones.




Fig. 1. Structure of sorgoleone 1.

As a result of a Bayer Grants4Ag project, we are employing both metabolomics and genomics approaches to evaluate a biodiverse collection of *S. bicolor* plus other unusual native Australian sorghum genotypes for their ability to a) produce significant quantities of root exudates with a focus on carbon capture, b) generate consistent root hair density and extensive primary roots, c) produce bioactive root exudates containing long chain hydroquinones with bioherbicidal and antimicrobial activity influencing nutrient cycling processes and d) produce exudates with altered composition both qualitatively and quantitatively. We share our research results in the upcoming presentation.

Keywords: sorgoleone, *Sorghum* spp., phytotoxicity, root exudation, PSII inhibitor, mode of action, carbon capture

References

- Weston L.A., Alsaadawi I.S., Baerson S.R. 2013. Sorghum allelopathy – from ecosystem to molecule. J Chem. Ecol. 39: 142–153. Weston L.A., Czarnota M.A. 2001. Activity and persistence of sorgoleone, a long chain hydroquinone produced by *Sorghum bicolor*. J. Crop Production 4, 363–377.
- Yang X., Scheffler B.E., Weston L.A. 2004. SOR1, a gene associated with bioherbicide production in sorghum root hairs. J. Exp. Botany 55: 2251–2259.
- Pan, P. et. al. 2016. *In vivo* assembly of the sorgoleone biosynthetic pathway and its impact on agroinfiltrated leaves of *Nicotiana benthamiana*. New Phytologist (2021) 230: 683–697.

* Presenting author. E-mail address: leweston@csu.edu.au (Leslie A Weston).

‘Two neutrons walk into a bar’. The power of the $^{35}\text{Cl}/^{37}\text{Cl}$ isotope effect in structure elucidation of natural products by NMR and mass spectrometry

Tadeusz F. Molinski^{a,b,*}, Neil Sturchio^c, Xiao Wang^a, Brendan Duggan^b

^a Department of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093, USA

^b Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093, USA

^c Department of Earth Sciences, University of Delaware, 255 Academy Street, Newark, Delaware 19716, USA

In the course of structure elucidation of chlorinated marine natural products (MNPs) from algae (*Rhodophyta*) and sponges (*Porifera*), we required a precise NMR method to locate Cl in the polyhalogenated molecular structures at the ‘nanomole-scale’. Powerful 2D NMR pulse sequences were developed that allowed detection of both $^1J_{\text{CH}}$ and $^nJ_{\text{CH}}$ ($n = 2,3$) cross peaks at C-Cl bonds that were ‘split’ by the $^{35}\text{Cl}/^{37}\text{Cl}$ isotope effect. Here, we show extension of the method, with complementary isotope-rationing mass spectrometry (IRMS) measurements, that reveal $^{35}\text{Cl}/^{37}\text{Cl}$ isotopic fractionation during the biosynthesis of chlorinated MNPs; a pattern that correlates with different modes of biosynthetic halogenation that create C-Cl bonds.




Fig. 1. (a) Mollenyne A. (b) bsHSQC $^{35}\text{Cl}/^{37}\text{Cl}$ isotope shifted C-9-H-9- cross-peak (c) slice through (b) and (c) deconvolution of ^{35}Cl and ^{37}Cl components.

Keywords: marine natural product, chlorine, trichloroleucine, NMR, mass spectrometry, *Dysidea herbacea*, *Porifera*, *Rhodophyta*

References

- Wang, X., Duggan, B.M. Molinski, T.F., 2015. Mollenynes B-E from the Marine Sponge *Spirastrella mollis*. Band- Selective HSQC for Discrimination of Bromo-chloro Regioisomerism in Natural Products. *J. Am. Chem. Soc.* 137, 12343-12351
- Wang, X., Duggan, B.M.; Molinski, T.F., 2017. Ultra-high Resolution Band Selective HSQC for Nanomole-Scale Identification of Chlorine-Substituted ^{13}C in Natural Products Drug Discovery. *Magn. Reson. Chem.* 55, 263-268.
- Sauri, J., Reibarkh, M., Zhang, T., Cohen, R.D., Wang, X., Molinski, T. F., Martin, G. E. Williamson, R.T. 2016. Band- Selective 2D HSQMBC: A Universal Technique for Detection and Measurement of $^{35,37}\text{Cl}$ Isotope Effects for ^{13}C Nuclei. *Org. Lett.* 18, 4786-4789.
- Wang, X, Heraty, L.J. Sturchio, N.C., Molinski, T.F. 2024. $^{35}\text{Cl}/^{37}\text{Cl}$ Fractionation in the Biosynthesis of Chlorinated Marine Natural Products. A Rubric for Interrogating Halogenation Mechanisms, submitted.

* Presenting author. E-mail address: tmolinski@ucsd.edu (Tadeusz F. Molinski).

Charting the chemical content of toxic and lethal French fungi through a multiply annotated molecular networking-based approach

Adrien Jagora^{a,*}, Jean-Michel Bellanger^b, Mehdi A. Beniddir^a, Pierre Le Pogam^a

^a Équipe “Chimie des Substances Naturelles” BioCIS, CNRS, Université Paris-Saclay, 17 Avenue des Sciences, 91400 Orsay, France

^b CEFE, CNRS, Univ Montpellier, EPHE, IRD, INSERM, 1919 Route de Mende, CEDEX 5, 34293 Montpellier, France

The chemical features underlying the toxic effects of mushrooms remain understudied and greatly vary from a species to another. While some syndromes (e.g. phalloidian syndrome) are chemically well characterized (Walton et al., 2018). Others, such as the resinoid, the paxillian or the rhabdomyolytic syndrome, are still poorly documented from a chemical standpoint. Often, the true nature of the toxic species is still debated (Rzymiski et al., 2018). The difficulties raised by mycotoxicologists often stem from a lack of chemical knowledge about macromycetes. A collaboration with the CEFE unit in Montpellier allowed us to raise a collection of more than 3,700 DNA-barcoded entries. This collection will serve as a basis for small-scale extractions of the main fungi responsible for the intoxication syndromes recorded in Europe, prior to acquiring a reference set of LC-MS/MS data. To help us choose the most relevant fungal species to include in a toxicological context, we relied on statistics from the Mycoliste (Bourgeois et al., 2017), that allows French poison control centers to centralize cases of fungal intoxication. We plan to generate molecular networks annotated with a syndromic tag using the GNPS platform (Wang et al., 2016). Merging metabolomics data with toxicological knowledge will enable to visualize the chemical space occupied by fungal toxins and hopefully contribute to answer some long-lasting questions. Besides, disseminating the MS/MS datasets should facilitate the detection of fungal-derived metabolites, either in a context of natural product chemistry targeting new analogues or in a context of analytical toxicology/forensics (Allard et al., 2018).

Keywords: macromycetes, toxins, analytical toxicology, molecular networks

References

- Walton, J., 2018. The Cyclic Peptide Toxins of Amanita and Other Poisonous Mushrooms. Springer International Publishing.
- Rzymiski, P., Klimaszyk, P., 2018. Is the Yellow Knight Mushroom Edible or Not ? A Systematic Review and Critical Viewpoints on the Toxicity of *Tricholoma equestre*. Compr Rev Food Sci Food Saf. 17, 1309-1324.
- Bourgeois et al., 2017. La Mycoliste : un outil d'aide à l'identification des champignons impliqués dans les intoxications humaines en France. Bilan de fonctionnement 2014-2015. Toxicol Anal et Clin. 29, S32-S33.
- Wang et al., 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. Nat. Biotechnol. 34, 828-837.
- Allard et al., 2018. Réseaux moléculaires en toxicologie medico-légale: application à une suspicion d'intoxication à *Tabernanthe iboga*. Toxicol Anal Clin. 30, 182.

* Presenting author. E-mail address: adrien.jagora@universite-paris-saclay.fr (Adrien Jagora).

Discovery of novel biopesticides from an agricultural disease nursery – a proof of concept study

Leah Gauthier^{a,b,*}, Adam Foster^b, Christopher W. Kirby^b, Brian Wagner^a

^a Department of Chemistry, University of Prince Edward Island, 550 University Avenue, Charlottetown, C1A 4P3, Prince Edward Island, Canada

^b Charlottetown Research and Development Center, Agriculture and Agri-Food Canada, 440 University Avenue, C1A 4N6, Charlottetown, Prince Edward Island, Canada

This presentation will discuss results from a proof-of-concept study suggesting that the combination of in-situ isolation using an ichip, and soil subjected to high disease pressure, results in the isolation of novel bacterial species that are promising biopesticide candidates and sources of novel natural product leads. This work revealed that a *Fusarium graminearum* disease nursery in PEI, Canada, which has been maintained at artificially high disease-levels to test resistant crops, harbours a significant number of microbes able to combat fungal pathogens including *F. graminearum*. Half of the isolates recovered were novel strains or species of bacteria and the majority exhibited antagonism against fungal pathogens as summarized in **Fig. 1**. Previously, ichip devices have been used to isolate novel microorganisms under physiological stress such as contaminated soils (Polrot et. al., 2022), or extreme temperatures (Zhao et al., 2023). To our knowledge environments under biotic stress have not been explored in such ways provide compelling evidence that soils experiencing high disease pressure are a valuable source of novel biopesticidal bacteria and antimicrobial compounds. Results from the isolation, genome sequencing, activity assays, and natural products screening experiments of ichip isolated bacteria will be discussed.




Figure 1. Plate antagonism assay isolates recovered from a *F. graminearum* nursery using an ichip where half of the isolates are novel strains or species, and the majority exhibit antagonism against *F. graminearum*.

Keywords: *Fusarium graminearum*, ichip, biocontrol, biopesticides, soil microbiota

References

- Polrot, A. Kirby, J. R. Olorunniji, F. J., Birkett, J. W., Sharples, G. P., 2022. iChip increases the success of cultivation of TBT-resistant and TBT-degrading bacteria from estuarine sediment. *World J Microbiol Biotechnol.* 38, 180.
- Zhao, J., Shakir, Y., Deng, Y., Zhang, Y., 2023. Use of modified ichip for the cultivation of thermo-tolerant microorganisms from the hot spring. *BMC Microbiol.* 23,56.

* Presenting author. E-mail address: lkgauthier@upe.ca (Leah Gauthier).

Deep ocean biodiscovery – diving into abyss

Darren Scobie^a, Artem Fetter^b, Adrian Glover^c, Muriel Rabone^c, Tammy Horton^d, Gagan Preet^e, Marcel Jaspars^e, Rainer Ebel^e, Joape Ginigini^f, Jon Copley^g, Katherine R. Duncan^{b,*}

^a University of Strathclyde, Strathclyde Institute of Pharmacy and Biomedical Sciences, Glasgow, UK

^b University of Newcastle, Newcastle University Biosciences Institute, Newcastle Upon Tyne, UK

^c Natural History Museum, London, UK

^d National Oceanography Centre, UK

^e University of Aberdeen, Department of Chemistry, Aberdeen, UK

^f University of South Pacific, Fiji

^g University of Southampton, Southampton, UK

The extreme variety of deep-sea habitats, encompassing ancient abyssal plains some of which date to over 180mya, to high-temperature hydrothermal vents, makes the deep-sea the perfect evolutionary test-lab for natural product research. Often, we think of deep-sea as void of life, but a recent study has uncovered 5,000 new species in Pacific deep-sea mining regions (Rabone et al., 2023) and we know that understanding diversity-ecosystem function is of primary importance in the face of biodiversity loss. These ecosystems have remained largely unexplored in terms of biodiversity and biomedical potential, yet over recent years they have gained significant interest due to the potential of deep-sea mining for metals such as cobalt and nickel. Here we aim to profile the biodiversity and biomedical (antibiotic) potential of deep-sea microorganisms. Microbial drug discovery in the ‘omics era relies on three key datasets, biosynthetic, chemical, and biological (activity). Yet, to integrate and interrogate these large and complex datasets remains a challenge and results in the low-throughput prioritization of only a few strains based on observed antibiotic activity. Despite this wealth of genomic and metabolomic data, linking metabolites to the BGC responsible for their production and to observed bioactivity is limited, slow (manual) and challenging. Here, approaches to combine data sets consisting of bacterial genomes (and their predicted BGCs), the chemical products of these same strains and their bioactivity profiles will be discussed. Several datasets of Actinobacteria genomes have been mined for BGCs and these strains cultured to generate metabolite extracts for comparative metabolomics (high resolution tandem mass spectrometry / molecular networking) and antibiotic screening (against a panel of clinically relevant pathogens). Driven by biological questions, tools have been developed to establish patterns across strains and learn relationships between BGC, spectral features and bioactivity – here we will discuss experimental data considerations and how ‘omics information is informing biodiscovery from the deep-sea.

Keywords: *Streptomyces*, antibiotics, metabolomics, genomics, deep-sea

References

Rabone, M. et al. 2023. How many metazoan species live in the world’s largest mineral exploration region? *Curr. Bio.* 33, P2383-2396.

* Presenting author. E-mail address: katherine.duncan@newcastle.ac.uk (Katherine R. Duncan).

Specialized metabolites of the herptile gut fungus, *Basidiobolus*

George F. Neuhaus^{a,*}, Ian A. Trautman^b, Richard M. Tehan^{a,c}, Lluvia Vargas-Gastélum^b, Alexander S. Romer^d, N. Reed Alexander^d, Marjan Ghotbi^e, Kylie C. Moe^d, Leila Shadmani^f, Javier F. Tabima^g, Jason E. Stajich^f, Donald M. Walker^d, Joseph W. Spatafora^b, Kerry L. McPhail^a

^a Department of Pharmaceutical Sciences, Oregon State University, Corvallis, OR, USA

^b Department of Botany and Plant Pathology, Oregon State University, Corvallis, OR, USA

^c School of Arts and Sciences, Utica University, Utica, NY, USA

^d Department of Biology, Middle Tennessee State University, Murfreesboro, TN, USA

^e GEOMAR-Helmholtz Centre for Ocean Research, The Research Division Marine Ecology, Kiel, Germany

^f Department of Microbiology and Plant Pathology, University of California, Riverside, CA, USA

^g Department of Biology, Clark University, Worcester, MA, USA

The zygomycete fungus *Basidiobolus* is often a prominent member in the herptile (amphibians and reptiles) gut microbiome. Draft genomes of *Basidiobolus* show evidence of horizontal gene transfer (HGT) of nonribosomal peptide synthetases (NRPS), some of which are predicted to originate in co-occurring gut bacteria (Tabima et al., 2020). It is suspected that the peptidic metabolites encoded play a role in modulation of the herptile gut microbiome. Here, untargeted LCMS²-based metabolomics using computational tools (GNPS, Sirius5, and CANOPUS) for structural class prediction revealed that 10 unique *Basidiobolus* isolates (derived from field collected herptile fecal pellets) produce cyclic peptides belonging to the same molecular family in laboratory culture (Vargas-Gastélum et al., 2024). An analysis of the antimicrobial activity of extracts from a gecko isolate of *Basidiobolus*, using the Bioactive Molecular Networks Project protocol, highlighted 16 mass features that correlated with the observed antimicrobial activity against a panel of human pathogens. Analysis of tandem mass spectra assigned these mass features as peptides and depsipeptides and isolation and structure elucidation efforts are currently underway. Furthermore, a laboratory model was developed that can establish stable infections of specific *Basidiobolus* isolates in adult wood frogs (*Lithobates sylvaticus*). This allows monitoring effects *Basidiobolus* has on microbiome community structure and metabolism. These experiments support the hypothesis that cyclic peptides from *Basidiobolus* can influence microbial growth and lay the groundwork for further study of the implications *Basidiobolus* has on herptile health.

Keywords: *Basidiobolus*, herptile microbiome, cyclicpeptides, horizontal gene transfer, antimicrobial

References

- Tabima, J.F., Trautman, I.A., Chang, Y., Wang, Y., Mondo, S., Kuo, A., Salamov, A., Grigoriev, I.V., Stajich, J.E., Spatafora, J.W., 2020. Phylogenomic Analyses of Non-Dikarya Fungi Supports Horizontal Gene Transfer Driving Diversification of Secondary Metabolism in the Amphibian Gastrointestinal Symbiont, *Basidiobolus*. *G3*, 10, 3417–3433. Vargas-Gastélum, L., Romer, A.S., Ghotbi M., Dallas J.W., Reed, A.N., Moe, K.C., McPhail, K.L., Neuhaus, G.F., Shadmani, L., Spatafora, J.W., Stajich, J.E., Tabima, J.F.; Walker, D.M., 2024. Herptile Gut Microbiomes: A Natural System to Study Multi-Kingdom Interactions between Filamentous Fungi and Bacteria. *mSphere*, e00475–23.

* Presenting author. E-mail address: neuhausg@oregonstate.edu (George F. Neuhaus).

Session II: Natural products in the era of emerging infectious diseases

Topical Lecture

From microbes to medicine: leveraging nature's arsenal for innovative anti-infectives

Rolf Müller

Helmholtz Institute for Pharmaceutical Research Saarland (HIPS), Saarbrücken, Germany

The global rise of antimicrobial resistance, mainly due to the mis- and overuse of antibiotics, is one of the most pressing issues of our time. To counteract this development, novel resistance-breaking antibiotics are urgently needed. Historically, the vast majority of antibiotics have been derived from microbial natural products. As compared to traditional bacterial producers, such as actinomycetes and bacilli, myxobacteria have been studied less extensively and thus harbor a large potential for the discovery of entirely new natural product scaffolds exhibiting promising bioactivities. Comparisons of myxobacterial metabolite profiles with the number of underlying biosynthetic gene clusters encoded in their large genomes show, that many compounds still remain unknown. Further, recent studies indicate that the order of myxobacteria likely comprises many more biodiverse representatives than previously assumed. According to metagenomics analyses, myxobacteria (including many underexplored representatives) are highly abundant in the soil microbiome, where they play a crucial role in soil nutrient and carbon cycling. Taken together with our recent genomic analyses, these findings suggest that the biosynthetic potential of myxobacteria is a long way from being exhausted.

We recently demonstrated that chemical diversity correlates with taxonomic distance in myxobacteria. Accordingly, we are more likely to isolate novel compound classes from strains which are phylogenetically distant from previously characterized strains as compared to closely related strains. This knowledge is applied to prioritize strains for natural product discovery, thus increasing the chance of discovering compound classes with yet unknown chemical structures and biological activities.

I will discuss recent results from our laboratory regarding the identification of novel bioactive NPs from myxobacteria based on different approaches, and show recent advances in the preclinical development of selected compound classes.

* Presenting author. E-mail address: rolf.mueller@helmholtz-hips.de (Rolf Müller)

Oral contributions

Observations and insights into the prolific and varied metabolites of the fascinating plant genus *Dalea* (Fabaceae)

Gil Belofsky^{a,*}, Hyojin Ahn^a, Caroline Cruz^a, Ericka DuBore^a, Ryan Galperin^a, Trevor Schultz^a, Dominique Wilcox^a, Maxwell Zapata^a, Christine E. Salomon^b, P. Clint Spiegel^c

^a Department of Chemistry, Central Washington University, Ellensburg, Washington 98926, United States

^b Center for Drug Design, University of Minnesota, Minneapolis, Minnesota, 55455, United States ^c Department of Chemistry, Western Washington University, Bellingham, Washington 98225, United States

Work on the phytochemistry of *Dalea* spp. continues to afford new phenolic metabolites with no sign of diminishing returns after two decades. Metabolites of *D. jamesii* and *D. nana* (Belofsky et al., 2023 and 2024), provide insights into in vitro structure-activity relationships of isoflavonoids toward methicillin-resistant *Staphylococcus aureus*, and other human pathogens. The prenylated isoflavans isolated exhibit MICs ≤ 9 mg/mL (aq. DMSO) that appear correlated to the number and type of these lipophilic groups (e.g., **1** and **2**, Fig. 1).

Prior work on pawhuskins A-C (Belofsky et al. 2004), k-opioid antagonists from *D. purpurea*, was unique among all *Dalea* spp. examined until current work on *D. candida* var. *oligophylla* afforded a new member of this class, pawhuskin D (**3**). The continued work on extractives of this species will be presented.




Fig. 1. Structures of antimicrobial isoflavans **1** and **2** and of pawhuskin D **3**.

Cumulative results clearly support the ecological observation that more structurally diverse metabolites, with more potent antimicrobial activities, are concentrated in the roots rather than aerial portions of *Dalea* spp. Details will be presented of our exclusive use of open-column chromatography, that makes such research accessible to students at a primarily undergraduate institution.

Keywords: *Dalea* spp., Fabaceae, isoflavanoid, pawhuskin, MRSA

References

- Belofsky, G., Ahn, H., Zapata, M., Wilcox, D., Salomon, C.E., Spiegel, P.C., 2023. Antimicrobial Isoflavans and Other Metabolites of *Dalea jamesii*. *Planta med.*, 89, 1-10.
- Belofsky, G., Cruz, C., Schultz, T., Zapata, M., Wilcox, D., Wasmund, B., Salomon, C.E.; Spiegel, P.C., 2024. Antimicrobial Isoflavans and Other Metabolites of *Dalea nana*. *Phytochem.* Submitted 1/4/24.
- Belofsky, G., French, A.F., Wallace, D.R., Dodson, S.L., 2004. New Geranyl Stilbenes from *Dalea purpurea* with *in vitro* Opioid Receptor Affinity. *J. Nat. Prod.*, 67, 26-30.

* Presenting author. E-mail address: belofskyg@cwu.edu (Gil Belofsky)

Target identification of 3-(phenethylamino)demethyl(oxy)aaptamine (PDOA) as an anti-dormant Mycobacterial substance.

Michihisa Matsuzaki^a, Atsushi Kimishima^a, Yusuke Shimada^a, Hiroaki Hase^a, Yuji Sumii^a, Kazutake Tsujikawa^a, Naoyuki Kotoku^{a,b}, Andi Setiawan^c, Masayoshi Arai^{a,*}

^a Graduate School of Pharmaceutical Sciences, Osaka University, Osaka, 565-0871, Japan

^b College of Pharmaceutical Sciences, Ritsumeikan University, Shiga, 525-8577, Japan

^c Faculty of Sciences, Lampung University, Bandar Lampung, 35145, Indonesia

Tuberculosis is an infectious disease caused by infection with *Mycobacterium tuberculosis*, resulting in approximately 1.3 million deaths annually. The presence of dormant *M. tuberculosis*, which is tolerance to anti-tuberculosis drugs, is considered a reason for requiring prolonged chemotherapy through multi-drug combinations. There is a need to discover new medicinal seeds effective against dormant *M. tuberculosis*. 3-(phenethylamino)demethyl(oxy)aaptamine (PDOA, **1**), isolated from Indonesian marine sponge *Aaptos* sp., was effective against dormant *Mycobacterium* species (Sumii et al., 2020). However, its target molecule is not yet identified. In this study, we synthesized the probe molecule of **1** and validated its utility as a probe molecule, as well as attempted to identify binding protein from the bacterial cell lysate. From the result of structure-activity relationships study using synthetic analogs of **1** (Sumii et al., 2023), PDOA probe (**2**) was designed and synthesized (Fig. 1). The functional evaluation of **2** was conducted using BSA. As a result, **2** bound to BSA only when subjected to UV irradiation. The Huisgen reaction between **2** and a biotin-labeled PEG linker with an azide group proceeded in an aqueous solvent. Furthermore, **2** having biotin was found to be capable of pulling down BSA using streptavidin beads. Therefore, we further attempted to identify the binding protein of **1** using **2** from cell lysate of *M. bovis* BCG. As a result, we succeeded in finding some proteins on SDS-PAGE gel that were not detected in the absence of **2** or when **1** was added to the reaction system as a competitor, but were selectively detected in the presence of **2**.




Fig. 1. Structures of PDOA (**1**) and PDOA probe (**2**)

Keywords: *Aaptos* sp., PDOA, anti-microbial, tuberculosis, target molecule

References

- Y. Sumii, N. Kotoku, C. Han, K. Kamiya, A. Setiawan, C. Vilchèze, W. R. Jacobs Jr., M. Arai, 2020. 3- (Phenethylamino)demethyl(oxy)aaptamine as an anti-dormant mycobacterial substance: Isolation, evaluation and total synthesis. *Tetrahedron Lett.*, 61(22), 151924.
- Y. Sumii, K. Kamiya, T. Nakamura, K. Tanaka, T. Kaji, N. Kotoku, M. Arai, 2023. Structure activity relationship study of 3-(Phenethylamino) demethyl(oxy)aaptamine as an anti-dormant mycobacterial substance to create probe molecule for detecting target protein. *Mar. Drugs*, 20(2), 98.

* Presenting author. E-mail address: araim@phs.osaka-u.ac. (Masayoshi Arai)

Revealing anti-SARS-CoV-2 natural products from traditional remedies using molecular networking

Andreas Wasilewicz^{a,*}, Denisa Bojkova^b, Mehdi A. Beniddir^c, Jindrich Cinatl Jr. ^b, Holger F. Rabenau^b, Ulrike Grienke^a, Judith M. Rollinger^a, Benjamin Kirchweger^{a,d}

^a University of Vienna, Department of Pharmaceutical Sciences, Division of Pharmacognosy, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^b University Hospital Frankfurt, Institute of Medical Virology, Paul-Ehrlich-Straße 40, 60596 Frankfurt am Main, Germany

^c Université Paris-Saclay, Équipe Chimie des substances naturelles, BioCIS, CNRS, 17, Avenue des Sciences, 91400, Orsay, France

^d Ludwig-Maximilians-Universität München, Department of Pharmacy, Pharmaceutical Biology, Butenandtstraße 5-13, 81377 Munich, Germany

Natural remedies have a long-standing tradition in the treatment of respiratory diseases. This ethnopharmacological knowledge provides a remarkable feature of natural product research, forshadowing potential new bioactivities. In this study, 179 extracts derived from 154 natural remedies traditionally used against respiratory infections and inflammation were tested for their anti- SARS-CoV-2 activity in a Caco-2 cell model (Wasilewicz et al., 2024). 13 extracts exhibited distinct anti-SARS-CoV-2 activity with IC₅₀ values below 50 µg/mL; notably, five extracts demonstrated IC₅₀ values below 5 µg/mL. In the search for constituents within the complex and highly biodiverse extract set, non-toxic anti-SARS-CoV-2 active extracts and some inactive extracts were dereplicated using a molecular network (MN) approach. For this purpose, MS² spectra of both active and inactive extracts were recorded and used for the generation of a feature-based MN (Nothias et al., 2020). The obtained MN analysis enabled the determination of compounds classes characteristic of active extracts. Thereby, in particular some flavonoid glycosides, triterpene saponins, sesquiterpenes and acetogenins were unravelled as putative bioactive metabolites from the screened extracts.

To summarize, the herein applied bioactivity-featured MN approach allowed to identify the most promising constituents from the hit extracts, thereby providing crucial chemical insights into known as well as novel structural features with anti-SARS-CoV-2 activity. Thus, this work serves as a basis for the targeted isolation of novel or known natural compounds with yet unexplored anti-SARS-CoV- 2 activities.

Keywords: SARS-CoV-2, ethnopharmacology, molecular networking

References

Nothias, L.F., Petras, D., Schmid, R., Dührkop, K., Rainer, J., Sarvepalli, A., Prostyuk, I., Ernst, M., Tsugawa, H., Fleischauer, M., et al. 2020. Feature-based molecular networking in the GNPS analysis environment. *Nat. Methods*. 17, 905-908.

Wasilewicz, A., Bojkova, D., Beniddir, M.A., Cinatl Jr, J., Rabenau, H.F., Grienke, U., Rollinger, J.M., Kirchweger, B. 2024. Molecular networking unveils anti-SARS-CoV-2 constituents from traditionally used remedies. *J. Ethnopharmacol.* 319, 117206.

Acknowledgments

Funding: FWF project P 34028

* Presenting author. E-mail address: andreas.wasilewicz@univie.ac.at (Andreas Wasilewicz)

Identification and SAR studies of indole-functionalized polyamines as antibiotic enhancers

Melissa M. Cadelis^{a,b,*}, Steven A. Li^b, Kenneth Sue^b, Tim Liu^b, Marie-Lise Bourguet-Kondracki^c, Marine Blanchet^d, Jean-Michel Brunel^d, Brent R. Copp^b

^a School of Medical Sciences, The University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

^b School of Chemical Sciences, The University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

^c Laboratoire Molécules de Communication et Adaptation des Micro-organismes, UMR 7245 CNRS, Muséum National d'Histoire Naturelle, 57 rue Cuvier (C.P. 54), 75005 Paris, France

^d Aix-Marseille Université, Centre de Recherches en Cancérologie de Marseille (CRCM), CNRS, UMR7258, 13385 Marseille, France

Antibiotics have long been pivotal in modern medicine, saving countless lives by effectively combating infectious diseases (Lewis, 2020). However, the rise of drug-resistant strains of bacteria coupled with a dearth of new antibiotic development presents a pressing need for novel approaches (Singh, 2014). One promising strategy involves identifying compounds that can augment the efficacy of existing antibiotics, particularly those losing effectiveness against resistant bacteria (Douafer et al., 2019).

Initial screening of compounds sourced from our in-house collection of purified marine natural products and their synthetic derivatives, identified a promising lead: an indole-functionalized polyamine (Fig. 1). This compound not only restored the activity of the antibiotic doxycycline against the drug-resistant Gram-negative bacterium *Pseudomonas aeruginosa* but also demonstrated intrinsic antibacterial properties. Subsequent Structure-Activity Relationship (SAR) studies yielded a range of novel analogues, many of which displayed potent enhancing effects on both doxycycline and erythromycin against *P. aeruginosa* and *Escherichia coli*. The forthcoming presentation will detail the results of these studies, shedding light on the mechanisms of action of the most potent compounds and their capacity to augment the efficacy of diverse antibiotics against drug-resistant bacterial strains.




Fig. 1. Representative structure of indole-functionalized polyamine

Keywords: Indole, polyamine, antimicrobial, adjuvant, SAR

References

- Douafer, H., Andrieu, V., Phanstiel, O., Brunel, J.M., 2019. Antibiotic Adjuvants: Make Antibiotics Great Again! J. Med. Chem. 62, 8665–8681.
- Lewis, K., 2020. The Science of Antibiotic Discovery. Cell. 181, 29–45.
- Singh, S.B., 2014. Confronting the Challenges of Discovery of Novel Antibacterial Agents. Bioorg. Med. Chem. Lett. 24, 3683–3689.

* Presenting author. E-mail address: m.cadelis@auckland.ac.nz (Melissa M. Cadelis)

Hope in hop

Gabriel Lefèvre^a, Anne-Sophie Paguet^a, Augustin Bildstein^a, Laetitia Bocquet^a, Jennifer Samaillie^a, Sophie Moureu^a, Imelda Raczekiewicz^b, Lowiese Desmarets^b, Charline Camuzet^b, Alexandra Hubert^c, Lucie Borkova^c, Justine Jacquin^a, Sevser Sahpaz^a, Caroline Rambaud^a, Christel Neut^a, Marie-Laure Fauconnier^a, Roland Molinié^a, Jean-Xavier Fontaine^a, Cécile Palaric^a, Ali Siah^a, Jérôme Muchembled^a, Mathilde Vandenberghe^a, Sylvie Chollet^a, Jean Dubuisson^b, Sandrine Belouzard^b, Mourad Elhabiri^c, Karin Séron^b, Céline Rivière^{a,*}

^a UMRt 1158 BioEcoAgro, Univ. Lille, JUNIA, UPJV, Univ. Liège, INRAE, Univ. Artois, Univ. Littoral Cote D'Opale, ICV—Institut Charles Viollette, F59650 Villeneuve D'Ascq, France

^b Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 – UMR9017 – Center for Infection and Immunity of Lille (CIIL), France

^c CNRS – Univ. Strasbourg – Univ. de Haute-Alsace, LIMA, CNRS UMR 7042, Equipe Chimie Bioorganique et Médicinale, ECPM, 25 Rue Becquerel, 67200 Strasbourg, France

Hop (*Humulus lupulus* L., Cannabaceae), a plant of economic and cultural interest, is cultivated for its female inflorescences, called hops or cones, used by the brewing industry and for their medicinal properties. Hops produce both an essential oil, mainly represented by non-oxygenated monoterpenes and sesquiterpenes, and original prenylated compounds, including chalcones and acylphloroglucinols (α and β acids) (Bocquet et al., 2018). The composition of the essential oil and the α acid content are particularly sought after by brewers. The evolution of the beer market, in particular the growing interest in craft beers with a trend toward strongly hopped beers, is impacting hop production and varietal selection (Paguet et al., 2022). Hops are also studied for their many pharmacological activities, including sedative, estrogenic, antioxidant, anti-inflammatory and antimicrobial activities (Bocquet et al., 2018). In this context, we focused our research on the chemical and genetic diversity of wild hops harvested in the North of France in order to underline their interest both for potential varietal selection and for the identification of new bioactive specialized metabolites. This presentation will also highlight the various work carried out by our teams on the antimicrobial potential of hop metabolites and their co-products for human and plant health.

Keywords: *Humulus lupulus* L., prenylated phenolic compounds, antimicrobial agents, co-products, wild hops

References

- Laetitia Bocquet, Sevser Sahpaz, Jean-Louis Hilbert, Caroline Rambaud, Céline Rivière, 2018. *Humulus lupulus* L., a very popular beer ingredient and medicinal plant: overview of its phytochemistry, its bioactivity, and its biotechnology. *Phytochem. Rev.* 17, 1047-1090.
- Anne-Sophie Paguet, Ali Siah, Gabriel Lefèvre, Sevser Sahpaz, Céline Rivière, 2022. Agronomic, genetic and chemical tools for hop cultivation and breeding. *Phytochem Rev.* 21, 667-708.

Acknowledgments

We thank the Hauts-de-France region, the European Union, the University of Lille and CNRS Innovation (CPER Alibiotech and BiHauts Eco de France, ANR Flash, FEADER, BQI and CNRS prematuration programs). Many thanks also to the different funders of PhD theses of Laetitia Bocquet (2015-2018), Anne-Sophie Paguet (2019-2022) and Augustin Bildstein (2022-2025): Hauts-de France Region, Univ. Lille, French Foundation of Brewery and Maltery, Junia- ISA, the Inter-branch organization of hops “Interhoublon” and “Syndicat des Brasseurs des Hauts-de-France”.

* Presenting author. E-mail address: celine.riviere@univ-lille.fr (Céline Rivière)

Hops (*Humulus lupulus* L.) compounds against two tropical infectious diseases: *Chikungunya* and *Oropouche* virus (Bunyavirales)

Tsvetelina Mandova^{a*}, Marielena Vogel Saivish^{b,c}, Gabriela de Lima Menezes^d, Umberto Laino Fulco^d, Fernando Batista Da Costa^e, Maurício Lacerda Nogueira^f

^a Gilson Purification, 22 rue Bourseul, 56890 Saint Avé, France

^b Research Laboratories in Virology, Department of Dermatological, Infectious, and Parasitic Diseases, Faculty of Medicine of São José do Rio Preto, São José do Rio Preto, SP, Brazil;

^c Brazilian Biosciences National Laboratory, National Center for Research in Energy and Materials (CNPEM), Campinas 13083-100, SP, Brazil;

^d Bioinformatics Multidisciplinary Environment, Graduate Program in Bioinformatics, Federal University of Rio Grande do Norte, Natal 59078-400, RN, Brazil;

^e AsterBioChem Research Team, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto 14040-020, SP, Brazil

^f Department of Pathology, University of Texas Medical Branch, Galveston, TX 77555, USA

Humulus lupulus L. (Cannabaceae), commonly named hop, is widely grown around the world for its use in the brewing industry. Its female inflorescences (hops) are particularly prized by brewers because they produce some secondary metabolites that confer bitterness, aromas, and antiseptic properties to the beer. These sought-after metabolites as prenylated phenolic compounds, mainly acylphloroglucinols (bitter acids) have shown numerous biological activities. We investigated whether purified hop components α - and β - acids and prenylflavonoids (xanthohumol) exhibit antiviral activity against Chikungunya virus (CHIKV) and Oropouche virus (OROV), two arboviruses that cause serious health problems in tropical and subtropical regions. The bioactive compounds were purified by the countercurrent separation method. For CHIKV, the antiviral activity was determined by the plaque reduction test. Almost all hop compounds demonstrated a promising post-treatment viral inhibition. β -acids in the highest tested concentration of 125 $\mu\text{g/mL}$ expressed the strongest virucidal activity ($\text{EC}_{50} = 15.21 \mu\text{g/mL}$), in a drug-addition experiment on Vero cells. Hypotheses for the mechanism of action concerning protein kinase C (PKC) transduction cascades were proposed. Same bioactive compounds from hops were also tested against OROV. The evaluation of the inhibitory potential on the viral cycle of OROV was performed through two complementary approaches. The first approach applied cell-based assay post-inoculation experiments to explore the inhibitory potential on the latest steps of the viral cycle, such as genome translation, replication, virion assembly, and virion release from the cells. The second part covered *in silico* methods evaluating the ability of these compounds to inhibit the activity of the endonuclease domain, which is essential for the transcription, binding, and cleaving of RNA. In conclusion, the β -acids beta acids showed strongest inhibitory potential in the post-treatment assay ($\text{EC}_{50} = 26.7 \mu\text{g/mL}$). The results observed from the docking and MM/PBSA analysis confirmed the affinity and among the three tested ligands, Lys92 and Arg33 exhibited the highest affinity with the protein.

Keywords: tropical infectious diseases, *Chikungunya*, *Oropouche*, hops compounds, countercurrent chromatography

References

Bocquet, L.; Sahpaz, S.; Hilbert, J.L.; Rambaud, C.; Rivière, C. 2018. *Humulus lupulus* L., a very popular beer ingredient and medicinal plant: Overview of its phytochemistry, its bioactivity, and its biotechnology. *Phytochem. Rev.*, 17, 1047– 1090.

* Presenting author. E-mail address: tmandova@gilson.com (Tsvetelina Mandova)

The phototherapeutic potential of fungal pigments

Bianka Siewert^{a,*}, Fabian Hammerle^a, Johannes Fiala^{a,b}, Arianna Serra, Anja Höck^a, Lesley Huymann^{a,b}, Yurii Husiev^c, Pamela Vrabl^b, Tom May^d, Michaela Lackner^e, Sylvestre Bonnet^c, Ursula Peintner^b

^a Department of Pharmacognosy, Institute of Pharmacy, University of Innsbruck, Innsbruck, Austria

^b Institute of Microbiology, University of Innsbruck, Innsbruck, Austria

^c Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333CC Leiden, The Netherlands

^d Royal Botanic Gardens Victoria, Melbourne, Australia

^e Hygiene and Medical Microbiology, Medical University Innsbruck, Innsbruck, Austria

Mushrooms are an underestimated source of natural photosensitizers, especially those that produce polyketide pigments. Since fungi share their ecological niche with other microorganisms, we hypothesized that some fungal pigments are photoantimicrobials. To test our hypothesis, fungal species of the genus *Cortinarius* producing colored fruiting bodies were explored (e.g., *Cortinarius sanguineus* or *C. holoxanthus*). The extracts were assessed among others against the yeast *Candida albicans* as well as the bacteria *Escherichia coli* and *Staphylococcus aureus* under different irradiation conditions ($\lambda = 428, 478, \text{ or } 528 \text{ nm}$, $H = 30 \text{ J cm}^{-2}$) and in the dark. A significant photoantimicrobial effect was discovered for several extracts ($c < 25 \mu\text{g/mL}$). Targeted isolation of the responsible pigments from *C. sanguineus* led to the identification of two new potent photoantimicrobials. One, dermocybin, was active against *S. aureus* and *C. albicans* under green light irradiation, while it was ineffective in the dark and non-toxic to cell lines. The other, emodin, was also active against *E. coli* in the low micromolar range. Inspired by the activity of these anthraquinones, we tested the potential of related polyhydroxy anthraquinones and showed that besides aloe-emodin and chrysophanol, trihydroxy anthraquinones are also potent photoantimicrobials.

In sum, here we will show the remarkable potential of fungal pigments against the gram-positive and gram-negative bacteria as well as several yeasts, causing unpleasant skin infections in humans.

Keywords: dermocybin *Cortinarii*, photoantimicrobials, anthraquinones

References

Fiala, J., Hammerle, F., Höck, A., Huymann, L., Vrabl, P., Husiev, Y., Bonnet, S., Peintner, U., and Siewert, B., 2023. J. Nat. Prod. 86 (10), 2247-2257, DOI: 10.1021/acs.jnatprod.2c01157

* Presenting author. E-mail address: bianka.siewert@uibk.ac.at (Bianka Siewert)

Villosane A, an allene containing bisabolane sesquiterpene with potent antiplasmodial activity isolated from cultures of the marine derived basidiomycete *Halocyphina villosa*

Frederick Boye Annang^a, Guiomar Pérez-Moreno^b, Cristina Bosch-Navarrete^b, Victor González- Menéndez^a, Caridad Díaz^a, Maria C. Ramos^a, Ignacio Pérez-Victoria^a, Jesús Martín^a, Thomas A. Mackenzie^a, Duane Choquesillo-Lasarte^c, Luis M. Ruiz-Pérez^b, Olga Genilloud^a, Dolores González-Pacanowska^b, Francisca Vicente^a, Fernando Reyes^{a,*}

^a Fundación MEDINA, Avda. del Conocimiento 34, 18016, Armilla, Granada, Spain

^b Instituto de Parasitología y Biomedicina "López-Neyra", Consejo Superior de Investigaciones Científicas (CSIC) Avda. del Conocimiento s/n, 18016, Armilla, Granada, Spain

^c Laboratorio de Estudios Cristalográficos, IACT, CSIC, University of Granada, Avenida de las Palmeras 4, 18100, Armilla, Granada, Spain

The chemotherapeutic armamentarium to fight parasitic diseases such as malaria and Chagas disease needs novel bioactive molecules with potential for development into effective drugs. Sustained efforts in antiparasitic high throughput screening of a collection of microbial extracts (Annang et al. 2015; Pérez-Moreno et al. 2016) coupled with bioassay-guided compound isolation led to the discovery of a new family of potent antiparasitic bisabolane sesquiterpenoids containing a hydroxamide functionality, villosanes A-C, from an extract of the marine derived basidiomycete *Halocyphina villosa* CF-090066. (+)-ESI-TOF mass spectrometry, nuclear magnetic resonance spectroscopy and X-ray diffraction were used to establish their structures and absolute configurations.

Villosane A, featuring a rare allene moiety in its structure was the most potent and most selective of the three compounds, exhibiting EC₅₀ values of 0.0055 and 6.2 µM against *Plasmodium falciparum* 3D7 and *Trypanosoma cruzi* Tulahuen C4 parasites, respectively. Villosanes B and C exhibited weaker EC₅₀ values of 0.062 and 4.83 µM, respectively, against *P. falciparum* and were both inactive against *T. cruzi* at 25 µM. Preclinical profiling of villosane A revealed non-inhibition of hERG, Nav

1.4 and Cav 1.5 ion channels at 50 µM, adequate in vitro metabolic stability, weak potential for drug- drug interaction and no in vivo toxicity at a dosage of 50 mg/kg in mice. Hence, villosane A, the major component of the extract, presents a promising potential for development into an effective antimalaria drug.

Data on the production, isolation, structures and antiparasitic properties of the villosane family of natural products will be presented in this communication.

Keywords: *Halocyphina villosa*, bisabolene sesquiterpenoids, allenes, bioassay-guided isolation, antiparasitic

References

- Annang, F. et al., 2015. High throughput screening platform for Natural Product-based drug discovery against three Neglected Tropical Diseases: human African trypanosomiasis, leishmaniasis and Chagas disease. *J. Biomol. Screen.* 20, 82-91.
- Pérez-Moreno, G., 2016. Discovery of New Compounds Active against *Plasmodium falciparum* by High Throughput Screening of Microbial Natural Products. *PLoS ONE* 11, e0145812

Acknowledgements

This work has been funded by the Instituto de Salud Carlos III (ISCIII) (RICET: RD16/0027/0014, RD16/0027/0015, and RD12/0018/0005), the Plan Nacional (SAF PID2019-109623RB-I00; SAF2016-79957-R) and the Junta de Andalucía (BIO-199).

* Presenting author. E-mail address: fernando.reyes@medinaandalucia.es (Fernando Reyes)

Naphthylisoquinoline alkaloids streamlining mycetoma drug discover by circumventing differential diagnosis

Sami Ahmed Khalid^{a,*}, Shereen Abd Algaffar^a, Nasir Tajuddeen^b, Blaise Kimbadi Lombe^{c,d}, Gerhard Bringmann^b

^a Faculty of Pharmacy, University of Science and Technology, Omdurman, Sudan

^b Department of Chemistry, Ahmadu Bello University, Zaria, Nigeria

^c Institute of Organic Chemistry, University of Würzburg, Würzburg, Germany

^d Faculty of Sciences, University of Kinshasa, 11 Kinshasa, Democratic Republic of the Congo

Mycetoma is a chronic, slowly developing, devastating, and progressive neglected tropical infection of the subcutaneous tissues caused by either bacteria (actinomycetoma) or fungi (eumycetoma), leading to life-long disabilities and placing a significant burden on the poorest members of the society. The lack of reliable, safe, fast, and affordable diagnostic methods for identifying the causative agent is a considerable problem that hampers prescribing appropriate medication. Current therapy of eumycetoma causative agents, such as *Madurella mycetomatis*, consists of long-term azole antifungal agents coupled with surgery, yet with unsatisfactory clinical outcomes. Actinomycetoma is more responsive to treatment with co-trimoxazole and amikacin (Relhan et al., 2017). Hence, there is a pressing need to discover novel broad-spectrum antimicrobial agents to circumvent the time- consuming and costly diagnosis. A series of 23 naphthylisoquinoline (NIQ) alkaloids and related naphthoquinones were subjected to *in vitro* screening against three fungal strains of *M. mycetomatis*, and two bacterial strains of *Actinomyces madurae*, and *A. syzygii* (Abd Algaffar et al., 2021, 2022). Seven NIQs, mostly dimers, showed promising *in vitro* activities against at least one strain, while the naphthoquinones did not inhibit any of the tested strains. A synthetic NIQ dimer, 8,8'''-O,O'-dimethylmichellamine A **1**, inhibited all tested fungal and bacterial strains (IC₅₀ = 2.81-12.07 µg/mL). Michellamine B **2**, an *in vitro* active dimeric NIQ inhibited a strain of *M. mycetomatis* and significantly enhanced the survival rate of *M. mycetomatis* infected-*Galleria mellonella* larvae at concentrations of 1 and 4 µg/mL, without being toxic to the uninfected *Galleria* larvae (Konings et al., 2021).




Fig. 1. Structure of 8,8'''-O,O'-dimethylmichellamine A **1** and michellamine B **2**

Keywords: eumycetoma, actinomycetoma, broad-spectrum agents, naphthylisoquinoline alkaloids

References

- Abd Algaffar SO, Verbon A, vande Sande WWJ, Khalid SA. 2021. Development and validation of an *in vitro* resazurin- based susceptibility assay against *Madurella mycetomatis*. *Antimicrob Agents Chemother.* 65:e01338-20.
- Abd Algaffar SO, Verbon A, Khalid SA, van de Sande WWJ. 2022. Development and validation of a resazurin assay for *in vitro* susceptibility testing of *Actinomyces madurae*: a common causative agent of actinomycetoma. *J Antimicrob Chemother.* 23;78(1):155-160.
- Konings M, Eadie K, Lim W, Fahal AH, Mouton J, Tesse N, van de Sande W. 2021. The synthetic synergistic cinnamon oil CIN-102 is active against *Madurella mycetomatis*, the most common causative agent of mycetoma. *PLoS Negl Trop Dis.* 15(6).
- Relhan V, Mahajan K, Agarwal P, Garg VK. 2017. Mycetoma: an update. *Indian J Dermatol.* 62(4):332.

* Presenting author. E-mail address: khalidseek@gmail.com (Sami Ahmed Khalid)

Antibacterial marinopyrroles and pseudilins act as protonophores

Chambers C. Hughes^{a,b,d,e,*} Gabriel Castro-Falcón^a, Jan Straetener^b, Jan Bornikoel^c, Daniela Reimer^a, Trevor N. Purdy^a, Anne Berscheid^b, Florence M. Schempp^a, Dennis Y. Liu^c, Roger G. Linington^c, Heike Brötz-Oesterhelt^{b,d,e}

^a Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California, San Diego, California 92093, USA

^b Department of Microbial Bioactive Compounds, Interfaculty Institute of Microbiology and Infection Medicine, University of Tübingen, 72076 Tübingen, Germany

^c Department of Chemistry, Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada ^d Cluster of Excellence EXC 2124: Controlling Microbes to Fight Infection, University of Tübingen, 72076 Tübingen, Germany

^e German Center for Infection Research, Partner Site Tübingen, 72076 Tübingen, Germany

Elucidating the mechanism of action (MoA) of antibacterial natural products is crucial to evaluating their potential as novel antibiotics. Marinopyrrole (**1**), pentachloropseudilin (**2**), and pentabromopseudilin (**3**) are densely halogenated, hybrid pyrrole-phenol natural products with potent activity against Gram-positive bacterial pathogens like *Staphylococcus aureus* (Fig. 1) (Hughes et al., 2008). However, the exact way they exert this antibacterial activity has not been established. In this study, we explore their structure–activity relationship, determine their spatial location in bacterial cells, and investigate their MoA (Castro-Falcón et al., 2024). We show that the natural products share a common MoA based on membrane depolarization and dissipation of the proton motive force (PMF) that is essential for cell viability. The compounds show potent protonophore activity but do not appear to destroy the integrity of the cytoplasmic membrane via the formation of larger pores or interfere with the stability of the peptidoglycan sacculus. Thus, our current model for the antibacterial MoA of marinopyrrole, pentachloropseudilin, and pentabromopseudilin stipulates that the acidic compounds insert into the membrane and transport protons inside the cell. This MoA may explain many of the deleterious biological effects in mammalian cells, plants, phytoplankton, viruses, and protozoans that have been reported for these compounds.




Fig. 1. Structure of marinopyrrole (**1**), pentachloropseudilin (**2**), pentabromopseudilin (**3**)

Keywords: halogens, antibiotics, protonophore, fluorescent microscopy, *Staphylococcus aureus*

References

Hughes CC, Prieto-Davo A, Jensen PR, Fenical W. The marinopyrroles, antibiotics of an unprecedented structure class from a marine *Streptomyces* sp. *Org Lett*. 2008;10:629-631.

Castro-Falcón G, Straetener J, Bornikoel J, Reimer D, Purdy TN, Berscheid A, Schempp FM, Liu DY, Linington RG, Brötz-Oesterhelt H, Hughes CC. Antibacterial marinopyrroles and pseudilins act as protonophores. *ACS Chem Biol*. 2024;19:743-752.

* Presenting author. E-mail address: chambers.hughes@uni-tuebingen.de (Chambers C. Hughes)

Session III: Biotransformation and biosynthesis of natural products

Topical Lecture

Flavoenzyme-driven structural diversification of bacterial polyketide and tropone natural products

Robin Teufel

Department of Pharmaceutical Sciences, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

Bacterial aromatic polyketides as well as tropone natural products are structurally diverse and often play crucial roles in mutualistic and antagonistic symbiotic interactions with eukaryotic hosts. The biosynthesis of their backbones relies on dedicated polyketide synthases and accessory enzymes in the case of the aromatic polyketides and on an unusual intertwining of primary and secondary metabolism for the tropones, in which the initial steps depend on enzymes from phenylacetic acid catabolism that give rise to a central tropone precursor. The structural diversification of these backbones are then mediated by tailoring enzymes that are specific to each type of natural product and producer strain. In my talk, I will highlight how structurally and functionally distinct flavoenzymes are employed during late-stage biosynthesis for the functionalization of both pentangular rubromycin-type polyketides as well as tropone scaffolds and thereby generate potent protein inhibitors, virulence factors and antibiotics.

Keywords: bacterial aromatic polyketides, tropone natural products, flavoenzymes

* Presenting author. E-mail address: robin.teufel@unibas.ch (Robin Teufel)..

Oral contributions

Streamlined chemoenzymatic synthesis of cyclic peptides by non-ribosomal peptide cyclases

Toshiyuki Wakimoto*

Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan

Macrocyclization of peptide improves their pharmaceutical properties such as cell-membrane permeability, target specificity and metabolic stability. However, regio- and chemoselective intramolecular cyclization remain challenging. In the biosynthesis of non-ribosomal cyclic peptides, macrocyclases such as thioesterases (NRPS-TEs) efficiently catalyze the intramolecular peptide cyclization in a regiospecific manner without the use of protecting groups. Thus, NRPS-TEs are promising biocatalysts for the efficient synthesis of macrocyclic peptides. NRPS-TEs generally require the thioester leaving groups on its substrate, which are usually installed by solution-phase coupling reactions during substrate synthesis. However, this step often generates epimerized products which necessitates the purification of desired peptide from diastereomeric mixture, which is notoriously difficult and time-consuming. In this study, we have developed a streamlined chemoenzymatic approach to synthesize cyclic peptides that bypasses the need for leaving group installation in solution phase (Kobayashi et al., 2023). Linear peptides with diol ester functionalities on C-terminus were synthesized on a solid support to circumvent the solution-phase installation of leaving groups. Cleavage of the resin-bound peptides yielded the diol esters with sufficient purity. The diol-activated peptides were efficiently cyclized in a head-to-tail manner by SurE, a representative penicillin-binding protein-type thioesterase which we previously discovered in biosynthesis of non-ribosomal peptide, surugamides (Kuranaga et al., 2018; Matsuda et al., 2019; 2020). Explorations of homologous wild type enzymes as well as rational protein engineering, have broadened the scope of the enzymatic macrolactamization (Kobayashi et al., 2023). This method will potentially accelerate the exploitation of NRPS-TEs as biocatalysts.




Fig. 1. Diol-activated peptides were synthesized by SPPS and cyclized by NRP cyclases.

Keywords: NRPS, cyclase, cyclic peptide, thioesterase, SPPS

References

- M. Kobayashi, K. Fujita, K. Matsuda, T. Wakimoto, *J. Am. Chem. Soc.* 2023, 145, 3270–3275.
- T. Kuranaga, K. Matsuda, A. Sano, M. Kobayashi, A. Ninomiya, K. Takada, S. Matsunaga, T. Wakimoto, *Angew. Chem., Int. Ed.* 2018, 57, 9447–9451.
- K. Matsuda, M. Kobayashi, T. Kuranaga, K. Takada, H. Ikeda, S. Matsunaga, T. Wakimoto, *Org. Biomol. Chem.* 2019, 17, 1058–1061.
- K. Matsuda, R. Zhai, T. Mori, M. Kobayashi, A. Sano, I. Abe, T. Wakimoto, *T. Nat. Catal.* 2020, 3, 507–515.

* Presenting author. E-mail address: wakimoto@pharm.hokudai.ac.jp (Toshiyuki Wakimoto).

Enzymatic synthesis of human phase II metabolites of flavonoids and other (poly)phenols

Barbora Petránková, Katerina Brodsky, Lucie Petrásková, Pavla Bojarová, Kateřina Valentová*

Institute of Microbiology of the CAS, Videňská 1083, 14200 Prague, Czechia

Standards of Phase II metabolites of flavonoids and other (poly)phenols, needed as standards to assess their bioavailability and biological activity, are generally not commercially available. Enzymatic biotransformation is a promising route for the synthesis of such metabolites, it offers advantages in terms of efficiency, selectivity and sustainability compared to conventional chemical methods. This contribution explores the applications of enzymes in the production of phase II metabolites including challenges such as substrate specificity, scalability, and optimization with particular focus on aryl sulfotransferases (ASTs). AST from *Desulfitobacterium hafniense* has been successfully used for the sulfation of various (poly)phenols (Valentová 2017, Valentová 2018, Káňová 2020, Kolaříková 2022, Petrásková 2022). New potential recombinant ASTs were then selected, produced in *E. coli*, purified, and characterized including kinetic parameters. Substrate specificity was tested using *p*-nitrophenyl sulfate and 4-methylumbelliferyl sulfate as donors and various flavonoids as acceptors, which differed mainly in the amount and positions of the -OH groups (chrysin, apigenin, luteolin, kaempferol, myricetin, quercetin and silydianin). The reactions were monitored using optimized HPLC methods. Enzymatic sulfation of polyphenols with bacterial ASTs is an effective method for the production of authentic metabolites.




Fig. 1. Sulfation of myricetin

The work was supported by projects NU21-02-00135 and 23-04654S.

Keywords: aryl sulfotransferase, sulfation, flavonoid, phenolic, metabolite

References

- Káňová, K., L. Petrásková, H. Pelantová, Z. Rybková, K. Malachová, J. Cvačka, V. Křen and K. Valentová, 2020. "Sulfated metabolites of luteolin, myricetin, and ampelopsin: Chemoenzymatic preparation and biophysical properties." *J. Agric. Food Chem.* 68, 11197-11206.
- Kolaříková, V., K. Brodsky, L. Petrásková, H. Pelantová, J. Cvačka, L. Havlíček, V. Křen and K. Valentová, 2022. "Sulfation of phenolic acids: Chemoenzymatic vs. chemical synthesis." *Int. J. Mol. Sci.* 23, 15171.
- Petrásková, L., K. Káňová, K. Brodsky, A. Hetman, B. Petránková, H. Pelantová, V. Křen and K. Valentová, 2022. "Sulfated phenolic substances: Preparation and optimized HPLC analysis." *Int. J. Mol. Sci.* 23, 5743.
- Valentová, K., K. Káňová, F. Di Meo, H. Pelantová, C. Chambers, L. Rydlová, L. Petrásková, A. Křenková, J. Cvačka, P. Trouillas and V. Křen, 2017. "Chemoenzymatic preparation and biophysical properties of sulfated quercetin metabolites." *Int. J. Mol. Sci.* 18, 2231.
- Valentová, K., K. Purchartová, L. Rydlová, L. Roubalová, D. Biedermann, L. Petrásková, A. Křenková, H. Pelantová, V. Holečková-Moravcová, E. Tesařová, J. Cvačka, J. Vrba, J. Ulrichová and V. Křen, 2018. "Sulfated metabolites of flavonolignans and 2,3-dehydroflavonolignans: Preparation and properties." *Int. J. Mol. Sci.* 19, 2349.

* Presenting author. E-mail address: kata.valentova@email.cz (Kateřina Valentová).

Fungi – important biocatalysts in production of natural pigments and fragrances

Filip Boratyński^{a,*}, El-Sayed R. El-Sayed^{a,b}, Ewa Szczepańska^a, Stefano Serra^c

^a Department of Food Chemistry and Biocatalysis, Wrocław University of Environmental and Life Sciences, Norwida 25, 50-375 Wrocław, Poland

^b Plant Research Department, Nuclear Research Center, Egyptian Atomic Energy Authority, Cairo, Egypt

^c Consiglio Nazionale delle Ricerche (C.N.R.), Istituto di Scienze e Tecnologie Chimiche, Via Mancinelli 7, 20131 Milano, Italy

Due to the awareness of consumers regarding the origin of food additives, as well as the growing demand in the industry for natural pigments and fragrances, the development of an alternative method for their synthesis is currently explored (Maurya et al., 2022). Biotechnological methods constitute the most attractive solution as they are more sustainable and environmentally friendly processes with an emphasis on the prevention of waste generation and avoidance of hazardous compounds. Fungi, among the microbial populations, could be used as the most efficient biotechnology agents for the production of colorant and aroma compounds due to their tolerance, flexibility, and cheap requirements of growth. Microbial pigments, in contrary to plant-based pigments, show several advantages including the ease of scaling up and harvest as well as not subjected to the nature vagaries. Herein, the application of endophytic fungi as an efficient production platform for natural pigments will be discussed. A promising strain isolated from leaves of *Origanum majorana* was identified as *Monascus ruber* SRZ112 produced several types of pigments, mainly rubropunctamine, monascin, ankaflavin, rubropunctatin, and monascorubrin (El-Sayed et al., 2022). In the area of flavors and fragrances different biosynthesis and biotransformation methods involving whole fungal cells will be presented: filamentous fungi transformation of ferulic acid to vanillin using different agro-industrial by-products; Basidiomycete-mediated synthesis of anisaldehyde, vanillin, veratraldehyde and piperonal (Serra et al., 2024). Our research provide results with both high knowledge value and applicability of the developed fungal-based transformations in the synthesis of natural pigments and flavors.

Acknowledgments: The presented research is part of the BioExplor project No. 2021/43/P/NZ9/02241 co-funded by the National Science Centre and the European Union Framework Programme for Research and Innovation Horizon 2020 under the Marie Skłodowska-Curie grant agreement no. 945339.

Keywords: fungi, *Monascus*, piperonal, vanillin, agro-industrial wastes

References

- Maurya R., Patel H., Bhatt D., Shakhreliya S., Gohil N., Bhattacharjee G., Lam N.L., Alzahrani K.J., Gyanchander E., Singh V. 2022 Microbial production of natural flavors and fragrances. In: Recent advances in food biotechnology, Kumar A., Patruni K., Singh V. (eds). Springer.
- El-Sayed R. El-Sayed; Gach J., Olejniczak T., Boratyński F., 2022 A New Endophyte *Monascus ruber* SRZ112 as an Efficient Production Platform of Natural Pigments Using Agro-industrial Wastes, Sci. Rep. 12, 12611.
- Serra S., Marzorati S., Szczepańska E., Strzała T., Boratyński F., 2024 Basidiomycota strains as whole-cell biocatalysts for the synthesis of high-value natural benzaldehydes, Appl. Microbiol. Biotechnol., 2024, 108, 1–12.

* Presenting author. E-mail address: filip.boratynski@upwr.edu.pl (Filip Boratyński).

First insights in the metabolic faith of paraconic acids from *Cetraria islandica*

Eduardo Villicaña González^a, Eva Sanchez Armengol^b, Annelies Breynaert^c, Kenn Foubert^c, Hermann Stuppner^a, Nina Hermans^c, Flavia Laffleur^b, Stefan Schwaiger^{a,*}

^a Institute of Pharmacy/Pharmacognosy, Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, CCB, Innrain 80/82, 6020, Innsbruck, Austria

^b Institute of Pharmacy/Pharmaceutical Technology/Research Group Drug Delivery, University of Innsbruck, Innrain 80-82, 6020, Innsbruck, Austria

^c Faculty of Pharmaceutical, Biomedical and Veterinary Sciences/Department of Pharmaceutical Sciences/Research Group NatuRAPT, University of Antwerp, Universiteitsplein 1, 2610, Wilrijk, Belgium

Cetraria islandica or Icelandic moss is a lichen employed in traditional European medicine to treat mainly respiratory diseases, but also some gastrointestinal complications such as gastric and duodenal ulcers, indigestion, diarrhoea, and dysentery. The activity is mainly related to the contained polysaccharides, but several other compounds can be detected in decoctions of the lichen thalli, representing a recommended formulation. Among depsides and depsidones, also paraconic acids (PA), with known *in vitro* pharmacological properties on the arachidonic acid pathway, can be detected up to 2.7 w/% of used lichen material, when quantified by LC-MS. Up to now, several PA have been identified from *C. islandica* including dihydroprotolichesterinic acid, roccellaric acid, protolichesterinic acid, lichesterinic acid **1** (Fig 1), and nephrosterinic acid (González et al., 2023; Fernandes et al., 2020).




Fig. 1. Structure of lichesterinic acid **1**.

Thus, this project aimed to analyse if PAs could potentially penetrate into the gut tissue by analysing the permeation and absorption of lichesterinic acid through and into a Caco-2 monolayer. A weak to moderate absorption profile was indicated by analysis of the Caco-2 monolayer model results (time points 0, 6, 24, and 48 hours) using two concentrations (50 and 100 μ M) of **1**, which prompted a thorough examination utilizing an *in vitro* gastrointestinal digesting model (GIDM). Due to the described antibacterial effects of PAs we also monitored the colon microbiota viability in the presence of PAs via quantification of colonies formed.

Keywords: paraconic acids, *C. islandica*, caco-2 monolayer, *in-vitro* gastrointestinal digestion model

References

- González, E.V., Schwaiger, S., Stuppner, H., 2023. Two-step isolation of the two major paraconic acids of *Cetraria islandica*. J of Separation Science 46, 2200805.
- Fernandes, R.A., Chaudhari, D.A., Jha, A.K., 2020. Evolution of Strategies in Paraconic Acids Synthesis. Asian Journal of Organic Chemistry 9, 1478–1501.

* Presenting author. E-mail address: stefan.schwaiger@uibk.ac.at (Stefan Schwaiger).

Biotransformation of favelines from *Cnidoscolus quercifolius* into a new bis-nor-diterpene with anti-melanoma potential

Raimundo Gonçalves de Oliveira Junior^{a,*}, David Hozain^a, Thomas Gaslonde^a, Séverine Amand^b, Jackson Roberto Guedes da Silva Almeida^c, Laurent Picot^d, Marie-Christine Lallemand^a, Didier Buisson^b

^a CiTCoM UMR 8038 CNRS, Paris Cité University, 75006 Paris, France

^b MCAM UMR 7245 CNRS, National Museum of Natural History (NMNH), 75005 Paris, France

^c NEPLAME, Federal University of São Francisco Valley, 56304-917 Petrolina, Brazil

^d LIENSs UMRi 7266 CNRS, La Rochelle University, 17000 La Rochelle, France

C. quercifolius is a Brazilian medicinal plant, rich in bis-nor-diterpenes known as favelines. These molecules have a tricyclic benzocycloheptenone basic skeleton and showed significant cytotoxic activity against chemoresistant human melanoma cells (A2058 cell line). Phyllacanthone, its major compound, exhibited pro-apoptotic activity associated with tubulin depolymerization, resulting in cytoskeleton disruption (Oliveira-Junior et al., 2022). To design new anti-melanoma favelines, we performed biotransformation assays on *C. quercifolius* crude extract using 12 fungi and bacterial strains from the microorganism collection of NMNH (Paris, France). All biotransformation reactions were carried out in citrate buffer (pH 5.0), and the formation of new products was monitored by HPLC-DAD and/or UHPLC-DAD-HRMS/MS every 24 hours. A combinatorial approach allowed the selection of *Mortierella isabellina* as the active strain, resulting in a major biotransformation product after 24 hours of reaction with the crude extract. This compound was obtained from the supernatant by liquid-liquid extraction and from the biomass by ultrasound-assisted extraction, and then purified by preparative HPLC. After HRMS and 1D and 2D NMR experiments, the new compound was identified as a phyllacanthone derivative hydroxylated at C-12 and C-14. Analytical biotransformation assays with the major favelines isolated from the crude extract (phyllacanthone and isofavelol) led to the same biotransformation compound, indicating that they probably served as substrates for *M. isabellina* (Fig. 1). Finally, the cytotoxic activity of the new faveline was assessed on A2058 cells using the MTT method, resulting in an $IC_{50} = 65.8 \mu M$.




Fig. 1. Biotransformation of phyllacanthone by the fungus *M. isabellina* NRRL 1757.

Keywords: bioconversion, diterpene, Euphorbiaceae, melanoma, skin cancer

References

Oliveira-Junior, R.G., Ferraz, C.A.A., Oliveira, A.P., Araújo, E.C.C., Prunier, G., Beaugeard, L., Groult, H., Picot, L., Alencar-Filho, E.B., Aouad, N., Rolim, L.A., Almeida, J.R.G.A., 2022. Chem. Biol. Interact. 355, 109849.

* Presenting author. E-mail address: raimundo.goncalves-de-oliveira-junior@u-paris.fr (Raimundo Gonçalves de Oliveira Junior).

Enterolactone and enterodiols produced by fermentation of Valerian root extract with human fecal samples

Maria-Eleni Grafakou^{a,*}, Eva-Maria Pferschy-Wenzig^a, Olaf Kelber^b, Rudolf Bauer^a

^a Institute of Pharmaceutical Sciences, Department of Pharmacognosy, University of Graz, Austria

^b Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt, Germany

Valeriana officinalis L. is commonly used to alleviate mild mental stress and sleep disorders. However, like in many herbal extracts, there remains a lack of scientific consensus regarding its active ingredients (Shinjyo et al., 2020). With the aim to investigate its potential gut microbiome-mediated biotransformation, we subjected valerian root extract to fermentation with human fecal microbiota from three healthy donors (24 h, anoxic, 37°C) (Pérez-Burillo et al., 2021). Annotation of the constituents present in the native preparation and of the metabolites formed during anaerobic fermentation was accomplished by UHPLC-HRMS. In the extract, lignans and caffeic acid derivatives were detectable as major constituents and valerenic acids, valepotriates and baldrinals as minor constituents. During fermentation with fecal samples, the phenylpropanoid derivatives were found to be readily catabolized by gut microbiota. The mammalian lignans enterolactone and enterodiols were found as the major metabolites in all donor samples. It is noteworthy that enterolactone was recently associated with lower prevalence of depressive symptoms (Cui et al., 2022) and sleep disorders (Sun et al., 2020). This indicates that the metabolic activation of *V. officinalis* constituents by the gut microbiota, leading to newly produced mammalian lignans, could contribute to the observed therapeutic effects of Valerian root preparations.

Conflicts of interest: The investigations and MEG have been funded by, and OK is fully employed by Steigerwald Arzneimittelwerk, Bayer Consumer Health.

Keywords: *Valeriana officinalis*, gut microbiota, biotransformation, enterodiols, enterolactone

References

- Cui, K., Luo, J., Zhang, L., Zhang, T., Feng, Z., Zhang, D., 2022. The association between urinary phytoestrogens and depressive symptoms. *Int. J. Food Sci. Nutr.* 73, 657–668.
- Pérez-Burillo, S., Molino, S., Navajas-Porras, B., et al., 2021. An in vitro batch fermentation protocol for studying the contribution of food to gut microbiota composition and functionality. *Nat. Protoc.* 16, 3186–3209.
- Shinjyo, N., Waddell, G., Green, J., 2020. Valerian root in treating sleep problems and associated disorders—a systematic review and meta-analysis. *J. Evid. Based Integr. Med.* 25, 1–31.
- Sun, J., Jiang, H., Wang, W., Dong, X., Zhang, D., 2020. Associations of urinary phytoestrogen concentrations with sleep disorders and sleep duration among adults. *Nutrients.* 12, 2103.

* Presenting author. E-mail address: maria.grafakou@uni-graz.at (Maria-Eleni Grafakou).

Biosynthesis of bicyclo[3.3.1]nonane derivatives in *Hypericum sampsonii*

Lukas Ernst^{a,*}, Hui Lyu^b, Pi Liu^c, Christian Paetz^b, Hesham MB Sayed^{a,d}, Tomke Meents^a, Hongwu Ma^c, Ludger Beerhues^a, Islam El-Awaad^{a,d}, Benye Liu^a

^a Technische Universität Braunschweig, Institute of Pharmaceutical Biology, Braunschweig 38106, Germany

^b Max Planck Institute for Chemical Ecology, NMR/Biosynthesis Group, Jena 07745, Germany

^c Chinese Academy of Sciences, Tianjin Institute of Industrial Biotechnology, Biodesign Center, Tianjin 300308, China

^d Assiut University, Faculty of Pharmacy, Department of Pharmacognosy, Assiut 71526, Egypt

Polycyclic polyprenylated acylphloroglucinols (PPAPs) with bicyclo[3.3.1]nonane cores are medicinally intriguing compounds, which exist in two constitutions (Yang et al., 2018). Prominent examples are hyperforin (type A) and garcinol (type B). Both PPAP types are present in *Hypericum sampsonii* (Hypericaceae). This presentation will focus on the identification and characterization of a pair of bifunctional prenyltransferases (*HsCPTa*, *HsCPTb*), which catalyze regiodivergent prenylative cyclizations to yield the PPAP regiomers 7-*epi*-nemorosone and 7-*epi*-clusianone (Fig. 1; Ernst et al., revision submitted).




Fig. 1. Inverted substrate binding modes in *HsCPTa* and *HsCPTb*, yielding type A and B bicyclo[3.3.1]nonane scaffolds

Molecular modelling and docking experiments predicted inverted substrate binding modes in the active site cavities of the two enzymes. Reciprocal mutagenesis demonstrated that the regiospecific binding modes are interconvertible by ninefold mutation. Detailed structural elucidation of the enzymatic products confirmed *endo* configurations of the transferred C-7 prenyl groups, required for biosynthesis of caged adamantane-type PPAPs characteristic of *H. sampsonii* (Hu et al., 2000).

Keywords: biosynthesis, acylphloroglucinols, *Hypericum sampsonii*, prenyltransferases

References

- Ernst, L., Lyu, H., Liu, P., Paetz, C., Sayed, H.M.B., Meents, T., Ma, H., Beerhues, L., El-Awaad, I., Liu, B., 2024. Regiodivergent biosynthesis of bridged bicyclononanes. *Nat. Commun.* (revision submitted).
- Yang, X.W., Grossman, R.B., Xu, G., 2018. Research progress of polycyclic polyprenylated acylphloroglucinols. *Chem. Rev.* 118, 3508–3558.
- Hu, L.H., Sim, K.Y., 2000. Sampsoniones A–M, a unique family of caged polyprenylated benzoylphloroglucinol derivatives, from *Hypericum sampsonii*. *Tetrahedron* 56, 1379–1386.

* Presenting author. E-mail address: lukas.ernst@tu-braunschweig (Lukas Ernst).

Unravelling the biosynthesis of phenylphenalenones: insights from metabolites and enzymes in *Musella lasiocarpa*

Hui Lyu^{a,*}, Lukas Ernst^b, Yoko Nakamura^{a,d}, Yu Okamura^f, Tobias G. Köllner^{c,d}, Katrin Luck^{c,d}, Benye Liu^b, Yu Chen^c, Ludger Beerhues^b, Jonathan Gershenzon^c, Christian Paetz^a

^a NMR/Biosynthesis Group, Max Planck Institute for Chemical Ecology, Germany

^b Technische Universität Braunschweig, Institute of Pharmaceutical Biology, Germany

^c Department of Biochemistry, Max Planck Institute for Chemical Ecology, Germany

^d Department of Natural Product Biosynthesis, Max Planck Institute for Chemical Ecology, Germany

^e Jiangsu Key Laboratory for the Research and Utilization of Plant Resources, Institute of Botany, Jiangsu Province and Chinese Academy of Sciences, China

^f Department of Insect Symbiosis, Max Planck Institute for Chemical Ecology, Germany

Phenylphenalenones (PPs) are complex polycyclic phytoalexins found in banana and plantain (Musaceae) plants. Previous studies have linked the disease susceptibility of commercial banana cultivars to low levels of PPs compared to disease-resistant wild bananas (Krishnamurthy et al., 2023). However, how plants synthesize PPs is still poorly understood.



Fig. 1. Characterized transformations to phenylphenalenone in *Musella lasiocarpa* (Musaceae).

We find that PPs are the major secondary metabolites of mature seeds of *M. lasiocarpa* and that seeds can be a model system to study PP biosynthesis (Lyu et al., 2023). Using transcriptomic data from *M. lasiocarpa* seeds at different developmental stages, we identified three biosynthetic genes involved in the formation of dihydrobisdemethoxycurcumin (Lyu et al., under review). Furthermore, through multiple rounds of feeding potential intermediates to *M. lasiocarpa* root protein extracts, we demonstrate the stepwise conversion of dihydrobisde-methoxycurcumin to the PP 4’hydroxylachnanthocarpone (Fig. 1). The discovery of a monocyclic diarylheptanoid as an unexpected PP intermediate opens a two-step cyclization route to the phenylphenalenone scaffold which contrasts the commonly hypothesized Diels–Alder mechanism.

Keywords: biosynthesis, phenylphenalenones, diarylheptanoids, Musaceae, natural products

References

- Krishnamurthy, P., Ravikumar, M. J., Arumugam Palanivelu, S., Pothiraj, R., Suthanthiram, B., Subbaraya, U., Morita, H., 2023. Phenylphenalenone-type phytoalexins in banana (*Musa* species): a comprehensive review for new research directions. *Phytochem. Rev.* 22, 187–210.
- Lyu, H., Chen, Y., Gershenzon, J., Paetz, C., 2023. Diarylheptanoid Derivatives (Musellins A–F) and Dimeric Phenylphenalenones from Seed Coats of *Musella lasiocarpa*, the Chinese Dwarf Banana. *J. Nat. Prod.* 86, 1571–1583.
- Lyu, H., Ernst, L., Nakamura, Y., Okamura, Y., Koellner, T., Luck, K., Liu, B., Chen, Y., Beerhues, L., Gershenzon, J., Paetz, C., 2024. Phenylphenalenones and Linear Diarylheptanoid Derivatives are Biosynthesized via Parallel Routes in *Musella lasiocarpa*, the Chinese Dwarf Banana. *Angew. Chem. Int. Ed.* Under review.

* Presenting author. E-mail address: hlyu@ice.mpg.de (Hui Lyu).

A synthetic biology platform for refactoring and combinatorial biosynthesis of anthracyclines

Rongbin Wang^a, Heli Tirkkonen^a, Khaled A. Shaaban^b, Mikko Metsä-Ketelä^a, S. Eric Nybo^{c,*}

^a Department of Life Technologies, University of Turku, FIN-20014 Turku, Finland

^b Department of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, Kentucky 40536, USA

^c Department of Pharmaceutical Sciences, College of Pharmacy, Ferris State University, Big Rapids, MI 49307, USA

Aromatic polyketides produced predominantly by actinobacteria typically exhibit antimicrobial, anticancer, and immunosuppressant activities (Hulst et al., 2022). Aromatic polyketides, such as antitumor compound doxorubicin, are biosynthesized by type II polyketide synthases (PKS), which carry out the iterative condensation of acyl-CoA precursors. Doxorubicin is a broad-spectrum anticancer agent used in the clinic. However, its use is diminished by cardiotoxicity. New investigations into cardiotoxicity-free doxorubicins, which do not induce DNA strand breakage, have reinvigorated interest in this class of natural products (Hulst et al., 2022). We have employed a biosynthetic approach to complement the synthetic studies recently reported. Developing standardized genetic tools to facilitate true “plug and play” experimentation could further enhance the combinatorial biosynthesis of these pathways. Here we present a metabolic engineering platform that allows rapid generation of anthracyclines for bioactivity testing³. Our modular BioBricks-based design architecture for (i) aglycone biosynthesis, (ii) tailoring steps, (iii) TDP-carbohydrate biosynthesis and (iv) resistance and carbohydrate transfer allows facile construction of any desired combination from existing genetic parts. Simultaneous development of a stable host chassis *S. coelicolor* M1152 Δ matAB and a multi-plasmid gene expression system has facilitated robust production of the natural products. Current progress on the iterative engineering of anthracycline pathways will be discussed (Nguyen et al., 2022, Wang et al., 2022, Tirkkonen et al., 2023).

Keywords: synthetic biology, actinomycetes, BioBricks, polyketide synthase, biosynthesis

References

- Hulst, M.B., Grocholski, T., Neefjes, J.J.C., van Wezel, G.P., Metsä-Ketelä, M., 2022. Anthracyclines: biosynthesis, engineering and clinical applications. *Nat Prod Rep.* 39, 814-841.
- Nguyen JT, Riebschleger KK, Brown KV, Gorgijevska NM, Nybo S.E., 2022. A BioBricks toolbox for metabolic engineering of the tetracenomycin pathway. *Biotechnol J.* 17, e2100371.
- Wang, R., Nguyen, J., Hecht, J., Schwartz, N., Brown, K.V., Ponomareva, L.V., Niemczura, M., van Dissel, D., van Wezel, G.P., Thorson, J.S., Metsä-Ketelä, M., Shaaban, K.A., Nybo, S.E., 2022. A BioBricks metabolic engineering platform for the biosynthesis of anthracyclines in *Streptomyces coelicolor*. *ACS Synth Biol* 11: 4193–4209.
- Tirkkonen H, Brown KV, Niemczura M, Faudemer Z, Brown C, Ponomareva LV, Helmy YA, Thorson JS, Nybo SE, Metsä-Ketelä M, Shaaban KA. 2023. Engineering BioBricks for Deoxysugar Biosynthesis and Generation of New Tetracenomycins. *ACS Omega.* 8, 21237-21253.

* Presenting author. E-mail address: EricNybo@Ferris.edu (S. Eric Nybo).

Discovery of new bacterial sesterterpene synthases and their products

Daniel Icenhour, Jeffrey Rudolf

Department of Chemistry, University of Florida, 165 Buckman Drive, Gainesville, FL 32611, USA

While terpenes remain the largest family of natural products, their structural diversity is what sets them apart from other families of natural products and confers a vast array of different biological activities. Sesterterpenes, made up of C₂₅ products, remains a relatively understudied subfamily that accounts for approximately 1,000 of the over 80,000 known terpenes, and even more staggering is that less than 25 of these compounds were isolated from bacteria (Rudolf et al., 2021). Here we describe a terpene production system in *E. coli* that we have developed to screen bacterial sesterterpene synthases and the isolation of their C₂₅ products. Our work here will help to characterize new bacterial sesterterpene synthases, isolate and elucidate the structures of new sesterterpenes.

Keywords: sesterterpenes, terpenoids, terpene synthases, biosynthesis, enzymes

References

Rudolf, J. D., Alsup, T. A., Xu, B., Li, Z., 2021. Bacterial terpenome. *Nat. Prod. Rep.* 38, 905-980.

* Presenting author. E-mail address: dicenhour@ufl.edu (Daniel Icenhour).

Colletofragarone A2 from the fungus *Colletotrichum* sp. promoted the degradation and aggregation of mutant p53

Sachiko Tsukamoto*

Department of Natural Medicines, Graduate School of Pharmaceutical Sciences, Kumamoto University, 5-1 Oe-honmachi, Kumamoto 862-0973, Japan

The tumour suppressor p53 regulates the transcription of genes for apoptosis, cell cycle arrest, and senescence. p53 are mutated in more than 50% of tumours, and mutant p53 aggregates in cells to promote cancer progression and lead to the loss of its tumour suppressor function. Therefore, compounds targeting mutant p53 are promising for cancer therapy and we search for compounds that show functional recovery of mutant p53 to wild-type like p53. Recently, we isolated colletofragarone A2 (1) and three new compounds colletoins A (2)–C (Fig. 1) from an endophytic fungus *Colletotrichum* sp. (Sadahiro et al., 2021). Compound 1 showed cytotoxicity with an IC_{50} value of 0.18 mM against SK-BR-3 (p53^{R175H}) cells and decreased tumour cell growth *in vivo* using xenograft model mice with HeCCT1 (p53^{R175H}) cells (Sadahiro et al., 2022). Compound 1 decreased mutant p53 levels in the cells, induced the degradation of mutant p53 by the proteasome, and promoted the accumulation of aggregated mutant p53 in the cells.




Fig. 1. Structures of colletofragarone A2 (1) and colletoin A (2).

Keywords: *Colletotrichum* sp., mutant p53, cancer

References

- Sadahiro, Y., Hitora, Y., Tsukamoto, S., 2021. Colletofragarone A2 and colletoins A–C from a fungus *Colletotrichum* sp. decrease mutant p53 levels in cells. *J. Nat. Prod.* 84, 3131–3137.
- Sadahiro, Y., Hitora, Y., Kimura, I., Hitora-Imamura, N., Onodera, R., Motoyama, K., Tsukamoto, S., 2022. Colletofragarone A2 inhibits cancer cell growth *in vivo* leads to the degradation and aggregation of mutant p53. *Chem. Res. Toxicol.* 35, 1598–1603.

* Presenting author. E-mail address: sachiko@kumamoto-u.ac.jp (Sachiko Tsukamoto).

Novel thiosugar sulfoniums, salacinol and neokotalanol, with antidiabetic activity obtained from plants of genus *Salacia*

Toshio Morikawa^{a,b,*}

^a Pharmaceutical Research and Technology Institute, Kindai University, 3-4-1 Kowakae, Higashi-osaka, Osaka, Japan

^b Antiaging Center, Kindai University, 3-4-1 Kowakae, Higashi-osaka, Osaka, Japan

During our studies characterizing functional substances from food resources for the prevention and treatment of lifestyle diseases, we isolated the active constituents, salacinol (1) and neokotalanol (2), and related thiosugar sulfoniums, from the roots and stems of the genus *Salacia* plants (Hippocrateaceae) such as *Salacia reticulata* Wight, *S. oblonga* Wall., and *S. chinensis* L., and observed their antidiabetic effects (Fig. 1) (Yoshikawa et al., 2002; Muraoka et al., 2011). These plant materials have been used traditionally in Ayurvedic medicine as a specific remedy at the early stage of diabetes, and have been extensively consumed in Japan, the United States, and other countries as a food supplement for the prevention of obesity and diabetes (Morikawa et al., 2021). Here, I would like to present our studies on the antidiabetic effects of plants from the genus *Salacia*, from basic chemical and pharmacological research to their application and development as new functional food ingredients (Morikawa, 2023).




Fig. 1. Structures of salacinol (1) and neokotalanol (2).

Keywords: *Salacia*, salacinol, neokotalanol, α -glucosidase inhibitor, diabetes

References

- Yoshikawa, M., Morikawa, T., Matsuda, H., Tanabe, G., Muraoka, O., 2002. Absolute stereostructure of potent α -glucosidase inhibitor, salacinol, with unique thiosugar sulfonium sulfate inner salt structure from *Salacia reticulata*. *Bioorg. Med. Chem.* 10, 1547-1554.
- Muraoka, O., Morikawa, T., Miyake, S., Akaki, J., Ninomiya, K., Pongpiriyadacha, Y., Yoshikawa, M., 2011. Quantitative analysis of neosalacinol and neokotalanol, another two potent α -glucosidase inhibitors from *Salacia* species, by LC-MS with ion pair chromatography. *J. Nat. Med.* 65, 142-148.
- Morikawa, T., Ninomiya, K., Tanabe, G., Matsuda, H., Yoshikawa, M., 2021. A review of antidiabetic active sulfoniums, salacinol and neokotalanol, from plants of the genus *Salacia*. *J. Nat. Med.* 75, 449-466.
- Morikawa, T., 2023. Pharmaceutical food science: search for bio-functional molecules obtained from natural resources to prevent and ameliorate lifestyle diseases. *Chem. Pharm. Bull.* 71, 756-765.

* Presenting author. E-mail address: morikawa@kindai.ac.jp (Toshio Morikawa).

3-epi-18 β -Glycyrrhetic acid and its 30-O-glucuronide involves the development of licorice-induced pseudoaldosteronism in humans with individual differences

Toshiaki Makino*

Department of Pharmacognosy, Graduate School of Pharmaceutical Sciences, Nagoya City University, Japan

Licorice is a crude drug used in traditional Japanese Kampo medicine and is also used as a sweetener. It occasionally causes pseudoaldosteronism (PsA) as a side effect. Its major symptoms are hypokalemia, hypertension, edema, and low plasma aldosterone levels. PsA is thought to be caused by some metabolites of glycyrrhizic acid (GL), a component of licorice. The development of PsA varies greatly among individuals, but the factors that determine the individual differences are still unknown. Overall, 78 patients who had taken Kampo medicines containing licorice were enrolled, and we obtained laboratory data including their serum potassium levels, plasma aldosterone concentrations, and residual blood and/or urine samples. Concentrations of GL metabolites in serum and urine samples were measured by LC-MS/MS. In the serum samples of the 78 participants, 18 β -glycyrrhetic acid (GA) was detected in 65 cases, 3-epi-GA in 47 cases, 3-oxo-GA in 63 cases, 18 β -glycyrrhetinyl-30-O-glucuronide (GA30G) in 62 cases, and 3-epi-GA30G in 3 cases. Among 29 urine samples obtained, GA30G in 27 cases, and 3-epi-GA30G in 19 cases. 3-epi-GA30G was a new GL metabolite, and the finding of 3-epi-GA, 3-oxo-GA, and 3-epi-GA30G in human samples was the first time. The appearances of 3-epi-GA in serum and 3-epi-GA30G in urine had high individual differences, and their concentrations significantly correlated with serum PsA markers. 3-epi-GA, 3-oxo-GA, GA30G, and 3-epi-GA30G inhibited human 11 β -hydroxysteroid dehydrogenase type 2 (11 β -HSD2) almost similar titers. These findings suggest that 3-epi-GA and/or 3-epi-GA30G would be associated with the cause of individual differences in the development of pseudoaldosteronism.

Keywords: licorice, pseudoaldosteronism, 3-epi-18 β -glycyrrhetic acid

* Presenting author. E-mail address: makino@phar.nagoya-cu.ac.jp (Toshiaki Makino).

Amway-Nutriline

Roses and buds, in the current state of botanical dietary supplements – a global perspective

Amit Chandra^{a,*}, Oran Kwon^b, Mark Blumenthal^c, Holly Johnson^d, Funda N. Yalcin^e, Fred Stevens^f

^a Amway Corporation, Innovation and Science, Michigan, 49355, USA

^b Ewha Womans University, Department of Nutritional Science and Food Management, Seoul, South Korea

^c American Botanical Council, Austin, Texas, 78723, USA

^d American Herbal Products Association, Silver Spring, Maryland, 20910, USA

^e Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy, Ankara, Turkey

^f College of Pharmacy, Linus Pauling Institute, Oregon State University, Corvallis, Oregon, USA

This oral session is dedicated to the 90th anniversary of the Nutrilite Dietary Supplement Brand from Amway. Nutrilite™ is one of the leaders in the dietary supplement industry in the world. This session will cover topics from pharmacognosy, and emerging technology related to consumers' health and wellbeing in the supplement industry. The session panel comprises of leaders in industry from Asia, Europe, and United States. ROSES represents the strengths and all the value-added developments that are currently happening and where this sector is headed in the future. BUDS are the emerging areas that can be seen as opportunities to improve or build on to convert into roses (evolution or continuous improvement). Topics such as botanical quality, efficacy, botanical authenticity, future of traditional medicine, AI-linked precision nutrition, sustainability, traceability, integrity, and impact of extreme weather will be covered in detail. A message to cross-pollinate the science of pharmacognosy with botanical ingredients and phytonutrients in commercial supplement products will be emphasized. For ICNPR (attendees and organization) it will be a unique opportunity to take a leadership role in industry and partner with industry cohorts for the enhancement of quality, safety, and efficacy of botanical products on a global stage. This session will have the speaker panel deliver individual talks followed by a moderated panel discussion and ending with an interactive discussion between panel and audience.

Keywords: botanicals, ai-nutrition, sustainability, ESCOP, traditional herbals, safety, quality, east and west

* Presenting author. E-mail address: amitabh_chandra@amway.com (Amit Chandra).



Society Session: GA

Welcome address

**Planta Medica Best Paper Award and PM Special Issue
„Celebrating Austrian Pharmacognosy”**

W. Schwabe Young Talent Price and Willmar Schwabe Award

Dr Cherry Lifeng Li, Baptist University, Hong Kong

GA Synergy – Award

Prof. Andreas Bender, University of Cambridge, United Kingdom

E. Stahl Bronze – Award

Dr Valentina Parisi, University of Salerno, Italy

Session I: Medicinal natural products: from bench to bedside

Topical Lecture

Will artificial intelligence become a game changer in natural product drug discovery?

Judith M. Rollinger

Division of Pharmacognosy, Department of Pharmaceutical Sciences, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

Nature has historically proven to deliver the smartest chemical motifs for drug discovery. Despite tremendous advances in instrumentation technologies, the exploration of natural sources is challenging. It requires not only deep knowledge and sophisticated technical equipment for getting natural compounds isolated, identified, and pharmacologically characterized; also aspects of sustainable supply and property rights of the underlying natural material have to be considered. On the other hand, already millions of data on physicochemical and biological properties of natural as well as synthetic compounds are in the public domain, which can be utilized by AI to streamline the experimental efforts for identifying underexplored natural hit compounds. Special care must be taken on the reliability of data and the AI method most suitable for the underlying research question. Here, I will present and compare some recently developed AI strategies implemented in natural product drug discovery workflows, which can be used to harness the advances in computational techniques for taking advantage from the huge amount of information available in the public domain. This ranges from virtual screening for hit identification, machine learning tools, molecular networking to a previously developed HNMR-MS-based biochemometric approach (Grienke, 2019). The effectiveness, strengths and limitations of the presented strategies will be demonstrated on most recent application examples for the identification of novel natural lead structures (Kirchweger, 2022; Langeder, 2023; Wasilewicz, 2023, 2024; Zwirchmayr 2023).

Funding: Wilhelm-Doerenkamp-Foundation, Switzerland (NATVANTAGE Grant 2018); Austrian Science Fund (FWF P34028)

Keywords: cheminformatics, biochemometry, molecular networking, SARS-CoV-2, traditional medicine

References

- Grienke, U., Foster, P.A., Zwirchmayr, J., Tahir, A., Rollinger, J.M., Mikros, E., 2019. ¹H NMR-MS-based heterocovariance as a drug discovery tool for fishing bioactive compounds out of a complex mixture of structural analogues. *Sci Rep.* 9, 11113.
- Kirchweger, B., Klein-Junior, L.C., Pretsch, D., Chen, Y., Cretton, S., Gasper, A.L., Vander Heyden, Y., Christen, P., Kirchmair, J., Henriques, A.T., Rollinger, J.M., 2022. Azepine-indole alkaloids from *Psychotria nemorosa* modulate 5-HT_{2A} receptors and prevent in vivo protein toxicity in transgenic *C. elegans*. *Front. Neurosci.* 16, 826289.
- Langeder, J., Döring K., Schmietendorf H., Grienke U., Schmidtke M., Rollinger J.M., 2023. ¹H NMR based biochemometric analysis of *Morus alba* extracts towards a multipotent herbal anti-infective. *J. Nat. Prod.* 86(1), 8-17.
- Wasilewicz, A., Kirchweger B., Bojkova D., Abi Saad M., Langeder J., Bütikofer M., Adelsberger S., Grienke U., Cinatl J., Petermann, O., Scapozza L., Orts J., Kirchmair J., Rabenau, H., Rollinger J.M.. Identification of natural products inhibiting SARS-CoV-2 by targeting the viral proteases: a combined in silico and in vitro approach. *J. Nat. Prod.* 86(2), 264-275.
- Wasilewicz A., Bojkova D., Benididir M.A., Cinatl J. Jr, Rabenau H.F., Grienke U., Rollinger J.M., Kirchweger B., 2024. Molecular networking unveils anti-SARS-CoV-2 constituents from traditionally used remedies. *J. Ethnopharmacol.* 117206.
- Zwirchmayr J., Cruz C.D., Grienke U., Tammela P., Rollinger J.M., 2023. Biochemometry Identifies ostruthin as pluripotent antimicrobial and anthelmintic agent from masterwort. *iScience.* 26(9), 107523.

* Presenting author. E-mail address: judith.rollinger@univie.ac.at (Judith M. Rollinger).

Oral contributions

Exploring lipidomic changes: the impact of St. John's wort extract ze 117 and escitalopram in a rat model of depression induced by chronic stress

Hendrik Bussmann^a, Swen Bremer^b, Jürgen Drewe^a, Hanns Häberlein², Sebastian Franken², Virginie Freytag^{c,d}, Georg Boonen^a, Veronika Butterweck^{a*}

^a Max Zeller Soehne AG, Seeblickstrasse 4, 8590 Romanshorn, Switzerland

^b Institute of Biochemistry and Molecular Biology, Medical Faculty, University of Bonn, Nussallee 11, 53115 Bonn, Germany

^c Division of Molecular Neuroscience, Medical Faculty, Department of Biomedicine, Birmannsgasse 8, 4055 Basel, Switzerland

^d GeneGuide AG, Birmannsgasse 8, 4055 Basel, Switzerland

Chronic stress plays a pivotal role in depression development, triggering hyperactivation of the hypothalamic-pituitary-adrenal (HPA) axis and subsequent elevation of glucocorticoids (GC) (Kim et al., 2022). Persistently heightened GC levels disrupt neuroplasticity and impact brain lipid metabolism (Pinto et al., 2022). This study utilized repeated corticosterone injections to induce depression-like behavior in male Sprague-Dawley rats. Rats were subjected to corticosterone injections (40 mg/kg, s.c.) for 22 consecutive days, co-administered with varying doses of the St. John's wort extract Ze 117 (30, 90, or 180 mg/kg, p.o.) or escitalopram (10 mg/kg, p.o.). Behavioral changes were assessed using a modified forced-swim test (FST) (Johnson et al., 2006). Corticosterone injections significantly reduced the latency to first immobility. Co-treatment with Ze 117 dose-dependently and significantly increased this latency to first immobility compared to corticosterone-only treated rats ($p < 0.001$). The lipidome of plasma and hippocampus samples was analyzed by shotgun mass spectrometry to explore the biochemical effects of corticosterone-induced stress and potential counter-regulation by antidepressants. Corticosterone-induced stress altered key lipid metabolites in plasma, while hippocampal samples exhibited changes primarily in glycerophospholipids, such as dose-dependent increases in lysophosphatidylethanolamines (LPEs) induced by Ze 117. These changes correlated significantly with the latency to first immobility. In conclusion, this study unveils significant behavioural and lipidome-modulating effects of Ze 117 and escitalopram in rat plasma and hippocampus, offering novel targets for clinical diagnosis and antidepressant intervention.

Keywords: *Hypericum perforatum*, escitalopram, chronic stress, forced swim test, lipidomics

References

- Kim, I.B., Lee, J.H., Park, S.C., 2022. The Relationship between stress, inflammation, and depression. *Biomedicines*. 10(8), 36009476.
- Pinto, B., Conde, T., Domingues, I., Domingues, M.R., 2022. Adaptation of lipid profiling in depression disease and treatment: a critical review. *Int. J. Mol. Sci.* 23(4), 35216147.
- Johnson, S.A., Fournier, N.M., Kalynchuk, L.E., 2006. Effect of different doses of corticosterone on depression-like behavior and HPA axis responses to a novel stressor. *Behav. Brain Res.* 168(2), 280-288.

* Presenting author. E-mail address: veronika.butterweck@zellerag.ch (Veronika Butterweck).

Proton spin network fingerprints for dereplication, pharmacophore identification and standardization of the botanical supplement *Centella asiatica*

Harianantenaina L. Rakotondraibe^{*a}, Sara Neiheisel^a, Caroline Wittman^a, Luke C. Marney^{b,c,d}, Jaewoo Choi^{c,d}, Liping Yang^{b,d}, Nora Gray^{d,5}, Jan F. Stevens^{c,d,f}, Claudia S. Maier^{b,c,d}, Amala Soumayanath^{d,e}

^a Division of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, The Ohio State University, Columbus, OH, USA^b

^b Department of Chemistry, Oregon State University, Corvallis, OR, USA

^c Linus Pauling Institute, Oregon State University, Corvallis, OR, USA

^d BENFRA Botanical Dietary Supplements Research Center, Oregon Health & Science University, Portland, OR, USA

^e Department of Neurology, Oregon Health & Science University, Portland, OR, USA

^f Department of Pharmaceutical Sciences, Oregon State University, Corvallis, OR, USA

Centella asiatica (Apiaceae) is a botanical dietary supplement that is used in Asia, Madagascar, and the United States to provide resilience to age related cognitive decline. This property has been supported by preclinical evidence including our work on NRF2 activation in HepG2-ARE cells and increased expression of the antioxidant response gene *Nrf2* *in vivo*. Similar to many botanical dietary supplements, lack of consistency of *Centella asiatica* (CA) formulations limits the validation of its medicinal benefits in clinical studies. Efforts have been made to develop liquid chromatography coupled with mass spectrometry (LC-MS) techniques to detect, derePLICATE and characterize metabolites in botanicals using different refinement of ionization techniques and mass libraries. For *C. asiatica*, comprehensive identification of chemical constituents present in active samples is still necessary to complement standardization and quality control efforts by populating mass libraries. Bioactivity has previously been associated with CA's triterpene and caffeoylquinic acid (CQA) and new CQAs are being identified. Our laboratory focuses on development of a method that helps identify proton spin network fingerprint (SNF) of CA metabolites that bear pharmacophores. After being identified, encrypted, and stored in an in-house database, proton SNFs are coupled with mass spectrometric data to accurately identify isomeric metabolites such as CQAs. This presentation will focus on the process of identification of pharmacophore containing chemical markers of *C. asiatica* provided by BENFRA center, by SNF coupled with two-dimensional NMR experiment as well as LC-MS spectrometric analyses. Application of SNF-coupled LC-MS in standardization and quality control will be also discussed.

Keywords: dietary supplement, *Centella asiatica*, proton spin network, LC-MS, standardization

* Presenting author. E-mail address: rakotondraibe.1@osu.edu (Harianantenaina L. Rakotondraibe).

Drug discovery pipeline reveals ellagic acid to reduce forgetting in *Caenorhabditis elegans* through specific Musashi inhibition

Tamara Balsiger^{a,*}, Attila Stetak^{b,c}, Robert Hagmann^a, Kim-Dung Huynh^b, Pablo Solis^d, Matthias Hamburger^a, Andreas Papassotiropoulos^{b,c}, Robin Teufel^a, Eliane Garo^a

^a Division of Pharmaceutical Biology, Department of Pharmaceutical Sciences, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

^b Research Cluster Molecular and Cognitive Neurosciences, Department of Biomedicine, University of Basel, Birmannsgasse 8, 4055 Basel, Switzerland

^c University Psychiatric Clinics, University of Basel, Wilhelm Klein-Strasse 27, 4002 Basel, Switzerland

^d Center for Pharmacognostic Research and Panamanian Flora (CIFLORPAN), Faculty of Pharmacy, University of Panama, Panama City 0801, Republic of Panama

Memory maintenance and forgetting are fundamental processes in our lives. However, increased forgetting due to aging or neurodegenerative conditions (e.g. Alzheimer's disease) dramatically impairs the life quality of millions of people worldwide. A recently discovered protein called Musashi (MSI) has emerged as a crucial player in promoting forgetting (Hadziselimovic et al., 2014). The fact that no efficient treatment for age- or disease-related forgetfulness is available, prompted us to search for new natural product-based MSI inhibitors. Our drug discovery pipeline included the screening of an in-house plant extract library (2576 extracts) with a newly developed MSI inhibitor assay, followed by HPLC-based activity profiling to localize the active compounds in these extracts. Targeted isolation of the active constituents of the *Freziera candicans* Tul. (Pentaphylacaceae) extract has elucidated flavonol glycosides and ellagitannins. While flavonols inhibited MSI at low micromolar concentrations, the series of ellagitannins demonstrated a strong structure-activity relationship, revealing ellagic acid (EA) as by far the most potent MSI inhibitor in the low nanomolar range. EA was further tested in an associative learning and memory experiment in *Caenorhabditis elegans* (*C. elegans*) and remarkably led to a concentration-dependent reduction of forgetting in short- and long-term associative memory tests. Further *in vivo* experiments confirmed EA to act specifically via MSI inhibition. EA was more potent *in vitro* and *in vivo* than the previously published (-)-gossypol (Mastrandreas et al., 2023). Additionally, results on the role of EA in the rescue of Alzheimer symptoms in a *C. elegans* model mediated through MSI inhibition will be presented.

Keywords: memory & forgetting, ellagic acid, Musashi inhibitors, *C. elegans*, Alzheimer disease

References

Hadziselimovic, N., Vukojevic, V., Peter, F., Milnik, A., Fastenrath, M., Fenyves, B., Hieber, P., Demougin, P., Vogler, C., De Quervain, D., Papassotiropoulos, A., Stetak, A., 2014. Forgetting is regulated via musashi-mediated translational control of the Arp2/3 complex. *Cell*. 156, 1153-1166.

Mastrandreas, P., Arnold, A., Boglari, C., De Quervain, D., Stetak, A., Papassotiropoulos, A., 2023. (-)-Gossypol inhibition of Musashi-mediated forgetting improves memory and age-dependent memory decline in *Caenorhabditis elegans*. *Mol. Neurobiol.* 60, 820-835.

* Presenting author. E-mail address: tamara.balsiger@unibas.ch (Tamara Balsiger).

Anti-inflammatory and cytoprotective polypharmacology of BNO 2103 reveals targeting of IKK-NF- κ B and p38-MK2-RIPK1 axes

Marija Milosevic^a, Alexander Magnutzki^a, Shah Hussain^a, Thomas Jakschitz^a, Theodor Braun^a, Bernhard Nausch^b, Martin Kragl^{b,*}, Michael Soeberdt^b, Günther K. Bonn^a, Lukas A. Huber^c, Taras Valovka^c

^a Austrian Drug Screening Institute GmbH, Innrain 66a, 6020 Innsbruck, Austria

^b Bionorica SE, Kerschensteinerstraße 11-15, 92318 Neumarkt in der Oberpfalz, Germany

^c Institute of Cell Biology, Biocenter, Medical University of Innsbruck, Innrain 80/82, 6020 Innsbruck, Austria

Urinary tract infections (UTIs) are among the most frequent infections, affecting approximately 150 million patients per year worldwide. Associated inflammation and tissue damage contribute significantly to symptoms, e.g., dysuria and urge. Canephron, an herbal medicinal product with anti-inflammatory (Künstle et al., 2013, Nausch et al., 2019), spasmolytic (Brenneis et al., 2012), anti-adhesive (Künstle et al., 2013), and anti-nociceptive (Nausch et al., 2019) activities, is approved for treatment of uncomplicated UTIs. To investigate cytoprotective and anti-inflammatory properties of the active component of Canephron N (BNO 2103: mixture of pulverized rosemary leaves, centaury herb, and lovage root), and underlying signaling mechanisms, a newly established NF- κ B-reporter kidney epithelial (HK-2) cell line, luminescent assays (NF- κ B activity), Western blots (signaling pathways and protein phosphorylation), and multiplex-immunoassays (Luminex[®] xMAP[®] technology, cytokine release) were used. BNO 2103 repressed NF- κ B activation (IC_{50} = 241.1 μ g/mL) by *lipopolysaccharide* (LPS) and tumor necrosis factor alpha (TNF α), inhibiting the phosphorylation and degradation of I κ B $\pm$. BNO 2103 also suppressed inflammation-specific S536 phosphorylation of p65 (NF- κ B protein) and transcription of pro-inflammatory cytokine genes. In contrast to other, specific NF- κ B inhibitors, BNO 2103 revealed a strong cytoprotective effect against TNF $\pm$ -induced cytotoxicity. Our data indicate that BNO 2103 acts through the p38 MAPK/MK2 axis, promoting RIPK1 phosphorylation at S320 while repressing S166 autophosphorylation and subsequent activation of RIPK1, which is required for apoptotic and necroptotic responses to TNF α . According to our findings, Canephron exhibits anti-inflammatory and cytoprotective effects mediated by a well-defined and highly specific mechanism.

Keywords: urinary tract, inflammation, NF- κ B, cytoprotection, herbal medicinal product

References

- Brenneis, C., Kuenstle, G., Haunschild, J., 2012. Spasmolytic Activity of Canephron[®] N on the Contractility of Rat and Human Isolated Urinary Bladder: 13th International Congress of the Society for Ethnopharmacology. September 2012, Graz, Austria.
- Künstle, G., Brenneis, C., Haunschild, J., 2013. Efficacy of Canephron[®] N against bacterial adhesion, inflammation and bladder hyperactivity. Eur. Urol. Suppl. 12:e671.
- Nausch, B., Pace, S., Pein, H., Koeberle, A., Rossi, A., Kuenstle, G., Werz, O., 2019. The standardized herbal combination BNO 2103 contained in Canephron N alleviates inflammatory pain in experimental cystitis and prostatitis. Phytomedicine. 60, 152987.

* Presenting author. E-mail address: martin.kragl@bionorica.de (Martin Kragl).

Verification of the medicinal value of *Patriniae Herba* for colon cancer using preclinical models

Huihai Yang^a, Grace Gar-Lee Yue^a, Tao Zheng^a, Kwan-Ho Wong^b, Dandan Hu^a, Man-Ho Tong^a, David Tai-Wai Lau^b, Hung Kay Lee^c, Pang-Chiu Shaw^{a,d}, Clara Bik-San Lau^{a,d,e,*}

^a Institute of Chinese Medicine and State Key Laboratory of Research on Bioactivities and Clinical Applications of Medicinal Plants, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^b Shiu-Ying Hu Herbarium, School of Life Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^c Department of Chemistry, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^d Li Dak Sum Yip Yio Chin R&D Centre for Chinese Medicine, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^e Department of Pharmacology and Pharmacy & School of Chinese Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong SAR, China

Patriniae Herba, referring to the perennial herbs such as *Patrinia villosa* (Thunb.) Juss. and *Patrinia scabiosaeifolia* Link (family Valerianaceae), is a folk medicine for treating gastrointestinal disorders over thousands of years in China. With the anti-tumor activities of *P. villosa* (PV) previously reported (He et al., 2017), our recent studies aimed to elucidate the underlying mechanism of the anti-tumor effects of PV aqueous extract in colon cancer using preclinical models. In order to obtain the genuine herbal material of PV, our research team established a systematic authentication procedure which includes morphology, chemical characterization and DNA markers (Wong et al., 2024). The PV aqueous extract (PVW) was prepared, chemically characterised and used for biological evaluation in colon cancer cell-based and animal studies. Our results showed that PVW exhibited cytotoxicity, apoptosis, cell cycle arrest and inhibitory effects on cell motility and migration in human and murine colon cancer cells via regulating ErbB, focal adhesion, and TGF- β signaling pathways. Furthermore, PVW suppressed tumor growth and metastasis in tumor-bearing mice via modulating the levels of immune cells (cytotoxic T cells and MDSCs) (Yang et al., 2023a). Besides, the compound 8,9-didehydro-7-hydroxydolichodial was firstly proven to be present in PVW and was shown to suppress colon cancer cells migration via increasing expression of tumor suppressor p53 (Yang et al., 2023b), thus it could be considered as an active component.

In conclusion, our findings scientifically verified the therapeutic value of the herb *Patrinia villosa* and provided evidences to support the clinical use in colon cancer patients.

Keywords: *Patrinia villosa*, Valerianaceae, *Patriniae herba*, authentication, colon cancer

References

- He, X., Luan, F., Zhao, Z., Ning, N., Li, M., Jin, L., Chang, Y., Zhang, Q., Wu, N., Huang, L., 2017. The Genus *Patrinia*: a review of traditional uses, phytochemical & pharmacological studies. *Am J Chin Med.* 45(4), 637-666.
- Wong, K.H., Zheng, T., Yue, G.G.L., Li, M.C., Wu, H.Y., Zhao, X.L., Chen, H.B., Lau, C.B.S., Shaw, P.C., Lau, D.T.W., 2024. Authentication of medicinal *Patrinia* species: an integration of morphological, chemical and molecular methods. *Scientific Reports* (under revision).
- Yang, H., Yue, G.G.L., Yuen, K.K., Gao, S., Leung, P.C., Wong, C.K., Lau, C.B.S., 2023a. Mechanistic insights into the anti-tumor and anti-metastatic effects of *Patrinia villosa* aqueous extract in colon cancer via modulation of TGF- β R1-smad2/3-E-cadherin and FAK-RhoA-cofilin pathways. *Phytomedicine.* 117, 154900.
- Yang, H., Zheng, T., Ku, C.F., Ngai, C.K., Yue, G.G.L., Lee, H.K., Lau, C.B.S., 2023b. Discovery of a heat-generated compound DHD derived from *Patrinia villosa* water extract with inhibitory effects on colon cancer cells viability and migration. *Frontiers in Chemistry.* 11, 1195883.

* Presenting author. E-mail address: cbslau@hku.hk (Clara Bik-San Lau).

***Symphytum* genus: novel insights into its phytochemistry and bioactivity**

Adriana Trifan^{a,*}, Monika E. Czerwińska^b, Sebastian Granica^c, Evelyn Wolfram^d, Krystyna Skalicka-Woźniak^e, Simon Vlad Luca^f

^a Department of Pharmacognosy-Phytotherapy, Faculty of Pharmacy, “Grigore T. Popa” University of Medicine and Pharmacy Iasi, Iasi, Romania

^b Department of Biochemistry and Pharmacogenomics, Medical University of Warsaw, Warsaw, Poland

^c Microbiota Lab, Centre for Preclinical Studies, Department of Pharmacognosy and Molecular Basis of Phytotherapy, Medical University of Warsaw, Warsaw, Poland

^d Phytopharmacy and Natural Products Research Group, Zurich University of Applied Sciences, Wädenswil, Switzerland

^e Department of Natural Products Chemistry, Medical University of Lublin, Lublin, Poland

^f Biothermodynamics, TUM School of Life Sciences, Technical University of Munich, Freising, Germany

With around 34 recognized species, the *Symphytum* genus (comfrey), Boraginaceae, has been traditionally used for its anti-inflammatory properties in treating musculoskeletal disorders (Trifan et al., 2020). This multi-methodological study aimed to i) map comfrey species' metabolic complexity using state-of-the-art spectro-chromatographic techniques; ii) isolate main phenolic markers by liquid-liquid chromatography (LLC); iii) assess their anti-inflammatory potential in lipopolysaccharide (LPS)-stimulated human neutrophils; iv) remediate pyrrolizidine alkaloids (PAs) from comfrey extracts using a polymer-based resin. LC-HRMS/MS-based metabolite profiling of roots and aerial parts of underutilized *Symphytum* species identified numerous phenolic acids, flavonoids, PAs, and organic and fatty acids (Luca et al., 2024). Four caffeic acid oligomers (rosmarinic acid, rabsidosiin, globoidans A and B) were isolated from *S. officinale* (comfrey) roots following a LLC approach. The four compounds significantly inhibited IL-1 β release in LPS-stimulated neutrophils, contributing to comfrey's overall anti-inflammatory activity (Trifan et al., 2020). Additionally, a survey across European countries identified these phenolics as phytochemical markers in comfrey roots (Trifan et al., 2021). Lastly, in light of PAs-induced toxicity, a polymer-based resin depletion procedure was employed for their removal from comfrey extracts. PA depletion did not alter the profile of the main phenolic markers while reducing PAs to below 2 ppm. Moreover, PA removal did not affect the inhibitory effects of comfrey extracts on the release of IL-1 β , IL-8, and TNF- α in LPS-stimulated neutrophils (Trifan et al., 2023). Thus, the *Symphytum* genus shows promise for drug development, with PA-depleted extracts being considered for cosmetic and pharmaceutical products.

Keywords: comfrey, Boraginaceae, pyrrolizidine alkaloids, anti-inflammatory

References

- Luca, S.V., Zengin, G., Kulinowski, L., Sinan, K.I., Skalicka-Woźniak, K., Trifan, A., 2024. Phytochemical profiling and bioactivity assessment of underutilized *Symphytum* species in comparison to *S. officinale*. *J. Sci. Food Agric.* 104(7), 3971-3981.
- Trifan, A., Skalicka-Woźniak, K., Granica, S., Czerwińska, M.E., Kruk, A., Marcourt, L., Wolfender, J-L., Wolfram, E., Esslinger, N., Grubelnik, A., Luca, S.V., 2020. *Symphytum officinale* L.: Liquid-liquid chromatography isolation of caffeic acid oligomers and evaluation of their influence on pro-inflammatory cytokine release in LPS-stimulated neutrophils. *J. Ethnopharmacol.* 262, 113169.
- Trifan, A., Wolfram, E., Esslinger, N., Grubelnik, A., Skalicka-Woźniak, K., Minceva, M., Luca, S.V., 2021. Globoidnan A, rabsidosiin and globoidnan B as new phenolic markers in European-sourced comfrey (*Symphytum officinale* L.) root samples. *Phytochem. Anal.* 32, 482-494.
- Trifan, A., Czerwińska, M.E., Zengin, G., Esslinger, N., Grubelnik, A., Wolfram, E., Skalicka-Woźniak, K., Luca, S.V., 2023. Influence of pyrrolizidine alkaloids depletion upon the biological activity of *Symphytum officinale* L. extracts. *J. Ethnopharmacol.* 303, 116010.

* Presenting author. E-mail address: adriana.trifan@umfiasi.ro (Adriana Trifan).

Insight into the mechanism of anti-migraine activity of butterbur (*Petasites hybridus* L.)

Łukasz Kulinowski*, Krystyna Skalicka-Woźniak

Department of Natural Products Chemistry, Medical University of Lublin, 1 Chodzki Str., 20-093 Lublin, Poland

Petasites hybridus (L.) Gaertn., B. Mey. & Scherb (Fig. 1) is a plant of Asteraceae genus. It is the most common butterbur species, which is used in migraine treatment as a clinically approved drug (Petadolex®) (Kulinowski et al., 2022). This presentation is intended to shed a light on mechanism of action of *P. hybridus* extracts and pure compounds in the treatment of migraine. The effectiveness of *P. hybridus* rootstocks extracts is proved in many clinical trials, but the side and mode of action are not fully understood. We will cover the recent reports regarding mode of anti-migraine activity of petasins, which are considering the active components of butterbur extracts.



Fig. 1. *Petasites hybridus* (L.) Gaertn., B. Mey. & Scherb

We will also discuss the pathophysiology and theories of the development of migraine pain, including vascular, neural, trigeminovascular, and the newest molecular theories. We will then consider the effects of butterbur and petasins in the context of those theories, according to recent research on its effects on e.g. TRPA1 and TRPV1 ion channels (Benemei et al., 2017), calcitonin-gene related peptide (CGRP) activity (Kleeberg-Hartmann et al., 2021), and inflammatory mediators (Arnold et al., 2015; Fiebich et al., 2005). In the end we will identify gaps and set the stage for the future research of the activity of butterbur in migraine.

Acknowledgements: This work was financially supported by the National Science Centre, Poland, no 2022/45/N/NZ7/02294.

Keywords: *Petasites hybridus*, butterbur, petasins, anti-migraine, TRP channels, CGRP

References

- Arnold, E., et al. 2015. Inhibition of cytosolic phospholipase A2 α (cPLA2 α) by Medicinal plants in relation to their phenolic content. *Molecules*. 20(8), 15033–15048.
- Benemei, S., et al. 2017. The anti-migraine component of butterbur extracts, isopetasin, desensitizes peptidergic nociceptors by acting on TRPA1 cation channel. *Br. J. Pharmacol.* 174(17), 2897–2911.
- Fiebich, B.L., et al. 2005. *Petasites hybridus* extracts in vitro inhibit COX-2 and PGE2 release by direct interaction with the enzyme and by preventing p42/44 MAP kinase activation in rat primary microglial cells. *Planta Med.* 71(1), 12–19.
- Kleeberg-Hartmann, J., et al. 2021. Petasin and isopetasin reduce CGRP release from trigeminal afferents indicating an inhibitory effect on TRPA1 and TRPV1 receptor channels. *J. Headache Pain.* 22(1).
- Kulinowski, Ł., et al. (2022). A review on the ethnobotany, phytochemistry, pharmacology and toxicology of butterbur species (*Petasites* L.). *J. Ethnopharmacol.* 115263.

* Presenting author. E-mail address: lukasz.kulinowski@umlub.pl (Łukasz Kulinowski).

Study of the prebiotic potential of *Althaea officinalis* L. root extract using the high throughput *ex vivo* technology SIFR^{*}

Pieter Van den Abbeele^a, Stef Deyaert^a, Konstanze Bedal^b, Olaf Kelber^{b,*}, Andreas Hensel^c, Aurélien Baudot^a

^a Cryptobiotix SA, Technologiepark-Zwijnaarde 82, 9052 Gent, Belgium

^b R&D, Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Havelstr. 5, 64295 Darmstadt, German

^c University of Münster, Institute of Pharmaceutical Biology and Phytochemistry, Corrensstraße 48. 48149 Münster, Germany

The extract of *Althaea officinalis* L. root is used in irritations of the pharyngeal mucosa and resulting dry cough, but also in gastrointestinal complaints. While the underlying mucosa protective and anti-inflammatory effects have been studied thoroughly over the last years, data on its effect on the gut microbiota are missing. Given that the extract constituents can be assumed to be not fully absorbed in the small intestine, its influence on the gut microbiota in the large intestine was intended to be studied. The impact of an extract (STW 42, AR) on the gut microbiome was investigated using the high-throughput, *ex vivo* SIFR^{*} technology (Cryptobiotix, Ghent, Belgium). Considering six subjects individually covered the broad spectrum of microbial compositions that occurs *in vivo*, in line with the so-called enterotypes, thus ensuring representative findings. Simulating a single intake, AR was dosed at 1.875 g/d and inulin (IN; used as positive control) was dosed at equivalent of 2 g/day. AR significantly increased total SCFA and significantly decreased pH. While the extent of the effects was lower than with IN, health-related propionate/butyrate was significantly increased, showing a specific effect on butyrate production, which suggests a product-specific impact on the microbiome. Potent treatment effects were noted at higher phylogenetic resolution (OTU level): AR exerted effects including the consistent stimulation of an OTU related to *Ruminiclostridium cellulolyticum*. Overall, the SIFR^{*} study demonstrated the prebiotic potential of marshmallow root extract (AR) by its influence on microbial metabolite production and microbial composition.

Keywords: *Althaea officinalis*, gut microbiota, cough and cold, gastrointestinal complaints

* Presenting author. E-mail address: olaf.kelber@bayer.com (Olaf Kelber).

Development of bio-based active materials for textiles and high-traffic objects

Emilie Stierlin^{a,*}, Job Tchoumtchoua^a, Etienne Maron^a, Caroline Piffet^b, Tamara Oroz^c, Leyre Hernández^c, Marta Mateo^c, Kyriaki-Marina Lyra^d, Zili Sideratou^d, Fotis Katsaros^d, Jean-Michel Thomassin^b

^a Extraction Department, CELABOR Scrl, Avenue du Parc 38, 4650, Herve, Belgium

^b Biomaterials Department, CELABOR Scrl, Avenue du Parc 38, 4650, Herve, Belgium

^c Lurederra Technological Centre, Area Industrial Perguita, Calle A, 31210 Los Arcos, Spain

^d Institute of Nanoscience and Nanotechnology, NCSR “Demokritos”, Aghia Paraskevi Attikis, Athens 15310, Greece

Infectious diseases pose a significant threat to both human health and the global economy, they account for more than 20% of global mortality and viruses are responsible for about one-third of these deaths. To date, we know of about 200 infectious diseases and about 80% of infectious diseases are transmitted by unclean hands touching contaminated surfaces. SUSAAN project is focused on development of new sustainable antiviral and antimicrobial coatings for textiles and high traffic objects made of plastics and metal, involving textile, bathrooms and switches manufacture industries. Rhubarb and horseradish (root and leaves) were both studied within this project due to their rich composition in anthraquinones and isothiocyanate, respectively. Produced extracts were purified and characterized by chromatography (e.g. UPLC-DAD-MS for anthraquinones, GC-MS for isothiocyanates) and their antimicrobial activity was studied against Gram-negative *Escherichia coli* bacteria and Gram-positive *Staphylococcus aureus* bacteria. The obtained antibacterial activities showed that the bio-based extracts exhibit good bacteriostatic activity but are not bactericidal. Therefore, these plant extracts have the potential to be used for the development of bio-based active nanomaterials.

Acknowledgements: This project was funded by the European Union Horizon Europe Programme for Research and Innovation under the Grant Agreement N° 101057988 – SUSAAN.

Keywords: Active biomaterial, natural products, infectious diseases, antimicrobial efficiency

References

Wong, C.S.-A., Lim, C.W., Mohammed H.I., Liew, K.Y., Tham, C.L., Tan, J.W., Chee, H.Y., 2022. Current Perspective of Plant-Based Diets on Communicable Diseases Caused by Viruses: A Mini Review. *Front. Nutr.* 9, 786972.

* Presenting author. E-mail address: emilie.stierlin@celabor.be (Emilie Stierlin).

Macroalgae derived natural product scavengers of reactive aldehyde and carbonyl species and their implications in slowing the progression of prevalent chronic diseases

George S. Hanna^{a,*}, Mark T. Hamann^b

^a Department of Public Health, Medical University of South Carolina, Charleston, SC, 29425

^b Department of Drug Discovery and Biomedical Sciences, Medical University of South Carolina, Charleston, SC 29425

Reactive Aldehyde and Carbonyl Species (RACS) arise from the degradation of essential primary metabolites—sugars and lipids. The resulting stress, derived from their ability to spontaneously react with proteins, DNA, and monoamine neurotransmitters, is a fundamental biological pressure that has driven the evolution of diverse mechanisms for minimizing its impact on organismal health. Relevant to human health, the complications that accompany metabolic disorders, such as diabetes, illustrate what happens when RACS stress exceeds the body's capacity to prevent excessive damage. These include elevated risks for cardiovascular disease, liver disease, kidney disease, neurodegeneration, and cancer, which are accompanied by a heightened vulnerability to infections—illustrating the broad need for targeted interventions. Presented here is a novel quantitative NMR assay for evaluating and screening natural products to discover scavengers of a prominent RACS, methylglyoxal. This tool was employed to explore the chemical diversity of marine macroalgae, with a focus on species that have a history in the diets of the World's healthiest populations and are subject to production by aquaculture. The results revealed the exceptional RACS scavenging capacity and AGE formation inhibition of phlorotannins, a class of molecules abundantly produced by Phaeophytes. Likewise, the guanylureas an unusual class of alkaloids found in Rhodophytes, were found to possess similar activity to the diabetes front line therapy, metformin—evoking questions about their impact on the longevity of populations that regularly consume them. Both illustrate the potential for macroalgal natural products to displace or outright replace widely used drugs and supplements.

Keywords: macroalgae, aquaculture, degenerative disease, aldehydes, diabetes


* Presenting author. E-mail address: hannag@musc.edu (George S. Hanna).

Mitochondrial modulators: the defenders

Emmanuel Makinde*, Linlin Ma, George D. Mellick, Yunjiang Feng

Griffith Institute for Drug Discovery, Griffith University, Qld 4111, Australia

Mitochondria, commonly known as the cell's energy core, are increasingly becoming attractive targets in the search for potent therapeutics against neurodegenerative diseases. Impaired mitochondrial function has been extensively cited as a common hallmark in the pathogenesis of several neurodegenerative diseases, especially Parkinson's disease (PD) (Annesley and Fisher, 2019; Makinde et al. 2023). Given the limitations associated with available treatments for mitochondrial dysfunction-related ailments, the search for new, potent alternative therapeutics has become imperative. In this work, we present the screening of 4224 fractions obtained from 384 Australian marine organisms and plant samples using a modified lead-like fractionation protocol to search for natural products with mitoprotective properties. Primary screening of the fractions using PD patient-sourced olfactory neurosphere-derived (hONS) cells with rotenone as a mitochondria stressor revealed 108 hits from 11 different biota. The potency of these hits was validated in another round of primary screening using human neuroblastoma (SH-SY5Y) cells with 6-hydroxydopamine as a mitochondria stressor. This was followed by a mitochondrial membrane potential assay, resulting in 35 active fractions from 8 biotas. The most active fractions were subjected to LC-Orbitrap MS analysis to identify constituent natural products. A total of 56 natural products were identified, 12 of which have been previously reported to have significant mitoprotective activity. Finally, we also report the isolation, characterization, and testing of metabolites from the top 3 active biotas, leading to the isolation of a new dimeric aptamine compound **1** (Fig. 1)



Bioassay-guided fractionation, identification and docking experiments of α -glucosidase inhibitors from *Geum kokanicum* root

Avishan Farzaneh^a, Somayeh Mojtavavi^b, Houman Kazemzadeh^c, Faezeh Sadat Hosseini^c, Mohammad-Reza Delnavazi^a, Hamidreza Monsef-Esfahani^a, Mohammad Ali Farzamarzi^b, Massoud Amanlou^c, Hamid-Reza Adhami^{a*}

^a Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

^b Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

^c Department of Medicinal Chemistry, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

Geum kokanicum Regel & Schmalh. ex Regel (Rosaceae) is a perennial herb used as a natural remedy for gastric dysfunctions such as diarrhea, dysentery, dyspepsia, and gastroenteritis. Despite some studies reporting the anti-diabetic activity of this plant, bioactive compounds have not been well identified. The present study was designed to isolate bioactive constituents from the root of *Geum kokanicum* with anti-diabetic properties (Farzaneh et al, 2022). The α -glucosidase inhibitory effect of various extracts (chloroform, ethyl-acetate, methanol) were investigated. Regarding a bio-assay guided procedure, the active compounds were isolated using different chromatography methods. The structure of the pure compounds were elucidated by analytical techniques and their IC₅₀ values were determined for α -glucosidase inhibition. In addition, a molecular docking study was used to evaluate the candidates for α -glucosidase inhibitory activity. Furthermore, 3, 3', 4'-tri-O-methylellagic acid (**1**), daucosterol (**2**) from the chloroform and (+) - catechin (**3**), casuarinin (**4**) and ellagic acid (**5**) from the ethyl-acetate fraction were identified. Evaluation of the compounds **1–5** toward α -glucosidase revealed that compounds **4**, **5** and **1** were the most potent and selective inhibitors of α -glucosidase enzyme with the IC₅₀ values of, 1.01 ± 0.30 , 30.00 ± 1.10 and 94.00 ± 0.21 μ M, respectively. Docking study was in harmony with *in-vitro* α -glucosidase inhibitory also revealed that compound **4** presented the lowest docking binding energy (-12.4 kcal/mol) by forming several strong hydrogen bonds in the active site of the α -glucosidase. The results of this study demonstrated the role of phenolic compounds on anti-dabetic activity of *Geum kokanicum* root.

Keywords: anti-diabetes, α -glucosidase, *Geum kokanicum*, casuarinin, ellagic acid

References

Farzaneh, A., Faramarzi, M.A., Delnavazi, M.R., Monsef-Esfahani, H.R., Adhami, H.R., 2022. *In Vitro* Anti-Diabetic and Anti-Oxidant Activities of *Geum* Species from Iran. Research Journal of Pharmacognosy. 9, 37-44.

* Presenting author. E-mail address: hr-adhami@tums.ac.ir (Hamid-Reza Adhami).

Session II: Natural product research and the challenges of climate change – ecological and environmental solutions

Topical Lecture

Marine chemical ecology and natural products research in our changing ocean

Valerie J. Paul

Smithsonian Marine Station, Fort Pierce, FL, USA

Macroalgae and benthic cyanobacteria are becoming increasingly abundant on coral reefs worldwide. The die-off of the sea urchin *Diadema antillarum* in Caribbean waters in the 1980s coincided with a dramatic increase in macroalgal biomass on coral reefs. More recently, marine heat waves are also contributing to increased macroalgal and cyanobacterial abundance on reefs. Algal natural products play important roles as chemical defenses against grazers such as sea urchins and herbivorous fishes. These compounds may also play a role in algal competition with corals and other benthic organisms. Experiments were designed to test interactions between chemically defended species of algae and benthic cyanobacteria and different life history stages of corals. Some extracts and isolated compounds from macroalgae and cyanobacteria negatively influenced the settlement of coral larvae. On reefs experiencing increased abundance of chemically defended macroalgae and benthic cyanobacteria, the rebuilding of coral populations may be inhibited by algal natural products. These chemical interactions are taking place under changing ocean conditions where increases in temperature and carbon dioxide concentrations are leading to changes in pH (ocean acidification) and warming. In contrast, some marine red algae (crustose coralline algae) and bacteria can induce settlement and metamorphosis of coral larvae. Studies of these settlement cues may lead to the development of compounds that are useful in coral restoration. Changing ocean conditions have also contributed to increased incidence of marine diseases, coral bleaching and hypoxia. How compounds in microbes and macroorganisms respond to these changing ocean conditions is an area of ongoing research interest.

Keywords: chemical defense, black band disease, stony coral tissue loss disease, coral settlement, climate change

References

- Gunasekera, S. P., Meyer, J. L., Ding, Y., Abboud, K. A., Luo, D., Campbell, J. E., Angerhofer, A., Goodsell, J. L., Raymundo, L. J., Liu, J., Ye, T., Luesch, H., Teplitski, M., Paul, V. J., 2019. Chemical and metagenomic studies of the lethal black band disease of corals reveal two broadly-distributed, redox-sensitive mixed polyketide/peptide macrocycles. *Journal of Natural Products* 82: 111-121.
- Paul, V. J., Kuffner, I. B., Walters, L. J., Ritson-Williams, R., Beach, K. S. and Becerro, M. A., 2011. Chemically mediated interactions between macroalgae *Dictyota* spp. and multiple life-history stages of the coral *Porites astreoides*. *Marine Ecology Progress Series* 426: 161-170.
- Ritson-Williams, R., Arnold, S. N. and Paul, V. J., 2020. The impact of macroalgae and cyanobacteria on larval survival and settlement of the scleractinian corals *Acropora palmata*, *A. cervicornis* and *Pseudodiploria strigosa*. *Marine Biology* 167:31
- Sneed, J. M., Meickle, T., Engene, N., Reed, S., Gunasekera, S. and Paul, V. J., 2017. Bloom dynamics and chemical defenses of benthic cyanobacteria in the Indian River Lagoon, Florida. *Harmful Algae* 69: 75-82.
- Ushijima, B., Gunasekera, S. P., Meyer, J. L., Tittl, J., Pitts, K., Thompson, S., Sneed, J., Ding, Y., Chen, M., Houk, J., Aeby, G., Häse, C. C., Paul, V. J., 2023. Chemical and genomic characterization of a potential probiotic treatment for stony coral tissue loss disease. *Communications Biology* 6:248.

* Presenting author. E-mail address: paul@si.edu (Valerie J. Paul).

Oral contributions

Search for anti-phytopathogenic compounds by studying biotic interactions between *Fusarium* sp. and the endophytic fungi of the palm tree model *Astrocaryum sciophilum*

Véronique Eparvier^{a,*}, Didier Stien^b, Jean-Luc Wolfender^{c,d}, David Touboul^e

^a Université Paris-Saclay, Institut de Chimie des Substances Naturelles, ICSN UPR2301, Avenue de la Terrasse, 91190 Gif-sur-Yvette, France

^b Sorbonne Université, Centre National de la Recherche Scientifique (CNRS), Laboratoire de Biodiversité et Biotechnologie Microbiennes, Laboratoire de Biodiversité et Biotechnologies Microbiennes (LBBM), Observatoire Océanologique, Banyuls-Sur-Mer, France

^c School of Pharmaceutical Sciences, University of Geneva, Centre Médical Universitaire (CMU), Geneva, Switzerland

^d Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, Centre Médical Universitaire (CMU), Geneva, Switzerland

^e Laboratoire de Chimie Moléculaire (LCM), CNRS UMR 9168, École Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France

Biopesticides from living microorganisms or compounds from natural substances are used for biological control as alternative pest management. In order to find microorganisms that naturally possess weapons against phytopathogens, we investigate the biological and chemical diversity of plant leaf endophytes. Endophytes are microorganisms living in association with plants and can improve host plants resistance against biotic stresses, especially phytopathogens aggressions (Arnold et al. 2003). The palm *Astrocaryum sciophilum* is the host plant model chosen in this work. Due to the longevity and the easy dating of its leaves (Charles-Dominique et al., 2003), we expect to find a highly competitive and diverse microbial community. 197 endophytes have been isolated from eight palm specimens. Firstly, we selected a population of 42 fungi species from the *Colletotrichum* genus due to their extracts exhibiting biological activities relevant to their role within our macroscopic host. Strains were classified using MALDI-ToF-MS lipid and protein fingerprinting along with chemotaxonomic network analysis. Using protein extracts, rapid chemotaxonomic classification correlated distinct chemotypes with observed biological activities. The molecules accountable for the observed activities across different chemotypes were annotated, isolated, and characterized (Barthélemy et al., 2020). Secondly, we focus on the chemical study of a *Xylaria* sp. strain, whose extract exhibited activity against *F. oxysporum*. Through comparison of the LC-MS/MS profiles of the extracts and employing molecular networking, we identified metabolites from both strains whose production is induced by competition. Additionally, we observed the biosynthesis of these compounds over time during competition using MS imaging (Barthélemy et al., 2021).

Keywords: endophyte, competition, phytopathogen, metabolites

References

- Arnold, A.E., Mejía, L.C., Kylo, D., Rojas, E.I., Maynard, Z., Robbins, N., Herre, E.A., 2003. Fungal endophytes limit pathogen damage in a tropical tree. *Proc Natl Acad Sci U S A*. 100, 15649-15654.
- Charles-Dominique, P., Chave, J., Marc-A. Dubois, De Granville, J.-J., Riera, B., & Vezzoli, C., 2003. Colonization Front of the Understorey Palm *Astrocaryum sciophilum* in a Pristine Rain Forest of French Guiana. *Global Ecol. Biogr.*, 12, 237–248.
- Barthélemy, M., Guérineau, V., Genta-Jouve, G., Ropy, M., Guillot, R., Pelissier, L., Wolfender, J.-L., Stien, D., Eparvier, V., Touboul, D., 2020. Identification and dereplication of endophytic *Colletotrichum* strains by MALDI-TOF Mass Spectrometry and Molecular Networking, *Sci. Rep.* 10, 19788.
- Barthélemy, M., Elie, N., Genta-Jouve, G., Stien, D., Touboul, D., Eparvier, V., 2021 Identification of antagonistic compounds between the palm tree endophytic fungus of BSNB-0294 (*Xylaria* sp.) and the phytopathogen *Fusarium oxysporum*. *J. Agri. Food Chem.* 69, 10893-10906.

* Presenting author. E-mail address: veronique.eparvier@cnrs.fr (Véronique Eparvier).

Using natural, greener, biodegradable and recyclable solvents to enhance lignin extraction from Acacia wood

Carla Varela^{a,b,c,*}, Catarina Fernandes^{a,d}, Solange Magalhães^a, Bruno Medronho^{d,e}, Luís Alves^a, M. Graça Rasteiro^a

^a CERES, Department of Chemical Engineering, University of Coimbra, 3030-790 Coimbra, Portugal

^b iCBR/CACC/FMUC, University of Coimbra, 3000-548 Coimbra, Portugal

^c CIBB, University of Coimbra, Portugal

^d MED – Mediterranean Institute for Agriculture, Environment and Development & CHANGE – Global Change and Sustainability Institute, University of Algarve, 8005-139 Faro, Portugal

^e FSCN, Surface and Colloid Engineering, Mid Sweden University, Sundsvall, Sweden

The increasing interest in sustainable materials has motivated the adoption of a circular economy approach, where residues are treated as useful resources to produce new materials of added value. In this respect, lignin has gained significant attention as an alternative renewable feedstock due to its abundant availability and broad range of interest and applications. However, its extraction from biomass is a challenging task given its complex structure. Among the different extraction methods available, deep eutectic solvents (DES) have shown promising results. Enhanced selectivity and extraction performance have been reported when using ternary mixtures with tuned properties (Fernandes et al., 2021). These mixtures present high solubility towards lignin and low cellulose dissolution making them highly selective and yielding high purity lignin. In addition, DES can be prepared with natural ingredients and be easily recovered at the end of the process with further reuse without compromising the extraction efficiency. New binary (lactic acid:choline chloride) and ternary DES mixtures (lactic acid:citric acid:choline chloride) for lignin extraction from Acacia wood were tested and are covered in this presentation. Not only DES physicochemical properties but also their extraction capacity was greatly affected by their composition and extraction conditions. As for DES recovery and reuse, it was observed that it did not negatively affect its efficiency or selectivity, up to 3 cycles of re-use, leading only to a small decrease in lignin's purity (~80%) (Figure 1). Therefore, the use of a natural-based and recyclable DES makes this process highly appealing for biomass fractionation with low environmental impact.




Fig. 1. Extraction yield and purity of freshly synthesized and reused DES.

Keywords: natural DES, greener wood extraction, recycling and reuse

Funding: This work was financially supported by the Portuguese Foundation for Science and Technology (FCT) via the Research and Technological Development Project (2022.06810.PTDC), the research PhD grant (2021.05991.BD) and the Strategic Research Centre Project (UIDB/00102/2020).

References

Fernandes, C., Melro, E., Magalhães, S., Alves, L., Craveiro, R., Filipe, A., Valente, A.J.M., Martins, G., Antunes, F.E., Romano, A., Medronho, B., 2021. New deep eutectic solvent assisted extraction of highly pure lignin from maritime pine sawdust (*Pinus pinaster* Ait.). *International Journal of Biological Macromolecules* 177, 294-305.

* Presenting author. E-mail address: carlalvarela@gmail.com (Carla Varela).

Development of biobased films from ‘Annurca’ apple cellulose dopped with phenolic compounds with novel characteristics

Sarmad Ahmad Qamar^{a,b,*}, Simona Piccolella^a, Yi Zhang^b, Severina Pacifico^a

^a Department of Environmental, Biological & Pharmaceutical Sciences and Technologies, University of Campania ‘Luigi Vanvitelli’, Via Vivaldi 43, 81100 Caserta, Italy

^b Department of Food Science, The Pennsylvania State University, University Park, PA, 16802, USA

The circular biomanufacturing of biodegradable and bioactive materials using agro-food byproducts is being promoted in food sector applications. Herein, the use of ‘Annurca’ apple waste is considered for the first time to introduce cellulose-based blended films using polyphenolic compounds to improve their bioactive functionality. The current strategy demonstrates ultrasound-assisted ethanolic extraction of bioactive compounds followed by the extraction of crystalline cellulose from the leftover biomass. The conversion of ‘Annurca’ cellulose into carboxymethyl cellulose (CMC) was conducted and processed for the development of films using casting technique. For this purpose, a combination of plasticizer (both glycerol and sorbitol, with 40% (w/w) individual contents) was used and heated for 1 h at 40–50°C. Polyphenolic extract (0.02%, w/w) was used in the casting solution to improve the functional properties of resulting films. Further, a control film was prepared without the addition of polyphenolic compounds. Both films were analyzed using Fourier transform infrared spectroscopy (FTIR). For visual appearance stereomicroscopic images was used, while the advanced surface morphological analysis was performed by scanning electron microscopy (SEM). Morphological analyses indicated crystalline nature of cellulose and CMC, while the resulting films were transparent and smooth with a uniform distribution of polyphenolic compounds. FTIR findings further supported that polyphenolic compounds are well dispersed in ‘Annurca’ CMC matrix, and no chemical reactions occurred during the film formation. Further, the newly developed film incorporating bioactive compounds showed excellent antioxidant and antimicrobial characteristics against both gram-positive and gram-negative bacteria. In conclusion, the present study describes a facile method to prepare ‘smart’ multifunctional biobased materials for different bio-sector applications.

Keywords: circular biomanufacturing, biobased materials, antioxidant, antimicrobial, waste utilization, bioactive compounds

* Presenting author. E-mail address: sarmad.qamar@unicampania.it (Sarmad Ahmad Qamar).

Potential and barriers of triterpenoid saponins as tomorrow's plant based bio-pesticides

Søren Bak*, Jan Günther, Pablo D. Cárdenas, Malboro Dervishi, Jincheng Shen, Rajeswari Gopal Geetha, Anja Fuglsang, Nina Cedergreen, Hans Chr. Bruun Hansen

University of Copenhagen, Department of Plant and Environmental Sciences, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark

Synthetic pesticides have increased yields, but come with a high cost for the environment, human health, and biodiversity. At the same time, climate change is leading to increased food insecurity, making crops more susceptible to fungal diseases and agricultural insect pest. Plant pests and diseases account for a reduction of between 20 and 40% of global crop yields per year. This calls for novel sustainable alternatives to the current use of synthetic pesticides for both conventional and organic farming. There are only a few biocontrol crop protections available on the market, with less than 5% of plant protection products sold worldwide being biocontrol agents. We can work with, and learn from, nature and find bio-solutions. Plants have co-evolved with biotic stress and therefore have the ability to defend themselves, typically by producing bioactive compounds. We have been focusing on how some triterpenoid saponins convey resistance to insect pest and others. Saponins are amphipathic molecular present in more than 100 plant families. While toxicity to humans is regarded as low, they have an untapped potential against plant pests. We are conducting mode-of-action studies to understand their toxicity at the molecular and at the ecosystems level, by combining 2D and 3D structure analysis with bioactivity assays to uncover structure-activity-relationships. Finally, we determine targeted vs non-target organism's responses to explore the potential of saponins as tomorrow's biocontrol agents.

References

- Günther, J, Erthmann, PØ, Bekzod Khakimov, Bak, S, 2022. Reciprocal mutations of two multifunctional β -amyrin synthases from *Barbarea vulgaris* shift α/β -amyrin ratios, *Plant Physiology*, 188: 1483-1495
- Liu Q, Khakimov B, Cárdenas PD, Cozzi F, Olsen CE, Jensen KR, Hauser TP, Bak S, 2019. The cytochrome P450 CYP72A552 is key to production of hederagenin-based saponins that mediate plant defense against herbivores. *New Phytol.* 222:1599-1609.
- Khakimov B, Poulsen VK, Erthmann PØ, Fukushima EO, Augustin JM, Olsen CE, Scholtalbers J, Volpin H, Andersen SB, Hauser TP, Muranaka T, Bak S, 2016. Identification and genome organization of saponin pathway genes from a wild crucifer, and their use for transient production of saponins in *Nicotiana benthamiana*. *The Plant Journal.* 84:478-490.
- Augustin JM, Kuzina V, Andersen SB, Bak S, 2011. Molecular activities, biosynthesis and evolution of triterpenoid saponins. *Phytochemistry.* 72:435-57.

* Presenting author. E-mail address: bak@plen.ku.dk (Søren Bak).

Purification of derivatives of 2-amino-3H-phenoxazin-3-one using CPC

Cristina Díaz-Franco^{a,*}, Carlos Rial^a, Stefan Schwaiger^b, José. M. G. Molinillo^a, Rosa M. Varela^a, Francisco A. Macías^a

^a Allelopathy Group, Department of Organic Chemistry, Institute of Biomolecules (INBIO), Campus de Excelencia Internacional (ceiA3), School of Science, University of Cadiz, Spain

^b Institute of Pharmacy/Pharmacognosy and Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, Innrain 80-82, 6020 Innsbruck, Austria

Aminophenoxazinones represent natural products, originating from a plant produced precursor (benzoxaziones), produced by members of the Poaceae family, which are further metabolized by microorganisms to form the additional aromatic ring. Among all identified downstream metabolites, this compound class has shown multiple biological activities, like a pronounced phytotoxic activity (Macías et al., 2005), via a novel mode of action, identified as HDAC inhibition. Additionally, they act as anti-inflammatories, anticancer agents, antibiotics, and immunologic regulators, to mention some (Zorrilla et al., 2021). To improve their physio-chemical properties and bioactivities, we synthesized derivatives of the natural product 2-amino-3H-phenoxazin-3-one (Figure 1), introducing different heteroatoms and functional groups to the scaffold at the carbons C-1, C-9, C-7, or C-8.




Fig. 1. Aminophenoxazinone scaffold.

Despite successful synthesis, purification of the reaction mixtures was challenging due to strong compound interactions with solid phases. We addressed this using Centrifugal Partition Chromatography (CPC). CPC (Pauli et al., 2008) is a support-free liquid–liquid chromatographic technique with mobile and stationary phases consisting of two immiscible liquids forming a two- phase solvent system, where compounds are distributed and transferred between the two phases according to their corresponding partition coefficients. The stationary phase is maintained in the rotor by centrifugal forces while the mobile phase is pushed through it via a pump. Because of the absence of solid support, CPC allows the full recovery of the sample. With this technique, we were able to purify different derivatives.

Keywords: allelopathy, natural products, CPC, aminophenoxazinones

References

- F.A. Macías, D. Marín, A. Oliveros-Bastidas et al. 'Structure–Activity Relationships (SAR) Studies of Benzoxazinones, Their Degradation Products and Analogues. Phytotoxicity on Standard Target Species (STS)' J. Agric. Food Chem., vol 53, pp 538-548, 2005, doi: 10.1021/jf0484071
- J. G. Zorrilla, C. Rial, D. Cabrera et al. 'Pharmacological Activities of Aminophenoxazinones' Molecules, vol. 56, pp 3453, 2021, doi:10.3390/molecules26113453
- G.F. Pauli, S.M. Pro and J.B. Friesen, 'Countercurrent Separation of Natural Products' J. Nat. Prod., vol. 7, pp 1489- 1508, 2008

*Presenting author. E-mail address: cristina.diaz@uca.es (Cristina Diaz-Franco).

Leveraging phytochemical variation in *Piper scintillans* Trel. to discover novel flavanone meroterpenes

Casey S. Philbin^{a,b,*}, Megan Burroughs^{a,b}, Lora A. Richards^{b,c}, Trevor M. Faske^c, Lee A. Dyer^{b,c}, Thomas L. Parchman^{b,c}, Christopher S. Jeffrey^{a,b}

^a Department of Chemistry, University of Nevada Reno, Reno, NV, USA

^b Hitchcock Center for Chemical Ecology, University of Nevada Reno, Reno, NV, USA

^c Department of Biology, University of Nevada Reno, Reno, NV, USA

Ecology-driven approaches to natural products discovery leverage phytochemical variation resulting from environmental and genetic factors to uncover structurally distinct novel molecules. The *Piper* genus produces phytochemicals from a diverse array of chemical classes that have varied biological activities and traditional uses (Salehi et al., 2019), and is a model genus for studying chemically mediated ecological interactions (Dyer and Palmer, 2004). A genomics and metabolomics survey of 100 *Piper scintillans* Trel. parent-offspring sets led to the discovery of a small group of genetically related individuals that produce a novel dioxane-linked flavonoid meroterpene, raduladioxanolide (**1**).




Fig. 1. Discovery of flavanone meroterpene raduladioxanolide from population-level phytochemical survey of *P. scintillans*.

Leaf cuttings and seeds were collected from parent plants within La Selva Biological Station, Costa Rica (Fig. 1A) and offspring were propagated from seed in a common garden. Foliar tissue from parent plants and their adult offspring were analyzed by UPLC-IM-QTOF and crude ¹H NMR. Cluster analysis revealed three distinct chemotypes (Fig. 1B), one of which (Fig. 1B, CT 2) contained a subset of parent-offspring “families” that produced a set of flavanone meroterpenoids including **1** (Fig. 1C). Absolute stereochemistry at flavanone C2 of **1** remains unknown but IM data suggest a 2:1 diastereomeric ratio (CCS = 215 Å², 240 Å²). We propose that raduladioxanolide results from the oxidative cyclization of sternbin and americanolide G as both were solely isolated from individuals that produce **1**. We have demonstrated how leveraging phytochemical variation within a plant species can reveal unexpected and structurally unique molecules, highlighting new avenues for the discovery of novel natural product architectures.

Keywords: piper, meroterpenes, flavonoids, metabolomics, chemical ecology

References

- Salehi, B.; Zakaria, Z.A.; Gyawali, R.; Ibrahim, S.A.; Rajkovic, J.; Shinwari, Z.K.; Khan, T.; Sharifi-Rad, J.; Ozleyen, A.; Turkdonmez, E.; et al., 2019. Piper Species: A Comprehensive Review on Their Phytochemistry, Biological Activities and Applications. *Molecules*. 24, 1364.
- Dyer, L.A. and Palmer, A.D. eds., 2004. *Piper: a model genus for studies of phytochemistry, ecology, and evolution* (Vol. 25). New York: Kluwer academic/Plenum publishers.

* Presenting author. E-mail address: cphilbin@unr.edu (Casey S. Philbin).

Invasive weeds: a treasure chest of medicinal chemicals from Palmer amaranth (*Amaranthus palmeri*)

Katherine M. Warpeha*

Department of Biological Sciences, University of Illinois at Chicago, 900 S Ashland Ave, Chicago, IL, USA

This presentation addresses the surprising finding that abundant chemicals of the weed Palmer Amaranth (*Amaranthus palmeri* L.) have medicinal utility. Palmer amaranth is an annual weed that threatens the yield of many crops in the USA, a plant native to the southwest and Mexico (Sauer, 1957). This study originally targeted herbicide development. It was discovered that under specific growth conditions and environmental challenges associated with global change, Palmer amaranth produces numerous chemicals, especially phenylpropanoids, useful for medicinal purposes, with some similar chemical types of our prior work in broccoli (Gurgul et al. 2023). With recent global change, Palmer amaranth has become a significant threat to crop yield, especially soybean. Metabolomics and lipidomics studies were conducted on specific growth stages of Palmer amaranth under stress to look for any chemicals of interest. Dozens of identified structures (reported in other unrelated plants) are in abundance that show promise for drug and nutraceutical development. Data indicate a living drug factory, despite global change, where controlled growth may be possible to benefit humankind.

Keywords: phenylpropanoids, Palmer amaranth, medicinal, weeds, stress

References

Sauer, J.D. 1957. Recent migration and evolution of the dioecious amaranths. *Evolution* 11:11–31.

Gurgul, A.A., Najjar, Y., Chee, A. An H., Che C-T., Park T.J., Warpeha K.M. 2023. Phenylpropanoid-enriched broccoli seedling extract can reduce inflammatory markers and pain behavior. *J. Transl. Med.* 21, 922.

* Presenting author. E-mail address: kwarpeha@uic.edu (Katherine M. Warpeha).

Ecotoxicity studies of natural sesquiterpene lactones on *C. elegans* as soil model organism for potential bioherbicides

Francisco J. Rodríguez-Mejías^{a,b,*}, Stefan Schwaiger^b, Minh Bui Hoang^b, Mark J. Horgan^b, Alexander K.H. Weiss^c, Francisco A. Macías^a

^a Allelopathy Group, Department of Organic Chemistry, Institute of Biomolecules (INBIO), Campus CEIA3, School of Science, University of Cádiz, C/ República Saharaui, 7, 11510, Puerto Real (Cádiz), Spain

^b Institute of Pharmacy/Pharmacognosy, Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, CCB, Innrain 80/82, 6020, Innsbruck, Austria

^c Research Institute for Biomedical Aging Research, University of Innsbruck, Rennweg 10, A-6020 Innsbruck, Austria

Sesquiterpene lactones (SLs) and their derivatives are recognized as potential bioherbicides for controlling weed growth in crop plantations intended for human consumption (Macías et al., 2019). This approach, rooted in natural products, offers an alternative to conventional herbicides, which often comprise persistent soil pollutants detrimental to water ecosystems. Various SLs have demonstrated efficacy as pre- and post-emergent bioherbicides against a range of monocot and dicot weeds (Macías et al., 2020). Additionally, their effectiveness against parasitic plants has been documented. However, research regarding their impacts on soil fauna remains limited.




Fig. 1. Relevant phytotoxic sesquiterpene lactones

Among numerous model organisms, *C. elegans* stands out as a particularly compelling nematode for analysis due to its ubiquitous presence in temperate soils, transparency, and approximate length of 1000 μm (Brenner, 1974). This presentation will delve into toxicity studies of various SLs families—eudesmane, guaiane, germanacrane, and dimers—on the *C. elegans* N2 ancestor strain. Evaluation will encompass toxicity assessments on both adult worms and eggs, alongside examination of degradation products post-bioassay to identify optimal candidates for bioherbicide applications. Costunolide, dehydrocostuslactone, and santamarine exhibited significant inhibition of *C. elegans* motility at 1000 μM , while showing no discernible effects at 300 μM , an advantageous trait considering their bioherbicidal properties, where they inhibit wheat coleoptile elongation by approximately 90% compared to the negative control at 100 M. Moreover, these compounds undergo rapid degradation within 24 hours, boasting a half-life of approximately 12 hours in culture media, thus mitigating environmental pollution risks.

Keywords: sesquiterpene lactones, *C. elegans*, ecotoxicology, phytotoxicity, agrochemistry

References

- Brenner, S., 1974. The genetics of *Caenorhabditis elegans*. *Genetics* 77, 71–94. <https://doi.org/10.1093/genetics/77.1.71>
- Macías, F.A., Durán, A.G., Molinillo, J.M.G., 2020. Allelopathy: The Chemical Language of Plants, in: *Progress in the Chemistry of Organic Natural Products*. Springer, Berlin/Heidelberg, Germany, pp. 1–84.
- Macías, F.A., Mejías, F.J., Molinillo, J.M., 2019. Recent advances in allelopathy for weed control: from knowledge to applications. *Pest Manag. Sci.* 75, 2413–2436.

* Presenting author. E-mail address: javi.rodriguezmejias@gm.uca.es (Francisco J. Rodríguez-Mejías).

Session III: Regulation for the use of natural products in food and drugs

Topical Lecture

Three-pronged strategy of TCM quality control: deep research, holistic standard and precision testing

De-an Guo*

National Research Center for TCM Standardization Technology, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China

It is of a tremendous challenge for the quality control of traditional Chinese medicines due to its extreme complexity largely caused by its multicomponent feature. TCM contains hundreds or even thousands of chemical components in a single herb, not to mention the compound formulas composed of a handful of even tens of herbs. This generates gigantic hurdle to clarify its chemical composition, which is the fundamental basis for the research of quality control, mechanism, pharmacological and toxicological actions, PK/PD, clinical trial, new drug discovery, etc. As to these challenges, we developed a three-pronged strategy dubbed “Deep Research, Holistic Standard and Precision Testing” to have intended a delivery of an ultimate solution. By implementing a quality-oriented deep research, mainly a integrative approach with chemical, metabolic and biological analysis namely “trinity analysis”, the quality markers will be clarified to facilitate the ensuing quality standard elaboration. Based on the comprehensive analysis, a holistic quality control model for herbal quality was developed and practiced in herbal drugs and their products therefrom such as dispensing granules, classic formulas and finished combination products. This type of research cemented for elaboration of quality standards. Due to the deficiencies of the developed quality standards, it is of great necessity to develop the cutting-edge precision testing techniques to compensate the inadequacy of quality standards to fully control the quality. In the past several decades, our research team has devoted to the herbal quality control by practicing the three-pronged approach. A set of herbal drugs, granules and the finished products were researched in depth, which will be exemplified in the lecture.

Keywords: Traditional Chinese medicines, quality control, trinity analysis, three-pronged strategy

* Presenting author. E-mail address: daguo@simmm.ac.cn (De-an Guo)

Oral contributions

Herbal Health Products in Europe – an overview on challenges and chances

Bernd Roether^{a,b,*}

^a BIONORICA SE, Germany

^b Society of Medicinal Plants and Natural Products Research (GA)

Herbal Health Products (HHP) account for more than 20% of the OTC market. HHP can be assigned to different product categories like medicinal products, botanicals, substance-based medical devices, cosmetics. Concurrently all these categories have to comply with a different legal base. The ongoing revision of the EU pharmaceutical legislation as well as the dissensus of European Commission and European Parliament not to implement Regulation (EC) No 1924/2006 on nutrition and health claims made on foods might have great impact on the market. On 20 February this year GA hosted an EU Summit on Herbal Health Products (HHP) in Brussels, bringing politics, industry and academia together to exchange minds on how a level playing field for these kind of products might be restored to guarantee reliable information and safe use to consumers and patients. This talk will highlight relevant issues and summarize essentials.

Keywords: (traditional) herbal medicinal products, food supplements, botanicals, EU Summit on Herbal Health Products

References

Directive 2001/83/EC of 6 November 2001 on the Community code relating to medicinal products for human use.

European Commission (2011). Memo/11/868: Questions and Answers on the list of permitted Health Claims. Brussels, 5 December. Available via: https://ec.europa.eu/commission/presscorner/detail/en/MEMO_11_868.

European Commission (2012). Discussion Paper on Health Claims on Botanicals used in Foods. European Commission (2012). Memo/12/346: Questions and answers on the list of permitted Health Claims on food products. Brussels, 16 May. Available via https://ec.europa.eu/commission/presscorner/detail/en/MEMO_12_346.

Regulation (EC) No 1924/2006 of the European Parliament and of the Council of 20 December 2006 on nutrition and health claims made on foods (Consolidated version: 13 December 2014).

Regulation (EC) No 1925/2006 of 20 December 2006 on the addition of vitamins and minerals and of certain other substances to foods.

* Presenting author. E-mail address: bernd.roether@bionorica.de (Bernd Roether)

Understanding psilocybin mushrooms and aiding in setting standards for a burgeoning industry in Jamaica

Rupika Delgoda^{a,*}, Kimberley Foster^b, Isaac Morrison^b, Marshall Tyler^b, William Irvine^a, Lavonne Wilson^a, Babumon M. Thankappan^a, Shemar Daniel^b

^a Caribbean Centre for Research in Bioscience, (formerly Natural Products Institute), University of the West Indies, Mona, Jamaica

^b Future Wellness, 2 St. George's Close, University of the West Indies, Mona, Jamaica

Psychedelic fungi are experiencing a surge in interest in recent years, particularly among patients under treatment for a variety of mental health issues. These include depression, anxiety disorder, addiction, post-traumatic stress disorder, and various psychiatric treatments within group or individualized therapy sessions. 'Magic mushrooms' biosynthesize the alkaloid psilocybin, which undergoes dephosphorylation following ingestion to its bioactive counterpart, psilocin. The expression of the tryptamine content (determined using HPLC) in 13 psilocybin containing species (within 3 genera) were explored in this study, under numerous growth conditions, including solid/liquid media, varying light/dark conditions, with and without additives, along with optimum conditions determined for drying and storage, following examinations of growth curves. Optimized growth conditions for *Psilocybe cubensis*, the most commonly cultivated species, confirms that an additive layer of gypsum and casing layer provides maximum yield (896.6 g/kg of dried substrate), with a biological efficiency of 89.6%, while also maintaining high total tryptamine expressions (>0.95%). Emanating fruiting bodies from this unique, legal, EU-GMP certified lab, has subsequently been encapsulated and tested (for mycotoxins, microbes, moisture, heavy metals, pesticides and stability) for use by medical practitioners in Jamaica. This presentation will detail our journey and further outline on-going biological activity investigations including the impact on the activities of heterologously expressed human microsomal Cytochromes P450 enzymes.

Keywords: psilocybin, psychoactive, cytochromes P450, mental health, drug interactions

* Presenting author. E-mail address: thejani.delgoda@uwimona.edu.jm (Rupika Delgoda)

Session: Animal Health Care (VET)

Topical Lecture

Nature's Pharmacy" in the context of insects – is there anything to be learned?

Michael Boppré

Forstzoologie und Entomologie, Albert-Ludwigs-Universität, 79085 Freiburg i.Br., Germany

Secondary natural products, plant secondary metabolites (PSMs) and those synthesised by animals, play an extraordinary role not only in all ecosystems, but also directly for humans. For millennia, we have been using many of these non-nutritional chemicals from what is often called "Nature's Pharmacy". When individuals suffering from health problems (e.g., parasites, infectious diseases) ingest compounds with medicinal effects through a specific behavioural activity which is independent of their daily needs and primary metabolism, and serves to cure or prevent infections or attacks by parasites or support well-being, it is usually referred to as "self-medication". This is distinct from "compensatory feeding", which involves nutritive substances and minerals.

Other animals also take advantage of Nature's Pharmacy by deliberately gathering compounds from particular natural sources with a specific behaviour that is distinct from the general activities of the species, independent of its nutritional needs and not essential for living. This behaviour may be related to a health problem, but it may also be independent of health status and serve to increase chances of survival and biological fitness in various ways. It is sometimes referred to as "pharmacophagy".

This presentation explores the commonalities and differences in the use of Nature's Pharmacy for self-medication *s.l. versus* pharmacophagy by humans, vertebrate wildlife, and especially insects. By adding ecological contexts to the discussion (including population ecology and sensory physiology), it attempts to clarify conceptual and terminological issues, identify gaps in general understanding, and suggest key aspects of future research.

Keywords: herbal medicine, self-medication, pharmacophagy, physical fitness, biological fitness, chemical defense

* Presenting author. E-mail address: boppre@fzi.uni-freiburg.de (Michael Boppré)

Oral contributions

Gut-health histomorphometry in fattening lambs fed with hazelnut skin and linseed

Talal Hassan^{a,*}, Elena Diaz Vicuna^a, Dorotea Ippolito^b, Francesca Barroero^a, Muhammad Irfan Malik^a, Martino Musati^c, Antonio Natalello^c, Claudio Forte^a, Maria Teresa Capucchio^a

^a Department of Veterinary Sciences, University of Turin, 10095 Grugliasco, Italy

^b National Institute of Health, Department of Food Safety, Nutrition and Veterinary Public Health, 00161 Roma, Italy

^c Department of Agriculture, Food and Environment, University of Catania, 95123 Catania, Italy

Agro-industrial byproducts in animal nutrition offer animal feed that is both economical and environmentally friendly (Vastolo et al., 2022). In particular, hazelnut skin (*Corylus avellana*), due to the phenolic compounds and oleic acid content, may affect the rumen microbiota and polyunsaturated fatty acids (PUFA) biohydrogenation, potentially modifying the rumen fermentation and morphology (Priolo et al., 2021). Moreover linseed (*Linum-usitatissimum*) has been demonstrated to be capable of increased accumulation of PUFA in meat (Andrés et al., 2014). To evaluate the potential effects of these natural products, 40 male lambs were randomly assigned to four dietary treatments: conventional (C), 8% linseed (LS), 15% hazelnut skin (HS), and a combination (7.5% HS + 4% LS) as a partial replacement of maize in the diet. The lambs (n=20) were assessed for growth performance, and ruminal/gut histomorphometry after a 60- day feeding period. The growth performance did not significantly differ among the groups. Ruminal ventral sac papillae were lengthy (0.508 ± 0.143 cm, $p < 0.05$) and exhibited strong positive association (0.610^{**} , $p < 0.04$) with HS+LS. The HS showed somewhat higher intestinal inflammation scores (3.275 ± 0.568 , $p < 0.05$) than the control group. These results would be fortified by the on-going rumen microbiota and VFA analysis. The pro-inflammatory effects by HS need further investigations to identify their mechanism. In summary, HS+LS/HS seem to have an impact on gut health and particularly on rumen histomorphometry. These findings emphasize the need for further assessment of new feed components to improve animal health outcomes, to maximize animal well-being.

Keywords: hazelnut/linseed, lambs, gut-health, VFA, histomorphometry

References

- Andrés, S., Morán, L., Aldai, N., Tejido, M.L., Prieto, N., Bodas, R., Giráldez, F.J. 2014. Effects of linseed and quercetin added to the diet of fattening lambs on the fatty acid profile and lipid antioxidant status of meat samples. *Meat Sci.* 97, 156–163.
- Priolo, A., Valenti, B., Natalello, A., Bella, M., Luciano, G., & Pauselli, M. 2021. Fatty acid metabolism in lambs fed hazelnut skin as a partial replacer of maize. *Anim. Feed. Sci. and Tech.*, 272, 114794.
- Vastolo, A., Calabrò, S., & Cutrignelli, M. I. 2022. A review on the use of agro-industrial CO-products in animals' diets. *Ital. J. of Anim. Sci.* 21, 577–594.

Funding

LIVE HAZE - Hazelnut industrial by-product inclusion in livestock chains in Italy” funded by the MUR Progetti di Ricerca di Rilevante Interesse Nazionale (PRIN) Bando 2020 - grant 2020244EWW

* Presenting author. E-mail address: talal.hassan@unito.it (Talal Hassan)

Malassezia pachydermatis sensitivity to plant extracts from the Brazilian Amazon Rain Forest

Ivana Barbosa Suffredini^{a,b,*}, Jefferson de Souza Silva^c, Sergio Alexandre Frana^{a,b}, Karen Cristina Comin Maldonado^a, Mateus Luís Barradas Paciencia^a, Liv Nunes Almeida Gomes^{a,b}, Marcelo Henrique Soller Ramada^c, Antonio Drauzio Varella^{a,d}

^a Center for Research in Biodiversity, Paulista University, 900 Paulista Ave., 13100-100 São Paulo, Brazil

^b Graduate Program in Environmental and Experimental Pathology, Paulista University, 1212 Dr. Bacelar St., 04026-002 São Paulo, Brazil

^c Amazônia University, 1544, Ville Roy Ave., 69307-7025 Boa Vista, Roraima, Brazil

^d Drauzio Varella Site (www.drauziovarella.com.br), São Paulo, Brazil

The present work covers the search for new active plant extracts to be used in the treatment of infections caused by *Malassezia pachydermatis*, one of the most important pathogenic agents related to canine otitis and dermatitis. The microorganism can be cross-species transmitted from pets to tutors, and its control shows epidemiologic importance. Our group screened more than 2,300 Brazilian plant extracts against *M. pachydermatis*, a pathogenic yeast that has both human and veterinarian importance, using sensitivity tests (Suffredini et al., 2023) such as disk diffusion (DDA) and microdilution broth (MBA) assays and bioautography to identify antifungal extracts, compared to Amphotericin B (AmB) solution. The extract obtained from the leaves of *Guatteria* sp. (Annonaceae) showed antifungal activity in the DDA (12.31 ± 0.32 mm; 100 mg/mL), compared to AmB (18.76 ± 0.65 mm); in the MDB (minimal inhibitory concentration, MIC, of 5 mg/mL; AmB – MIC of 78 μ g/mL). Bioautography showed that fractions eluted to $R_f = 0.5$ showed antifungal activity. Analytical thin-layer chromatography (TLC) was performed and indicated the presence of alkaloids and phenolic compounds; preparative TLC was performed to assess gas-chromatography mass spectrometry analyses. Results indicated the presence of gamma-sitosterol **1**, 3',5'-dimethoxyacetophenone **2**, and palmitic acid **3** in the fractions (Fig 1).




Fig. 1. Structure of gamma-sitosterol **1**, 3',5'-dimethoxyacetophenone **2**, and palmitic acid **3** The extract obtained from the Annonaceae species showed antifungal activity, which may be related to the alkaloids that are present in the active fractions, combined or not with **1**, **2**, and **3**.

Keywords: Antarctic Continent, apoptosis, Brazilian Rain Forests, cytotoxicity, wound healing

References

Suffredini, I.B., et al., 2023. Screening Amazon rainforest plant extracts for antimicrobial activity: a 15-year commitment to the Brazilian biodiversity. *Front. Antib.* 2, 1122400. DOI:10.3389/frabi.2023.1122400.

Financial support acknowledgments: CNPq, CAPES, FAPESP

* Presenting author. E-mail address: ivana.suffredini@docente.unip.br (Ivana Barbosa Suffredini)

Orangutan herbal remedies: lessons for human health

Ivona Foitová^{a,*}, Radka Pecková^a, Karel Doležal^b, Wisnu Nurcahyo^c, Milan Olšanský^d

^a Masaryk University, Department Botany and Zoology, Faculty of Science, Kotlarska 2, Brno 61137 Brno, Czech Republic

^b Institute of Experimental Botany CAS, CZ-78371 Olomouc, Czech Republic

^c Gadjah Mada University, Department of Parasitology, Faculty of Veterinary Medicine, Jalan Fauna 2, Yogyakarta 55281, Indonesia

^d Foundation “UMI Saving of Pongidae”, Brno, Czech Republic

Both human and animal health are continually threatened by infectious diseases. The “One Health concept” has highlighted the strong interrelation between human and animal health in recent years. Understanding factors affecting individual health is crucial for determining the underlying health and resilience of the environment. The health of both individuals and ecosystems depends on homeostasis, the balancing of dynamic processes through continuous interaction and feedback within the integrated system.

Parasites play a significant role in individual, species, and ecosystem health. Given the phylogenetic similarities between humans and orangutans, studying orangutan feeding behavior is relevant, particularly regarding specific plant foods consumed to combat parasitic infections. Orangutans, the only ape species in Southeast Asia, face critical endangerment, listed as such by the IUCN.

Self-medication in primates sheds new light into the complex interactions of the animal, plant and parasite. While primates consume a variety of non-nutritional plant compounds and nutrient-poor bark, little is understood about the potential medicinal benefits of such ingestion. Our recent research confirms self-medication in orangutans (*Pongo pygmaeus*) for the first time, based on pharmacological analyses.

Several plants in the orangutan diet hold potential medicinal value, some used in traditional human medicine. This presentation will explore plants within the orangutan diet that demonstrate effectiveness against parasites through *in vitro* testing, such as *Piper betle*, *Archidendron fagifolium*, *Mimosa sp.*, *Knema laurina*. Our findings, along with descriptions of plant species exhibiting novel antiparasitic effects, may inform drug development efforts to combat infectious diseases in both human and animal populations.

Keywords: self-medication, anti-parasitic activity, orangutan, *in vitro* tests

References

Morrogh-Bernard, H.C., Foitová, I., Yeen, Z., Wilkin, P., de Martin, R., Rárová, L., Doležal, K., Nurcahyo, W., Olšanský, M. Scientific Reports, (2017); 7: 1-7.

Acknowledgements

Project is supported by Grant Agency of the Czech Republic (23-06571S) and by the “UMI – Saving of Pongidae” Foundation project “Parasites and Natural Antiparasitics in the Orang-utan”.

* Presenting author. E-mail address: foitova@sci.muni.cz (Ivona Foitová)

Willow (*Salix acmophylla*) leaf and twig extracts inhibit sporulation of coccidia (*Eimeria* spp.) in goats

Manal Haj-Zaroubi^{a,b}, Nariman Mattar^a, Sami Awabdeh^c, Alex Markovics^d, Joshua D. Klein^e, Hassan Azaizeh^{f,*}

^a Institute of Applied Research, The Galilee Society, P.O. Box 437, Shefa-Amr 20200, Israel

^b Department of Natural Resources and Environmental Management, Faculty of Management, University of Haifa, Haifa, Israel

^c National Agricultural Research Center (NARC), Amman, Jordan

^d Department of Parasitology, Kimron Veterinary Institute, P O. Box 12, Bet Dagan, 50250, Israel

^e Department of Natural Resources, Institute of Plant Sciences, Agricultural Research Organization, Volcani Center, Rishon Le Ziyon, 7505101, Israel

^f Tel Hai College, Department of Environmental Sciences & Water Science, Upper Galilee 12208, Israel

Willow (*Salix acmophylla*) trees, found worldwide, contain secondary metabolites that are valuable as dietary supplements for animal feed and as anti-parasites (Awwad et al., 2021). This study investigated the ethanol extracts from leaves and branches of three willow genotypes for their secondary metabolite composition and their potential to inhibit *Eimeria* sp. sporulation, a major concern in ruminant health. The extracts were characterized for phenolic compounds, flavonoids, and salicylic acid using HPLC. The total phenolic content of willow leaves and branches was similar in the three different genotypes. The total flavonoid content of the branches was significantly higher than that of leaves of the same genotype; however, the salicylic acid content was significantly higher in leaves than in branches. Importantly, all extracts exhibited significant inhibition of *Eimeria* sporulation, where over 70% inhibition was obtained at concentrations as low as 750 ppm. The sporulation inhibition of branch or leaf extracts exceeded 80% for leaves and 90% for branches at concentrations above 1250 ppm. HPLC analysis identified the flavonoids rutin and hyperoside in both leaves and branches, but their concentrations, along with those of salicin were higher in leaves. Conversely, p-coumaric acid was detected in branches but not in leaves. The study highlights the potential of using *Salix* extracts as bioactive compounds for biological control of coccidiosis in ruminants and emphasizes the importance of genotype-specific variations in secondary metabolite profiles. We conclude that all parts of *S. acmophylla* can provide secondary metabolites that act as a coccidiostat to treat *Eimeria* in goats.

Keywords: Ethanol extracts; Plant secondary metabolites; Coccids sporulation; *Eimeria* sp.; Willow genotypes

References

Awwad, S., Markovics, A., Halahlih, F., Yazbak, A., Haj-Zaroubi, M., Muklada, H., Klein, J.D., Azaizeh, H. 2021. Effect of irrigation water source on secondary metabolites in *Salix acmophylla* and their potential to impair exsheathment of gastro-intestinal nematodes. Ann. Agric. Crop. Sci. 6(3): 1079. <https://doi.org/10.26420/annagriccropsci.2021.1079>.

* Presenting author. E-mail address: hazaizeh@yahoo.com (Hassan Azaizeh)

Hot Pepper (*Capsicum annuum*) extract: not only a flavor enhancer, but also a useful fish feed additive

Sevdan Yılmaz, Sebahattin Ergün

Department of Aquaculture, Faculty of Marine Sciences and Technology, Çanakkale Onsekiz Mart University, Çanakkale, Türkiye

The present study investigated the effects of hot pepper (*Capsicum annuum*) extract (CAO) supplementation on growth performance, serum biochemical parameters, immune responses, histology, gene expression levels, gut microbiome, and sensory characteristics of rainbow trout. Five isonitrogenous and isoenergetic diets were formulated to contain hot pepper extract at levels of 0.7% (C7), 1.4% (C14), 2.1% (C21), and 2.8% (C28). Fish were fed experimental diets for 30 days (experiment one) and 45 days (experiment two). It was determined that significant improvements were achieved in growth performance, serum biochemical parameters and expression levels of genes responsible for immunity along with some humoral immune responses. Additionally, dietary CAO induced changes in the intestinal histological structure, as evidenced by an increased number of goblet cells in the CAO-supplied groups.

The sensory attributes of cooked trout fillets demonstrated that capsicum's pungent odor and taste were evident in the C14 group and subsequently increased with higher dietary CAO levels. Importantly, our results indicate, for the first time, that dietary CAO at levels of 7–14 g/kg can enhance the growth of rainbow trout without adversely affecting sensory characteristics or gut health.

Keywords: Rainbow trout, *Oncorhynchus mykiss*, immune responses, gene expression, biochemistry

* Presenting author. E-mail address: sergun@comu.edu.tr (Sebahattin ERGÜN).

Session IV: A.I.M.S. Cluster: Prospecting aquatic and terrestrial natural biological resources for biologically active compounds

Oral contributions

Enhancing bio-based compound diversity through advanced biotechnological approaches: The SECRETed project

Manuel Salvador^{a,*} on behalf of SECRETed consortium^b

^a Biotechnology division, IDENER.AI, Seville, Spain

^b <https://www.secreted.eu/partners/>

SECRETed, an EU-H2020 funded initiative, leads in aquatic biotechnology, targeting innovation in agrochemical, pharmaceutical, cosmetic, and chemical sectors. This multidisciplinary consortium utilizes the genetic diversity of biosurfactants and siderophores from marine and extremophilic microorganisms to create novel hybrid molecules with tailored properties. By leveraging existing microbial collections and advanced Machine Learning algorithms, SECRETed has developed a comprehensive microbial amphiphilic compound database, decoding genetic blueprints to enhance chemical diversity and sustainable industrial applications. Significant progress in data collection and analysis has been achieved through natural language processing and graph convolutional neural networks (Qin et al., 2021), enriching the database's depth. A top-down approach using novel QSAR/QSPR models that identified over 3700 molecules from natural product databases, based on their chelating or surfactant abilities. In addition, a bottom-up approach updated the MIBiG database (Terlouw et al., 2023) with retrosynthetic details on 473 compounds linked to 417 biosynthetic gene clusters (BGCs), significantly advancing our understanding of microbial biosynthesis. In parallel, screening of natural producers has been performed towards identification of appropriate microbial hosts, which were combined taking advantage of metabolic engineering methods to achieve growth coupled production of custom industry-driven biosurfactants and siderophores. Fermentation and downstream processing of selected microbial hosts has been scaled up, while their target metabolites (biosurfactants and siderophores) are currently being tested on end-user applications. The applications include encapsulation of enzyme Q10 (cosmetics and nutraceuticals) and essential oils (agrochemicals), siderophore-based antibiotics production, pharmaceuticals with cytotoxic effects and RNAi-based drug delivery formulations for eye diseases.

Keywords: biosurfactants, siderophores, green chemistry, systems and synthetic biology

Acknowledge: This work has been funded by H2020-FNR-11-2020: SECRETed (Grant number: 101000794).

References

- Qin, S., Jin, T., Van Lehn, R.C., Zavala, V.M., 2021. Predicting Critical Micelle Concentrations for Surfactants Using Graph Convolutional Neural Networks. *J. Phys. Chem. B.* 125(37), 10610-1062066.
- Terlouw, B.R., et al., 2023. MIBiG 3.0: a community-driven effort to annotate experimentally validated biosynthetic gene clusters. *Nucleic Acids Research.* 51(D1), D603-D610.

* Presenting author. E-mail address: manuel.salvador@idener.ai (Manuel Salvador de Lara).

Marine biodiversity as sustainable resource of disease-suppressive microbes and bioprotectants for aquaculture and crop diseases “MARBLES”

Fernando Reyes^{a,*} on behalf of the MARBLES consortium^b

^a Fundación MEDINA, Avda. del Conocimiento 34, 18016, Armilla, Granada, Spain

^b <https://marblesproject.eu/team/consortium/>

The marine environment remains a largely untapped and poorly understood source of natural products with potential for application in the pharmaceutical and food industries. Challenges include finding and identifying the bacteria and their molecules, extracting their active ingredients in a sustainable, effective way, and the commercial and regulatory difficulties of bringing a discovery from the lab to market. Currently there is an urgent need to sustainably harness new compounds with pharmaceutical and nutritional applications. MARBLES project addresses this need by exploring and exploiting the microbiomes of marine ecosystems including sponges, microalgae and fish, which rely on their microbiomes and microbial natural products for disease resistance. MARBLES is obtaining novel bioactive compounds and microbes that act as bioprotectants in aquaculture, agriculture and pharmaceuticals. MARBLES ecology-based bioprospecting strategy focuses on unique host-microbe interactions in marine environments. Partners' existing microbial collections and new ones generated during MARBLES are being harnessed for the discovery of novel natural products and microbial communities. For this, MARBLES is using a systems-wide genomics approach to uncover the bioactive agents in disease-suppressive microbiomes. Also, MARBLES is exploring host- and microbe-derived chemicals that elicit production of bioactive compounds to revitalise drug screening. The expected outcomes will be sustainable production of bioprotectants (microbes and consortia and bioactive natural products, their derivatives and elicitors) with potential to fight infectious diseases in the agrochemical, aquaculture and healthcare industries. MARBLES contributes to UN SDG 2, 3, 12, 13 and 14, as well as to current UN processes (BBNJ, DSI, SynBio).

Keywords: marine bioprospecting, microbiome, sustainable aquaculture and agriculture, bioactive natural products, healthy ecosystems

Acknowledge: This work has been funded by H2020-FNR-11-2020: MARBLES (Grant number: 101000392).

References

<https://marblesproject.eu/>

* Presenting author. E-mail address: fernando.reyes@medinaandalucia.es (Fernando Reyes).

Sustainable skincare ingredients from plant cells with scientifically proven efficacy

Amir Akhgari^{a,*}, Alain Goossens^{b,c,*}, Richard M. Twyman^d, Heiko Rischer^a, Kirsi-Marja Oksman-Caldentey^a

^a VTT Technical Research Centre of Finland Ltd., Tekniikantie 21, FIN-02150 Espoo, Finland

^b Department of Plant Biotechnology and Bioinformatics, Ghent University, Ghent, Belgium

^c Center for Plant Systems Biology, VIB, Ghent, Belgium

^d TRM Ltd, PO Box 493, Scarborough, YO11 9FJ, United Kingdom

The global cosmetics market was valued at \$375.3 billion in 2022 and is expected to reach \$560.5 billion by 2030. Manufacturers are focusing on natural ingredients to meet consumer demands for bioactive and sustainable cosmetic products. However, the sector is facing technical and economic challenges due to the lack of raw materials and insufficient scientific evidence of efficacy. *InnCoCells* is an EU-funded project aiming to address these issues by revolutionizing the way cosmetic ingredients are manufactured from underutilized plant resources, leading to sustainable products with scientifically proven effects. There are 17 partners in the *InnCoCells* consortium, bringing together academic and industry researchers from nine EU member states as well as the UK and Norway. The project is coordinated by VTT (Finland) and has a budget of €7.9 million. *InnCoCells* is developing pilot-scale production platforms based on plant cell suspension cultures, hairy roots, whole plants and agrifood side-streams, as well as innovative downstream processing methods to generate bioactive extracts. This includes a cascade biorefinery concept that uses by-products and biowaste to extract value-added ingredients. The safety and efficacy of the extracts is tested using a broad and diverse panel of enzymatic and cell-based assays, and the active ingredients are identified by detailed metabolic profiling. The most promising candidates will be tested in human volunteers and will ultimately be marketed by our industrial partners. As a case study, we discuss chili (*Capsicum* spp.) cell cultures for the production of bioactive metabolites with antimicrobial and antioxidant activities that are suitable as cosmetic ingredients.

Keywords: plants and plant cell cultures, bioassays, cosmetic ingredients, sustainability, underutilized plants resources

Acknowledge: This work has been funded by H2020-FNR-11-2020: InnCoCells (Grant number: 101000373).

* Presenting author. E-mail address: alain.goossens@psb.vib-ugent.be (Alain Goossens), amir.akhgari@vtt.fi (Amir Akhgari).

New trends in pharmacognosy and cosmeceuticals

Pharmacognosy in the digital era: exciting opportunities offered by metabolomics and accessible digitised data

Jean-Luc Wolfender^{a,b,*}, Arnaud Gaudry^{a,b}, Adriano Rutz^{a,b}, Luis Quiros-Guerrero^{a,b}, Olivier A. Kirchhoffer^{a,b}, Laurence Marcourt^{a,b}, Bruno David^d, Antonio Grondin^d, Emerson Ferreira Queiroz^{a,b}, Louis-Félix Nothias^{a,b}, Pierre-Marie Allard^{a,b,c}

^a Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, CMU, 1211 Geneva, Switzerland

^b School of Pharmaceutical Sciences, University of Geneva, CMU, 1211 Geneva, Switzerland

^c Department of Biology, University of Fribourg, 1700 Fribourg, Switzerland

^d Green Mission Pierre Fabre, Branche Phytochimie et Biodiversité, Institut de Recherche Pierre Fabre, 3 Avenue Hubert Curien, BP 13562, 31562 Toulouse, France

With the advent of omics, digital sciences, and artificial intelligence (AI), a real revolution has taken place in life sciences and is also affecting pharmacognosy. Today, detailed metabolome data of various organisms can be readily obtained, accelerating our understanding and offering a more comprehensive perspective on Earth's chemodiversity (Rutz et al. 2022). In metabolomics, the challenges remain significant, and several questions need to be addressed: How reliable are metabolite annotations based on LC-MS profiling of natural extracts? Massive qualitative information is obtained, how is it related to amounts of active ingredients? Is it possible to deduce 3D structure, key to bioactivity, from metabolite profiling data? As part of our research, we obtained metabolomic information on thousands of highly biodiverse extracts of both plant and fungal origin (Allard et al. 2022). We are trying to understand how the massive amount of information collected and the taxonomic links can be used to answer questions about confidence/redundance in MS annotation (Rutz et al. 2023). On the other hand, the resulting management of such massive data represents a further challenge. With the development of knowledge graphs (KG) based on semantic web tools (Gaudry et al. 2024) it is possible to organize this data and make it searchable by advanced queries (Gaudry et al. 2022). We will address the practical challenges associated with automating the assessment of extract chemical composition. We will discuss the potential of KG and AI in analyzing metabolomics data within the context of pharmacognosy findings, aiming to foster new discoveries.

Funding: The authors are thankful to the Swiss National Science Foundation (SNSF) for the financial support of this project (no. CRSII5_189921/1)

Keywords: metabolomics, LC-MS, knowledge graph, metabolite annotation, semantic web

References

- Allard, P.M., Gaudry, A., Quiros-Guerrero, L.M., Rutz, A., Dounoue-Kubo, M., Walker, T.W.N., Defosse, E., Long, C., Grondin, A., David, B., Wolfender, J.L., 2022. Open and reusable annotated mass spectrometry dataset of a chemodiverse collection of 1,600 plant extracts. *GigaScience* 12, 1-9.
- Gaudry, A., Huber, F., Nothias, L.F., Cretton, S., Kaiser, M., Wolfender, J.L., Allard, P.M., 2022. MEMO: mass spectrometry-based sample vectorization to explore chemodiverse datasets. *Front. Bioinform.* 2, 842964.
- Gaudry, A., Pagni, M., Mehl, F., Moretti, S., Quiros-Guerrero, L.M., Cappelletti, L., Rutz, A., Kaiser, M., Marcourt, L., Queiroz, E.F., Ioset, J.R., Grondin, A., David, B., Wolfender, J.L., Allard, P.M., 2024. A sample-centric and knowledge-driven computational framework for natural products drug discovery. *ACS Cent. Sci.* 10, 494-510.
- Rutz, A., Sorokina, M., Galgonek, J., Mietchen, D., Willighagen, E., Gaudry, A., Graham, J.G., Stephan, R., Page, R., Vondrasek, J., Steinbeck, C., Pauli, G.F., Wolfender, J.L., Bisson, J., Allard, P. M., 2022. The LOTUS initiative for open knowledge management in natural products research. *Elife* 11, e70780.
- Rutz, A., Wolfender, J.L., 2023. Automated composition assessment of natural extracts: untargeted mass spectrometry-based metabolite profiling integrating semiquantitative detection. *J. Agric. Food Chem.* 71, 18010-18023.

* Presenting author. E-mail address: jean-luc.wolfender@unige.ch (Jean-Luc Wolfender).

Nanophytocosmetics – a new trend in cosmeceuticals

Ilkay Erdogan Orhan^{a,b,*}

^a Department of Pharmacognosy, Faculty of Pharmacy, Gazi University, 06330 Ankara, Turkey

^b Turkish Academy of Sciences (TÜBA), Vedat Dalokay Street, No. 112, 066670, Ankara, Turkey

Cosmeceuticals are a subclass term of cosmetics with vast product miscellany and increased share in the world cosmetic market. Natural and/or organic cosmetics are receiving more attention from consumers all over the world concerning their safety and efficacy. Nanocosmetics with herbal substances refer to cosmetic products that incorporate nanoparticles along with natural herbal ingredients. These nanoparticles are typically on the nanometer scale, which allows for enhanced delivery of active herbal compounds into the skin or hair follicles, leading to improved efficacy and targeted delivery. The herbal substances used in these nanocosmetics can include extracts from plants, essential oils, and pure natural compounds as active ingredients. These products are often marketed as offering benefits such as increased hydration, antioxidant protection, anti-inflammatory properties, and overall skin health improvement. However, further research is needed to fully understand the safety and efficacy of nanocosmetics with herbal substances, as well as their potential long-term effects on human health and the environment. Consequently, the research and development of novel phyto-based cosmeceuticals *via* extensive screening studies on plant extracts and pure natural substances have been conducted by our group using *in vitro* (enzyme inhibition, *etc.*), *in silico* (molecular docking and toxicity screening), and cell-based assays. Some active ingredients identified and formulated for skin bleaching, anti-wrinkle, and anti-aging purposes by us up to date have been as the extracts from *Cotinus coggygria*, *Geranium glaberrimum*, *Garcinia mangostana*, *etc.*, in the forms of niosome, nanofiber or nanoemulsion formulations. In this connection, an anti-acne formulation based on a number of plant extracts tested against *Propionibacterium acnes* has also been developed by our group, which led to several patent applications. For wound healing, we have been studying nanofiber formulations loaded with some plant extracts. Our continuing analyses on discovering novel, effective, and safe natural ingredients with possible cosmeceutical use have so far yielded 3 patents and 5 patent applications. In this talk, our current upshots on phyto-based nanocosmeceuticals will be underlined.

Keywords: nanocosmetics, phytocosmetics, natural products, skin bleaching, anti-aging

* Presenting author. E-mail address: iorhan@gazi.edu.tr (Ilkay Erdogan Orhan).

Nitrogenous flavonoid derivatives production by reacting with amino acids

Haolin Zhang¹, Petra Högger², Randolph Arroo³, Mohamed A. Farag⁴, Jianbo Xiao^{5,*}

¹ Institute of Chinese Medical Sciences, University of Macau, Macao

² Institut für Pharmazie und Lebensmittelchemie, Universität Würzburg, 97074 Würzburg, Germany

³ De Montfort University – Leicester School of Pharmacy, The Gateway, Leicester LE1 9BH, UK

⁴ Mohamed A. Farag Pharmacognosy Department, Faculty of Pharmacy, Cairo University, Cairo 11562, Egypt

⁵ Universidade de Vigo, Nutrition and Bromatology Group, Department of Analytical and Food Chemistry, Faculty of Sciences, 32004, Ourense, Spain

A class of nitrogenous derivatives of flavonoids was detected using *in vitro* cell culture medium and *in vivo* mice produced without enzymatic catalysis have been consistently overlooked in literature. In this study, 39 flavonoids were incubated in Dulbecco's modified eagle's medium (DMEM) at 37° for 2 h to explore the reaction mechanism behind nitrogenous derivatives from flavonoids. Baicalein, scutellarein, DMY, GC, EGC, and EGCG were found to produce corresponding nitrogenous derivatives in both DMEM and mixed amino acid solution. The nitrogen source of these 6 flavonoid nitrogenous derivatives was revealed to be amino acid. The reaction site with amines in these flavonoid nitrogenous derivatives was identified as OH of the pyrogallol moiety in flavonoids *via* LC-MS/MS and NMR. This pyrogallol group was a key motif being first oxidized into quinone, further, to react with Strecker degradation of amino acids to yield N-flavonoids and corresponding aldehydes. Reaction optimization revealed that a slightly alkaline environment accelerates flavonoid nitrogenous derivatives formation by promoting the formation of flavonoid quinone. These results provide the first mechanistic evidence for the *in vitro* generation of flavonoid nitrogenous derivatives yet to be tested using *in vivo* assay.

Keyword: Nitrogenous flavonoids; baicalein; dihydromyricetin; catechin; pyrogallol; amino acids

Acknowledgement

This study was supported by Ramón y Cajal grant (RYC2020-030365-I) and Xunta de Galicia for supporting the program (Excelencia-ED431F2022/01).

SIF Italian Society of Phytochemistry and Sciences of Medicinal, Food and Perfume Plants

Investigating structural features and bioactivity of natural products by NMR and computational techniques

Maria Giovanna Chini*

Department of Biosciences and Territory, University of Molise, Contrada Fonte Lappone, Pesche 86090 (IS), Italy

Marine and terrestrial species provide an inexhaustible source of active secondary metabolites (Atanasov, Zotchev et al. 2021). These entities feature a vast array of 3D structures, particular biochemical properties, and other molecular characteristics, which make them a reliable starting point for the identification and development of potential lead compounds in the field of drug development. The correct elucidation of the structural features of already reported Natural Products (“old NPs”) and unprecedented (“new NPs”) ones represents the primary critical step for accelerating their complete biochemical and pharmacological characterization. Considering the complex structural architecture which often characterizes these NPs (especially if compared to synthetic molecules), the elucidation of their 2D structure and the precise 3D arrangements is required to provide essential insights into ligand-target interaction, which can drive a rational lead optimization process. Combining NMR spectroscopy with modern computational techniques (Chini, Urbani et al. 2020, De Vita, Terracciano et al. 2020) is one of the most effective approaches to achieve this task. In this context, the DFT/NMR integrated method is a valid support for clarifying the configuration and conformational profile of active secondary metabolites, shedding light on their molecular basis of mechanism of action. Also, the robust and efficient Inverse Virtual Screening (IVS) approach, developed and optimized by us (Chini, Lauro et al. 2021), can be helpful in achieving the target identification task (De Vita, Chini et al. 2020, De Vita, Chini et al. 2023), guiding subsequent biological investigations and pharmacological tests and for speeding-up, when possible, the drug repurposing. Moreover, the complete comprehension of NPs structural determinants responsible for their biological activity could be used for designing and synthesizing novel molecular platforms through computer-aided modifications of their original active skeletons (De Vita, Chini et al. 2020). This could lead to semi-synthetic derivatives with enhanced potency and selectivity towards specific targets involved in pathological events, such as inflammation and cancer.

Keywords: natural products, bioactive compounds, structural elucidation, NMR, computational tools

References

- Atanasov, A.G., Zotchev, S.B., Dirsch, V.M., Supuran, C.T., 2021. Natural products in drug discovery: advances and opportunities. *Nat. Rev. Drug Discov.* 20, 200-216.
- Chini, M.G., Lauro, G., Bifulco, G., 2021. Addressing the target identification and accelerating the repositioning of anti-inflammatory/anti-cancer organic compounds by computational approaches. *Eur. J. Org. Chem.* 2021, 2966-2981.
- Chini, M.G., Urbani, D., Dambruoso, P., Riccio, R. and Bifulco, G., 2020. Elucidating heteroatom influence on homonuclear $^4J_{(H,H)}$ coupling constants by DFT/NMR approach. *Magn. Reson. Chem.* 58, 566-575.
- De Vita, S., Chini, M.G., Lauro, G., Bifulco, G., 2020. Accelerating the repurposing of FDA-approved drugs against coronavirus disease-19 (COVID-19). *RSC Adv.* 10, 40867-40875.
- De Vita, S., Terracciano, S., Bruno, I., Chini, M.G., 2020. From natural compounds to bioactive molecules through NMR and *in silico* methodologies. *Eur. J. Org. Chem.* 2020, 6297-6317.
- De Vita, S., Chini, M.G., Bifulco, G., Lauro, G., 2023. Target identification by structure-based computational approaches: Recent advances and perspectives. *Bioorg. Med. Chem. Lett.* 83.

* Presenting author. E-mail address: mariagiovanna.chini@unimol.it (Maria Giovanna Chini).

The epidemic of antibiotic-resistant: a call to research and development of new natural antimicrobial molecules

Giuliana Donadio^{a,*}, Raffaella Nocera^a, Valentina Parisi^a, Emanuele Rosa^a, Fabrizio Dal Piaz^b, Nunziatina De Tommasi^a

^a Department of Pharmacy, University of Salerno, 84084 Fisciano, Italy

^b Department of Medicine and Surgery, University of Salerno, 84082 Baronissi, Italy

Presently, the world is in the midst of an emerging crisis of antibiotic resistance for microbial pathogens (Akram et al., 2023). The global spread of this phenomena is associated with misuse and overuse of existing antimicrobials in humans, animals and plants, which is accelerating the development and spread of antimicrobial resistance. Scientists and pharmaceutical companies must urgently focus on a common mission: the search for new antibiotics compounds possibly acting through unexploited mechanisms and molecular target(s). Plants represent a promising source of bioactive compounds (Rajput et al., 2020). Based on these considerations, a study aimed at shedding light on the effects of manool, a labdane diterpene from *Salvia officinalis* L. (Lamiaceae) with full blown antimicrobial activity, on the dental pathogen *Streptococcus mutans* (MIC 6 µg/ mL), which is considered a new Gram-positive model organism, was undertaken. Drug Affinity Responsive Target Stability (DARTS) (Lomenick et al., 2011) allowed the identification of transporters ATP-binding cassette (ABC), a large superfamily of integral membrane proteins involved in the absorption of nutrients, as manool target. To validate the inhibition of ABC proteins, AA and sugar uptake assays were performed by MS and NMR analyses. The obtained results offer two possible scenarios: the direct or indirect binding of manool to the ATP site, common to the majority of identified proteins. Furthermore, expression proteomics and metabolomic studies were also conducted in *S. mutans* treated or not with manool, allowing to define the pathways primary affected by diterpene treatment.

Keywords: bioactive compounds, antibiotic resistant, plants, drug affinity responsive target stability

References

- Akram, F., Imtiaz, M., ul Haq, I., 2023. Emergent crisis of antibiotic resistance: A silent pandemic threat to 21st century. *Microb. Pathog.* 174, 105923.
- Rajput, M., Kumar, N., 2020. Medicinal plants: A potential source of novel bioactive compounds showing antimicrobial efficacy against pathogens infecting hair and scalp. *Gene Rep.* 21, 100879.
- Lomenick, B., Jung, G., Wohlschlegel, J. A., Huang, J., 2011. Target identification using drug affinity responsive target stability (DARTS). *Curr. Protoc. Chem. Biol.* 3, 163-180.

* Presenting author. E-mail address: gdonadio@unisa.it (Giuliana Donadio).

Metabolomic patterns in response to abiotic stimuli of *Cladonia* lichens from dry grasslands

Manuela Mandrone^{a,*}, Augusto Hubaide Nozella^a, Gabriele Gheza^b, Juri Nascimbene^b, Ilaria Chiochio^a, Chiara Vallese^c, Ferruccio Poli^a

^a Department of Pharmacy and Biotechnology, University of Bologna, Via Irnerio 42, 40126 Bologna, Italy

^b BIOME Lab, Department of Biological, Geological and Environmental Sciences, University of Bologna, Via Irnerio 42, 40126 Bologna, Italy

^c Department of Earth, Environmental and Life Sciences, University of Genova, Corso Europa 26, 16132 Genova, Italy

Lichens are complex organisms, defined as a self-sustaining ecosystem generated from the symbiosis between photobionts and mycobionts (Hawksworth and Martin, 2020). This study focuses on 4 species of genus *Cladonia* (*C. foliacea*, *C. polycarpoides*, *C. rangiformis*, *C. rei*), endowed with numerous bioactivities (Adenubi et al, 2022). Each species was sampled monthly for one year, in 5 different sites alongside dry grasslands of the Ticino River valley (Italy). The objectives were: 1) to acquire and elucidate (for the first time) the ¹H NMR profile of these species, 2) to picture intraspecies metabolomic differences, and 3) interspecies metabolomic variation due to environmental factors. By chemometrics were obtained validated models, and the species were clustered not only based on their biomarkers (i.e. usnic, protocetatic, fumarprotocetatic, rangiformic, norrangiformic, sekikaic and homosekikaic acids), but also for the enrichment in primary metabolites. For instance, fumarate and sucrose were more concentrated in *C. polycarpoides*. *C. rangiformis* had the highest concentration of arabinitol, acetate and alanine, while glucose and trehalose were more abundant in *C. rei*. Several intraspecies differences, both related to time-point and harvesting site, were also detected. Moreover, the metabolome of all species resulted significantly affected by the temperature gradient. Arabinitol emerged as one of the most important metabolites produced in response to chilling. The overall results offer a metabolomics-based discrimination method for *Cladonia* species, bringing insights both on when to harvest these lichens in view of a specific medicinal use, and on how they might adapt to environmental changes.

Keywords: *Cladonia*, lichens, ¹H NMR-profiling, metabolomics, lichen-climate adaptation

References

Hawksworth, D.L., Martin G., 2020. Lichens redefined as complex ecosystems. *New Phytol.* 227, 1281.

Adenubi, O.T., Famuyide, I.M., McGaw, L.J., Eloff, J.N., 2022. Lichens: an update on their ethnopharmacological uses and potential as sources of drug leads. *J. Ethnopharmacol.* 115657.

* Presenting author. E-mail address: manuela.mandrone2@unibo.it (Manuela Mandrone).

Natural products as active substances and structure components of delivery systems: the case of nanovesicles

Giulia Vanti*, Lucia Grifoni, Maria Camilla Bergonzi, Anna Rita Bilia

Department of Chemistry, University of Florence, Via Ugo Schiff 6, Sesto Fiorentino, Florence, Italy

We report the development of nanovesicles made of natural active substances used as bilayer architecture components, replacing the conventional bilayer fluidity modulators such as cholesterol or acting as co-modulators. These nanovesicles can be easily loaded with other actives to improve their biopharmaceutical properties. Different cases will be presented. Escinosomes are deformable nanovesicles made of phosphatidylcholine and escin, a triterpene saponin performing both biologically active components and bilayer modulator of fluidity, successfully reported for the delivery of natural drugs, i.e., berberine chloride for cutaneous administration (Vanti et al., 2019) and andrographolide for subcutaneous route (Vanti et al., 2022). Ascosomes represent a further innovative nanocarrier made of phosphatidylcholine and derivatives of L-ascorbic acid, i.e., ascorbyl octanoate and ascorbyl decanoate, investigated as bioactive constituents of the vesicle bilayer (Risaliti et al., 2020). They were also studied as potential nanocarriers for the skin delivery of khellin, a natural furanochromone with various applications in skin pathologies, and selected as a lipophilic drug model. Moreover, escinosomes and ascosomes were gelled using cellulose derivatives. The obtained nanovesicles-loaded hydrogels combined the benefits of a controlled release and improved transdermal permeability of loaded active ingredients thanks to the nanovesicle components, with optimized viscosity properties thanks to the polysaccharide matrix. Both nanosystems demonstrated a good safety profile in rats. A further study developed nanovesicles based on resins of *Cistus creticus* L. and *Pistacia lentiscus* L. These nanovesicles were loaded with the essential oils of *Origanum dictamnus* L. and *Salvia fruticosa* Mill. All these species are from the Balkan Peninsula, selected because of their ethnomedical uses related to skin diseases. However, formulating essential oils and resins in stable dosage forms is mandatory to overcome their low chemical stability and improve their therapeutic use.

Keywords: nanovesicle, escin, ascorbyl derivatives, resin, *Pistacia lentiscus*, berberine, andrographolide, khellin, *Origanum dictamnus*, *Salvia fruticosa*, essential oil

Acknowledgments

This research was funded by the project EthnoHERBS (H2020-MSCA-RISE-2018, Grant Agreement No. 823973). The financial support provided by the MUR - Dipartimenti di Eccellenza 2023-2027 (DICUS 2.0) to the Department of Chemistry “Ugo Schiff” of the University of Florence is acknowledged.

References

- Risaliti, L., Ambrosi, M., Calamante, M., Bergonzi, M.C., Nostro, P. L., Bilia, A.R., 2020. Preparation and characterization of ascosome vesicles loaded with khellin. *J. Pharm. Sci.*, 109, 3114-3124.
- Vanti, G., Bani, D., Salvatici, M.C., Bilia, A.R., 2019. Development and percutaneous permeation study of escinosomes, escin-based nanovesicles loaded with berberine chloride. *Pharmaceutics* 11, 682.
- Vanti, G., Capizzi, M., Di Cesare Mannelli, L., Lucarini, E., Bergonzi, M.C., Ghelardini, C., Bilia, A.R., 2022. Escinosomes: safe and successful nanovesicles to deliver andrographolide by a subcutaneous route in a mice model of oxaliplatin-induced neuropathy. *Pharmaceutics* 14, 493.

* Presenting author. E-mail address: giulia.vanti@unifi.it (Giulia Vanti).

Session I: Cutting edge technologies in natural product drug discovery, formulation and development

Topical Lecture

Mass spectrometry in natural products from Brazil: from understanding chemical signaling to pre-clinical studies

Norberto Peporine Lopes

NPPNS, Departamento de Ciências Biomoleculares, Faculdade de Ciências Farmacêuticas de Ribeirão Preto-Universidade de São Paulo, Brasil

Plants, animals and microorganisms emit, detect, and respond to an amazing number of chemical signals. Advances in mass spectrometry have enabled natural products analyses at several levels, including complex metabolomic analysis. In this talk we will describes the most recent analytical developments in metabolomics applying examples of the Brazilian biodiversity. We also present studies with plants, frogs, and microorganism, which have allowed us to elucidate different biological activities and to advance in the understanding of the eco-physiological function of secondary metabolites. Finally, we describe the selected substances subjected to pre-clinical studies where MS imaging techniques allowed new alternatives for innovation processes.

Acknowledgment

FAPESP, CAPES e CNPq

* Presenting author. E-mail address: npelopes@gmail.com (Norberto Peporine Lopes)

Oral contributions

Ion channels: beyond native mass spectrometry

Ronald J. Quinn*, Yushu Gu, Miaomiao Liu, Linlin Ma

Griffith Institute for Drug Discovery, Griffith University, Brisbane, Queensland 4111, Australia

Ion channels represent a significant class of drug targets. The inwardly rectifying potassium Kir4.2 channel plays a crucial role in regulating membrane potentials and maintaining potassium homeostasis. Kir4.2 has been implicated in various physiological processes, including insulin secretion, gastric acid regulation, and the pathogenesis of central nervous system diseases. Transient receptor potential melastatin 2 (TRPM2) is a calcium-permeable, nonselective cation channel. It is involved in many pathological and physiological processes, making it a potential therapeutic target for various diseases, including Alzheimer's disease, Parkinson's disease, and cancers.

Native mass spectrometry has been established as a robust platform for investigating membrane protein-lipid interactions (Laganowsky et al., 2014). Native MS has been proven capable of revealing thermodynamics and allostery of individual lipid binding events (Cong et al., 2016). Nativeomics is an approach integrating native MS with small molecule fragmentation, to identify bound ligands released after native MS. The ligand is fragmented for identification through database searching (Gault et al., 2020).

We developed a mass spectrometry approach to directly detect the ligand after capture of the protein-ligand complex (Fig. 1). We firstly demonstrate the detection of the known ligands, polymyxin B and clotrimazole, respectively. Screening of a natural product library comprising 2000 compounds, with each pool containing 100 compounds, each compound at a concentration of 10 μ M, was screened against Kir4.2 and TRPM2 to identify ginsenoside Rg1 and ginsenoside F3, respectively.




Fig. 1. Collision-induced affinity selection mass spectrometry (CIAS-MS) can be utilized to directly identify ligands binding to ion channels.

CIAS-MS circumvents the need for high mass detection typically associated with mass spectrometry of large membrane proteins.

Keywords: CIAS-MS, TRPM2, Kir4.2, ion channels, collisional induced activation

References

- Laganowsky, A., Reading, E., Allison, T., Ulmschneider, M. B., Degiacomi, M. T., Baldwin A. J., Robinson C. V., 2014. Membrane proteins bind lipids selectively to modulate their structure and function. *Nature* 510, 172–175.
- Cong, X., Liu, Y., Liu, W., Liang, X., Russell, D. H., Laganowsky, A., 2016, Determining Membrane Protein–Lipid Binding Thermodynamics Using Native Mass Spectrometry. *J. Am. Chem. Soc.* 138, 4346–4349.
- Gault, J., Liko, I., Landreh, M. Shutin, D. Bolla, J. R. Jefferies, D. Agasid, M. Yen, H.-Y. Ladds, M. J. G. W. Lane, D. P. Khalid, S. Mullen, C. Remes, P. M. Huguet, R. McAlister, G. Goodwin, M. Viner, R. Syka, J. E. P.. Robinson, C. V Combining native and ‘omics’ mass spectrometry to identify endogenous ligands bound to membrane proteins. *Nat. Methods* 2020, 17, 505–508.

* Presenting author. E-mail address: r.quinn@griffith.edu.au (Ronald J Quinn)

High-throughput cell phenotype screening coupled with ion mobility spectrometry of American *Aconitum*

Yi Zhao^{a,b}, Dennis Y. Liu^c, Trevor N. Clark^c, Roger G. Linington^c, Edward J. Kennelly^{a,b,*}

^a Department of Biological Sciences, Lehman College, The City University of New York, Bronx, 10468 NY, United States

^b Biology Ph.D. Program, The Graduate Center, The City University of New York, New York, 10016 NY, United States

^c Department of Chemistry, Simon Fraser University, V5A 1S6 British Columbia, Canada

Aconitum plants have been widely used as traditional medicine in Asian countries for centuries. However, there are few records about how American *Aconitum* has been used medicinally, despite its phylogenetic closeness to medicinally important Asian species (Luo et al., 2005). This study aims to discover the bioactive compounds in American *Aconitum* based on a systematic molecular network strategy integrating both mass spectrometric data and a high-throughput phenotype screening assay, cell painting. The chemical profile in different parts of two American *Aconitum* species was obtained by ion mobility spectrometry technique and compared with non-American *Aconitum* species. Image analysis and feature extraction were performed on four different concentrations of *Aconitum* extracts assayed in cell painting, resulting in 2,090 unique morphological features per extract. In conjunction with the TargetMol library of 4,400 compounds with known mechanisms of action, 198 unique hierarchical clusters were established after a feature selection strategy known as Fast Correlation-Based Filtering, which reduced the number of features to 429. An overall activity heuristic called CP score was calculated for each sample. After integrating CP score and spectrometric data, a molecular network containing higher CP scores was constructed and the compounds with high activity were targeted and being identified. The molecular network showed that American *Aconitum* is more active than non-American species in the cell painting assay.

Keywords: *Aconitum*, cell painting, mass spectrometry, molecular network, drug discovery

References

Luo, Y., Zhang, F., Yang, Q.-E., 2015. Phylogeny of *Aconitum* subgenus *Aconitum* (Ranunculaceae) inferred from ITS sequences. *Plant Syst. Evol.* 252 (1–2), 11–25.

* Presenting author. E-mail address: edward.kennelly@lehman.cuny.edu (Edward J. Kennelly)

Antibiotic discovery from a bacterial colony: isolation of new peptidic antibiotics using robotics, 3D printed bioassays, and MS-based bioinformatics

Jung-Ho Lee^a, Jin Yi Tan^a, Rui Ma^b, Linh Nguyen^{a,c}, Shahebraj Kahn^b, Mallique Qader^b, Enock Mpofu^b, Gauri Shetye^b, Nyssa K. Krull^a, Mario Augustinović^a, Sesselja Omarsdottir^d, William Edmonson^e, Jonathon Rodriguez^e, Sanghyun Cho^{a,b}, Scott G. Franzblau^{a,b}, Brian T. Murphy^{a,b,*}

^a Dept. of Pharmaceutical Sciences, College of Pharmacy, University of Illinois at Chicago, USA

^b Institute for Tuberculosis Research, College of Pharmacy, University of Illinois at Chicago, USA

^c Inst of Marine Biochemistry, Vietnam Academy of Science and Technology, Hanoi, Vietnam

^d Faculty of Pharmaceutical Sciences, University of Iceland, Reykjavík, Iceland

^e Boys & Girls Clubs of Chicago, USA

Traditional front end antibiotic discovery methods such as microbial library generation, strain fermentation, and in vitro screening of natural product fractions have afforded most antibiotics currently used in the clinic (Newman et al, 2020). However, in the past few decades these methods have been far less likely to yield new therapeutic leads (Koehn et al, 2005). Further, as pharmaceutical companies continue to divest from antibiotic discovery it is becoming increasingly important for academic labs to fill this gap. In order to address this, our lab has developed several methods designed to increase the throughput and efficiency of academic antibiotic discovery programs. These methods focus on analysis of *single bacterial colonies* in the initial steps of a discovery program and include i) use of front-end, untargeted robotic colony picking; ii) development of novel, reusable, 3D printed high-throughput agar competition bioassays; and iii) integration of *i* and *ii* with our open-access MS-based bioinformatics platform as a means to prioritize hits and reduce sample redundancy. This pipeline led to the discovery of a rare class of tetronate polyketide antibiotics active against *Staphylococcus aureus* and *Borrelia burgdorferi* (Lee et al, 2023), in addition to novel peptides that exhibit activity against *S. aureus* (led by a middle school student in partnership with a community center in Chicago). The discovery of these new antibiotics from two strains will be presented in the context of introducing an innovative, efficient front end antibiotic discovery pipeline designed for academic labs.

Keywords: antibiotics, bacteria, MALDI-TOF, robotics, bioinformatics

References

- Koehn, F.E. and Carter, G.T. The evolving role of natural products in drug discovery. *Nat. Rev. Drug Disc.* 2005, 4, 206.
- Lee, J.-H. et al. Discovery of a new antibiotic demethoxytetrone using a Dual-sided Agar Plate Assay (DAPA). *ACS Inf. Dis.* 2023, 9, 1593.
- Newman, D.J. and G.M. Cragg, Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J Nat Prod*, 2020. 83, 770.

* Presenting author. E-mail address: btmurphy@uic.edu (Brian T. Murphy)

Bioactive triterpenic acid metabolism investigation through mass spectrometry approaches

Eleni V. Mikropoulou^{a,*}, Charalampia Amerikanou^b, Andriana C. Kaliora^b, George Dedoussis^b, Maria Halabalaki^a

^a Division of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771, Athens, Greece

^b Department of Nutrition and Dietetics, School of Health Science and Education, Harokopio University, 17671 Athens, Greece

Despite the well-established therapeutic value of natural products, their valorization by the pharmaceutical industry in recent years has fallen behind. One example of this phenomenon is the natural resin of the endemic Mediterranean shrub *Pistacia lentiscus* var. *Chia*, a unique PDO product produced exclusively in the southern part of the Greek island of Chios. Chios mastic gum's (CMG) ethnopharmacological background dates back to antiquity, while modern research and a monograph published by the EMA further validate its traditional use. CMG's bioactivity has mainly been attributed to its unique triterpenoids, and most notably to its characteristic compounds, namely masticadienonic and isomasticadienonic acids; nevertheless, CMG has been largely limited to the supplement stage, with its isolated constituents having been poorly investigated (Pachi et al., 2020).

Taking the above into consideration, the current research aims to shed light on this unique phytotherapeutic product, with data obtained from *in vitro*, *in vivo* and human trials, with special focus on bioavailability and metabolism aspects. Through the exploitation of recent technological advances in the fields of mass spectrometry and metabolomics, we attempt to divulge the triterpenoids' fate within the gastrointestinal tract, as well as to propose structure – activity relationships that could be further applied to the versatile triterpenic skeleton in general. Overall, our results suggest that the presence of circulating triterpenic acid metabolites in plasma might hint to the idea that the true active forms of CMG's constituents are none other than biotransformation products, created by Phase I and II metabolic reactions within the host's organism.

Keywords: mastic, mass spectrometry, metabolism, metabolomics, triterpenic acids

References

Pachi, V.K., Mikropoulou, E.V., Gkiouvetidis, P., Siafakas, K., Argyropoulou, A., Angelis, A., Mitakou, S., Halabalaki, M., 2020. Traditional uses, phytochemistry and pharmacology of Chios mastic gum (*Pistacia lentiscus* var. *Chia*, Anacardiaceae): A review. *J. Ethnopharmacol.* 254, 112485.

* Presenting author. E-mail address: e.mikropoulou@pharm.uoa.gr (Eleni V. Mikropoulou)

Applications of ChemGPS-NP in natural product research – exploring the Outer Rim

Anders Backlund*, Erik Jacobsson, Ulf Göransson

Department of Pharmaceutical Bioscience – Research group Pharmacognosy, Uppsala University Biomedical Centre, Husargatan 3, SE-751 24 Uppsala

The concept of chemography, *i.e.* navigating chemical space, has over the last decade been applied to a number of studies of natural products.

It was early demonstrated that proximity in the ChemGPS-NP eight-dimensional physico-chemical property space can be interpreted as a measure of molecular similarity (Rosén et al., 2009). In subsequent studies it was further concluded that this measure provided additional information to that of other commonly used methods *e.g.* structure fragment based molecular fingerprints (Buonfiglio et al., 2015).

One of the most appealing aspects of chemography is that patterns can be readily visualized. This can also be combined with methods to define volumes from asymmetric ‘clouds’ of compound representations. Many studies up to date have been focused on small, low molecular weight, druglike molecules. However, we are now exploring the outer rim of this chemical property space including sets of polypeptides, polyketides and other larger and highly aberrant natural products.

In this study some implications of such exploration are demonstrated and discussed.




Fig. 1. ChemGPS-NP-based analysis for a series of larger natural product molecules, including representatives of groups of polypeptides, combined with low molecular weight compounds from the Maybridge screening set (gray). In the score plot of the three dimensions PC1 (red box) describes size and shape; PC2 (yellow box) the aromatic- and conjugation-related properties; and PC3 (green box) lipophilicity, polarity, and H-bond capacity.

Keywords: Chemography, ChemGPS-NP, polypeptides, polyketides, hyper-dimensionality

References

- Buonfiglio, R.; Engkvist, O.; Várkonyi, P.; Henz, A.; Vikeved, E.; Backlund, A.; Kogej, T. J., 2015. Investigating Pharmacological Similarity by Charting Chemical Space. *J. Chem. Info. Mod.* 55, 2375-2390.
- Rosén, J.; Gottfries, J.; Muresan, S.; Backlund, A.; Oprea T., 2009. Novel Chemical Space Exploration via Natural Products. *J. Med. Chem.* 52, 1953-1962.

* Presenting author. E-mail address: anders.backlund@uu.se (Anders Backlund)

Addressing the challenge of protein target identification of bioactive phytochemicals with a yeast-3-hybrid screening

Simone Moser*

Institute of Pharmacy, Department of Pharmacognosy, University of Innsbruck, Center for Chemistry and Biomedicine (CCB), Innrain 80-82, 6020 Innsbruck Austria

Although phytochemicals are nature's treasure box and are intensely studied for their biological effects in human cells, knowledge on direct protein interactions is scarce. A small molecule may have strong biological effects, but its protein target and molecular mechanism may remain unknown. These are, however, essential for developing the compound as a possible drug candidate (Bunnage *et al.*, 2015). Indeed, affected networks and downstream or upstream signaling pathways are often the missing link to completely understand the mechanisms leading to the activity of a small molecule and its cellular effects.

Un-biased target identification, or 'target fishing', is a challenging and often laborious task (Tabana *et al.*, 2023). Not only does the interaction partner need to be identified and selected out of a vast pool of potential hits obtained *via* diverse screening methods, the confirmation and validation of a putative target requires expertise in a variety of fields, depending on the target.

A potent yet complementary method to classical chemical proteomic approaches consists in the yeast three-hybrid (Y3H) system. The Y3H system was first described by Licitra and Liu in 1996 (Licitra *et al.*, 1996), and offers several benefits over affinity based methods for finding protein targets, one of the most important one is the fact that it is an *in vivo* method. In our lab, we have recently set up an improved Y3H system, and have demonstrated its power with the identification of a previously un-known target of the widely used contraceptive ethinylestradiol (Wang *et al.*, 2022).

We are now ready to apply the Y3H system as target identification platform for natural products, and are further planning to improve the versatility and usefulness of the system towards natural products.

Keywords: target identification, yeast-3-hybrid, phytochemicals

References

- Tabana, Y., Babu, D., Fahlman, R. *et al.* Target identification of small molecules: an overview of the current applications in drug discovery. *BMC Biotechnol* 23, 44 (2023).
- Bunnage, M., Gilbert, A., Jones, L. *et al.* Know your target, know your molecule. *Nat Chem Biol* 11, 368–372 (2015). <https://doi.org/10.1038/nchembio.1813>
- Licitra, E.J., Liu, J.O. A three-hybrid system for detecting small ligand–protein receptor interactions. *PNAS* 93 (23) 12817–12821 (1996).
- Wang P, Klassmüller T, Karg CA, Kretschmer M, Zahler S, Braig S, Bracher F, Vollmar AM, Moser S. Using the yeast three-hybrid system for the identification of small molecule-protein interactions with the example of ethinylestradiol. *Biol Chem.* 2022 Feb 28;403(4):421–431.

* Presenting author. E-mail address: simone.moser@uibk.ac.at (Simone Moser)

Remediation of plant extracts using liquid-liquid chromatography: removal of pesticides and tetrahydrocannabinol from hemp extracts as a case study

Simon Vlad Luca*, Mirjana Minceva

Biothermodynamics, TUM School of Life Sciences, Technical University of Munich, 85354 Freising, Germany

Liquid-liquid chromatography (LLC) is a preparative separation technique where the upper and lower phases of a biphasic solvent system are used as the mobile and stationary phases. Typically, LLC is employed for the purification of target compounds, such as bioactive natural products (Luca et al., 2020, 2021). In this work, we propose a model-based approach for the remediation of plant extracts with LLC. As a case study, we have selected the removal of intrinsic (i.e., tetrahydrocannabinol/THC) and extrinsic (i.e., pesticides) toxicants from hemp extracts. By taking advantage of a fully predictive thermodynamic model (COSMO-RS), a considerable number of biphasic solvent systems were initially screened. After validation in shake-flask experiments, three solvent systems were chosen: heptane/methanol/water (I), heptane/acetone/water (II), and heptane/acetonitrile/water (III). For pesticide remediation, a classification system ranking 59 pesticides in terms of their difficulty to be removed from hemp extracts was proposed. It was shown that the pesticide classification system can be used to select the appropriate solvent system for removing the majority of the contaminating pesticides. For THC remediation, various LLC operating modes were evaluated. Batch elution mode (conventional pulse injection) provided a THC-free hemp extract with a good productivity, whereas trapping multiple dual mode showed a lower productivity than batch. The sequential (continuous) mode afforded a THC-free extract with a similar composition to the batch mode and offered the advantage of continuous operation and superior productivity. The proposed approach is general and can be explored for the remediation of other contaminants from natural products.

Keywords: *Cannabis sativa* L., cannabidiol, tetrahydrocannabinol, pesticides,

References

- Luca, S.V., Roehrer, S., Kleigrew, K., Minceva, M., 2020. Approach for simultaneous cannabidiol isolation and pesticide removal from hemp extracts with liquid-liquid chromatography. *Ind. Crops Prod.* 155, 112726.
- Luca, S.V., Bruesewitz, M., Minceva, M., 2023. Evaluation of advanced liquid-liquid chromatography operating modes for the tetrahydrocannabinol-remediation of hemp extracts. *Chem. Eng. Sci.* 282, 119279.

Acknowledgments

This work was financed by the Federal Ministry of Education and Research (BMBF, Germany, no. 01DS19029B) and the Federal Ministry for Economic Affairs and Energy (BMW, Germany, no. 16KN092721).

* Presenting author. E-mail address: vlad.luca@tum.de (Simon Vlad Luca)

nmrXiv enhances the utility of NMR spectroscopic data of natural products

Venkata Chandrasekhar Nainala^{a,*}, Nisha Sharma^a, Noura Rayya^a, Stefan Kuhn^c, Hamed Musallam^a, Guido F. Pauli^b, Christoph Steinbeck^a

^a *Cheminformatics and Computational Metabolomics, Friedrich Schiller University, Lessingstr. 8, 07743 Jena, Germany*

^b *Pharmacognosy Institute and Dept. of Pharmaceutical Sciences, University of Illinois Chicago College of Pharmacy, Chicago, IL 60612, U.S.A*

^c *Institute of Computer Science, University of Tartu, Ülikooli 18, 50090 Tartu, Estonia*

Nuclear magnetic resonance (NMR) spectroscopy is a key tool in natural product (NP) analysis for the identification of unknowns, dereplication of knowns, metabolomic analysis, determination of relative and absolute configuration, and relative and absolute quantitation even of unweighable samples. Following the development of various first-generation databases for NMR data, recent efforts aim at broader scientific communities, seeking to enable direct data depositions connected to publications and projects. Our nmrXiv project, hosted at nmrxiv.org, addresses the particular needs of the NP and metabolomics communities by fostering FAIR principles of original data sharing, open licences, as well as structure and feature search capabilities.

A host of remarkable features makes nmrXiv stand out. At its core, nmrXiv serves as a versatile and long-term data archive, fostering collaboration and transparency in the scientific community. The advanced search and retrieval functionalities of nmrXiv empower researchers to access relevant NMR data and expedite their work. nmrXiv strongly emphasises open access and advocates for open science approaches that enhance rigor and reproducibility and drive innovation. Utilising state-of-the-art web tools, the nmrXiv platform fosters collaboration and knowledge sharing among researchers by creating an interconnected community rather than being repository-/storage-only. Stringent measures for data standardisation and quality control ensure the reliability and reproducibility of deposited data. Integration with popular NMR data formats and Electronic Lab Notebooks streamlines data processing and entry and promotes consistency. An interactive NMR spectra visualisation tool, NMRium [1], provides instrument-independent data processing capabilities and facilitates NMR data interpretation. Compliance with metadata standards (DataCite and BioSchemas) enhances the utility of NMR datasets.

nmrXiv is fully accessible to the global NP research community. It offers authors a streamlined submission tool for raw NMR data associated with publications, helping them to comply with contemporary FAIR requirements set forth by major publishers and funding agencies. Our contribution will explain the key tools available in nmrXiv and show how specifically NP researchers can benefit from engaging in both sharing of their own data as well as analyzing and mining data shared by fellow scientists.

Keywords: raw NMR data initiative, FAIR principles, data sharing

References

- [1] Patiny L, Musallam H, Bolaños A, Zasso M, Wist J, Karayilan M, Ziegler E, Liermann JC, Schlörer NE. NMRium: Teaching nuclear magnetic resonance spectra interpretation in an online platform. *Beilstein J Org Chem.* 2024 Jan 5;20:25-31. doi: 10.3762/bjoc.20.4. PMID: 38213841; PMCID: PMC10777324.

* Presenting author. E-mail address: chandu.nainala@uni-jena.de (Venkata Chandrasekhar Nainala)

Natural product-inspired chemical biology: development of highly sensitive labeling reagents for amino acids and scarce natural products

Hideaki Kakeya*

Graduate School of Pharmaceutical Sciences, Kyoto University, 606-8501 Kyoto, Japan

Scarce natural products that possess unique biological activities have been ideal drug leads for decades. Among them medium-sized peptide molecules have received much attention as good motifs in drug design as they have intermediate properties between those of small molecule pharmaceuticals and antibodies. However, their identification and structural determinations are problematic owing to sample amount limitation.

Inspired by an extremely rare marine natural products yaku'amides, our highly sensitive labeling reagents L-FDVDA (1-fluoro-2,4-dinitrophenyl-5-L-valine-*N,N*-dimethylethylene-diamineamide) and L-FDLDA (1-fluoro-2,4-dinitrophenyl-5-L-leucine-*N,N*-dimethylethylene-diamineamide) that are powerful tools for detection of chiral amino acids were recently developed. By fusion of the Marfey's reagents with the key C-terminus structural motif of yaku'amides, the detection sensitivities of the reagents L-FDVDA and L-FDLDA for proteinogenic amino acids and non-proteinogenic amino acids such as all structural isomers and enantiomers of aminobutyric acid were drastically enhanced in LC-MS analysis, whose method named "the highly sensitive-advanced Marfey's method" (Kuranaga et al., 2020). Additionally, our method enabled to separate and identify the D/L-forms of short-chain peptides as well as the amyloid b fragment peptides with racemised and isomerised aspartic acid residues (Ozaki et al., 2023).

Our sensitivity-enhancement design concept was also applicable to the development of reagents for labeling saccharides and reactivity-guided isolation of electrophilic scarce natural products. we succeeded in detection of presaccharothriolide Z, the 10-membered microbial macrolide anticancer produced by a rare actinomycete *Saccharothrix* sp. A1506, using our original molecular probe A, and accomplished the total synthesis of presaccharothriolide Z (Kuranaga et al., 2021).




Fig. 1. Structures of L-FDVDA, L-FDLDA, probe A, and presaccharothriolide Z

Keywords: labeling reagents, LC-MS, amino acids, scarce natural products, chemical biology

References

- Kuranaga, T. et al. 2020. Highly sensitive labeling reagents for scarce natural products. *ACS Chem. Biol.* 15, 2499-2506.
- Kuranaga, T. et al. 2021. Identification and total synthesis of an unstable anticancer macrolide pre-saccharothriolide Z produced by *Saccharothrix* sp. A1506. *Org. Lett.* 23, 7106-7111.
- Ozaki, M. et al. 2023. Separation of amyloid b fragment peptides with racemized and isomerized aspartic acid residues using an original chiral resolution labeling reagent. *Analyst*, 148, 1209-1213.

* Presenting author. E-mail address: scseigyo-hisyo@pharm.kyoto-u.ac.jp (Hideaki Kakeya)

Role of Metabolomics in natural product discovery and chemical ecology

Neha Garg*

School of Chemistry and Biochemistry, Georgia Institute of Technology, 950 Atlantic Drive, Atlanta, Georgia 30332-2000, United States

Microbes whether commensal or pathogenic respond to the surroundings via small molecule natural products. Many of these small molecules are inspired by the biochemical modification of simple molecules that are an integral part of our cell biology, yet their discovery is challenging. This presentation will cover the development and application of mass spectrometry-based untargeted metabolomics methods for the discovery and dereplication of small molecule natural products mediating microbe-drug, microbe-microbe, and microbe-host interactions. By coupling untargeted metabolomics with specific microbial phenotypes such as response to antimicrobial compounds, chemoinformatic methods such as molecular networking, sub-structure and chemical class identification, isotope labeling in cell culture, we will describe discovery of a strain specific pathway involved in inactivation of the antibiotic trimethoprim and the production of homogentisic acid derived natural products from a human mucus-dwelling microbial pathogen. Second application of these methods will describe the discovery of a siderophore degrading pathway mediating an interaction between coral mucus-dwelling commensal and pathogenic microbes. Finally, challenges and opportunities of mass spectrometry-based metabolomics will be discussed in discovery of natural products and their role in chemical ecology across biological systems.

* Presenting author. E-mail address: neha.garg@chemistry.gatech.edu (Neha Garg)

Session IIA: Total synthesis and biomimetic synthesis of natural products

Topical Lecture

***Olea europaea*: a compelling source of health promoting biomolecules and an inspiration for the semi-synthetic design of bioactive analogues**

Leandros A. Skaltsounis

Division of Pharmacognosy and Natural Products Chemistry, National & Kapodistrian University of Athens, Athens 15771, Greece

Olea europaea and its main products olive oil and olive drupes, have attracted scientific attention in recent years because of its biological activities and its attribution in many aspects of human health. Although olive oil primarily consists of oleic acid (up to 80 %) and other fatty acids, some minor phenolic compounds, comprising the 1-2 % of the total content, are generally considered to be responsible for the various health benefits of olive oil. The most characteristic compounds in this group are the simple phenylethyl alcohols such tyrosol, hydroxytyrosol, the seco-iridoids oleacein and oleocanthal, the decarboxylated aglycons of oleuropein and ligstroside respectively. These two compounds, and especially oleocanthal, have been identified as the agents responsible for the pungency of extra virgin oil. Oleocanthal's recent discovery as COX inhibitor, with similar effect to that of ibuprofen, has dramatically increased its interest both for the study of biological properties but also for the development of new non-steroidal anti-inflammatory drugs based on its structure. Additionally, according to many data, both compounds demonstrated promising anticancer and neuroprotective activities with no toxic effects. Prompted by the outstanding interest of these high-value natural compounds we have developed of a concise and scalable procedure for the synthesis of various analogues. The synthesis is performed by a convenient biomimetic and stereo-controlled approach, starting from oleuropein, isolated from olive leaves, an abundant biorenewable raw material and hydroxytyrosol the major compound of olive mill waste water.

References

- Skaltsounis, A.L., Argyropoulou, A., Aligiannis, N., Xynos, N., 2015. Recovery of high added value compounds from olive tree products and olive processing, 334-357, in olive and olive oil bioactive constituents. Ed. D. Boskou, AOCS PRESS.
- Beauchamp, G.K.; Keast, R.S.; Morel, D.; Lin, J.; Pika, J.; Han, Q.; Lee, C.H.; Smith, A.B.; Breslin, P.A., 2005. Ibuprofen-like activity in extra-virgin olive oil. *Nature* 437, 45046.
- Vougogiannopoulou, K., Lemus, C., Halabalaki, M., Pergola, C., Werz, O., Smith, A.B., Michel, S., Skaltsounis, A.L., Deguin, B., 2014. One-step semisynthesis of oleacein and the determination as a 5-lipoxygenase inhibitor. *J. Nat. Prod.* 77, 441-5.
- Skaltsounis, A.L., Kostakis, I., Gaboriaud-Kolar, N., Christoforidou, N., Sarikaki, G., 2021. Process for the production of oleocanthal, oleacein and their analogues. EP3838885A1.
- Sarikaki, G., Christoforidou, N., Gaboriaud-Kolar, N., Smith, A.B., Kostakis, I., Skaltsounis, A.L., 2020. Biomimetic synthesis of oleocanthal, oleacein, and their analogues starting from oleuropein, a major compound of olive leaves. *J. Nat. Prod.* 83, 1735-39.

* Presenting author. E-mail address: skaltsounis@pharm.uoa.gr (Leandros A. Skaltsounis).

Synthetic homoisoflavonoids to treat wet-stage macular degeneration

Elisha Griffin^a, Dulcie Mulholland^a, Sianne Schwikkard^b, Timothy Corson^c

^a University of Surrey, Guildford, Surrey, GU2 7XH

^b Kingston University, Kingston-upon-Thames, KT1 2EE, UK

^c University Of Toronto, 144 College Street, Toronto, ON, M5S 3M2, Canada

The occurrence of ocular angiogenesis resulting in choroidal neovascularisation (CNV) is the key process that leads to the wet-stage form of macular degeneration (wet AMD). This debilitating disease is the main cause of blindness in the aging population, but many patients are resistant to current therapies (Schwikkard et al., 2019). Existing treatments for wet AMD target the pro-angiogenic vascular endothelial growth factor (VEGF) pathway and consist of monoclonal antibodies and decoy receptors to VEGF. Due to the high molecular weights of these biologics, treatments are administered via intravitreal injection, a painful and costly process associated with undesirable side effects such as ocular haemorrhage (Tolentino, 2011). Thus, the discovery of a small molecule biologic to treat wet AMD could provide a less invasive route of administration, and potentially tackle the problems associated with current therapies. Homoisoflavonoids are a class of naturally occurring compounds often isolated from plants in the Hyacinthaceae family, whose *Eucomis* genus initially gained attention from researchers due to its use by African traditional healers (Mkhumbeni et al., 2022). Both naturally derived and fully synthetic homoisoflavonoids have previously been shown to exhibit anti-angiogenic effects against human retinal endothelial cells (HRECs) *in vitro* (Schwikkard et al., 2019). This provides an exciting opportunity for the potential to prevent CNV. Here, we describe the full syntheses of non-naturally derivatised homoisoflavonoids, many of which exhibited anti-angiogenic effects, with growth inhibitory concentrations (GI₅₀'s) as low as 1.48 µM against the proliferation of HRECs *in vitro*.




Fig 1. Synthetic homoisoflavonoids and their GI₅₀'s against the proliferation of HRECs.

Keywords: full chemical synthesis, homoisoflavonoid, macular degeneration, anti-angiogenic activity

References

- Mkhumbeni N., Pillay M., Mtunzi F, Motaung K.S.C., 2022. Effect of *Eucomis autumnalis* on the osteogenic differentiation of adipose-derived stem cells. *Tissue Eng. Part A*. 28, 136-149.
- Schwikkard, S., Whitmore, H., Sishtla, K et al., 2019. The antiangiogenic activity of naturally occurring and synthetic homoisoflavonoids from the Hyacinthaceae (sensu APGII). *J. Nat. Prod.* 82, 1227-1239.
- Tolentino, M., 2011. Systemic and ocular safety of intravitreal anti-VEGF therapies for ocular neovascular disease, *Surv. Ophthalmol.* 56, 95-113.

*Presenting author. E-mail address: eg00494@surrey.ac.uk (Elisha Griffin).

Minor Amaryllidaceae alkaloids as inspiration for development of highly selective butyrylcholinesterase inhibitors: design, synthesis, and biological evaluation

Lucie Cahlíková^{a,*}, Filip Pidany^a, Jan Korábečný^b, Daniela Suchánková^a, Jana Křoustková^a

^a Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmacy, Charles University, Czech Republic

^b Biomedical Research Center, University Hospital Hradec Kralove, Czech Republic

Plants of the family Amaryllidaceae have a long history of usage as herbal remedies all over the world to cure different ailments and diseases. Since the isolation of the first Amaryllidaceae alkaloid (AA), lycorine, more than 600 AAs have been isolated and studied in terms of their biological activities. Within the last 10 years, more than 100 AAs of different structural types have been isolated in our research group from the genera *Zephyranthes*, *Narcissus*, *Nerine*, and others. Isolated compounds have been tested for biological activity connected with the potential treatment of neurodegenerative, oncological, and infectious diseases. Newly isolated dimer alkaloids from *Narcissus pseudonarcissus* cv. Carlton, named carltonine A and B, showed strong selective inhibition activity against butyrylcholinesterase (BuChE) in nanomolar range ($IC_{50} = 31\text{--}910\text{ nM}$) (Mamun et al., 2020). Unfortunately, these alkaloids are present in plant material only in trace amounts, and they cannot be isolated for either more detailed biological investigation or commercial use. Thus, we decided to preserve some of the crucial structural requirements from carltonine A/B that are plausibly responsible for high BuChE inhibition activity, i.e. the 4-[2-(benzylamino)ethyl]phenol moiety, and modify other molecular regions to elucidate SAR. Up to date, more than 120 compounds, structurally inspired from minor AAs carltonine A and B, have been designed, synthesized, and evaluated for their acetylcholinesterase (AChE) and BuChE inhibition properties. Compounds that revealed BChE inhibition below 100 nM were selected for detailed biological investigation. The CNS-targeted profile of the most active compounds was confirmed theoretically by calculating the BBB score algorithm, these data were corroborated by determining the permeability *in vitro* using PAMPA-assay for the most active derivatives. Moreover, a crystallographic study was performed to inspect the binding mode of compound **87**, revealing essential interactions between **87** and hBChE active site (Pidany et al., 2023).

Keywords: Amaryllidaceae, butyrylcholinesterase, PAMPA-assay, crystallographic study

References

- Mamun, A.A., Marikova, J., Hulcova, D., Janousek, J., Safratova, M., Novakova, L., Kucera, T., Hrabanova, M., Kunes, J., Korabecny, J., Cahlikova, L., 2020. Amaryllidaceae alkaloids of belladine-type from *Narcissus pseudonarcissus* cv. Carlton as new selective inhibitors of butyrylcholinesterase. *Biomolecules* 10, 800.
- Pidany, F., Kroustkova, J., Mamun A.A., Suchankova, D., Brazzolotto, X., Nachon, F., Chantegreil, F., Dolezal, R., Pulkrabkova, L., Muckova, L., Hrabanova, M., Finger, V., Kufa, M., Soukup, O., Jun, D., Jenco, J., Kuns, J., Novakova, L., Korabecny, J., Cahlikova, L., 2023. Highly selective butyrylcholinesterase inhibitors related to Amaryllidaceae alkaloids- Design, synthesis, and biological evaluation. *Eur. J. Med. Chem.* 252, 115301.

* Presenting author. E-mail address: cahlikova@faf.cuni.cz (Lucie Cahlíková).

Naturally based prototypes for vector-control: synthesis and modification of carlina oxide scaffold


Eleonora Spinozzi^{a,*}, Marta Ferrati^a, Alessia Piergentili^a, Cecilia Baldassarri^a, Paolo Rossi^b, Loredana Cappellacci^a, Filippo Maggi^a, Riccardo Petrelli^a

^a Chemistry Interdisciplinary Project (ChIP), School of Pharmacy, University of Camerino, Via Madonna Delle Carceri, 62032 Camerino, Italy

^b School of Bioscience and Veterinary Medicine, University of Camerino, Via Gentile III Da Varano, 62032 Camerino, Italy

Vector-borne diseases cause public health concern worldwide and vector-control represents an everlasting challenge (Chala and Hamde, 2021). Thus, the research for novel effective tools is encouraged and compounds from botanical sources are gaining more interest as alternatives to conventional insecticides (Isman, 2020). Carlina oxide (**1**), isolated from the roots of *Carlina acaulis*

L. (Asteraceae), resulted effective against insect vectors, agricultural and stored-product pests (Spinozzi et al., 2023). Nevertheless, the plant species protection in several countries and the lack of a cultivation system are the main limiting factors for the exploitation of this molecule. This presentation will cover the development of a synthetic protocol to produce **1** (Fig. 1) through a one- step Sonogashira coupling and the synthesis of analogues featuring modifications to the chemical scaffold of the parental compound. These modifications were applied on the furan moiety and on the propynyl linker.



1

Fig. 1. Carlina oxide (**1**) scaffold.

The compounds were tested for acute toxicity on larvae of *Aedes albopictus* (Skuse) and *Anopheles stephensi* Liston and their safety profile was assessed on human keratinocytes (HaCaT) through MTT assay. Among the compounds tested, the analogue bearing the butynyl linker resulted more effective on the two insect species (LC₅₀ of 5.8 and 3.1 µg/mL on *Ae. albopictus* and *An. stephensi*, respectively) than its natural counterpart also displaying lower cytotoxicity (IC₅₀ > 100 µg/mL). The results demonstrated the possibility of synthesizing **1** and producing analogues with improved bioactivity and reduced cytotoxicity on keratinocytes.

Keywords: Vector-control, *Carlina acaulis* L., carlina oxide, synthesis, larvicidal

References

- Chala, B., Hamde, F., 2021. Emerging and re-emerging vector-borne infectious diseases and the challenges for control: A review. *Front. Public Health* 9, 1-10.
- Isman, M.B., 2020. Botanical insecticides in the twenty-first century—fulfilling their promise?. *Annu. Rev. Entomol.* 65, 233-249.
- Spinozzi, E., Ferrati, M., Cappellacci, L., Caselli, A., Perinelli, D. R., Bonacucina, G., ...& Benelli, G., 2023. *Carlina acaulis* L. (Asteraceae): Biology, phytochemistry, and application as a promising source of effective green insecticides and acaricides. *Ind. Crops Prod.* 192, 116076.

* Presenting author. E-mail address: eleonora.spinozzi@unicam.it (Eleonora Spinozzi).

Design and anti-inflammatory evaluation of ellagitannin-derived postbiotic metabolite conjugated with serotonin and dopamine

Maciej Korczak^{a,*}, Sheyda Bahiraii^b, Martina Redl^b, Piotr Roszkowski^c, Sebastian Granica^d, Judith M. Rollinger^b, Elke Heiss^b, Jakub P. Piwowski^a

^a Microbiota Lab, Medical University of Warsaw, Warsaw, Poland

^b Department of Pharmaceutical Sciences, University of Vienna, Vienna, Austria

^c Faculty of Chemistry, University of Warsaw, Warsaw, Poland

^d Department of Pharmaceutical Biology, Medical University of Warsaw, Warsaw, Poland

Urolithin A (UA) is a postbiotic metabolite produced by gut microbiota following oral administration of ellagitannin-rich plants, and it is known for its significant anti-inflammatory, anti-cancer, and anti-aging biological activities (García-Villalba et al., 2022). However, UA conjugation with glucuronic acid *in vivo* leads to the production of metabolites with reduced biological activity (Bobowska et al., 2021). In response to UA's limitations, a series of UA derivatives (UADs) have been synthesized through conjugation with nonsteroidal anti-inflammatory drugs, with their identity, stability, anti-inflammatory efficacy, and intestinal bioavailability being thoroughly assessed (Korczak et al., 2022, 2023). Recently, the synthesis of novel UADs linked to neurotransmitters, specifically serotonin (UASer) and dopamine (UADop), was proposed. The purity and identity of obtained conjugates were verified through HPLC-MS/MS and NMR. The cytotoxicity and anti-inflammatory effects of these novel UADs were examined using immortalized bone marrow-derived macrophages. qPCR experiments showed that the mRNA levels of pro-inflammatory markers such as IL-1 β , IL-6, iNOS, and TNF- α were significantly reduced by both UASer and UADop. Their influence on NF- κ B signaling was also investigated using a NF- κ B LUC – HEK reporter cell line. Additionally, preliminary *in vivo* studies on *Caenorhabditis elegans* were conducted to assess, among others, toxicity and potential life span extension by these newly synthesized compounds. In conclusion, the structural modification of natural products-derived postbiotic metabolites is presented as a promising approach, serving as a foundation for further investigations into their application in neuroinflammation.

Project financially supported by Polish National Science Centre grant Preludium Bis No. UMO- 2019/35/O/NZ7/00619

Keywords: ellagitannins, urolithin A, gut microbiota, neurotransmitters, inflammation

References

- Bobowska, A., Granica, S., Filipek, A., Melzig, M.F., Moeslinger, T., Zentek, J., Kruk, A., Piwowski, J.P., 2021. Comparative studies of urolithins and their phase II metabolites on macrophage and neutrophil functions. *Eur. J. Nutr.* 60, 1957–1972.
- García-Villalba, R., Giménez-Bastida, J.A., Cortés-Martín, A., Ávila-Gálvez, M.Á., Tomás-Barberán, F.A., Selma, M.V., Espín, J.C., González-Sarriás, A., 2022. Urolithins: a comprehensive update on their metabolism, bioactivity, and associated gut microbiota. *Mol. Nutr. Food Res.* 2101019.
- Korczak, M., Roszkowski, P., Granica, S., Piwowski, J.P., 2022. Conjugates of urolithin A with NSAIDs, their stability, cytotoxicity, and anti-inflammatory potential. *Sci. Rep.* 1–15.
- Korczak, M., Roszkowski, P., Skowrońska, W., Żołdak, K.M., Popowski, D., Granica, S., Piwowski, J.P., 2023. Urolithin A conjugation with NSAIDs inhibits its glucuronidation and maintains improvement of Caco-2 monolayers' barrier function. *Biomedicine and Pharmacotherapy* 169.

* Presenting author. E-mail address: maciej.korczak@wum.edu.pl (Maciej Korczak).

Session IIB: Genetic engineering of food crops and medicinal plants

Topical Lecture

Unravelling the fitness programs that steer plant metabolism allows creating novel tools for the engineering of plant-based production of bioactive metabolites

Alain Goossens^{a,b,*}

^a *Department of Plant Biotechnology and Bioinformatics, Ghent University, 9052 Ghent, Belgium*

^b *Center for Plant Systems Biology, 9052 Ghent, Belgium*

Nature has invented strict, yet not fully understood, regulatory networks that control plant metabolism. These networks safeguard plant fitness in a continuously changing environment. By investigating the reprogramming of plant metabolism by developmental and environmental cues, we aim to advance our fundamental understanding of the mechanisms that steer plant metabolism. We specifically focus on jasmonate (JA), the phytohormone that steers the delicate balance between growth and defense programs across the plant kingdom, including the production of bioactive specialized metabolites. In parallel, this enables us to unlock plant specialized metabolism for numerous human applications given that our findings serve simultaneously as a novel resource for engineering tools that facilitates the creation of (plant-based) synthetic biology platforms for the sustainable production of high-value plant metabolites. I will exemplify this by reporting on our recent work on the biosynthesis of bioactive metabolites in model, crop and medicinal plants.

Keywords: plant metabolism, biosynthesis of bioactive metabolites

* Presenting author. E-mail address: alain.goossens@psb.vib-ugent.be (Alain Goossens)..

Oral contributions

Towards the engineering of the plant endoplasmic reticulum for sustainable production of specialized metabolites

Elke Clicque^{a,b,*}, Alain Goossens^{a,b}

^a Department of Plant Biotechnology and Bioinformatics, Ghent University, Technologiepark 71, B-9052 Ghent, Belgium

^b VIB Center for Plant Systems Biology, Technologiepark71, B-9052 Ghent, Belgium

Plants are among the most creative survivors on the planet. They have optimized their own defence mechanisms against attackers, typically by producing a species-specific arsenal of specialized metabolites. These metabolites not only play vital roles within the plants themselves but also offer a multitude of applications as pharmaceuticals, fragrances etc. However, industrial production of these valuable compounds faces many bottlenecks and often requires intensive adaptations. To facilitate sustainable large-scale production of plant metabolites, we aim to develop a universal tool to boost plant-based production of specialized metabolites. Our strategy is based on recent findings in yeast, in which engineering of the endoplasmic reticulum (ER), which harbours a main part of the enzymatic machinery for many metabolite biosynthesis pathways, results in an enlarged ER size and an increased metabolite accumulation (Arendt et al., 2017). By interfering with the phospholipid biosynthesis pathway, a higher production of building blocks for the ER-membrane and ultimately an enlarged ER were achieved. Our project aims to translate the findings from yeast to plants. Thus far, we have obtained an increased accumulation of phospholipid building blocks in tomato (*Solanum lycopersicum*) roots, by knocking out the *PHOSPHATIDIC ACID PHOSPHOHYDROLASE* (PAH) genes. Currently, we are investigating whether this correlates with an increased production of bioactive specialized metabolites. Additionally, we are also exploring the potential of ER engineering by overexpressing other key phospholipid biosynthetic enzymes. With our obtained *in planta* results, we believe that ER engineering holds great promise for a more efficient and sustainable production of plant specialized metabolites.

Keywords: *Solanum lycopersicum*, endoplasmic reticulum, phospholipids, engineering, roots

References

Arendt, P., Miettinen, K., Pollier, J., De Rycke, R., Callewaert, N., Goossens, A., 2017. An endoplasmic reticulum-engineered yeast platform for overproduction of triterpenoids. *Metab. Eng.* 40, 165-175.

* Presenting author. E-mail address: elcli@psb.vib-ugent.be (Elke Clicque).

Session IIC: Medicinal natural products: from bench to bedside

A new cytotoxic diphenazine from a deep-sea fungus *Cystobasidium laryngis*

Hee Jae Shin^{a,b,*}, Hwa-Sun Lee^{a,c}, Jong Soon Kang^d, Min Ah Lee^{a,c}, Chang-Su Heo^{a,b}

^a Marine Natural Products Laboratory, Korea Institute of Ocean Science and Technology, 385 Haeyang-ro, Yeongdo-gu, Busan 49111, Republic of Korea

^b Department of Marine Biotechnology, University of Science and Technology, 217 Gajung-ro, Yuseong-gu, Daejeon 34113, Republic of Korea

^c Department of Chemistry, Pukyong National University, 45 Yongso-ro, Nam-gu, Busan 48513, Republic of Korea

^d Laboratory Animal Resource Center, Korea Research Institute of Bioscience and Biotechnology, 30 Yeongudanji-ro, Ochang-eup, Cheongwon-gu, Cheongju 28116, Republic of Korea

We report here the isolation, structure elucidation, and anticancer and anti-neuroinflammatory activities of phenazostatin J (**1**) from *Cystobasidium laryngis* isolated from a deep-sea sediment sample taken at the Indian Ocean Ridge along with semi-synthesis of **1** and *in vivo* test result against the human stomach cancer line (NUGC-3). The *Cystobasidium* genus has been reported as one of the uncommon fungi in deep-sea environments (Wei et al., 2018). In addition, no studies were conducted on natural products from *Cystobasidium* sp. until we first reported new metabolites from the strain in 2019 (Lee et al., 2019). As part of our ongoing search for bioactive compounds from an extract of culture broth of *C. laryngis*, we additionally isolated six new diphenazine derivatives, phenazostatins E–J, and evaluated their anti-neuroinflammatory and cytotoxic activities (Lee et al., 2022). Among the isolated new diphenazines, only phenazostatin J (**1**) exhibited strong anti-neuroinflammatory ($IC_{50} = 0.30 \mu M$) and cytotoxic ($IC_{50} = 7.7 nM$) activities against the NUGC-3 cell line.




Fig. 1. Structure of phenazostatin J.

To conduct an *in vivo* test against the NUGC-3 (stomach) cell line, we successfully semi-synthesized **1** using saphenic acid which is a monomer of **1**. Phenazostatin J (**1**) was administered intraperitoneally at different doses (1, 2, and 4 mg/kg) using a nude mouse subcutaneous model. An *in vivo* study demonstrated that **1** suppressed tumor growth in nude mice carrying NUGC-3 cells without significant secondary adverse effects. Tumor sizes and weights were reduced when treated with 4 mg/kg of **1**, with similar results to doxorubicin (2 mg/kg), a positive control. These results suggested that **1** has therapeutic potential as an anticancer candidate that deserves further investigation.

Keywords: *Cystobasidium laryngis*, phenazostatin J, semi-synthesis, cytotoxicity, *in vivo* test

References

- Lee, H.-S., Kang, J. S., Choi, B.-K., Lee, Y.-J., Lee, J., Shin, H. J., 2019. Phenazine derivatives with anti-inflammatory activity from the deep-sea sediment-derived yeast-like fungus *Cystobasidium laryngis* IV17-028. *Mar. Drugs* 17, 482.
- Lee, H.-S., Kang, J.S., Cho, D.Y., Choi, D. K., Shin, H.J., 2022. Isolation, structure determination, and semisynthesis of diphenazine compounds from a deep-sea-derived strain of the fungus *Cystobasidium laryngis* and their biological activities. *J. Nat. Prod.* 85, 857–865.
- Wei, X., Guo, S.S., Gong, L.F., He, G.Y., Pang, K.L., Luo, Z.H., 2019. Cultivable fungal diversity in deep-sea sediment of the East Pacific Ocean. *Geomicrobiol. J.* 35, 790–797.

* Presenting author. E-mail address: shinhj@kiost.ac.kr (Hee Jae Shin).

Safety and efficacy of topical Arnica tincture for the treatment of uncomplicated human cutaneous leishmaniasis

Sara M. Robledo^{a,*}, Any C. Garcés^b, Franziska M. Jürgens^c, Thomas J. Schmidt^c

^a PECET-Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia

^b Grupo de Investigación Clínica-PECET, Sede de Investigación Universitaria, Universidad de Antioquia, Medellín, Colombia

^c University of Münster, Institute of Pharmaceutical Biology and Phytochemistry (IPBP), PharmaCampus - Corrensstrasse 48, D-48149 Münster, Germany

Cutaneous leishmaniasis (CL) is classified as a neglected tropical disease for which new effective and safe therapeutic options are required. Arnica tincture (AT) is a herbal medicinal preparation with anti-inflammatory activity which is used traditionally for the topical treatment of blunt injuries among other uses. Its main bioactive constituents are sesquiterpene lactones (STLs) of the helenalin and 11 α ,13-dihydrohelenalin types. In our previous work, AT showed promising antileishmanial effects *in vitro* and *in vivo* (Robledo et al., 2022) and Glutathione conjugation plays a major role in the metabolism of these STLs (Jürgens et al., 2022a). Furthermore, *in vivo* dermal absorption, pharmacokinetic, and toxicity studies with STLs from AT were also reported (Jürgens et al., 2022b). As part of our continuing search for new alternatives to treat CL, an open-label, randomized, non-comparative phase Ib/II clinical trial was performed (NCT05094908). Adult volunteers with a parasitologically confirmed diagnosis of CL were randomly allocated to receive AT 3 times per day either for 4 or 6 weeks. Up to now (Feb. 2024), 7 out of 9 subjects screened have been included in the study. Cure has already been observed in all subjects in the 4- and 6-weeks group respectively (Fig. 1). Four patients so far have finished the 6-month follow-up period without showing relapses. The patients still under observation also demonstrate promising results. Furthermore, AT proved to be safe, and the adverse events reported are local, around the area of application of the tincture, and of mild intensity.



Fig. 1. Appearance of CL before and after treatment with AT

In conclusion, AT appears to be an efficacious, safe and well-tolerated intervention to treat CL.

Funding: This research was funded by the Wilhelm Doerenkamp-Foundation, grant specification: NATVANTAGE RESEARCH GRANT 2018.

Keywords: *Leishmania braziliensis*, clinical trial, sesquiterpene lactones, therapeutical response, adverse effects

References

- Robledo, S.M., Murillo, J., Arbeláez, N., Montoya, A., Ospina, V., Jürgens, F.M., Vélez, I.D., Schmidt, T.J., 2022. Therapeutic efficacy of *Arnica* in hamsters with cutaneous leishmaniasis caused by *Leishmania braziliensis* and *L. tropica*. *Pharmaceutics* (Basel). 15, 776.
- Jürgens, F.M., Behrens, M., Humpf, H.U., Robledo, S.M., Schmidt, T.J., 2022a. *In vitro* metabolism of helenalin acetate and 11 α ,13-dihydrohelenalin acetate: natural sesquiterpene lactones from *Arnica*. *Metabolites*. 12, 88.
- Jürgens, F.M., Robledo, S.M., Schmidt, T.J. 2022b. Evaluation of pharmacokinetic and toxicological parameters of arnica tincture after dermal application *in vivo*. *Pharmaceutics*. 14, 2379.

* Presenting author. E-mail address: sara.robledo@udea.edu.co (Sara M. Robledo).

Mesembryanthemum tortuosum improves Parkinson's Disease symptoms in a zebrafish larvae model

Keagile Lepule^{a,*}, Maxleene Sandasi^{a,b}, Alvaro Viljoen^{a,b}

^a Department of Pharmaceutical Sciences, Tshwane University of Technology, Pretoria, South Africa

^b SAMRC Herbal Drugs Research Unit, Tshwane University of Technology, Pretoria, South Africa

Parkinson's disease (PD) is a progressive neurodegenerative disease characterised by the loss of dopaminergic neurons in the brain. The exact cause of PD is still not fully understood however, oxidative stress, inflammation, and genetic and environmental factors have been implicated (Olanow and Tatton, 1999). The pathological repertoire of PD can be induced in zebrafish larvae using reserpine (Puttonen et al., 2017). Currently, PD has no cure, however, medicinal plants such as *Mesembryanthemum tortuosum* with documented central nervous system properties may provide some benefit. The aim of the study was to evaluate the neuroprotective and neurorestorative effects of *M. tortuosum* extracts on reserpine-induced deficits, using a zebrafish larvae model. To achieve this, methanol and alkaloid-rich extracts were prepared from *M. tortuosum* raw material, while Zembrin[®] (commercial extract of *M. tortuosum*), the two alkaloids (mesembranol and Δ^7 mesembrenone) and L-dopa (positive control) were commercially sourced. Zebrafish larvae were treated with the extracts and the effect on locomotion, reactive oxygen species (ROS) and total glutathione (tGSH) was monitored (Fig. 1).




Fig. 1. The model for neuroprotective and neurorestorative treatment regimen

The results demonstrate that Zembrin[®] exhibited superior anti-PD activity by improving locomotion, reducing ROS and subsequently increasing tGSH in both the neurorestorative and neuroprotective assays. The alkaloid-rich extract did not improve locomotion but increased tGSH. The individual alkaloids enhanced movement, although not to the same extent as L-dopa and Zembrin[®]. *Mesembryanthemum. tortuosum* extracts and alkaloids improve reserpine-induced PD like effects in a zebrafish larvae model.

Keywords: *Mesembryanthemum tortuosum*, parkinson's disease, reserpine, zebrafish larvae, locomotion

References

- Olanow, C., Tatton, W., 1999. Etiology and pathogenesis of Parkinson's disease. Annu. Rev. Neurosci. 22, 123-144.
- Puttonen, H.A., Semenova, S., Sundvik, M., Panula, P., 2017. Storage of neural histamine and histaminergic neurotransmission is VMAT2 dependent in the zebrafish. Sci. Rep. 7, 3060-3012.

*Presenting author. E-mail address: LepuleKH@tut.ac.za (Keagile Lepule).

Tailoring plants as photosynthetic production chassis for small molecules

Connor Tansley^{a,b,*}, Mahima^b, Melissa Salmon^b, Daria Golubova^b, Honghao Su^b, Nicola Patron^{a,b}

^a Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA

^b Engineering Biology, Earlham Institute, Norwich Research Park, UK NR4 7UZ

Plant natural products are utilised in many industries including as dyes, scents, pharmaceuticals and agrochemicals. Chemical synthesis has provided easy and cheap access to some molecules but remains challenging or uneconomical for many molecules. In the past decade, aided by advances in genomics and synthetic biology, biosynthetic pathways have been elucidated and rebuilt in so-called ‘chassis organisms’. To date, most work has been done in microbes, in which it has been shown that yields can be maximised by engineering precursor availability, balanced expression of pathway genes, and removing enzymes that compete for substrates or derivatise intermediates (Cravens et al., 2019). The model plant *Nicotiana benthamiana* is an increasingly attractive organism for the heterologous biosynthesis of high-value, biologically active molecules (Stephenson et al., 2019), however little has been done to optimise production. Previously, our lab observed that the relative positions of genes within synthetic constructs affects both transient and transgenic expression as well as product yields (Kallam et al., 2023). Further, genome editing techniques can be used to remove molecules and enzymes that interfere with production and purification techniques (Dudley et al., 2022; Vollheyde et al., 2023). In this study, we investigate how the expression of synthetic genes are affected by their neighbours (Fig.1A), and how knowledge of enzyme kinetics and rate limiting steps can be used to improve construct design. We also describe the production of genome-edited plants with multiplexed mutations (Fig.1B), tailored for triterpene production.




Fig. 1. Optimising natural product biosynthesis in plant chassis (A) The expression of synthetic genes is affected by transcription of upstream genes and can inform optimal construct designs (B) Genome editing of competing metabolic pathways.

Keywords: metabolic engineering, *Nicotiana benthamiana*, gene expression, synthetic biology, triterpenes

References

- Cravens, A., Payne, J., Smolke, C.D., 2019. Synthetic biology strategies for microbial biosynthesis of plant natural products. *Nat. Commun.* 10(1), 2142.
- Dudley, Q.M., Jo S, Guerrero, D.A.S., Chhetry M., et al., 2022. Reconstitution of monoterpene indole alkaloid biosynthesis in genome engineered *Nicotiana benthamiana*. *Commun Biol.* 5(1), 949.
- Vollheyde, K., Dudley, Q.M., Yang, T., Oz, M.T., Mancinotti, D., Fedi, M.O., Heavens, D., Linsmith, G., Chhetry, M., Smedley, M.A., Harwood, W.A., Swarbreck, D., Geu-Flores, F. Patron, N.J., 2023. An improved *Nicotiana benthamiana* bioproduction chassis provides novel insights into nicotine biosynthesis. *New Phytol.* 240, 302-317.
- Kallam, K., Moreno-Giménez, E., Mateos-Fernández, R., Tansley, C., Gianoglio, S., Orzaez, D. Patron, N.J., 2023. Tunable control of insect pheromone biosynthesis in *Nicotiana benthamiana*. *Plant Biotechnol. J.* 21, 1440-1453.
- Stephenson MJ, Reed J, Patron NJ, Lomonossoff G., Osbourn A., 2019. Engineering tobacco for plant natural product production. *Comprehensive Natural Products III (Third Edition) Chemistry and Biology* 6, 244-262.

*Presenting author. E-mail address: connor.tansley@earlham.ac.uk (Connor Tansley).

Session III: Natural cosmetics, nutraceuticals, and food supplements

Topical Lecture

How to harness the potential of eutectic solvents for the formulation and enhancement of natural ingredients

Leslie Boudesocque-Delaye^{a,*}, Alexis Verger^b, Laura Divoux^{a,b}, Solène Odou^a, Roxane Grard^b, Iron Mike Ardeza^{a,b}, Xavier Perse^b, Salima Bouderbala^b, Charlotte Pradel^b, Alexandra Despres^c, Emilie Munnier^b

^a SIMBA EA 7502 Université de Tours, Faculté de Pharmacie, 31 avenue Monge 37200 Tours, France

^b UPR CNRS CBM, Département NMNS, Faculté de Pharmacie, 31 avenue Monge 37200 Tours, France

^c RCP Design Global, Avenue Marcel Dassault, 37000 Tours, France

Natural deep eutectic solvents (NaDES) are being studied as an alternative to conventional solvents for preparing cosmetic ingredients from biomasses. These solvents are prepared by mixing plant cellular constituents, such as sugars, or amino acids. Their unique hydrogen bond network provides them high extraction power and the ability to stabilize sensitive active ingredients. To provide innovative natural cosmetic active ingredients, NaDES used must comply with international cosmetic regulations (Europe, China), which excludes NaDES based on choline chloride. Judiciously chosen NaDES-based extracts could be included directly in cosmetic products, making the solvent removal step in the extraction process unnecessary. Therefore, it is necessary to investigate the impact of various types of NaDES on the stability of active ingredients and the properties of cosmetic products. We will present the extraction and formulation of two examples of cosmetic active ingredients in a eutectic environment. First, we will examine an example of phycobiliproteins enriched extract, derived from microalgal biomasses. These extracts were prepared using an innovative solid/liquid/liquid extraction/purification approach (Hilali et al. 2024). The purified fractions were then incorporated into a gel formulation, demonstrating the successful stabilization of fragile phycobiliproteins within the gel. Additionally, sensory properties were enhanced. The following section presents an active ingredient from marigold (*Calendula officinalis*) that contains flavonoids and carotenoids. This ingredient is obtained through an innovative mechanical process that does not involve heating and is then incorporated into a cream formulation. (Boudesocque et al., 2024). These studies showcase the potential of eutectic solvents for extracting and formulating cosmetic ingredients.

Keywords: Eutectic solvent, natural cosmetic, natural active ingredient, sensory, stability

References

Boudesocque-Delaye L, Ardeza IM, Verger A, Grard R, Théry-Koné I, Perse X, Munnier E. Natural Deep Eutectic Solvents as a Novel Bio-Based Matrix for Ready-to-Use Natural Antioxidants-Enriched Ingredients: Extraction and Formulation Optimization. *Cosmetics*. 2024; 11(1):17.

Soukaina Hilali, Louise Van Gheluwe, Mervé Yagmur, Laura Wils, Myriam Phelippe, Barbara Clément-Larosièrre, Bénédicte Montigny, Johan Jacquemin, Emilie Thiery, Leslie Boudesocque-Delaye, NaDES-based biorefinery of *Spirulina* (*Arthrospira platensis*): A new path for sustainable high value-added metabolites, *Separation and Purification Technology*, 2024, 329, 125123.

* Presenting author. E-mail address: leslie.boudesocque@univ-tours.fr (Leslie Boudesocque-Delayea)

Oral contributions

A multi-target approach to evaluate the efficacy of a capsiate rich fraction of *Capsicum annuum* fruit against typeII diabetes and obesity

Shabana I. Khana^{a,b,*}, Hee-Sung Chae^a, Charles L. Cantrell^c, Robert L. Jarret^d, Ikhlas A. Khan^{a,b}

^a National Center for Natural Products Research, School of Pharmacy, University of Mississippi, MS 38677, USA

^b Department of Biomolecular Sciences, School of Pharmacy, University of Mississippi, MS 38677, USA

^c Natural Products Utilization Research Unit, USDA-ARS, University, MS 38677, USA

^d Plant Genetic Resources Unit, USDA-ARS, Griffin, GA 30223, USA

The fruits of a nonpungent cultivar of *Capsicum annuum* are rich in capsiate which is a nonpungent analog of capsaicin. This study was carried out to investigate the effects of a capsiate-rich fraction of *C. annuum* (germplasm 509-45-1) fruits (CR) on various nuclear receptors involved in multiple signaling pathways related to carbohydrate and lipid metabolism. The effects of CR on the muscular glucose uptake and adipogenesis were also determined to evaluate its anti-diabetic and anti-obesity properties.

At a concentration of 100 µg/mL, CR caused the activation of PPARα, PPARγ, LXR and NRF2 in hepatic cells (2 to 4-fold increase). CR was effective in decreasing adipogenesis and caused a 23% decrease in lipid accumulation in adipocytes. Moreover, the adipogenic effect induced by rosiglitazone was also antagonized by CR. Further studies in differentiated muscle cells indicated an increase of glucose uptake by 30%. Gene profiling studies showed a decrease in C/EBPα and an increase of KLF4 expression.

In conclusion, the study revealed a strong agonistic action of CR on multiple nuclear receptors which was associated with enhanced glucose uptake and decreased fat accumulation indicating a potential utility of the capsiate-rich fraction of *C. annuum* as a natural product that could ameliorate the symptoms of diabetes and obesity without the adverse effect of pungency. As a multiple agonist of PPARα, PPARγ and LXR, CR may also prevent the undesired adipogenic effects of antidiabetic drugs that are full PPARγ agonists. Further studies are warranted in animal models of TypeII diabetes and obesity.

Keywords: *Capsicum annuum*, capsiate, nuclear receptors, adipogenesis, glucose uptake

* Presenting author. E-mail address: skhan@olemiss.edu (Shabana I. Khana)

***Cimicifuga racemosa* extract Ze 450 mediates resilience against oxidative stress, inflammatory responses and UV-induced aging through metabolic rewiring**

Madeline Günther^{a,b,*}, Jürgen Drewe^c, Georg Boonen^c, Veronika Butterweck^c, Carsten Culmsee^{a,b}

^a Institute of Pharmacology and Clinical Pharmacy, Biochemisch-Pharmakologisches Centrum Marburg, Philipps-University Marburg, Karl-von-Frisch-Straße 2, 35032 Marburg, Germany

^b Center for Mind, Brain, and Behavior, 35032 Marburg, Germany

^c Preclinical Research, Max Zeller Soehne AG, Romanshorn, Switzerland

Mitochondria are the key organelles of energy metabolism and play critical roles in neurodegenerative diseases, in (neuro-)inflammation and also in processes of aging. Impaired mitochondrial metabolism featuring dysfunctional oxidative phosphorylation (OXPHOS) is regarded as a hallmark in paradigms of cellular dysfunction and death. Here, we investigated the effects of the *Cimicifuga racemosa* extract Ze 450 in settings of mitochondrial disorders with specific focus on metabolic alterations mediated by the extract.

Employing a systems approach with proteomics, metabolomics and metabolic flux characterization, we examined the effects of Ze 450 on the overall metabolic phenotype in different model systems of disease. Ze 450 demonstrated a robust ability to induce a metabolic switch in hippocampal HT22 cells, in RAW 264.7 macrophages and primary mouse embryonic fibroblasts. This shift was associated with a decrease in mitochondrial respiration and an increase in glycolytic activity to meet the cellular energy demands. Importantly, the metabolic rewiring was responsible for the resilience against oxidative stress, but also for the prevention of the LPS-induced inflammatory response in mouse macrophages. Moreover, the preserved mitochondrial integrity and reduced ROS levels following reduced mitochondrial respiration mediated by Ze 450, mitigated UV-induced cell death and senescence in MEF cells. Strikingly, we identified the glutamine-derived α -ketoglutarate anaplerotic flux to be critical for the beneficial effects of Ze 450 against ferroptosis and inflammation.

Our results highlight the importance of mitochondrial function and dynamics in regulating cell fate decisions and anti-inflammatory functions, but also the potential of *Cimicifuga racemosa* extract Ze 450 to interfere these processes.

Keywords: Ze 450, *Cimicifuga racemosa*, metabolic reprogramming, mitochondria, ferroptosis, inflammation, uv-induced cellular damage

* Presenting author. E-mail address: madeline.guenther@pharmazie.uni-marburg.de (Madeline Günther)

Phenolic-rich *Cratoxylum cochinchinense* leaves extract: a promising natural cosmeceutical

Sheri-Ann Tan^{a,*}, Zhi Xuan Lee^a, Mary Gaik Hong Khoo^b

^a Department of Bioscience, Faculty of Applied Sciences, Tunku Abdul Rahman University of Management and Technology, Setapak, Kuala Lumpur 53300, Malaysia

^b Natural Products Division, Forest Research Institute Malaysia (FRIM), Kepong, Selangor 52109, Malaysia

Human skin is frequently subjected to oxidative stress due to reactive oxygen species (ROS) and the lack of intracellular antioxidants. UV overexposure is one of the primary causes of oxidative stress, which accelerates inflammatory responses and the skin aging processes. Our previous work revealed that *Cratoxylum cochinchinense* leaves extract (CCE) possessed high total phenolic content along with strong antioxidative activities (Tan et al., 2022). In addition, mangiferin, a phenolic compound, was identified as the main phytochemical in CCE (Fig. 1).



Fig. 1. The leaves extract of *Cratoxylum cochinchinense* contained mangiferin as the main phytochemical

Hence, the aim of this study was to assess the potential of CCE as an anti-aging and sun protective agent. The anti-aging ability of CCE was examined using *in vitro* anti-tyrosinase and anti-elastase activity, while sunscreen ability was determined using *in vitro* SPF method. The anti-inflammatory efficacy of CCE was evaluated via a protein denaturation assay using bovine serum albumin (BSA). The plant extract was able to reduce tyrosinase activity by $38.95 \pm 0.86\%$ at $60 \mu\text{g/mL}$. Moreover, CCE at similar concentration demonstrated anti-elastase activity as effective as the positive control, ascorbic acid, by reducing elastase enzymatic activity by $62.11 \pm 4.17\%$. Moreover, a dose-dependent increment of SPF values was observed for CCE indicating its efficiency as a sunscreen agent. Anti-inflammatory effect was also demonstrated by CCE via inhibition of BSA denaturation. In conclusion, *Cratoxylum cochinchinense* leaves extract is indeed a promising cosmeceutical agent that could be incorporated into anti-aging skin care products.

Keywords: *Cratoxylum cochinchinense*, anti-aging, sun protective, anti-inflammatory, cosmetics

Reference

Tan, S.-A., Yam, H. C., Cheong, S. L., Chow, Y. C., Bok, C. Y., Ho, J. M., Lee, P. Y., Gunasekaran, B. 2022. Inhibition of *Porphyromonas gingivalis* peptidyl arginine deiminase, a virulence factor, by antioxidant-rich *Cratoxylum cochinchinense*: *In vitro* and *in silico* evaluation. Saudi J. Biol. Sci., 29(4), 2573–2581.

* Presenting author. E-mail address: tansw@tarc.edu.my (Sheri-Ann Tan)

Dual ^{13}C NMR and LC-MS² analysis for chemical profiling of Polynesian plant extracts

Marion Chambon^{a*}, Anthony Budai^b, Pedro Lameiras^b, Raimana Ho^a, Gaëtan Herbette^c, Agathe Martinez^b, Nicolas Borie^b, Simon Remy^b, Jean-Hugues Renault^b, Phila Raharivelomanana^a

^a UMR 241 SECOPOL, Université de Polynésie française, BP 6570, Faaa, Tahiti, French Polynesia

^b Université de Reims Champagne-Ardenne, CNRS, ICMR UMR7312, 51097 Reims, France

^c Aix Marseille Université, CNRS, Centrale Méditerranée, FSCM, Spectropole, Service 511, Campus Saint-Jérôme, Marseille, France

Five Polynesian plants among the most traditionally used in cosmetopeia and pharmacopeia were studied: *Calophyllum inophyllum*, *Gardenia taitensis*, *Curcuma longa*, *Cordia subcordata*, and *Ficus prolixa*. Their extracts were obtained by eco-extraction performed by ultrasound-assisted extraction (UAE) using ethanol:water (7:3, v/v) solvent. Liquid-liquid partition of crude extracts using stepwise gradients (heptane, EtOAc, n-butanol, and water) led to four extracts, and EtOAc extracts were fractionated by centrifugal partition chromatography (CPC).

The dereplicative chemical profiling method, combining ^{13}C NMR and LC-MS² analysis, was performed on CPC fractions. NMR analysis were carried out and hierarchical clustering analysis (HCA) was performed on ^{13}C signals with the Permut Matrix application. The resulting map included several clusters corresponding to molecular structures (Hubert *et al.*, 2014). Custom databases were created with VersaDB (Cordonnier *et al.*, 2024) and used to assign compound structures for each cluster. Right structures were then confirmed manually by 2D NMR analysis. LC-MS² data were processed via MZmine 3. Molecular networks were built using GNPS Feature-based molecular networking (Nothias *et al.*, 2020) and annotation was performed with SIRIUS.

The intelligent use of these two complementary methods allowed accelerated and efficient characterization of main compounds in complex mixtures. The known chemical composition of plant extracts such as *C. longa* and *C. inophyllum* allowed the validation of this process. Compounds identified and annotated for *F. prolixa* and *C. subcordata* were described for the first time in these two indigenous Polynesian tree extracts.

This powerful strategy led to the quick characterization of cosmetics with antioxidant and inflammatory properties.

Keywords: NMR; LC-MS²; centrifugal partition chromatography; molecular networking; dereplication

References

- Cordonnier, J., Remy, S., Renault, J. H., Nuzillard, J. M., 2024. Versa DB: Assisting ^{13}C NMR and MS/MS Joint Data Annotation Through On-Demand Databases. *Chemistry-Methods*. 4, e202300020.
- Hubert, J., Nuzillard, J. M., Purson, S., Hamzaoui, M., Borie, N., Reynaud, R., Renault, J. H., 2014. Identification of natural metabolites in mixture: a pattern recognition strategy based on ^{13}C NMR. *Analytical chemistry*. 86, 2955-2962.
- Nothias, L.F., Petras, D., Schmid, R., Dührkop, K., Rainer, J., Sarvepalli, A., ... & Dorrestein, P.C., 2020. Feature- based molecular networking in the GNPS analysis environment. *Nature methods*. 17, 905-908.

* Presenting author. E-mail address: marion.chambon@doctorant.upf.pf (Marion Chambon)

Chemical composition and nutritional value of the Sonora gum

Araceli-Perez Vasquez^a, Vanya Meneses^a, Laura Flores-Bocanegra^a, Valeria Reyes Pérez^a, Robert Bye^b, Rachel Mata^{a,*}

^a Facultad de Química, Universidad Nacional Autónoma de México, Cd. Mx. 04510, México

^b Instituto de Biología, Universidad Nacional Autónoma de México, Cd. Mx. 04510, México

The Sonora gum is an exudate produced by female insect *Tachardiella fulgens* Cockerell (Kerriidae), a host of the tree *Coursetia glandulosa* A. Gray (Fabaceae). The hard exudate (lac) encases the insect bodies and eggs and is valued as a medicinal and flavoring agent in the Northern gastronomy of Mexico. Previous investigations revealed the presence of lacaic acids in the lac, an unusual type of anthraquinones (Bisulca et al., 2018)

From an aqueous extract of the gum, a few specialized metabolites were isolated and characterized by spectrometric and spectroscopic analyses, including emodin, lacaic acid B, along with two new nerolidol derivatives, namely (10S,2E,6E)-10-hydroxy-2,6,10-trimetildodeca-2,6,11-trienoic acid and (2E,6E,10Z)-12-hydroxy-2,6,10-trimetildodeca-2,6,10-trienoic acid.

Headspace-solid-phase microextraction technique was employed to determine the volatile composition of the Sonora gum using four different coated fibers. The volatilome was characterized by a high content of vinyl-γ-valerolactone, lilac aldehyde D, α-curcumene, and nerolidol.

According to the Association of Official Analytical Chemists standard methods (AOAC, 2023), proximate analysis was carried out to determine the nutrient composition of the gum. The average protein, fat, carbohydrate, moisture, ash, and fiber contents were found to be 14.3, 27, 0.5, 0.9, and 53 %, respectively. The energy value was 200 kcal/100g. The gum was characterized by a high phosphorous content, a key element in the energetic metabolism; the other minerals were present in the amount required for a balanced diet. Niacin and thiamine were the most important vitamins. Thus, the Sonora gum is a good source of natural minerals and other constituents essential in the human diet.

Keywords: *Tachardiella fulgens*, *Coursetia glandulosa*, (10S,2E,6E)-10-hydroxy-2,6,10-trimetildodeca-2,6,11-trienoic acid and (2E,6E,10Z)-12-hydroxy-2,6,10-trimetildodeca-2,6,10-trienoic acid

References

- AOAC, Association of Official Analytical Chemists. International Official Methods of Analysis, Washington, DC, 22nd edition; 2023.
- Bisulca, C., Pool, M., & Odegaard, N., 2017. A survey of plant and insect exudates in the archaeology of Arizona. J. Archaeol. Sci. Rep.15, 272-281.

Acknowledgments

This work was supported by a grant of DGAPA IN203523

* Presenting author. E-mail address: rachel@unam.mx (Rachel Mata)

Cannabinoids and olive oil secoiridoids: a fascinating case of synergy

Evangelos Dadiotis^{a,e,*}, Lida Evmorfia Vagiaki^{b,c,e}, Aikaterini Papakonstantinou^d, Meng Cui^e, Vangelis Mitsis^f, Eleni Melliou^a, Diomedes Logothetis^e, Haralabia Boleti^d, Kyriaki Sidiropoulou^{b,c}, Prokopios Magiatis^a

^a Department of Pharmacy, National and Kapodistrian University of Athens, Greece

^b Department of Biology, University of Crete, Greece

^c Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology, Greece

^d Intracellular Parasitism Laboratory, Department of Microbiology, Hellenic Pasteur Institute, Athens, Greece

^e Department of Pharmaceutical Sciences, School of Pharmacy, Bouvé College of Health Sciences, Northeastern University, Boston, USA

^f Ekati Alchemy Lab SL, Barcelona, Spain

This study presents an investigation of the synergistic pharmacological effects of cannabinoids, from *Cannabis sativa*, and secoiridoids, found in olive oil. These two categories of natural compounds share common pharmacological targets. TRPA1 channels, are upregulated in numerous pathological conditions such as cancer and neurodegenerative diseases (Bosson et al., 2017; Takahashi et al., 2018), are a potential synergistic target for these compounds. The study initially explored the synergistic modulation of TRPA1 channels by cannabinoids and secoiridoids, revealing enhanced efficacy through different binding sites interactions. Specifically, a mixture of oleocanthal and cannabidiol in a ratio of 3:1 yielded the best results. *In vitro* MTT assay was utilized to investigate the synergistic cytotoxic effect of these compounds on various cancer cell lines, including breast, liver, and melanoma lines. The identified synergistic effect based on the coefficient of drug interactions ($CDI < 0.7$) led to the *in vivo* evaluation of the antitumor effect in MDA-MB-231 xenograft model of cancer (50 mg/kg CBD/ 5 mg/kg secoiridoids). Electrophysiological recordings in hippocampal brain slices of mice revealed a synergistic action in the decrease of seizure-like activity upon simultaneous perfusion of the two compounds. Additionally, employment of electrophysiological recordings uncovered an increase in long-term potentiation suggesting an enhancement in synaptic plasticity following their perfusion. *In vivo* behavioral experiments using an Alzheimer's disease mouse model (5xFAD), indicated improved performance in several cognitive tasks. The acute oral toxicity of a mixture of oleocanthal and cannabinoids was also evaluated in mice, with the combination being nontoxic in doses up to 300 mg/Kg.

Keywords: cannabinoids, secoiridoids, cannabidiol, oleocanthal, TRPA1

References

- Bosson, A., Paumier, A., Boisseau, S., Jacquier-Sarlin, M., Buisson, A., Albrieux, M., 2017. TRPA1 channels promote astrocytic Ca^{2+} hyperactivity and synaptic dysfunction mediated by oligomeric forms of amyloid- β peptide. *Mol Neurodegener* 12, 53.
- Takahashi, N., Chen, H.-Y., Harris, I.S., Stover, D.G., Selfors, L.M., Bronson, R.T., Deraedt, T., Cichowski, K., Welm, A.L., Mori, Y., Mills, G.B., Brugge, J.S., 2018. Cancer Cells Co-opt the Neuronal Redox-Sensing Channel TRPA1 to Promote Oxidative-Stress Tolerance. *Cancer Cell* 33, 985-1003.

* Presenting author. E-mail address: edadioti@pharm.uoa.gr (Evangelos Dadiotis)

***Lignosus rhinocerus* extract ameliorates glutamate-induced neuronal toxicity**

Tewin Tencomnao^{a,b,*}, Nattaporn Pattarachotanant^{a,b}, Anchalee Prasansuklab^c, Panthakarn Rangsinth^d, George Pak-Heng Leung^d, Siriporn Chuchawankul^e

^a Natural Products for Neuroprotection and Anti-Ageing (Neur-Age Natura) Research Unit, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^b Department of Clinical Chemistry, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^c College of Public Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^d Department of Pharmacology and Pharmacy, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China

^e Department of Transfusion Medicine and Clinical Microbiology, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

Lignosus rhinocerus (LR) extract has been widely used in traditional folk medicine. However, its potential in anti-neurodegeneration has not yet been fully elucidated. So, this study was aimed at investigating the protective effects of LR extract on glutamate-induced toxicity. Cell viability, immunofluorescence staining, flow cytometry and Western blot analysis were performed *in vitro* using mouse hippocampal neuronal HT-22 cells. Moreover, the ability of likely active ingredients from LC-MS binding to NMDA receptor was analysed using *in silico* docking. Both ethyl acetate (LR-EA) and ethanolic (LR-EE) extracts of LR effectively decreased glutamate-induced neuronal apoptosis and intracellular ROS accumulation. Both extracts could significantly attenuate mitochondrial dysfunction and mitophagy overactivation corresponding to the decreased mitochondrial depolarization, the increased ATP production, and the modulation of mitophagy related proteins (LC3 and p62/ SQSTM1) and LC3 colocalization. Moreover, both extracts possessed neuroprotective effect, demonstrated by significant suppression of glutamate-induced ER stress through the decrease of calcium influx, the ratio of p-ERK/ERK, and CHOP expression. In addition, phytochemicals of both extracts and their competence on inhibiting NMDAR were resolved. *In silico* analysis, 4 phytochemicals (canthin-6-one, magnoshinin, royleanone, and cernuine) from LR-EA and 2 phytochemicals (cernuine and levopimaric acid) from LR-EE revealed that their binding to NMDAR was more efficient than the binding of NMDAR antagonist, memantine. Likewise, cernuine could be the possible active constituent in both LR extracts with the strongest binding energy against NMDAR. In conclusion, LR is potentially classified as a neuroprotectant with cernuine as an energetic component for the prevention of glutamate-triggered neuronal toxicity.

Keywords: *Lignosus rhinocerus*, glutamate-triggered neuronal toxicity, HT-22 cells, Neuroprotectant, cernuine

* Presenting author. E-mail address: tewin.t@chula.ac.th (Tewin Tencomnao)

Novel health potential of isomers of 13²-hydroxypheophytin a from *Arthrospira platensis*

C. Filipe Henriques^{a,*}, Ana C. Fonseca^{a,b}, Mariana Reis^c, Ralph Urbatzka^a

^a Interdisciplinary Centre of Marine and Environmental Research, Biodiscovery for Health, 4400-208 Matosinhos, Porto, Portugal

^b University of Porto, Faculty of Sciences, 4169-007 Porto, Portugal

^c Interdisciplinary Centre of Marine and Environmental Research, Blue Biotechnology and Ecotoxicology and Health, 4400-208 Matosinhos, Porto, Portugal

Spirulina is widely commercialized for its fitness and health benefits, however the compounds responsible for said activity are unknown. This work covers a bioassay-guided isolation and structural elucidation of bioactive natural products from *Arthrospira platensis*. Zebrafish larvae were used for bioactivity screening throughout the isolation process. Structure elucidation of obtained most active isolates revealed chlorophyll a derivative isomers 13²-R- hydroxypheophytin a (**1**) and 13²-S-hydroxypheophytin a (**2**).

A DCM:MeOH (2:1) extract of *Arthrospira platensis* was submitted for subsequential fractionations using flash chromatography and HPLC with Hex:EtOAc mixtures. Each step was intercalated with lipid reducing assays until complete isolation of **1** and **2**. For structure elucidation, HRESIMS/MS and 2D NMR spectroscopy were employed using COSY, HSQC and ROESY for stereochemistry determination.

The bioactivity of both compounds was compared for lipid reduction and 2 food uptake assays in zebrafish larvae, and for the reversion of steatosis in fatty acid overloaded HepG2 cells. Both compounds reduced active feed uptake with similar IC₅₀ however, only compound **1** presented activity in passive food uptake (IC₅₀ = 22.5 µM), indicating putative hormonal and neuronal interactions. Isomer **2** presented instead 12x more lipid reducing potential with IC₅₀ = 1.18 µM, and 2x steatosis reversion capacity (IC₅₀ = 4.8 µM) than isomer **1**.

In summary, differences in bioactivity were revealed for both isomers of 13²- hydroxypheophytin a indicating the importance of conformational change in their mechanism of action. This knowledge will be important for future development of potential health applications.

Keywords: *Arthrospira platensis*, chlorophylls, obesity, appetite, steatosis

Acknowledgements:

This work was financially supported by “Pacto da Bioeconomia azul” (Project no. C644915664-00000026) withing the WP5 Algae Vertical, funded by Next Generation EU European Fund and the Portuguese Recovery and Resilience Plan (PRR).

* Presenting author. E-mail address: filiphenriquesbiotec@outlook.com (C. Filipe Henriques)

Green extraction *via* NaDES of waste and by-products of the Aglianico cv. vinery supply chain

Lara Comune*, Claudia Gravina, Simona Piccolella, Severina Pacifico

Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "Luigi Vanvitelli", Via Vivaldi 43, 81100 Caserta, Italy

In recent years, as part of sustainable development policies, the revaluation of end-of-life products has become more and more increased. Indeed, their effective reuse becomes pivotal as a starting point for the obtainment of high-added-value products (Caponio et al., 2022), especially in those supply chains (e.g., vinery), in which high amounts of waste are produced during the whole year, both in the field and transformation processes. In this context, although anthocyanins from the grape pomace have been previously proposed as active ingredients in cosmetics, herein a sustainable approach was used to recover them from the leaves and grape pomace of *Vitis vinifera* L. cv. Aglianico. Thus, the waste/by-products were subjected to maceration with Natural Deep Eutectic Solvents (NaDES), made by citric acid and sucrose (Paiva et al., 2014). The polyphenolic fraction, obtained after the removal of NaDES, was chemically analyzed using UHPLC-HR-MS/MS tools, and the biological activities were evaluated in cell-free and cell-based systems. Radical scavenging capacity was highlighted through ABTS and DPPH tests, while the cytotoxic screening, carried out on HaCaT cell line by MTT test, evidenced the absence of toxicity. Thus, wound healing properties were evaluated, suggesting a valuable applicability in the cosmeceutical sector.

Keywords: *Vitis vinifera* L., waste valorization, UHPLC-HRMS, NaDES, cosmeceuticals.

References

- Caponio, F., Piga, A., Poiana, M., 2022. Valorization of food processing by-products. *Foods* 11, 3246.
- Paiva, A., Craveiro, R., Aroso, I., Martins, M., Reis, R.L., Duarte, A.R.C., 2014. Natural Deep Eutectic Solvents – Solvents for the 21st Century. *ACS Sustainable Chem. Eng.* 2, 5, 1063–1071.

*Presenting author. E-mail address: lara.comune@unicampania.it (Lara Comune)

Bioassay-guided isolation of potential cancer chemopreventive agents from the fruits of *Rhodomyrtus tomentosa*

Chuanchom Khuniad^{a,b*}, Lutfun Nahar^c, Kenneth J. Ritchie^a, Satyajit D. Sarker^a

^a Centre for Natural Products Discovery (CNPD), School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Liverpool L3 3AF, United Kingdom ^bDepartment of Thai Traditional Medicine, Faculty of Health and Sports Science, Thaksin University, Phatthalung, 93210, Thailand

^c Laboratory of Growth Regulators, Palacký University and Institute of Experimental Botany, The Czech Academy of Sciences, Šlechtitelů 27, 78371 Olomouc, Czech Republic

Based on cancer morbidity and mortality rates, cancer is a major threat to human health.. Undoubtedly, cancer chemoprevention is one of the most important strategies for cancer control. *Rhodomyrtus tomentosa* (Aiton) Hassk. (Myrtaceae) is a Thai medicinal plant used traditionally for various diseases. Notably, phytochemicals from this species possess antioxidant, anti-inflammatory, anticancer and antimicrobial activities (Nwabor *et al.*, 2022). This study aimed to identify phytochemicals with cancer chemopreventive effect from *R. tomentosa* fruits following a bioassay-guided isolation approach. Dried ground fruits were sequentially Soxhlet-extracted with *n*-hexane, DCM and MeOH. Chemopreventive potential of crude extracts, fractions and isolated compounds was evaluated for Nrf2 induction potential using a cell-based luciferase assay in the AREc32 cell line. The MeOH and DCM extracts were the most active extracts with a 7.1-fold and 6.7-fold induction of luciferase activity (relative to control), respectively. As a result, the active MeOH and DCM extracts were selected for fractionation using SPE and VLC, respectively. Compounds were isolated from the most active fractions using TLC and reversed-phase HPLC. Structural elucidation of isolated compounds was achieved by extensive 1D and 2D NMR spectroscopic analyses. 3-*O*-Methylellagic acid 3'-*O*- α -rhamnopyranoside, a polyphenolic glycoside (Fig. 1), was isolated from the active MeOH fraction 2. Studies on the chemopreventive potential of this compound are under way. Similarly, several other compounds have also been isolated and their structures are under investigation. All these results will be presented during the oral presentation.




Fig. 1. Structure of 3-*O*-methylellagic acid 3'-*O*- α -rhamnopyranoside

Keywords: *Rhodomyrtus tomentosa*, myrtaceae, chemopreventive effect, bioassay-guided isolation

References

Nwabor, O.F., Singh, S., 2022. Systematic review on *Rhodomyrtus Tomentosa* (Aiton) Hassk: A potential source of pharmacological relevant bioactive compounds with prospects as alternative remedies in varied medical conditions. *Int. J. Pharm. Sci. Nanotech.* 15, 5875-5890.

* Presenting author. E-mail address: C.Khuniad@2020.ljmu.ac.uk (Chuanchom Khuniad)

Honeypot strategy for parasite weed control: An offer they cannot refuse

Francisco A. Macías*, Jesús G. Zorrilla, Carlos Rial, Antonio Cala, Francisco J. Rodríguez-Mejías, Rosa M. Varela, José M. G. Molinillo

Allelopathy Group, Department of Organic Chemistry, School of Science, Institute of Biomolecules (INBIO), University of Cadiz, C/ Avenida República Saharaui, nº 7, 11510 Puerto Real, (Cádiz), Spain

Parasitic weeds pose a significant threat to agricultural productivity, necessitating the development of sustainable control methods. The use of traditional herbicides has led to the emergence of resistant weed varieties, highlighting the urgent need for eco-friendly and efficient alternatives. Allelopathy presents promising avenues for exploration, such as the honeypot strategy (Macías et al., 2020), which focused part of our continuing research on this topic. A primary approach involved the discovery of bioactive metabolites produced by host plants (biocommunicators), leading to the identification and study of promising compounds isolated from host species, such as strigolactones, sesquiterpene lactones, or α -tomatine. Implicit in this task was the development of a simple and reliable UHPLC-MS/MS method for the simultaneous analysis of strigolactones in exudates and extracts. This method was tested with limits of quantifications ranging from 0.05 to 0.96 $\mu\text{g/L}$. A second approach focused on synthesizing parasitic plant stimulants as strigolactone mimics and analogous. Encouraging results were achieved by adding a butenolide ring to phthalimides, isolated sesquiterpene lactones, and diterpenoids as starting materials, optimizing previously reported synthetic strategies (Zorrilla et al., 2022). The isolated and synthesized compounds demonstrated stimulating activity in bioassays with seeds from various *Orobanch*e and *Phelipanche* species. Furthermore, a selection among the most promising compounds was encapsulated, yielding favorable results in bioassays. These findings introduce novel structures with the potential to serve as active ingredients in pre-emergence herbicides based on natural products, thereby stimulating further exploration and research into their efficacy and applicability.

Keywords: parasitic weeds, honeypot strategy, strigolactones, organic synthesis, encapsulations

References

- Macías, F.A., Durán, A.G., Molinillo, J.M.G., 2020. Allelopathy: The chemical language of plants. *Prog. Chem. Org. Nat. Prod.* 112, 1-84.
- Zorrilla, J.G., Rial, C., Varela, R.M., Molinillo, J.M.G., Macías, F.A., 2022. Strategies for the synthesis of canonical, non-canonical and analogues of strigolactones, and evaluation of their parasitic weed germination activity. *Phytochem. Rev.* 21, 1627-1659.

* Presenting author. E-mail address: famacias@uca.es (Francisco A. Macías).

A (sub)cellular mapping of natural substances

Miroslav Strnad^{a,*}, Vladimír Skalický^a, Aleš Pěňčík^a, Ioanna Antoniadi^b, Karin Ljung^b, Ondřej Novák^a

^a *Laboratory of Growth Regulators, Palacký University and Institute of Experimental Botany of the Czech Academy of Sciences, CZ-771 00 Olomouc, Czech Republic*

^b *Umeå Plant Science Centre, Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, SE-90183 Umeå, Sweden*

Phytohormones, preferentially auxins and cytokinins (CKs), control most aspects of plant growth, development and plasticity. Their distribution in plants has been described, but the importance of cell- and subcellular-type specific phytohormone homeostasis remains undefined. Recently, we revealed auxin and cytokinin distribution maps showing their different organelle-specific allocations within the Arabidopsis plant cells. More specifically, we revealed the intracellular distribution of auxin and CKs in nuclei, chloroplasts, mitochondria, endoplasmic reticulum (ER) and vacuoles. To achieve that, we introduced a breakthrough technique, so-called Fluorescence-Activated multi-Organelle Sorting (FAMOS), for the simultaneous fractionation of four organelle populations from a single sample. This time-efficient method is based on flow-cytometric principles, utilizes organelle-specific fluorophores and considers hormone metabolism stability. We combined our recently developed FAMOS tool with ultra-sensitive mass spectrometry-based methods. Apart from the four organelles isolated via FAMOS, vacuoles have been collected by density-gradient centrifugation to complete the subcellular hormone map. Altogether, this provides a unique approach for organelle-specific metabolic profiling of natural substances, auxin and CKs, combined with other “omics” applications such as proteomics and metabolomics. Shedding light on regulatory processes at the subcellular level will allow a deeper understanding of plant developmental processes and cellular homeostasis maintenance, as well as in intra- and intercellular communication.

Keywords: natural substances, phytohormones, auxin and cytokinin, (sub)cellular localization

References

- Novák, O., Napier, R. Ljung, K., 2017. Zooming in on plant hormone analysis: tissue- and cell-specific Approaches. *Ann. Rev. Plant Biol.* 68, 323-348.
- Pěňčík, A., Casanova-Sáez, R., Pilařová, V., Žukauskaitė, A., Pinto, R., Luis Micol, J. et al. 2018. Ultra-rapid auxin metabolite profiling for high-throughput mutant screening in Arabidopsis. *J. Exp. Bot.* 69, 2569-2579.
- Skalický, V., Antoniadi, I., Pěňčík, A., Chamrád, I., et al. 2023. Fluorescence-activated multi-organelle mapping of subcellular plant hormone distribution. *Plant J.* 116, 1825-1841.

* Presenting author. E-mail address: miroslav.strnad@upol.cz (Miroslav Strnad).

Utilizing natural products for weight management and longevity promotion

Milen I. Georgiev^{a,b,*}, Liliya V. Mihaylova^{a,b}, Martina S. Savova^{a,b}, Monika N. Todorova^a

^a Laboratory of Metabolomics, Institute of Microbiology, Bulgarian Academy of Sciences, 139 Ruski Blvd., 4000 Plovdiv, Bulgaria

^b Department of Plant Cell Biotechnology, Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

Obesity presents a significant challenge to modern society, imposing a substantial burden on health systems and individuals alike. Nowadays, more than a billion people globally are considered obese, a condition that does not solely refer to weight management but rather to mitigating its far-reaching consequences on healthspan and lifespan (Savova et al., 2023). Individuals diagnosed with obesity are predisposed to comorbidities including type 2 diabetes, cardiovascular disease, and metabolic complications often referred to as metabolic syndrome. Most importantly, obesity is associated with a reduction in disease-free years, an excess risk of premature death, and accelerated aging (Savova et al., 2023; Todorova et al., 2024a). On the other hand, we live in a rapidly aging world, with an expectancy of more than a 2 billion people aged 65 years or older by 2050, alongside a rising proportion of age-related diseases. Thus, both obesity and ageing are, *per se*, different sides of the same coin and represent healthcare burden on our society (Savova et al., 2023; Todorova et al., 2024a). The development of strategies targeting both of these processes now becomes a challenge for science. Central to this pursuit is the recognition that metabolic health serves as a cornerstone for both healthy weight maintenance and prolonged lifespan (Todorova et al., 2024a). Additionally, key molecular pathways attributed to nutrient signalling that are implicated in obesity progression, intersect with those fundamental to longevity, suggesting potential shared targets for intervention (Todorova et al., 2024a). Utilizing the model organism *Caenorhabditis elegans*, along with molecular pharmacology approaches, our research focuses on the discovery of natural products that target these shared pathways (Savova et al., 2021; 2023; Todorova et al., 2024b). Through our approach, we aim not only to mitigate the adverse effects of obesity but also to uncover novel strategies for promoting healthy aging and longevity (Todorova et al., 2024a).

Keywords: obesity, aging, longevity, natural products

References

- Savova M.S., Vasileva L.V., Mladenova S.G., Amirova K.M., Ferrante C., Orlando G., Wabitsch M., Georgiev M.I., 2021. *Ziziphus jujuba* Mill. leaf extract restrains adipogenesis by targeting PI3K/AKT signaling pathway. *Biomed. Pharmacother.* 141, 111934.
- Savova M.S., Mihaylova L.V., Tews D., Wabitsch M., Georgiev M.I., 2023. Targeting PI3K/AKT signaling pathway in obesity. *Biomed. Pharmacother.* 159, 114244.
- Todorova, M.N., Savova, M.S., Mihaylova, L.V., Georgiev, M.I., 2024a. Nurturing longevity through natural compounds: Where do we stand, and where do we go? *Food Frontiers*. 5, 267-310.
- Todorova, M.N., Savova, M.S., Mihaylova, L.V., Georgiev, M.I., 2024b. Icaritin improves stress resistance and extends lifespan in *Caenorhabditis elegans* through hsf-1 and daf-2-driven hormesis. *Int. J. Mol. Sci.* 25, 352.

* Presenting author. E-mail address: milengeorgiev@gbg.bg (Milen I. Georgiev).

Treatment strategies of pulmonary fibrosis using natural products

Yun-Sil Lee*

School of Pharmaceutical Sciences, Ewha Womans University, Seoul, Korea

Naturally occurring compounds have been shown to be non-toxic over wide dose ranges and are inexpensive and effective for treating various diseases. Idiopathic pulmonary fibrosis (IPF) is a chronic and irreversible lung disease characterized by limited treatment options, creating an urgent need for advancements in identifying new therapeutic targets. Currently, the Food and Drug Administration has only approved pirfenidone and nintedanib for IPF treatment. Although these drugs can delay disease progression and alleviate symptoms, they do not offer a cure for IPF or improve survival rates. Furthermore, they come with side effects like thrombocytopenia, gastrointestinal discomfort, and dermatological reactions. Consequently, there is a pressing medical need to develop new drugs targeting novel pathways to treat IPF. In this presentation, I will primarily focus on exploring the involvement of novel treatment targets in pulmonary fibrosis and showed the treatment strategy using natural products.

Keywords: pulmonary fibrosis, natural products, novel targets, non-toxic

References

- Jeon, S.K., Jin, H., Kim, J.M., Hur, Y., Song, E.J., Lee, Y.J., Na, Y., Cho, J., Lee, Y.S., 2023. The miR-15b-Smurf2-HSP27 axis promotes pulmonary fibrosis. *J. Biomed. Sci.* 30, 2.
- Choi, S.K., Hwang, S., Jeon, S.K., Yoo, H., Lee, J., Shin, J., Na, Y., Kwon, Y., Lee, Y.S., 2023. Design, synthesis, and biological evaluation of novel HSP27 inhibitors to sensitize lung cancer cells to clinically available anticancer agents. *Bioorg. Chem.* 130, 106260.
- Yoo, Y., Jeon, S.K., Won, H., Jeong, M., Cho, Y.S., Hwang, E.S., Na, Y., Cho, J., Lee, Y.S., 2023. Drug like HSP27 cross linkers with chromenone structure ameliorates pulmonary fibrosis. *Front. Pharmacol.* 14, 1203033.
- Jin, H., Yoo, Y.J., Kim, Y., Kim, Y., Cho, J., Lee, Y.S., 2020. Radiation-induced lung fibrosis: preclinical animal models and therapeutic strategies. *Cancers* 12, 1562.
- Kim, J.Y., Jeon, S., Yoo, Y.J., Hin, H., Won, H.Y., Hwang, E.S., Lee, Y.J., Na, Y., Cho, J., Lee, Y.S., 2019. The HSP27-mediated I κ B α -NF κ B signaling axis promotes radiation-induced lung fibrosis. *Clin. Cancer Res.* 25, 5364-5395.

* Presenting author. E-mail address: yslee0425@ewha.ac.kr (Yun-Sil Lee).

Coding metabolic profile of medicinal herbs using 2D-HPTLC

Kyung Ho Lee^{a,b}, Minsik Choi^{b,c}, Young Pyo Jang^{a-c,*}

^a Department of Integrated Drug Development and Natural Products, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

^b Department of Pharmacognosy, College of Pharmacy, Kyung Hee University, Seoul 02447, Republic of Korea

^c Department of Biomedical and Pharmaceutical Sciences, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

The technology for classifying organisms through their genetic information is now universally utilized through gene coding (Zhu et al., 2022). However, classification technology using the metabolome of organisms is still in its very early stages. This is because it is challenging to find appropriate analytical techniques and methods that can effectively and universally represent and utilize metabolome profiles. In this presentation, we propose a systematic coding system that can universally code the metabolites of medicinal plants using 2-D HPTLC technology, and apply this system for the origin identification and quality control of medicinal plants. We determine the optimal 2-D development conditions that can comprehensively separate plant metabolites of various structures and obtain reproducible 2-D HPTLC chromatograms in the same analytical environment, including the development conditions. By visualizing the obtained chromatogram in various ways, we aim to encode the presence or absence of bands in each region of the TLC plate and assign a 2D-HPTLC code to each plant. In this presentation, we will introduce this system, discuss potential issues, and propose solutions to overcome them.

Keywords: species identification, metabolic coding, 2D-HPTLC, quality control

References

Zhu, Shuang. et al., 2022. DNA barcoding: an efficient technology to authenticate plant species of traditional Chinese medicine and recent advances. *Chinese Med.* 17, 112.

* Presenting author. E-mail address: ypjang@khu.ac.kr (Young Pyo Jang).

Status and prospect of natural deep eutectic solvents in current natural products research

Young Hae Choi^{a,*}, Samir Scandar^{a,b}, Natali Rianika Mustafa^a, Maria Carla Marcotullio^b, Robert Verpoorte^a

^a *Natural Products Laboratory, Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE, Leiden, The Netherlands*

^b *Department of Pharmaceutical Sciences, University of Perugia, 06122, Perugia, Italy*

Mixing salts and/or organic compounds can significantly reduce the melting point, converting them into liquids even at extremely low temperatures. The exploration of ionic liquids (ILs) began in the early 20th century with the investigation of the physical properties of ethylammonium nitrate. However, it is only in recent years that ILs and deep eutectic solvents (DES) have been reconsidered by chemical engineering for their potential as environmentally friendly mediums in chemical processing. Despite being hailed as promising candidates for green processing, ILs and DES are often composed of ingredients, predominantly synthesized chemically, that pose toxicity risks to both humans and the environment. Overcoming this challenge remains imperative for the widespread adoption of these solvents. Inspired by synthetic ILs, our research group has postulated the existence of a third type of liquid, distinct from water and lipids, hypothesizing that metabolites abundant in cells may give rise to such a medium (1, 2). Analysis of plant metabolomics data collected over several years revealed a striking similarity with synthetic ILs. This revelation challenges the long-standing notion that water is the sole medium in nature. This hypothesis could provide explanations for numerous scientific inquiries, including the low solubility of natural products, the mysteries surrounding biosynthesis, transportation, and storage of non-water-soluble metabolites, among others. Drawing from these insights and the mechanisms of ILs or DES, we propose the existence of a diverse array of natural deep eutectic solvents (NADES) in nature, comprising ILs or DES whose constituents are derived from natural sources. As an initial step to substantiate this concept, we experimented with various combinations of these potential candidates, resulting in the discovery of over 200 formulations yielding viscous liquids. Since the initial publication on NADES in 2011, a plethora of NADES have found applications across diverse fields of natural product research, including extraction for enhancing the concentration of bioactive molecules, serving as green reaction media for organic or enzymatic synthesis, stabilizing labile compounds, and formulating pharmaceuticals. In this presentation, I aim to showcase the diverse applications of natural products, focusing particularly on recent advancements such as their utilization as diffusion matrices for plant volatiles, while also addressing associated limitations.

Keywords: eutectic solvents, NADES, green technology, natural products

References

- Choi, Y.H., Van Spronsen, J., Dai, Y., Verberne, M., Hollmann, F., Arends, I.W.C.E., Witkamp, G.-J., Verpoorte, R., 2011. Are natural deep eutectic solvents the missing link in understanding cellular metabolism and physiology? *Plant Physiol.* 156, 1701-1705.
- Dai, Y., Van Spronsen, J., Witkamp, G.-J., Verpoorte, R., Choi, Y.H., 2013. Natural deep eutectic solvents as new potential media for green technology. *Anal. Chim. Acta* 766, 61-68.

* Presenting author. E-mail address: y.choi@chem.leidenuniv.nl (Young Hae Choi).

Discovery and biosynthesis of specialized metabolites from human gut microbiome

Chung Sub Kim*

School of Pharmacy and Department of Biopharmaceutical Convergence, Sungkyunkwan University, Suwon 16419, Republic of Korea

Gut microbes are considered as important contributors to human physiology and disease states, and they in the intestinal tract produce small molecule metabolites, which regulate diverse physiological responses (Donia and Fischbach, 2015). However, most of current research topics are focused on well-known gut bacteria, such as *Escherichia coli*, *Akkermansia muciniphila*, *Bacillus cereus*, *Clostridium* spp., *Enterococcus* spp., *Lactobacillus* spp., *Bacteroides* spp., and others. Other infrequent gut bacteria have relatively been neglected and therefore the majority of their metabolites remain structurally and functionally uncharacterized. *Aneurinibacillus aneurinilyticus* is a human gut bacterium, which might be involved in intestinal disease pathogenesis such as irritable bowel syndrome (Rajilić-Stojanović et al., 2011). This human bacterium has also been found from mouse lung tumors and showed probiotic effect in fish (Jin et al., 2018). Although they imply potential correlation between this bacterium and human health, the bacterial small molecule mediators remain largely unknown. In this presentation, we discover an angiogenesis inhibitory metabolite termed the “aneuristatin”, which possesses unusual pyrrolo[1,2-a]pyrazine scaffold, from *A. aneurinilyticus* ATCC 12856 along with seven structurally related metabolites. We showed that aneuristatin is biosynthesized from three L-tyrosine molecules by stable isotope feeding experiment, *in vitro* reconstitution, and organic synthesis. Aneuristatin showed anti-angiogenesis activity in the mouse and zebra fish models. Our studies provide a molecular foundation for *A. aneurinilyticus*’s potential ability to prevent and treat hypervascularized diseases including cancer.

Keywords: human microbiome, gut bacteria, *Aneurinibacillus aneurinilyticus*, pyrrolo[1,2-a]pyrazine, anti-angiogenesis

References

- Donia, M.S., Fischbach, M.A., 2015. Small molecules from the human microbiota. *Science* 349, 1254766.
- Jin, C., Lagoudas, G.K., Zhao, C., Bullman, S., Bhutkar, A., Hu, B., Ameh, S., Sandel, D., Liang, X.S., Mazzilli, S., Whary, M.T., Meyerson, M., Germain, R., Blainey, P.C., Fox, J.G., Jacks, T., 2018. Commensal microbiota promote lung cancer development via $\gamma\delta$ T Cells. *Cell* 176, 998–1013.
- Rajilić-Stojanović, M., Biagi, E., Heilig, H.G.H.J., Kajander, K., Kekkonen, R.A., Tims, S., de Vos, W.M., 2011. Global and deep molecular analysis of microbiota signatures in fecal samples from patients with irritable bowel syndrome. *Gastroenterology* 141, 1792–1801.

* Presenting author. E-mail address: chungsub.kim@skku.edu (Chung Sub Kim).

Posters

Marine organisms, fungi, microbes and plants as chemical factories

S1.P1 Anti-cancer effect of *Willughbeia edulis* Roxb. stem extract in hepatocellular carcinoma cells through triggering apoptosis and G2/M cell cycle arrest

Thi Tuyet Nhung Nguyen^{a,b,c,*}, Yu-Chi Lin^b, Ching-Kuo Lee^a, Jing-Jy Cheng^{a,b}

^a Ph.D. Program in Clinical Drug Development of Herbal Medicine, College of Pharmacy, Taipei Medical University, Taipei, Taiwan

^b National Research Institute of Chinese Medicine, Taipei, Taiwan

^c Faculty of Pharmacy, Lac Hong University, Dong Nai, Vietnam

The medicinal plant *Willughbeia edulis* Roxb. (Apocynaceae), native to Vietnam, has a long history of medicinal uses by traditional doctors in the treatment of liver cancer (Chi, 2012). Our research aims to investigate the *in vitro* cytotoxic effect against liver cancer cells of *W. edulis* (WE) stem extract and apply Bioactive Molecular Networking (BMN) for guided isolation of bioactive components from WE stem extract. Both the crude extract and butanol layer (WEB) showed selective and significant cytotoxic effects on HepG2 cells, with IC₅₀ values of 27.1 ± 2.7 and 26.06 ± 1.3 $\mu\text{g/mL}$, respectively. Furthermore, the cell cycle progression was analyzed by utilizing a flow cytometer following the treatment of cells with 16 and 32 $\mu\text{g/mL}$ of crude extract and WEB for 24 or 48 hours. The results found that they induced apoptosis and G2/M cell cycle arrest in HepG2 cells. Then WEB was fractionated using Diaion HP-20 chromatography column, yielding five fractions. These fractions were subjected to LC-MS/MS analysis and screened cytotoxicity assays. The obtained mass spectrum and cytotoxicity data were applied to bioactive molecular networking workflow (Nothias et al., 2018), leading to the isolation of five known compounds from the butanol layer of the stem of *W. edulis*. These isolates, cinchonain Ia, cinchonain Ib, and cinchonain IIa, with high bioactive scores in the molecular network, exhibited potent cytotoxic effects in HepG2 cells with IC₅₀ values of 39.4 ± 3.2 , 38.5 ± 2.6 , and 17.0 ± 1.5 μM , respectively.

Keywords: *Willughbeia edulis* Roxb., liver cancer, G2/M cell cycle arrest, bioactive molecular networking

References

Chi, Vo Van. 2012. Dictionary of Vietnamese Medicinal Plants. Medical Publishing House 1,2.

Nothias, Louis-Félix, Mélissa Nothias-Esposito, Ricardo Da Silva, Mingxun Wang, Ivan Protsyuk, Zheng Zhang, Abi Sarvepalli, Pieter Leyssen, David Touboul, and Jean Costa. 2018. Bioactivity-Based Molecular Networking for the Discovery of Drug Leads in Natural Product Bioassay-Guided Fractionation. *Journal of Natural Products* 81 (4): 758–67.

* Presenting author. E-mail address: d339109002@tmu.edu.tw (Nguyen, Thi Tuyet Nhung)

S1.P2 Chemical constituents isolated from leaves of seashore plant *Aglaia rimosa*

Jing-Ru Weng^{a,*}, Jui-Hsin Su^b, Ping-Jyun Sung^b, Wei-Yu Lin^c, Zhen-Hong Han^a, You-Ying Chen^b, Chun-Lin Chen^d

^a Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804, Taiwan, ROC

^b National Museum of Marine Biology and Aquarium, Pingtung 944, Taiwan, ROC

^c Department of Pharmacy, Kinmen Hospital, Kinmen 891, Taiwan, ROC

^d Department of Biological Science, National Sun Yat-sen University, Kaohsiung 804, Taiwan, ROC

Multiple pieces of evidence have shown that there are various types of compounds including amides, flavonoids, lignans, steroids, triterpenoids, rocaglamide, and dammarane in the *Aglaia* genus, and some of these compounds exhibit diverse biological activities (Harneti et al., 2021; Bueno Pérez et al., 2014). Eleven compounds, stigmasta-5,22E-diene-3 β ,7 β -diol (**1**), stigmasta-5,22-diene-3 β ,7 α -diol (**2**), stigmasta-5-ene-3 β ,7 α -diol (**3**), 7 β -hydroxysitosterol (**4**), 5 α ,8 α -epidioxy-24-methyl-cholesterol-6,22-dien-3 β -ol (**5**), 4',5,7-tri-O-methyl-kaempferol (**6**), 5,6-desmethylenedioxy-5-meth-oxy-aglallactone (**7**), yangambin (**8**), eudesmin (**9**), 2,4-dimethoxy-6-hydroxyl benzoic acid (**10**) and dehydroodorin (**11**), were isolated from the acetone extract of leaves of the seashore plant *A. rimosa*, a member of the Meliaceae family. Among these compounds, six compounds (**2** and **5-9**) were subjected to anticancer activity testing against oral cancer cells (SCC2095), breast cancer cells (MCF-7), and gastric cancer cells (SC-M1). Compound **5** exhibited cytotoxic activity against MCF-7 cells (IC₅₀ = 11.2 $\pm$ 1.3 μ M), while **6** demonstrated cytotoxicity against SCC2095, MCF-7, and SC-M1 cell lines (IC₅₀ = 16.3 $\pm$ 2.9, 24.8 $\pm$ 0.9, and 14.6 $\pm$ 0.7 μ M, respectively) (Fig. 1). These findings highlight the potential of compounds from *A. rimosa* as promising candidates for further exploration in cancer therapy.




Fig. 1. Structures of compounds **5** and **6**.

Keywords: *Aglaia rimosa*, cytotoxicity, oral cancer cells, breast cancer cells, gastric cancer cells

References

- Bueno Pérez, L., Still, P.C., Naman, C.B., Ren, Y., Pan, L., Chai, H.B., Carcache de Blanco, E.J., Ninh, T.N., Thanh, B. V., Swanson, S.M., Soejarto, D. D., Kinghorn, A. D., 2014. Investigation of Vietnamese plants for potential anticancer agents. *Phytochem. Rev.*, 13, 727-739.
- Harneti, D., Supratman, U., 2021. Phytochemistry and biological activities of *Aglaia* species. *Phytochemistry*, 181, 112540.

* Presenting author. E-mail address: jrweng@mail.nsysu.edu.tw (Jing-Ru Weng)

S1.P3 Fragment-guided genome mining leads to unprecedented structural complexity in natural products

Yalong Zhang^a, Lin Wu^a, Tyler A. Kerr^b, Yi Tang^{a,b,*}

^a Department of Chemical and Biomolecular Engineering, University of California, Los Angeles, Los Angeles, California 90095, USA

^b Department of Chemistry and Biochemistry, University of California, Los Angeles, Los Angeles, California 90095, USA

Here we reported the discovery and characterization of an unprecedented 6/12/6/6/6/6/6/6 fused octacyclic cyclophane-containing alkaloid, octacyclin A (1), by fragment-guided genome mining. The biosynthesis pathway of 1 involving 14 genes was completely unraveled by a combination of heterologous expression, feeding experiments and biochemical assays, showing that the complicated ring system was constructed by two single-module non-ribosomal peptide synthetases (NRPSs) and three cytochrome P450s. Briefly, NRPS OcyN with a noncanonical terminal condensation (CT) domain showed a novel function by modifying the piperazine cyclophane moiety to form a tetrahydroquinolin-4(1H)-one moiety. Subsequently, two P450s OcyM and OcyL mediated cascade cyclization with a cryptic protection process to forge the octacyclic system.

Keywords: oxidative cyclization, genome mining

* Presenting author. E-mail address: yitang@g.ucla.edu (Yi Tang)

S1.P4 OSMAC approach, the tool to unlock the biosynthetic potential of deep-sea actinobacteria

Sofia Correia^{a,*}, Inês Ribeiro^b, Andreia Braga-Henriques^c, Ralph Urbatzka^b, Lada Lukić Bilela^d, Pedro Leão^b, Maria F. Carvalho^b

^a ICBAS - Institute of Biomedical Sciences Abel Salazar, University of Porto, Porto, Portugal

^b CIIMAR- Interdisciplinary Centre of Marine and Environmental Research, Matosinhos, Portugal

^c The Regional Secretariat for the Sea and Fisheries, Government of the Azores, Horta, Portugal

^d UNSA - University of Sarajevo, Faculty of Science, Sarajevo, Bosnia and Herzegovina

Research on natural products has played a major role in the discovery of new drugs. The class Actinomycete comprises a group of microorganisms highly prolific in the production of NP (natural products) with several relevant bioactivities (Murtaza et al., 2023). It is known that actinobacteria include in their genome multiple biosynthetic gene clusters (BGC), meaning that a single strain may be capable of producing several NP. In the recent years, the OSMAC (One Strain - Many Compounds) strategy has revealed very promising in activating the expression of dormant BGC by varying culture conditions (Le Loarer et al., 2023), and its application has allowed the discovery of novel bioactive compounds (Hug et al., 2018; Hussain et al., 2017). In this study, the OSMAC approach was applied to four actinobacterial strains isolated from deep-sea samples collected at Madeira and Azores islands, in order to find novel bioactive compounds with pharmaceutical relevance. Three different culture conditions were applied to each strain and the capacity of these conditions to induce different bioactivities was analyzed. The results showed that the conditions applied stimulated significant antimicrobial and anticancer activities in the analyzed strains. The bioactive actinobacterial extracts were subjected to a metabolomics analysis, included mass spectrometry-based dereplication and molecular networking analyses. These results revealed the presence of 5 possible new compounds. We are currently performing guided mass isolation of these potential new NP. This study shows that the OSMAC approach is a promising strategy for the discovery of new molecules with biotechnological potential.

Keywords: Actinomycete, OSMAC approach, Bioprospecting, secondary metabolites, pharmaceutical potencial

References

- Hug, J.J., Bader, C.D., Remškar, M., Cirnski, K., Müller, R., 2018. Concepts and methods to access novel antibiotics from actinomycetes. *Antibiotics* 7, 44.
- Hussain, A., Rather, M.A., Dar, M.S., Aga, M.A., Ahmad, N., Manzoor, A., Qayum, A., Shah, A., Mushtaq, S., Ahmad, Z., Hassan, Q.P., 2017. Novel bioactive molecules from *Lentzea violacea* strain AS 08 using one strain-many compounds (OSMAC) approach. *Bioorg. Med. Chem. Lett.* 27, 2579-2582.
- Le Loarer, A., Dufossé, L., Bignon, J., Frédérick, M., Ledoux, A., Fouillaud, M., Gauvin-Bialecki, A., 2023. OSMAC Method to Assess Impact of Culture Parameters on Metabolomic Diversity and Biological Activity of Marine-Derived Actinobacteria. *Mar. Drugs* 22, 23.
- Murtaza, M., Abrol, V., Nehra, E., Choudhary, P., Singh, S.K., Jaglan, S., 2024. Biodiversity and Bioactive Potential of Actinomycetes from Unexplored High Altitude Regions of Kargil, India. *Indian J. Microbiol.* 64, 110-124.

* Presenting author. E-mail address: sifcorreia98@gmail.com (Sofia Correia)

S1.P5 Antiprotozoal activity of alkaloid fractions and isolation of four new steroidal alkaloids from *Buxus obtusifolia*

Justus Mukavi^{a,*}, Marcel Kaiser^b, Pascal Mäser^b, Leonidah Kerubo Omosa^c, Thomas J. Schmidt^a

^a Institute of Pharmaceutical Biology and Phytochemistry (IPBP), University of Münster, PharmaCampus, Corrensstraße 48, D-48149 Münster, Germany

^b Swiss Tropical and Public Health Institute, Kreuzstraße 2, CH-4123 Allschwil, Switzerland

^c Department of Chemistry, Faculty of Science and Technology, University of Nairobi, P.O. Box 30197-00100, Nairobi, Kenya

As part of our continuing search for new antiprotozoal compounds derived from Buxaceae family (Szabó et al., 2021), we investigated the dichloromethane extract from the leaves of the East African *Buxus obtusifolia* (Mildbr.) Hutch. *B. obtusifolia* is an evergreen shrub utilized ethnomedicinally in many parts of E. Africa to treat chest-related ailments (Beentje et al., 1994). Fractionation of the alkaloid enriched fraction by centrifugal partition chromatography (CPC) yielded a total of 16 sub-fractions. The anti-protozoal activity of the crude extract, the alkaloid-enriched fraction, and the 16 CPC subfractions was assessed against *Trypanosoma brucei rhodesiense* (*Tbr*) and *Plasmodium falciparum* (*Pf*) using established standard protocols (Szabó et al., 2021). The alkaloid fraction displayed stronger activities against *Tbr* and *Pf* (IC₅₀ = 8.07 and 0.69 µg/mL, respectively), compared to the crude extract (IC₅₀ = 16.05 and 1.08, respectively). Sub-fractions 3 (IC₅₀ = 0.76 µg/mL) and 4 (IC₅₀ = 0.77 µg/mL) demonstrated the highest antitrypanosomal activity. Additionally, promising antiplasmodial effects with IC₅₀ values ranging from 0.93 – 1.89 µg/mL were observed for fractions 3 to 16. Bioassay-guided separation of CPC sub-fractions 3 – 5 using prep-HPLC resulted in the isolation of four new steroidal alkaloids including cycloprotobuxoline-C (1), cycloprotobuxoline-C N-oxide (2), 3-aminocycloprotobuxoline-C (3), dehydrocyclovirobuxeine-B (4) along with the known cyclovirobuxeine-A (5) (Atta-ur-Rahman et al., 2001) (Fig. 1). *In vitro* antiprotozoal efficacy studies of the isolated compounds are underway.




Fig. 2: Structures of compounds 1 – 5 isolated from *Buxus obtusifolia* (Mildbr.) Hutch.

Keywords: *Buxus obtusifolia*, Buxaceae, steroidal alkaloids, *Trypanosoma brucei rhodesiense*, *Plasmodium falciparum*

References

- Atta-ur-Rahman, Parveen, S., Khalid, A., Farooq, A., Choudhary, M. I., 2001. Acetyl and butyrylcholinesterase-inhibiting triterpenoid alkaloids from *Buxus papillosa*. *Phytochemistry*, 58, 963–968.
- Beentje, H., Adamson, J., & Bhandari, D., 1994. Kenya trees, shrubs, and lianas. National Museums of Kenya. <https://agris.fao.org/agris-search/search.do?recordID=US201300288929>
- Szabó, L.U., Kaiser, M., Mäser, P., Schmidt, T.J., 2021. Antiprotozoal Nor-Triterpene Alkaloids from *Buxus sempervirens* L. *Antibiotics*, 10, 696.

* Presenting author. E-mail address: justusmukavi32@gmail.com (Justus Mukavi)

S1.P6 Genome mining-based discovery of new glycosylated terpenoids from *Micromonospora* sp. isolated from *Leontopodium nivale* subsp. *alpinum* rhizosphere

Jaime Felipe Guerrero-Garzón^{a,*}, Una Bajric^a, Martin Zehl^b, Rasmus Kirkegaard^{c,d}, Sergey B. Zotchev^a

^a Division of Pharmacognosy, University of Vienna, Vienna 1090, Austria

^b Department of Analytical Chemistry, Faculty of Chemistry, University of Vienna, Vienna 1090, Austria

^c Centre for Microbiology and Environmental Systems Science, University of Vienna, Vienna, Austria ^d Joint Microbiome Facility of the Medical University of Vienna and the University of Vienna, Vienna, Austria

Representatives of the bacterial genus *Micromonospora* have been isolated from rhizosphere soil of the rare medicinal plant *Leontopodium nivale* subsp. *alpinum* (Oberhofer et al., 2019). Genome sequencing and analysis of one of the *Micromonospora* isolates led to the identification of a gene cluster presumably specifying biosynthesis of a glycosylated terpenoid. Transformation-associated recombination-assisted assembly of this cluster in yeast and its heterologous expression in an engineered *Streptomyces* host resulted in the production of a novel brasiliardin-related terpenoid which, however, was not glycosylated. At the same time, targeted screening of extracts from the native *Micromonospora* host strain by means of LC-MS allowed the identification of glycosylated congeners of this terpenoid. The latter suggested that the heterologous *Streptomyces* host is deficient in either biosynthesis of particular sugars or in expression of the cluster-specific glycosyltransferases. To alleviate this problem, several plasmids containing different cassettes for the biosynthesis of activated deoxysugars were introduced into the recombinant *Streptomyces* host harboring the terpenoid cluster. Analysis of the extracts from these recombinant strains revealed the production of several novel glycosylated terpenoid analogs. Currently, different approaches are being explored aimed at producing various glycosylated terpenoids via combinatorial biosynthetic engineering of this gene cluster.

Keywords: genome mining, glycosylated terpenoids, *Micromonospora*, combinatorial biosynthetic engineering, heterologous expression

References

Oberhofer, M., Hess, J., Leutgeb, M., Gössnitzer, F., Rattei, T., Wawrosch, C., Zotchev, S.B., 2019. Exploring actinobacteria associated with rhizosphere and endosphere of the native alpine medicinal plant *Leontopodium nivale* subspecies *alpinum*. *Front. Microbiol.* 8;10:2531.

*Presenting author. E-mail address: jaime.felipe.guerrero.garzon@univie.ac.at (Jaime Felipe Guerrero-Garzón)

S1.P7 Changing paradigms in natural product discovery: A molecule to microbe approach

Paul R. Jensen*, Tadeusz Molinski, Alexander Bogdanov

Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA USA

Traditional approaches to microbial natural product discovery take a microbe first approach in which individual strains are cultured in the lab and bioassays used to guide compound isolation. While once productive, the limitations to this approach are now well documented and include the recognition that only a small percentage of the bacteria present in the environment are readily obtained in culture. We have developed a new approach to microbial natural product discovery in which compounds are isolated directly from the environments in which they are produced thus bypassing the initial need for laboratory cultivation. This culture independent approach, which we call the Small Molecule In Situ Resin Capture (SMIRC) technique, is agnostic to the biological source of the compounds and requires no up-front knowledge of cultivation requirements or the cues needed to induce biosynthesis. SMIRC deployments in marine habitats yielded extensive chemical diversity including compounds previously reported from marine plants, invertebrates, and bacteria. Mining compounds that could not be identified has yielded unprecedented carbon skeletons and demonstrated that sufficient yields can be obtained for NMR-based structure elucidation. These results, along with “omic” efforts to identify the producing organisms, suggest a path forward to access chemical space natural product diversity without the need for cultivation.

Keywords: natural product discovery

* Presenting author. E-mail address: pjensen@ucsd.edu (Paul R. Jensen)

S1.P8 Bioactive scalarane sesterterpenoids and polybrominated diphenyl ethers from the two sponges *Hippospongia* sp. and *Dysidea* sp.

Chen You-Ying^a, Sung Ping-Jyun^a, Weng Jing-Ru^b, Chun-Lin Chen^c, Su Jui-Hsin^{a,*}

^a National Museum of Marine Biology and Aquarium, Pingtung, Taiwan

^b Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung, Taiwan

^c Department of Biological Sciences, National Sun Yat-sen University, Kaohsiung, Taiwan

One new scalarane sesterterpenoid (**1**), along with four known sesterterpenoids (**2–5**), were isolated from the sponge *Hippospongia* sp. Five known polybrominated diphenyl ethers natural compounds (**6–10**) were isolated from the sponge *Dysidea* sp. The structure of marine natural products **1–10** was established by NMR spectral data analysis. Moreover, the structure of brominated diphenyl ethers **6**, **9** and **10** was further confirmed by X-ray single-crystal diffraction analysis (Fig. 1).



Fig. 1. Molecular structures of **9** and **10** based on X-ray analysis.

The study assessed the cytotoxicity of compounds **1–10** against human cholangiocellular carcinoma (HuCCT1) and human colonic adenocarcinoma (SW620). The findings indicated that sesterterpenoids **2** and **5** demonstrated substantial cytotoxic effects on these cancer cell lines. They significantly reduced cell viability and promoted apoptosis in a dose-dependent manner. Furthermore, the application of these compounds was associated with an increased expression of apoptosis-related proteins, particularly caspase-3, caspase-7, and PARP, highlighting their potential as therapeutic agents.

Keywords: sesterterpenes, polybrominated diphenyl ethers, sponge, cytotoxicity

* Presenting author. E-mail address: x2219@nmmba.gov.tw (Su Jui-Hsin)

S1.P9 New triterpenoid saponins from the aerial parts of *Glinus oppositifolius* and their anti-adipogenic activity.

Kim-Hoa Phi^{a,b}, Min-Ha Park^c, Seulah Lee^d, Man Hyung Koo^e, Sung Suk Suh^c, Ui Joung Youn^{a,b,*}

^a Division of Life Sciences, Korea Polar Research Institute, Incheon 21990, Republic of Korea

^b University of Science and Technology (UST), Daejeon 34113, Republic of Korea

^c Department of Bioscience, Mokpo National University, Jeonnam 58554, Republic of Korea

^d College of Life Sciences, Kyung Hee University, Yongin 17104, Republic of Korea

^e Research Unit of Cryogenic Novel Material, Korea Polar Research Institute, Incheon 21990, Republic of Korea

Glinus oppositifolius L., Molluginaceae family, has a long-standing history of utilization as both a vegetable and a medicinal agent across numerous countries (Burkill, 1985). As a part of our ongoing study to find natural anti-adipogenic agents from Vietnamese plants, an alcoholic extract from the aerial parts of *G. oppositifolius* was found to exhibit an inhibitory effect on adipogenic cells. This presentation will cover the isolation of six new triterpenoid saponins (**1**, **2**, **6-9**) along with three known compounds (**3-5**) (Fig. 1). The structure elucidation was performed by a combination of extensive NMR analysis, HRFABMS, and DFT calculations.




Fig. 1. Structures of isolated compounds (**1-9**) from *G. oppositifolius*.

Through the adipogenesis assay, compounds **5** and **7** significantly suppressed preadipocyte differentiation of 3T3-L1 cells with IC₅₀ values at 35.4, 32.7, and 40.9 μM, respectively, compared to that of epigallocatechin-3-gallate (EGCG) previously reported as a positive control (Lee et al., 2013).

Keywords: *Glinus oppositifolius*, hopane-type saponin, anti-adipogenic activity, 3T3-L1, obesity

References

- Lee, M., Song, J. Y., Chin, Y.-W., & Sung, S. H. (2013). Anti-adipogenic diarylheptanoids from *Alnus hirsuta* f. *sibirica* on 3T3-L1 cells. *Bioorganic & Medicinal Chemistry Letters*, 23(7), 2069–2073.
- Burkill, H. M. (1985). *The Useful Plants of West Tropical Africa* (2nd ed.). Royal Botanic Gardens, Kew.

* Presenting author. E-mail address: phikimhoa246@gmail.com / pkh246@kopri.re.kr (Kim-Hoa Phi)

S1.P10 Chlorine-containing polyacetoxylbriaranes from the octocoral *Junceella fragilis*

Ping-Jyun Sung^{a,b,*}, Hai Nhat Do^{a,b}, Jui-Hsin Su^{a,b}

^a National Museum of Marine Biology and Aquarium, Pingtung 944401, Taiwan

^b Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804201, Taiwan

Chemical screening of the octocoral *Junceella fragilis* has led to the isolation of five chlorinated briarane-type diterpenoids, including three known metabolites, gemmacolide X (1), frajunolide I (2), and fragilide F (3), along with two new analogs, 12 α -acetoxylfragilide F (4) and 12 α -acetoxyljunceellin (5) (Fig. 1).




Fig. 1. Structures of gemmacolide X (1), frajunolide I (2), fragilide F (3), 12 α -acetoxylfragilide F (4), 12 α -acetoxyl- and junceellin (5); ORTEP revealing the structures of 1 and 2.

Single-crystal X-ray diffraction analysis established the complete absolute stereochemistry of natural products 1 and 2 as depicted, while the structures of new compounds 4 and 5 were ascertained with 2D NMR experiments. An evaluation of the conformation of frajunolide I (2) was carried out. Briaranes 4 and 5 exhibit cytotoxicity toward MG63 human osteosarcoma cells at concentration of 10 μ M.

Keywords: *Junceella fragilis*, Ellisellidae, octocoral, cytotoxicity, osteosarcoma

References

- Liaw, C.-C., Shen, Y.-C., Lin, Y.-S., Hwang, T.-L., Kuo, Y.-H., Khalil, A. T., 2008. Frajunolides E–K, briarane diterpenes from *Junceella fragilis*. J. Nat. Prod. 71, 1551–1556.
- McFadden, C.S., van Ofwegen, L.P., Quattrini, A.M., 2022. Revisionary systematics of octocorallia (cnidaria: anthozoa) guided by phylogenomics. Bull. Soc. Syst. Biol. 1, 8735.
- Qi, X., Zhang, X., Meng, J., Wu, J., Cheng, W., Huang, J., Lin, W., 2023. Briarane-type diterpenoids, the inhibitors of osteoclast formation by interrupting Keap1-Nrf2 interaction and activating Nrf2 pathway. Eur. J. Med. Chem. 246, 114948.

* Presenting author. E-mail address: pjsung@nmmba.gov.tw (Ping-Jyun Sung)

S1.P11 Apoptotic agents from endolichenic fungi against chemoresistant cancers

Hind Makhoulfi^{a,*}, Yves Champavier^b, Aline Pinon^a, Jennifer Saliba^c, Ingrid Arnaudin^c, Guillaume Chemin^a, Lengo Mambu^a

^a University of Limoges, LABCIS laboratory, UR22722, F-87000 Limoges, France

^b University of Limoges, BISCEm, UAR2015 US42, F-87000 Limoges, France

^c University of La Rochelle, LIENSs laboratory, UMRi 7266, F17000 LA Rochelle, France

In 2020, cancer emerged as a prominent global cause of death, with breast, lung and colon cancers being the most prevalent. Overcoming chemoresistance represents a major challenge in cancer treatment. A potential breakthrough lies in endolichenic fungi, non-pathogenic fungi found in lichens, which offer new bioactive compounds with anticancer potential. Endolichenic fungi (ELF) were previously isolated from lichens collected in the Nouvelle Aquitaine region (France) and a screening of 20 endolichenic extracts against the human colorectal cancer cell line (HT-29) highlighted their antiproliferative potential, with IC₅₀ values ranging from 2 to 60 µg/mL (MTT assay). EtOAc extracts from strains PA08 and XC04 were the most active, with IC₅₀ values below 6 µg/mL against HT-29. However, only the extract PA08S was active on triple-negative human breast cancer (MDA-MB-231), with an IC₅₀ of 4.1 µg/mL. Solid-state fermentation of PA08 and XC04 on appropriate medium yielded extracts of 5 g EtOAc each. Evaluation of the antiproliferative activity of their sub-extracts highlighted the potency of PA08S-H and PA08S-M on HT-29 with IC₅₀ values of 16 and 3 µg/mL respectively. For XC04P, the two precipitates and the intermediate fractions from the liquid-liquid partition displayed IC₅₀ values on HT-29 of 4.9, 4.3 and 1.1 µg/mL respectively. The main compound isolated from XC04P-I was active with IC₅₀ of 1.5 and 3.7 µM on HT-29 and MDA-MB-231 cells respectively. Purification of bioactive fractions is underway to isolate further bioactive compounds, and their bioactivity will be investigated through apoptosis studies on MDA-MB231 and HT-29 cells.

Keywords: lichens, endolichenic fungi, antiproliferative activity, colorectal cancer, triple negative breast cancer

References

- Ferlay J., Colombet M., Soerjomataram I., Parkin D.M., Piñeros M., Znaor A., Bray F., 2021, Cancer statistics for the year 2020: An overview. *International Journal of Cancer*. 149, 778–789.
- Aurelie L., Marion M., Aline P., Bertrand L., Marion G., C. Imbert., T.S. Ouk., P. Jargeat., L. Mambu., 2019 Antiproliferative and antibiofilm potentials of endolichenic fungi associated with the lichen *Nephroma laevigatum*. *Journal of Applied Microbiology*, 126, 1044–1058.
- Gao H., Zou J., Li J., Zhao H., 2016, Endolichenic Fungi: A Potential Treasure Trove for Discovery of Special Structures and Bioactive Compound. *Elsevier*, 48, 347–397.

* Corresponding author. E-mail address: hind.makhoulfi@etu.unilim.fr (Hind Makhoulfi)

S1.P12 *Aetokthonos hydrillicola* – an epiphytic cyanobacterium with surprising toxic potential

Markus Schwark^a, José A. Martínez Yerena^b, Steffen Breinlinger^a, Tabitha J. Phillips^c, Susan B. Wilde^b, Jan Mareš^c, Timo H. J. Niedermeyer^{a,d,*}

^a Institute of Pharmacy, Martin-Luther-University Halle-Wittenberg, Hoher Weg 8, 06120 Halle (Saale), Germany

^b Biology Centre of the Czech Academy of Sciences, Institute of Hydrobiology, Na Sádkách 702/7, České Budějovice 37005, Czech Republic

^c Warnell School of Forestry and Natural Resources, University of Georgia, 180 E Green Street, Athens, GA 30602-2152, USA

^d Institute of Pharmacy, Freie Universität Berlin, Königin-Luise-Straße 2+4, 14195 Berlin, Germany

Cyanobacteria are infamous producers of toxins. While the toxic potential of planktonic cyanobacterial blooms is well documented, the effects of toxigenic benthic and epiphytic cyanobacteria are an understudied threat. In a previous study, we showed that the freshwater epiphytic cyanobacterium *Aetokthonos hydrillicola* produces a neurotoxin, aetokthonotoxin (AETX, **1**), causing the fatal disease Vacuolar Myelinopathy that affects a wide array of wildlife in the southeastern United States (Breinlinger et al., 2021).

In an assay for cytotoxicity, however, we found the crude extract of the cyanobacterium to be much more potent than pure AETX, prompting further investigation. This led to the isolation and structure elucidation of the aetokthonostatins (AESTs, **2**), linear peptides belonging to the dolastatin compound family, featuring a unique modification of the C-terminal phenylalanine-derived moiety. We confirmed that AEST potently impacts microtubule dynamics and can bind to tubulin like dolastatin 10, and we showed that AEST inhibits reproduction of the nematode *C. elegans*. Bioinformatic analysis and biochemical studies revealed the AEST biosynthetic gene cluster encoding a nonribosomal peptide synthetase / polyketide synthase accompanied by a unique tailoring machinery (Schwark et al., 2023).




Fig. 1. Structures of aetokthonotoxin **1** and aetokthonostatin **2**

Keywords: *Aetokthonos hydrillicola*, cyanotoxin, dolastatin, biosynthesis

References

Breinlinger, S., Phillips, T.J., [...], Niedermeyer, T.H.J., Wilde, S.B., 2021. Hunting the eagle killer: A cyanobacterial neurotoxin causes vacuolar myelinopathy. *Science* 6536, eaax9050.

Schwark, M., Martínez Yerena, J.A., [...], Mareš, J., Niedermeyer, T.H.J., 2023. More than just an Eagle Killer: The freshwater cyanobacterium *Aetokthonos hydrillicola* produces highly toxic dolastatin derivatives. *PNAS* 120, e2219230120.

* Presenting author. E-mail address: timo.niedermeyer@fu-berlin.de (Timo H. J. Niedermeyer)

S1.P13 Cellulamides, a new family of marine-sourced peptides from a seaweed-associated *Cellulosimicrobium*

Mariana Girão^{a,b,*}, José Murillo-Alba^c, Jesús Martín^c, Ignacio Pérez-Victoria^c, Ralph Urbatzka^a, Pedro N. Leão^a, Maria F. Carvalho^{a,b}, Fernando Reyes^c

^a CIIMAR - Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Porto, Portugal

^b ICBAS - School of Medicine and Biomedical Sciences, University of Porto, Portugal

^c Fundación MEDINA, Granada, Spain

Mining the secondary metabolism of underexplored Actinomycetota taxa from unique marine sources represents a valuable route to uncover novel chemistry. These bacteria often live in symbiosis with other organisms, as seaweeds, prompting a prolific – yet poorly explored – hotspot for the synthesis of new molecules (Girão *et al.*, 2019). In this work we report the isolation, structure elucidation and bioactivity screening of cellulamides A-C (**1-3**), a family of novel linear peptides isolated from cultures of the seaweed-associated strain *Cellulosimicrobium funkei* CT-R177. The host macroalgae *Codium tomentosum* was collected in the northern Portuguese coast and, as part of a bioprospection program to explore its associated actinobacterial community, CT-R177 was isolated, taxonomically identified and screened for the production of antimicrobial and anticancer compounds. Dereplication of CT-R177 crude extract using LC/HRMS unveiled a putative novel natural product, cellulamide A (**1**), that was isolated following mass spectrometry-guided fractionation. Throughout the chromatographic process two other analogs, cellulamides B and C (**2, 3**), were also obtained and chemically characterized. Using a combination of 1D/2D NMR spectroscopy and HRMS, the structure of the novel peptides was elucidated. Cellulamide A was subjected to a set of bioactivity screenings, with no positive results. More assays are being conducted to uncover its potential biotechnological application. Cellulamides represent the first family of natural products reported from the Actinomycetota genus *Cellulosimicrobium* – a bacterial group known for their industrial-value in cellulose degradation with unreported secondary metabolite production – showcasing not only the potential of less-explored taxa but also host-associated marine strains for novel chemistry discovery.

Keywords: Actinomycetota, *Cellulosimicrobium*, linear peptides, natural products, seaweed- associated;

References

Girão, M., 2019. Actinobacteria isolated from *Laminaria ochroleuca*: a source of new bioactive compounds. *Frontiers in Microbiology*, 10, 683.

* Presenting author. E-mail address: mariana.martins@ciimar.up.pt (Mariana Girão)

S1.P14 A high-performance thin layer chromatography (HPTLC) method to identify lichen compounds and highlight their antibacterial activity

Alice Brissoit*, Flavia Bollaro, Isabelle Rouaud, Françoise Lohezic-Le Dévéhat, Marylène Chollet-Krugler, Sophie Tomasi, Joël Boustie

Rennes University, ISCR, CNRS, UMR 6226, CorInt team, Rennes, France

Lichens are symbiotic organisms that produce unique specialized metabolites. Phenols and terpenoids are the major ones and are often biomarkers used for lichen recognition (Lohezic-Le Dévéhat, 2012). Moreover, some of them are described for their bioactivities (Kello et al., 2023). This explains the challenge of the present work, which consists on the development and optimization of a new efficient analytical method to identify lichen compounds and verify the presence of biological activities. One analytical technique that meets both these objectives is HPTLC. Indeed, it allows the user to quickly visualize the major compounds in a lichen extract. Moreover, it is an interesting way to analyze UV absorbing compounds (phenols) and also those that are not visible under ultraviolet wavelengths (terpenoids), thanks to a derivatization step that reveals the compounds as colorful spots (Gerlach et al., 2018). Two different HPTLC methods (terpenoids/phenols) were developed in different modes (gradient/isocratic). Hundreds of standards were analyzed, enabling the creation of databases in order to gather all information (R_f , color of the spots before and after derivatization step, UV spectra data) and facilitate the compounds identification in a lichen extract. Once the compounds were well separated and recognized, bioautographies were performed on new plates. Culture medium and bacteria (*Bacillus subtilis*) were poured onto the plates after the migration step. After incubation and derivatization (methyl thiazolyl tetrazolium bromide), the zone of inhibition was measured to highlight the antibacterial activity of the identified lichen compounds. Results will be discussed in the presentation.

Keywords: lichens, HPTLC, identification, bioautography, antibacterial

References

- Gerlach A.D.C.L. et al, 2018, The Use of Anisaldehyde Sulfuric Acid as an Alternative Spray Reagent in TLC Analysis Reveals Three Classes of Compounds in the Genus *Usnea* Adans. (Parmeliaceae, lichenized Ascomycota), *Preprints*, DOI: 2018020151.<https://doi.org/10.20944/preprints201802.0151.v1>
- Kello M. et al, 2023, Screening Evaluation of Antiproliferative, Antimicrobial and Antioxidant Activity of Lichen Extracts and Secondary Metabolites In Vitro, *Plants*, Volume 12 (3), DOI: <https://doi.org/10.3390/plants12030611>
- Lohezic-Le Dévéhat F., 2012, La chromatographie sur Couche Mince appliquée aux lichens, *Bulletin de l'Association Française de Lichénologie*, Volume 37(2).

* Presenting author. E-mail address: alice.brissoit.these@gmail.com (Alice Brissoit)

S1.P15 Potent antiproliferative dimeric biaryl-cyclohexapeptides from *Lentzea flaviverrucosa*

Shugeng Cao^{a,b,*}, Chunshun Li^{a,b}, Yifei Hu^c, Xiaohua Wu^a, Spencer D. Stumpf^c, Yunci Qi^c, John M. D'Alessandro^c, Keshav K. Nepal^c, Ariel M. Sarotti^d, Joshua A.V. Blodgett^c

^a Department of Pharmaceutical Sciences, Daniel K. Inouye College of Pharmacy, University of Hawaii at Hilo, Hilo, HI 96720, USA

^b Cancer Biology Program, University of Hawaii Cancer Center, Honolulu, HI 96813, USA

^c Department of Biology, Washington University in St Louis, St Louis MO 63122, USA

^d Instituto de Química Rosario (CONICET), Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario 2000, Argentina

Rare actinomycetes represent an underexploited source of new bioactive compounds. Using a targeted metabologenomic approach we identified piperazyl compounds in the rare actinomycete *Lentzea flaviverrucosa* DSM 44664. These efforts to identify molecules that incorporate piperazate building blocks resulted in the discovery and structural elucidation of two dimeric biaryl-cyclohexapeptides, petrichorins A (1) and B. Petrichorin B (2) is a symmetric homodimer similar to the known compound chloptosin, but petrichorin A (1, Fig. 1) is unique among known piperazyl cyclopeptides because it is an asymmetric heterodimer. Due to the structural complexity of petrichorin A, solving its structure required a combination of several standard chemical methods plus in silico modeling, strain mutagenesis, and solving the structure of its biosynthetic intermediate petrichorin C (3) for confident assignment. Furthermore, we found that the piperazyl cyclopeptides comprising each half of the petrichorin A heterodimer are made via two distinct nonribosomal peptide synthetase (NRPS) assembly lines, and the responsible NRPS enzymes are encoded within a contiguous biosynthetic supercluster on the *L. flaviverrucosa* chromosome. Requiring promiscuous cytochrome p450 crosslinking events for asymmetric and symmetric biaryl production, petrichorins A and B exhibited potent in vitro activity against A2780 human ovarian cancer, HT1080 fibrosarcoma, PC3 human prostate cancer, and Jurkat human T lymphocyte cell lines with IC50 values at low nM levels.




Fig. 1. Structure of petrichorin A 1.

Keywords: *Lentzea flaviverrucosa*, dimeric biaryl-cyclohexapeptides, nonribosomal peptide synthetase, p450, NMR, antiproliferative activity

References

- Li, C., Hu, Y., Wu, X., Stumpf, S.D., Qi, Y., D'Alessandro, J.M., Nepal, K.K., Sarotti, A.M., Cao, S., Blodgett, J.A.V., 2022. Discovery of unusual dimeric piperazyl cyclopeptides encoded by a *Lentzea flaviverrucosa* DSM 44664 biosynthetic supercluster. *Proc Natl Acad Sci U S A*. 119(17), e2117941119.

* Presenting author. E-mail address: scao@hawaii.edu (Shugeng Cao)

S1.P16 A validated RP-HPLC-DAD method for analyzing flavonoids in Caatinga Brazilian green propolis from *Mimosa tenuiflora* (Willd.) Poir. produced by *Apis mellifera* L.

Ninh The Son^{a,b}, Victor Pena Ribeiro^a, Jairo Kenupp Bastos^{a,*}

^a School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Avenida Professor Doutor Zeferino Vaz, S/N, 14040-903, Ribeirão Preto-SP, Brazil

^b Institute of Chemistry, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

There are several types of propolis in Brazil produced by *Apis mellifera* (Santos et al., 2021), and the Caatinga biome green propolis stands out for its high content of flavonoids (Son et al., 2022). Hence, the presentation will cover the flavonoids isolated from Brazilian green propolis of Caatinga *Mimosa tenuiflora*, and the development of a reliable RP-HPLC quantitative method using a Shim-pack VP-ODS column (250 × 4.6 mm i.d., 5 µm) with nonlinear gradient elution and UV visualization at 280 nm. Additionally, a sample preparation method for extracting flavonoids using 96% ethanol and caffeic acid as the internal standard was developed.

The developed method allowed an excellent detection response, in which the LOD (limit of detection) and LOQ (limit of quantification) reached ranges of 0.65–2.08 µg/mL and 1.97–6.31 µg/mL, respectively. The maximum RSD (relative standard deviation) was 4.61%. Thirteen flavonoids, comprising santin (1), ermanin (2), sakuranetin (3), quercetin 3-methyl ether (4), viscosine (5), eriodictyol 5-O-methyl ether (6), isokaempferide (7), kaempferide (8), penduletin (9), quercetagenin 3,6,7-trimethyl ether (10), cirsimaritin (11), 3,3'-O-dimethylquercetin (12), and luteolin (13), were quantified (Fig. 1).




Fig. 1. Chemical structures of compounds 1–13.

The developed method met all the parameters set by the international guidelines for developing analytical methods. The current analytical method is reliable for the quality control of *M. tenuiflora* green propolis and its related products.

Keywords: *Apis mellifera*, *Mimosa tenuiflora*, Caatinga, Fabaceae, flavonoids, RP-HPLC-DAD

References

- Santos, M.F.C., Oliveira, L.C., Ribeiro, V.P., Soares, M.G., Moraes, G.O.I., Sartori, A.G.D.O., Rosalen, P.L., Bastos, J.K., Alencar, S.M.D., Veneziani, R.C.S., Ambrósio, S.R., 2021. Isolation of diterpenes from *Araucaria* sp Brazilian brown propolis and development of a validated high-performance liquid chromatography method for its analysis. *J Sep Sci.* 44, 3089–97.
- Son, N.T., Ribeiro, V.P., Bastos, J.K., 2022. Flavonoids from green propolis of the Northeastern Brazilian Caatinga *Mimosa tenuiflora* (Willd.) Poir.: A chemotaxonomic aspect. *Biochem. System. Ecol.* 104, 104473.

* Presenting author. E-mail address: jkbastos@fcfrp.usp.br (Jairo Kenupp Bastos)

S1.P17 Functional oligosaccharides derived from the degradation of natural *Dendrobium* polysaccharides by human gut microbes regulate physiological factors associated with type-2 diabetes

Ming-Che Cheng^a, Hsi-Mei Chen^a, Bao-Jiun Bai^a, Tsu-Yuan Chang^a, Meng-Chin Chen^a, Yen-Ting Lin^b, Chi-Kung Ni^b, Mei-Kuang Lu^c, Hui-Kang Liu^c, Chia-Chuan Chang^{a,*}

^a School of Pharmacy, National Taiwan University, Taipei, Taiwan, R.O.C

^b Institute of Atomic and Molecular Sciences, Academia Sinica, Taipei, Taiwan, R.O.C

^c National Research Institute of Chinese Medicine, Ministry of Health and Welfare, Taipei, Taiwan, R.O.C

Dendrobium, a noble traditional Chinese herb medicine, has been found to be rich in glucomannan and possesses significant bioactivities in regulating blood sugar and blood pressure (Pan et al., 2014). Herein, we selected a unique hybrid species, *Dendrobium Taiseed Tosnobile* (DTT), which is derived from the crossbreeding of *D. tosaense* and *D. nobile*, as our research material. The acetylated glucomannan, isolated from DTT, consists of mannose to glucose at a ratio of 5.5:1 and has an average molecular weight of 194 kDa. After administering isolated DTT polysaccharide (DTTPS) to diabetic model mice for 7 weeks, notable improvements in body weight, appetite, and postprandial blood glucose levels of the mice were observed. Furthermore, the abundance of probiotics such as *Akkermansia*, *Lactobacillus*, and *Bifidobacterium* significantly increased in mice fed with DTTPS compared to those in the control group. To further investigate the mechanism of action of DTTPS, *Bacteroides ovatus* was chosen to ferment DTTPS under carbon source-depleted conditions. In this study, small molecular weight acetylated glucomannan, ranging from 1-2 kDa, were successfully isolated. These oligosaccharides demonstrated potent activities toward promoting GLP-1 secretion and inhibiting DPP-4 degradation, along with moderate inhibitory activity against α -glucosidase, revealing a comprehensive function in regulating physiological factors associated with type-2 diabetes.

Keywords: *Dendrobium* spp., glucomannan, gut microbes, GLP-1, DPP-4

References

Pan, L.-H., Li, X.-F., Wang, M.-N., Zha, X.-Q., Yang, X.-F., Liu, Z.-J., Luo, Y.-B., Luo, J.-P., 2014. Comparison of hypoglycaemic and antioxidative effects of polysaccharides from four different *Dendrobium* species. *Int. J. Biol. Macromol.* 64, 420-427.

* Presenting author. E-mail address: D10423101@ntu.edu.tw (Ming-Che Cheng)

S1.P18 Discovering original antiplasmodial compounds through mass-guided exploration of the alkaloid diversity in the *Strychnos* genus

Olivier Bonnet^{a,*}, Mehdi A. Beniddir^b, Pierre Champy^b, Charline Leroy^a, Carla Hamann^a, Allison Ledoux^a, Alembert Tiabou Tchinda^{a,c}, Luc Angenot^a, Michel Frédérich^a

^a CIRM, Department of Pharmacognosy, University of Liège, Avenue Hippocrate 15, 4000 Liège, Belgium

^b Équipe « Chimie des Substances Naturelles », BioCIS, CNRS, Université Paris-Saclay, 17 avenue des Sciences, 91400 Orsay, France

^c Laboratory of Phytochemistry, Center for Research on Medicinal Plants and Traditional Medicine, Institute of Medicinal Research and Medicinal Plants Studies, Yaoundé P.O. Box 13033, Cameroon

Several monoterpene indole alkaloids from various plants of the *Strychnos* genus have shown promising antiplasmodial activity. In front of the parasites' growing resistance to existing treatments, the alkaloid content from 44 crude extracts of 28 *Strychnos* species are explored using the molecular networking technique with the aim of identifying new antimalarial compounds. Thanks to public MS/MS spectral databases and the antiplasmodial activities of studied crude extracts, clusters containing mainly unknown metabolites potentially active against malaria were highlighted and targeted (Fig. 1) (Bonnet *et al.*, 2022a).




Fig. 1. One cluster from GNPS-annotated molecular network of 44 crude extracts. (Colors of antiplasmodial activities: Dark green= < 5 µg/mL; Light green= 5-15 µg/mL; Yellow= 15-30 µg/mL; Orange= 30-50 µg/mL; Red: > 50 µg/mL).

Exploring the annotated molecular network led to reveal unexpected presence of strychnine in seven species (Bonnet *et al.*, 2022b), and to select *Strychnos usambarensis* leaves and *Strychnos longicaudata* trunk barks for in-depth studies. Using bio- and mass-guided fractionation, several unknown metabolites were isolated, and are currently being identified using NMR and MS/MS data.

Keywords: *Strychnos*, alkaloids, molecular networking, mass spectrometry, malaria

References

- Bonnet, O., Beniddir, M. A., Champy, P., Degotte, G., Mamede, L., Desdemoustier, P., Ledoux, A., Tchinda, A. T., Angenot, L., Frédérich, M., 2022a. Unveiling antiplasmodial alkaloids from a cumulative collection of *Strychnos* extracts by multi-informative molecular networks. *Frontiers in Molecular Biosciences*, 9:967012.
- Bonnet, O., Beniddir, M. A., Champy, P., Kagisha, V., Nyirimigabo, A., Hamann, C., Jgerenaia, G., Ledoux, A., Tchinda, A. T., Angenot, L., Frédérich, M., 2022b. Exploration by molecular networking of *Strychnos* alkaloids reveals the unexpected occurrence of strychnine in seven *Strychnos* species. *Toxicon*, 215, 57–68.

* Presenting author. E-mail address: Olivier.Bonnet@uliege.be (Olivier Bonnet)

S1.P19 Characterization and antimicrobial activity of extracts from microalgae *Chlorococcum* sp. with green organic solvents

Elia Lio^{a,b,*}, Martina Dramis^a, Gianluca Ottolina^a, Giancarlo Aldini^b, Francesco Secundo^a

^a CNR, Istituto di Scienze e Tecnologie Chimiche, Via Mario Bianco 9, 20131 Milan, Italy

^b University of Milan, Department of Pharmaceutical Sciences, Milano, Italy

The processes involved in extracting and characterizing natural compounds are crucial for enhancing the environmental sustainability of industrially relevant products, particularly when aimed at discovering novel chemical structures for drug development. Various extraction methods utilizing environmentally friendly organic solvents (Byrne et al., 2016) are employed to obtain bioactive compounds from natural sources. Microalgae, rich in bioactive compounds such as carotenoids, polyphenols, phytosterols, and tocopherols (Sarkar et al., 2020), are gaining attention from numerous agro-food and pharmaceutical industries as a source of natural products. This study focuses on extracting the lipophilic fraction from the microalgae *Chlorococcum* sp. and identifying compounds with antimicrobial properties against *Escherichia coli*, *Bacillus megaterium*, and *Bacillus subtilis* through principal component analysis (PCA) of GC-MS spectra of the whole extracts. The extraction process relies on green organic solvents, aiming to demonstrate their efficacy as replacements for conventional organic solvents like chloroform (Patras et al., 2011). The proposed approach seeks to offer alternative methods to the laborious extraction and material-consuming steps typically involved in isolating and characterizing individual compounds. Antimicrobial tests were conducted using the agar well diffusion method (Valgas et al., 2007) and the broth dilution method (Rubin, 2013), revealing significant antimicrobial activity in the extracts. Overall, the study underscores the potential of microalgae as a valuable source of bioactive compounds and investigates the processes involved in their extraction, characterization, and assessment of antimicrobial activity.

Financial support from the European Union – Next Generation EU in the frame of the “National Biodiversity Future Center”(CUPn°B83C22002930006) is gratefully acknowledged.

Keywords: microalgae, green organic solvent, extractions, antimicrobial activity, extract characterization, PCA

References

- Byrne, F.P., Jin, S., Paggiola, G., Petchey, T.H.M., Clark, J.H., Farmer, T.J., Hunt, A.J., Robert McElroy, C., Sherwood, J., 2016. Tools and techniques for solvent selection: green solvent selection guides. *Sustainable Chemical Processes*, 4(1). Patras, A., Brunton, N.P., Downey, G., Rawson, A., Warriner, K., Gernigon, G., 2011. Application of principal component and hierarchical cluster analysis to classify fruits and vegetables commonly consumed in Ireland based on in vitro antioxidant activity. *Journal of Food Composition and Analysis*, 24, 250–256.
- Rubin, J.E. (2013). Antimicrobial Susceptibility Testing Methods and Interpretation of Results. In *Antimicrobial Therapy in Veterinary Medicine* (pp. 11–20). Wiley.
- Sarkar, S., Manna, M.S., Bhowmick, T.K., Gayen, K., 2020. Priority-based multiple products from microalgae: review on techniques and strategies. *Critical Reviews in Biotechnology*, 40, 590–607.
- Valgas, C., Souza, S.M. de, Smânia, E.F.A., Smânia Jr., A., 2007. Screening methods to determine antibacterial activity of natural products. *Brazilian Journal of Microbiology*, 38, 369–380.

* Presenting author. E-mail address: elia.lio@unimi.it (Elia Lio)

S1.P20 Exploration of new biological activity of *Mikania micrantha*, a widespread weed in Asia

Grace Gar-Lee Yue^{a,*}, Tao Zheng^a, Wai-Fung Lee^a, Julia Kin-Ming Lee^a, Clara Bik-San Lau^{a,b}

^a Institute of Chinese Medicine and State Key Laboratory of Research on Bioactivities and Clinical Applications of Medicinal Plants, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^b Department of Pharmacology and Pharmacy & School of Chinese Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong SAR, China

Mikania micrantha Kunth (family Asteraceae), a fast-growing perennial herbaceous vine, has been regarded as a widespread weed in Asia. However, originates from the tropical South and Central America, *M. micrantha* (MM) has been used as medicinal plant. Pharmacological activities such as anti-inflammatory, antioxidant, antiviral and cytotoxicity towards various cancer cells have been previously reported (But et al., 2009; Dong et al., 2017; Rios et al. 2014). This study aimed to identify the active fraction of MM which may account for the cytotoxic activity in human cancer cells, using bioassay-guided fractionation. The aerial parts of MM were extracted with 80 % ethanol under reflux, followed by sequential partition with petroleum ether, ethyl acetate and *n*-butanol. The active fraction was then subjected to column chromatography to yield subfractions for further evaluation on the cell viability of different human cancer cells and normal fibroblasts Hs27 using MTT assay. Results showed that the ethyl acetate fraction significantly reduced cell viability of MDA-MB-231 (breast), HT-29 (colon) and EC-109 (esophageal) cancer cells with IC₅₀ of 12.9, 21.5 and 46.9 mg/mL, respectively, but not Hs27 (IC₅₀ > 200 mg/mL) cells, suggesting its selective cytotoxicity. Moreover, in MDA-MB-231 and HT-29 cells, the ethyl acetate fraction induced G2 phase arrest of cell cycle in a concentration dependent manner. The subfraction E2 maintained selective cytotoxic activity in MDA-MB-231 cells (IC₅₀ at 46.8 mg/ml). Our findings demonstrated the potential of this subfraction in suppressing breast tumor growth, which warrant further *in vivo* investigations to verify the new bioactivity of this weed.

Keywords: *Mikania micrantha*, Asteraceae, cytotoxicity, breast cancer, colon cancer

References

- But, P.P., He, Z.D., Ma, S.C., Chan, Y.M., Shaw, P.C., Ye, W.C., Jiang, R.W., 2009. Antiviral constituents against respiratory viruses from *Mikania micrantha*. J Nat Prod. 72, 925-928.
- Dong, L.M., Jia, X.C., Luo, Q.W., Zhang, Q., Luo, B., Liu, W.B., Zhang, X., Xu, Q.L., Tan, J.W., 2017. Phenolics from *Mikania micrantha* and their antioxidant activity. Molecules. 22, 1140.
- Ríos, E.V., León, A., Chávez, M.I., Torres, Y., Ramírez-Apan, M.T., Toscano, R.A., Bravo-Monzón, Á.E., Espinosa- García, F.J., Delgado, G., 2014. Sesquiterpene lactones from *Mikania micrantha* and *Mikania cordifolia* and their cytotoxic and anti-inflammatory evaluation. Fitoterapia. 94, 155-163.

* Presenting author. E-mail address: graceyue@link.cuhk.edu.hk (Grace Gar-Lee Yue)

S1.P21 Preclinical study on the anti-proliferative effects and underlying mechanisms of brousoflavonol F in colon cancer

Yiying Zhu^{a,*}, Grace Gar-Lee Yue^a, Chun Kwok Wong^a, Wei-Lie Xiao^b, Clara Bik-San Lau^{a,c}

^a Institute of Chinese Medicine and State Key Laboratory of Research on Bioactivities and Clinical Applications of Medicinal Plants, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^b Key Laboratory of Medicinal Chemistry for Natural Resource, Ministry of Education, Yunnan University, Yunnan, China

^c Department of Pharmacology and Pharmacy & School of Chinese Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong SAR, China

Brousoflavonol F (BFF), a natural prenylated flavonol isolated from *Macaranga indica* Wight, has been reported to possess cytotoxicities against various cancer cells, including MCF7, A549, HepG2 and Hela (Segun et al., 2019; Vu et al., 2021). However, there is no report on the anti-proliferative activity of BFF in colon cancer. This study investigated on the anti-tumor effect of BFF and its underlying mechanisms in colon cancer. The anti-proliferative effects of BFF on human colon cancer HCT116 and LoVo cells were evaluated using MTT, BrdU, and colony formation assays, cell cycle and cell apoptosis analysis. Besides, network pharmacology analysis was used to predict the effects of BFF on the expression of key proteins in intracellular signaling pathways, with validation using western blot. The anti-tumor effect of BFF was also examined in HCT116 tumor-bearing nude mice. Our results showed that BFF significantly exhibited anti-proliferative activities in both HCT116 (2.5-7.5 μ M) and LoVo (5-10 μ M) cells via inducing apoptosis and cell cycle arrest at the G0/G1 phase. Further investigation revealed that BFF (1.25-5 μ M) inhibited cell proliferation by downregulating the expression of HER2, RAS, p-BRAF, p-MEK and p-Erk. In addition, intraperitoneal administration of BFF (10 mg/kg) suppressed tumor growth and expression of Ki-67 and CD31 in the tumor tissues of HCT116 tumor-bearing mice. This is the first report on the *in vitro* and *in vivo* anti-proliferative effects of BFF in colon cancer, through regulation of HER2-RAS-MEK-Erk pathway. Our findings support further research development of BFF as an anti-cancer agent for colon cancer.

Keywords: brousoflavonol F, *Macaranga indica*, colon cancer, anti-proliferative, anti-tumor

References

Segun PA, Ogbole OO, Ismail FMD, Nahar L, Evans AR, Ajaiyeoba EO., 2019. Bioassay-guided isolation and structure elucidation of cytotoxic stilbenes and flavonols from the leaves of *Macaranga barteri*. Fitoterapia. 134, 151–157.

Vu LTN, Anh LT, Cuc NT, Nhiem NX, Tai BH, Van Kiem P., 2021. Prenylated flavonoids and other constituents from *Macaranga indica*. Nat Prod Res. 35, 2123–2130.

* Presenting author. E-mail address: 1155189460@link.cuhk.edu.hk (Yiying Zhu)

S1.P22 Isolates of *Pavonia multiflora* as alternative control of the phytopathogen *Fusarium solani piperis*

Jessyca Aparecida Paes Dutra^a, Jéssica Raquel Borges Monteiro^{b,d}, Weverson Cavalcante Cardoso^c, Valdemar Lacerda Júnior^d, Rita de Cássia Ribeiro Gonçalves^b, Warley de Souza Borges^d, Rodrigo Rezende Kitagawa^{b,d,*}

^a School of Pharmaceutical Science of Sao Paulo State University (UNESP), Araraquara, Sao Paulo, Brazil

^b Federal University of Espirito Santo (UFES), Graduate Program of Pharmaceutical Sciences, Health Sciences Center, Vitoria, Espirito Santo, Brazil

^c Federal University of Espirito Santo (UFES), Department of Biological Sciences, Human and Natural Sciences Center, Vitoria, Espirito Santo, Brazil

^d Federal University of Espirito Santo (UFES), Graduate Program of Chemistry, Exact Sciences Center, Vitoria, Espirito Santo, Brazil

Fusarium species, responsible for a significant proportion of crop fungal diseases, act a substantial threat to global agriculture, resulting in considerable economic losses. Environmental and health concerns associated with synthetic fungicides have prompted the exploration of natural products as alternative control measures. Antifungal compounds derived from the endemic Brazilian species *Pavonia multiflora* (Malvaceae) emerge as ecologically friendly alternatives for *Fusarium* control. In this study, *in vitro* assays were conducted to screen eight *P. multiflora* isolates (Lopes et al., 2016) and identify compound(s) with fungicidal efficacy and safe cytotoxic profile. Four isolates: vanillic acid (MFC = 200 µg.mL⁻¹), loliolide (MFC=400 µg.mL⁻¹), vomifoliol (MFC=200 µg.mL⁻¹), and 4,5- dihydroblumenol A (MFC = 400 µg.mL⁻¹) exhibited fungicidal capacity, remaining biocompatible up to 50 µg.mL⁻¹ with no significant metabolic changes in basal cells (Raw 267.4). However, distinct cytotoxic profiles were observed after exposure to a metabolizing system (Fraction S9) (Gonçalves et al., 2016) under HepG2 cells. Vanillic acid showed decrease of cytotoxicity (IC50 54.31 to 226.3 µg.mL⁻¹), while other isolates displayed increase of cytotoxicity. Given the use of only Phase I metabolism cofactors, a more precise evaluation, including Phase II cofactors and mutagenicity assays, is required in future works. However, this study showed these compounds hold promise as fungicides, particularly vanillic acid and it marks the first reporting of antifungal and cytotoxic activities of isolate 4,5 - dihydroblumenol A. Considering the long-term consequences of synthetic pesticides (carbendazim and tebuconazole), this work paves the way for safe and effective alternatives of *Fusarium* species control.

Keywords: phytopathogen, fungicides, natural products, cytotoxicity, vanillic acid

References

- Gonçalves, R.C.R., Kitagawa, R.R., Varanda, E.A., Raddi, M.S.G., Leite, C.A., Sponchiado, S.R.P., 2016. Effect of biotransformation by liver S9 enzymes on the mutagenicity and cytotoxicity of melanin extracted from *Aspergillus nidulans*. *Pharmaceutical Biology*, 54, 1014–1021.
- Lopes, L. G., Tavares, G. L., Thomaz, L. D., Sabino, J. R., Borges, K. B., Vieira, P. C., Veiga, T. A. M., Borges, W.S., 2016. Taraxerol 4-Methoxybenzoate, an *in vitro* Inhibitor of Photosynthesis Isolated from *Pavonia multiflora* A. (Malvaceae). *Chemistry & Biodiversity*, 13, 284–292.

* Presenting author. E-mail address: rodrigo.kitagawa@ufes.br (Rodrigo Rezende Kitagawa)

S1.P23 Co-culturing *Streptomyces albidoflavus*, an epiphyte isolated from *Lavandula angustifolia* Mill., increased the anti-*Porphyromonas gingivalis* effect in exo-metabolome extracts

Rosario Flores-Vallejo*, Jan Maarten van Dijk

Department of Medical Microbiology, University of Groningen, University Medical Center Groningen, Hanzeplein 1, 9700 RB Groningen, The Netherlands

Background: *Porphyromonas gingivalis* (Pg) is a keystone pathogen in periodontitis, which affects ~20% of the global population and represents a societal burden of USD\$298 billion/year (Listl et al., 2015). Common treatments are tissue debridement and antibiotics administration. However, between 13% to 76% of the clinical isolates display high antibacterial resistance (Ardila et al., 2020). **Aim:** Our aim is to identify novel anti-Pg agents by bioprospecting the phytomicrobiota of medicinal plants, an underexplored niche of microbial and chemical diversity (Newman and Cragg, 2020). **Methods:** Bacterial endophytes and epiphytes from *L. angustifolia* were isolated from 2021 to 2023. A screening panel of 20 isolates was characterized and confronted against Pg ATCC 33277 using our modified version of the “agar-overlay antibacterial assay” tested in different culture media. Then, ethyl-acetate extracts obtained from co-culturing the bioactive strains together with Pg were tested anaerobically in the microdilution assay to determine the Minimum Inhibitory Concentration (MIC). **Results:** An epiphytic *Streptomyces albidoflavus* isolate (Sa_LaEp14) grown in supplemented Potato Dextrose media presented the highest bioactivity against Pg (inhibition halo: 1.2 cm) (Fig. 1). Extracts from co-cultures with Pg showed a stronger anti-Pg effect (MIC: 0.25 µg/mL) than extracts from pure Sa_LaEp14 cultures (MIC: 2 µg/mL).




Fig. 1. Modified “agar overlay assay” showing Sa_LaEp14 (seeded in the middle) forming an inhibition halo against Pg.

Conclusions: This is the first report demonstrating that co-culturing Pg and the epiphyte Sa_LaEp14 increases the production of anti-Pg secondary metabolites by Sa_LaEp14. Currently, the identity of the bioactive secondary metabolites is under analysis.

Keywords: phytomicrobiota, lavender, co-culture, *P. gingivalis*, antibacterial

References

- Ardila, C. M., Bedoya-García, J. A., 2020. Antimicrobial resistance of *Aggregatibacter actinomycetemcomitans*, *Porphyromonas gingivalis* and *Tannerella forsythia* in periodontitis patients. J. Glob. Antimicrob. Resist. 22:215-218.
- Listl, S., Galloway, J., Mossey, P. A., Marcenes, W., 2015. Global economic impact of dental diseases. J. Dent. Res. 94: 1355-1361.
- Newman, D.J., and Cragg, G.M., 2020. Plant endophytes and epiphytes: Burgeoning sources of known and “unknown” cytotoxic and antibiotic agents? Planta Med. 86, 891-905.

* Presenting author. E-mail address: r.c.flores.vallejo@umcg.nl (Rosario Flores-Vallejo)

S1.P24 Tyrosinase and hyaluronidase inhibitors in *Eleutherococcus divaricatus*

Jakub Gębalski^{a,*}, Maciej Strzemiński^b, Magdalena Wójcik^b, Daniel Załuski^a

^a Department of Pharmaceutical Botany and Pharmacognosy of the Ludwik Rydygier Medical College in Bydgoszcz, Nicolaus Copernicus University in Torun, Marie Curie-Skłodowska 9, 85-094 Bydgoszcz, Poland

^b Department of Analytical Chemistry, Medical University of Lublin, Chodźki 4a, 20-093 Lublin, Poland

For centuries, plants have been a major source of medicines for the treatment of many diseases. The development of modern research methods has made it possible to isolate the active components of plants and develop much more effective and safer drugs (Katiyar et al. 2012). Currently available compounds, which inhibit tyrosinase or hyaluronidase, are characterized by a few unfavorable properties such as low persistence, poor absorption, or carcinogenicity (Shakya, 2016). The aim of the present study was to determine the usefulness of *Eleutherococcus divaricatus* in the treatment of tyrosinase- and hyaluronidase-related diseases. These enzymes represent potential therapeutic targets in the inflammatory-related diseases. Based on the results, *E. divaricatus* inhibited tyrosinase and hyaluronidase activity. The ethyl acetate fraction showed the strongest inhibition (bovine hyaluronidase - IC₅₀=27.5 µg/ml, fungal tyrosinase - IC₅₀=65.5 µg/ml, respectively). The extract was also active against human hyaluronidase from plasma (55.83 µg/ml). In both cases, the extract inhibited the enzyme stronger than aescin, used as the positive control. The fraction was rich in rosmarinic acid (26.62 mg/g) and caffeic acid (24.02 mg/g). The results support the traditional use of the roots in enzymes-related diseases. The use of ethyl acetate fraction demonstrates its inhibiting effect especially towards the hyaluronidase activity, which may additionally indicate its anti-inflammatory property. Thus, the results mean that the fraction may be used in inflammatory-related diseases.

Keywords: *Eleutherococcus divaricatus*, Araliaceae, hyaluronidase, tyrosinase, human serum

References

- Katiyar, C., Gupta, A., Kanjilal, S., Katiyar, S. 2012. Drug discovery from plant sources: An integrated approach. *Ayu*, 33, 10.
- Shakya, A.K. 2016. Medicinal plants: Future source of new drugs. *International journal of herbal medicine*, 4, 59-64.

* Presenting author. E-mail address: jakub.gebalski@cm.umk.pl (Jakub Gębalski)

S1.P25 Four sesquiterpene-acylated flavonol glycosides from the sprouts of *Astragalus membranaceus*

So-Ri Son^{a,*}, Jaeyoung Kwon^b, Hak Cheol Kwon^b, Dae Sik Jang^a

^a Department of Biomedical and Pharmaceutical Sciences, Graduate School, Kyung Hee University, Seoul, Republic of Korea

^b KIST Gangneung Institute of Natural Products, Korea Institute of Science and Technology (KIST), Gangneung, Republic of Korea

Sprouting as a cultivation technique has shown promise in enhancing the metabolite profile of medicinal plants, leading to potential new applications in functional foods (Benincasa et al., 2019). The present research investigated *Astragalus membranaceus* Bunge, a medicinal plant widely utilized for its roots in Oriental medicine, focusing on its sprouts which have not been extensively studied for their secondary metabolites (Hou et al., 2023). Initially, feature-based molecular networking workflow was employed to distinguish the metabolites in *A. membranaceus*' sprouts from those found in its roots. Next, compounds were annotated by matching the MS/MS spectrum with those in public databases, leading to the discovery of compounds not previously reported. Then, the targeted isolation and structure elucidation of these compounds from *A. membranaceus*' sprouts were carried out. As a result, four unreported sesquiterpene-acylated flavonol glycosides (**1–4**) were identified as compounds specific to *A. membranaceus* sprouts (Fig.1.). The chemical structures of compounds **1–4** were determined through a comprehensive analysis of their spectroscopic data (UV, HR-MS, 1D- and 2D-NMR, and CD), along with hydrolysis.




Fig. 1. Sesquiterpene-acylated flavonol glycosides from the sprouts of *A. membranaceus*

Keywords: *Astragalus membranaceus* Bunge, sprouts, flavonol glycosides

References

- Benincasa, P., Falcinelli, B., Lutts, S., Stagnari, F., Galieni, A., 2019. Sprouted grains: A comprehensive review. *Nutrients* 11, 421.
- Hou, M., Leng, Y., Shi, Y., Tan, Z., Min, X., 2023. *Astragalus membranaceus* as a Drug Candidate for Inflammatory Bowel Disease: The Preclinical Evidence. *The American Journal of Chinese Medicine*, 51, 1501-1526.

* Presenting author. E-mail address: allosori@khu.ac.kr (So-Ri Son)

S1.P26 Effects of Astraflavonoids A and D, flavonol glycosides from the aerial parts of *Astragalus membranaceus*, on TNF- α -induced human dermal fibroblasts

Mingoo Jun^{a,*}, So-Ri Son^a, Kang Sub Kim^b, Sullim Lee^b, Dae Sik Jang^c

^a Department of Biomedical and Pharmaceutical Sciences, Graduate School, Kyung Hee University, Seoul, Republic of Korea

^b College of Korean Medicine, Gachon University, Seongnam, Republic of Korea

Astragalus membranaceus, a commonly used medicinal herb, is primarily valued for its root, while the aerial parts are typically discarded (Guo et al., 2022). To utilize this agricultural waste, the present study focused on the anti-skin aging properties of the aerial parts of *A. membranaceus* and their major constituents. First, a LC-MS analysis identified flavonoid glycosides as the major constituents in the aerial parts of *A. membranaceus* extract. Following this, two significant acylated flavonol glycosides—astraf flavonoids A and D (**1** and **2**; Fig. 1.)— were isolated, with astraf flavonoid D (**2**) being a previously unreported compound. The chemical structures of the isolates were elucidated through HR-MS, UV, IR, 1D- and 2D-NMR analysis, acid hydrolysis, and comparison with existing literature (Hao et al., 2016). Subsequent experiments revealed that both astraf flavonoids A and D (**1** and **2**) have antioxidative properties, reducing ROS and MMP-1 level stimulated by TNF- α in NHDF. Specifically, astraf flavonoid A (**1**) also inhibited the expression of pro-inflammatory cytokines (IL-6 and IL-8). Additionally, the expression of inflammatory mediators (NF- κ B and COX-2) was suppressed by astraf flavonoid A (**1**). These findings indicate that astraf flavonoids A and D (**1** and **2**) from the aerial parts of *A. membranaceus* are active principles to possess anti-skin aging properties, particularly astraf flavonoid A (**1**), as a potential inhibitor for inflammatory skin diseases.




Fig. 1. Structures of compounds **1** and **2**

Keywords: *Astragalus membranaceus* Bunge, reactive oxygen species, MMP-1, COX-2

References

- Guo, L., Sun, Y., Ping, X., Liu, J., Wang, X., Qin, N., 2022. Chemical composition and antibacterial activity of ethyl acetate extract of *Astragalus membranaceus* aerial parts. *Journal of Food Safety*. 42, e12947.
- Hao, J., Li, J., Li, X., Liu, Y., Ruan, J., Dong, Y., Zhang, Y., 2016. Wang, T. Aromatic constituents from the stems of *Astragalus membranaceus* (Fisch.) Bge. var. *Mongholicus* (Bge.) Hsiao. *Molecules*. 21, 354.

* Presenting author. E-mail address: mingoojun0104@khu.ac.kr (Mingoo Jun)

S1.P27 Harnessing the microbial richness of *Ips perturbatus* for novel antimycotic agents

Nicolas de Aguiar Justus, Nirasha Ataputta, Mitz Baylosis, Kalindi D. Morgan*

University of Northern British Columbia, Department of Chemistry and Biochemistry, 3333 University Way, Prince George, BC, V2N 4B6, Canada

This presentation will provide an exploration of the potential of the bacterial symbionts of northern spruce engraver beetle (*Ips perturbatus*) as novel antimicrobial sources. Previously complex interplay between this beetle and its associated microorganisms has demonstrated biological novelty. For example, a new species of fungus, *Leptographium fruticetum* sp. nov., was identified from *Ips perturbatus* collected in northern British Columbia and Yukon Territory, with a possibly specific symbiotic relationship between the beetle and this fungus highlighted (Alamouti et al., 2006a). Additional studies have revealed the presence of a diverse range of ophiostomatoid fungi associated with *Ips perturbatus* (Alamouti et al., 2006b), along with a significant chemical component to the holobiont's interaction with its environment (Holsten et al., 2001). This variety of ecological interaction involving insects, plants and microorganisms has previously been shown to be associated with rich bacterial chemistry (Beemelmans et al., 2016). Nevertheless, the capacity of the bacteria within this holobiont to produce unique chemistry has not been explored. This presentation describes our work as we begin to unravel the potential chemistry of this microbiome. As part of our continuing search for new antimycotics active against both plant and human pathogens, the analysis of the bacterial community of *Ips perturbatus* collected in the Yukon Territory demonstrated a community rich in Actinobacteria. Examination of the resulting isolates with bioassays, spectroscopic and genomic data, continues to drive our interest in the bacteria of this underexplored microbiome.

Keywords: microbiome, antimicrobial, bioactive compounds, insect-associated bacteria

References

- Alamouti, S. M., Kim, J.-J., Breuil, C., 2006a. A new *Leptographium* species associated with the northern spruce engraver, *Ips perturbatus*, in western Canada. *Mycologia* 98 1, 149–60.
- Alamouti, S. M., Kim, J.-J., Humble, L., Uzunovic, A., and Breuil, C., 2006b. Ophiostomatoid fungi associated with the northern spruce engraver, *Ips perturbatus*, in western Canada. *Antonie van Leeuwenhoek* 91, 19–34.
- Beemelmans, C., Guo, H., Rischer, M., and Poulsen, M., 2016. Natural products from microbes associated with insects. *Beilstein J. Org. Chem.* 12, 314–327.
- Holsten, E., Burnside, R., and Seybold, S., 2001. Verbenone interrupts the response to aggregation pheromone in the northern spruce engraver, *Ips perturbatus* (Coleoptera: Scolytidae), in south-central and interior Alaska. *Journal of the Entomological Society of British Columbia* 98, 251–256.

* Presenting author. E-mail address: kalindi.morgan@unbc.ca (Kalindi D. Morgan)

S1.P28 Insights into the mechanism of antibacterial action for prenylated (iso)flavonoids from Fabaceae: a systematic study on inhibition and permeabilization of Methicillin-resistant *Staphylococcus aureus*

Janniek Ritsema^{a,*}, Nynke Kramer^b, Wouter de Bruijn^a, Maurice Franssen^c, Jean-Paul Vincken^a, Carla Araya-Cloutier^a

^a Laboratory of Food Chemistry, Wageningen University & Research, Bornse Weiland 9, 6709 WG Wageningen, The Netherlands

^b Toxicology Chair group, Wageningen University & Research, Stippeneng 4, 6708 WE Wageningen, The Netherlands

^c Laboratory of Organic Chemistry, Wageningen University & Research, Stippeneng 4, 6708 WE Wageningen, The Netherlands

Prenylated (iso)flavonoids are plant-derived compounds with potent antimicrobial activity against Methicillin-resistant *Staphylococcus aureus* (MRSA). They primarily act on the bacterial membrane and disrupt membrane integrity by fast permeabilization (Araya-Cloutier et al., 2018; Sun et al., 2022). Very potent antimicrobial diprenylated (iso)flavonoids, however, did not show membrane permeabilization in *Listeria monocytogenes* (Araya-Cloutier et al., 2018). So far, the permeabilization capacity of prenylated (iso)flavonoids is only assessed at singular concentrations, making it impossible to establish dose-response relationships and accurately compare permeabilization potencies. In this study antimicrobial activity and permeabilization capacity of 36 different prenylated (iso)flavonoids were assessed against MRSA. Antimicrobial activity was determined by the broth microdilution assay and membrane permeabilization by measuring the uptake of propidium iodide using fluorescence microscopy and spectroscopy. Dose-response curves were established and used to extract effective concentrations (EC₁₀). No relationship between antimicrobial activity and permeabilizing capacity was found for compounds having good antimicrobial activity (minimum inhibitory concentration (MIC) ≤ 50 µg/mL). Luteone (**1**) (EC₁₀: 23±7 µg/mL) and neobavaisoflavone (**2**) (EC₁₀: 28±8 µg/mL) showed most potent permeabilization but were not the most potent antimicrobials (MIC of 25 and 38 µg/mL, respectively). Molecular descriptors analyses revealed that permeabilization (EC₁₀) could not be explained by relative hydrophobic surface area, as found for *Listeria monocytogenes*. Conversely, polar surface area was positively correlated to pEC₁₀, suggesting that membrane permeabilization occurs through interactions with polar phospholipid headgroups. These results highlight the complexity of prenylated (iso)flavonoids' antimicrobial action, suggesting other (membrane) effects besides permeabilization.

Keywords: prenylated (iso)flavonoids, methicillin-resistant *Staphylococcus aureus*, antimicrobial activity, permeabilization, propidium iodide uptake

References

- Araya-Cloutier, C., Vincken, J.-P., van de Schans, M. G. M., Hageman, J., Schaftenaar, G., den Besten, H. M. W., & Gruppen, H., 2018. QSAR-based molecular signatures of prenylated (iso)flavonoids underlying antimicrobial potency against and membrane-disruption in Gram positive and Gram negative bacteria. *Scientific reports*, 8(1), 9267.
- Sun, L., Tang, Z., Wang, M., Shi, J., Lin, Y., Sun, T., Zou, Z., Weng, Z. (2022). Exploration of Antimicrobial Ingredients in *Psoralea corylifolia* L. Seed and Related Mechanism against Methicillin-Resistant *Staphylococcus aureus*. *Molecules*, 27(20).

* Presenting author. E-mail address: janniek.ritsema@wur.nl (Janniek Ritsema)

S1.P29 *Porphyridium cruentum* as a chemical factory: valorization of high-value phycobiliproteins and lipids using an innovative solid-liquid-liquid method based entirely on eutectic solvents

Solène Odou^{a,*}, Louise VanGheluwe^a, Alia Chevalley^b, Myriam Phelippe^b, Leslie Boudesocque-Delaye^a

^a EA 7502 SIMBA, Pharmacy Faculty, University of de Tours, 31 Avenue Monge, 37200 Tours, France

^b Aqua Eco Culture, 7 Rue d'Armor Maroué, 22400 Lamballe, France

Microalgae are being studied as sustainable factories for sourcing pharmaceutical, cosmetics, and nutraceutical molecules (Sonani et al., 2014). As an example, *Porphyridium cruentum*, a red microalgae, is a valuable source of metabolites with high added value, such as free fatty acids and pigments like phycobiliproteins. Natural Deep Eutectic Solvents (NaDES) are a new environmentally friendly substitute for petrochemical solvents. NaDES are green, non-toxic, economical, and bio-based solvents formed from plant metabolites such as sugars (Dai et al, 2013). We recently developed an innovative solid/liquid/liquid (SLL) strategy for one-step extraction and purification of metabolites using a combination of polar and nonpolar NaDESs (Fig. 1) (Hilali et al., 2024).




Figure 1 : Workflow for extracting *P. cruentum* using two different processes.

A screening process was conducted on a large library of NaDES to identify the most suitable polar-nonpolar NaDES partners. Compared to the classical solid/liquid approach, the SLL process significantly increases the extractive performance of non-polar NaDES. Among all NaDES combinations screened, we highlighted good combinations to improve phycobiliprotein stability over time or to maximize lipid and phycoerythrin recovery.

Keywords: *Porphyridium cruentum*, solid-liquid-liquid extraction, phycobiliproteins, free fatty acids, NaDES

Reference

- Dai, Y., Witkamp, G.-J., Verpoorte, R., Choi, Y. H., 2013, Natural Deep Eutectic Solvents as a New Extraction Media for Phenolic Metabolites in *Carthamus tinctorius* L. *Anal. Chem.* 85, 6272–6278.
- Hilali, S., Van Gheluwe, L., Yagmur, M., Wils, L., Phelippe, M., Clément-Larosi re, B., Montigny, B., Jacquemin, J., Thiery, E., Boudesocque-Delaye, L., 2024, NaDES-based biorefinery of *Spirulina* (*Arthrospira platensis*): A new path for sustainable high value-added metabolites. *Sep. Purif. Technol.* 329, 125123
- Sonani, R. R., Singh, N. K., Kumar, J., Thakar, D., Madamwar, D., 2014, Concurrent Purification and Antioxidant Activity of Phycobiliproteins from *Lyngbya* Sp. A09DM: An Antioxidant and Anti-Aging Potential of Phycoerythrin in *Caenorhabditis Elegans*. *Process Biochem.* 49 (10), 175766p.

* Presenting author. E-mail address: solene.odou@univ-tours.fr (Sol  ne ODOU).

S1.P30 Formicine ant venoms put to the acid test - novel antimicrobial peptide families revealed by proteotranscriptomics

Lukas Koch^{a,b,*}, Simon Tragust^c, Timo H.J. Niedermeyer^b

^a RG Pharmacognosy, Institute of Pharmacy, Martin-Luther-University Halle-Wittenberg, Hoher Weg 8, 06120 Halle (Saale), Germany

^b RG Pharmaceutical Biology, Institute of Pharmacy, Freie Universität Berlin, Königin-Luise-Str. 2+4, 14195 Berlin, Germany

^c RG General Zoology, Institute of Biology, Martin-Luther-University Halle-Wittenberg, Hoher Weg 8, 06120 Halle (Saale), Germany

Research on venoms of ants (Hymenoptera: Formicidae) dates back to Samuel Fisher's first report on the isolation of formic acid from wood ants (*Formica* sp.) in 1670. Since, studies have revealed a plethora of natural products from ant venoms, including alkaloids, terpenoids, and peptides with neurotoxic, insecticidal, or antimicrobial activities (Touchard et al., 2016). Among ants, venom sprays containing concentrated formic acid are exclusively found in the Formicinae subfamily. Formicine ants use their multipurpose venoms in predation and defense, for disinfection in brood- and self-grooming, and, when swallowed, gut microbial control. (Pull et al., 2018; Tragust et al., 2020) Despite their biological importance, little is known about the composition of formicine ant venoms, and the occasionally presumed presence of uncharacterized peptide natural products has never been substantiated (Flury, 1919; Osman and Brander, 1961). Pursuing this enigma, we analyzed the venom of the carpenter ant *Camponotus nicobarensis* combining mass spectrometry and RNA sequencing. Next to various lipids, we identified three novel peptides in the venom. These peptides, the formicitoxins, are encoded by two distinct gene families exclusive to formicine ants. Intriguingly, they share the tendency to adopt amphipathic alpha-helical structures – a common feature of many antimicrobial peptides. Indeed, we found two formicitoxins to inhibit growth of the entomopathogenic fungi *Metarhizium robertsii* and *Beauveria bassiana*. In conclusion, we for the first time characterize peptides from a formicine ant venom. Crucially, these may function as antimicrobials in external immune defense and thus represent an adaptation to the microbial challenges threatening colony health.

Keywords: *Camponotus nicobarensis*, Formicidae, venom, peptides, antifungal

References

- Flury, F., 1919. Über die Bedeutung der Ameisensäure als natürlich vorkommendes Gift. Ber. Dtsch. Pharm. Ges. 29 (9), 650–669.
- Osman, M.F.H.; Brander, J., 1961. Weitere Beiträge zur Kenntnis der chemischen Zusammensetzung des Giftes von Ameisen aus der Gattung *Formica*. Z. Naturforsch. B, 16 (11), 749–753.
- Pull, C.D.; Ugelvig, L.V.; Wiesenhofer, F.; Grasse, A.V.; Tragust, S.; Schmitt, T.; Brown, M.J.F.; Cremer, S.; Tautz, D., 2018. Destructive disinfection of infected brood prevents systemic disease spread in ant colonies. eLife 7, e32073.
- Touchard, A.; Aili, S.R.; Fox, E.G.P.; Escoubas, P.; Orivel, J.; Nicholson, G.M.; Dejean, A., 2016. The Biochemical Toxin Arsenal from Ant Venoms. Toxins 8 (1): 30.
- Tragust, S.; Herrmann, C.; Häfner, J.; Braasch, R.; Tilgen, C.; Hoock, M.; Milidakis, M.A.; Gross, R.; Feldhaar, H., 2020. Formicine ants swallow their highly acidic poison for gut microbial selection and control. eLife 9, e60287.

* Presenting author. E-mail address: lukas.koch@fu-berlin.de (Lukas Koch)

S1.P31 Antibacterial activity of a novel callus culture line from *Cistus creticus*

Carmen Laezza^{a,*}, Mariavittoria Verrillo^a, Vincenzo D'Amelia^b, Maria Manuela Rigano^a

^a Department of Agricultural Sciences, University of Naples Federico II, Via Università 100, 80055 Portici, Italy

^b Institute of Bioscience and Bioresources, National Research Council, Via Università 100, 80055 Portici, Italy

Callus cultures are currently the most used biotechnological platform used to provide year-round and low cost production of natural compounds. This method can also greatly contribute to the study of specific biosynthetic pathway, speeding up the process. In the wide-ranging plant kingdom, *Cistus creticus* (Cistaceae) is a source of specialized metabolites mainly represented by catechins, monoterpenes and sesquiterpenes (Zalegh et al., 2021). Accordingly, this species has long been reported to be a medicinal plant and to be marketed as herbal infusions and dietary supplements (Viapiana et al., 2017). This is specifically due to the antimicrobial properties of *C. creticus* extract tested against several viruses (Ebola virus, HSV-1, PI-3, and dengue virus) and bacteria (*Acinetobacter baumannii* and *Pseudomonas aeruginosa*) causing severe human diseases (Zalegh et al., 2021). Our work aimed to develop a novel callus culture line from *C. creticus* hypocotyl and explore the potential of its extract against foodborne pathogens (Fig. 1. (B)).




Fig. 1. (A) *Cistus creticus* in vitro plant; (B) Hypocotyl-derived calli; (C) Antibacterial activity of hypocotyl-derived callus extract compared to an antibiotic mixture.

As shown in Fig. 1. (C), the antibacterial activity of the methanolic extract was tested against *Salmonella enterica* (Gram negative) and *Listeria monocytogenes* (Gram positive) by the disk diffusion method. The results demonstrated that the hypocotyl-derived callus extract was more than twice as effective as the antibiotic mixture (combination of ampicillin and clavulanic acid) in reducing bacterial growth. This confirms the outstanding potential of cell cultures in the production of natural phytochemicals. Further investigations need to be performed to better elucidate the specific molecules that might be involved in this function.

Keywords: Cistaceae, cell cultures, specialized metabolites, antimicrobial potential

References

- Viapiana, A., Konopacka, A., Waleron, K., Wesolowski, M., 2017. *Cistus incanus* L. commercial products as a good source of polyphenols in human diet. Ind. Crops Prod. 107, 297–304.
- Zalegh, I., Akssira, M., Bourhia, M., Mellouki, F., Rhallabi, N., Salamatullah, A.M., Alkaltham, M.S., Khalil Alyahya, H., Mhand, R.A., 2021. A review on *Cistus* sp.: phytochemical and antimicrobial activities. Plants 10, 1214.

* Presenting author. E-mail address: carmen.laezza@unina.it (Carmen Laezza)

S1.P32 Secondary metabolites produced by *Nigrospora sphaerica* associated with the invasive weed *Cenchrus ciliaris*

Maria Michela Salvatore^a, Maria Teresa Russo^a, Susan Meyer^b, Angela Tuzi^a, Marina DellaGreca^a, Marco Masi^a, Anna Andolfi^{a,*}

^a Department of Chemical Sciences, University of Naples Federico II, via Cinthia 4, 80126 Naples, Italy

^b Department of Geosciences, Southern Utah University, Cedar City, UT 84721, USA

Due to the urgent need for new strategies in managing the agricultural pests, the production of phytotoxic compounds from fungal pathogens could represent a new frontier for weed management. Phytopathogenic fungi can be employed as biocontrol agents representing good alternatives for weed management that allow reducing hazardous effects of synthetic pesticides (Masi et al. 2017A and B; Siciliano et al., 2023). *Nigrospora sphaerica* is known to be a plant pathogen, causing diseases in various plants including fruits and vegetables. In this respect, the pathogenic association of this fungus with invasive weeds suggests a possible production of phytotoxic compounds which could help in the development of new weed control strategies. A screening of secondary metabolites produced in mycelial and culture filtrate extracts of a foliar strain of *N. sphaerica* isolated from buffelgrass (*Cenchrus ciliaris*) was conducted after observing an intense activity of the organic extract on buffelgrass seed germination and subsequent seedling growth. The present communication reports the structural and stereostructural elucidation of a new triol, nigrosphaeritriol (**1**) and a new lactol, nigrosphaerilactol (**2**), together with the identification of several known compounds. Some of them are reported in Fig. 1.




Fig 1. Structures of nigrosphaeritriol, nigrosphaerilactol, nigrosphaerilactone, (S)-hydroxybutyrolactone, lupinolactone, nigrosporione A (**1-6**).

Keywords: weed, bioherbicides, specialized metabolites, Integrate Pest Management, buffelgrass

References

- Masi, M., Meyer, S., Clement, S., Pescitelli, G., Cimmino, A., Cristofaro, M., Evidente, A., 2017A. Chloromonilinic acids C and D, phytotoxic tetrasubstituted 3-chromanonacrylic acids isolated from *Cochliobolus australiensis* with potential herbicidal activity against buffelgrass (*Cenchrus ciliaris*). J. Nat. Prod. 80, 2771-2777.
- Masi, M., Meyer, S., Górecki, M., Mandoli, A., Di Bari, L., Pescitelli, G., Cimmino, A., Cristofaro, M., Clement, S., Evidente, A., 2017B. Pyriculins A and B, two monosubstituted hex-4-ene-2,3-diols and other phytotoxic metabolites produced by *Pyricularia grisea* isolated from buffelgrass (*Cenchrus ciliaris*). Chirality 29, 726-736.
- Siciliano, A., Zorrilla, J.G., Saviano, L., Cimmino, A., Guida, M., Masi, M., Meyer, S., 2023. Insights into the ecotoxicology of radicinin and (10S,11S)-(-)-epi-pyriculol, fungal metabolites with potential application for buffelgrass (*Cenchrus ciliaris*) biocontrol. Toxins 15, 405.

* Presenting author. E-mail address: andolfi@unina.it (Anna Andolfi)

S1.P33 Anti-inflammatory activity of a methanol extract of *Adiantum capillus-veneris* L.

Linru Mou^a, Bernd L. Fiebich^b, Caroline Maake^c, Francesca Scotti^a, Michael Heinrich^a

^a Research Group "Pharmacognosy and Phytotherapy, School of pharmacy, University College London, London, United Kingdom

^b Neuroimmunology and Neurochemistry Research Group, Department of Psychiatry and Psychotherapy, Medical Center–University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany

^c Institute of Anatomy, University of Zürich, Zürich, Switzerland

Four highly programmed phases are involved in wound healing, namely, haemostasis, inflammation, proliferation, and remodelling. A normal inflammation process is necessary for removing contaminated factors. However, prolonged inflammation occurs in certain circumstances, such as incomplete clearance of micro-organisms. This often leads to the transition from acute wounds to chronic wounds, resulting in a severe impact on patient's quality of life (Guo et al., 2010).

The fern *Adiantum capillus-veneris* L. (Pteriaceae) has been used traditionally as an analgesic for relieving headache, preventing hair loss, and treating asthma (Dehdari et al., 2018). In our previous research, the methanol extracts of *A. capillus-veneris* promoted proliferation and migration of human HaCaT keratinocytes at low concentrations. As part of our continuing research, we explored its anti-inflammatory activity on LPS-stimulated RAW 264.7 mouse macrophages. Using ELISA, we found that the methanol extract significantly inhibited LPS-induced release of three inflammation-related cytokines (TNF- α , IL-6, IL-1 β), and three chemokines (CXCL2, CXCL10, CCL2) in a concentration-dependant manner (0.1, 1, 10, 50 μ g/mL). None of the tested concentrations showed cytotoxic effects in RAW 264.7 cells. In addition, Western blot analyses showed that LPS-activated mitogen-activated protein kinases (MAPK) p38 MAPK and ERK1/2 were also inhibited by the methanol extracts of *A. capillus-veneris* in RAW 264.7 cells. Here, we provide first evidence that the anti-inflammatory activity of *A. capillus-veneris* observed in vivo may be mediated by its inhibitory effect on release of CXCL10, CCL2, CXCL2 and IL-1 β , and inhibition of activated p38 MAPK and ERK1/2 kinases under inflammatory conditions.

Keywords: medicinal plants, *Adiantum capillus-veneris* L., anti-inflammatory activity, wound healing

References

- Dehdari, S. and Hajimehdipour, H., 2018. Medicinal Properties of *Adiantum capillus-veneris* Linn. In Traditional Medicine and Modern Phytotherapy: A Review Article. Iranian Journal of Public Health, 47(2), 188.
- Guo, S. and DiPietro, L. A., 2010. Factors Affecting Wound Healing. Journal of Dental Research, 89(3), 219.

Acknowledgment

We gratefully acknowledge the support provided by Velux Stiftung for our research program.

* Presenting author. E-mail address: linru.mou.20@ucl.ac.uk (Linru Mou)

S1.P34 Fatty acid profile of lipids from microalgae grown on media with hydrolysed agro-food biowaste

Carlo Esposito^{a,*}, Elia Lio^a, Giancarlo Aldini^a, Francesco Secundo^b

^a Department of Pharmaceutical Sciences, University of Milan, Via Mangiagalli 25, 20133, Milan, Italy

^b National Research Council (C.N.R), Institute of Chemical Sciences and Technologies, Via Bianco Mario, 9; Milan, Italy

The growing demand for natural products for the nutraceutical, pharmaceutical and cosmetic industries has led to investigate micro and macroalgae as a source of bioactive compounds of interest, including carotenoids, polyphenols and fatty acids (Levasseur et al., 2020). Moreover, as microalgae often grow in mixotrophic conditions, enriching the culture media with a carbon source coming from biowaste, is an interesting perspective to develop sustainable processes. This presentation will concern the study of the fatty acid profiles obtained from the microalgae *Chlamydomonas reinhardtii* (CC-503 and CC-5163) and *Chlorococcum* sp. The microalgae were grown in mixotrophic conditions, employing TAP or BG-11 media containing variable percentages (5% and 15%) of hydrothermally hydrolysed biowaste coming from the agri-food industry or the environment such as orange peels, hemp or the algae *Cladophora glomerata*. In a typical procedure, the microalgae biomass was harvested after 10 days from inoculum by centrifugation and washed with water to eliminate media residues and finally freeze-dried. The fatty acids fractions were obtained by rupture and extraction of the microalgae biomasses (2-3 mgs) in two-steps carried out with 0.5 mL 0.035 M KOH in methanol (keeping the sample in these conditions overnight) and sonication for 5 min (step 1), then 0.1 mL of 2% H₂SO₄ in methanol is added (Step 2). Next, the sample was extracted with 0.6 mL of petroleum ether and analysed by gas chromatography (GC-FID)(Cooney et al., 2009). The microalgae growth and the fatty acids profiles were influenced by the addition of the biowaste hydrolysate, which significantly increased the percentage of unsaturated fatty acids, C16:2, C16:3, C18:2 and C18:3.

Financial support from the European Union – Next Generation EU in the frame of the “National Biodiversity Future Center”(CUPn°B83C22002930006) is gratefully acknowledged.

Keywords: microalgae, lipid, mixotrophy growth, *Chlamydomonas reinhardtii*, *Chlorococcum* sp

References

Cooney, M., Young, G., Nagle, N., 2009. Extraction of bio-oils from microalgae. Separation & Purification Reviews, 38(4), 291-325.

Levasseur, W., Perré, P., Pozzobon, V., 2020. A review of high value-added molecules production by microalgae in light of the classification. Biotechnology advances, 41, 107545.

* Presenting author. E-mail address: carlo.esposito@unimi.it (Carlo Esposito)

S1.P35 Cytotoxic cyclopeptides from the fungus *Gymnopus fusipes* (Omphalotaceae)

Andrea Nagyné Nedves^a, Márton Weber^b, Miklós Dékány^b, Csenge Anna Felegyi-Tóth^a, Suratno Suratno^a, Ágoston Zelei^a, Áron Szigetvári^b, Enikő Bodó^c, Zoltán Béni^b, Attila Ványolós^{a,*}

^a Department of Pharmacognosy, Semmelweis University, Üllői u. 26, H-1085 Budapest, Hungary

^b Spectroscopic Research Department, Gedeon Richter Plc., Gyömrői út 19-21, H-1103 Budapest, Hungary

^c University of Nyíregyháza, Agricultural and Molecular Research and Service Institute, H-4400 Nyíregyháza, Sóstói út 31./B, Hungary

In our ongoing research focused on identifying fungal natural products with anticancer potential, we have discovered novel cyclic octadecapeptides from the mushroom *Gymnopus fusipes* (Bull.: Fr.) Gray. Commonly referred to as spindleshank, *G. fusipes* is a parasitic species (Camy et al., 2003) within the Omphalotaceae family, native to woodlands of Europe and Asia. The mushroom material was freeze-dried followed by extraction using methanol. The resultant crude extract was subjected to normal-phase flash chromatography. The ultimate purification was accomplished through reversed-phase HPLC, leading to the novel gymnopeptides C and D, besides the previously known gymnopeptides A and B (Ványolós et al., 2016). The structure elucidation of the isolated metabolites was performed by extensive NMR and MS analysis. The absolute configurations of the amino acid residues were examined using HPLC–MS. Initially, the cyclopeptides underwent acidic hydrolysis, followed by derivatization of the resultant amino acid mixture with Marfey's reagent. Subsequently, HPLC–MS analysis was conducted, wherein the retention times corresponding to the accurate mass values of the derivatized amino acids in the extract ion chromatograms were compared with those observed for the derivatized reference L,D-amino acid pairs. The antiproliferative properties of the isolated peptides were determined by means of a 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay on human cancer cell lines. Gymnopeptides A-D demonstrated very strong cell growth inhibitory activity with nanomolar IC₅₀ values.

Acknowledgements: The authors are grateful for the support of the National Research, Development and Innovation Office, Hungary (grant: OTKA NKFIH FK-146487).

Keywords: *Gymnopus fusipes*, Omphalotaceae, cyclic peptides, cytotoxicity, fungus

References

- Camy, C., Delatour, C., Caël, O., Marçais, B. 2003. Inoculation of mature pedunculate oaks (*Quercus robur*) with the root rot fungus *Collybia fusipes*: Relationships with tree vigour and soil factors. Eur. J. Plant Pathol. 109, 545–553.
- Ványolós, A., Dékány M., Kovács, B., Krámos, B., Bérdi, P., Zupkó, I., Hohmann, J., and Béni, Z. 2016. Gymnopeptides A and B, Cyclic Octadecapeptides from the Mushroom *Gymnopus fusipes*. Org. Lett. 18, 2688–2691.

* Presenting author. E-mail address: vanyolos.attila@semmelweis.hu. (Attila Ványolós)

S1.P36 Cytotoxic hederagenin saponins from the seeds of *Oxybasis rubra* (L.) S. Fuentes, Uotila & Borsch (Amaranthaceae)

Irma Podolak, Karolina Grabowska*, Adam Mynarski, Dagmara Wróbel-Biedrawa, Agnieszka Galanty

Department of Pharmacognosy, Faculty of Pharmacy, Jagiellonian University Medical College, 9 Medyczna Str., 30-688 Kraków, Poland

This work reports isolation, structure elucidation and cytotoxic activity evaluation of two triterpene saponins (**1,2**) from *Oxybasis rubra* (L.) S. Fuentes, Uotila & Borsch (Amaranthaceae). The species (syn. *Chenopodium rubrum* L.) is an annual plant, widely distributed in temperate zone, with characteristic shiny leaves. It has been mainly used as a dye, and as a food ingredient, but is also a folk medicinal anticancer remedy (Kokanova-Nedialkova et al., 2009).

Our on-going studies on triterpene saponins from various plant sources are focused on the evaluation of their cytotoxic potential. As Amaranthaceae is a known source of saponin-plants we decided to analyze *O. rubra*, which is a phytochemically under-explored species. MeOH extracts from different plant parts were initially tested for cytotoxic activity (Mynarski et al. 2018). Chromatographic separation of fruit extract (MPLC, CC) led to the isolation of two compounds, which, following extensive NMR spectroscopy and HRESIMS, have been identified as 3-O- β -D-glucopyranosyl(1 $\rightarrow$ 3)- β -D-glucopyranosyl] hederagenin **1** (a new structure) and 3-O- β -D-glucopyranosyl(1 $\rightarrow$ 3)- β -D-glucopyranosyl] hederagenin 28-O- β -D-glucopyranosyl ester **2** (Fig. 1).




Fig. 1. Structures of hederagenin saponins **1** and **2** from *Oxybasis rubra*

Both saponins were tested on a wide panel of 11 human cancer cell lines using colorimetric LDH assay. Normal human cells were included to assess selectivity of action. In general, the bidesmosidic saponin was less active compared to the monodesmoside. Both saponins showed moderate activity and were most active against lung cancer A549 (IC₅₀ 8.24 and 11.8 μ g/mL) while best selectivity was observed for **2** (IC₅₀ 21.09 μ g/mL and >100 μ g/mL, against cancer PC3 and normal prostate PNT2 cells, respectively).

Keywords: *Oxybasis rubra*, Amaranthaceae, hederagenin saponins, cytotoxicity

References

- Kokanova-Nedialkova, Z., Nedialkov, P.T., Nikolov, S.D., 2009. The genus *Chenopodium*: phytochemistry, ethnopharmacology and pharmacology. *Pharmacogn. Rev.* 3, 280–306.
- Mynarski, A., Pietrzak, W., Galanty, A., Dawiec, E., Nowak, R., Podolak, I., 2018. Phenolic Acid LC/MS Profile of *Chenopodium rubrum* and Evaluation of Cytotoxic Activity. *Nat. Prod. Commun.* 13, 855–857.

* Presenting author. E-mail address: karolina1.grabowska@uj.edu.pl (Karolina Grabowska)

S1.P37 Untargeted and targeted LC-MS metabolomic analyses of putative chemical defenses in *Taenaris* butterflies

Belkes N. Stambouli^{a,b,*}, Ella T. Vardeman^{a,b,c}, Edward J. Kennelly^{a,b}, David J. Lohman^{b,d,e}

^a Department of Biological Sciences, Lehman College, City University of New York, Bronx, NY 10468, USA

^b PhD Program in Biology, The Graduate Center, City University of New York, 365 Fifth Ave., NY 10016, USA

^c The New York Botanical Garden, 2900 Southern Blvd, The Bronx, NY 10458, USA

^d Department of Biology, City College of New York, 160 Convent Ave., New York, NY 10031, USA

^e Entomology Section, National Museum of Natural History, Rizal Park, Manila, 1000, Philippines

The potential chemical defense mechanisms of *Taenaris* butterflies, a group of 25 species found in New Guinea and nearby islands, was investigated. While the chemical defense of herbivorous insects in temperate regions are reasonably well-studied (Agrawal, 2017), tropical taxa, including *Taenaris*, have received limited attention. *Taenaris* butterflies exhibit characteristics of being chemically defended (aposematic), such as mimicry by other species (Lohman et al., 2020). We examined the hypothesis (Parsons, 1984, 1998) that adult *Taenaris* butterflies are defended by cycad-derived compounds, including cycasin, macrozamin, and methylazoxymethanol acetate, acquired through larval or adult feeding. Contrary to previous findings, our comprehensive metabolomic analyses fail to detect these compounds in *Taenaris* samples. This suggests that cycad-derived compounds are not present in *Taenaris* butterflies, though compound degradation, instrument limitations, or sample handling may have prevented detection. A multivariate analysis highlighted metabolomic differences between *Taenaris* and their close relative *Faunis eumeus*, which prompted further exploration into taxon-specific defensive chemical features. Field observations on Yapen Island suggest that *Taenaris* adults may acquire defensive compounds via pharmacophagy from sources other than cycads, challenging published observations suggesting that species rely on cycad-derived toxins. These findings contribute to our understanding of the ecological and evolutionary dynamics of butterfly communities in tropical regions.

Keywords: chemical defense, Cycadaceae, cycasin, LC-MS metabolomics, mimicry

References

- Agrawal, A., 2017. *Monarchs and Milkweed: A Migrating Butterfly, a Poisonous Plant, and their Remarkable Story of Coevolution*. Princeton: Princeton University Press, 296 pp.
- Lohman, D. J., Sarino, and D. Pegg. 2020. Syntopic *Elymnias agondas aruana* female forms mimic different *Taenaris* model species (Papilionoidea: Nymphalidae: Satyrinae) on Aru, Indonesia. *Treubia* 47,1-12.
- Parsons, M., 1984. Life histories of *Taenaris* (Nymphalidae) from Papua New Guinea. *J. Lepid. Soc.* 50, 261–268 Parsons, M., 1998. *The Butterflies of Papua New Guinea*. San Diego Acad. Press

*Presenting author. E-mail address: bstambouli@gradcenter.cuny.edu (Belkes Stambouli)

S1.P38 Multimodal analysis of oak wood metabolites for a comprehensive understanding of the ageing process in French typical spirits barrels

Aline Cournut^{a,b,*}, Quentin P. Vanbellingen^a, Anaïs Demaye^b, Andreas Römpf^c, David Touboul^d, Alain Brunelle^a, Véronique Eparvier^b

^a Laboratoire d'Archéologie Moléculaire et Structurale, LAMS UMR8220, CNRS, Sorbonne Université 75005 Paris, France

^b Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, UPR 2301, 91190 Gif-sur-Yvette, France

^c Bioanalytical Sciences and Food Analysis, University of Bayreuth, Bayreuth, Germany

^d LCM, CNRS, Ecole Polytechnique, Institut Polytechnique de Paris, 91120 Palaiseau, France

Oak wood is vital in the barrel ageing process of spirits, impacting their flavor and quality (Canas, 2017). To optimize this process, understanding the dynamics of wood metabolites is crucial. In our study, we employed a comprehensive and multimodal analytical approach to investigate the spatial distribution of central and specialized metabolites in oak wood. To achieve this, we conducted a detailed exploration using various analytical techniques such as Mass Spectrometry in tandem, Laser Desorption/Ionization, and Time-of-Flight Secondary Ion Mass Spectrometry (Vanbellingen et al., 2016), to study the distribution of metabolites in oak wood according different parameters. We analyzed metabolite profiles of samples from freshly cut or toasted oak compared to a 150-year-old oak stave. By mapping the distribution of lignins, polysaccharides, lipids, and phenolic compounds at the subcellular level and across the stave thickness, we gained a comprehensive understanding of the metabolite dynamics within the wood. One notable finding was the significant decline in polysaccharide fragment ions observed in the 150-year-old oak stave. This decrease suggests potential degradation of polysaccharides over time, which could have implications for the ageing process of spirits in oak barrels. In addition, investigation of the spatial distribution of inorganic compounds within the oak wood revealed unique patterns that could influence oak growth and durability. Overall, our study provides a fresh perspective on the shifts in metabolites within oak wood over time. The insights gained from this research can contribute to the development of refined barrel aging techniques, ultimately leading to better spirit quality.

Keywords: oak wood, TOF-SIMS, LC-MS, LDI-MS, molecular network

References

- Canas, S. 2017. Phenolic composition and related properties of aged wine spirits: influence of barrel characteristics. A review. *Beverages*. 3, 55.
- Vanbellingen, Q. P., Fu, T., Bich, C., Amusant, N., Stien, D., Della-Negra, S., Touboul, D., Brunelle, A. 2016. Mapping *Dicorynia guianensis* Amsh. wood constituents by submicron resolution cluster-TOF-SIMS imaging. *J. Mass Spectrom.* 51, 412–423.

* Presenting author. E-mail address: aline.cournut@cnrs.fr (Aline Cournut)

S1.P39 Exometabolites from the Mediterranean sponge *Agelas oroides*: where is oroidin?

Morgane Mauduit, Stéphane Greff, Thierry Pérez, Charlotte Simmler*

Institut Méditerranéen de Biodiversité et d'Écologie Marine et Continentale (IMBE), UMR CNRS 7263, IRD 237, Aix Marseille Université, Avignon Université, Endoume Marine Station, rue de la batterie des lions, 13007 Marseille, France

Sponges are filter-feeding sessile marine animals that are keystone species of various benthic ecosystems. They are known to produce a plethora of structurally diverse metabolites. Some of them are part of their toolbox for defense and communication strategies. When released, these (exo)metabolites may contribute to distant species interactions (Hay 2009; Santonja et al. 2018). In the course of our studies on the chemical diversity of exometabolites from sponges, we wanted to know whether *Agelas oroides* (Schmidt, 1864), a rather dominant species of Mediterranean marine ecosystems, releases its characteristic bromo-pyrrole alkaloids. A series of captures and enrichment experiments were performed in aquaria and in situ (Mauduit et al. 2023). MS-based metabolomics revealed the paucity of bromo-pyrrole alkaloids as exometabolites. Oroidin was not detected in seawater despite being the most abundant compound in *A. oroides* extract (Fig. 1), and as opposed to published results with *Agelas conifera* (Richelle-Maurer et al. 2003).




Fig. 1. Stacked chromatograms of *A. oroides* extract containing 615% w/w of oroidin.

Such findings prompted us to explore the fate of oroidin from the sponge biomass to the seawater. The presentation will delve into these experimentations, and illustrate how examining organisms in their ecosystems can lead to new discoveries even for extensively studied species.

Keywords: sponges, exometabolites, oroidin, biotransformation, MS-based metabolomics

References

- Hay, M.E., 2009. Marine Chemical Ecology: Chemical Signals and Cues Structure Marine Populations, Communities, and Ecosystems. Annual Review of Marine Science 1, 193–212.
- Mauduit, M, Derrien M, Grenier M, Greff S, Molinari S, Chevaldonné P, Simmler C, and Pérez T. 2023. *In Situ* Capture and Real-Time Enrichment of Marine Chemical Diversity. ACS Central Science 9, 2084–2095.
- Richelle-Maurer E, De Kluijver M.J, Feio S, Gaudêncio S, Gaspar H, Gomez R, Tavares R, Van de Vyver G, Van Soest R.W.M. 2003. Localization and Ecological Significance of Oroidin and Scepterin in the Caribbean Sponge *Agelas conifera*. Biochemical Systematics and Ecology 31 (10): 1073–91.
- Santonja, M, Greff S, Le Croller M, Thomas O.P, Pérez T. 2018. Distance Interaction between Marine Cave-Dwelling Sponges and Crustaceans." Marine Biology 165, 1–15.

* Presenting author. E-mail address: charlotte.simmler@imbe.fr (Charlotte Simmler)

S1.P40 Anti-platelet aggregation and vasorelaxant activity of rapanone from *Ardisia crenata* Sims

Dagmara Wróbel-Biedrawa^{a,*}, Monika Kubacka^b, Karolina Grabowska^a, Agnieszka Galanty^a, Irma Podolak^a

^a Department of Pharmacognosy, Faculty of Pharmacy, Jagiellonian University Medical College, 9 Medyczna Str., 30-688 Kraków, Poland

^b Department of Pharmacology, Faculty of Pharmacy, Jagiellonian University Medical College, 9 Medyczna Str., 30-688 Kraków, Poland

The results of the study that will be presented concern the isolation and biological evaluation of naturally occurring benzoquinone, rapanone, from leaves of *Ardisia crenata* Sims (Primulaceae). This shrub can achieve 1-2 m of height, has characteristic dark green leaves with tightly waved edges and originally red berries. *A. crenata* occurs naturally in Southern and Eastern Asia and in Europe it is cultivated as a greenhouse plant due to its ornamental value (Larson et al., 2023; Lim et al., 2012). As rapanone (Fig.1) was recognized as an anti-inflammatory and antioxidant agent, we decided to check its potential value in non-communicable diseases which are characterized by chronic inflammation, such as e.g. cardiovascular disorders (Geto et al., 2020). Thus, its anti-platelet and vasorelaxant activity was evaluated. Rapanone has been isolated from the air-dried leaves of *A. crenata* using extraction protocol developed by us previously (Wróbel-Biedrawa et al., 2022). Ethyl acetate UAE extract was separated by chromatographic methods (CC, pTLC) to obtain rapanone. The purity and identity of the isolate was checked by LC-MS/MS and ¹³C NMR.




Fig.1. Structure of rapanone.

Rapanone was tested in an *in vitro* aggregation test and was shown to inhibit collagen induced platelet aggregation with $IC_{50}=40.2\pm18.6\ \mu M$. Furthermore, in an *ex vivo* assessment of the effect in the rat aorta precontracted with phenylephrine, the compound exhibited concentration-dependent (in the range of 0.3 – 100 μM) relaxation in endothelium intact aortic rings, reaching maximal effect at the level of almost 50%.

Keywords: *Ardisia crenata*, Primulaceae, anti-platelet, vasorelaxation, cardiovascular diseases

References

- Geto, Z., Molla, M.D., Challa, F., Belay, Y., Getahun, T., 2020. Mitochondrial dynamic dysfunction as a main triggering factor for inflammation associated chronic non-communicable diseases. *J. Inflamm. Res.* 97-107.
- Larson, D.A., Chanderbali, A.S., Maurin, O., Gonçalves, D.J., Dick, C. W., Soltis D.E., Fritsch P. W., Clarkson J.J., Grall A., Davies N.M.J., Larridon I., Kikichu I.A.B.S., Forest F., Baker W.J., Smith S.A., Utteridge, T.M., 2023. The phylogeny and global biogeography of Primulaceae based on high-throughput DNA sequence data. *Mol. Phylogenet. Evol.* 182, 107702.
- Lim, T.K., Lim, T.K., 2012. *Ardisia crenata*. Edible medicinal and non-medicinal plants: Volume 4, Fruits. 65-71. Wróbel-Biedrawa, D., Galanty, A., Zagrodzki, P., Podolak, I., 2022. Optimization of extraction conditions and cytotoxic activity of rapanone in comparison to its homologue, embelin. *Molecules.* 27(22), 7912.

* Presenting author. E-mail address: dagmara.wrobel-biedrawa@uj.edu.pl (Dagmara Wróbel-Biedrawa)

S1.P41 Unlocking bioactivity of endophytic fungi from Mokrzański Forest in Wrocław, Poland

El-Sayed R. El-Sayed^{a,b,*}, Filip Boratyński^b, Joanna Grzelczyk^c

^a Department of Food Chemistry and Biocatalysis, Wrocław University of Environmental and Life Sciences, Norwida 25, 50-375 Wrocław, Poland

^b Plant Research Department, Nuclear Research Center, Egyptian Atomic Energy Authority, Cairo, Egypt

^c Institute of Food Technology and Analysis, Lodz University of Technology, Stefanowskiego 2/22, 90-537, Lodz, Poland

This presentation will cover the isolation of fungal endophytes from some plant species of a local forest in Wrocław, Poland, and the testing of several biological activities of their extracts. Currently, the search for new bioactive compounds with innovative modes of action and chemistry is desperately needed to discover new drug leads (Miethke et al. 2021). Indeed, the increased emergence of drug-resistant microbes is a dramatic example of such pressing needs. Generally, forests are unique habitats where the growing plants face extreme environmental conditions, such as nutrient deficiency, high oxidative stress, etc (Terhonen et al. 2019). Therefore, the associated endophytic fungi of these plants are more metabolically active than their counterparts, and as a result, may biosynthesize novel bioactive chemical scaffolds (El-Sayed et al. 2021). With this view, the presented research was conducted for the isolation of fungal endophytes of twelve different plant species viz., *Larix decidua*, *Pinus sylvestris*, *Corylus avellana*, *Alnus glutinosa*, *Quercus robur*, *Fagus sylvatica*, *Picea abies*, *Populus tremula*, *Sorbus aucuparia*, *Acer platanoides*, *Betula pendula*, and *Robinia pseudoacacia*. A total of 46 endophytic fungi were isolated and tested for bioactive metabolites with antifungal, antibacterial, antioxidant, and anti-enzymatic (monoamine oxidase A, acetyl, and butyl cholinesterase inhibitors) properties in their extracts. Upon screening, several fungal isolates showed promising activities. Extracts with activities were fractionated by TLC and then re-tested. Current work is to chemically identify compounds responsible for these activities. The presented research also explains the high value of fungal endophytes from forest plants as untapped sources of bioactive metabolites.

Acknowledgments: The presented research is part of the BioExplor project No. 2021/43/P/NZ9/02241 co-funded by the National Science Centre and the European Union Framework Programme for Research and Innovation Horizon 2020 under the Marie Skłodowska-Curie grant agreement no. 945339.

Keywords: endophytic fungi, bioactivities, cytotoxicity, antimicrobial, anti-enzymatic

References

Miethke, M., Pieroni, M., Weber, T. et al., 2021. Towards the sustainable discovery and development of new antibiotics.

Nat Rev Chem 5, 726–749.

Terhonen, E., Blumenstein, K., Kovalchuk, A., Asiegbu, F.O., 2019. Forest tree microbiomes and associated fungal endophytes: Functional roles and impact on forest health. *Forests* 10, 42.

El-Sayed, E.R., 2021. Discovery of the anticancer drug vinblastine from the endophytic *Alternaria alternata* and yield improvement by gamma irradiation mutagenesis. *J Appl Microbiol* 131, 2886–2898.

* Presenting author. E-mail address: elsayed.ali@upwr.edu.pl; sayed_zahran2000@yahoo.com (El-Sayed R. El-Sayed)

S1.P42 HME-DDS-applied astaxanthin nanocapsule suppresses DNCB-induced atopic dermatitis-like inflammation in BALB/c mice

Tae-Young Gil^a, Ha-Yeon Sim^b, Suji Ryu^c, Ha Yeon Lee^c, Jong-Suep Baek^{c,d}, Dae Geun Kim^e,
Hyo-Jin An^{a,b,*}

^a Department of Oriental Pharmaceutical Science, College of Pharmacy, Kyung Hee University, Seoul 02447, Republic of Korea

^b Department of Integrated Drug Development and Natural Products, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

^c Department of Bio-Health Convergence, Kangwon National University, Chuncheon 24341, Republic of Korea

^d BeNatureBioLab, Chuncheon 24206, Republic of Korea

^e LED Agri-bio Fusion Technology Research Center (LAFTRC), Jeonbuk National University, Jeonbuk 54596, Republic of Korea

Astaxanthin is a carotenoid as a natural product be found in various bacteria or microalgae. Among the different resources, microalga *Haematococcus pluvialis* is known for the most potential source for the biological production. Astaxanthin has broad latent abilities on skin health; anti-aging, anti-oxidant, and anti-inflammatory. Since the skin plays a crucial role as the outermost barrier against external surroundings, keeping the skin health is one of the important ways to keep the immunological balance. Though the high capacity on the skin biology and related application of astaxanthin, its poorly low solubility and difficult processes lower the possibility of utilization. The authors applied hot melt extrusion (HME)-drug delivery system (DDS) on the natural product (HDA) leading high contents and usages of astaxanthin. The authors investigated the effects of HDA on 2,4-dinitrochlorbenzene (DNCB)-induced atopic dermatitis (AD)-like skin inflammation in an animal model for 4 weeks. Comparison between HDA and non-process raw astaxanthin material (NRA) was conducted. Repeated topicalization of DNCB caused the pathological symptoms on the dorsal skin including edema, redness, and dryness. Mice were orally administered with HDA showed relatively low itching scratching behavior frequency and less scaly areas than NRA-treated ones. Also, high skin barrier index was shown in HDA treated group than in the group with NRA; index was measured by transepidermal water loss and severity of symptoms. With these results, the authors assumed the HDA showed ameliorative effects on DNCB-induced AD mouse model comparing to the NRA.

Acknowledgments: This research supported by Korea Institute of Marine Science & Technology Promotion (KIMST) funded by the Ministry of Oceans and Fisheries (grant # RS-2023-00254674).

Keywords: *Astaxanthin Nanocapsule, HME-DDS, DNCB, atopic dermatitis, skin barrier*

References

- Davinelli S, Nielsen ME, Scapagnini G. Astaxanthin in Skin Health, Repair, and Disease: A Comprehensive Review. *Nutrients*. 2018 Apr 22;10(4):522. doi: 10.3390/nu10040522. PMID: 29690549.
- Mularczyk M, Michalak I, Marycz K. Astaxanthin and other Nutrients from *Haematococcus pluvialis*-Multifunctional Applications. *Mar Drugs*. 2020 Sep 7;18(9):459. doi: 10.3390/md18090459. PMID: 3290661
- Simpson EL. Atopic dermatitis: a review of topical treatment options. *Curr Med Res Opin*. 2010 Mar;26(3):633-40. doi: 10.1185/03007990903512156. PMID: 20070141.

* Presenting author. E-mail address: hjan@khu.ac.kr (Hyo-Jin An)

S1.P43 Spermidine alkaloids of the native Australian plant *Acacia auriculiformis*

Laura Coe^{a,*}, Joanne Blanchfield^a, David Leach^b, James De Voss^a

^a School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD, 4072, Australia

^b Integria Healthcare, Eight Mile Plains, QLD, 4113, Australia

Australian Aboriginal culture is one of the oldest enduring cultures on Earth. This ethnomedicinal knowledge offers a curated library of traditionally used and medicinally active plants (Mani et al. 2021). Our group is exploring the phytochemical profile of a range of native Australian plants used in traditional Aboriginal medicine. *Acacia auriculiformis* from the family *Fabaceae/Leguminosae* is a small tree native to northern regions of Australia and parts of Indonesia and Papua New Guinea (Boland et al., 1990). More commonly known as the 'ear-pod wattle', the pods and leaves have been used traditionally by Aboriginal communities to prepare an antiseptic skin wash and to relieve itching skin conditions. Crushed plant material may also be thrown in to watering holes to kill fish for later collection (Smith et al., 199q; Marrfurra, 1995). Our preliminary investigations indicated that the leaves and seed pods were rich in alkaloids that had not previously been explored. We now report the isolation and structure elucidation of a family of alkaloids using 1D and 2D ¹H, ¹³C and ¹⁵N NMR techniques and high-resolution mass spectrometry. These compounds contain an unusual substituted perhydro-1,5-diazocine-2-one structure (Fig. 1) and are previously unreported.




Fig. 1. General structure of isolated perhydro-1,5-diazocine-2-one compound family.

Keywords: *Acacia auriculiformis*, ethnopharmacology, alkaloids, diazocine

References

- Mani, J.S., Johnson, J.B., Hosking, H., Ashwath, N., Walsh, K.B., Neilsen, P.M., Broszczak, D.A., Naiker, M., 2020. Antioxidative and therapeutic potential of selected Australian plants: A review. *J. Ethnopharmacol.* 268, 113580.
- D.J. Boland; Pinyopusarerk, K.; McDonald, M. W.; Jovanovic, T.; Booth, T. H., The habitat of *Acacia auriculiformis* and probable factors associated with its distribution. *J. Trop. For. Sci.* 1990, 3, 159-180.
- Smith, N.M., Ethnobotanical Field Notes From The Northern Territory, Australia. *J. Adelaide Bot. Gard.* 1991, 14 (1), 1- 65.
- Marrfurra, P. A., M; Wawul, M; Kumunerrin, T; Adya, H; Kamarrama, K; Kanintyanyu, M; Waya, T; Kannyi, M; Wightman, G; Williams, L., 1995. Ngan'gikurunggurr and Ngangiwumirri ethnobotany: Aboriginal plant use from the Daly River area, Northern Australia. Conservation Commission of the Northern Territory: Darwin, 1995.

* Presenting author. E-mail address: l.coe@uq.net.au (Laura J. Coe)


S1.P44 New bioactive peptaibols from a soil fungus, *Trichoderma strigosum* (Hypocreaceae)

Yun Seo Park^{a,*}, Jeyong Jung^a, Jin Woo Lee^b, Sang Hee Shim^a

^a Natural Products Research Institute, College of Pharmacy, Seoul National University, Seoul 08826, Republic of Korea

^b College of Pharmacy, Duksung Women's University, Seoul 01369, Republic of Korea

The genus *Trichoderma*, commonly found in soil, is known to produce cytotoxic peptides with unique amino acid sequences, known as peptaibols. In an effort to discover bioactive secondary metabolites from fungi, *Trichoderma strigosum* was isolated from soil in north part of Seoul. Large scale culture for chemical investigation followed by LC-MS-guided isolation led to isolation five new lipopeptaibols (**1-5**) and eight new 19-residue peptaibols (**8-15**) along with two known lipopeptaibols, lipovelutibols C (**6**) and D (**7**). The planar structures of the newly discovered peptaibols (**1-5**, **8-15**) were elucidated through a comprehensive analysis of 1D, 2D NMR spectral data and UPLC-MS/MS data. Additionally, the determination of absolute configuration for new peptaibols (**1-5**, **8-15**) was achieved using the advanced Marfey's method and GTC (2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl isothiocyanate) derivatization. Co-cultivation with a plant pathogen, *Fusarium* sp. increased the production of peptaibols, supporting that *Trichoderma* in soil might help to protect plants from external pathogens. In addition, *T. strigosum* extracts exhibited a remarkable 72.4% inhibition against the triple-negative breast cancer cell line, MDA-MB-231 in a preliminary cytotoxicity test. Detailed discussion on biological activities of the isolated peptaibols (**1-15**) will be presented.



Keywords: *Trichoderma strigosum*, peptaibols, UPLC-MS/MS, advanced Marfey's method, cytotoxicity

References

- Qingfeng, G.; Lei, S.; Xinyang, W.; Dandan, L.; Zhenhua, Y.; Juanjuan, Z.; Gang, D.; Lin, C., 2023. Structures and Biological Activities of Secondary Metabolites from the *Trichoderma* genus (Covering 2018-2022). *J. Agric. Food Chem.* 71, 37, 13612-13632
- Sarath P. D, S.; Dongdong, W.; Chang-Kwon, K.; Emily A, S.; David E, D.; Patrick A, A.; Karen L, W.; Andrew G, S.; Deborah K, M.; Robert H, C.; Curtis J, H.; John A, B., 2022. Tolypocladamides A-G: Cytotoxic Peptaibols from *Tolypocladium inflatum*. *J. Nat. Prod.* 85, 6, 1603-1616.

* Presenting author. E-mail address: asdddb@snu.ac.kr (Yun Seo Park)

S1.P45 Evaluation of bioisosteric 6,5-fused heterocyclic salvinorin A analogues

Ashlyn Standley*, Ava Dickens, Adam Keasling

Department of Chemistry, Andrew College, Cuthbert, GA 39840, USA

Salvinorin A (**1**) is a neoclerodane diterpenoid isolated from the leaves of *Salvia divinorum*. It represents a pharmacologically unique natural product in that it is a subtype-selective opioid receptor agonist with high binding affinity towards the kappa opioid receptor (Roth et al., 2002). Additionally, it exhibits a comparatively safe physiological profile, being well tolerated in clinical trials, and is supported by a growing body of literature identifying potentially useful clinical applications, including: antinociceptive, antiaddictive, and neuroprotective effects (Moreira et al., 2023). These potential applications support the continued characterization of the structure-activity-relationship of salvinorin A through the development of analogues as essential molecular probes.




Fig. 1. Structure of salvinorin A with the C(22)-substituted position indicated.

In this study, we evaluated a series of C(22)-6,5-fused heterocyclic salvinorin A analogues by exploring the effect of bioisosteric replacement through incorporation of divalent substitutions into the fused five-member ring. *In vitro* evaluation consisted of competitive radioligand binding affinity at delta, kappa, and mu opioid receptors, followed by functional [³⁵S]GTP[γS] binding activity assays (Leon et al., 2013). This resulted in analogues exhibiting mix-opioid receptor binding affinity and functional activity (EC₅₀ < 200 nM). However, a declining trend in both binding affinity and functional efficacy was observed in order of benzofuran > indole > benzothiophene in C(22)-substitution when compared to salvinorin A. These results support further pharmacological characterization, especially *in vivo* assessment, to ascertain the potential physiological consequences of these exhibited trends.

Keywords: *Salvia divinorum*, salvinorin A, opioid, SAR

References

- Leon, F., Gao, J., Dale, O.R., Wu, Y., Habib, E., Husni, A.S., Hill, R.A., Cutler, S.J. 2013. Secondary metabolites from *Eupenicillium parvum* and their *in vitro* binding affinity for human opioid and cannabinoid receptors. *Planta Med.* 79, 1756-1761.
- Moreira, C.V.L., Faria, A.L., Silva, D.P.B., Ghedini, P.C., Martins, J.L.R., Keasling, A.W., Zjawiony, J.K., Pandey, P., Doerksen, R.J., Napalitano, H.B., Rocha, F.F., Costa, E.A., Fajemiroye, J.O. 2023. Heteroaromatic salvinorin A analogues (P-3l) elicits antinociceptive and anxiolytic-like effects. *Fitoterapia.* 167, 105488.
- Roth, B.L., Baner, K., Westkaemper, R., Siebert, D., Rice, K.C., Steinberg, S., Ernsberger, P., Rothman, R.B. 2002. Salvinorin A: a potent naturally occurring nonnitrogenous kappa opioid selective agonist. *PNAS USA.* 99, 11934-11939.

* Presenting author. E-mail address: standley0147@andrewcollege.edu (Ashlyn Standley)

S1.P46 Two novel saturated alicyclic carboxylic acids in *Ginkgo biloba* (Ginkgoaceae) leaf extract EGb 761

Steffen Reinhard, Julia Lutz, Vincent Schüler, Žarko Kulić*

Preclinical Research and Development, Dr. Willmar Schwabe GmbH & Co. KG, 76227 Karlsruhe, Germany

We report about the two saturated alicyclic carboxylic acids (1*S*,3*R*,4*S*)-3,4-dihydroxycyclo-hexane-1-carboxylic acid (**1**) and (1*R*,3*R*,4*R*,5*R*)-3,4,5-trihydroxycyclohexane-1-carboxylic acid ((1*R*)- dihydroshikimic acid) (**2**) (Fig. 1). The two compounds were purified from *Ginkgo biloba* leaf extract EGb 761[®] by preparative HPLC. Molecular formulae were determined by LC-HRMS and structural constitutions and relative stereochemical configurations were elucidated by NMR spectroscopy. Absolute stereochemical configurations were determined by stereoselective synthesis and hydrogenation of shikimic acid to afford **1** and **2**, respectively, followed by CD spectroscopy. The relative concentration of the two compounds in 5 batches of proanthocyanidin reduced EGb 761[®] were determined by HPLC coupled to a charged aerosol detector to be 2.6 % and 2.9 % for **1** and **2**, respectively. So far, these two substances have been described in literature as metabolites of quinic acid formed by microorganisms (Whiting et al., 1971) or rodents (Brewster et al., 1978). Our study is the first report of the two substances being present in EGb 761[®] and plant extracts in general to the best of our knowledge.




Fig. 1. Structures of the two saturated alicyclic carboxylic acids in EGb 761[®].

Keywords: *Ginkgo biloba* leaf extract, Ginkgoaceae, structure elucidation, stereochemistry, alicyclic organic acids

Conflicts of interest: All authors are employees or were interns (JL) of Dr. Willmar Schwabe GmbH & Co. KG, Germany, manufacturer of *Ginkgo* leaf extract EGb 761[®].

References

- Whiting G.C., Coggins R.A. The role of quinate and shikimate in the metabolism of lactobacilli. *Antonie van Leeuwenhoek* 1971; 37: 33–49.
Brewster D., Jones R.S., Parke D.V. The metabolism of shikimate in the rat. *Biochem J* 1978; 170 (2): 257–264.

* Corresponding author. E-mail address: zarko.kulic@schwabe.de (Žarko Kulić)

S1.P47 Microalgae: versatile microbial factories for natural product synthesis

Elia Lio^{a,b,*}, Gandolfi Stefano^a, Jacopo Pains^a, Carlo Esposito^{a,b}, Martina Dramis^a, Floriana Basile^a, Gianluca Ottolina^a, Francesco Secundo^a

^a Istituto di Scienze e Tecnologie Chimiche “Giulio Natta”, CNR, Via Mario Bianco 9, 20131 Milan, Italy

^b University of Milan, Department of Pharmaceutical Sciences, Milano, Italy

Recent advancements in microbial biotechnology have underscored the potential of photosynthetic microorganisms, such as microalgae, for industrial substance production. This study focused on optimizing the growth conditions of the microalga *Chlorella sorokiniana*, under both autotrophic and mixotrophic conditions (Perez-Garcia et al., 2015), employing orange peel extract and spent mushroom substrate to stimulate biomass growth. The utilization of orange peel extract led to a 17- fold increase in biomass compared to autotrophic conditions. Furthermore, efforts were made to valorize the biomass by extracting the lipophilic fraction using green organic solvents like dimethyl carbonate and identifying compounds with antimicrobial properties through principal component analysis (PCA) of GC-MS spectra of raw extracts. Remarkable antimicrobial activities were observed primarily against *Escherichia coli* and *Bacillus megaterium*. Emphasis was placed on elucidating the impact of various growth conditions on the quantity and composition of lipids, particularly polyunsaturated fatty acids, with significant implications for healthcare and biotechnological applications (Lakshimi et al., 2023). However, extracts obtained from biomass grown under mixotrophic conditions exhibited a reduction in fatty acid content compared to those grown under autotrophic conditions. In conclusion, this study highlights the potential of microalgae to produce bioactive molecules, with particular attention to optimizing growth conditions and valorizing biomass for antimicrobial and nutritional applications.

Acknowledgments: Financial support from Progetti di Ricerca@CNR, Delibera 197/2021 del 21/12/2021, Progetto CENOMA CUP n° B33C21000200005, is gratefully acknowledged.

Keywords: microalgae, cyanobacteria, omega-3, extractions, extract characterization

References

- Lakshimi, V. I., Kavitha, M., 2023. New Insights into Prospective Health Potential of ω -3 PUFAs. *Current Nutrition Reports*, 12, 813-829.
- Perez-Garcia, O., Bashan, Y., 2015. Microalgal heterotrophic and mixotrophic culturing for bio-refining: from metabolic routes to techno-economics. *Algal Biorefineries: Volume 2: Products and Refinery Design*, 61-131.

* Presenting author. E-mail address: elia.lio@unimi.it (Elia Lio)

S1.P48 Chemical and biological characterization of species of the genus *Weinmannia* (Cunoniaceae) endemic to Reunion Island for valorization purposes

Shamsia Pithon^{a,*}, Thomas Petit^{a,b}, Simon Remy^c, Jean-Hugues Renault^c, Yanis Caro^{a,b}

^a Natural Products Chemistry and Biotechnology Laboratory (CHEMBIOPRO), University of Reunion Island, 97410 Saint - Pierre, France

^b Health, Safety and Environment Department, University Institute of Technology, University of Reunion Island, 97410 Saint - Pierre, France

^c Reims Institute of Molecular Chemistry (UMR CNRS 7312), University of Reims Champagne- Ardenne

This study focuses on the chemical and biological characterization of species from the *Weinmannia* genus endemic to Reunion Island. These plants belong to the Cunoniaceae family and two species are present on this territory: *W. tinctoria* or “tan rouge” and *W. mauritiana* or “petit bois de tan” (“*Weinmannia* L. | Plants of the World Online” n.d.). The “tan rouge” is an emblematic plant of Reunion Island and was used traditionally in tannery and for dyeing natural fabrics for the red color of its bark (De Lanessan, 1886). The lack of knowledge on the chemical and biological characterization of endemic Reunionese species of the *Weinmannia* genus makes them plants with a high potential for innovation in the search for original active substances. As part of the chemical and biological characterization of *Weinmannia* species, a library of extracts has been created from different plant organs collected. Then, the purification of the natural substances present in the fractions of interest derived from the plant was carried out. Moreover, the chemical profiling of plant extracts and fractions following a dereplication workflow based on both NMR and LC-HRMS2 data (Hubert et al., 2017) has been achieved, and additionally, biological assays has been carried out in parallel. Finally, to characterize the tinting strength, the coordinates in CIEL*a*b international color system from the plant extracts has been measured. By improving the knowledge of the specialized metabolites produced by these plants, this project will contribute to developing and conserving the natural plant resources of the southwestern Indian Ocean.

Keywords: *Weinmannia* spp, Cunoniaceae, characterization, chemical profiling, bioactivities

References

- De Lanessan, J.-L. 1886. L'Expansion Coloniale de la France: étude économique, politique et Géographique sur les Etablissements Français d'Outre-Mer. In Félix Alcan.
- Hubert, J., Nuzillard, J.-M., and Renault, J.-H. 2017. Dereplication strategies in natural product research: How many tools and methodologies behind the same concept? *Phytochem. Rev.* **16**(1): 55–95.
- Weinmannia* L. | Plants of the World Online. (n.d.). Available from <http://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:328438-2> [accessed 9 February 2024].

* Presenting author. E-mail address: shamsia.pithon@univ-reunion.fr (Shamsia Pithon)

S1.P49 Enhancing caffeoylquinic acids production in *Limbarda crithmoides* (L.) Dumort. through yeast extract elicitation in liquid root cultures

Luísa Custódio^a, Zoltán Cziáky^b, Viana Castañeda-Loaiza^a, Maria João Rodrigues^{a,*}

^a Centre of Marine Sciences, University of Algarve, Faculty of Sciences and Technology, Ed. 7, Campus of Gambelas, 8005-139 Faro, Portugal

^b Agricultural and Molecular Research and Service Institute, University of Nyíregyháza, Nyíregyháza, Hungary

In this research, we developed a liquid culture of adventitious roots of *Limbarda crithmoides* (L.) Dumort., aiming at generating extracts overproducing hepatoprotective caffeoylquinic acids through yeast extract (YE) elicitation. The initiation of adventitious roots occurred in leaves grown *in vitro* on MS medium enhanced with 1 mg/L indole-3-butyric acid (IBA). The growth curve and synthesis of secondary metabolites (phenolics, flavonoids, and hydroxycinnamic acids) were monitored weekly to ascertain the ideal time for harvesting. Then, the roots were subjected to elicitation using different concentrations of YE – 50, 100, and 200 mg/L for four weeks. Ethanol extracts were assessed for their hepatoprotective capabilities against ethanol-induced damage and chemically analyzed via high- performance liquid chromatography coupled to electrospray ionization mass spectrometry (HPLC- ESI-MS/MS). Roots from greenhouse-grown plants served as a comparison. The addition of IBA led to the successful development of adventitious roots, peaking in biomass and phenolic content during the fourth week. Application of YE enhanced the concentration of hydroxycinnamic acids, especially chlorogenic acid and di-O-caffeoylquinic acid isomers. The extracts demonstrated marked liver- protective effects with 50 mg/L YE elicitation. Therefore, the use of yeast extract to stimulate liquid root cultures of *L. crithmoides* proves to be an effective method for enhancing the production of caffeoylquinic acids, providing scientific validation for the traditional medicinal claims of liver protection attributed to this plant species.

Acknowledgments: This research was funded by Portuguese national funds from the FCT - Foundation for Science and Technology, through projects UIDB/04326/2020, UIDP/04326/2020, and LA/P/0101/2020. VCL received a PhD grant from FCT (2020.04541.BD), MJR benefited from an FCT program contract (UIDP/04326/2020), and LC was supported by the FCT Scientific Employment Stimulus (CEECIND/00425/2017).

Keywords: golden samphire, Asteraceae, plant tissue culture, yeast extract

* Presenting author. E-mail address: mjrodrigues@ualg.pt (Maria João Rodrigues)

S1.P50 Daphnane diterpenoids from the fruits of *Daphne pseudomezereum*

Kouharu Otsuki*, Takashi Kikuchi, Wei Li

Faculty of Pharmaceutical Sciences, Toho University, Miyama 2-2-1, Funabashi, Chiba 274-8510, Japan

Daphnane diterpenoid possess a 5/7/6 (A/B/C) *trans*-fused tricyclic structure, which is exclusively distributed in plants of the Thymelaeaceae and Euphorbiaceae families (Otsuki et al. 2023). Daphnane diterpenoids have been demonstrated to exhibit a wide range of biological activities, such as antineoplastic, antiviral, analgesic, and pesticidal effects. *Daphne pseudomezereum* A. Gray (Thymelaeaceae) is a deciduous shrub mainly distributed in Japan, China, and Korea. The fruits of this plant are known to be poisonous and may cause diarrhea and vomiting if accidentally eaten. Previous chemical investigations of *D. pseudomezereum* have been conducted on the leaves and reported the isolation of coumarins, flavonoids, lignans and their glycosides. As part of our research for searching bioactive diterpenoids from the plants of Thymelaeaceae family, herein, we report the isolation and structural elucidation of daphnane diterpenoids from the fruits of *D. pseudomezereum*. A methanolic extract of the fruits of *D. pseudomezereum* was partitioned between EtOAc and H₂O, respectively. The EtOAc fraction was fractionated by Diaion HP-20 column chromatography and preparative HPLC, to afforded ten daphnane diterpenoids (**1–10**), including four new compounds (**1–4**). Their structures were determined by extensive spectroscopic analyses. The new daphnanes were structurally characterized by 1,2-dihydro structures for the A-ring or 4,6- and 4,7-ether structures for the B-ring. This study reveals the presence of daphnane diterpenoids in *D. pseudomezereum* for the first time, contributing to further understanding of bioactive diterpenoids in plants of the Thymelaeaceae family.

Keywords: *Daphne pseudomezereum*, Thymelaeaceae, diterpenoid, daphnane, phytochemical investigation

References

Otsuki, K., Li, W., 2023. Tiglane and daphnane diterpenoids from Thymelaeaceae family: chemistry, biological activity, and potential in drug discovery. J. Nat. Med. 77, 625-643.

* Presenting author. E-mail address: kouharu.otsuki@phar.toho-u.ac.jp (Kouharu Otsuki)

S1.P51 Nephroprotective *Ganoderma* sp. extracts towards cisplatin *in vitro*: chemical composition, metabolomics study and potential impact on cancer cells

Sébastien Sinaeve^{a,*}, Cedric Delporte^{a,b}, Cécile Husson^c, Marie-Hélène Antoine^c, Stéphane Welte^d, Joëlle Nortier^c, Caroline Stévigny^a

^a RD3-Pharmacognosy, Bioanalysis and Drug Discovery Unit, Faculty of Pharmacy, Université libre de Bruxelles, Brussels, Belgium

^b Analytical Platform of the Faculty of Pharmacy (APFP), Faculty of Pharmacy, Université libre de Bruxelles, Boulevard du Triomphe, 1050 Brussels, Belgium

^c Laboratory of Experimental Nephrology, Faculty of Medicine, Université libre de Bruxelles, Brussels, Belgium

^d Laboratoire des Sciences Végétales et Fongiques, UFR Pharmacie, Université Lille 2, Lille, France

This work aims to study the chemical composition, identify the bioactive compounds through metabolomics analyses and investigate the potential impact of nephroprotective extracts on lung cancer cells. For this, methanolic extracts of 5 mushrooms species from the *Ganoderma* P. Karst genus were first studied *in vitro* on Human Kidney (HK-2) cells as a 1 h pretreatment prior to cisplatin (20 μ M) exposure (Sinaeve et al., 2022). The chemical content was then analyzed by a phytochemical screening, the determination of total triterpenes and the phenolic compounds content, TLC-MS dereplication, metabolomics study (LC-MS), identification of potential bioactive compounds (LC- MS/MS) and estimation of their nephroprotective effect *in vitro*. Finally, the potential impact of cisplatin on lung cancer cells (A549) growth was established in the presence of the nephroprotective extracts and the bioactive compounds. Results have shown that methanolic extracts of *G. parvigibbosum* Welte & Courtecuisse and *G. tuberculosum* Murill. (10 μ g/mL), and their combination (5 + 5 μ g/mL) were significantly nephroprotective on HK-2 cells, mainly through an inhibition of the apoptosis and its signaling pathway. The phytochemical screening and chemical content study demonstrated a high content of triterpenes. The metabolomics study resulted in the detection of 9 discriminant compounds. Among them, 6-hydroxycoumarin was identified for the first time in these species, and exhibited, at 10 μ g/mL, a significant protective effect on HK-2 cells exposed to cisplatin. Finally, the nephroprotective extracts were tested on A549 cells and were found to potentiate cisplatin anti-cancer activity. The evaluation of 6-hydroxycoumarin on A549 cells is undergoing.

Keywords: cisplatin, metabolomic, nephroprotective, cancer, *Ganoderma*

Reference

Sinaeve, S.; Husson, C.; Antoine, M.-H.; Welte, S.; Stévigny, C.; Nortier, J. Nephroprotective Effects of Two *Ganoderma* Species Methanolic Extracts in an *In Vitro* Model of Cisplatin Induced Tubulotoxicity. *J Fungi* (Basel) 2022, 8, 1002.

* Presenting author. E-mail address: sebastien.sinaeve@ulb.be (Sébastien Sinaeve)

S1.P52 Phytochemical differentiation of wild thyme from thyme

Andressa Blainski^{a,*}, Cynthia Suarez^a, René Roth-Ehrang^a, Dimitris Michailidis^b, Aikaterini Argyropoulou^b

^a Finzelberg GmbH & Co. KG, Koblenzer Straße 48-56, 56626 Andernach, Germany

^b PharmaGnose S.A., 57th km Athens-Lamia National Road, 32011 Oinofyta, Greece

The *Thymus* L. is a large genus with about 350 species that are used as food and medicine. The most popular ones are Thyme (*Thymus vulgaris* L./*Thymus zygis* L.) and Wild Thyme (*Thymus serpyllum* L.) that are traditionally used against productive coughs mainly. In addition, recent studies with Wild Thyme extract have shown intestinal anti-inflammatory effects, metabolic improvements and modulation of the gut microbiota in rodents (Algieri et al., 2014; Ruiz-Malagón et al., 2022a, 2022b). A clinical study has confirmed the gastrointestinal benefits and the modulation of the Firmicutes/Bacteroidetes ratio in healthy volunteers (Knaub et al., 2022). To make sure that the intended species is used, distinguishing characteristics are of particular importance. In addition to pharmacopoeial morphological examination, our current study aims to differentiate between Wild Thyme and Thyme by phytochemical methods. The results identified characteristic HPTLC- fingerprints on flavonoids and on polar constituents of aqueous and methanolic extracts (Meier, 2011). The LC-HRMS identified more than 40 common substances, but one flavonoid (m/z 447.0933, C₂₁H₂₀O₁₁) was present only in Thyme, while another (m/z 463.0882, C₂₁H₂₀O₁₂) only in Wild Thyme. Salvianolic acid K was detected in both species but as different isomers. In addition, a flavonoid (m/z 299.0; C₁₆H₁₂O₆) was present only in the Wild Thyme aqueous extract. The different Thymol:Carvacrol ratio (16:1 for Thyme and 1:1.2 for Wild Thyme) also distinguishes the two species. These results corroborate that the evaluated *Thymus* species differ not only morphologically but chemically, and support the differentiation in laboratory routines.

Keywords: *Thymus serpyllum* L., *Thymus vulgaris* L., *Thymus zygis* L., phytochemical profile

References

- Algieri, F., Rodriguez-Nogales, A., Garrido-Mesa, N., Zorrilla, P., Burkard, N., Pischel, I., Sievers, H., Benedek, B., Feistel, B., Walbroel, B., Rodrigues-Cabezas, M.E., Galvez, J., 2014. Intestinal anti-inflammatory activity of the Serpylli herba extract in experimental models of rodent colitis. *J. Crohns Colitis* 8, 775-788.
- Knaub, K., Schon, C., Suarez, C.G., Pischel, I., 2022. Effects of a Food Supplement with a Wild Thyme (*Thymus serpyllum* L.) Extract on Gut Health and the Microbiome in Humans: A Randomized, Double-Blinded, Placebo- Controlled Clinical Trial. *Food and Nutrition Sciences* 13, 931-949.
- Meier, M., 2011. Einsatzmöglichkeiten der HPTLC-MS Kopplung bei der Untersuchung von Heilpflanzen. Diplomarbeit an der Naturwissenschaftlichen Fakultät der Karl-Franzens-Universität Graz.
- Ruiz-Malagón, A.J., Rodríguez-Sojo, M.J., Hidalgo-García, L., Molina-Tijeras, J.A., García, F., Pischel, I., Romero, M., Duarte, J., Diez-Echave, P., Rodríguez-Cabezas, M.E., Rodríguez-Nogales, A., Galvez, J., 2022a. The Antioxidant Activity of *Thymus serpyllum* Extract Protects against the Inflammatory State and Modulates Gut Dysbiosis in Diet- Induced Obesity in Mice. *Antioxidants* 11, 1073.
- Ruiz-Malagón, A.J., Rodríguez-Sánchez, M.J., Rodríguez-Sojo, M.J., Vezza, T., Pischel, I., Algieri, F., Rodríguez- Cabezas, M.E., Rodríguez-Nogales, A., Gálvez, J., 2022b. Intestinal anti-inflammatory and visceral analgesic effects of a Serpylli herba extract in an experimental model of irritable bowel syndrome in rats. *Front. Pharmacol.* 13:967644.

* Presenting author. E-mail address: andressa.blainski@finzelberg.com (Andressa Blainski)

S1.P53 Chemical ecology in marine fungi: diversity and dynamics of pyran-2-ones in a mussel- derived *Penicillium restrictum*

Van-Tuyen Le^a, Thibaut Robiou du Pont^a, Fabrice Fleury^b, Emmanuel Gentil^a, Gregory Genta-Jouve^c, Laurence Meslet-Cladière^d, Samuel Bertrand^a, Olivier Grovel^{a,*}

^a Nantes Université, ISOMER, UR 2160, F-44000 Nantes, France

^b Nantes Université, UFIP UMR CNRS 6286, 44322, Nantes, France

^c LEEISA, USR 3456, Université De Guyane, 97334 Cayenne, French Guiana

^d Univ Brest, INRAE, LUBEM, F-29280 Plouzané, France

Very little is known about chemical interactions between fungi and their marine mollusk hosts. Here, we investigated the metabolome of a *Penicillium restrictum* strain isolated from the blue mussel *Mytilus edulis* collected on the Loire estuary, France. Following the OSMAC approach, effects of salinity and of a mussel-derived medium on the metabolic expression of this strain were analyzed. An MS-based untargeted metabolomics study highlighted the high chemical diversity of pyran-2-ones that was found related to the presence of mussel lyophilizate in the culture medium. MS- and UV-guided purification allowed to isolate thirteen pyran-2-ones including five new compounds (Fig. 1), whose complete structure was elucidated using NMR, ECD and DP4 calculations (Le et al., 2021). A both untargeted and targeted time-scale metabolomics study has then been performed on the strain, which showed the dynamics of specialized metabolism with a variety of compounds produced at an early, intermediate or late stage of growth. One compound appearing at the very late stage was successfully isolated and characterized as (2*E*)-5-acetoxy-3-methoxy-2-pentenoic acid (Fig. 1), an unreported precursor of pyran-2-ones. Biosynthesis was further explored using a dynamic molecular networking approach, which highlighted the sequential biosynthetic steps leading to pyran-2-ones and the specific induction of some analogs when colonies reached confluence. These results illustrate the utility in using host-derived media for the discovery of new marine natural products and that combining metabolomics and dynamic molecular networking is a new approach to explore microbial biosynthesis, chemo-diversity and chemical induction.




Fig. 1. Structures of new fungal pyran-2-ones.

Keywords: *Penicillium restrictum*, pyran-2-ones, metabolomics, dynamic molecular networking, ecological cultures

References

Le V.T., Bertrand S., Robiou du Pont T., Fleury F., Caroff N., Bourgeade-Delmas S., Gentil E., Logé E., Genta-Jouve G., Grovel O., 2021. Untargeted metabolomics approach for the discovery of environment-related pyran-2-ones chemodiversity in a marine-sourced *Penicillium restrictum*. Mar. Drugs, 19, 378.

* Presenting author. E-mail address: olivier.grovel@univ-nantes.fr (Olivier Grovel)

S1.P54 Development and optimization of different eco-extraction and analytical methods on *Cannabis sativa* L.

Pichon Valentin^{a,*}, Pinault Emilie^{a,b}, Louvet François^c, Ducret Gilles^d, Viana Marylène^a, Mambu Lengo^a

^a Laboratoire LABCIS, UR 22722, Faculté de Pharmacie, 2 rue du Docteur Marcland, 87025 Limoges cedex, France

^b BISCEmplatform, US 42 Inserm -UAR 2015 CNRS, Univ. Limoges, CHU Limoges, 87025 Limoges, France

^c ENSIL-ENSCI Formation: Céramique Industrielle, ESTER Université de Limoges, 87068 Limoges, France

^d Centre Lab, Z.A. de la Granderaie, rue de Vernet, 23000 Guéret, France

The cannabinoids, predominantly found in the glandular trichomes of *Cannabis sativa* L., have garnered significant scientific interest due to their diverse pharmacological properties (Pacher et al., 2006). These include analgesic, anti-inflammatory and psychoactive effects (Russo et al., 2011). The access to these compounds depends on the extraction methods used, which can vary in their yield and metabolite profile for the same plant. This study was focused on the development and optimization of green extraction methods in the laboratory scale and their transferability to industry. A variety of green extraction techniques were employed (Tiago et al. 2022) and results were compared according to their mass yields, total contents (polyphenols, flavonoids, chlorophylls) and quantifications. Ethanol was used as the reference solvent. The use of ultrasounds, pressure, microwaves methods significantly improved mass yield and desired molecules. The HRLC-MS/MS allowed for the monitoring of cannabinoid concentrations, facilitating the selection of the most appropriate technique for the industrial scale. In addition to these results, a statistical approach was applied and made it easier to sort the data, to compare and obtain the best compromises for metabolites of pharmaceutical interest. The notion of desirability was added and highlighted the possibility to choose the condition which minimizing undesirable molecules (e.g. degradation products) while maximizing interesting molecules and enhance entourage effect (e.g. polyphenols, terpenes). Among the various options tested, ASE at 75°C emerged as the most effective process (65% of global desirability) providing a balance between mass yields, total contents and concentration of cannabinoids.

Keywords: *Cannabis sativa* L., eco-extractions, total contents, quantification, statistic description

References

- Russo, E.B. 2011, Taming THC: potential cannabis synergy and phytocannabinoid-terpenoid entourage effects. *British Journal of Pharmacology*, 163: 1344-1364.
- Pacher P., Bátkai S. and Kunos G., 2006, The endocannabinoid system as an emerging target of pharmacotherapy. *Pharmacological Reviews*, 58(3): 389-462.
- Tiago F. J., Paiva A., Matias A. A. and Duarte A.R.C., 2022, Extraction of Bioactive Compounds From *Cannabis sativa* L. Flowers and/or Leaves Using Deep Eutectic Solvents. *Frontiers in Nutrition*, 9, 892314.

* Presenting author. E-mail address: valentin.pichon@etu.unilim.fr (Valentin Pichon)

S1.P55 Impact of biomass hydration on the valorization of non-polar metabolites from *Spirulina* using eutectic solvents

Mervé Yagmur^{a,*}, Alia Chevalley^b, Bénédicte Montigny^c, Johan Jacquemin^d, Leslie Boudesocque-Delaye^a

^a EA7502 SIMBA, Faculty of Pharmacy, University of Tours, 31 avenue Monge, 37200 Tours, France

^b AquaEcoCulture, 7 rue d'Armor Maroué, 22400 Lamballe, France

^c EA 6299 PCM2E, University of Tours, Parc de Grandmont, 37200 Tours, France

^d MSN Department, Mohammed VI Polytechnic University, Lot 660, Hay Moulay Rachid, Ben Guerir 43150, Morocco

Arthrospira platensis (Spirulina), is a cyanobacteria that contains non-polar value-added metabolites for nutraceuticals and cosmetics, such as carotenoids or free fatty acids (Wils et al., 2021). This work focuses on the use of custom-made Natural Deep Eutectic Solvents (NaDES), which are alternative biobased solvents, to target non-polar metabolites of *Spirulina* (Wils et al., 2021). Biomass pre-treatment is crucial to ensure data reproducibility and stability before scaling up any process. We then investigated different biomass preparations: fresh or flaked biomass provided by our industrial partner. Initial extraction tests were performed using fresh biomass. Even though promising levels of target compounds were obtained, the extraction recovery was not reproducible in all experiments. This lack of consistency is linked to the instability of fresh biomass over time, as revealed by further analysis (Fig. 1).



Fig. 1. Spirulina preparations after cultivation and biomass stability.

To tackle this challenge, extraction experiments were conducted using rehydrated *Spirulina* flakes with 20% dry matter instead of fresh biomass. This change prevented technical issues related to structural changes in the fresh biomass over time, ensuring the collection of reproducible data (Fig. 1). To investigate the effects of rehydration on extraction performance, experiments were conducted to determine metabolite recovery as a function of water rates ranging from 0 to 80%. These analyses show that it is possible to modulate the profile of extracted compounds depending on the rehydration rate used. This opens up various options for optimization prior to industrial scaling up.

Keywords: Microalgae, *Spirulina* (*Arthrospira platensis*), green chemistry, NaDES, extraction

References

Wils, L., Leman-Loubière, C., Bellin, N., Clément-Larosièrre, B., Pinault, M., Chevalier, S., Enguehard-Gueiffier, C., Bodet, C., Boudesocque-Delaye, L., 2021. Natural deep eutectic solvent formulations for spirulina: Preparation, intensification, and skin impact. *Algal Res.* 56, 102317.

* Presenting author. E-mail address: merve.yagmur@univ-tours.fr (Mervé Yagmur)

S1.P56 Chemical investigation of the beetle *Agrilus cyanescens* (Buprestidae)

Isaiah I. Korostil, Jessica L. Pepe, Stephen T. Deyrup*

Department Chemistry and Biochemistry, Siena College, Loudonville, NY 12211, USA

There are over 400,000 described species of beetles with many still unknown, making them the most diverse group of macroorganisms on earth. Many beetles have been shown to biosynthesize potentially active molecules with the potential to act as new drug leads. *Agrilus cyanescens* is metallic wood-boring beetle (Buprestidae) native to Europe and Asia that is an invasive species in North America. To investigate the chemistry of adult *A. cyanescens*, we analyzed their chemical extracts via NMR spectroscopy and UPLC-HRMS. It was determined that *A. cyanescens* contains a mixture of buprestins (Fig. 1), defensive compounds known from other metallic wood-boring beetles. Interestingly, *A. cyanescens* contained all the known buprestins except for buprestin H, for which no evidence could be found. Studies of the defensive chemistry of invasive species are important to help understand the ecology of these organisms.




Fig. 1. Structures of buprestins A, B, and D-G.

Keywords: *Agrilus*, insect, defensive chemistry, buprestins, invasive species

* Presenting author. E-mail address: sdeyrup@siena.edu (Stephen T. Deyrup)

S1.P57 New lipopeptides from the marine bacteria *Pseudovibrio denitrificans*

Gyu Sung Lee^a, Chung Sub Kim^{a,b,*}

^a Department of Biopharmaceutical Convergence, Sungkyunkwan University, Suwon 16419, Korea

^b School of Pharmacy, Sungkyunkwan University, Suwon 16419, Korea

Microorganisms inhabiting distinctive and underexplored environments represent a rich source of bioactive compounds characterized by unique structures. Similarly, marine bacteria have garnered considerable interest due to their potential to yield unprecedented bioactive compounds. In this study, we isolated eight lipopeptides (**1–8**) from the seawater-derived bacterium *Pseudovibrio denitrificans* KCTC 62704. The chemical structures of the new compounds (**3–8**) were characterized by spectroscopic and spectrometric data interpretation, including NMR (¹H and ¹³C NMR, ¹H–¹H COSY, HSQC, HMBC, and NOESY) and HRESIMS analysis. Experimental ECD data analysis was conducted to assign the absolute configurations of the new compounds. We propose that these lipopeptides are synthesized from hexanoic acid, L-phenylalanine, and additional building blocks by stable-isotope feeding experiments. AntiSMASH analysis of the genome of *P. denitrificans* KCTC 62704 revealed the potential biosynthetic gene cluster for the isolated lipopeptides.

Acknowledgements: This work was supported by the National Research Foundation of Korea (NRF) grants funded by the Korean government (MSIT) (Nos. 2021R1C1C1011045, 2022R1A6A1A03054419), by the BK21 FOUR Project.

Keywords: marine bacteria, *Pseudovibrio denitrificans*, lipopeptide, biosynthesis

* Presenting author. E-mail address: chungsub.kim@skku.edu (Chung Sub Kim)

S1.P58 The Arabian Peninsula: A largely untapped resource for biotechnology

Ryan M. Young, Carsten Wolff

Biotechnology Research Center, Technology Innovation Institute. Abu Dhabi, United Arab Emirates

The Arabian Peninsula is a desert environment surrounded by three seas, namely the Arabian Gulf, Arabian Sea and Red Sea. The Arabian Gulf is a semi-enclosed shallow sea with mean depth of about 35 m. Because of the arid climate there, the Gulf's evaporation greatly exceeds freshwater input from precipitation and river runoff causing Gulf waters to be hypersaline. With vast expanses of desert dominating the landscape, temperatures can soar to extreme levels, often surpassing 50°C degrees Celsius during the peak of summer increase the sea temperatures from 22 to 35°C. The relentless sun beats down on the arid terrain, resulting in high levels of UV radiation. The combination of blistering heat, high salinity and relentless UV rays makes the Arabian Peninsula's climate one of the most unforgiving on the planet, shaping the lifestyle and survival strategies of many organisms. Extreme climates have long been recognized as catalysts for genetic mutations, driving adaptation and evolutionary change in organisms inhabiting such environments. The extremities of temperature, pressure, salinity, and radiation encountered in extreme climates can induce mutations in the genetic material of living organisms, allowing them to better survive and thrive in these hostile conditions. These mutations may confer novel traits that enhance an organism's ability to withstand the challenges posed by extreme environments, such as heat-resistant enzymes or mechanisms for preserving cellular integrity in high-salinity environments. Extremophiles, organisms that thrive in extreme conditions, provide compelling examples of the genetic adaptations that arise from exposure to extreme climates. Studying extremophiles and their genetic makeup holds promise for diverse applications, including biotechnology, medicine, and environmental remediation, as these organisms offer valuable insights into the mechanisms of adaptation and resilience in the face of extreme challenges. Herein, we discuss our strategy for collecting/isolating macro and microorganisms from these extreme marine and terrestrial environments. Additionally, we highlight our methodology to activate silent biosynthetic gene clusters in order to upregulate/activate the production of secondary metabolites as well as evaluate these compounds for their medical or agricultural potential. Compounds showing biotechnological potential are purified and elucidated through HRMS and 1D/2D NMR spectroscopy.

Keywords: extremophiles, Arabian Gulf, biotechnology, polyextremophiles

References

- Ibrahim, H.D., Xue, P. and Eltahir E.A.B. 2020. Multiple Salinity Equilibria and Resilience of Persian/Arabian Gulf Basin Salinity to Brine Discharge. *Front. Mar. Sci.* 7, 573
- Schultz J., Modolon F, Peixoto R.S. and Rosado A.S. 2023 Shedding light on the composition of extreme microbial dark matter: alternative approaches for culturing extremophiles. *Front. Microbiol.* 14:1167718
- Merino N., Aronson H.S., Bojanova D.P., Feyhl-Buska J., Wong M.L., Zhang S. and Giovannelli D. 2019 Living at the Extremes: Extremophiles and the Limits of Life in a Planetary Context. *Front. Microbiol.* 10, 780.

* Presenting author. E-mail address: ryan.young@tii.ae (Ryan M. Young)

S1.P59 Valorisation of north atlantic macroalgae: antimicrobial activity of phlorotannins

Adrian H. Kopf^{a,*}, Anna Hammer^a, René Dirks^b, Carla Araya-Cloutier^a

^a Laboratory of Food Chemistry, Wageningen University and Research, The Netherlands

^b Wageningen Food Safety Research, Wageningen University and Research, The Netherlands

Multidrug-resistant bacteria are a severe threat to society. Discovery of novel antimicrobials with distinctive mechanisms of action is pertinent. Marine environments, specifically, seaweeds (macroalgae) are excellent sources to look for new bioactive compounds (Silva et al., 2020). Due to the harsh environment that seaweeds grow in, they produce a variety of bioactives with unique chemical scaffolds (Pérez et al., 2016). In this work, fourteen species of seaweed harvested in the North Atlantic (North Sea) waters were screened for their antimicrobial potential. Crude extracts were obtained by ultrasound-assisted extraction using sequential solvents of varying polarity. Their antimicrobial activity was assessed in a particularly sensitive agar-well diffusion assay against bacteria *Bacillus cereus*, *B. pumilus*, *B. subtilis*, *Micrococcus luteus*, and *Yersinia ruckeri* (NAT- screen, Pikkemaat et al., 2008). At a concentration of 1 mg/mL, 57 out of the 85 extracts showed antimicrobial activity, with the largest zones of inhibition (~7 mm) detected in highly apolar (hexane) and polar (methanol) solvent extracts. The most potent antimicrobial activity was observed for brown seaweed *Ascophyllum nodosum*, *Fucus vesiculosus*, *Sargassum pallidum*, and *Undaria pinnatifida*. As a highlight, methanol extracts (*A. nodosum* and *F. vesiculosus*) were further investigated by activity-guided fractionation through purification with FLASH chromatography and analysis by RP- UHPLC-PDA-ESI-MS. Bioactive fractions were found to be enriched in phlorotannins (polymers of phloroglucinol units **1**, such as fucophlorethol A **2**, Fig. 1.) (Visser et al., 2017). We developed a highly sensitive pipeline for non-targeted screening and characterization of antimicrobial compounds in seaweeds, for the valorisation of this abundant marine resource.




Fig. 1. Structures of phloroglucinol **1**, and fucophlorethol A **2**.

Keywords: antimicrobials, macroalgae (Seaweeds), NAT-screening, bioactive polymers, phlorotannins

References

- Anne M. Vissers, Augusta Caligiani, Stefano Sforza, Jean-Paul Vincken, Harry Gruppen, 2017. Phlorotannin Composition of *Laminaria digitata*. *Phytochemical Analysis*, 28, 487-495.
- Aurora Silva, Sofia A. Silva, M. Carpena, P. Garcia-Oliveira, P. Gullón, M. Fátima Barroso, M.A. Prieto, J. Simal- Gandara, 2020. Macroalgae as a Source of Valuable Antimicrobial Compounds: Extraction and Applications. *Antibiotics*, 9, 642.
- María José Pérez , Elena Falqué, Herminia Domínguez, 2016. Antimicrobial Action of Compounds from Marine Seaweed. *Marine Drugs*, 14, 52.
- Mariël G. Pikkemaat, Sabrina Oostra-van Dijk, Jan Schouten, Michel Rapallini, Harry J. van Egmond, 2008. A new microbial screening method for the detection of antimicrobial residues in slaughter animals: The Nouws antibiotic test (NAT-screening). *Food Control*, 19, 781-789.

* Presenting author. E-mail address: adrian.kopf@wur.nl (Adrian H. Kopf)

S1.P60 Assessment of the antioxidant potential of leaf extracts from twelve *Sambucus nigra* varieties and their phytochemical characteristic

Sara Wójcik^a, Elżbieta Studzińska-Sroka^{a,*}, Magdalena Paczkowska-Walendowska^a, Piotr Szulc^b, Judyta Cielecka-Piontek^a

^a Department of Pharmacognosy and Biomaterials, Poznan University of Medical Sciences, 3 Rokietnicka Str., 60-806 Poznan, Poland

^b Department of Agronomy, Poznan University of Life Sciences, 11 Dojazd Str., 60-632 Poznan, Poland

Elderberry flowers and fruits are a medicinal raw material known in phytotherapy (Stępień et al., 2023); however, the leaves have not yet been used in medicine. The study aimed to compare the antioxidant activity and phytochemical screening of 70% methanolic extracts obtained from the leaves of twelve elderberry varieties collected from the experimental station of the Central Center for Research on Cultivated Plant Varieties in Słupia Wielka (Poland). In this study, the ability of the extracts to scavenge free radicals (DPPH, ABTS), their properties to reduce transition metal ions (CUPRAC, FRAP), and the ability to chelate Fe²⁺ ions were tested. The total content of polyphenols and flavonoids (Studzińska-Sroka et al., 2022, 2021) and the content of rutin, isoquercetin and neochlorogenic acid (Paczkowska-Walendowska et al., 2021) were examined. To statistically evaluate the relationship between properties, principal component analysis was performed. The obtained results showed that the extract from the Black Beauty variety had the strongest antioxidant potential, which was correlated with the highest content of total polyphenols and a high content of total flavonoids. The analysis of the main components showed that the antioxidant potential of all tested extracts is strongly correlated with the total polyphenol content, moderately dependent on the content of isoquercetin and neochlorogenic acid, and weakly dependent on the content of rutin. Moreover, the biplot graph shows that among the analysed varieties, Samyl, Obelisk, and Sabbo are characterized by the lowest antioxidant activity and low content of the tested active compounds.

Keywords: black elderberry leaves, PCA, phytochemical screening, biological activity *in vitro*

References

- Paczkowska-Walendowska, M., Gościński, A., Szymanowska, D., Szwajgier, D., Baranowska-Wójcik, E., Szulc, P., Dreczka, D., Simon, M., Cielecka-Piontek, J., 2021. Blackberry leaves as new functional food? Screening antioxidant, anti-inflammatory and microbiological activities in correlation with phytochemical analysis. *Antioxidants* 10, 1945.
- Stępień, A.E., Trojnik, J., Tabarkiewicz, J., 2023. Health-Promoting Properties: Anti-Inflammatory and Anticancer Properties of *Sambucus nigra* L. Flowers and Fruits. *Molecules* 28, 6235.
- Studzińska-Sroka, E., Galanty, A., Gościński, A., Wieczorek, M., Kłaput, M., Dudek-Makuch, M., Cielecka-Piontek, J., 2021. Herbal Infusions as a Valuable Functional Food. *Nutr.* 13, 4051.
- Studzińska-Sroka, E., Majchrzak-Celińska, A., Bańdurska, M., Rosiak, N., Szwajgier, D., Baranowska-Wójcik, E., Szymański, M., Gruszka, W., Cielecka-Piontek, J., 2022. Is Caperatic Acid the Only Compound Responsible for Activity of Lichen *Platismatia glauca* within the Nervous System? *Antioxidants* 11, 2069.

* Presenting author. E-mail address: elastudzinska@ump.edu.pl (Elżbieta Studzińska-Sroka)

S1.P61 Unlocking the potential of *Geissospermum* trees: exploring bioactive alkaloids for neurodegenerative diseases

Marcela Šafratová*, Lubomír Opletal, Jana Křoustková, Lucie Cahlíková

Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmacy in Hradec Králové, Charles University, Czech Republic

The genus *Geissospermum*, belonging to the Apocynaceae family, consists of Amazonian trees native to Brazil, primarily found in the eastern region. Indigenous tribes utilize aqueous and ethanolic extracts from the bark of these trees for various ailments such as malaria, cancer, and bacterial infections. *Geissospermum vellosii*, a species within this genus, is particularly notable for its abundance in indole and β -carboline alkaloids. Despite limited phytochemical research on this genus, isolation of alkaloids remains largely unexplored. The biological activity of extracts obtained from *G. vellosii* closely mirrors traditional decoction uses. A preliminary screening study assessing cholinesterase inhibition of the bark extract revealed significant activity against huBuChE (with an IC₅₀ value of 0.37 ± 0.05 μ g/mL). Furthermore, analysis via gas chromatography-mass spectrometry (GC/MS) and thin-layer chromatography (TLC) identified at least 12 alkaloids. The primary ethanolic extract was obtained from 40 kg of dried, crushed bark, and subsequent preparation of the alkaloidal extract involved different solvents (diethyl ether and chloroform) based on polarity. The purified diethyl ether extract (53 g) underwent separation via column chromatography, resulting in 16 fractions. Following purification and crystallization processes, five compounds have been isolated thus far. The inhibitory activity against recombinant human AChE and BuChE, as well as GSK-3 β , of these isolated alkaloids, was evaluated along with their ability to penetrate the blood-brain barrier.

Keywords: *Geissospermum vellosii*, Apocynaceae, cholinesterase activity, alkaloids

* Presenting author. E-mail address: safratom@faf.cuni.cz (Marcela Šafratová)

S1.P62 Bioguided-isolation of antimicrobial compounds against phytopathogens and human pathogens from lichen-associated bacteria

Bollaro Flavia^{a,b,*}, Derbaj Ghada^a, Rouaud Isabelle^a, Dreux-Zhiga Assia^b, Berthon Jean-Yves^b, Defois-Fraysse Clémence^b, Tranchimand Sylvain^a, Tomasi Sophie^a

^a Univ. Rennes, CNRS, ISCR – UMR 6226, 35043 Rennes, France

^b Greencell, Biopole Clermont Limagne, 1 Rue Henri Mondor, 63360 Saint-Beauzire, France

The World Health Organization places Antimicrobial Resistance (AMR) to be one of the greatest threats to global health (Murray *et al.*, 2022). The emergence of AMR in humans, animals and plants highlights the need for a "One Health" strategy (Despotovic *et al.*, 2023). This project goal is the investigation of the antimicrobial potential of a collection of 500 strains from the microflora of several lichen samples of *Rhizocarpon geographicum* against two phytopathogens: *Erwinia amylovora* (Piqué *et al.*, 2015), *Venturia inaequalis* (González-Domínguez, Armengol and Rossi, 2017) and against three human bacterial pathogens (*Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*). Lichen samples were collected from coastal and terrestrial areas, allowing them to undergo various biotic and abiotic stresses. Moreover, lichens, as holobionts, constitute micro- ecosystems that shelter unidentified microbial communities which could take part in the holobiont's defense against pathogens (Miral *et al.*, 2022). The first step of our study involved culturing 286 lichen-associated bacteria in a suitable medium, collecting the supernatant and pellet at an appropriate phase for the production of specialized metabolites and evaluating their activities against the targeted pathogens. Initial results performed on 150 freeze-dried supernatants highlighted the inhibitory activities of 10 supernatants against *E. amylovora*, 2 against *V. inaequalis*, 4 against *P. aeruginosa* and 1 against *E. coli*. Four of the most active strains were selected for production optimisation and large-scale cultivation leading to bioguided fractionation. The results of this fractionation to isolate bioactive compounds will be discussed.

Keywords: antimicrobial resistance, biocontrol, lichen microflora, bioguided fractionation, *Rhizocarpon geographicum*

References

- Despotovic, M. et al. 2023. 'Reservoirs of antimicrobial resistance in the context of One Health', *Current Opinion in Microbiology*, 73, p. 102291.
- González-Domínguez, E., Armengol, J. and Rossi, V. 2017. 'Biology and Epidemiology of *Venturia* Species Affecting Fruit Crops: A Review', *Frontiers in Plant Science*, 8, p. 1496.
- Miral, A. et al. 2022. 'Rhizocarpon geographicum Lichen Discloses a Highly Diversified Microbiota Carrying Antibiotic Resistance and Persistent Organic Pollutant Tolerance', *Microorganisms*, 10, 1859.
- Murray, C.J.L. et al. 2022. 'Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis', *The Lancet*, 399, 629–655.
- Piqué, N. et al. (2015) 'Virulence Factors of *Erwinia amylovora*: A Review', *International Journal of Molecular Sciences*, 16, 12836–12854.

* Presenting author. E-mail address: flavia.bollaro@univ-rennes.fr (Flavia Bollaro)

S1.P63 Cardiometabolic effects on obese mice of the Mediterranean seagrass *Cymodocea nodosa* hydroalcoholic extract

Maria Vitiello^{a,*}, Marinella De Leo^{a,b,c,d}, Alessandra Braca^{a,b,c,d}, Lara Testai^{a,b,c,d}, Paola Nieri^{a,b,c,d}, Giada Benedetti^a, Lorenzo Flori^a, Virginia Menicagli^e, Claudio Lardicci^{b,c,d,f}, Elena Balestri^{c,e}

^a Department of Pharmacy, University of Pisa, Via Bonanno 33, 56126 Pisa, Italy

^b Interdepartmental Research Center of Marine Pharmacology “MarinePharma”, University of Pisa, Via Bonanno 6, 56126 Pisa, Italy

^c Interdepartmental Research Center Nutrafood “Nutraceuticals and Food for Health”, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

^d CISUP, Centre for Instrumentation Sharing, University of Pisa, Lungarno Pacinotti 43/44, 56126 Pisa, Italy

^e Department of Biology, University of Pisa, Via Derna 1, 56126 Pisa, Italy

^f Department of Earth Sciences, University of Pisa, Via S. Maria 53, 56126 Pisa, Italy

Cymodocea nodosa (Ucria) Asch. (Cymodoceaceae) is a dioecious medium-size, fast-growing seagrass colonizing coastal waters of the Mediterranean and eastern Atlantic Ocean (Garrido et al., 2013). Due to its high plasticity, it can colonize different substrates and environments, tolerating salinity fluctuations and climate changes (Chefaoui et al., 2015). Flavonoids, steroids, polyamines, diarylheptanoids, and sulfated polysaccharides with interesting potential biological activity (Ben Abdallah Kolsi et al., 2017) have been previously extracted from its plant tissue (Smadi et al., 2017). This study aims to investigate the phytochemical profile of a leaf hydroalcoholic extract of *C. nodosa* collected in the Ligurian Sea (Italy) and its activity in a murine model of cardiometabolic disorder. UHPLC coupled to an Orbitrap-based HR-MS led to the characterization of phenolic bioactive substances including flavonoid glycosides (quercetin and isorhamnetin derivatives) and chicoric acid as the most represented compound (1.58 ± 0.3 mg/g fresh weight). Mice fed with an obesogenic diet were supplemented with the extract for 9 weeks and at the end of the experimental protocol a containment of body weight gain (increase of 32 ± 2 vs $49.6 \pm 8.7\%$) and an improvement of glycaemic profile, compared to animals supplemented with placebo, were observed. Glucose levels were 115 ± 4 vs 138 ± 10 mg/dl and HbA1c levels were 32 ± 0.4 vs 29.8 ± 0.8 mmol/mol. These results suggest that the extract obtained has beneficial effects in the prevention of metabolic disorders associated with obesogenic diet. Further experiments will be carried out to explore the mechanisms involved in this beneficial profile.

Keywords: *Cymodocea nodosa*, phenols, chicoric acid, LC-HR-ESI/MS, cardiometabolic disorder

References

- Ben Abdallah Kolsi, R., Ben Salah, H., Jardak, N., Chaaben, R., El Feki, A., Rebai, T., Jamoussi, K., Allouche, N., Belghith, H., Belghith, K., 2017. Effects of *Cymodocea nodosa* extract on metabolic disorders and oxidative stress in alloxan-diabetic rats. *Biomed. Pharmacother.* 89, 257–267.
- Chefaoui, R.M., Assis, J., Duarte, C.M., Serrão, E.A., 2015. Large-scale prediction of seagrass distribution integrating landscape metrics and environmental factors: The case of *Cymodocea nodosa* (Mediterranean–Atlantic). *Estuar. Coast* 39, 123–137.
- Garrido, M., Lafabrie, C., Torre, F., Fernandez, C., Pasqualini, V., 2013. Resilience and stability of *Cymodocea nodosa* seagrass meadows over the last four decades in a Mediterranean lagoon. *Estuar. Coast. Shelf Sci.* 130, 89–98.

* Presenting author. E-mail address: maria.vitiello@phd.unipi.it (Maria Vitiello)

S1.P64 Obtaining natural and sustainable blue dyes from the fungus *Terana coerulea* (Lam.) for textile application – The TEXTBLUESTYLE project

Isabel P. Fernandes^{a,*}, Miguel Gonçalves^a, Hugo Bogalho^b, Yara Goulart^{c,d}, Sandrina A. Heleno^{c,d}, Ricardo Calhelha^{c,d}

^a Tree Flowers Solutions, Lda. Edifício Brigantia Ecopark, 5300-358, Bragança, Portugal

^b Bioreishi, Quinta da Fonte Velha, nº 936, 2440 234 Golpilheira, Batalha, Portugal

^c Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia 5300-253 Bragança, Portugal. ^dLaboratório para a Sustentabilidade e Tecnologia em Regiões de Montanha, Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

Amongst the several categories of synthetic dyes, the azo is one of high concern as once under specific conditions, these dyes decompose generating aromatic amines classified as carcinogenic. This toxicity has led to legal impositions at European level, resulting in application restrictions in several industrial sectors (Keshava et al., 2023). Among the possible natural dyes' sources available, higher fungi stand out due to their ability to synthesize bioactive molecules and pigments, like the cobalt blue pigment generated by the fungus *T. coerulea*. Hence, the TextBlueStyle project aims at obtaining blue, stable, and natural dyes by applying green extraction techniques and using water as solvent, following the work performed within our research group (Maisterra et al., 2017). The obtained extracts are being characterized regarding antioxidant (DPPH, FRAP and reducing power methods), antimicrobial activity and toxicity. Previous results obtained pointed out that cytotoxicity of *T. coerulea* aqueous extract against non-tumor porcine liver cells primary culture, occurred at concentrations higher than 400 µg/ml, the maximum concentration tested (Maisterra et al., 2017), suggesting the low toxic character of the obtained extract. This work is still ongoing, being expected to achieve extracts with different shades of blue as well as antioxidant activity suitable for textile functionalization. This would allow to consider further development with the production of technical textile products, in a sustainable and eco-innovation context.

Acknowledgements: To Foundation for Science and Technology (FCT, Portugal) for financial support to CIMO (UIDB/00690/2020 and UIDP/00690/2020) and SusTEC (LA/P/0007/2021); national funding by FCT, P.I., through the institutional scientific employment program-contract for S.A. Heleno and R.C. Calhelha. To Promove Pilotos Inovadores Fundação „La Caixa” for the financial support of TEXTBLUESTYLE (PL23-00065) project.

Keywords: *Terana coerulea*, antioxidant extract, natural blue dye, textile application

References

- Keshava, C., Nicolai, S., Vulmir, S.V., Cruz, F.A., Ghoreishi, N., Knueppel, S., Lenzner, A., Tarnow, O., Vanselow, J.T., Schulz, B., Williams, A.J., Pirow, R., 2023. Application of systematic evidence mapping to identify available data on the potential human health hazards of selected market-relevant azo dyes. *Environ. Int.*, 176, 107952.
- Maisterra, M., Castro, M.A., Centeno, L.M., Calhelha, R.C., Ferreira, I.C.F.R., Garcia, P.A., 2017. Cytotoxic terphenyl neolignans from fungus *Terana coerulea*: new natural corticins D and E, and revised structure for corticin A. *Nat. Prod. Com.*, 12(5), 695-698.

* Presenting author. E-mail address: isabelfernandes@treeflowerssolutions.com (Isabel P. Fernandes)

S1.P65 Using low levels of antimicrobials and antiseptics as elicitors for environmental isolates to activate or re-activate antimicrobial producing biosynthetic gene clusters

Amy McLeman*, Emma Murphy, Adam P. Roberts

Liverpool School of Tropical Medicine, Department of Tropical Disease Biology, Liverpool Life Sciences Accelerator, Liverpool, L7 8XZ, UK

This presentation will describe the results of screening >2000 environmental isolates against six antimicrobials at low levels to determine elicitation of antimicrobial activity. The overall aim of this study being the recovery of antimicrobial activity that has been lost from environmental bacteria following an initial agar plate based antimicrobial production assay. We compare the antimicrobial activity of >2000 environmental isolates against *Escherichia coli*, *Micrococcus luteus*, MRSA and *Candida albicans* with and without one of six elicitors: tetracycline (0.1 µg/ml), ampicillin (0.1 µg/ml), clindamycin (0.5 µg/ml), ciprofloxacin (0.01 µg/ml), CTAB (0.08 µg/ml) and Triclosan (0.003 µg/ml). These experiments showed a 25% success rate in eliciting antimicrobial production from the environmental isolates. As part of our wider antimicrobial discovery activity, we have




Fig. 1. The effect cell free supernatant from isolate X when elicited by triclosan (blue) and when not elicited (red) on the growth of susceptible *E. coli*. The effect of the triclosan concentration on *E. coli* growth is shown (green) and normal *E. coli* growth (purple).

assembled a large library of environmental microbes (currently >70,000), which we are screening for antimicrobial activity against the WHO priority pathogens *E. coli*, *C. albicans* and *Candida auris* (WHO 2017, WHO 2022). We found that approximately 40% of active isolates lose activity on secondary agar plate-based assays so aimed to re-activate antimicrobial production using the elicitors described above. We found a 23% elicitation on agar assays using tetracycline, CTAB and triclosan; however, this reduced to only 0.7% (1 isolate) elicitation in broth (Fig. 1). Sequencing and subsequent bioinformatics on the whole genome of this isolate showed likely biosynthetic gene clusters and the potential antimicrobials being produced.

Keywords: elicitation, antimicrobial production, environmental isolates, microbial

References

- WHO, 2017. Prioritization of pathogens to guide discovery, research and development of new antibiotics for drug-resistant bacterial infections, including tuberculosis. <https://iris.who.int/bitstream/handle/10665/311820/WHO-EMP-IAU-2017.12-eng.pdf?sequence=1>
- WHO, 2022. WHO fungal priority pathogens list to guide research, development and public health action. <https://iris.who.int/bitstream/handle/10665/363682/9789240060241-eng.pdf?sequence=1>

* Presenting author. E-mail address: amy.mcleman@lstm.ac.uk (Dr. Amy McLeman)

S1.P66 Biotransformation of the secondary metabolites from *Picralima nitida* and their antibacterial activities

F.D. Mabou^{a,b}, I. Rouaud^b, S. Ferron^b, A. Sauvager^b, L. Bousarghin^c, D. Ngnokam^a, S. Tomasi^b

^a Univ. Dschang, Department of Chemistry, Dschang, Cameroun

^b Univ. Rennes, CNRS, ISCR – UMR 6226, 35043 Rennes, France

^c INSERM, Univ. Rennes, INRAE, CHU Rennes, Nutrition Metabolisms and Cancer (NuMeCan), UMR 1317, Biosit, MRic/ISFR, 35000 Rennes, France

This presentation will focus on the antibacterial activities of secondary metabolites from *Picralima nitida* (Apocynaceae) and their biotransformed derivatives. *Picralima nitida* Durand and Hook is a medicinal plant from the genus *Picralima* and family Apocynaceae. It is found in tropical African countries such as Ivory Coast, Nigeria, Uganda, and Gabon (Erharuyi et al., 2014; Olajide et al., 2014). This plant is used in traditional medicine for the treatment and management of malaria, abscesses, hepatitis, pneumonia, diabetes, and hypertension (Erharuyi et al., 2014; Teugwa et al., 2013). Several studies have previously shown that various extracts of this plant are good sources of phytochemicals such as glycosides, alkaloids, triterpenes, flavonoids, polyphenols, saponins, and tannins (Akinwunmi et al., 2019; Creed et al., 2020; Erharuyi et al., 2014). As such, in our continuous effort to discover new bioactive compounds, the fruits of *Picralima nitida* were collected in Cameroon. The air-dried powder was extracted with MeOH. The separation and isolation of compounds from the obtained extract were performed using classical chromatography methods (column chromatography, flash-chromatography) while the structures of the isolates were elucidated by interpretation of their spectroscopic data and by comparison with literature. Biotransformation was performed on the major compounds using some bacteria belonging to the *Bacillus*, *Nocardia* and *Streptomyces* genera. The antibacterial activity of natural compounds and their biotransformed derivatives was carried out using the broth microdilution method against human bacterial pathogens.

Keywords: *Picralima nitida*, Apocynaceae, biotransformation, antibacterial activities

References

- Akinwunmi, K.F., Amadi, C.V., 2019. Assessment of antioxidant and antidiabetic properties of *Picralima nitida* seed extracts. *Journal of Medicinal Plants Research* 13, 9-17.
- Creed, S.M., Gutridge, A.M., Argade, M.D., Hennessy, M.R., Friesen, J.B., Pauli, G.F., van Rijn, R.M., Riley, A.P., 2020. Isolation and Pharmacological Characterization of Six Opioidergic *Picralima nitida* Alkaloids. *Journal of Natural Products*.
- Erharuyi, O., Falodun, A., Langer, P., 2014. Medicinal uses, phytochemistry and pharmacology of *Picralima nitida* (Apocynaceae) in tropical diseases: a review. *Asian Pacific Journal of Tropical Medicine* 7, 1-8.
- Erharuyi, O., Falodun, A., Langer, P., 2014. Medicinal uses, phytochemistry and pharmacology of *Picralima nitida* (Apocynaceae) in tropical diseases: A review. *Asian Pacific Journal of Tropical Medicine* 1-8.
- Olajide, O.A., Velagapudi, R., Okorji, U.P., Sarker, S.D., Fiebich, B.L., 2014. *Picralima nitida* seeds suppress PGE2 production by interfering with multiple signalling pathways in IL-1 β -stimulated SK-N-SH neuronal cells. *Journal of Ethnopharmacology* 152, 377-383.
- Teugwa, C.M., Mejiato, P.C., Zofou, D., Tchinda, B.T., Boyom, F.F., 2013. Antioxidant and antidiabetic profiles of two African medicinal plants: *Picralima nitida* (Apocynaceae) and *Sonchus oleraceus* (Asteraceae) *BMC Complementary and Alternative Medicine* 13, 175.

* Presenting author. E-mail address: [Florence Mabou \(florence.mabou@univ-rennes.fr\)](mailto:florence.mabou@univ-rennes.fr)

S1.P67 Natural products isolation from subterranean fungi of Wind Cave National Park (US)

Kirk P. Manfredi*, Justin Peters, Joshua Sebree, Elizabeth Guevara

Department of Chemistry and Biochemistry, University of Northern Iowa, Cedar Falls, IA 50614 USA

Since 2019 undergraduate students and faculty from the University of Northern Iowa have been active in a NASA sponsored research program to explore Wind Cave National Park (South Dakota, USA) as a terrestrial analog for a possible exobiological environment off earth. The subsurface lakes within Wind Cave contain fewer microbial cells (5×10^3 cells/mL) than nearly all other bodies of water on earth. It's thought that these isolated, nutrient-limited lakes may be an analog for such oceans on the moons of Saturn. One of the goals of this multi disciplined research project is to develop a genetic map of the Wind Cave microbial system. To accomplish this goal, microbes have been collected from the air, aqueous, and terrestrial environments from many areas of the cave not accessible to the general public. Though not part of the original scope of this project, our lab viewed these subterranean microbes as possible sources of novel, biologically active natural products. To date, we have purified and identified over 90 fungal isolates from the cave. This presentation will summarize the species diversity that has been isolated from the cave to date. Additionally, the antimicrobial activity of isolated fungal cultures and the structural characterization of compounds will be presented.

Keywords: antimicrobial activity, fungal isolate, cave microbes

* Presenting author. E-mail address: manfredi@uni.edu (Kirk P. Manfredi)

S1.P68 Chemical and biological study of *Glutamicibacter* sp. NCA-315 isolated from the sinkhole Tza- Itzá in the Yucatan Peninsula

Carlos D. Hernández-Granillo^a, Oswaldo P. Martínez-Rodríguez^a, Alejandra Prieto-Davó^b, Rodolfo García-Contreras^c, William Fenical^d, Mario Figueroa^{a,*}

^a Departamento de Farmacia, Facultad de Química, Universidad Nacional Autónoma de México, Coyoacán, CDMX, 04510, México

^b Unidad de Química SISAL, Facultad de Química, Universidad Nacional Autónoma de México, Sisal, Yucatán, 97356, México

^c Departamento de Microbiología y Parasitología, Facultad de Medicina, Universidad Nacional Autónoma de México, Coyoacán, CDMX, 04510, México

^d Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, 92093, USA

The Yucatan Peninsula in Mexico is an enormous reservoir of unique microbes for bioprospecting purposes (Suárez-Moo et al., 2022). In this work, the chemical and biological potential of a bacterium identified as *Glutamicibacter* sp. NCA-315, isolated from sediment of the sinkhole Tza-Itzá in the Anillo de Cenotes (Ring of Sinkholes) Reserve in the Yucatan Peninsula, was explored. The EtOAc extract of the supernatant of the strain cultivated in DSC-ASW media showed $\geq 50\%$ inhibition against *E. faecalis*, MSSA, and MRSA. The chemical study of the extract yielded the bile acids cholic acid (**1**), 7-oxo-3,12-dihydroxycholic acid (**2**), glycocholic acid (**3**), and glycodeoxycholic acid (**4**) (Fig. 1). Compounds **3** and **4** inhibited 40% and 60% of the biofilm formation of MRSA, respectively, at 64 $\mu\text{g/mL}$ (Martínez-Rodríguez et al., 2023). Finally, WGS was completed for the bacterium and BGC analysis was performed.

Acknowledgments: DGAPA-PAPIIT IN203923, DGAPA-PASPA; CONAHCyT (Estancia Sabática 2023); Fulbright- García Robles.




Fig. 1. *Glutamicibacter* sp. NCA-315 and isolated bile acids.

Keywords: *Glutamicibacter* sp., Tza-Itzá sinkhole, bile acids, MRSA biofilm inhibition

References

- Suárez-Moo, P., Remes-Rodríguez, C.A., Márquez-Velázquez, N.A., Falcón, L.I., García-Maldonado, J.Q., Prieto-Davó, A. 2022. Changes in the sediment microbial community structure of coastal and inland sinkholes of a karst ecosystem from the Yucatan Peninsula. *Sci. Rep.* 12, 1110.
- Martínez-Rodríguez, O.P., García-Contreras, R., Aguayo-Ortiz, R., Figueroa M. Antimicrobial and antibiofilm activity of fungal metabolites on methicillin-resistant *Staphylococcus aureus* (ATCC 43300) mediated by SarA and AgrA. *Biofouling*, 39, 830-837.

* Presenting author. E-mail address: mafiguer@unam.mx (Mario Figueroa)

S1.P69 Comparative analysis of chemical variations across plant parts, seasons, and species in three *Osmanthus* plants

Kyungha Lee^{a,*}, Mina Lee^b, Kyo Bin Kang^a

^a College of Pharmacy, Sookmyung Women's University, Seoul 04310, Korea

^b College of Pharmacy, Sunchon National University, Suncheon 57922, Korea

The genus *Osmanthus* is distributed mainly in East Asia, and is famous for *Osmanthus fragrans* (OF) having a strong scent. Due to its value as an ingredient of perfume, phytochemical and pharmacological studies on this genus have been focused on OF. Due to the lack of other species in this genus, we performed a comparative study on the chemical composition of three plants, OF, *Osmanthus fragrans* var. *aurantiacus* (OFA), and *Osmanthus* × *fortunei* (OxF). To this end, LC- MS/MS analysis of samples by parts (twig, leaf, and flower) and months (May, June, July, September, and October) of the three species was conducted. The differences between the parts were evident. The flavonoid class predominated in the leaves, while the lignan and phenylethanoid classes were dominant in the twig samples. Furthermore, the chemical dissimilarity within the same parts was observed. In the leaves, there was a tendency for OxF to exhibit a higher prevalence of flavonoid glycosides with two sugars rather than aglycones. For the twigs, OFA showed a different trend from the other species; lignans were notably shown in leaves of OFA, whereas these compounds were mainly detected in twigs of the other two species. Moreover, the phenolic compounds were prominently in OFA compared to the others. We assessed the ABTS and DPPH radical scavenging activity to know the relationship between chemical composition and physiological activity. Antioxidant capacity decreased until July and then increased after September. Consistent with this trend, the phenylethanoid class compounds mainly detected in the twigs of OFA exhibited high bioactivity scores. Therefore, it is considered most effective to collect OFA twigs after September to maximize the antioxidant activity of *Osmanthus* spp.

Keywords: *Osmanthus* spp., LC-MS/MS, metabolomics, antioxidant activity

* Presenting author. E-mail address: khlee21@sookmyung.ac.kr (Kyungha Lee)

S1.P70 Siderophores-mediated antagonism of specialized metabolites in fungal-bacterial interactions

Huong T. Pham^{a,*}, Young Woon Lim^b, Kyo Bin Kang^a

^a College of Pharmacy, Sookmyung Women's University, Seoul 04310, Korea

^b School of Biological Sciences, Seoul National University, Seoul 08826, Korea

Siderophores are specialized microbial metabolites that chelate iron and play multiple roles in the survival strategies of microorganisms. In a coculture model of 86 *Penicillium* species and *Bacillus subtilis*, we find that siderophores have an additional function of releasing toxic metabolites from partner organisms during competitive interactions. In expectation of the cocultivation-induced new metabolites to have antibacterial activity, the LC-MS/MS were analyzed to observe the changes in metabolomes. However, surprisingly, various secondary metabolites, including verruculogen and its analogs, were reduced by coculture while bacillibactin, a siderophore produced by *B. subtilis*, was significantly induced. Through further assays, we hypothesized that an iron-deficient condition modulated by bacillibactin was caused leading to a reduction in the amount of metabolite production of *Penicillium*. As these reduced compounds were antimicrobial, we speculated that the siderophore could contribute to the coexistence as a cross-protector by observing the phenotypes of the designed microbial consortia. In summary, our results propose that through iron-limitation mechanisms, bacillibactin inhibits the antimicrobial compounds and thereby indirectly protects other members within microbial communities.

Keywords: bacillibactin, siderophore, *Penicillium*, *Bacillus subtilis*

* Presenting author. E-mail address: phamthihuong@sookmyung.ac.kr (Pham Thi Huong)

S1.P71 Absolute configuration of unisolated epimers of pseurotin D from the marine-derived fungus *Penicillium janczewskii*

Yu Kyung Choi^{a,*}, Gyu Sung Lee^b, Yeongseo Kim^a, Men Thi Ngo^a, Jin Wook Cha^d, Chung Sub Kim^{b, c}, Kyo Bin Kang^a

^a College of Pharmacy and Research Institute of Pharmaceutical Sciences, Sookmyung Women's University, Seoul 04310, Republic of Korea

^b Department of Biopharmaceutical Convergence, Sungkyunkwan University, Suwon 16419, Republic of Korea

^c School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea

^d Natural Product Informatics Research Center, KIST Gangneung Institute of Natural Products, Gangneung 25451, Republic of Korea

Halogen supplements affect microbial secondary metabolism in both direct and indirect ways. During our comparative analysis on metabolic change of *Penicillium janczewskii* by different halogen supplements, we found that the addition of iodide resulted in a significant increase in pseurotin D (**1**). We deduced that iodide promoted activation of the carbon-carbon double bond in the last non- enzymatic SN2' reaction step of pseurotin D biosynthesis. The absolute configuration of **1** was deduced from the biosynthesis pathway, without spectroscopic evidence (Tsunematsu et al., 2014). Consequently, we isolated from *P. janczewskii* and determined the absolute configuration of **1** using advanced analytical technologies. Each chiral center except for C-13 was determined to be 5*S*, 8*S*, 9*R*, and 10*R* through NOESY, CD spectra, and the modified Mosher's method. Our analysis revealed that **1** exists as a mixture of diastereomers. Here, we utilized a gradient-enhanced multiplet-selective targeted-observation NMR experiment (GEMSTONE) and ultraselective heteronuclear polarization transfer (UHPT) to determine the NMR chemical shift values of each epimer (Cha et al., 2023) and applied CP3 to establish the absolute configuration.




Fig. 1. Structure of pseurotin D (**1**).

Keywords: *Penicillium janczewskii*, pseurotin D, halogen, diastereomer, GEMSTONE

References

- Tsunematsu, Y., Fukutomi, M., Saruwatari, T., Noguchi, H., Hotta, K., Tang, Y., Watanabe, K., 2014. Elucidation of pseurotin biosynthetic pathway points to trans-acting C-methyltransferase: Generation of chemical diversity. *Angew Chem*, 126, 8615-8619.
- Cha, J. W., Kim, M. S., Park, J. S., 2023. A Single-Scan Ultraselective Heteronuclear Polarization Transfer Method for Unambiguous Complex Structure Assignment. *Angew Chem*, 135, e202304196.

* Presenting author. E-mail address: ykchoi@sookmyung.ac.kr (Yu Kyung Choi)

S1.P72 A solution from the past: exploring the untapped potential of museum specimens to preserve secondary metabolites

Sam Afoullouss^a, Jennifer Netjes^a, Matthew Simpson^a, Micah Harris^a, Laurent Calcul^c, Dale Chaput^b, William Keel^d, Bill J. Baker^a

^a University of South Florida, Department of Chemistry, Tampa, FL, USA

^b University of South Florida, Proteomics and Mass Spectrometry Core Facility, Tampa, FL, USA

^c University of South Florida, Chemical Purification Analysis and Screening core facility (CPAS), Tampa, FL, USA

^d National Museum of Natural History, Smithsonian Institution, USA

A fundamental pillar of natural products focused drug discovery is that biodiversity results in chemical diversity. To this end, marine natural products researchers have relied on sample collection trips to remote locations, searching for rare and unstudied species. The high cost of sampling trips, increased restriction of sampling permits and ecosystem degradation has resulted in multiple hurdles to acquiring new samples. One solution to these issues comes to us from the past, in the form of museum marine invertebrate collections. For hundreds of years, specimens from around the world, covering a diverse range of species, have been consolidated in museums, primarily stored in 90% ethanol. This preserving solution may be an untapped source of new natural products, from specimens with rich scientific history including varying collection depths, dates and location, high-level taxonomical assignment, and genetic studies. In collaboration with the Smithsonian Museum, this study explores the ability of this ethanol solution to preserve secondary metabolites over time from two distinct taxa: the tunicate *Synoicum* and the glass sponge *Aphrocallistes*, collected between 1945- 2016. Ethanol preserving solution was concentrated and analyzed using untargeted LC-tandem MS/MS (orbitrap). LC-MS² data was processed through MZmine3's targeted feature detection workflow, identifying palmerolide A (**1**) and aphrocallistin B (**2**), in samples of *Synoicum* and *Aphrocallistes*, respectively. Multiple reaction monitoring (MRM) with a LC-MS QqQ was used to confirm and quantify the presence of **1** and **2**, using standards from our in-house library of marine metabolites. Patterns in chemical composition of each sample was explored by untargeted LC-IMS- QToF combined with GNPS and SIRIUS spectral libraries.

Keywords: drug discovery, mass spectrometry, metabolomics, marine natural products

* Presenting author. E-mail address: samafoullouss@usf.edu (Sam Afoullouss)

S1.P73 Discovery of a brominated Ahp-containing cyclic depsipeptide from the marine cyanobacterium *Moorena* sp. collected in Panama

Byeol Ryu^{a,*}, Nicole E. Avalon^a, Lena Gerwick^a, William H. Gerwick^{a,b}

^a Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California San Diego, 9500 Gilman Drive, La Jolla, California 92093, USA

^b Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California San Diego, 9500 Gilman Drive, La Jolla, California 92093, USA

Bromine-containing natural products are recognized for their potential in drug discovery and as toxins due to their notable bioactivity. Marine organisms represent abundant reservoirs of structurally diverse halogenated natural products. With the goal of identifying novel brominated secondary metabolites from marine cyanobacteria, we surveyed several extracts using mass spectrometry and the query language (MassQL). Consequently, a brominated Ahp-containing cyclic depsipeptide was successfully isolated from *Moorena* sp. collected from the Las Perlas islands in Panama. In parallel, derivatives of this compound were identified through MS/MS molecular networking. Their structures were elucidated by a combination of MS and NMR-based methods. A genome sequence of the producing organism was also obtained, and this enabled the identification of the putative biosynthetic gene cluster responsible for the production of these new Ahp-containing cyclic depsipeptides. Curiously, no candidate gene for introducing the bromine atom was located in the cluster, suggesting that a *trans*-acting brominase may be responsible for its production.

Keywords: *Moorena* sp., brominated natural product, Ahp-containing cyclic depsipeptide

* Presenting author. E-mail address: b2ryu@ucsd.edu (Byeol Ryu)

S1.P74 The utilization of LED lamps to enhance metabolite production in plants – fagopyrins from buckwheat

Marta Hornyák^{a,*}, Monika Kula-Maximenko^b, Ulf Göransson^c, Blazej Slazak^{a,c}

^a W. Szafer Institute of Botany, Polish Academy of Sciences, Lubicz 46, 31-512 Cracow, Poland

^b The Franciszek Górski Institute of Plant Physiology, Polish Academy of Sciences, Niezapominajek 21, 30-239 Cracow, Poland

^c Pharmacognosy, Department of Pharmaceutical Biosciences, Uppsala University, Box 574, 751 23 Uppsala, Sweden

Light is one of the main signals detected by plants that influence plant development and metabolism. Metabolic processes in plants are sensitive to the ratio of blue (B) and red (R) light (Kami et al., 2010). In recent years, Light-Emitting Diodes (LEDs) have become the most prevalent light sources utilized in Controlled Environmental Agriculture (CEA), offering the capability to manipulate light spectra (Engler and Krarti, 2021). Fagopyrins (**1**) are photosensitizers present in common buckwheat (*Fagopyrum esculentum* Moench). These compounds have antifungal and antibacterial properties. There are known six forms of **1** (A-F). The primary form is protofagopyrin (**2**) which after irradiation converts to **1**. The highest content of **1** was found in buckwheat flowers (Kim and Hwang, 2020), specifically in their pistils (Hornyák et al., 2023). Our study aimed to investigate whether the LED spectrum modification may enhance the production of **1** and **2** in buckwheat pistils. The content of **1** and **2** in extracts from buckwheat pistils was analyzed using liquid chromatography-mass spectrometry (LC-MS). Common buckwheat can grow in CEA conditions using the LED lamp as the sole light source (Hornyák et al. 2022). Three LED lamps emitting wavelengths corresponding to red (R), blue (B), and red-blue (RB) light were tested. The results demonstrate that RB light stimulates the production of **1** and **2** in F form while R light had a promoting effect on production of **1** and **2** in E form in buckwheat pistils.

Keywords: *Fagopyrum esculentum*, light spectrum, photosensitizer, pistil

References

- Engler, N., Krarti, M. 2021. Review of energy efficiency in controlled environment agriculture. *Renew. Sustain. Energy Rev.*, 141, 110786.
- Hornyák M., Prystupa I., Kula-Maximenko M., Nilsson A., Andren P.E., Göransson U., Ślazak B. 2023. Friend or foe? - Photosensitive fagopyrins in buckwheat flowers. 15th International Symposium on Buckwheat. Puławy, 2-8 lipca. Książka abstraktów: 48. ISBN 978-83-7562-398-7.
- Hornyák, M., Dziurka, M., Kula-Maximenko, M., Pastuszak, J., Szczerba, A., Szklarczyk, M., Płazek, A. 2022. Photosynthetic efficiency, growth and secondary metabolism of common buckwheat (*Fagopyrum esculentum* Moench) in different controlled-environment production systems. *Sci. Rep.*, 12, 257.
- Kami, C., Lorrain, S., Hornitschek, P., Fankhauser, C. 2010. Light-regulated plant growth and development. *Curr. Top. Dev. Biol.*, 91, 29-66.
- Kim, J., Hwang, K. T. 2020. Fagopyrins in different parts of common buckwheat (*Fagopyrum esculentum*) and Tartary buckwheat (*F. tataricum*) during growth. *J. Food Compos. Anal.*, 86, 103354.

* Presenting author. E-mail address: m.hornyak@botany.pl (Marta Hornyák)

S1.P75 From bugs to drugs: genomic characterisation and bioinformatic analysis of *Streptomyces* strains from the bark beetle microbiome

Anne K. C. Goll*, Arne Bergfeld^a, Andreas Bechthold^a

Institute for Pharmaceutical Sciences, Department for Pharmaceutical Biology and Biotechnology, University of Freiburg, Germany

Natural product research in *Streptomyces* is crucial in the battle against antibiotic resistance. Although there has been extensive research on soil-associated *Streptomyces*, there is a growing interest in exploring insect-associated strains as well. Insect microbiomes, such as those found in bark beetles, host unique *Streptomyces* with distinct production profiles that are shaped by symbiotic interactions. Studying those *Streptomyces* can reveal new biosynthetic pathways and secondary metabolites that hold promise for developing new antimicrobial substances. In this context, three novel *Streptomyces* strains (*Streptomyces* sp. ITFR-6, ITFR-16 and ITFR-21) were isolated from the microbiome of the Central European bark beetle species *Ips typographus*, collected in the Black Forest, Germany. The aim of this study was to screen the strains using a genome mining approach to identify putative biosynthetic gene clusters (BGCs) and associate them with compounds released into production media. Methods used for identifying the *Streptomyces* strains included isolating them on selective agar plates and sequencing partially conserved genes, such as 16S rRNA and GyrB. Subsequently, whole genome sequencing of the three novel strains was performed. Comprehensive genome analysis was conducted using advanced genome mining techniques, such as antiSMASH, BiG-SCAPE, MIBiG, and KEGG-KAAS-MinPath. This approach allowed exploration of the biosynthetic potential of these microorganisms at the genomic level, revealing a large number of promising biosynthetic gene clusters encoding for secondary metabolites. Crude extracts were also purified and screened for bioactive compounds, as the three novel strains have previously demonstrated antibiotic properties. In this context various aspects and new developments will be discussed.

Keywords: *Streptomyces*, genome mining, RiPPs, bark beetle, insect microbiome

*Presenting author. E-mail address: anne.goll@pharmazie.uni-freiburg.de (Anne K. C. Goll)

S1.P76 A microscale cultivation platform integrating OSMAC to explore metabolomic changes and antimicrobial activity of a fungal strain collection

Alexandre Bory^{a,b,*}, Marine Cosseron^{a,b}, Alexandre Luscher^c, Sylvain Schnee^d, Thilo Köhler^c, Katia Gindro^d, Jean-Luc Wolfender^{a,b}

^a School of Pharmaceutical Sciences, University of Geneva, CMU, Rue Michel-Servet 1, Geneva, Switzerland

^b Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, CMU, Rue Michel-Servet 1, Geneva, Switzerland

^c Department of Microbiology and Molecular Medicine, University of Geneva, Rue Michel-Servet 1, Geneva, Switzerland

^d Mycology Group, Research Department Plant Protection, Agroscope, Route de Duiller 60, Nyon, Switzerland

Antimicrobial resistance is a major global health threat. Therefore, it is imperative to discover new antimicrobials with new modes of action. In this context, natural products derived from microorganisms have proven to possess very potent activities (Dai et al., 2020). However, their cultivation is a slow process and the frequent isolation of known active compounds is a limitation when screening natural extracts. To circumvent this problem, we recently developed a drug discovery platform compatible with the OSMAC approach (Bode et al., 2002). Cultivation of fungal strains, extraction, bioassays and metabolomics are performed in a standardized 96-well plate format thus allowing high-throughput and compatibility across the screening and analytics platforms. In a proof of concept study, sixty fungal strains were cultured in triplicates in eight different liquid media. Extracts were enriched by solid phase extraction and systematically submitted to antimicrobial bioassays and metabolomic profiling. Around 1'500 fungal extracts were generated and screened for antimicrobial and antivirulence activities. The highly reproducible results exhibited hits, mostly in triplicate, which were either spread across different media or limited to a specific medium; hence highlighting the power of the OSMAC approach to increase the hit rate. The full dataset was also analyzed by UHPLC-HRMS/MS allowing for the construction of a massive molecular network which integrates the bioactivity results (Olivon et al., 2017). The integration of the massive data generated allows efficient prioritization of strains and conditions for subsequent upscaling and targeted isolation of active candidates.

Keywords: OSMAC, antimicrobials, antivirulence, metabolomics, fungi

References

- Bode, H.B., Bethe, B., Höfs, R., Zeeck, A., 2002. Big Effects from Small Changes: Possible Ways to Explore Nature's Chemical Diversity. *ChemBioChem* 3, 619–27.
- Dai, J., Hand, R., Xud, Y., Lie, N., Wanga, J., Danc, W., 2020. Recent Progress of Antibacterial Natural Products: Future Antibiotics Candidates. *Bioorganic Chemistry* 101 (August): 103922.
- Olivon, F., Allard, P.M., Koval, A., Righi, D., Genta-Jouve, G., Neyts, J., Apel, C., Pannecouque, C., Nothias, L.F., Cachet, X., Marcourt, L., Roussi, F., Katanaev, V.L., Touboul, D., Wolfender, J.L., Litaudon, M., 2017. Bioactive Natural Products Prioritization Using Massive Multi-Informational Molecular Networks. *ACS Chem. Bio.* 12, 2644–51.

* Presenting author. E-mail address: alexandre.bory@unige.ch (Alexandre Bory)

S1.P77 Enhanced production of hexylaconitic anhydride compounds from endolichenic fungus *Xylaria* sp. by co-culture with *Chaetomium* sp.

Fotios A. Lyssaios^{a,*}, Azucena González-Coloma^b, M. Fe. Andrés^b, Carmen E. Díaz^a

^a Instituto de Productos Naturales y Agrobiología, IPNA-CSIC, Avda. Astrofísico F. Sánchez, nº 3, 38206. Tenerife Spain

^b Instituto de Ciencias Agrarias, ICA-CSIC, Serrano 115-bis, 28006, Madrid, Spain

Endolichenic fungi form a diverse group of microscopic fungi that live asymptotically in the lichen thalli (Agrawal et al., 2020). In nature microorganism co-exist in the same ecological habitat interacting with each other in a symbiotic or antagonistic manner. Secondary metabolites play a key role in these interactions acting as chemical signals for communication or competition of the limited nutrients, triggering various defense mechanisms. These stress factors may increase the production of known compounds or activate silent biosynthetic pathways resulting to new compounds (Knowles et al., 2022). Therefore, by mimicking the natural environment, we studied the mixed fermentation of two endolichenic fungi, *Xylaria* sp. and *Chaetomium* sp., isolated from the lichen, *Hypogymnia tubulosa*. Co-culturing these strains led to increase the total yield of the extract and a significant change in the production of secondary metabolites of *Xylaria* sp.. HPLC-MS profiles of co-culture were different compared to the monocultures resulting in a stimulation of hexylaconitic anhydride metabolites while piliformic acid derivatives decreased significantly. These metabolites were isolated from the extract and identified by mass spectrometry and NMR analysis.



Fig. 1. Morphological interaction observed among endolichenic strains in Petri dish.

The co-culture extract showed strong nematocidal effect against the plant-parasite *Meloidogyne javanica* while the control monocultures were not active. This enhancement was mainly attributed to major production of hexylaconitic anhydride. This compound was found also to be active against the fungi (*Botrytis cinerea*, *Alternaria alternata*) and aphids (*Myzus persicae*, *Rhopalosiphum padi*). These findings suggest the potential of this metabolite, along with the co-culture extract, as promising biocontrol agents for agricultural applications.

Keywords: endolichenic fungi, co-culture, *Xylaria* sp., hexylaconitic anhydride, biopesticide

References

- Agrawal, S., Deshmukh S. K., Reddy M. S, Prasad R., Goel M., 2020. Endolichenic fungi: A hidden source of bioactive metabolites., Afr. J. Bot. 134, 163-186.
- Knowles, S. L., Raja, H. A., Roberts, C. D., Oberlies, N. H., 2022. Fungal-fungal co-culture: a primer for generating chemical diversity. Nat Prod Rep. 39(8), 1557–1573.

* Presenting author. E-mail address: f.lyssaios@ipna.csic.es (Fotios A. Lyssaios)

S1.P78 Bioprospection of bioactive compounds from fungi in Cameroon: current status and outlook

Simeon Fogue Kouam*

Department of Chemistry, Higher Teacher Training College, University of Yaounde I, P.O. Box 47, Yaounde, Cameroon

Natural bioactive compounds are considered the cornerstone for the development of high-value products with applications in medicine, agriculture and the food industry. Plants have been long studied for their bioactive constituents and the use of microorganisms as an alternative appears to be a promising strategy for offering compounds with high therapeutic potential. Among microorganisms, fungi have played an important role in the production of antibiotics and other drugs for the treatment of infectious and non-infectious diseases. The diversification of fungi and their gene clusters involved in the biosynthesis of secondary metabolites indicate their unlimited potential to produce a plethora of molecules. Although various African cultures have long used fungi as medicines, they remain relatively unexplored from a chemical standpoint. Recent scientific efforts have targeted the bioprospecting of fungi in Cameroon for the isolation of bioactive constituents (Kenla et al. 2013, Happi et al. 2015, Teponno et al. 2017, Anoumedem et al. 2020, Mountessou et al. 2023).

We will highlight the current state of chemical research on fungi from Cameroon and special attention given to their investigation under a project funded by the Alexander von Humboldt Foundation called Humboldt Research Hub- CECANAPROF.

Keywords: fungi, bioactive compounds, Cameroon, CECANAPROF

References

- Kenla TJN, Tatong MDK, Talontsi FM, et al., 2013. Si-enterobactin from the endophytic *Streptomyces* sp. KT-S1-B5 – a potential silicon transporter in Nature? Chem Commun 49, 7641.
- Happi GM, Kouam SF, Talontsi FM, et al (2015). A new dimeric naphtho- γ -pyrone from an endophytic fungus *Aspergillus niger* AKRN associated with the roots of *Entandrophragma congoëse* collected in Cameroon. Z Naturforsch B 70:625–630.
- Teponno R, Noumeur S, Helaly S, et al., (2017) Furanones and anthranilic acid derivatives from the endophytic fungus *Dendrothyrium variisporum*. Molecules 22, 1674.
- Anoumedem EGM, Mountessou BYG, Kouam SF, et al., 2020. Simplicilones A and B isolated from the endophytic fungus *Simplicillium subtropicum* SPC3. Antibiotics 9, 753.
- Mountessou BYG, Anoumedem EGM, Kemkuignou BM, et al., 2023. Secondary metabolites of *Diaporthe* sp., isolated from the Cameroonian medicinal plant *Trema guineensis* (Schumach. & Thonn.) Ficalho. Beilstein J Org Chem 19, 1555– 1561.

* Presenting author. E-mail address: kfogue@yahoo.com (Simeon Fogue Kouam)

S1.P79 Natural product biosynthesis regulation in fungi. What can system biology tells us?

Delphine Nègre^{a,b}, Abdelhalim Larhlmi^b, Florence Thomas Giraud^{a,b}, Clément Poulain^{a,b}, Samuel Bertrand^{a,b,*}

^a Laboratoire des Sciences du Numérique de Nantes-UMR6004, Université de Nantes, France

^b Institut des Substances et Organismes de la Mer, ISOMer, Nantes Université, UR 2160, Nantes, France

Microorganisms is an unlimited reservoir of Natural Products (NPs) that are waiting to be discovered (Carroll *et al.* 2022). To get access to a higher chemical diversity, one possibility remains to modify culture condition thus activating silent biosynthetic pathways. This strategy called “One Strain Many Compounds” – OSMAC – is however very time consuming (Romano *et al.* 2018). Thus, one may consider using alternative strategies to anticipate NP production regulation. We are currently exploring the possibility to use System Biology approaches to explore such regulation using the fungus *Penicillium rubens* as a model organism (Guzmán-Chávez *et al.* 2018). Approaches based on Genome Scale Metabolic Networks (GSMN) are widely used to interpret metabolic regulation within an organism. A GSMN represents a model of an organisms based on genome encoded biological reactions. However, still many limitations exist to be able to use of GSMN modelling. One of the main limitations lays in the absence of NP biosynthetic pathways in the available GSMN (Nègre *et al.* 2023). After manual integration of various NPs biosynthetic pathways in the available GSMN of *P. rubens* (Agren *et al.* 2013), strategies to explore biosynthetic pathways regulation was performed to explore the impact of simple culture condition modification on NP production. Such “dry laboratory” conditions are currently being explored in “wet laboratory” experiments. In conclusion, the System Biology can be applied to NP chemistry as an alternate strategy to explore NP production regulation by fungi. Further in-depth study needs to be explored to innovative approach.

This project is funded by the ANR-18-CE43-0013-01 - FREE-NPs project.

Keywords: annotation, dereplication, LC-HRMS/MS

References

- Agren, R., L. Liu, S. Shoaie, W. Vongsangnak, I. Nookaew, *et al.*, 2013. The RAVEN Toolbox and Its Use for Generating a Genome-scale Metabolic Model for *Penicillium chrysogenum*. PLoS Computational Biology 9, e1002980. Carroll, A. R., B. R. Copp, R. A. Davis, R. A. Keyzers, M. R. Prinsep, 2022. Marine natural products. Natural Product Reports 39, 1122-1171.
- Guzmán-Chávez, F., R. D. Zwahlen, R. A. L. Bovenberg, A. J. M. Driessen, 2018. Engineering of the filamentous fungus *Penicillium chrysogenum* as cell factory for natural products. Frontiers in Microbiology 9, 2768.
- Nègre, D., A. Larhlmi, S. Bertrand, 2023. Reconciliation and evolution of *Penicillium rubens* genome-scale metabolic networks–What about specialised metabolism? PLoS ONE 18, e0289757.
- Romano, S., S. Jackson, S. Patry, A. Dobson, 2018. Extending the “One Strain Many Compounds” (OSMAC) principle to marine microorganisms. Marine Drugs 16, 244.

* Presenting author. E-mail address: samuel.bertrand@univ-nantes.fr (Samuel Bertrand)

S1.P80 Exploration of the clusioid clade metabolome through molecular networking: In the search for new pharmacophoric scaffolds of fungal UPR inhibitors

Thomas Charpentier^{a,*}, Mehdi A. Beniddir^b, Marc Litaudon^c, Cécile Apel^c, Guillaume Viault^a, Karine Leblanc^b, Antoine Grivard^a, Nadège Blon^a, Anne-Marie Le Ray Richomme^a, Nelly Bataillé-Simonneau^d, Emmanuel Pajot^e, Thomas Guillemette^d, Pascal Richomme^a

^a Université Angers, SONAS, SFR QUASAV, F-49000 Angers, France

^b Université Paris-Saclay, CNRS, BioCIS, Équipe "Chimie des Substances Naturelles" 17 Avenue des Sciences, 91400 Orsay, France

^c Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, UPR 2301, 91198 Gif-sur-Yvette, France

^d Université Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, 49070 Beaucouzé, France

^e Green Impulse SAS, 1 rue Alexandre Fleming, 49000 Angers, France

Exploring alternative approaches and seeking innovative solutions for crop protection is a major challenge for the next decades (Almeida et al., 2019). In this context we have developed a new strategy targeting compensatory fungal pathways, to increase the sensitivity of fungi to plant defenses and therefore controlling phytopathogenic infections (Charpentier et al., 2023). Among these pathways, the Unfolded Protein Response (UPR) appears as a promising target for the development of new antifungal drug (Joubert et al., 2011). Recently, polyhydroxylated and prenylated xanthenes have been identified as potent inhibitors of the IRE1 protein, the unique effector in fungal UPR pathway (Charpentier et al., 2023). To gain a deeper understanding of structure-activity relationships of these inhibitors, phytochemistry study have been undertaken. However, these xanthenes are most often isolated as complex mixtures extracted from different parts of plants belonging to the clade clusioid. Therefore, as a dereplication strategy, feature-based molecular networking (FBMN) (Nothias *et al*, 2020) was applied to map the chemical space of clusioid plant extracts and a MS/MS spectra database of 130 xanthenes with different scaffolds and know active compounds, has been constituted. As will be shown this data base allowed us to annotate one FBMN as well as to explore chemical diversity of xanthenes clusters in order to select nodes of interest.

Keywords: xanthenes, Molecular Networking, MS/MS spectra database, UPR inhibitor

References

- Almeida, F., Rodrigues, M.L., Coelho, C., 2019. The still underestimated problem of fungal diseases worldwide. *Front. Microbiol.* 10, 214.
- Charpentier, T., Viault, G., Le Ray Richomme, A-M., Bataillé-Simonneau, N., Helesbeux, J-J., Blon, N., Bastide, F., Marchi, M., Aligon, S., Bruguère, A., Dinh, C. P., Benbelkacem, Z., Dallery, J-F, Simoneau, P., Richomme, P., Guillemette, T., 2023. Natural Products Targeting the Fungal Unfolded Protein Response as an Alternative Crop Protection Strategy. *J. Agric. Food Chem.* 71, 13706-13716.
- Joubert, A., Simoneau, P., Campion, C., Bataille-Simonneau, N., Iacomini-Vasilescu, B., Poupard, P., François, J. M., Georgeault, S., Sellier, E., Guillemette, T., 2011. Impact of the Unfolded Protein Response on the Pathogenicity of the Necrotrophic Fungus *Alternaria Brassicicola* : Impact of the UPR on *Alternaria brassicicola* Virulence. *Mol. Microbiol.* 79, 1305-1324.
- Nothias, L.-F., Petras, D., Schmid, R., *et al*, 2020. Feature-based molecular networking in the GNPS analysis environment. *Nat. Methods* 17, 905-908.

* Presenting author. E-mail address: thomas.charpentier@univ-angers.fr (Thomas Charpentier)

S1.P81 Differences in chemical compounds of selected strains of *Hericium erinaceus*

Jaroslav Matejka*, Petr Marsik, Jan Stastny, Ivan Jablonsky, Tomas Skala, Zdenka Kahankova, Pavel Kloucek

Department of Food Science, Czech University of Life Sciences Prague, Kamýcká 129, 165 00 Praha-Suchbát, Czech Republic

Hericium erinaceus, a medicinal mushroom, has been known in Chinese traditional medicine for its positive effects on human health. It has been used to treat a wide range of health problems or as a prevention of various diseases including Alzheimer's and Parkinson's disease, ischemic stroke or stomach disorders. *H. erinaceus* have neuroprotective properties, anticancer effects, antiglycemic, antidepressant, antimicrobial and antioxidant effects (Thongbai et al., 2015). The aim of our work was to determine the differences in the metabolome of selected strains of *H. erinaceus*. Six samples from Asia, Belgium and Czech Republic were used for the analysis. After metabolite extraction, their abundance was determined by LC/HRMS, and results were evaluated by multivariate analysis. From the reduced set of the most significant features, 13 of them were selected by comparison with published data, which were tentatively identified on the basis of MS/MS spectra and extracted ion chromatograms. By statistical comparison, we found that the strains of the genus *Hericium* show significant differences in metabolic profile, which corresponded with their origin. In the strains of Czech origin, the substances hericerin A, which has anticancer effects (Li et al., 2015), and hericenone J, which has been found to have neurotrophic activity (Ueda et al., 2008), were more represented. The biological activity of the observed strains was then tested for their anti-inflammatory effect in THP-1 cell culture in vitro. The test proved, that *H. erinaceus* extracts inhibited the production of inflammatory markers in this model system.

Keywords: *Hericium erinaceus*, bioactive substances, biological activity, LC-HRMS

References

- Li, W., Zhou, W., Kim, E., Shin, S.H., Kang, H.K., Kim, Y.H., 2015. Isolation and identification of aromatic compounds in Lion's Mane Mushroom and their anticancer activities. *Food Chemistry*. 170, 336-342.
- Thongbai, B., Rapior, S., Hyde, K. D., Wittstein, K., & Stadler, M., 2015. *Hericium erinaceus*, an amazing medicinal mushroom. *Mycological Progress*. 14, 1-23.
- Ueda, K., Tsujimori, M., Kodani, S., Chiba, A., Kubo, M., Masuno, K., Sekiya, K., Kawagishi, H., 2008. An endoplasmic reticulum (ER) stress-suppressive compound and its analogues from the mushroom *Hericium erinaceum*. *Bioorganic & Medicinal Chemistry*. 16, 9467-9470.

* Presenting author. E-mail address: matejkaj@af.czu.cz (Jaroslav Matejka)

S1.P82 Genome mining paired with stable isotope labeling allows for discovery of new cyanopeptolins in cyanobacterium *Nostoc* sp. UIC10890

Isabel M. Chauvin^{*}, Kelsey L. Alexander, Jimmy Orjala

Department of Pharmaceutical Sciences, College of Pharmacy, University of Illinois Chicago, 833 S. Wood Street, Chicago, IL, 60612, USA

Cyanobacteria, also known as blue-green algae, are a phylum of gram-negative photosynthetic autotrophs that can be found in almost every biome of the world. Their secondary metabolites are diverse in chemical structure and many have been shown to have therapeutic relevance. They are also capable of nitrogen fixation, which influences their production of nitrogen-containing compounds such as non-ribosomal peptides (NRPs). Traditional workflows for natural product drug discovery are usually bioassay-guided to prioritize activity, however advances in technology as well as lowering costs of sequencing offer the alternative of using a genome mining-guided workflow to prioritize the discovery of novel structures and specific compound classes. The use of genome mining on a *Nostoc* sp. cyanobacterial strain, UIC10890, led to the discovery of an NRP biosynthetic gene cluster encoding for the production of a new cyanopeptolin. Cyanopeptolins (CPs) are depsipeptides characterized by the presence of 3-amino-6-hydroxy-2-piperidone in the third position, often with side chains and N-methylation occurring at other positions. ¹⁵N stable isotope labeling followed by comparative metabolomics using tandem mass spectrometry (MS/MS) allowed for the detection and subsequent isolation of two new CPs from UIC10890. The structures of the CPs were confirmed through NMR spectroscopy. This poster will present the new CPs, their biosynthetic gene cluster, and the workflow used to isolate them. This research demonstrates a genetics-guided approach drug discovery and how it can be used to expedite dereplication as well as discover novel compounds.

Keywords: cyanobacteria, cyanopeptolin, genome mining, biosynthesis, mass spectrometry

^{*} Presenting author. E-mail address: ichauv2@uic.edu (Isabel M. Chauvin)

S1.P83 Diversity of *Origanum onites* essential oils in the ecological regions of Eastern Mediterranean

Dimitrios Mertzanidis^{a,c}, Alexandros Nakas^{b,c,*}, Andreana N. Assimopoulou^{b,c}, Stella Kokkini^{a,c}

^a Laboratory of Systematic Botany and Phytogeography, School of Biology, Aristotle University of Thessaloniki, Thessaloniki 54124, Greece

^b Laboratory of Organic Chemistry, School of Chemical Engineering, Aristotle University of Thessaloniki, Thessaloniki 54124, Greece

^c Center for Interdisciplinary Research and Innovation, Natural Products Research Centre of Excellence (NatPro-AUTH), Aristotle University of Thessaloniki, Thessaloniki 57001, Greece

Origanum onites L. is a very popular aromatic herb widely used in everyday cooking and traditional medicine since antiquity (Hanlidou et al., 2004). The native range of the species is restricted in Eastern Mediterranean; it extends from South and West Turkey, via South Greece and islands, up to Sicily (POWO, 2024). The essential oils (EOs) of *O. onites* have been studied for a number of potential applications in the pharmaceutical and food industry, such as antioxidant, antimicrobial, antifungal and health protective agents (Kintzios, 2002; Canli et al., 2023). Data from 37 years of scientific literature and new headspace GC-MS analyses were examined to assess the EOs diversity of wild *O. onites* plants. Information concerning the content in EO and its composition – focused on main oil components: γ -terpinene, *p*-cymene, carvacrol, thymol and linalool – were collected from different localities scattered along the total species range, and discussed in relation to six ecological regions of Eastern Mediterranean (EEA, 2024). The highest values of oil content have been recorded in the ecological region of Aegean & West Turkey sclerophyllous and mixed forest (up to 7.0% v/d.w., island of Chios). The EOs from the localities of the three Mediterranean ecological regions are characterized by the dominant occurrence of carvacrol (up to 92.7% of the total oil, island of Kos). The oil composition in the southeastern part of species range, which belongs to the more humid and colder Mediterranean-Continental ecological regions, is much more diversified, and characterized by either carvacrol, linalool and/or thymol.

Acknowledgements: This research has been co-financed by the European Union and Greek national funds through the Operational Program Competitiveness, Entrepreneurship and Innovation, under the call RESEARCH – CREATE – INNOVATE (project code: T1EDK-04174).

Keywords: oregano, *Origanum onites*, ecological regions, climatic regions, headspace GC-MS

References

- Canli, K., Bozyel, M.E., Turu, D., Benek, A., Simsek, O., Altuner, E.M., 2023. Biochemical, antioxidant properties and antimicrobial activity of steno-endemic *Origanum onites*. *Microorganisms* 11, 1987.
- European Environmental Agency, 'Digital map of European ecological regions', <https://www.eea.europa.eu/data-and-maps/figures/dmeer-digital-map-of-european-ecological-regions>; Accessed February 06, 2024.
- (<https://www.eea.europa.eu/legal/copyright>). Copyright holder: European Topic Centre on Biological Diversity (ETC/BD).
- Hanlidou, E., Karousou, R., Kleftoyanni, V., Kokkini, S., 2004. The herbal market of Thessaloniki (N Greece) and its relation to the ethnobotanical tradition. *J. Ethnopharmacol.* 91, 281-299.
- Kintzios, S.E., 2002. *Oregano: the genera Origanum and Lippia*. Medicinal and aromatic plants - industrial profiles. Taylor and Francis Group, New York.
- POWO. Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. <https://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:453331-1>; Accessed February 06, 2024.

* Presenting author. E-mail address: anakas@cheng.auth.gr (Alexandros Nakas)

S1.P84 Comparative GC/MS analysis of essential oils from two varieties of *Vitex negundo* L.

Olha Korablova^a, Iryna Levchuk^b, Mariia Shanaida^{c,*}

^a M.M. Gryshko National Botanical Garden, National Academy of Sciences of Ukraine, Sadovo-Botanichna Str. 1, 01014 Kyiv, Ukraine

^b Scientific and Research Center for Products Testing, State Enterprise "UkrMetrTestStandart", Metrology Str. 4, 03143 Kyiv, Ukraine

^c Department of Pharmacognosy and Medical Botany, I. Horbachevsky Ternopil National Medical University, Voli Str. 1, 46001 Ternopil, Ukraine

The genus *Vitex* L. includes several hundred species of woody plants (Islam et al., 2024). Some *Vitex* species are of significant scientific interest as they are widely used in traditional medicine of many countries. However, only fruits of *V. agnus-castus* L. are used in the conventional medicine (European Pharmacopoeia, 11th ed., 2023). Phytochemical analysis of different *Vitex* representatives is a very promising direction for further research, especially because genetic prerequisites and climatic conditions have a great influence on the biosynthesis of bioactive compounds (Shanaida et al., 2021). In this study, the gas chromatography-mass spectrometry (GC/MS) method was used to analyze the composition of essential oils from the flowering shoots of two *Vitex negundo* varieties: *V. negundo* var. *heterophylla* (Franch.) Rehd. and *V. negundo* var. *cannabifolia* L. The raw materials were harvested in 2022 under cultivation of plants in M.M. Gryshko National Botanical Garden (Kyiv, Ukraine). The GC/MS study revealed that the predominant components of *V. negundo* var. *heterophylla* essential oils were α -terpineol (13.58%), terpinyl acetate (11.37%), borneol (10.90%), 1,8-cineol (eucalyptol) (9.52%), *beta*-farnesene (8.98%), bicyclogermacrene (7.98%) and terpinen-4-ol (7.39%). The following terpenoids prevailed in the essential oil of *V. negundo* var. *cannabifolia*: borneol (58.36%), *trans*-sabinen hydrate (6.96%), eugenol (6.22%), α -terpinol (5.70%), terpinyl acetate (4.51%) and 1,8-cineol (3.61%). As the revealed predominant volatile compounds have the well-defined therapeutic properties, the essential oils of the studied *V. negundo* varieties could be considered for further investigations of their pharmacological effects.

Keywords: chromatographic analysis, flowering shoots, volatile compounds, *Vitex negundo* var. *heterophylla*, *Vitex negundo* var. *cannabifolia*

References

Islam, Z., Caldeira, G.I., Caniça, M., Islam, N., Silva, O., 2024. *Vitex* Genus as a Source of Antimicrobial Agents. *Plants*. 13(3), 401.

European Pharmacopoeia (11th ed.). 2023. <https://pheur.edqm.eu/home>

Shanaida, M., Holenko, A., Korablova, O., 2021. Gas-chromatographic determination of volatile compounds from the flowering shoots of some *Vitex* L. representatives. *PharmacologyOnLine*. 3, 1265-1274.

* Presenting author. E-mail address: shanayda@tdmu.edu.ua (Mariia Shanaida)

S1.P85 Molecular interactions in the ocean microbiome: tracking fungal metabolites modifying *Prorocentrum lima* physiology and toxin production

Olivier Grovel^{a,*}, Olivier Berry^a, Fabienne Herve^b, Simon Tanniou^b, Ruth Julianna Gutierrez^{a,b}, Enora Briand^b, Samuel Bertrand^{a,c}

^a Nantes Université, Institut des Substances et Organismes de la Mer, ISOMer, UR 2160, Nantes, France

^b IFREMER, PHYTOX, F-44311 Nantes, France

^c Nantes Université, Laboratoire des Sciences du Numérique de Nantes-UMR6004, Nantes, France

Harmful algal blooms are associated with detrimental effects on their environment and have a negative impact on associated human economic activities (shellfish farming) (Grattan et al., 2016). Toxin production is still a poorly explained phenomenon that could be related with microbial chemical induction. In this project, we worked on an original interaction model between the bloom forming toxic microalga *Prorocentrum lima*, and its co-isolated fungus *Aspergillus pseudoglaucus* (Berry et al., 2023). We observed that toxin production by the microalga was up-regulated in co- culture conditions and that the fungal crude extract yielded the same effect. Applying both a bioactivity and LC-HRMS/MS-guided strategy (including molecular networking and dereplication), fungal molecules altering *P. lima* growth and its production of diarrhetic toxins were tracked. From the initial crude extract, some fractions which were mainly composed of anthraquinones and echinulin derivatives seemed to interfere with normal metabolism of the microalga. Two other fractions were shown to lower or increase toxin quantities, due to the presence of polar anthraquinone derivatives and some unknown compounds. These fractions underwent further purification work to isolate their major molecules whose structures were solved by NMR experiments. All these isolated molecules were evaluated on *P. lima* physiology. This study showed that *A. pseudoglaucus* compounds can exert different effects on *P. lima* physiology, such as increased or decreased growth, or variation of toxin concentration and/or excretion in the culture medium. Further investigations are ongoing to investigate their respective and global role in the consortium co-culture.

Keywords: *Prorocentrum lima*, *Aspergillus pseudoglaucus*, microalgal toxins, fungal natural products, chemical interactions

References

Berry, O., E. Briand, A. Bagot, M. Chaigné, L. Meslet-Cladière, et al., 2023. Deciphering interactions between the marine dinoflagellate *Prorocentrum lima* and the fungus *Aspergillus pseudoglaucus*. *Env. Microbiol.* 25, 250-267. Grattan, L.M., S. Holobaugh, J.G. Morris, 2016. Harmful algal blooms and public health. *Harmful Algae* 57, 2-8.

* Presenting author. E-mail address: samuel.bertrand@univ-nantes.fr (Samuel Bertrand)

S1.P86 Chemical mediation of algal microbiome across temperate and subtropical regions

Skylar Carlson^{a,*}, Savannah Pierce^a, Melany Puglisi-Weening^b

^a Chemistry Department, University of the Pacific, Stockton, CA, USA

^b Department of Pharmaceutical Science, Chicago State University Chicago, IL, USA

The surface microbiomes of plants and animals plays a critical role in maintaining the health of the host. As climate change warms and acidifies the ocean, alterations in the microbiome of marine algae and ideal conditions for opportunistic pathogens could lead to an unprecedented loss of algal biodiversity (Minich et al., 2018; van der Loos et al., 2019). We examined the role algal metabolites play in shaping the microbiome across two climates. Algal species, from two subtropical sites in Florida and southern California and two temperate sites in northern California, were swabbed for surface associated bacteria (SAB) and extracted (CA) or extracted and partitioned (FL). These metabolite mixtures were then screened for SAB growth-promotion or inhibition activity. Our experiments from California display site specific bacterial growth modulation. Bacteria from the same location displayed neutral activity (9% promotion, 20% inhibition) but bacteria from other sites were strongly modified by the algal extracts (27% and 43%). Chloroform partitions from Florida algae inhibited the growth of 24% of FL SAB. This demonstrates a specific relationship between these partitions and FL SAB, compared to the other two partitions. Across phyla, we observed neutral activity in 3 partitions (no strong effect on any bacteria). Comparing across the two climates, more inhibition was observed from the warmer climates suggesting the presence of stronger defenses against potential pathogens. These results demonstrate that algal chemistry plays a role in shaping SAB population structure warranting future studies to identify the metabolites responsible for the observed inhibition.

Keywords: signaling, algae, climate change, partitions, microbiome

References

- Minich, J.J., Morris, M.M., Brown, M., Doane, M., Edwards, M.S., Michael, T.P., and Dinsdale, E.A. 2018. Elevated temperature drives kelp microbiome dysbiosis, while elevated carbon dioxide induces water microbiome disruption. *PLoS One* 13, e0192772.
- van der Loos, L. M., Eriksson, B.K., Salles, J.F. 2019. The macroalgal holobiont in a changing sea. *Trends in Microbiology* 27, 635-650.

* Presenting author. E-mail address: scarlson1@pacific.edu (Skylar Carlson)

S1.P87 Marine actinobacterium as a source of natural products against multiple myeloma

Emre Gezer^{a,b,c,*}, Eyüp Bilgi^c, Erdal Bedir^c, Muriel Cuendet^{a,b}

^a School of Pharmaceutical Sciences, University of Geneva, 1211 Geneva, Switzerland

^b Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, 1211 Geneva, Switzerland

^c Department of Bioengineering, Faculty of Engineering, Izmir Institute of Technology, 35430, Urla, Izmir, Turkey

Multiple myeloma is a hematologic malignancy characterized by the abnormal proliferation of clonal plasma cells. Patients frequently experience relapse and develop resistance to the currently available drugs. Therefore, there is a need to find new treatment options, and the investigation of sources of natural products is of interest (Yu et al., 2022). The present study evaluated the antiproliferative activity of compounds produced by a marine actinobacterium, *Streptomyces cacaoi*, in various multiple myeloma cell lines. The fermentation conditions were first optimized using a statistical experimental design to enhance the production of bioactive secondary metabolites (Gezer et al., 2022). Subsequently, the antiproliferative activity of the EtOAc extracts was tested in RPMI 8226 multiple myeloma cells at a concentration of 45 µg/mL. Incubation temperature (25-35°C) and water type (distilled vs seawater) strongly affected the production of bioactive compounds. Extracts obtained after fermentation in distilled water at 25°C were the most active. Then, the effect of some inorganic chemicals was also evaluated. Surprisingly, KNO₃, which is frequently used in media, caused a significant decrease in the antiproliferative activity of the extracts. Finally, three polyethers isolated from an extract obtained with optimized conditions showed antiproliferative activity with IC₅₀ values in the low micromolar concentrations against the multiple myeloma cell lines RPMI 8226 and KMS-12-BM, as well as their drug-resistant counterparts. In conclusion, this study demonstrates the critical impact of fermentation conditions on producing bioactive natural products against multiple myeloma cancer cell lines.

Keywords: multiple myeloma, *Streptomyces cacaoi*, optimization, fermentation, cytotoxicity

References

- Gezer, E., Üner, G., Küçüksolak, M., Kurt, M.Ü., Doğan, G., Kırmızıbayrak, P.B., Bedir, E., 2022. Undescribed polyether ionophores from *Streptomyces cacaoi* and their antibacterial and antiproliferative activities. *Phytochemistry*. 195, 113038.
- Yu, C.-C., Li, Y., Cheng, Z.-J., Wang, X., Mao, W., Zhang, Y.-W., 2022. Active components of traditional chinese Medicinal Material for multiple myeloma: Current evidence and future directions. *Frontiers in Pharmacology*. 13, 818179.

* Presenting author. E-mail address: emre.gezer@unige.ch (Emre Gezer)

S1.P88 Parishin derivatives from the aerial parts of *Sedirea japonica*

Yoon Seo Jang*, Gang Min Noh, Ki Hyun Kim

School of Pharmacy, Sungkyunkwan University (SKKU), Suwon 16419, Republic of Korea

Sedirea japonica (Orchidaceae) is found in Eastern Asian countries such as Korea, Japan, and China. Their attractive beauty and ornamental value have long been appreciated. In Eastern culture, *S. japonica* was commonly given as a gift (Lee et al., 2021). However, indiscriminate harvesting led to the endangered status of *S. japonica* in Korea and Japan (Suetsugu et al., 2013). Under this circumstance, only a few phytochemical studies of this plant have been conducted. In our natural product research to discover new metabolites, eight compounds including five new parishin derivatives, sedishins A-E (1-5) and three phenolic compounds (6-8) were successfully isolated from the aerial parts of *S. japonica* through LC/MS guided isolation approach. The structures of new compounds (1-5) were characterized through one-dimensional (1D) and two-dimensional (2D) NMR (¹H-¹H COSY, HSQC, HMBC) spectral data analysis, combined with high-resolution electrospray ionization mass spectrometry (HR-ESI-MS) analysis. All isolated compounds will be further evaluated for their biological activity.




Fig. 1. Structure of sedishin A.

Keywords: *Sedirea japonica*, Orchidaceae, parishin derivatives, sedishins A-E, structural elucidation

References

- Lee, K. C.; An, J.; Hwang, J. E.; Kim, P. B.; Park, H. B.; Kim, S.; Park, H. J.; Lee, C. W.; Lee, B. D.; Kim, N. Y. 2021. Effects of light condition on growth and physiological characteristics of the endangered species *Sedirea japonica* under RCP 6.0 climate change scenarios. *Plants (Basel)*, 10, 1891.
- Suetsugu, K., Tanaka, K., 2013. Pollination of *Sedirea japonica* (Orchidaceae) by *Bombus diversus diversus* (Hymenoptera: Apidae). *Eur. J. of Entomol.* 110, 545-548.

* Presenting author. E-mail address: bbj0423@gmail.com (Yoon Seo Jang)

S1.P89 Nitrobenzoyl sesquiterpenoids from a Hawaiian marine-derived fungus FM451

Chen Huo*, Ki Hyun Kim

School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea

Marine-derived fungi represent a vast reservoir of novel compounds, owing to their extensive chemical diversity and demonstrated ability to produce natural products with therapeutic and pharmacological potential (Hasan et al., 2015). In a screening test, the methanolic extract of a Hawaiian marine fungal strain FM451 was found to exhibit potent antimicrobial activity against *Staphylococcus aureus*. LC/MS-guided isolation of the methanolic extract of the fungus led to the isolation of six α -pyrone-derived compounds, including one undescribed compound (**1**) and five known derivatives (**2–5**), as well as six nitrobenzoyl sesquiterpenoids, including one unreported compound (**6**) (Fig. 1) and five known derivatives (**7–11**). Structural elucidation of the new compounds was achieved through extensive NMR spectroscopic methods and high-resolution electrospray ionization mass spectrometry (HR-ESI-MS) data analysis. Furthermore, their absolute configurations were unambiguously determined via theoretical electronic circular dichroism (ECD) calculation and DP4+ analysis. All of these compounds will be evaluated for their antimicrobial activities against *S. aureus*.




Fig. 1. Structure of the new nitrobenzoyl sesquiterpenoid **6**.

Keywords: marine-derived fungus, nitrobenzoyl sesquiterpenoid, antimicrobial activity

References

Hasan, S., Ansari, M. I., Ahmad, A., Mishra, M., 2015. Major bioactive metabolites from marine fungi: A Review. *Bioinformation*. 11, 176–181.

* Presenting author. E-mail address: huochen_0213@163.com (Chen Huo)

S1.P90 Steroid derivatives from cannabis-associated endophytic fungus *Hypoxylon perforatum*

Chen Huo*, Seo Yoon Lee, Ki Hyun Kim

^a School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea

Endophytic fungi, residing within plants, serve as prolific producers of novel and structurally diverse bioactive compounds, which hold promise as potential therapeutics for addressing major global diseases (Manganyi et al., 2020). *Cannabis sativa* L., commonly known as marijuana or hemp, is a versatile plant species with a long history of human cultivation for medicinal, industrial, and recreational purposes (Bonini et al., 2018). *Hypoxylon perforatum* was isolated from *C. sativa* as an intriguing endophytic fungus. Three new steroid derivatives (**1–3**) were isolated and characterized from the cultivation of cannabis-associated endophytic fungus *H. perforatum* (Fig. 1). The structural elucidation of these compounds was achieved through a comprehensive approach, combining 1D and 2D NMR spectroscopy techniques (including HSQC, HMBC, COSY, and NOESY) with high-resolution electrospray ionization mass spectrometry (HR-ESI-MS) data analysis. Additionally, the determination of their absolute configurations was confidently established through theoretical electronic circular dichroism (ECD) calculations and DP4+ analysis. The investigation into the bioactivity of these newly isolated steroid derivatives is ongoing.




Fig. 1. Structure of new steroid derivative **1**.

Keywords: endophytic fungi, *Hypoxylon perforatum*, steroid, absolute configurations

References

- Manganyi, M. C., Ateba, C. N., 2020. Untapped potentials of endophytic fungi: A review of novel bioactive compounds with biological applications. *Microorganisms*. 8, 1934.
- Bonini, S. A., Premoli, M., Tambaro, S., Kumar, A., Maccarinelli, G., Memo, M., Mastinu, A., 2018. Cannabis sativa: A comprehensive ethnopharmacological review of a medicinal plant with a long history. *J. Ethnopharmacol.* 227, 300–315.

* Presenting author. E-mail address: huochen_0213@163.com (Chen Huo)

S1.P91 Sesquiterpene-phthalide hybrids from cannabis-associated endophytic fungus *Zopfiella* sp.

Seo Yoon Lee*, Ki Hyun Kim

School of Pharmacy, Sungkyunkwan University (SKKU), Suwon 16419, Republic of Korea

Cannabis sativa is regulated as a drug in many countries due to its addictive properties. Endophytic fungi that interact with *C. sativa* plants have been investigated, which is expected to produce bioactive compounds. A total of 27 endophytic fungi were isolated from *C. sativa* and identified using ITS rDNA sequences. Among them, the endophytic fungus *Zopfiella* sp. was first isolated from *C. sativa*. LC-MS-based GNPS networking analysis has shown that *Zopfiella* sp. may produce unique compounds when grown in media containing *C. sativa* stem powder. As part of our ongoing research to discover bioactive natural products, we have isolated four new sesquiterpene-phthalide hybrids (**1-4**) (Fig. 1), along with five known compounds (**5-9**) including steroids and phthalide derivatives from the cultivation of cannabis-associated endophytic fungus *Zopfiella* sp. The structures of four new compounds were elucidated through analysis of 1D and 2D (¹H-¹H COSY, HSQC, HMBC and NOESY) NMR, HR-ESIMS, optical rotation, as well as ECD calculation. All isolated compounds will be further evaluated for their biological activity.




Fig. 1. Structure of new sesquiterpene-phthalide hybrids **1**.

Keywords: *Cannabis sativa*, *Zopfiella* sp., endophytic fungi, sesquiterpene-phthalide hybrids

* Presenting author. E-mail address: allag8201@naver.com (Seo Yoon Lee)

S1.P92 A new cyclic peptide and two new ellagic acid glycoside derivatives from *Jatropha podagrica* leaves

Seo Yoon Lee*, Se Yun Jeong, Ki Hyun Kim

School of Pharmacy, Sungkyunkwan University (SKKU), Suwon 16419, Republic of Korea

Jatropha podagrica, a succulent plant belonging to the Euphorbiaceae family, contains various phytochemicals, which is why it has been used as a traditional medicine for analgesic, tonic, and purgative purposes (Jang et al., 2023). The genus *Jatropha* is a combination of the Greek words 'jatos' (doctor) and 'trophe' (food), implying its medicinal uses (Kumar et al., 2008). *Jatropha* species have been used to treat various diseases in Asia and Latin America and are reported to have minimal side effects (Cowan et al., 1999). As a part of our continuing research to discover bioactive natural products, we successfully isolated one new cyclic peptide (**1**) (Fig. 1) and two new ellagic acid glucoside derivatives (**2-3**), along with fourteen known compounds (**4-17**) including megastigmane glycosides, and phenolic derivatives. The structures of three new compounds (**1-3**) were determined through comprehensive analysis of 1D and 2D (¹H-¹H COSY, HSQC, HMBC and NOESY) NMR, HR-ESIMS, and optical rotation, as well as ECD calculation. The absolute configuration of compound (**1**) was assigned using the advanced Marfey's method. All the isolated compounds, except for compound (**14**), were identified from *J. podagrica* for the first time. All isolated compounds will be further evaluated for their biological activity.




Fig. 1. Chemical structure of new cyclic peptide **1**

Keywords: *Jatropha podagrica*, Euphorbiaceae, cyclic peptide, ellagic acid glycoside, Marfey's method

References

- Cowan, M. M., 1999. Plant products as antimicrobial agents. Clin. Microbiol. Rev. 12(4), 564-582.
- Jang, Y.S., Lee, D.E., Ju, D.U., Jeong, S.Y., Ko, Y.J., Pang, C., Kang, K.S., Gwon, H.J., Yoo, H.M., Kim, K.H., 2023. Antiviral effects of secondary metabolites from *Jatropha podagrica* leaves against the pseudotyped virus of SARS-CoV- 2 Omicron. Plants, 12(23), 3942.
- Kumar, A., Sharma, S., 2008. An evaluation of multipurpose oil seed crop for industrial uses (*Jatropha curcas* L.): A review. Ind. Crops. Prod. 28(1), 1-10.

* Presenting author. E-mail address: allag8201@naver.com (Seo Yoon Lee)

S1.P93 Discovery of new sesqui-lignans from *Salix gracilistyla* leaves and their neuroprotective effects

Yeo Rang Cho*, Da Eun Lee, Jae-Cheol Lee, Ki Hyun Kim

School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea

Salix gracilistyla, commonly known as pussy willow, is primarily distributed along river streams in Eastern Asia, including South Korea. Various parts of *S.gracilistyla*, such as the leaves, stems, and bark, have been used in traditional medicine for treating skin diseases, wounds, and arthritis, and as a painkiller. Previous pharmacological studies on *S.gracilistyla* reported that these extracts showed anti-inflammatory activity by inhibiting NO production in LPS-stimulated macrophages (Ryu et al.,2003). NO-induced inflammation can increase oxidative stress in the brain, potentially triggering neurodegenerative disorders (Yang et al.,2013). In our ongoing research to identify new bioactive compounds from medicinal plants, the ethanolic extract of *S. gracilistyla* leaves was chemically investigated, and two new sesqui-lignans (**1-2**) were isolated (Fig. 1), along with 18 known compounds (**3-20**). The structures of the new compounds were determined by analyzing 1D- and 2D- NMR spectroscopic data and HR-ESIMS. Their absolute configuration was determined through comparison of their experimental circular dichroism (CD) data with calculated electronic circular dichroism (ECD) spectra. In this study, we evaluated the neuroprotective effects of all the isolated compounds from *S. gracilistyla* on iPSC-derived cortical neurons.



Fig 1. Chemical structure of compound 1.

Keywords: *Salix gracilistyla*, Salicaceae, lignans, ECD, neuroprotective effects

References

- Ryu, J.H., Ahn, H., Kim, J.Y., Kim, Y.K, 2003. Inhibitory activity of plant extracts on nitric oxide synthesis in LPS- activated macrophages. *Phytother. Res.* 17, 485–489.
- Yang, H., Lee, S., Sung, S., Kim, J., Kim, Y., 2013. Neuroprotective compounds from *Salix pseudo-lasiogyne* twigs and their anti-amnesic effects on scopolamine-induced memory deficit in mice. *Planta Med.* 79, 78-82.

* Presenting author. E-mail address: yr6755@g.skku.edu (Yeo Rang Cho)

S1.P94 Lipopeptides from the bioluminescent bacterium *Photorhabdus luminescens*

Yeo Rang Cho*, Seung Hye Cho, Ki Hyun Kim

School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea

Photorhabdus luminescens is a Gram-negative, entomopathogenic, and bioluminescent bacterium that establishes a mutualistic relationship with entomophagous nematodes. When these nematodes invade insects, they enter the hemolymph and release *P. luminescens* from their gut (Blackburn et al., 1998). Previous pharmacological studies reported that *P. luminescens* produces antimicrobial substances with antifungal or antibacterial activity to prevent the invasion of other microorganisms, supporting the growth and reproduction of the nematodes (Bowen et al., 1998). As part of our ongoing natural product research aimed at identifying bioactive new compounds, we conducted chemical investigations on the EtOAc extract of *P. luminescens* cultivation using LC/MS, HPLC, and TLC- based analyses. This process resulted in the isolation of two new lipopeptides, photorhamides A (1) and B (2) (Fig. 1), together with six known compounds (3-8). The structures of the new compounds were confirmed by analyzing their 1D- and 2D-NMR data and HR-ESIMS. The entire peptide sequence was determined through the interpretation of HSQC, HMBC, and NOESY signals, and the absolute configuration was established using Marfey's method. The biological activities of all the isolated compounds will be further evaluated.




Fig 1. Structure of photorhamide A (1)

Keywords: *Photorhabdus luminescens*, lipopeptides, antimicrobial activity, Marfey's method

References

- Blackburn, M., Golubeva, E., Bowen, D., French-Constant, R. H., 1998. A novel insecticidal toxin from *Photorhabdus luminescens*, toxin complex a (Tca), and its histopathological effects on the midgut of *Manduca sexta*. Appl. Environ. Microbiol. 64, 3036-3041.
- Bowen, D. J., Ensign, J. C., 1998. Purification and characterization of a high-molecular-weight insecticidal protein complex produced by the entomopathogenic bacterium *Photorhabdus luminescens*. Appl. Environ. Microbiol. 64, 3029- 3035.

* Presenting author. E-mail address: yr6755@g.skku.edu (Yeo Rang Cho)

S1.P95 A new phenolic glycoside from the leaves of *Salix chaenomeloides* and identification of anti- skin active compounds

Kyung Ah Kim^{a,*}, Sullim Lee^b, Ki Hyun Kim^a

^a School of Pharmacy, Sungkyunkwan University (SKKU), Suwon 16419, Republic of Korea

^b Department of Life Science, College of Bio-Nano Technology, Gachon University

Salix chaenomeloides Kimura, commonly known as Korean King willow, is a deciduous shrub and tree belonging to the Salicaceae family. It is native to eastern Asia, including regions of Korea, China, Japan, and Russia. For thousands of years, members of the genus *Salix* spp. have been recognized as healing herbs, used for the treatment of fever, inflammation, and pain relief (Freischmidt et al., 2012). The current study aimed to investigate the bioactive natural products from *S. chaenomeloides* leaves. Phytochemical investigation of an EtOH extract of *S. chaenomeloides* leaves led to the isolation of a new phenolic glycoside, chaenomelin (1) (Fig. 1), together with twelve known phenolic compounds (2–13) from the EtOAc fraction. The chemical structure of the new compound (1) was established by detailed analysis of its 1D- and 2D-NMR (¹H-¹H COSY, HSQC, and HMBC), HR-ESIMS, and chemical reactions.




Fig. 1. Structures of compounds 1 and 2.

The anti-skin aging effects of the isolates were examined using TNF- α -stimulated normal human dermal fibroblasts (NHDFs). Among these 13 compounds, tremuloidin (2) and tremulacin (3) inhibited TNF- α induced ROS generation ($p < 0.001$ at 100 μ M). Inhibition of ROS generation by tremuloidin (2) and tremulacin (3) led to the suppression of MMP-1 secretion and protection against collagen degradation. Particularly, the observed changes in TNF- α induced responses in NHDFs by tremuloidin (2) were found to correlate with the inhibition of COX-2, MAPK activation, and HO-1 induction. These findings offer evidence that tremuloidin (2) protects skin tissues from damage induced by oxidative stress accumulation.

Keywords: *Salix chaenomeloides*, phenolic compound, chaenomelin, tremuloidin, anti-skin activity

References

Freischmidt, A., Jürgenliemk, G., Kraus, B., Okpanyi, S.N., Müller, J., Kelber, O., Weiser, D., Heilmanna, J., 2012. Contribution of flavonoids and catechol to the reduction of ICAM-1 expression in endothelial cells by a standardised Willow bark extract. *Phytomedicine* 19, 245-252.

* Presenting author. E-mail address: ruddk5480@naver.com (Kyung Ah Kim)

S1.P96 Transcriptome mining, immunochemistry, and MSMS *de novo* sequencing – phosphorylated peptides from marine sponge *Stryphnus fortis*

Błażej Ślęzak^{a,b,*}, Lakmini Kosgahakumbura^{a,c}, Ulf Göransson^b, Chamari Hettiarachchi^c, Paco Cárdenas^{b,d}, Sunithi Gunasekera^b

^a W. Szafer Institute of Botany, Polish Academy of Science, 46 Lubicz St., 31-512, Kraków, Poland

^b Pharmacognosy, Department of Pharmaceutical Biosciences, Uppsala University, P.O. Box 591, 751 24 Uppsala, Sweden

^c Department of Chemistry, University of Colombo, Colombo 03, Sri Lanka

^d Museum of Evolution, Uppsala University, Norbyvägen 16, SE-752 36 Uppsala, Sweden

Stryphnus fortis (order: Tectractinellida, family: Ancorinidae) is a deep-sea sponge from the North Atlantic (Cárdenas, 2010). We could probably find cysteine-rich peptides in this sponge as it is closely related to the genus *Asteropus* from which such peptides – asteropsins – were isolated (Takada et al., 2006). Phosphorylation is one of proteins' most common types of posttranslational modifications (PM). However, it is rarely observed in cysteine-rich peptides and its physiological relevance remains elusive. Phosphorylation complicates the sequencing. It is difficult to observe in mass spectrometry as it is unstable and undergoes a process called beta elimination resulting in a loss of water (Tinette et al., 2007). It is also not apparent in standard NMR. In the current study, we describe the discovery of phosphorylated peptides from the marine sponge *S. fortis*. Moreover, it is an example of the application of transcriptome mining, immunochemistry, and mass spectrometry for the detection and characteristics of phosphorylation. In the initial experiments, we found two peptides with MW of 3330.385 Da and 3516.472 Da, named later named *P*-stryphnines A, and B. We established that they contain 6 cysteines. Knowing these characteristics and assuming similarity to asteropsins, we mined the transcriptome for sequences fitting the general pattern. These resulted in finding sequences matching theoretical MW -18 Da (Fig. 1). We confirmed the sequence and ruled out other PMs. The phosphorylation was detected using anti-phosphoserine/threonine antibody in a dot-blot experiment and the modification was pinpointed to S10 by MSMS *de novo* sequencing marine sponges, posttranslational modification.



Fig. 1. Sequence alignment of *P*-stryphnines A, and B Keywords: knottin,

References

- Cárdenas, P., 2010. Phylogeny, Taxonomy and Evolution of the Astrophorida (Porifera, Demospongiae).
- Takada, K., Hamada, T., Hirota, H., Nakao, Y., Matsunaga, S., van Soest, R.W.M., Fusetani, N., 2006. Asteropine A, a Sialidase-Inhibiting Conotoxin-like Peptide from the Marine Sponge *Asteropus simplex*. Chem. Biol. 13, 569–574.
- Tinette, S., Feyereisen, R., Robichon, A., 2007. Approach to systematic analysis of serine/threonine phosphoproteome using Beta elimination and subsequent side effects: intramolecular linkage and/or racemisation. J. Cell. Biochem. 100, 875–882.

* Presenting author. E-mail address: b.slazak@botany.pl (Błażej Ślęzak)

S1.P97 Distribution of cyanochelins, cyanobacterial beta-hydroxy aspartate lipopeptides capable of iron chelation

Štenclová, L.^{a,b,*}, Falcao, B. P.^{c,d}, Galica, T.^d, Martínez Yerena, J.A.^{b,c,d}, Mareš, J.^{b,d}, Hrouzek, P.^d

^a Institution of Pharmacy, Free University of Berlin, Königin-Luise-Str. 2+4, Berlin 14195, Germany

^b Institute of Hydrobiology, Biology Centre CAS, Na Sádkách 702/7, České Budějovice 37005, Czechia

^c Faculty of Science, University of South Bohemia, Branišovská 1760, České Budějovice 37005, Czechia

^d Institute of Microbiology, CAS, Novohradská 237, Třeboň, 37901, Czechia

Deficiency of biologically accessible iron e.g. in the arid-soil environment is one of the limiting factors for thriving of the microorganisms inhabiting such areas. Cyanobacteria as an important contributor in terms of primary production and excretion of bioactive compounds in the arid-soil- crust communities developed a coping mechanism, the secretion of siderophore compound enabling the ferric ions chelation. Cyanochelins, lipopeptides bearing two β -hydroxy-aspartate residues as chelating agents, were recently discovered in three phylogenetically distant cyanobacterial strains (Galica et al., 2021). Genome mining for biosynthetic gene clusters though indicates their widespread presence. Our metagenomic (Illumina) and secondary metabolite research (HPLC-MS) on environmental soil-crust samples discovered novel cyanochelin compounds and their putative producers from the genus *Gloeotheca* (Microcystaceae), as well as novel producers of known cyanochelin B from genus *Phormidesmis* (Leptolyngbyaceae). Research on our isolated producers and additional strains from algae collections showed wider distribution of cyanochelin gene clusters in genomes of those two groups of common inhabitants of soil-crust consortia. The production of another novel cyanochelin compound was confirmed e.g. in cyanobacteria in genus *Myxocorys* (Leptolyngbyaceae) and putative responsible biosynthetic gene clusters were identified. The intriguing feature of cyanochelins is the ability to undergo UV-dependent photolysis, release the bioavailable ferrous ions and presumably provide them to the microbial community, which together with their distribution suggest their potential ecological importance. Bioinformatic analyses of our metagenomes, targeted at cyanochelin transporters, discovered that the recipients could occur among both cyanobacterial and other bacterial taxa. In ongoing cocultivation experiments, characterisation of cyanochelin-dependent cohabitants can tell us more about processes of iron cycling in the soil- crusts of arid regions.

Keywords: arid soils, cyanobacteria, cyanochelins, iron, siderophores

References

Galica T, Borbone N, Mareš J, Kust A, Caso A, Esposito G, Saurav K, Hájek J, Řeháková K, Urajová P, Costantino V, Hrouzek P. 2021. Cyanochelins, an overlooked class of widely distributed cyanobacterial siderophores, discovered by silent gene cluster awakening. *Appl Environ Microbiol* 87, e03128-20.

* Presenting author. E-mail address: stenclova.lenka@gmail.com (Lenka Štenclová)

S1.P98 Syringin and phillygenin - natural compounds with a potential role in preventing of foam cells formation in human atherosclerotic plaque

Agnieszka Filipek*, Małgorzata Kołtun-Jasion, Agnieszka Sadowska, Anna Kiss

Department of Pharmaceutical Biology, Faculty of Pharmacy, Medical University of Warsaw, Banacha 1, 02-097 Warsaw, Poland

The presentation will cover an explanation of the biological properties of two natural compounds: syringin and phillygenin in inhibit cholesterol-induced differentiation of macrophage into foam cells (CD36 receptor) and increase cholesterol efflux from macrophage (ABCA1 transporter) (Chistiakov et al. 2016). Syringin is a phenylpropanoid glycoside isolated from the bark of *Syringa vulgaris*, (Oleaceae). Phillygenin is a lignan obtained mainly from the fruits and flowers of *Forsythia x intermedia* (Oleaceae). In previous studies, both compounds showed strong anti-inflammatory and antioxidant properties (Filipek, et al. 2019). Syringin and phillygenin were isolated from plants at the Department of Pharmaceutical Biology of the Medical University of Warsaw. Human monocyte- derived macrophages were incubated (24h) with syringin or phillygenin (10 mM, 20 mM, 50 mM) and cholesterol (20 mM). Oil-red-O staining showed that phillygenin and syringin in dose-dependent manner significantly ($p<0.001$) reduced lipid deposits in macrophages. Cytometric analysis showed that for syringin, the greatest decrease in CD36 receptor expression was determined, from 20% (10 μ M) to 80% (50 μ M) compared to the cholesterol-stimulated control ($p<0.001$). Phillygenin inhibited the expression of the CD36 receptor, from 5% (10 μ M) to 25% (50 μ M), compared to the stimulated control ($p<0.05$). The CD36 receptor down-regulation pathway was PPR-g-dependent ($p<0.05$). Western Blot analysis showed that phillygenin caused a statistically significant ($p<0.001$) increase in the expression of the ABCA1 receptor by 2.5-fold (10 μ M), 3-fold (20 μ M) and 4-fold (50 μ M), respectively, compared to the cholesterol-stimulated control. Also, in the case of syringin, a statistically significant increase (50%) in the expression of ABCA1 transporter proteins was observed. The ABCA1 transporter activation pathway was HO-1-dependent ($p<0.05$).

Keywords: syringin, phillygenin, foam cells, CD36, ABCA1

References

- Chistiakov, D.A., Bobryshev, Y.V., Orekhov, A.N. 2016. Macrophage-mediated cholesterol handling in atherosclerosis. J Cell Mol Med. Vol. 20, 17-28.
- Filipek, A., Wyszomierska, J., Michalak, B., Kiss A.K. 2019. *Syringa vulgaris* bark as a source of compounds affecting the release of inflammatory mediators from human neutrophils and monocytes/ macrophages. Phytochemistry Letters. Vol. 30, 309-313.

* Presenting author. E-mail address: agnieszka.filipek@wum.edu.pl (Agnieszka Filipek)

S1.P99 Deciphering monoterpene indole alkaloid metabolism in *Heliconius* butterfly: a MS-based computational chemistry to describe plant-host interactions

Inès Houari^{a,b,*}, Coralie Audoin^a, Grégory Genta-Jouve^b

^a Laboratoires Clarins, 5 rue Ampères, Pontoise 95300, France

^b Laboratoire Écologie, Évolution, Interaction des Systèmes Amazoniens (LEEISA), 275 route de Montabo, 97300 Cayenne, Guyane française

French Guiana renowned for its rich tropical biodiversity, notably exhibits a remarkable interplay between its Amazon vegetation and preserved fauna. Within this intricate ecosystem, butterflies of the *Heliconius* genus establish a distinct association with select plant species (See passion vines, Morrison et al., 2023). Extracts from *Heliconius* butterflies provided a unique dataset for developing a metabolite identification workflow through computational chemistry. Employing liquid chromatography–high-resolution tandem mass spectrometry, a molecular network was constructed. Thanks to the Lotus library (Rutz et al., 2022), in-silico spectral libraries were generated, highlighting compounds found in *Heliconius* like the notable presence of harmalan (**1**). Employing this library failed to produce results, but upon including **1** in the metadata folder, we surprisingly observed its clustering with three Monoterpene Indole Alkaloids (MIAs). Specifically targeting one of them, ajmalicine (**2**), identified with Sirius and In Silico DataBase (ISDB), characteristic fragments of this molecule were identified and a high cosine score similarity during annotation was observed. Additionally, **1** comprises a MIAs' characteristic 144.0808 m/z fragment shared with **2** (Nakabayashi et al., 2020). The presence of such molecules, although absent in *Passiflora* plants (Yang Ning et al., 2023), leads us to consider the sequestration of metabolites through the caterpillar's nutrition on a different plant. Consequently, we investigated native plants from Guyana known to contain MIAs, and identified *Uncaria guianensis* and *Rauvolfia tetraphylla* as potential host plants. This fascinating prospect of such interaction also suggest a possibility of de novo biosynthesis.



Fig. 1. Structure of harmalan **1**, ajmalicine **2** and a photo of *Heliconius erato*

Keywords: *Heliconius* butterflies, molecular networking, plant-host interactions, MIAs

References

- Morrison, C.R., C. Nguyen, C., Gilbert, L.E., 2023. The role of host plant usage and the accumulation of toxic secondary chemical compounds across the life cycle of a passion vine specialist butterfly. *Ecol. Ent.* 48, 389–395.
- Nakabayashi, R., Mori, T., Takeda, N., Toyooka, K., Sudo, H., Tsugawa, H., Saito, K., 2020. Metabolomics with ¹⁵N Labelling for Characterizing Missing MIAs in Plants. *Anal. Chem.* 92, 5670–5675.
- Rutz, A., Sorokina, M., Algonac, J., Mietchen, D., Willighagen, E., Gaudry, A., Graham, J.G., Stephan, R., Page, R., Vondrášek, J., Steinbeck, C., Pauli, G.F., Wolfender, J.-L., Bisson, J., Allard P.-M., 2022. The LOTUS initiative for open knowledge management in natural products research. *eLife* 11, e70780.
- Yang Ning, L., Rahman Azmi, A.A.A., Syamsumir, D.F., Ismail, W.I.W., Maulidiani, M., 2023. Phytochemical screening, TLC profile and ¹H NMR analysis of *Passiflora foetida* extracts. *UMT JUR.* Vol 5, No 2, 65–74.

* Presenting author. E-mail address: ines.houari@clarins.com (Inès Houari)

S1.P100 Aporphine alkaloids from *Phaeanthus lucidus* Oliv. and their α -glucosidase inhibitory activity

Passakorn Teerapongpisan^a, Virayu Suthiphasilp^b, Pakit Kumboonma^c, Tharakorn Maneerat^{a,d}, Thidarat Duangyod^{d,e}, Rawiwan Charoensup^{d,e}, Phunrawie Promnart^f, Raymond J. Andersen^g, Surat Laphookhieo^{a,d,*}

^a Center of Chemical Innovation for Sustainability (CIS) and School of Science, Mae Fah Luang University, Chiang Rai, 57100, Thailand

^b Department of Industrial Technology and Innovation Management, Faculty of Science and Technology, Pathumwan Institute of Technology, Bangkok, 10330, Thailand

^c Department of Applied Chemistry, Faculty of Science and Liberal Arts, Rajamangala University of Technology Isan, Nakhon Ratchasima 30000, Thailand

^d Medicinal Plant Innovation Center of Mae Fah Luang University, Chiang Rai, 57100, Thailand

^e School of Integrative Medicine, Mae Fah Luang University, Chiang Rai, 57100, Thailand

^f School of Science, Mae Fah Luang University, Chiang Rai, 57100, Thailand

^g Departments of Chemistry and Earth, Ocean & Atmospheric Sciences, University of British Columbia, 2036 Main Mall, Vancouver, BC V6T 1Z1, Canada

Phaeanthus lucidus Oliv. is a medium-sized tree distributed in Southern Thailand and Peninsular Malaysia. To our knowledge, there have been no reports on phytochemical investigations and biological activities from *P. lucidus*. This presentation will cover the isolation, structure elucidation, and α -glucosidase inhibitory activity of compounds isolated from the twig and leaf extracts of *P. lucidus*. In addition, molecular docking simulations of α -glucosidase inhibition of active compounds were included in this presentation. The twig and leaf extracts of *P. lucidus* were fractionated by various column chromatography to yield 14 compounds and seven of them were identified as new aporphine alkaloids (Teerapongpisan et al., 2023; Teerapongpisan et al., 2024). Their structures were elucidated using extensive NMR spectroscopy and HRESIMS. Compounds 1-3 were analyzed and resolved by chiral HPLC to yield the (Ra) and (Sa) atropisomers, whose absolute configurations were determined by ECD calculations. Compound 1 showed the best α -glucosidase inhibitory activities with IC₅₀ values of 6.7 μ M.




Fig. 1. Structures of new compounds isolated from *P. lucidus*.

Keywords: *Phaeanthus lucidus*, Annonaceae, α -glucosidase inhibitory, Aporphine alkaloids

References

- Teerapongpisan, P., Suthiphasilp, V., Kumboonma, P., Maneerat, T., Duangyod, T., Charoensup, R.; Andersen, R.J., Laphookhieo, S. 2023. Phaeanthuslucidines A-D, dimeric aporphine alkaloid derivatives from *Phaeanthus lucidus* Oliv. *Phytochemistry* 212, 113717.
- Teerapongpisan, P., Suthiphasilp, V., Kumboonma, P., Maneerat, T., Duangyod, T., Charoensup, R.; Promnart, P., Laphookhieo, S. 2024. Phaeanthuslucidines A-D, dimeric aporphine alkaloid derivatives from *Phaeanthus lucidus* Oliv. *Phytochemistry* 220, 114020.

* Presenting author. E-mail address: surat.lap@mfu.ac.th (Surat Laphookhieo)

S1.P101 Tychonemazole, a linear depsipeptide, from the cyanobacterium *Tychonema bourrellyi*

Nandani Balloo^a, Tatsufumi Okino^{a,b,*}

^a Graduate School of Environmental Science, Hokkaido University, Kita10-Nishi5, Kita-ku, 060-0810 Sapporo, Japan

^b Faculty of Environmental Earth Science, Hokkaido University, Kita10-Nishi5, Kita-ku, 060-0810 Sapporo, Japan

Many sponge-derived compounds are structurally similar to compounds isolated from cyanobacteria. Some are synthesized by symbiotic cyanobacteria in sponges, but some are derived from different bacteria. Citronamides A and B were isolated from the sponge *Citronia astra*, which contains many symbiotic cyanobacterial cells (Carroll et al., 2009). During our OSMAC approach for cyanobacteria, we discovered similar compounds to citronamides from the cyanobacterium *Tychonema bourrellyi*. Tychonemazole (**1**) was found as a major compound even in a culture condition without iron supplementation. Therefore, we attempted to isolate **1** by LC-MS-guided isolation approach. Analyses by 2D NMR including 1,1-ADEQUATE revealed a structure of linear depsipeptide, **1** containing Ile, Thr, glutaric acid, hydroxy oxo-histidine, an amino methoxyphenyl heptanoic acid moiety. The latter two moieties are unique residues in the compounds related to citronamides. A configurational analysis of these moieties is under progress.




Fig. 1. Structure of tychonemazole **1**.

Keywords: cyanobacteria, sponge, 1,1-ADEQUATE, oxo-histidine, OSMAC

Reference

Carroll, A.R., Duffy, S., Avery, V.M., 2009. Citronamides A and B, tetrapeptides from the Australian sponge *Citronia astra*. J. Nat. Prod. 72, 764-768.

* Presenting author. E-mail address: okino@ees.hokudai.ac.jp (Tatsufumi Okino)

S1.P102 Isocoumarin derivatives from cannabis-associated endophytic fungus *Phialemonium* sp. SUNP1021

Se Yun Jeong*, Seo Yoon Le, Ki Hyun Kim^a

School of Pharmacy Sungkyunkwan University, Suwon 16419, Republic of Korea

Cannabis sativa is renowned for its psychoactive metabolites. Because of its regulation in Korea, our attention shifted to the endophytic fungus *Phialemonium* sp. (Cephalothecaceae), isolated from *C. sativa* since we anticipated that this fungus would produce novel metabolites, as determined by LC/MS analysis. *Phialemonium* sp., a well-known endophytic fungus (Su et al., 2016) has not been extensively investigated for its metabolites. The chemical investigation of the methanol extract of *Phialemonium* sp. cultivated in cannabis powder dextrose agar (CDA) resulted in the isolation and identification of 12 isocoumarin derivatives (**1–12**), facilitated GNPS analysis. The planar structures of novel isocoumarins (**1–8**) were established by comprehensive analysis of 1D and 2D (¹H-¹H COSY, HSQC, and HMBC) NMR spectra and high-resolution electrospray ionization mass spectrometry (HR-ESI-MS) data. Their absolute configurations were fully established by computational methods, including DP4+ analysis and ECD calculations. Especially, the chemical structure of compound **8** was validated through X-ray crystallography. The biological activities of isolated compounds will be further evaluated.




Fig. 1. Structure of compound **8** and its X-ray structure.

Keywords: *Cannabis sativa*, cannabis-associated endophytic fungus, *Phialemonium* sp., Cephalothecaceae, isocoumarin

References

Su, L., Deng, H., Niu, Y. C., 2016. *Phialemoniopsis endophytica* sp. nov., a new species of endophytic fungi from *Luffa cylindrica* in Henan, China. Mycol. Prog. 15, 1-7.

* Presenting author. E-mail address: jseyun12@gmail.com (Se Yun Jeong)

S1.P103 Novel polyketide metabolites, skeletocutins M–Q, from *Tyromyces chioneus* and their antibacterial activity against *Staphylococcus aureus*

Se Yun Jeong*, Ki Hyun Kim

School of Pharmacy Sungkyunkwan University, Suwon 16419, Republic of Korea

Tyromyces chioneus (Polyporaceae) is widely distributed throughout temperate boreal pine forests across Asia, Europe, and North America (Chander et al., 2018). The extract of *T. chioneus* has been reported to possess potent antimicrobial activity against pathogenic microorganisms and notable antioxidant properties (Manjusha et al., 2020). Nonetheless, the bioactive metabolites of *T. chioneus* have not been thoroughly investigated. The skeletocutins from wood-inhabiting Basidiomycete, *Skeletocutis* sp. (Polyporaceae) have been reported to show antimicrobial activity against Gram- positive bacteria and hepatitis C virus (Chepkirui et al., 2019). As part of our extensive research on Korean wild mushrooms, a chemical investigation of ethyl acetate extract of *T. chioneus* cultivated in potato dextrose broth (PDB) resulted in the isolation and identification of 8 polyketide metabolites (1–8). The chemical structures of novel compounds (1–5), skeletocutins M–Q, were fully established via detailed analysis of 1D and 2D (¹H-¹H COSY, HSQC, and HMBC) NMR spectra and HR-ESIMS data.



Fig. 1. Structure of compound 4

The antibacterial activity of isolated polyketide metabolites (1–8) against *Staphylococcus aureus* was assessed by determining the inhibitory effect of PepZ leucine aminopeptidase which is crucial for the survival of *S. aureus* inside the host cell. All the tested compounds inhibited the PepZ leucine aminopeptidase of *S. aureus*. Specifically, compounds 4 and 5 displayed the most potent inhibition of PepZ. These findings demonstrate that skeletocutins identified from *T. chioneus* can selectively kill intracellular *S. aureus* with minimal perturbation of normal microbiota.

Keywords: *Tyromyces chioneus*, skeletocutins, antibacterial activity, leucine aminopeptidase, *Staphylococcus aureus*

References

- Chander, H., Pathania, J., 2018. Notes on occurrence of anti HIV-1 medicinal macrofungi *Tyromyces chioneus* in Hamirpur district, Himachal Pradesh. CPUH-Res. J. 3, 187-190.
- Chepkirui, C., Cheng, T., Sum, W.C., Matasyoh, J.C., Decock, C., Praditya, D.F., Wittstein, K., Steinmann, E., Stadler, M., 2019. Skeletocutins A-L: Antibacterial agents from the Kenyan wood-inhabiting Basidiomycete, *Skeletocutis* sp.. J. Agric. Food Chem. 67, 8468-8475.
- Manjusha, P.G.S., 2020. Molecular identification and isolation of antimicrobial, phytochemical, antioxidant properties of different solvent extracts from wild mushrooms. Saudi J. Pathol. Microbiol. 5, 98-107.

* Presenting author. E-mail address: jseyun12@gmail.com (Se Yun Jeong)

S1.P104 praca wycofana

S1.P105 Usnic acid enantiomers in the lichen-forming fungal family Parmeliaceae

Maonian Xu^{a,*}, Garima Singh^b, Starri Heidmarsson^c, Elín S. Ólafsdóttir^a

^a Faculty of Pharmaceutical Sciences, University of Iceland, Hofsvallagata 53, 107 Reykjavik, Iceland

^b Department of Biology, University of Padova, Via U. Bassi, 58/B, 35121 Padova, Italy

^c Northwest Iceland Nature Research Centre, 550 Sauðárkrúkur, Iceland

Enantiomers of usnic acid in lichens have potent antimicrobial activity, and the lichens in the genus *Usnea* (Parmeliaceae, Ascomycota) containing the (+)-usnic acid have been used as herbal medicines (Xu et al., 2016). However, there are multiple genera (e.g. *Flavocetraria*, *Alectoria*, etc.) in the family Parmeliaceae producing usnic acid, but they were not selected for medicinal uses. We hypothesize that they may produce the other enantiomer – (-)-usnic acid that might be less favorable than the (+)-usnic acid in bioactivities. The study aimed to characterize the usnic acid enantiomeric composition in the lichen-forming fungal family Parmeliaceae in a fungal phylogenetic context. A validated chiral HPLC method was used for the enantiomeric separation of enantiomers (Xu et al., 2022), and fungal phylogenetic tree was constructed using three fungal genetic markers (i.e. ITS, MCM7 and RPB1).




Fig. 1. Structures of usnic acid enantiomers: (-)-usnic acid (left) and (+)-usnic acid (right)

Our preliminary data show interesting phylogenetic patterns of usnic acid enantiomeric production: the cetrarioid clade is rich in (-)-usnic acid, while the parmelioid clade in (+)-usnic acid. We also performed phylogenetic analyses on the biosynthetic genes and proposed the biosynthetic pathways for both enantiomers. Ongoing antimicrobial assay tests indicate that the (+)-usnic acid is more potent than (-)-usnic acid in antibacterial activity.

Keywords: usnic acid, stereoisomers, lichens, Parmeliaceae

References

- Xu M, Oppong-Danquah E, Wang X, Oddsson S, Abdelrahman A, Pedersen SV, Gylfason AE, Snorraddottir BS, Christensen A, Tasdemir D, Jameson CJ, Murad S, Andersson OS, Magnusson KP, de Boer H, Thorsteinsdottir M, Omarsdottir S, Heidmarsson S, Olafsdottir ES. 2022. Novel methods to characterise spatial distribution and enantiomeric composition of usnic acids in four Icelandic lichens. *Phytochemistry*. 200:113210.
- Xu M, Heidmarsson S, Olafsdottir ES, Buonfiglio R, Kogej T, Omarsdottir S. 2016. Secondary metabolites from cetrarioid lichens: chemotaxonomy, biological activities and applications. *Phytomedicine* 23:441-459.

* Presenting author. E-mail address: maonian@hi.is (Maonian Xu)

S1.P106 The potential of Amazonian microorganisms in drug discovery

Ivan Lucas de Freitas Migliorini^a, Ana Ligia Leandrini de Oliveira^b, Mônica Tallarico Pupo^{a,*}

^a Department of Pharmaceutical Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Av. do Café, s/n, 14040-903 Ribeirão Preto - SP, Brazil

^b School of Pharmaceutical Sciences, Federal University of Amazonas, Av. General Rodrigo Otávio, 6200, 69077-000 Manaus - AM, Brazil

The discovery of antibiotics from soil microorganisms in the mid-20th century revolutionized medicine, with recent research underscoring continued interest in natural products, exemplified by the discovery of the antibiotic teixobactin (Ling et al, 2015). In parallel, the Amazon's rich and untapped biodiversity faces threats from climate change and deforestation, stimulating further studies. As part of our continuous interest in microbial natural products from different Brazilian biomes, we have collected soil specimens from the Amazon Forest in the Anavilhanas National Park. Ninety Actinobacterial strains were isolated, and 24% of them showed bioactivity in antimicrobial assays against *Candida albicans*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. Among these, Actinomycetota AMYS3.17 was selected for chemical studies, leading to the annotation of eurocidins D (1) and E (2). (Fig. 1). Eurocidins have been relatively underexplored in research, with only eurocidins D (1) and E (2) being extensively characterized. Molecular networking analysis (GNPS) (Wang et al, 2016) indicates that bacterial strain AMYS3.17 can produce over 20 compounds resembling eurocidin, including novel analogues putatively annotated through mass spectrometry. This highlights the significant potential of this microorganism for discovering new polyene compounds.




Fig. 1. Structures of eurocidin D (1) and eurocidin E (2)

Keywords: Actinomycetota, soil, microorganisms, natural products, Amazon Forest

References

Ling, L.L., Schneider, T., Peoples, A.J., Spoering, A.L., Engels, I., et al, 2015. A new antibiotic kills pathogens without detectable resistance. *Nature* 517, 455-459.

Wang, M., Carver, J.J., Phelan V.V., Sanchez L.M., Garg N., et al, 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat. Biotechnol.* 34, 828-837.

* Presenting author. E-mail address: mtpupo@fcrp.usp.br (Mônica T. Pupo)

S1.P107 From kitchen to lab: unveiling methyl 5-vinylnicotinate and other compounds formation during cooking with olive oil

Aristea Deli, Evangelos Dadiotis, Eleni Melliou*, Prokopios Magiatis

Laboratory of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Greece

The current study investigated the fate of the phenolic compounds found in extra virgin olive oil (EVOO) during common Mediterranean cooking practices, specifically sautéing and frying. A range of plant-based (onion, garlic, pepper, mushroom, carrot) and animal-based (cheese, bacon, egg) foods were fried in varying food-to-oil ratios to assess this impact. Key findings include the formation of a rare pyridine alkaloid, methyl 5-vinylnicotinate, identified through ^1H -NMR and MS techniques for the first time as a food ingredient. The compound was isolated from the olive oil matrix after cooking using low-pressure adsorption chromatography and preparative thin-layer chromatography. Further, a potential formation mechanism was proposed, involving the reaction of EVOO's phenolic compounds oleomissional or/and oleokoronal with nitrogen sources in the food, catalyzed by heat (Fig. 1). The semi-synthesis of methyl 5-vinylnicotinate was also performed using two methods: one involving the reaction of the dry polar extract of extra virgin olive oil with ammonium hydroxide, and the other one using pure oleomissional, extracted from olive tree leaves, also reacting with ammonium hydroxide. The synthesized compound was purified and isolated, highlighting a novel synthetic methodology. Additionally, hydroxytyrosol acetate and tyrosol acetate were produced during the above reaction and were correlated for the first time with cooking of EVOO.




Fig. 1. Methyl 5-vinylnicotinate and hydroxytyrosol acetate formation during frying with EVOO rich in oleomissional.

This research provides crucial insights into the chemical transformations occurring in Mediterranean cooking with EVOO, opening avenues for understanding the health impacts of these culinary practices. Future research will focus on evaluating the potential benefits or risks associated with methyl 5-vinylnicotinate consumption.

Keywords: EVOO, methyl 5-vinylnicotinate, Mediterranean cuisine, olive oil phenols

* Presenting author. E-mail address: emelliou@pharm.uoa.gr (Eleni Melliou)

S1.P108 C15 acetogenins and sesquiterpenes with anti-inflammatory activity from red algae of the genus *Laurencia* collected in the Red Sea

Mohamed A. Tammam^{a,b}, Aikaterini Koutsaviti^a, Susana Agusti^c, Carlos M. Duarte^c, Efstathia Ioannou^a, Vassilios Roussis^{a,*}

^a Section of Pharmacognosy and Chemistry of Natural Products, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771 Athens, Greece

^b Department of Biochemistry, Faculty of Agriculture, Fayoum University, 63514 Fayoum, Egypt

^c Red Sea Research Center (RSRC), King Abdullah University of Science and Technology (KAUST), 23955-6900 Thuwal, Saudi Arabia

Among red algae, the genus *Laurencia* (Rhodomelaceae), comprising approx. 140 accepted species distributed in tropical, subtropical and temperate coastal waters, is one of the richest sources of new secondary metabolites in the marine environment (Harizani et al., 2016). Up to now, almost 1,200 secondary metabolites, mostly halogenated C15 acetogenins and terpenes often displaying unprecedented carbon skeletons, have been reported from *Laurencia* sp. and mollusks of the genus *Aplysia* that feed on them. Among the isolated metabolites, a high number has displayed antiviral, anti-inflammatory, antibacterial, antifungal, cytotoxic, antiproliferative, antifeedant, ichthyotoxic, antifouling and insecticidal activity (MarinLit, 2024). In the framework of our investigations, aiming for the isolation of new bioactive marine metabolites, we studied the chemical profiles of two populations of *Laurencia* collected from Hurghada reef in Egypt and Rose reef off Thuwal in Saudi Arabia. Algal specimens were extracted with mixtures of CH₂Cl₂/MeOH and the organic residues were submitted to a series of chromatographic separations, leading to the isolation of 32 different secondary metabolites, including 12 new C15 acetogenins and 5 new sesquiterpenes. The structures and the relative configurations of the isolated natural products were established on the basis of extensive analysis of their NMR and MS data, while the absolute configuration was determined by single-crystal X-ray diffraction analysis. The anti-inflammatory activity of the isolated compounds was evaluated by measuring suppression of nitric oxide release in TLR4-activated RAW 264.7 macrophages in culture.

Keywords: marine natural products, *Laurencia* sp., red algae, anti-inflammatory activity

References

Harizani, M., Ioannou, E., Roussis, V., 2016. The *Laurencia* Paradox: An Endless Source of Chemodiversity. In *Progress in the Chemistry of Organic Natural Products*, Volume 102. Kinghorn, A.D., Falk, H., Gibbons, S., Kobayashi, J. (Eds). Springer International Publishing: Cham, Switzerland, pp. 91–252.

MarinLit. A Database of the Marine Natural Products Literature. Available online: <http://pubs.rsc.org/marinlit/> (accessed on February 2024).

* Presenting author. E-mail address: roussis@pharm.uoa.gr (Vassilios Roussis).

S1.P109 Diterpenes featuring the spatane, seco-spatane and prenylcubebane carbocycles from the invasive brown alga *Rugulopteryx okamurae*

Dikaia Varla^a, Aikaterini Koutsaviti^a, Kyriaki Papadopoulou^a, Susana Agusti^b, Carlos M. Duarte^b, Vassilios Roussis^a, Olga Tzakou^a, Efstathia Ioannou^{a,*}

^a Section of Pharmacognosy and Chemistry of Natural Products, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771 Athens, Greece

^b Red Sea Research Center (RSRC), King Abdullah University of Science and Technology (KAUST), 23955-6900 Thuwal, Saudi Arabia

Brown algae of the family Dictyotaceae, widely distributed in the tropical and subtropical regions of the Atlantic, Pacific and Indian Ocean, in the Sea of Japan, as well as in the Mediterranean Sea, are a prolific source of natural products. Approximately 700 secondary metabolites, mainly sesquiterpenes, diterpenes and meroterpenoids, have been isolated from taxa of this algal family, exhibiting a wide spectrum of biological activities, such as cytotoxic, antibacterial, antifungal, antiviral, antifouling, antiparasitic and anti-inflammatory activity (MarinLit, 2024). *Rugulopteryx okamurae* is an invasive member of this family, causing severe environmental and economic problems on the western Mediterranean coasts. Therefore, research is mandated in order to reduce the biomass and transform a problem into an opportunity for obtaining new valuable products. In this context, we investigated the chemical profiles of two populations of *R. okamurae* collected from deep and shallow waters from the coasts of Cadiz. Following extraction of the algal biomass, the organic residues were submitted to numerous chromatographic separation steps to afford up to now 25 metabolites, mostly diterpenes featuring the spatane, seco-spatane and prenylcubebane skeletons. Among them, 7 diterpenes are new natural products. The structures and relative configurations of the isolated metabolites were elucidated on the basis of extensive analyses of their NMR and MS spectroscopic data. The evaluation of the bioactivity of the isolated compounds is currently ongoing.

Keywords: marine natural products, *Rugulopteryx okamurae*, brown algae, bioactivity

References

MarinLit. A Database of the Marine Natural Products Literature. Available online: <http://pubs.rsc.org/marinlit/> (accessed on February 2024).

* Presenting author. E-mail address: eioannou@pharm.uoa.gr (Efstathia Ioannou)

S1.P110 Metabolic and biosynthetic potential of microbes isolated from North American snake venom

Katherine L. Lev^{a,b,*}, Matthew Holding^{a,f}, Anders Kiledal^g, Meredith Holm^{a,e}, David H. Sherman^{a,b,c,d,e}

^a Life Sciences Institute, University of Michigan, Ann Arbor, MI, USA

^b Program in Chemical Biology, University of Michigan, Ann Arbor, MI, USA

^c Department of Medicinal Chemistry, University of Michigan, Ann Arbor, MI, USA

^d Department of Microbiology & Immunology, University of Michigan, Ann Arbor, MI, USA

^e Department of Chemistry, University of Michigan, Ann Arbor, MI, USA

^f Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA

^g Department of Earth and Environmental Sciences, University of Michigan, Ann Arbor, MI, USA

Animal venoms are a well-established source of pharmacologically active peptides, which has led to a long-standing assumption that venom glands are sterile micro-environments (Calvete et al., 2009). However, advances in genome sequencing and bioinformatics tools have stimulated renewed interest in exploring venom microbiota (Esmaeilshirazifard et al., 2022). Moreover, new efforts seek to probe the diversity of venom-derived microbes and their evolutionary responses (i.e., antimicrobial compound production) to life in venom-associated organs (Ul-Hasan et al., 2019). Due to extreme conditions, including low nutrient availability, low pH, and the presence of antimicrobial peptides, venom microbes are hypothesized to harbor uniquely untapped biosynthetic potential as an evolutionary response to their hostile environment. We collected venom samples from rattlesnake species found in the Southwestern United States. Venom was collected via a standard sterile technique. We obtained control samples of oral microbiota to compare community composition. Liquid venom and oral swabs were frozen prior to transport on dry ice. Metagenomic DNA was extracted and sequenced via Illumina short-read sequencing. Following this, metagenome-assembled genomes (MAGs) were binned and annotated using an in-house bioinformatic pipeline. Additionally, venom samples were diluted in sterile H₂O before plating on agar plates and in liquid culture in varying nutrient media. Resulting mixed bacterial cultures with distinct morphological differences suggested many microbial lineages inhabiting snake venom. Genomic DNA from individual colonies was isolated for Illumina short-read sequencing. Using bioinformatics tools, we mined assembled genomes of our bacterial isolates to predict biosynthetic gene clusters (BGCs) that encode production of specialized metabolites.

Keywords: microbiome, venom, genomics, metabolomics, blood-clotting

References

- Calvete, J.J., Sanz, L., Angulo, Y., Lomonte, B., Gutiérrez, J.M., 2009. Venoms, Venomics, Antivenomics. *FEBS Lett.* 583, 1736–1743.
- Esmaeilshirazifard, E., Usher, L., Trim, C., Denise, H., Sangal, V., Tyson, G. H., Barlow, A., Redway, K. F., Taylor, J. D., Kremyda-Vlachou, M., Davies, S., Loftus, T. D., Lock, M. M. G., Wright, K., Dalby, A., Snyder, L. A. S., Wuster, W., Trim, S., Moschos, S. A., 2022. Bacterial Adaptation to Venom in Snakes and Arachnida. *Microbiology Spectrum* 0, e02408-21.
- Ul-Hasan, S., Rodríguez-Román, E., Reitzel, A. M., Adams, R. M. M., Herzig, V., Nobile, C. J., Saviola, A. J., Trim, S. A., Stiers, E. E., Moschos, S. A., Keiser, C. N., Petras, D., Moran, Y., Colston, T. J., 2019. The Emerging Field of Venom- Microbiomics for Exploring Venom as a Microenvironment, and the Corresponding Initiative for Venom Associated Microbes and Parasites (IVAMP). *Toxicon* 4, 100016.

*Presenting author. Email address: katlev@umich.edu (Katherine Lev)

S1.P111 Functionalized diterpenes featuring the cembrane skeleton from soft corals of the genus *Sarcophyton* collected in the Red Sea

Mohamed A. Tammam^{a,b}, Stefanos Kikionis^{a,*}, Susana Agusti^c, Carlos M. Duarte^c, Vassilios Roussis^a, Efstathia Ioannou^a

^a Section of Pharmacognosy and Chemistry of Natural Products, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771 Athens, Greece

^b Department of Biochemistry, Faculty of Agriculture, Fayoum University, 63514 Fayoum, Egypt

^c Red Sea Research Center (RSRC), King Abdullah University of Science and Technology (KAUST), 23955-6900 Thuwal, Saudi Arabia

On a global scale, coral reefs are regarded as the most spectacular and diverse marine ecosystems on the planet, hosting hundreds of thousands of species, many of which are currently undescribed. Among them, the coral reefs of the Red Sea are considered the best-developed reefs in the western Indian Ocean. Soft corals of the genus *Sarcophyton* (Sarcophytidae) have been proven to be a prolific source of bioactive natural products, having afforded more than 850 metabolites to date (MarinLit, 2024). They are especially rich in sesquiterpenes, diterpenes, steroids and ceramides, exhibiting a wide range of biological activities, such as antimicrobial, antiviral, antidiabetic, anti-inflammatory, neuroprotective and antitumor activities (Elkhawas et al., 2020). In the course of our investigations targeting the detection and isolation of new marine metabolites with bioactivity, we explored the chemical profiles of two morphologically distinct populations of *Sarcophyton* collected by SCUBA diving from the coral reefs near Al Lith on the south-west coast of Saudi Arabia. After exhaustive extraction of the fresh animal tissues with mixtures of CH₂Cl₂/MeOH, the organic residues underwent laborious chromatographic separations to yield 38 different secondary metabolites, mostly cembranoids, in pure form. Among them, 7 are new natural products. The structures and the relative configurations of the isolated natural products were determined on the basis of thorough analysis of their NMR and MS data. When possible, the absolute configuration was established by single-crystal X-ray diffraction analysis. The evaluation of the biological activity of the isolated compounds is currently in progress.

Keywords: marine natural products, *Sarcophyton* sp., soft corals, bioactivity

References

MarinLit. A Database of the Marine Natural Products Literature. Available online: <http://pubs.rsc.org/marinlit/> (accessed on February 2024).

Elkhawas, Y.A., Elissawy, A.M., Elnaggar, M.S., Mostafa, N.M., Al-Sayed, E., Bishr, M.M., Singab, A.N.B., Salama, O.M., 2020. Chemical diversity in species belonging to soft coral genus *Sarcophyton* and its impact on biological activity: A review. *Mar. Drugs* 18, 41.

* Presenting author. E-mail address: skikionis@pharm.uoa.gr (Stefanos Kikionis)

S1.P112 Neuroprotective activity of *Swertia iberica* (Gentianaceae) on BV-2 microglia cells

Fatmanur Tunç^{a,*}, Kübra Uzun^b, Funda Nuray Yalçın^c, Yavuz Bülent Köse^a

^a Department of Pharmaceutical Botany, Faculty of Pharmacy, Anadolu University, 26470, Tepebaşı, Eskişehir, Türkiye

^b Department of Pharmacognosy, Faculty of Pharmacy, Erciyes University, 38280, Talas, Kayseri, Türkiye

^c Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University, 06230, Altındağ, Ankara, Türkiye

The *Swertia* L., belonging to the Gentianaceae family, includes 166 species worldwide (http-1), and 2 species growing in Türkiye (http-2). *Swertia* exhibits a various pharmacological activities, including anticancer, antipyretic, analgesic, antidiabetic, antibacterial, antioxidant, hepatoprotective, and neuroprotective. *Swertia* species contain xanthenes, iridoids/secoiridoids, flavonoids, alkaloids, and terpenoids (Kshirsagar et al., 2019). The aerial parts of *S. iberica* were extracted with methanol and partitioned by *n*-hexane, *n*-butanol, and water, respectively. The *n*-butanol extract was chromatographed over a polyamide column eluting with stepwise H₂O–MeOH gradient to obtain fractions. Alzheimer's disease, a progressive neurodegenerative disease, affects learning and memory resulting from cholinergic dysfunction. Scopolamine has been employed to induce Alzheimer's disease-like pathology *in vivo* and *in vitro* through the alteration of cholinergic systems (Puangmalai et al., 2017). *S. iberica* extracts were applied to BV-2 microglia cells at 200 µg/mL concentrations to 3.12 µg/mL. Sodium valproate was used as a positive control. BV-2 cells were pretreated with extracts and fractions. 1 hour later, scopolamine (4mM) was applied. After 24 hours of incubation, the cell viability was evaluated using the MTT method. All tested extracts showed neuroprotective effects but, the water extract was found to be significant at the doses of 12.5-50 µg/mL (*p*<0.01). The isolation and structure elucidation studies to determine the active compounds are ongoing.

Keywords: *Swertia iberica*, cytotoxicity, Gentianaceae

References

http-1: <https://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:30002722-2> http-2: <https://bizimbitkiler.org.tr/v2/hiyerarsi.php?c=Swertia>

Kshirsagar, P. R., Jagtap, U. B., Gaikwad, N. B., & Bapat, V. A. (2019). Ethanopharmacology, phytochemistry and pharmacology of medicinally potent genus *Swertia*: An update. South African Journal of Botany, 124, 444-483.

Puangmalai, N., Thangnipon, W., Soi-Ampornkul, R., Suwanna, N., Tuchinda, P., & Nobsathian, S. (2017). Neuroprotection of N-benzylcinnamide on scopolamine-induced cholinergic dysfunction in human SH-SY5Y neuroblastoma cells. Neural Regeneration Research, 12, 1492.

*Presenting author. E-mail address: fatmanurt@anadolu.edu.tr (Fatmanur TUNÇ)

S1.P113 Tyrosinase inhibitory and cytotoxic activities of two *Inula* species (Asteraceae)

Zeynep Gülcan^{a,*}, Fatmanur Tunç^a, Nadire Özenver^b, Burak Temiz^c, Funda Nuray Yalçın^b, Yavuz Bülent Köse^a

^a Department of Pharmaceutical Botany, Faculty of Pharmacy, Anadolu University, 26470, Tepebaşı, Eskişehir, Türkiye

^b Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University, 06230, Altındağ, Ankara, Türkiye

^c Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University, 26470, Tepebaşı, Eskişehir, Türkiye

The *Inula* L. genus, belonging to the Asteraceae family, includes 128 species worldwide (<http-1>) and 32 taxa in Türkiye (<http-2>). *n*-hexane, ethyl acetate, methanol extracts of *Inula ensifolia* L. (Basionym: *Pentanema ensifolium* (L.) D.Gut.Larr.) and *Inula montbretiana* DC. were prepared from the aerial parts. The tyrosinase inhibition properties of the extracts were assessed following the method described by Likhitwitayawuid and Sritularak (2001) with slight modifications. Extracts were tested at 1 mg/mL and 0.5 mg/mL concentrations. Notably, at a concentration of 0.5 mg/mL, significant tyrosinase inhibition was observed solely in the methanolic extract of *I. montbretiana*. This extract exhibited a remarkably high tyrosinase inhibition activity (16.40 ± 0.48 %), compared to the positive control kojic acid (16.52 ± 0.9 %). Also, at a concentration of 1 mg/mL, the most potent tyrosinase inhibition was observed in the methanolic extract of *I. ensifolia* species (23.10 ± 2.49 %). The methanol extracts of the plants were evaluated in terms of their cytotoxic activity on MCF-7 and MDA-MB-231 breast cancer cell lines using the resazurin reduction assay (Kuate et al., 2013). The methanolic extract of *I. montbretiana* exhibited a higher cytotoxic activity profile on MDA-MB-231 and MCF-7 breast cancer cells with IC₅₀ values of 73.28 and 237.04 mg/mL, respectively, compared to those of *I. ensifolia*. The IC₅₀ value of the methanolic extract of *I. ensifolia* on MDA-MB-231 cells was 393.06 mg/mL, while it did not inhibit cell viability of MCF-7 cells at the tested concentrations (31.25 to 500 mg/mL).

Keywords: *Inula*, tyrosinase, cytotoxicity, activity, Asteraceae

References

<http-1>: <https://wofplantlist.org/taxon/wfo-4000019106-2023-12?page=1> <http-2>: <https://bizimbitkiler.org.tr/v2/hiyerarsi.php?c=Inula>

Kuate, V., Tchakam, P.D., Wiench, B., Ngameni, B., Wabo, H.K., Tala, M.F., Moungang, M.L., Ngadjui, B.T., Murayama, T., Efferth, T., 2013. Cytotoxicity and modes of action of four naturally occurring benzophenones: 2,2',5,6'- tetrahydroxybenzophenone, guttiferone E, isogarcinol and isoxanthochymol. *Phytomedicine*. 20, 528-536.

Likhitwitayawuid, K., Sritularak, B., 2001. A new dimeric stilbene with tyrosinase inhibitory activity from *Artocarpus gomezianus*. *J. Nat. Prod.* 64, 1457-1459.

* Presenting author. E-mail address: zgulcan@anadolu.edu.tr (Zeynep Gülcan)

S1.P114 Linking soil chemistry to Parkinson's disease pathogenesis: novel neurodegenerative compounds from Actinomycetes

Timothy J. Bushman^{a,*}, Osagie Emokpae^a, Jennifer L. Thies^{a,b}, Lena Seyfarth^a, Emily McEachern^c, Elizabeth Gokie^a, Brandon J. Kim^a, Kim A. Caldwell^a, Lukasz Ciesla^a

^a Department of Biological Sciences, The University of Alabama, Tuscaloosa, Alabama 35487, USA

^b UNC School of Medicine, UNC Chapel Hill, Chapel Hill, North Carolina 27514, USA

^c Department of Chemistry, The University of Alabama, Tuscaloosa, Alabama 35487, USA

With the increase in the global average age there has been a corresponding rise in age-related diseases such as Parkinson's Disease. One of the known risk factors for sporadic PD is the rural lifestyle, although the driving environmental factors are not fully understood. Prior research has demonstrated that an extract of rural soil-dwelling *Streptomyces venezuelae* resulted in neuronal death in *Caenorhabditis elegans* through disruption of mitochondrial complex-I and formation of reactive oxygen species (Caldwell et al., 2009). We have isolated three distinct small-molecular weight metabolites from the *S. venezuelae* neurodegenerative fraction via bio-guided fractionation using a *C. elegans* neurodegeneration model. The metabolite structure will be known by Summer 2024. These metabolites will be screened on human iPSC mid-brain dopaminergic neurons later this spring as well as a human blood-brain barrier transwell model. High-resolution MS/MS data will be uploaded to the Global Natural Products Social Molecular Networking (GNPS), to determine their presence in other samples. We also determined that the actinomycete metabolite piperazine A has the ability to cross the human blood-brain barrier and cause dopaminergic neuron death within *C. elegans*. We are further working on the identification of other neurodegenerative compounds from rural soils (such as fungal prenylated diketopiperazines), as well as screening a collection of previously established neurodegenerative *Streptomyces* strains from Alabama soils to create an MS/MS library (Watkins et al., 2016). This project seeks to uncover a potential contributor to the incidence of PD within rural communities due to specialized metabolites within the soil.

Keywords: Parkinson's disease, *Streptomyces*, blood-brain barrier, iPSCs, neurodegeneration

References

- Caldwell, K.A., Tucci, M.L., Armagost, J., Hodges, T.W., Chen, J., Memon, S.B., Blalock, J.E., DeLeon, S.M., Findlay, R.H., Ruan, Q., Webber, P.J., Standaert, D.G., Olson, J.B., Caldwell, G.A. 2009. Investigating bacterial sources of toxicity as an environmental contributor to dopaminergic neurodegeneration. PLoS ONE 4, e7227.
- Watkins, A.L., Ray, A.R., Roberts, L., Caldwell, K.A., Olson, J.B. 2016. The prevalence and distribution of neurodegenerative compound-producing soil *Streptomyces* spp. Sci. Rep. 2016, 6, 22566.

* Presenting author. E-mail address: tjbushman@crimson.ua.edu (Timothy Bushman)

S1.P115 Isolation and bioactivity evaluation of sesquiterpenes from soft corals of the genus *Lemnalia* collected in the Red Sea

Aikaterini Koutsaviti^a, Georgios Spyropoulos^a, Mohamed A. Tammam^{a,b}, Marie Kvasnicová^c, Gabriel Gonzalez^c, Tereza Štenclová^c, Susana Agusti^d, Carlos M. Duarte^d, Lucie Rarová^c, Miroslav Strnad^c, Vassilios Roussis^{a,*}, Efstathia Ioannou^a

^a Section of Pharmacognosy and Chemistry of Natural Products, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771 Athens, Greece

^b Department of Biochemistry, Faculty of Agriculture, Fayoum University, 63514 Fayoum, Egypt

^c Faculty of Science, Palacký University, Slechtitelu 27, CZ-78371 Olomouc, Czech Republic

^d Red Sea Research Center (RSRC), King Abdullah University of Science and Technology (KAUST), 23955-6900 Thuwal, Saudi Arabia

Over the last decades, soft corals have been proven to be a rich source of biologically active compounds, featuring a wide range of chemical structures. Soft corals of the genus *Lemnalia* (Neptheidae) are particularly rich in sesquiterpenes and nor-sesquiterpenes featuring various carbon skeletons, including the nardosinane and neolemnane carbocycles, that have demonstrated a broad spectrum of biological activities, such as cytotoxic, anti-inflammatory and antiviral (MarinLit, 2024; Wu et al., 2018). In the framework of our ongoing research towards the isolation of new bioactive marine metabolites, we investigated the chemistry of two populations of *Lemnalia* collected by SCUBA diving from the coral reefs of Al Jadir and Shi'b Sulaym near Al Lith on the south-west coast of the Kingdom of Saudi Arabia. Exhaustive extraction of the animal tissues with mixtures of CH₂Cl₂/MeOH afforded the organic residues that were subjected to several chromatographic steps to allow up to now for the isolation of 48 different sesquiterpenes, among which 6 are new natural products. The structures of the isolated compounds were determined on the basis of thorough analysis of 1D- and 2D-NMR and MS data. The metabolites isolated in sufficient amounts were evaluated in vitro in human tumor and non-cancerous cell lines for a number of biological activities, including their cytotoxic, anti-inflammatory, anti-angiogenic, and neuroprotective activities, as well as for their effect on androgen receptor-regulated transcription.

Keywords: marine natural products, *Lemnalia* sp., soft corals, biological activity

References

- MarinLit. A Database of the Marine Natural Products Literature. Available online: <http://pubs.rsc.org/marinlit/> (accessed on February 2024).
Wu, Q., Sun, J., Chen, J., Zhang, H., Guo, Y. W., Wang, H., 2018. Terpenoids from marine soft corals of the genus *Lemnalia*: Chemistry and biological activities. Mar. Drugs 16, 320-334.

* Presenting author. E-mail address: roussis@pharm.uoa.gr (Vassilios Roussis)

S1.P116 Production of natural products by using plant cell cultures as chemical factories

*Gilbert Gorr, David A. Ullisch, Thomas Leibold

Phyton Biotech GmbH, Alter Postweg 1, 22926 Ahrensburg, Germany

Reliable production of plant-based natural products can be challenging. Numerous issues have an impact on supply chains associated with natural harvest. Plant to plant variety in yields and profiles, endangered plant species which are difficult to cultivate, and climate changes are some of the limitations to be considered. In contrast, plant cell culture is known to be an attractive alternative supply solution. Plant cell culture technology is an excellent tool to enrich specific structures out of closely related compounds thus reducing impurities in a significant manner. Furthermore, and superior to synthetic biology approaches, plant cells can produce combinations of molecules thereby allowing matching with extract profiles identified in the whole plant. In fact, it has been shown that the production of secondary metabolites with plant cell fermentation can not only be economically attractive, but at the same time allows the supply of the substance of interest in high quality. Independently from the natural environment a plant cell fermentation process, the supply needs for active compounds can be fulfilled in terms of reliability, consistency, quality and quantity. Examples of successful production of natural products through plant cell culture technology will cover a diversity of secondary metabolites, including thapsigargin, ingenol mebutate, paclitaxel and QS-21. Presenting these case studies will also include confirmation of bioequivalence of plant cell culture-derived QS-21 profile to the product isolated from the bark extract (Lv et al., 2024). Furthermore, we will highlight the use of paclitaxel purified from *Taxus chinensis* cell culture cultivated at the 75000 L scale in pharmaceutical development (Dizerega et al., 2024).

Keywords: plant cell culture, paclitaxel, thapsigargin, ingenol mebutate, QS-21

References

- Lv, X., Martin, J., Hoover, H., Joshi, B., Wilkens, M., Ullisch, D.A., Leibold, T., Juchum, J.S., Revadkar, S., Kalinovska, B., Keith, J., Truby, A., Liu, G., Sun, E., Haserick, J., DeGnore, J., Conolly, J., Hill, A.V.S., Baldoni, J., Kensil, C., Levey, D., Spencer, A.J., Gorr, G., Findeis, M. and Tanne, A., 2024. Chemical and biological characterization of vaccine adjuvant QS-21 produced via plant cell culture. doi.org/10.1016/j.isci.2024.109006.
- Dizerega, G., Maulhardt, H.A., Verco, S.J., Marin, A.M., Baltezor, M.J., Mauro, S.A. and Iacobucci, M., 2024. Intratumoral injection of large surface area microparticle taxanes in carcinomas increases immune effector cell concentrations, checkpoint expression, and synergy with checkpoint inhibitors: A review of preclinical and clinical studies. doi:10.1007/s40487-024-00261-y.

*Presenting author. E-mail address: gilbert.gorr@phytonbiotech.com (Gilbert Gorr)

S1.P117 Synergistic approaches in natural products discovery: metabolomic, genomic, and bioactivity- guided isolation of new compounds from the cyanobacterium *Phormidium* sp. LEGE 15488

Leonor Ferreira^{a,b,*}, Ana C. Fonseca^{a,b}, João Morais^a, Ralph Urbatzka^a, Vitor Vasconcelos^{a,b}, Mariana Reis^a

^a CIIMAR - Interdisciplinary Centre of Marine and Environmental Research, Terminal de Cruzeiros do Porto de Leixões, Avenida General Norton de Matos, S/N, 4450-208, Matosinhos, Portugal

^b FCUP - Faculty of Sciences, University of Porto, Rua do Campo Alegre, S/N, 4169-007 Porto, Portugal

Cyanobacteria are a diverse group of photosynthetic prokaryotes explored for their potential to produce bioactive compounds with diverse applications. However, discovering natural products in cyanobacteria is challenging due to slow growth rates for biomass production and low secondary metabolites yields. More recently, the development of genomic and metabolomics tools has increased the efficiency of natural products research, and the combination of traditional and new methodologies delivers more successful results. *Phormidium* sp. LEGE 15488 (strain from the Blue Biotechnology and Ecotoxicology Culture Collection, LEGE-CC) was identified as a producer of novel bioactive compounds during a previous cyanobacterial natural products screening, which combined anticancer assays with untargeted metabolomics dereplication (Ferreira et al., 2021). Its genome was sequenced and analysed with antiSMASH, revealing numerous Nonribosomal Peptides/Polyketides biosynthetic gene clusters. Herein, we will present the genome mining and the isolation of novel bioactive metabolites from *Phormidium* sp. LEGE 15488. After large-scale cultivation, the compound isolation was guided by mass spectrometry and bioactivity results. The structure of a novel pseudospumigin- like compound was elucidated by 1D and 2D NMR experiments, along with LC-HRESIMS/MS. At 10 μ M this compound reduced 53% of appetite in the zebrafish feeding assay with fluorescent labelled *Paramecia bursaria*, which is an interesting pathway for obesity treatment. This work emphasizes the significance of integrating multiple strategies to discover novel cyanobacterial natural products with potential therapeutic applications.

Acknowledgements. This work is supported by national funding from Fundação para a Ciência e Tecnologia (FCT) by the strategic fund UIDB/04423/2020 and UIDP/04423/2020 and FCT grant 2022.11979.BD.

Keywords: cyanobacteria, metabolomics, genomics, pseudospumigins, obesity

References

Ferreira, L., Morais J., Preto M., Silva, R., Urbatzka R., Vasconcelos V., Reis M., 2021. Uncovering the Bioactive Potential of a Cyanobacterial Natural Products Library Aided by Untargeted Metabolomics. *Mar Drugs*. 19, 633.

* Presenting author. E-mail address: lferreira@ciimar.up.pt (Leonor Ferreira)

S1.P118 Peptide toxins from ribbon worms (*Nemertea*)

Erik Jacobsson^{a,*}, Quentine Laborde^a, Håkan S. Andersson^b, Ulf Göransson^a

^a Division of Pharmacognosy, Department of Pharmaceutical Bioscience, Faculty of Pharmacy, Uppsala University, Sweden

^b Department of Medical Biochemistry and Biophysics, Karolinska Institute, Sweden

This presentation will cover the field of ribbon worm peptide toxins. The ribbon worms, *Nemertea*, is a phylum of mostly marine animals, comprising some 1350 species. They vary in size from millimeters to 50 meters; some of them are the top predators/scavengers in their habitats. Ribbon worms secrete toxin-loaded mucus for protection and use an eversible proboscis to hunt prey. The mucus from some species has been investigated chemically. Some contain small molecule toxins (tetrodotoxin, TTX (Asakawa et al. 2003)), while others contain peptide toxins (nemertides a-c, parborlysins 1-7, toxin-A, and neurotoxins B I-IV (Göransson et al. 2019)). The peptide toxins have been attributed two main functions: nemertides a-c and neurotoxins B I-IV exhibit ion channel activity, while toxin-A and parborlysins are cytolytic. With increasing ribbon worm genetic data available in scientific databases (two genomes, and more than 100 transcriptomes), the number of putative peptide toxins and toxin families in ribbon worms is rising rapidly. Here, we present the current status of peptide toxins in ribbon worms.

Keywords: *Nemertea*, ribbon worm, peptide toxin, marine

References

Asakawa, M., Toyoshima, T., Ito K., Bessho K., Yamaguchi C., Tsunetsugu S., Shida Y., et al. 2003. Paralytic Toxicity in the Ribbon Worm *Cephalothrix* Species (*Nemertea*) in Hiroshima Bay, Hiroshima Prefecture, Japan and the Isolation of Tetrodotoxin as a Main Component of Its Toxins. *Toxicon* 41, 747–53.

Göransson, U., Jacobsson, E., Strand, M., and Andersson H.S. 2019. The Toxins of Nemertean Worms. *Toxins* 11 (2).

* Presenting author. E-mail address: erik.jacobsson@uu.se (Erik Jacobsson)

S1.P119 Exploring the pharmaceutical potential of the Antarctic fungus *Epicoccum nigrum* isolated from *Kallymenia antarctica* seaweed

Hosana Maria Deboni^{a,*}, Ana Carolina Jordão^a, Lorena Rigo Gaspar^b, Ludmilla Tonani Carvalho^c, Márcia Regina von Zeska Kress^c, Thaiz Rodrigues Teixeira^d, Conor Caffrey^d, Pio Colepicolo^e

^a Department of Biomolecular Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

^b Department of Pharmaceutical Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

^c Department of Clinical, Toxicological and Bromatological Analysis, School of Pharmaceutical Sciences, University of São Paulo, São Paulo, SP, Brazil

^d Center for Discovery and Innovation in Parasitic Disease, Skaggs School of Pharmacy and Pharmaceutical Sciences, University California San Diego, CA, USA

^e Department of Biochemistry, Chemistry Institute, University of São Paulo, São Paulo, SP, Brazil

The Earth's last frontiers, the poles, host extreme climatic conditions, triggering organisms to develop survival strategies, including biosynthesizing unique chemical compounds. Exploring organisms in these regions offers potential for discovering novel bioactive pharmaceutical compounds (Santos et al., 2021). In this context, we studied *Epicoccum nigrum*, an endophytic fungus isolated from the Antarctic seaweed *Kallymenia antarctica*. Our objectives involved the chemical and biological analysis of secondary metabolites produced by *E. nigrum*, including isolation/elucidation of the extract's major constituents, also assessing their biological potential. For this purpose, *E. nigrum* was cultured on parboiled rice and seawater. Ethyl acetate was used for extraction. Fractionation by liquid chromatography yielded nine fractions (A-I). All samples were tested for antifungal, antiparasitic and photoprotective activities. Four compounds were isolated and identified: epicoccarine A (**1**), epicoccarine B (**2**), beauvericin (**3**) and triterpene S19159 (**4**). Fractions E-G showed significant activity against fungal species, as did **1**, **3** and **4**. Fractions C-I and compound **1** exhibited high toxicity against *T. brucei* parasites but also human HEK293 cells, presenting low viability. Fractions F and G also displayed photoprotective potential, absorbing UV radiation with no degradation after UVA exposure. However, they showed phototoxic potential. One compound (pending identification) exhibited broad UV absorption with the advantage of no phototoxicity. Further study will involve co- cultivation with Actinomycetes to explore the potential for producing new compounds when challenged. Preliminary tests suggest that *E. nigrum* produces colorful compounds in proximity to other organisms. In conclusion, this work indicates that *E. nigrum* has potential as a source of diverse natural products with promising bioactivities.

Keywords: marine natural products, *Epicoccum nigrum*, photoprotection, antifungal, antiparasitic

References

Santos G.S., Teixeira T.R., Colepicolo P., Deboni H.M., 2021. Natural Products from the Poles: structural Diversity and Biological Activities. Rev. Bras. Farmacogn. 31, 531–560.

* Presenting author. E-mail address: hosana@fcfrp.usp.br (Hosana Maria Deboni)

S1.P120 Antiparasitic activity of the antarctic fungus *Aspergillus unguis* metabolites against *Trypanosoma brucei*

Márcia G. O. Lopes^{a,*}, Thaiz Rodrigues Teixeira^b, Conor Caffrey^b, Gustavo Santos^c, Anibal Freitas^c, Pio Colepicolo^d, Hosana Maria Deboni^c

^a Department of Pharmaceutical Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

^b Center for Discovery and Innovation in Parasitic Disease, Skaggs School of Pharmacy and Pharmaceutical Sciences, University California San Diego, CA, USA

^c Biopharmacy and Natural Products Laboratory of Bahia, Salvador, BA, Brazil

^d Department of Biochemistry, Chemistry Institute, University of São Paulo, São Paulo, SP, Brazil ^e Department of Pharmaceutical Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

The Antarctic exhibits extreme climatic conditions prompting organisms to evolve unique survival mechanisms, such as the synthesis of distinct chemical compounds. Investigating polar organisms presents significant opportunities for uncovering innovative bioactive pharmaceutical agents (Teixeira et al., 2019). Considering this, we suggested investigating *Aspergillus unguis*, which was obtained from the Antarctic seaweed *Palmaria decipiens*. The study aimed to conduct chemical and biological analyses of secondary metabolites generated by *A. unguis*, including the identification and characterization of the primary constituents of the extract. Additionally, biological activity against *Trypanosoma brucei* was evaluated. To achieve this, *A. unguis* was cultivated on parboiled rice and seawater. Ethyl acetate was employed for extraction, and subsequent fractionation via liquid chromatography resulted in the isolation of nine fractions (A-I). The full sample collection (4 mg) was tested for antiparasitic activity, and until now, one compound from D fraction was isolated and identified as ergosterol (1). According to the results, the crude extract and fractions C, D, E, F and H were promising against *Trypanosoma brucei*, ranging between 97 and 100% inhibition. Furthermore, the crude extract and the C and D fractions retained very high cell viability in HEK293 cells (CC50 µg/mL >10). Fraction D was submitted to GC-MS analyses which demonstrated that ergosterol was the main compound present in the fraction. Given the above, the most promising fractions will be further separated, and the compounds isolated and identified. In summary, this study highlights the potential of *A. unguis* as a source of diverse natural products with promising bioactivities against the parasitic organism *Trypanosoma brucei*.

Keywords: marine natural products, Antarctic, *Aspergillus unguis*, *Trypanosoma brucei*, antiparasitic

References

Teixeira, T.R., Santos, G.S., Turatti, I.C.C., Paziani, M.H., Kress, M.R.Z., Colepicolo, P., Deboni, H.M., 2019. Characterization of the lipid profile of Antarctic brown seaweeds and their endophytic fungi by gas chromatography– mass spectrometry (GC–MS). *Polar Biol.* 42, 1431–1444.

* Presenting author. E-mail address: marcialopes.qoam@usp.br (Márcia Gracielle de Oliveira Lopes)

S1.P121 Unlocking nature's pharmacy : *Iris pseudacarus* from Ireland as potential source for antimicrobial and anti-inflammatory compounds

Özlem Erol^{a,*}, Shipra Nigar^b, Mohammed A. Selo^b, Helen Sheridan^b, Young H. Choi^a

^a Natural Products Laboratory, Institute of Biology, Leiden University, Sylviusweg 72, 2333BE Leiden, The Netherlands

^b NatPro Centre, School of Pharmacy and Pharmaceutical Sciences, Trinity College Dublin, Dublin 02, D02 PN40 Dublin, Ireland

Iris pseudacarus, commonly known as Yellow Iris or Yellow Flag Iris, holds significant botanical and ethnopharmacological importance, particularly in Irish folklore. This flowering plant species, prevalent in Irish Bogs, is renowned for its diverse phytochemical composition, including flavonoids, isoflavonoids, triterpenes, phenolics, and alkaloids. Various studies have highlighted its pharmacological potential, showcasing activities such as antiviral, antibacterial, and anti-tumor properties. In this study, part of a broader project on unlocking nature's pharmacy from bogland species in Ireland, we focused on profiling *I. pseudacarus*. Through the analysis of 16 dried plant materials (leaves, stems, and roots) using HPTLC, LC-MS, and NMR, we aimed to identify unique compounds and explore their biological activities. HPTLC bioautography assays combine chromatographic separation with in situ biological activity determination, allowing the direct and rapid isolation of active compounds in complex plant extracts. Our investigation led to the isolation of three antimicrobial compounds - syringic acid, luteolin-7-O-glucoside, and liquiritigenin - from *I. pseudacarus* leaf and root extracts, guided by HPTLC bioautography. These compounds exhibited antimicrobial activity against *S. aureus* and were further evaluated for their anti-inflammatory and cytotoxic properties. Overall, our findings contribute to understanding the pharmacological potential of *I. pseudacarus* and underscore the significance of utilizing traditional botanical resources for drug discovery and development.

Keywords: *Iris pseudacarus*, HPTLC bioautography, antimicrobial and anti-inflammatory activity

References

- Nürk, N.M., Madriñán, S., Carine, M.A., Chase, M.W., Blattner, F.R., 2013. Molecular phylogenetics and morphological evolution of St. John's wort (*Hypericum*; Hypericaceae). *Mol. Phylogenetics Evol.* 66, 1-16.
- Reece, R.J., Maxwell, A., 1991. DNA gyrase: structure and function. *Crit. Rev. Biochem. Mol. Biol.* 26, 335-375.
- Robson, N.K., 2021. Studies in the genus *Hypericum* L.(Hypericaceae). *Phytotaxa.* 72, 1-111.
- Soepadmo, E., Saw, L.G., Chung, R.C.K. (Eds.). 2002. The Tree Flora of Sabah and Sarawak, Volume 4. Forest Research Institute of Malaysia, Selangor Darul Ehsan.

* Presenting author. E-mail address: o.erol@biology.leidenuniv.nl (Özlem Erol)

S1.P122 Antitrichomonal activity of secondary metabolites of *Alternaria* sp., a fungal endophyte of alpine Edelweiss

Elahe Mirzaei Moghadam^{a,*}, Sofia Karavergou^a, Tobias Kiss^a, Ursula Fürnkranz^b, Hubertus Haas^c, Mostafa Alilou^a

^a Unit of Pharmacognosy, Institute of Pharmacy, Center for Molecular Biosciences (CMBI), University of Innsbruck, Innsbruck, Austria

^b Pilzambulatorium, Schlösselgasse 19, Vienna, Austria

^c Institute of Microbiology, Universität Innsbruck, Innsbruck, Austria

Sexually transmitted infections (STIs) are considered a major global health concern, caused by bacteria, viruses or parasites. (Rowley et al., 2019). Among them, Trichomoniasis, with an estimated annual incidence of 300 million cases worldwide, is one of the most common sexually transmitted infections, caused by the unicellular protozoan parasite *Trichomonas vaginalis*. Resistance of the parasite to the existing therapies has increased the quest for the discovery of novel antitrichomonal substances (Natto et al., 2021). *Alternaria* species exhibit saprophytic, endophytic or pathogenic lifestyles. They have been studied thoroughly due to the production of toxins and unusual secondary metabolites, classified into the groups of nitrogen-containing compounds, quinones, and pyranones, displaying significant pharmacological activities against human pathogens (Lou et al., 2013). This investigation aimed to isolate and identify the bioactive metabolites of *Alternaria* sp., an endophytic fungus isolated from the leaves of alpine edelweiss (*Leontopodium alpinum*), and assessment of their antitrichomonal activities. Bioprospecting of *Alternaria* sp. resulted in purification of two known pyranone derivatives, alternariol (1) and alternariol-9-methyl ether (2), one unknown pyranone-type derivative (3), two new perylene quinone derivatives (4) and (5) and two unknown diastereomers of tetramic-acid containing metabolites (6) & (7). The structural characterization was performed using 1D- & 2D-NMR and HRMS, and their absolute configurations were established using electronic circular dichroism (ECD). Pure compounds were evaluated against *T. vaginalis*, all of which illustrated moderate to promising bioactivities. According to the results obtained, *Alternaria* sp. is considered a promising source of unexplored compounds with potential antitrichomonal activity.

Keywords: *Alternaria*, endophytes, metabolomics, structural characterization, *Trichomonas vaginalis*

References

- Lou, J., Fu, L., Peng, Y., Zhou, L., 2013. Metabolites from *Alternaria* fungi and their bioactivities. *Molecules* 18, 5891- 5935.
- Natto, M.J., Hulpia, F., Kalkman, E.R., Baillie, S., Alhejeli, A., Miyamoto, Y., Eckmann, L., Van Calenbergh, S., de Koning, H.P., 2021. Deazapurine nucleoside analogues for the treatment of *Trichomonas vaginalis*. *ACS infect dis.* 7, 1752-1764.
- Rowley, J., Vander Hoorn, S., Korenromp, E., Low, N., Unemo, M., Abu-Raddad, L. J., Chico, R. M., Smolak, A., Newman, L., Gottlieb, S., Thwin, S. S., Broutet, N., Taylor, M. M., 2019. Chlamydia, gonorrhoea, trichomoniasis and syphilis: global prevalence and incidence estimates, 2016. *Bull World Health Organ.* 97, 548–562.

* Presenting author. E-mail address: Elahe.Mirzaei-Moghadam@uibk.ac.at (Elahe Mirzaei Moghadam)

S1.P123 Isolation and structural elucidation of bioactive cycloartane glycosides from *Astragalus glycyphyllos*

Monica Scognamiglio*, Acácio Raphael A. Moraes, Antonio Fiorentino

Dipartimento di Scienze e Tecnologie Ambientali, Biologiche e Farmaceutiche, Università degli Studi della Campania "Luigi Vanvitelli", Via Vivaldi 43, 81100 Caserta, Italy

Species belonging to the *Astragalus* genus (Fabaceae family) have been used for millennia for treating various diseases, including cancer. Compounds isolated from plants belonging to this genus have been shown to exert significant anticancer activity in vitro and in animal models (Graziani et al., 2019a). Cycloartane saponins isolated from *Astragalus boeticus* have been shown to preferentially inhibit cell growth in colorectal cancer cell models resistant to epidermal growth factor receptor inhibitors (Graziani et al., 2019b). In this study, two further *Astragalus* species, namely *Astragalus hamosus* and *Astragalus glycyphyllos*, were taken into consideration as potential sources of bioactive compounds. To speed-up the discovery process, the plant material was subjected to a preliminary screening using a metabolomics approach. NMR-based profiling showed the presence of cycloartanes only in the *A. glycyphyllos* samples. The plant material of the latter was therefore extracted and separated using chromatographic techniques, to obtain the pure compounds. Structural elucidation of these compounds was carried out by extensive 1D and 2D-NMR analyses. Based on the knowledge acquired through NMR profiling, it was possible to design a custom-made time- and resource-saving isolation procedure, consisting only of a few chromatographic steps. It was, therefore, possible to rapidly isolate and characterize nine cycloartane glycosides, which were identified as cycloastragenol and cycloanthogenin derivatives. Six of the isolated compounds also contained an unusual oxalyl moiety and, to the best of our knowledge, eight of them were reported for the first time. Ongoing studies are dedicated to the evaluation of the anticancer potential of the compounds.

Keywords: *Astragalus glycyphyllos*, Fabaceae, metabolomics, NMR, cycloartane glycosides

References

- Graziani, V., Scognamiglio, M., Esposito, A., Fiorentino, A., D'Abrosca B. 2019a. Chemical diversity and biological activities of the saponins isolated from *Astragalus* genus: focus on Astragaloside IV. *Phytochemistry Reviews*, 18, 1133- 1166.
- Graziani, V., Esposito, A., Scognamiglio, M., Chambery, A., Russo, R., Ciardiello, F. Troiani, T., Potenza, N., Fiorentino, A., D'Abrosca B. 2019b. Spectroscopic characterization and cytotoxicity assessment towards human colon cancer cell lines of acylated cycloartane glycosides from *Astragalus boeticus* L. *Molecules*, 24, 1725.

* Presenting author. E-mail address: monica.scognamiglio@unicampania.it (Monica Scognamiglio)

S1.P124 Screening marine bacterial compounds for adhesion blockers of enteropathogenic *Escherichia coli*

Tuomas Pylkkö^{a,*}, Yannik K. Schneider^b, Päivi Tammela^a

^a Department of Pharmaceutical Biosciences, University of Helsinki, Norway

^b UiT, the Arctic University of Norway, Tromsø, Norway

Due to the rise of antimicrobial resistance and little recent progress in the discovery of new antimicrobials, novel chemical structures for treating bacterial infections are desperately needed. In the past, secondary metabolites of bacteria with larger genomes are more likely to carry biosynthetic gene clusters, and led to the discovery of most antibiotics. Furthermore, many compounds inhibiting virulence have been similarly discovered in screening collections of natural products and microbial metabolites. Therefore, we studied the antibacterial and virulence inhibiting activity of marine actinobacteria gathered on an earlier research cruise around Svalbard, in the Arctic Sea. For the first time, these bacteria were successfully cultivated and grown in liquid media and were then processed by extraction and fractionation using FLASH chromatography. The bioactivity of these fractions was then assessed against the human intestinal pathogen, enteropathogenic *Escherichia coli* (EPEC), utilizing both classic growth inhibition and virulence related assays. We have previously set up an EPEC effector translocation assay and a cell culture-based coinfection model using high-content microscopy. EPEC uses the type three secretion system to inject virulence-causing molecules (effectors), which ultimately lead to abnormal actin condensation, often called actin pedestals, and to the effacement of microvilli from the intestinal epithelium. Inhibiting these processes could effectively inhibit the virulence of the pathogen without largely damaging normal flora, or causing harsh selectional pressure on the pathogen, ultimately leading to antibiotic resistance. According to the screening results, several of the fractions reduced the readouts of these assays in a concentration- dependent manner. Some of these acted by suppressing growth, but some fractions, however, were able to diminish EPEC pathogenicity in our assays without inhibiting growth through as-yet undiscovered mechanisms of action. Work is on-going to determine the composition of the fractions as well as the precise mechanism of action by which these phenotypic changes occur.

Keywords: enteropathogenic *Escherichia coli*, marine actinobacteria

References

Kaper J.B., Nataro J.P., Mobley H.L.T., 2004 Pathogenic *Escherichia coli*. Nat. Rev. Microbiol. Feb;2(2):123–40. Pylkkö T., Ilina P., Tammela P., 2021 Development and validation of a high-content screening assay for inhibitors of enteropathogenic *E. coli* adhesion. J. Microbiol. Methods 106201.

Schneider Y.K., Hagestad O.C., Li C., Hansen E.H., Andersen J.H., 2022 Selective isolation of Arctic marine actinobacteria and a down-scaled fermentation and extraction strategy for identifying bioactive compounds. Front Microbiol. Nov 21;13:1005625.

* Presenting author. E-mail address: tuomas.pylkko@helsinki.fi (Tuomas Pylkkö)

S1.P125 NMR-based metabolomics approach and anti-proliferative activity of *Olea europaea* products

Brigida D'Abrosca^{a,*}, Vincenzo Carafa^b, Giovanna Valentino^a, Lucia Altucci^b, Antonio Fiorentino^a

^a Department of Environmental Biological and Pharmaceutical Sciences and Technologies, DiSTABiF University of Campania "Luigi Vanvitelli", Via Vivaldi 43, 81100 Caserta, Italy

^b Department of Precision Medicine, University of Campania "Luigi Vanvitelli" S. Andrea delle Dame - Via L. De Creschio, 7 - 80138 - Napoli, Italy

Olive (*Olea europaea* L.), belongs to the Oleaceae family, own a significant economic value, as well as nutritional and medicinal properties. Today, 98% of olive are cultivated in Mediterranean area (Antoniou and Hull 2021) and virgin olive oil represent a key food of the Mediterranean diet. However, the value of olive tree extends to its products as olives and leaves. Employed in traditional medicine since ancient times, recent research has highlighted the anticancer properties of olive extracts, attributed to their unique chemical composition rich in specialized metabolites (Scognamiglio 2012). Polyphenols, secoiridoid glycosides, and triterpenoids isolated from olive demonstrate efficacy across various *in vitro* and *in vivo* cancer models. In this way, this work aims to characterize the molecular and functional effects of 21 selected cultivars of *Olea europaea* on three cancer cell lines belonging to a group of "poor prognosis cancers": acute myeloid leukemia (AML), Glioblastoma (GBM) and Pancreatic adenocarcinoma (PA) for which to date there are no evidence in literature. A new screening method combining a NMR-based metabolomic approach (Esposito 2021) with MTT assays was employed to identify olive drupe crude extracts with anti-proliferative properties. The most active extract(s) will be purified to obtain enriched fraction and/or pure compounds for evaluate the involvement of several molecular players in the so-called alternative regulated cell death (RCD) pathways. This presentation will cover analyses of 21 cultivars of olives collected in the experimental agricultural company of Campania Region "Improsta". Dried drupes have been extracted directly with phosphate buffer in D2O and methanol-d4 (1:1) assisted by ultrasound and an aliquot was directly analyzed by Nuclear Magnetic Resonance (NMR). With the same methodology (1.67 g dried drupes), and after hexane extraction and SPE purification, in order obtain enriched in specialized metabolites, have been prepared crude extracts for cell viability test. The results highlighted that Oliviello and Racioppa cultivars are very rich in verbascoside, Grossale Oliva Bianca in oleocanthal, Romanella and Marinese cultivars in DHPEA-EA. Study of modulation of cell proliferation and cell death operated by crude extract of olive showed different effect on three cancer cell lines selected. In conclusion, this study increases the knowledge about *Olea europaea* cultivars diversity in term of metabolite profiling and bioactivity.

Keywords: *Olea europaea*, NMR, metabolomics, cytotoxicity, DNA fragmentation

References

- Antoniou, C., Hull J. 2021. The anti-cancer effect of *Olea europaea* L. Products: a Review. Curr Nutr Rep 10, 99-124. Scognamiglio, M., D'Abrosca, B., Pacifico, S., Fiumano, V., De Luca, P.F., Monaco, P., Fiorentino, A. 2012. Polyphenol characterization and antioxidant evaluation of *Olea europaea* varieties cultivated in Cilento National Park (Italy). Food Res. Intern. 46, 294–303.
- Esposito, A., De Luca, P.F., Graziani, V., D'Abrosca, B., Fiorentino A., Scognamiglio M. 2021. Phytochemical Characterization of *Olea europaea* L. Cultivars of Cilento National Park (South Italy) through NMR-Based Metabolomics. Molecules. 26, 3845.

* Presenting author. E-mail address: brigida.dabrosca@unicampania.it (Brigida D'Abrosca)

S1.P126 Potential nematicidal agents from a bioactive fraction of the fungus *Trichoderma parceramosum*

Carlo Raucci^{a,*}, Maria Chiara Zonno^b, Claudio Fabio Altomare^b, Trifone D'Addabbo^c, Antonio Evidente^b, Brigida D'Abrosca^a

^a Department of Environmental Biological and Pharmaceutical Sciences and Technologies, DiSTABiF University of Campania Luigi Vanvitelli, Via Vivaldi 43, 81100 Caserta, Italy

^b Institute of Sciences of Food Production, National Research Council, Via Amendola 122/O, 70126 Bari, Italy

^c Institute for Plant Protection, National Research Council of Italy, Via Amendola 122/D 70126 Bari, Italy

Endophytic fungi are a category of fungi that reside inside the tissues of host plants without causing evident damage; this symbiotic relationship can be beneficial for both participants. One of the main advantages for plants is that endophytic fungi provide protection against pathogenic organisms (Selim et al., 2012). For example, fungi can produce substances that inhibit the growth of pathogens or directly compete with them for resources, reducing the incidence of diseases (Rajesh et al., 2016). *Trichoderma* spp. secretes a range of specialized metabolites, many of which have demonstrated a broad-spectrum of biological properties in *in vitro* assays. For example, the koniginins, polyketides obtained from *Trichoderma koningii*, in an *in vitro* assay, exhibited strong activity against plant pathogens (Ba-Yi et al., 2015). In the latter years, the potential of *Trichoderma* metabolites against parasitic nematodes has also been highlighted (Moo-Koh et al., 2022). In this context, the focus of this work is the evaluation of unexplored *Trichoderma parceramosum* as a potential source of nematicidal agents. The hydroalcoholic extract of *T. parceramosum* solid culture was extracted first with *n*-hexane and successively with dichloromethane. This latter extract, through chromatographic techniques, afforded eight fractions which were tested for their nematicidal activity against *Meloidogyne incognita*, one of the most damaging pests in vegetable crops worldwide. From more active fractions, phenolic compounds and heterocyclic lactones have been isolated. The structural characterization of the isolated compounds was performed by 1D-NMR and 2D-NMR investigations.

Keywords: endophytic fungi, *Trichoderma parceramosum*, NMR, nematicidal activity, *Meloidogyne incognita*

References

- Ba-Yi, L., Jing, L., Xiao-Xue, Z., Yu-Hui, C., Yin-He, Y., Xiao-Nian, L., Ying, Z., Pei-Ji, Z., 2015. Koninginins L and M, two polyketides from *Trichoderma koningii* 8662. *Phytochem. Lett.* 11, 1-4.
- Moo-Koh, F.A., Cristóbal-Alejo, J., Andrés, M.F., Martín, J., Reyes, F., Tun-Suárez, J.M., Gamboa-Angulo, M., 2022. In Vitro Assessment of Organic and Residual Fractions of Nematicidal Culture Filtrates from Thirteen Tropical *Trichoderma* Strains and Metabolic Profiles of Most-Active". *J. Fungi (Basel)* 8, 82.
- Rajesh, R.W., Rahul, M.S., Ambalal, N.S., 2016. *Trichoderma*: A significant fungus for agriculture and environment. *Afr. J. Agric. Res.* 11, 1952-1965.
- Selim, K.A., El-Beih, A.A., AbdEl-Rahman, T., El-Diwany, A.I., 2012. Biology of endophytic fungi. *Curr. Res. Environ. Appl. Mycol.* 2, 31-82.

* Presenting author. E-mail address: carlo.raucci@unicampania.it (Carlo Raucci)

S1.P127 Antimicrobial secondary metabolites of *Trochila* sp., a new endophytic fungus of *Lilium carnolicum*

Mostafa Alilou^{a,*}, Ursula Fürnkranz^b, Isidor Happacher^c, Hubertus Haas^c

^a Unit of Pharmacognosy, Institute of Pharmacy, Center for Molecular Biosciences (CMBI), University of Innsbruck, Innsbruck, Austria

^b Pilzambulatorium, Schlüsselgasse 19, Vienna, Austria

^c Institute of Microbiology, Universität Innsbruck, Innsbruck, Austria

Endophytes are group of microorganisms, which protect their host plants against various abiotic and biotic stress factors. These microorganisms are underexplored sources of new biologically active natural products against not only phytopathogens, but also against bacteria, fungi, viruses, and protozoans that affect human and animals (Strobel et al., 2003). In a screening campaign for the discovery of antimicrobial natural products against a human pathogen *Trichomonas vaginalis* and a phytopathogen *Botrytis cinerea*, we have identified a fungal hit, *Trochila* sp., an undescribed species isolated from leaves of *Lilium carnolicum*, an endemic *Lilium* species found in the alpine region of Austria. In this study we aimed for dereplication, and identification of secondary metabolites produced by a rice culture of *Trochila* sp., using Feature-Based Molecular Networking of the GNPS platform, as well as SIRIUS for in silico annotations. By inspecting the network clusters and annotations results, we were able to dereplicate and perform targeted isolation of two dibenzofuran derivatives, usnic acid and isousnic acid (**1-2**), two unknown and two previously reported pyridine type alkaloids, (**3-6**), and a known dibenzoquinone, oosporein (**7**), and annotated potentially new metabolites through analysis of the MN clusters. The structures of the isolated compounds were elucidated using 1D and 2D-NMR spectroscopy and their absolute configurations established utilizing DP4+ probability and circular dichroism calculations. Ultimately, the biological activity of isolates was evaluated against three aforementioned pathogens. Our results shed new light on the metabolite profile of the genus *Trochila* as a promising source of bioactive natural products.

Keywords: *Trochila*, endophytes, metabolomics, structural characterization, *Trichomonas vaginalis*

References

Strobel, G., Daisy, B., 2003. Bioprospecting for Microbial Endophytes and Their Natural Products. Microbiol. Mol. Biol. Rev. 67, 491–502.

* Presenting author. E-mail address: mostafa.alilou@uibk.ac.at (Mostafa Alilou)

S1.P128 Seasonal variation of dimeric flavonoids in ginkgo (*Ginkgo biloba* L.) leaves

Iva Jurčević Šangut, Dunja Šamec*

Department of Food Technology, University North, Trg Dr. Žarka Dolinara 1, 48 000 Koprivnica, Croatia

Ginkgo (*Ginkgo biloba* L.) is a well-known medicinal plant that is rich in various bioactive compounds. To date, 110 different flavonoids have been identified in ginkgo, including flavonoid dimers known as biflavonoids. Biflavonoids have recently gained much attention as potential molecules with biological activity such as antiviral and antimicrobial activity as well as effective molecules for the treatment of neurodegenerative and metabolic diseases and anti-cancer molecules. As part of our ongoing project, we developed a method for the separation and quantification of biflavonoids in ginkgo leaves using HPLC-DAD (Kovač Tomas et al., 2023). The biflavonoids present in ginkgo leaves are amentoflavone, bilobetin, isoginkgetin, ginkgetin and sciadopitysin (Fig. 1).




Fig. 1. Structure of ginkgo bioflavonoids.

During the 2022 season, from May to November, we collected leaves from ginkgo trees every month, 7 stages in total. The leaves were freeze-dried and we analysed the content of individual biflavonoids. In the period from May to November, we observed an increased accumulation of biflavonoids, for most of them by more than 300%, indicating that fall leaves are richer in biflavonoids than green leaves. Only in the case of amentoflavone we did not detect an increase in content. A particularly large increase in biflavonoids occurred between the leaves collected in June and July, which could indicate extensive biosynthesis of biflavonoids during this period. Further investigations will reveal possible factors related to the biosynthesis of biflavonoids.

Keywords: *Ginkgo biloba* L., biflavonoids, seasonal variation

References

- Liu, L., Wang, Y., Zhang, J., Wang, S. 2021. Advances in the chemical constituents and chemical analysis of *Ginkgo biloba* leaf, extract, and phytopharmaceuticals. J. Pharm. Biomed. Anal. 193,113704.
- Kovač Tomas M, Jurčević I, Šamec D (2023) Tissue-Specific Profiling of Biflavonoids in Ginkgo (*Ginkgo biloba* L.). Plants 12(1), 147.

* Presenting author. E-mail address: dsemec@unin.hr (Dunja Šamec)

S1.P129 Efficient production of some bioactive depsides by *Aronia* × *prunifolia* (Rosaceae) microshoot agitated *in vitro* cultures as the effect of precursor feeding strategy

Paweł Kubica, Agnieszka Szopa, Halina Ekiert*

Chair and Department of Pharmaceutical Botany, Faculty of Pharmacy, Jagiellonian University, Collegium Medicum, 9 Medyczna Str., 30-688 Cracow, Poland

Aronia × *prunifolia* (Marsh.) Rehd. (Rosaceae) is rich source of phenolic acids (PhAs) - antioxidants important in phytotherapy and also in phytocosmetology (Szopa et. al., 2017). (PhAs) could be produced in high amounts in plant *in vitro* cultures, also in established by our team microshoot cultures of *A. × prunifolia* (Ekiert et al., 2022). The aim of the present study was the testing of precursor feeding strategy on the production of (PhAs) in agitated cultures of *A. × prunifolia*. The cultures were maintained in Murashige-Skoog medium (1mg/l BAP and 1mg/l NAA) (Murashige et al., 1962). Precursors of simple (PhAs) - phenylalanine, cinnamic acid, benzoic acid and of depsides - caffeic acid at five concentrations (0.1, 0.5, 1.0, 5.0 and 10.0 mmol/L) were fed into the medium at point "0" (culture initiation) and independently on the 10th day of growth cycle. After the 20th day of growth cycle, the contents of compounds were determined in methanolic extracts of biomasses by HPLC (Ellnain-Wojtaszek et al., 1999). All extracts contained 4 depsides (chlorogenic, neochlorogenic, rosmarinic and cryptochlorogenic acids) and 4 simple (PhAs) - protocatechuic, vanillic, caffeic and syringic acids. Chlorogenic and neochlorogenic acids, dominated in all extracts (max. 388.39 and 263.54 mg/100g d.w., respectively). The total content of all compounds was the highest after the addition at point "0" of cinnamic acid (5 mmol/l) and caffeic acid (10 mmol/l) which caused a 2.68- and 2.49-fold increase in the contents of (PhAs) - 603.03 and 558.48 mg/100g d.w., respectively. The obtained results documented efficacy of the precursor feeding strategy in enhancing the production of bioactive depsides in microshoot agitated cultures of *A. × prunifolia* and have a potential application value.

Keywords: *Aronia* × *prunifolia*, *in vitro* cultures, biogenetic precursors, depsides, HPLC

References

- Ekiert, H., Kubica, P., Kwiecień, I., Jaferník, K., Klimek-Szczykutowicz, M., Szopa, A., 2022. Cultures of medicinal plants *in vitro* as a potential rich source of antioxidants. Ekiert, H., Ramawat, K., Arora, J. (Eds.), 2022. References Series in Phytochemistry. Plant Antioxidants and Health. Springer Nature, Switzerland.
- Ellnain-Wojtaszek, M., Zgórká, G., 1999. High-performance liquid chromatography and thin-layer chromatography of phenolic acids from *Ginkgo biloba* L. leaves collected within vegetative period. J. Liq. Chrom. Rel. Tech. 22, 1457-1471. Murashige, T., Skoog, F., 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. 15, 473-497.
- Szopa, A., Kokotkiewicz, A., Kubica, P., Banaszcak, P., Wojtanowska-Krośniak, A., Krośniak, M., Marzec-Wróblewska, U., Badura, A., Zagrodzki, P., Bucinski, A., Luczkiewicz, M., Ekiert, H., 2017. Comparative analysis of different groups of phenolic compounds in fruit and leaf extracts of *Aronia* sp.: *A. melanocarpa*, *A. arbutifolia*, and *A. × prunifolia* and their antioxidant activities. Eur. Food Res. Technol. 243, 1645-1657.

* Presenting author. E-mail address: halina.ekiert@uj.edu.pl (Halina Ekiert)

S1.P130 Acridone alkaloids and lignans from the leaves of *Luvunga scandens* (Rutaceae)

Nguyen-Bien-Thuy Bui^{a,b,c,*}, Thanh-Hoa Vo^c, Chia-Ching Liaw^b, Chih-Hua Chao^{d,e}, Hui-Chi Huang^f, Ching-Yeu Chen^g, Yu-Chi Lin^b, Mei-Chuan Chen^a

^a Ph.D. Program in Clinical Drug Development of Herbal Medicine, College of Pharmacy, Taipei Medical University, Taipei, Taiwan

^b National Research Institute of Chinese Medicine, Taipei, Taiwan

^c School of Medicine, Vietnam National University Ho Chi Minh City, Ho Chi Minh, Vietnam

^d School of Pharmacy, China Medical University, Taichung, Taiwan

^e Chinese Medicine Research and Development Center, China Medical University Hospital, Taichung, Taiwan

^f Department of Chinese Pharmaceutical Sciences and Chinese Medicine Resources China Medical University, Taichung, Taiwan

^g Department of Physical Therapy, Tzu-Hui Institute of Technology, Pingtung, Taiwan

Luvunga scandens (Roxb.) Buch. Ham. is a species with a restricted range can be found in tropical rain forests in Vietnam and Thailand. This species is one of just twelve in the world that belong to the genus *Luvunga*, which has a wide range of chemical components such as coumarins, triterpenes, flavonoids, and alkaloids; in addition, exhibiting a variety of biological properties as cytotoxicity, antibacterial, anti-inflammatory, etc (Nguyen et al., 2017); (Nguyen et al., 2021). In the ongoing pursuit of novel phytochemistry and bioeffects chemicals from genus *Luvunga*, we made collections of *L. scandens* from Vietnam. Together with 13 known acridone alkaloids (Fig. 1) and 19 known lignans, the ethanol extract of the leaves delivered a previously unknown acridone alkaloid and 9 new lignans.




Fig. 1. Structural framework of acridone alkaloid

The structures of isolated compounds were identified by detailed spectroscopic data, including 2D- NMR (COSY, HMQC, HMBC, and NOESY), HRESIMS, X-ray crystallographic analysis, and comparison with reported data (Tran et al., 2019). All new isolates' radical scavenging activity, anti- inflammatory and cytotoxicity activities were evaluated.

Keywords: *Luvunga scandens*, Rutaceae, cridone alkaloid, lignan, cytotoxicity

References

- Nguyen, H.M.; Nguyen, N.Y.T.; Chau, N.T.N.; Nguyen, A.B.T.; Tran, V.K.T.; Hoang, V.; Le, T.M.; Wang, H.-C.; Yen, C.-H. 2021. Bioassay-Guided Discovery of Potential Partial Extracts with Cytotoxic Effects on Liver Cancer Cells from Vietnamese Medicinal Herbs. *Processes*. 9, 1956.
- Nguyen, T. P., Minh, N. P., Dat, T. B., Le, T. D., Tran Nguyen, M. A., Pham, T. N. T., Tri, D. M. 2017. Limonoid from the rhizomes of *Luvunga scandens* (Roxb.) Buch. Ham. *Nat. Prod. Res.* 31, 2281-2285.
- Tran, N. M. A., Do, T. H. T., Truong, L. H., Le, D. T., Phan, M. N., Pham, N. K. T., Nguyen, T. P. 2019. Acridone alkaloids from the rhizomes of *Luvunga scandens* (Roxb.) Buch. Ham. *Nat. Prod. Res.* 33, 2176-2181.

* Presenting author. E-mail address: d339111002@tmu.edu.tw (Nguyen-Bien-Thuy Bui)

S1.P131 *Schisandra henryi* bioreactor grown *in vitro* culture as a valuable raw material – biotechnological, phytochemical and biological studies

Karolina Jaferník^a, Paweł Kubica^a, Michał Dziurka^b, Łukasz Kulinowski^c, Izabela Korona-Główniak^d, Hosam O. Elansary^e, Piotr Waligórski^b, Krystyna Skalicka-Woźniak^c, Agnieszka Szopa^{a,*}

^a Department of Pharmaceutical Botany, Medical College, Jagiellonian University, Medyczna 9 str, 30-688 Kraków, Poland

^b Polish Academy of Sciences, The Franciszek Górski Institute of Plant Physiology, Niezapominajek 21 str, 30-239 Kraków, Poland

^c Department of Natural Products Chemistry, Medical University of Lublin, Chodźki 1 str, 20-093 Lublin, Poland

^d Department of Pharmaceutical Microbiology, Medical University of Lublin, 1 Chodźki str., 20-093 Lublin, Poland

^e Department of Plant Production, College of Food & Agriculture Sciences, King Saud University, P.O. Box 2460 str, Riyadh 11451, Saudi Arabia

Schisandra henryi C. B. Clarke is endemic to the Yunnan Province in China and is a plant species known in traditional Chinese medicine (Szopa et al., 2019). *In vitro* culture of *S. henryi* microshoot propagation (acc. to Jaferník et al., 2020) in Plantform bioreactors (temporary immersion systems, TIS) was investigated. The phytochemical analyses of dibenzocyclooctadiene lignans, aryltetralin lignans and neolignans were carried out using UHPLC-MS/MS, and HPLC-DAD methods for qualitative and quantitative determinations. The dominant compounds were isolated using CPC chromatography. The highest content of compounds was found in microshoots extracts (max. 543.99 mg/100g DW). The major compound was schisantherin B (390.16 mg/100g DW). The anti-inflammatory activity tests (COX-1, COX-2, sPLA2 and LOX-15 inhibition), showed the strong activity against COX-1 and -2 (for 177 mg/mL inhibition percentage was 76% and 66%, respectively). The studies showed also the high antioxidant power (FRAP, CUPRAC and DPPH assays), and good antiproliferative activities (Jurkat, MCF-7, HT-29 and HeLa lines) of extracts. Antimicrobial studies against gram+ and gram- bacteria and fungi proved especially high activity towards *H. pylori* (MIC and MBC 0.625 mg/mL). To sum up, the bioreactor grown *in vitro* culture of *S. henryi* may be a cost-effective, innovative alternative to the rare naturally grown medicinal-raw material.

Keywords: plant biotechnology, *in vitro* cultures, PlantForm temporary immersion system, dibenzocyclooctadiene lignans, pharmacological activity

References

- Szopa, A.; Barnaś, M.; Ekiert, H. Phytochemical Studies and Biological Activity of Three Chinese *Schisandra* Species (*Schisandra sphenanthera*, *Schisandra henryi* and *Schisandra rubriflora*): Current Findings and Future Applications. *Phytochemistry Reviews* 2019, 18, 109–128.
- Jaferník, K.; Szopa, A.; Barnaś, M.; Dziurka, M.; Ekiert, H. *Schisandra henryi* C. B. Clarke *in vitro* Cultures: A Promising Tool for the Production of Lignans and Phenolic Compounds. *Plant Cell Tissue Organ Cult* 2020, 143, 45–60.

* Presenting author. E-mail address: a.szopa@uj.edu.pl (Agnieszka Szopa)

S1.P132 The phenolic profile and antioxidant potential of different types of *Schisandra henryi* *in vitro* cultures

Karolina Jafernika^a, Paweł Kubica^a, Marta Sharafan^b, Aleksandra Kruk^c, Magdalena Anna Malinowska^b, Sebastian Granica^c, Agnieszka Szopa^{a,*}

^a Chair and Department of Pharmaceutical Botany, Jagiellonian University, Medical College, Medyczna 9 Str, 30-688 Cracow, Poland

^b Department of Organic Chemistry and Technology, Faculty of Chemical Engineering and Technology, Cracow University of Technology, Warszawska 24 Str, 31-155 Cracow, Poland

^c Department of Pharmacognosy and Molecular Basis and Phytotherapy, Medical University of Warsaw, Banacha 1 Str, 02-097 Warsaw, Poland

Schisandra henryi C. B. Clarke is an endemic species, used as medicinal plant in Asia, but it is practically unknown in European and American countries. The small number of studies on the *S. henryi* has been focused on the phytochemical profiling and biological activity (Szopa et al., 2019). The aim of the study was to investigate the polyphenol profile and to assess the antioxidant power of the extracts from the *in vitro* cultures of *S. henryi*. Under the study the different types of *in vitro* cultures were tested: agar microshoot and callus, microshoot in bioreactor Plantform, agitated microshoot and suspension cultures (Jafernika et al., 2020). The UHPLC-DAD-ESI-MS³ and HPLC- DAD methods were used for the analyses of extracts. The total content of polyphenols (TPC) was measured using the Folin-Ciocalteu method. The antioxidant potential was assessed using 2,2- diphenyl-1-picrylhydrazine hydrate (DPPH), ferric reducing antioxidant power (FRAP) and ferrous ion chelating (FIC) assays. In the extracts the procyanidins, phenolic acids (neochlorogenic acid, caffeic acid, protocatechuic acid) and catechin were estimated. The highest content of phenolic compounds was found for microshoot agar culture (max. total content: 229.87 mg/100 g DW) and agitated culture (max. total content: 22.82 mg/100 g DW). The max. TPC was equal 1240.51 mg GAE/100g DW (agar microshoot culture). The highest antioxidant potential was indicated for agar microshoot culture (90.08% of inhibition and 59.31 nM/L TEAC, respectively). To sum up, the performed studies are innovative and showed that extracts from *S. henryi* microshoot cultures are the valuable source of polyphenolic compounds, which are characterized by a very strong antioxidant potential.

Keywords: plant biotechnology, *in vitro* cultures, phenolic compounds, antioxidant potential

References

- Szopa, A.; Barnaś, M.; Ekiert, H. Phytochemical Studies and Biological Activity of Three Chinese *Schisandra* Species (*Schisandra sphenanthera*, *Schisandra henryi* and *Schisandra rubriflora*): Current Findings and Future Applications. *Phytochemistry Reviews* 2019, 18, 109–128.
- Jafernika, K.; Szopa, A.; Barnaś, M.; Dziurka, M.; Ekiert, H. *Schisandra henryi* C. B. Clarke *in Vitro* Cultures: A Promising Tool for the Production of Lignans and Phenolic Compounds. *Plant Cell Tissue Organ Cult* 2020, 143, 45–60.

* Presenting author. E-mail address: a.szopa@uj.edu.pl (Agnieszka Szopa)

S1.P133 Isolation and characterisation of stigmasterol and β -sitosterol from *Stachytarpheta jamaicensis* (L) Vahl (Verbenaceae) leaves of Eastern Nigeria

Odoh Uchenna Estella^{a,*}, Payare L. Sangwan^b, Santosh Kumar Rath^c

^a Department of Pharmacognosy and Environmental Medicines, Faculty of Pharmaceutical Sciences, University of Nigeria, Nsukka, Nigeria

^b Bioorganic Chemistry Division, CSIR-Indian Institute of Integrative Medicine, Canal Road Jammu, 180001, India

^c School of Pharmaceuticals and Population Health Informatics, Faculty of Pharmacy, DIT University, Dehradun, Uttarakhand-248009, India

Stachytarpheta jamaicensis (L.) Vahl (Verbenaceae) is a potent medicinal plant commonly used in treatment of infections in West African (1). This study was to isolate and characterize the bioactive constituents from the leaves of *Stachytarpheta jamaicensis*. The powdered leaves were extracted with dichloromethane: methanol (1:1) using cold maceration method. Column chromatography of the crude extract lead to a number of fractions. TLC fingerprinting and the spraying reagent (Concentrated H₂SO₄ and vanillin in methanol) were used to identify the fraction containing phytosterols. The isolation and purification afforded white crystalline powder which was subjected to physical, chemical and spectral identification by IR, ¹H-NMR, 2D-NMR and ¹³C-NMR. The compound was identified as a mixture of stigmasterol and β -sitosterol (Fig. 1).




Fig. 1: Structures of stigmasterol and β -sitosterol.

Stachytarpheta jamaicensis contains stigmasterol and β -sitosterol which are responsible for the various therapeutic activities of the plant.

Keywords: *Stachytarpheta jamaicensis*, stigmasterol, β -sitosterol, characterization

*Presenting author Email address: uchenna.odoh@unn.edu.ng (Odoh, Uchenna Estella)

S1.P134 UHPLC-HRMS profiling of sulphated constituents in different plant parts of *Salvadora persica*

Eva-Maria Pferschy-Wenzig^{a,*}, Ahmed M.A. Mazen^b

^a University of Graz, Institute of Pharmaceutical Sciences, Pharmacognosy, Beethovenstraße 8, 8010 Graz, Austria

^b Botany and Microbiology Department, Faculty of Science, Sohag University, Sohag, Egypt

Salvadora persica L. (Salvadoraceae) is traditionally used for oral health. Chewing sticks prepared from its roots and twigs are a widespread traditional oral hygiene device in Middle Eastern, African and Asian countries, and the leaves are used to alleviate gum problems (Khatak et al, 2010; Ahmad and Rajagopal, 2014). The plant is known to contain flavonoids, phenolic acids, volatile oil, alkaloids, and some sulphur-containing compounds (Aumeeruddy et al, 2018). In our previous work, we detected numerous sulphated constituents in *S. persica* leaf extracts (Kobetitsch et al., 2024). This study aimed to systematically profile sulphated constituents in methanolic *S. persica* leaf, twig and root extracts. Analysis was performed by UHPLC-QExactive-MS/MS, and sulphates were tracked on the basis of the MS/MS-fragment m/z 96.960 $[\text{HSO}_4]^-$. Around 60 peaks with sulphate moiety were detected. Almost all of them displayed an additional m/z 241.002 $[\text{hexosylsulphate-H}_2\text{O}]^-$ fragment, indicating glycosylsulphate structures. The profile of the root extract was dominated by the only detected glucosinolate glucotropaeolin, which also occurred in the leaves and twigs at lower levels. 15 peaks were annotated as aromatic (phenyl, benzyl-, phenethyl-) glycosylsulphates; two among them that occurred in all three extracts were annotated as isomers of the known benzylglycosylsulphate salvadoside (Kamel et al., 1992). 24 peaks that displayed 19 C-atoms in their molecular formula were annotated as megastimane glycosylsulphates, detected for the first time in our previous work (Kobetitsch et al., 2024). They displayed highest levels and variety in leaves, followed by stems and roots. Their isolation and structural elucidation are under progress.

Keywords: *Salvadora persica*, UHPLC-HRMS, secondary metabolites

References

- Ahmad H, Rajagopal K., 2014. *Salvadora persica* L. (Meswak) in dental hygiene. The Saudi Journal for Dental Research. 5, 130–134.
- Aumeeruddy MZ, Zengin G, Mahomoodally MF., 2018. A review of the traditional and modern uses of *Salvadora persica* L. (Miswak): Toothbrush tree of Prophet Muhammad. J Ethnopharmacol. 213, 409–444.
- Kamel MS, Ohtani K, Assaf MH, Kasai R, El-Shanawani MA, Yamasaki K, Ali AA, Tanaka O. 1992. Lignan glycosides from stems of *Salvadora persica*. Phytochemistry. 31:2469–2471.
- Khatak M, Khatak S, Siddiqui AA, Vasudeva N, Aggarwal A, Aggarwal P, 2010. *Salvadora persica*. Pharmacogn Rev. 4, 209–214.
- Kobetitsch, S., Gierlikowska, B., Kunert, O., Mazen, A.M.A., Raab, P., Kretschmer, N., Donolo, C., Pirker, T., Bauer, R., Kiss, A.K., Pferschy-Wenzig, E.M., 2024. *Salvadora persica* leaves: Phytochemical profile and *in vitro*- inhibitory activity on inflammatory mediators implicated in periodontal disease. Pharmaceutical Biology, under review.

* Presenting author. E-mail address: eva-maria.wenzig@uni-graz.at (Eva-Maria Pferschy-Wenzig)

S1.P135 Koninginins DS A-C, novel polyketides from *Trichoderma koningii*, a soil fungus collected from medicinal herbal garden

Dahyeon Lee^a, Eun-Sook Kim^a, Jae Sang Han^b, Bang Yeon Hwang^b, Jin Woo Lee^{a,*}

^a College of Pharmacy, Duksung Women's University, Seoul 01369, Republic of Korea

^b College of Pharmacy, Chungbuk National University, Cheongju 28160, Republic of Korea

Most of sources of foods, health supplements and natural medicines consumed by humans are cultivated in the soil. So, beneath the ground, soil fungi may influence bioactive metabolites (or their biosynthesis pathway) of medicinal plants or microorganisms through their interactions. Based on this assumption and curiosity, we collected soil samples from herbs garden in Duksung Women's University where various medicinal plants having used for Korean traditional therapeutics grow every year. Totally, 10 fungi were isolated from the soil sample of herb garden, which are expected to produce bioactive compounds with novel scaffolds. Among these, the first strain (DS1-1) was identified as *Trichoderma koningii*. *Trichoderma* is one of the most common genera of soil fungi including various species known for their biological effects such as anti-cancer, anti-microbial and biological control agents (Cutler et al., 1989). Among this genus, *T. koningii* has been known as rich sources of koninginins (polyketides) and peptaibols, which have been reported various biological activities (Lang et al., 2015). As part of our continuing study for the discovery of soil fungi-derived novel bioactive molecules, three new koninginins, koninginins DS A-C (**1-3**), together with 6 known koninginin derivatives (**4-9**) and two known peptaibols (**10-11**), were isolated from the EtOAc extract of *T. koningii* fermented on oat-based cereal, Cheerios[®] (Fig. 1). Their relative structures were determined by *J* value and chemical shifts comparing with previously reported known molecules, and biosynthetic consideration. The elucidation of the absolute chemistry is remained, so we will establish by comparison experimental ECD spectrum with calculated one.




Fig. 1. Structures of isolated metabolites **1-11** from *T. koningii*

Keywords: soil fungi, *Trichoderma*, *T. koningii*, koninginins, peptaibols

References

- Cutler, H. G., Himmelsbach, D. S., Arrendale, R. F., Cole, P. D., Cox, R. H., 1989, Koninginin A: A novel plant growth regulator from *Trichoderma koningii*, Agricultural and Biological Chemistry 10, 2605-2611.
- Lang, B., Li, J., Zhou, X., Chen, Y., Yang, Y., Li, X., Zeng, Y., Zhao, P., 2015. Koninginins L and M, two polyketides from *Trichoderma koningii* 8662. Phytochemistry Letters, 11, 1-4.

* Presenting author. E-mail address: jwlee22@duksung.ac.kr (Jin Woo Lee)

S1.P136 Anti-parkinson pyridone alkaloids from an endolichenic fungus, *Thielavia* sp. CNC14

Jin Won Choi^{a,*}, Chaesun Kwon^b, Jin Woo Lee^c, Jae-Seon Hur^d, Min-kyoo Shin^b, Sang Hee Shim^a

^a Natural Products Research Institute, College of Pharmacy, Seoul National University, Seoul 08826, Republic of Korea

^b College of Pharmacy and Research Institute of Pharmaceutical Sciences, Seoul National University, Seoul 08826, Republic of Korea

^c College of Pharmacy, Duksung Women's University, Seoul 01369, Republic of Korea

^d Korean Lichen Research Institute, Sunchon National University, Suncheon 57922, Republic of Korea

4-Hydroxy-2-pyridone alkaloids have attracted much attention in the scientific community due to their distinctive structures as well as diverse biological effects. In an effort to discover this class of compounds from fungi, endolichenic fungal extracts were screened by analysis of LC-UV-MS profiles, which afforded a strain, *Thielavia* sp. CNC14 with characteristic UV pattern for 4-hydroxy-2-pyridone alkaloids using our in-house library. This result prompted us to culture the strain in large scale, followed by several steps of chromatographic methods, which led to the isolation of four new (**1–4**) and ten known (**5–14**) compounds. The structures of the isolated compounds were elucidated by spectroscopic methods such as NMR and MS together with ECD calculation. Stereochemistry of the new compounds (**1–4**) were established using ROESY and comparison of their electronic circular dichroism spectra with those of calculated data. Interestingly, compounds **1** and **2** were cyclized in opposite direction to that of tolypyridone A (**5**) and predicted to be biosynthesized from reduced tolypyridone C (**7**) through hetero-Diels-Alder reaction. Moreover, compounds **4** and **5** had reduction and dehydration in comparison to compound **7**. Among the isolated compounds, **4** significantly protected neuronal cells against treatment with MPP⁺, a Parkinsonian neurotoxin, in an *in vitro* Parkinson's disease model.




Fig. 1. Structures of isolated new compounds **1–4**

Keywords: endolichenic fungi, 4-hydroxy-2-pyridone alkaloid, anti-parkinson

* Presenting author. E-mail address: climax0629@snu.ac.kr (Jin Won Choi)

S1.P137 The anticancer activity and DNA synthesis inhibition potential of *Teucrium sandrasicum* and *Teucrium divaricatum* subsp. *graceum*

Nagehan Saltan^{a,*}, Derya Osmaniye^b, Yavuz Bulent Kose^a

^a University of Anadolu, Faculty of Pharmacy, Department of Pharmaceutical Botany, Eskisehir, 26470, Turkey

^b University of Anadolu, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Eskisehir, 26470, Turkey

This presentation will cover the evaluation of the cytotoxic activities of different extracts (methanol, ethanol, water and n-hexane) prepared from *Teucrium sandrasicum* O.Schwarz and *Teucrium divaricatum* subsp. *graceum* (Celak.) Bornm. *T. sandrasicum*, one of the species belonging to the genus *Teucrium*, is a species endemic to Turkey. *T. sandrasicum*, a perennial plant that grows in the serpentine regions of Muğla Sandras Mountain, is used among the people in our country as a diuretic, diaphoretic, tonic, antipyretic, antidiabetic, antispasmodic and cholagogic (Aksoy-Sagirli et al., 2015). *T. divaricatum* subsp. *graceum*. is distributed on rocky, stony, gravelly, calcareous and arid slopes. It is popularly known as ‘Böceotu’ in Turkey (Guner et al., 2012). *Teucrium* is used for the treatment of coughs, abdominal pain complaints, diseases of the urinary tract, gallbladder, and kidneys, for fevers, colds and stomach disorders, as well as for an external cicatrisant (Formisano et al., 2010; Asghari et al., 2020). There is a lot of interest in the discovery and development of herbal raw materials that can efficiently cause apoptosis and have cytotoxicity in tumor cells. In previous studies, extracts and compounds of different species within the genus *Teucrium* have been evaluated for cytotoxicity and anticancer activity in various cell lines (Stankovic et al., 2011). The cytotoxic effects of different extracts of the plants were evaluated MTT assay against different cell lines including C6, A549, MCF-7, and NIH3T3. Methanol extract of air-dried aerial parts of *T. divaricatum* subsp. *graceum* (TDM) exhibited significant cytotoxicity (IC₅₀=17.09 µg/mL) against MCF-7 cell line. BrdU (5-bromo-20-deoxyuridine) cell proliferation method was applied to analyze the effects of this extract on the proliferation of MCF-7 cells. Three concentrations, IC₅₀/2, IC₅₀, and 2xIC₅₀, were studied. It was determined that the percentage of DNA synthesis inhibition of TDM extract increased depending on the concentration and incubation time.

Keywords: *Teucrium*, cytotoxicity, DNA synthesis inhibition, Lamiaceae

References

- Aksoy-Sagirli, P., Ozsoy, N., Ecevit-Genc, G., Melikoglu, G., 2015. *In vitro* antioxidant activity, cyclooxygenase-2, thioredoxin reductase inhibition and DNA protection properties of *Teucrium sandrasicum* L. Ind. Crops Prod. 74, 545– 550.
- Asghari, A.A., Mokhtari-Zaer, A., Niazmand, S., Mc Entee, K., Mahmoudabady, M., 2020. Anti-diabetic properties and bioactive compounds of *Teucrium polium* L. Asian Pac. J. Trop. Biomed. 10, 433.
- Formisano, C., Napolitano, F., Rigano, D., Arnold, N.A., Piozzi, F., Senatore, F., 2010. Essential oil composition of *Teucrium divaricatum* Sieb. ssp. *villosum* (Celak.) Rech. fil. growing wild in Lebanon. J. Med. Food. 13, 1281– 1285. Guner, A., Aslan, S., Ekim, T., Vural, M., Babac, M.T. (Eds.) 2012. Türkiye Bitkileri Listesi (Damarlı Bitkiler). Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul.
- Stankovic, M.S., Curcic, M.G., Zizic, J.B., Topuzovic, M.D., Solujic, S.R., Markovic, S.D., 2011. *Teucrium* plant species as natural sources of novel anticancer compounds: antiproliferative, proapoptotic and antioxidant properties. Int. J. Mol. Sci. 12, 4190-4205.

* Presenting author. E-mail address: ndagdeviren@anadolu.edu.tr (Nagehan Saltan)

S1.P138 Kronopolitides A-D, a new family of 36-membered polyol macrolides with antiparasitic activity

Mercedes Pérez-Bonilla^{a,*}, Marina Sánchez-Hidalgo^a, Ignacio González^a, Jesús Martín^a, David Shum^b, Joo Hwan No^b, Jean-Robert Ioset^c, Fernando Reyes^a, Olga Genilloud^a

^a Fundación MEDINA, Avda. del Conocimiento 34, 18016 Armilla, Granada, Spain

^b Institut Pasteur Korea, 16 Daewangangyo-ro 712 beon-gil, Bundang-gu, Seongnam-si, Gyeonggi-do, 13488 Rep. of Korea

^c Drugs for Neglected Diseases initiative, 15 Chemin Camille-Vidart, 1202 Geneva, Switzerland

The neglected tropical diseases Leishmaniasis and American Trypanosomiasis (Chagas disease) caused by the parasites *Leishmania* spp. and *Trypanosoma cruzi*, respectively, cause thousands of deaths worldwide, and in recent years they have also emerged as a health concern in developed countries. New therapeutic solutions are required due to increasing resistance and adverse effects of existing treatments (No, 2016; Pérez-Molina et al., 2018). Natural products (NPs) have been historically a rich reservoir of new bioactive compounds, originating new drugs used today in clinical practice (Newman et al., 2020). With the aim of discovering new NP scaffolds with novel mechanism of action (MoA) against *Leishmania* and *T. cruzi*, 40K representative microbial extracts from MEDINA's NP collection have been tested. The acetone extract of a strain identified by 16S rRNA as *Streptomyces kronopolitis* (CA-143054) displayed activity against *T. cruzi*. Bioassay-guided isolation using SP207ss resin column and preparative reversed-phased HPLC led to the isolation of four novel polyketide macrolides, kronopolitides A-D. The planar structures of the new family of 36-membered polyol macrolides were determined using a combination of 1D and 2D NMR data, HRMS, and ESI-qTOF-MS/MS. The absolute configuration of their chiral centers was established using NOE, *J*-based configuration analysis, comparison with NMR data of structurally related macrolides, and the bioinformatic analysis of the biosynthetic gene cluster. The antiparasitic activity, measured as the number of host cells infected with parasite in their cytoplasmic region, displayed IC₅₀ values ranging from 0.16-1.27 μ M for *T. cruzi*, being kronopolitide D the most potent compound. To characterize their MoA, the new macrolides will be profiled in an innovative high content imaging screen parasite painting assay.

Keywords: *Streptomyces kronopolitis*, *Trypanosoma cruzi*, *Leishmania*, polyketide macrolides, kronopolitides

Acknowledgements: This research was funded by "la Caixa" Foundation grant number HR20-00584. We also thank UK aid, UK, the Dutch Ministry of Foreign Affairs (DGIS), the Netherlands; and the Federal Ministry of Education and Research (BMBF) through KfW, Germany, for supporting via DNDi the bioassays run at Institut Pasteur Korea.

References

- Newman, D.J., Cragg, G.M., 2020. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J. Nat. Prod.* 83, 770–803.
- No, J. H., 2016. Visceral leishmaniasis: Revisiting current treatments and approaches for future discoveries. *Acta Trop.* 115, 113-123.
- Pérez-Molina, J.A., Molina, I., 2018. Chagas disease. *Lancet* 391, 82-94.

* Presenting author. E-mail address: mercedes.perez@medinaandalucia.es (Mercedes Pérez-Bonilla)

S1.P139 A correlative metabologenomics approach for the elucidation of the biosynthetic pathway of 3-(3-Furyl)alanine-containing endolides produced in *Stachylidium bicolor*

Tim Berger*, Raphael Reher

Institute for Pharmaceutical Biology and Biotechnology, Department of Pharmacy, Philipps - University Marburg, Robert-Koch-Straße 4, 35037 Marburg, Germany

The marine sponge-derived fungus *Stachylidium bicolor* 293 K04 is a prolific producer of bioactive natural products, particularly the rare 3-(3-furyl)-alanine containing cyclic tetrapeptides known as endolides. Leveraging molecular networking integrated with targeted MassQL queries we've recently unveiled the intricate amino acid makeup of numerous endolides, hinting at their biosynthesis through a combinatorial process (Berger et al., 2024). The production of 3-(3-furyl)-alanine moieties in rhizonin A was recently proven to originate from oxidative cleavage and rearrangement of tyrosine in the endofungal symbiont *Mycetohabitans endofungorum*, catalyzed by a dioxygenase (Rhzb, Ehinger et al., 2023). This revelation raises intriguing questions about the potential bacterial contribution to endolide production, especially considering reports of *Burkholderia contaminans* within *S. bicolor* (Almeida et al., 2018). Despite our efforts, confirming this hypothesis has proven elusive. On the contrary, after sequencing and assembly of the genome of *Stachylidium bicolor* 293 K04 we found the putative biosynthetic gene cluster for endolide production. Furthermore, while feeding experiments with the fungus were generally successful (incorporation of ¹³C-labeled phenylalanine, and leucine in respective endolides) in our case 3-(3-furyl)-alanine did not show ¹³C- enrichment in LC-MS experiments after feeding ¹³C-labeled-tyrosine, suggesting an alternative biosynthesis. Feeding experiments with [U-¹³C] glycerol, however, point to a biosynthesis of the unusual amino acid via the shikimate pathway and a putatively new cyclic precursor intermediate (El Maddah et al., 2016) Through a correlative metabologenomics approach, we're tirelessly working to unravel the enigma surrounding the origin of endolides—whether they stem from fungal or bacterial sources.

Keywords: *Stachylidium bicolor*, cyclic tetrapeptide, 3-(3-furyl)alanine, biosynthesis

References

- Almeida, C., Silva Pereira, C., Gonzalez-Menendez, V., Bills, G., Pascual, J., Sánchez-Hidalgo, M., Kehraus, S., Genilloud, O., 2018. Unveiling Concealed Functions of Endosymbiotic Bacteria Harbored in the Ascomycete *Stachylidium Bicolor*. *Appl. Environ. Microb.* 84 (15).
- Berger, T., Alenfelder, J., Steinmüller, S., Heimann, D., Gohain, N., Petras, D., Wang, M., Berger, R., Kostenis, E., Reher, R., 2024. A MassQL-Integrated Molecular Networking Approach for the Discovery and Substructure Annotation of Bioactive Cyclic Peptides. *J. Nat. Prod.*
- Ehinger, F. J., Niehs, S. P., Dose, B., Dell, M., Krabbe, J., Pidot, S. J., Stinear, T. P., Scherlach, K., Ross, C., Lackner, G., Hertweck, C., 2023. Analysis of Rhizonin Biosynthesis Reveals Origin of Pharmacophoric Furylalanine Moieties in Diverse Cyclopeptides. *Angew. Chem. Int. Edit.* 62 (42).
- El Maddah, F., Kehraus, S., Nazir, M., Almeida, C., König, G. M., 2016. Insights into the Biosynthetic Origin of 3-(3- Furyl)alanine in *Stachylidium* Sp. 293 K04 Tetrapeptides. *J. Nat. Prod.* 79 (11), 2838–2845.

* Presenting author. E-mail address: tim.berger@pharmazie.uni-marburg.de (Tim Berger)

S1.P140 Discovery and structure elucidation of novel antagonistic cyclic lipopeptides for the control of fungal agricultural pathogens

Leah Gauthier^{a,b,*}, Tim Dumonceaux^c, Brian Wagner^a, Christopher W. Kirby^b

^a Department of Chemistry, University of Prince Edward Island, 550 University Avenue, Charlottetown, Prince Edward Island, C1A 4P3, Canada

^b Charlottetown Research and Development Center, Agriculture and Agri-Food Canada, 440 University Avenue, Charlottetown, C1A 4N6, Prince Edward Island, Canada

^c Saskatoon Research and Development Centre, Agriculture and Agri-Food Canada, 107 Science Place, Saskatoon, S7N 0X2, Canada

Novel antifungal cyclic lipopeptides (CLPs) produced by a *Pseudomonas tensinigenes* strain have been found to be related to Amphisin (1), and its family of CLPs (Gotze, S. et al., 2017) with applications as biopesticides. While pest and pathogen management products help ensure crop quality and yield in conventional farming, the widespread over-use of synthetic pesticides have been found to have many detrimental non-target effects. These include decreasing soil fertility, contaminating water systems and triggering the development of fungicide-resistant pathogen strains. Biopesticides, which consist of microorganisms and their natural products (NPs) are a less harmful alternative to synthetic pesticides as the NPs already exist in the environment with developed degradation pathways, eliminating the environmental concerns associated with synthetic products. A microbial biopesticide screening study of using thin-layer chromatography direct bioautography (TLC-DB) assays resulted in the discovery and subsequent structure elucidation of 2, and 3. Work involving the discovery and structure elucidation of these compounds by TLC-DB, mass spectrometry and nuclear magnetic resonance will be discussed.




Figure 1. The proposed structures of two novel biopesticide candidates, 1, and 2, elucidated by nuclear magnetic resonance and mass spectrometry.

Keywords: Cyclic lipopeptides, biopesticides, NMR, Mass spectrometry, structure elucidation.

References

Gotze, S., Herbst-Irmer, R., Klapper, M., Górls, H., Schneider, K. R. A., Barnett, R., Burks, T., Neu, U., Stallforth, P., 2017. Structure, Biosynthesis, and Biological Activity of the Cyclic Lipopeptide Anikasin. *ACS Chem Biol.* 12, 2498-2502.

* Presenting author. E-mail address: lkgauthier@upei.ca (Leah Gauthier)

S1.P141 Hydrogen sulfide is a key mediator in the antioxidant effects of osthole on mouse corpus cavernosum and liver

Elif Alan-Albayrak*, Ozan Mert, Gulcan Demir, Gulnur Sevin

Department of Pharmacology, Faculty of Pharmacy, Ege University, 35040, Izmir, Türkiye

Osthole is a major bioactive coumarin found in *Cnidium monnieri* and is also isolated from *Prangos* species traditionally used as aphrodisiacs in Türkiye (Albayrak et al., 2023). Erectile dysfunction (ED), affecting 52% of men, is exacerbated by risk factors like diabetes and hypertension, which cause endothelial dysfunction by increasing oxidative stress (Agarwal et al., 2006). Given the ineffectiveness of PDE-5 inhibitors in cases of endothelial dysfunction, there is a need for treatments independent of the endothelium. Hydrogen sulfide (H₂S), produced from L-cysteine mainly in the liver, enhances erectile function endothelium-independently. Osthole demonstrates antioxidant effects by increasing antioxidants (catalase, etc.) and activating the Keap1/Nrf2 pathway, similar to H₂S mechanisms. Therefore, we aimed to investigate the osthole-H₂S relationship for the first time. We examined the effects of osthole (30 µM, 30 min) on H₂S production by measuring real-time H₂S synthesis using an H₂S microsensor in the presence/absence of pyrogallol (0.1 mM, 5 min)-induced oxidative stress in mouse corpus cavernosum (MCC) and liver (ML). Additionally, we evaluated the role of H₂S in the antioxidant effects of osthole by measuring reactive oxygen species (ROS) using chemiluminescence assays (luminol/lucigenin) in MCC/ML. Osthole increased L-cysteine-induced H₂S synthesis in healthy MCC/ML ($P < 0.01$, ANOVA, $n = 6$) and inhibited the pyrogallol-induced reduction in H₂S production in MCC/ML ($P < 0.05$, ANOVA, $n = 6$). Osthole decreased the pyrogallol-induced increase in ROS levels ($P < 0.01$, ANOVA, $n = 6$), these effects were inhibited by the H₂S synthesis inhibitor aminooxyacetic acid ($P < 0.05$, ANOVA, $n = 6$). These results indicate that osthole protects MCC/ML from oxidative stress by increasing H₂S synthesis. Osthole could be an effective drug candidate, especially in oxidative-stress-induced endothelial dysfunction-related ED and liver diseases.

This project is supported by the Ege University Office of Scientific Research Projects (Grant number: 23603).

Keywords: oxidative stress, hydrogen sulfide, erectile function, corpus cavernosum, liver

References

- Agarwal, A., Nandipati, K.C., Sharma, R.K., Zippe, C.D., Raina, R., 2006. Role of oxidative stress in the pathophysiological mechanism of erectile dysfunction. *J Androl.* 27, 335-47.
- Albayrak, G., Demir, S., Kose, F.A., Baykan, S., 2023. New coumarin glycosides from endemic *Prangos heyneana* H. Duman & M.F. Watson. *Nat Prod Res.* 37, 227-239.

* Presenting author. E-mail address: elifalan1293@gmail.com (Elif Alan-Albayrak)

S1.P142 Isoimperatorin induces relaxation in mice corpus cavernosum via increasing hydrogen sulfide (H₂S) synthesis

Emine Nur Ozbek^a, Nazlican Belen^a, Elif Alan-Albayrak^a, Tugce Demiroz^b, Gokay Albayrak^{c,*}, Serdar Demir^b, Gulnur Sevin^a, Gunay Yetik Anacak^d, Sura Baykan Ozturk^b

^a Department of Pharmacology, Faculty of Pharmacy, Ege University, 35040, Izmir, Türkiye

^b Department of Pharmaceutical Botany, Faculty of Pharmacy, Ege University, 35040, Izmir Türkiye

^c Department of Pharmaceutical Botany, Faculty of Pharmacy, Izmir Katip Celebi University, 35620, Izmir, Türkiye

^d Department of Pharmacology, Faculty of Pharmacy, University of Acibadem Mehmet Ali Aydınlar, 34752, Istanbul, Türkiye

Prangos sp. roots have been used traditionally as aphrodisiac in Anatolia. Hydrogen sulfide(H₂S), as a novel multi-targeted gasotransmitter with antioxidant, anti-inflammatory, and vasorelaxant effects, plays a regulatory role in erectile function(Srilatha et al., 2007). Previously, we showed that chloroform extracts of *Prangos heyneiae* roots (PHC) induced relaxation on mice corpus cavernosum(MCC) through H₂S(Alan et al., 2022). However, the active compound responsible for the relaxant effect of *P. heyneiae* remained ambiguous. Thus, we aimed to investigate whether the major compound has a relaxing effect on MCC and the role of the H₂S mechanism in its effects. Previously, we determined isoimperatorin(ISO), a furanocoumarin, as the major compound in PHC by HPLC(Arzu et al., 2023) and then the compound was isolated and identified in another study using column chromatography, NMR, and LC-MS(Albayrak et al., 2023). The relaxant effect of ISO was tested in MCC strips by using a myograph in the presence/absence of H₂S synthesis inhibitor aminooxyacetic acid(AOAA, 10⁻²M). Furthermore, time-dependent H₂S formation was measured in L-cysteine(L-cyst, 10mM)-stimulated conditions in the presence/absence of AOAA in MCC homogenates by H₂S sensor. ISO (10⁻⁷-10⁻⁴g/mL) caused relaxation in MCC(*P*<0.001, Anova, n=4-6). AOAA inhibited ISO-induced relaxations(*P*<0.001, Anova, n=6-7). ISO significantly increased L- cyst-stimulated H₂S formation in the MCC homogenates(*P*<0.01, Anova, n=6-8) and this augmentation was significantly inhibited by AOAA(*P*<0.001, Anova, n=4-5). Our study demonstrated that ISO induces relaxation on MCC through H₂S pathway for the first time. As a major compound of PHC, ISO could be responsible for the relaxant effect of this plant. ISO could be a potential drug candidate as an H₂S inducer for the treatment of ED.

Keywords: *Prangos heyneiae*, isoimperatorin, erectile function, hydrogen sulfide

References

- Alan, E., Albayrak, G., Sevin, G., Yetik-Anacak, G., Baykan, S., 2022. Relaxation mechanisms of chloroform root extracts of *Prangos heyneiae* and *Prangos uechtritzi* on mouse corpus cavernosum. *Andrologia*, 54(11), e14604.
- Albayrak G., Demir S., Kose-Aydin F., Baykan S., 2023. New coumarin glycosides from endemic *Prangos heyneiae* H. Duman & MF Watson. *Nat. Prod. Res.*, 37(2), 227-239.
- Arzu, E., Albayrak G., Erguc E., Ecrin E., Tan I., Baykan S., 2024. Investigation of cytotoxic and apoptotic effects of *Prangos heyneiae* H. Duman M.F. Watson extracts on Hepg2 cells. *J. Fac. Pharm. Ankara*, 48(1), 46-55.
- Srilatha, B., Adaikan, P. G., Li, L., Moore, P. K., 2007. Hydrogen sulphide: a novel endogenous gasotransmitter facilitates erectile function. *J. Sex. Med.*, 4(5), 1304-1311.

*Presenting author. E-mail address: albayrakgokay@gmail.com (Gokay Albayrak)

S1.P143 Purple willow (*Salix purpurea* L.) varieties as a source of valuable herbal raw material with a high content of salicylic glycosides

Paweł Sulima^{a,*}, Aleksandra Owczarek-Januszkiewicz^b, Agnieszka Kicel^b, Jerzy Andrzej Przyborowski^a

^a Department of Genetics, Plant Breeding and Bioresource Engineering, University of Warmia and Mazury in Olsztyn, Plac Łódzki 3/432, 10-724 Olsztyn, Poland

^b Department of Pharmacognosy, Medical University of Lodz, 1 Muszyńskiego St., 90-151 Lodz, Poland

Purple willow (*Salix purpurea* L.) bark is considered as valuable herbal raw material, which is characterized by analgesic, antipyretic, anti-inflammatory, antirheumatic, antibacterial and antimicrobial properties. The medicinal properties arise from the unique composition of therapeutic substances, such as salicylic glycosides (SG), flavonoids, phenolic acids, and tannins (Wichtl 2004, Sulima and Przyborowski 2019). Currently, the bark used in the production of natural aspirin, referred to medicinal products manufactured from willow bark, is primarily sourced from various randomly growing forms in highly diverse and uncontrolled conditions. This results in the material's heterogeneity and contamination. Therefore, field cultivation of selected willow varieties under strictly controlled conditions appears to be a simple and effective solution to the above problem (Bubner et al. 2018, Sulima et al. 2021). In this study we conducted a qualitative and quantitative chromatographic analysis of three registered varieties of purple willow (Aspi, Aspira, Cortexa) with the use of UHPLC methods. In the plant material, approximately 40 different polyphenolic compounds were detected, of which six were identified as SG. The presence of salicin and salicortin was confirmed through parallel analyses using standard substance testing. Salireposide, tremulacin, and disalicylortin 1 and 2 were identified based on the comparison of their spectral data. The obtained results showed the high potential of all studied varieties of *S. purpurea*. A high concentration of SG was observed in the bark of ASPIRA ($145.47 \pm 4.18 \text{ mg g}^{-1}$), ASPI ($133.22 \pm 4.41 \text{ mg g}^{-1}$) and CORTEXA ($104.93 \pm 1.69 \text{ mg g}^{-1}$). In this study, the content of SG in leaves, wood and young shoots was also examined. The results indicated the possibility of using whole, several months old shoots as a herbal raw material of *S. purpurea*. This is a completely novel proposal that provides an opportunity for easier acquisition of larger quantities of herbal raw material from purple willow varieties cultivated on arable lands.

Keywords: *Salix purpurea*, herbal raw material, salicylic glycosides

References

- Bubner, B., Köhler, A., Zaspel, I., Zander, M., Förster, N., Gloger, J.C., Ulrichs, Ch., Schneck, V. 2018. Breeding of multipurpose willows on the basis of *Salix daphnoides* Vill., *Salix purpurea* L. and *Salix viminalis* L. Appl. Agric. Forestry Res. 68, 53–66.
- Sulima, P., Przyborowski, J.P. 2019. Purple willow (*Salix purpurea* L.) and its potential uses for the treatment of arthritis and rheumatism. In: Watson, R.R., Preedy, V.R. (Ed.), Bioactive food as dietary interventions for arthritis and related inflammatory diseases. Academic Press, 535–551.
- Sulima, P., Kuszewska, A., Przyborowski, J.A. 2021. Are *Salix purpurea* L. genotypes from natural locations promising candidates for the production of high-quality herbal raw materials under controlled conditions? Ind. Crops Prod. 171, 113982.
- Wichtl, M. (Ed.). 2004. Herbal drugs and phytopharmaceuticals: a handbook for practice on a scientific basis. 3rd ed. London: CRC press.

* Presenting author. E-mail address: pawel.sulima@uwm.edu.pl (Paweł Sulima)

S1.P144 Exploring myxobacteria for bioactive compounds against malaria and neglected tropical diseases

Abdelhalim Babiker Mahmoud^{a,b,c,*}, Marcel Kaiser^d, Rolf Müller^{a,b}

^a Department of Microbial Natural Products, Helmholtz Institute for Pharmaceutical Research Saarland (HIPS), Saarbrücken, Germany

^b Department of Microbial Drugs, Helmholtz Centre for Infection Research (HZI), Braunschweig, Germany

^c Faculty of Pharmacy, University of Khartoum, Sudan

^d Swiss Tropical and Public Health Institute, Basel, Switzerland

Tropical parasitic diseases such as malaria, human African trypanosomiasis, Chagas disease, and leishmaniasis are major health hazards affecting more than a billion people worldwide, mainly in developing countries. Currently available drugs are problematic given their limited efficacy, serious adverse effects and emergence of drug resistance. Thus, there is an urgent need for new antiparasitic drugs. Natural products, particularly those deriving from microbial origins, still play a dominant role in lead discovery and development against neglected tropical diseases. In this work, we aimed to systemically identify and isolate new scaffolds of myxobacteria secondary metabolites that could serve as potential leads for antiparasitic drug development. A subset of our library of extracts and pure compounds driven from myxobacteria was screened for *in vitro* antiparasitic activity against *Trypanosoma brucei rhodesiense*, *Trypanosoma cruzi*, *Leishmania donovani*, and *Plasmodium falciparum*. Here we communicate the results of using different approaches that combine bioactivity- and structure-guided isolation as well as metabolome mining for the characterization of bioactive molecules. Also, we report the remarkable antiprotozoal activity of some myxobacterial compounds that are described for the first time.

Keywords: myxobacteria, trypanosomatids, malaria, bioactivity-guided isolation

* Presenting author. E-mail address: halim.mahmoud@helmholtz-hips.de (Abdelhalim Babiker Mahmoud)

S1.P145 Dereplication of hidden saponins in bioactive extracts

Ana M. Simonet*, Alexandra G. Durán, Francisco A. Macías

Allelopathy Group, Department of Organic Chemistry, Institute of Biomolecules (INBIO); School of Science, University of Cádiz, Puerto Real, Cádiz, Spain

Saponins have been highlighted for their biological properties, including anticancer, antibacterial, antifungal, or anti-inflammatory. They are formed by a complex two-part structure, a hydrophobic core named aglycone and a hydrophilic moiety constituted by a sugar chain linked together by a *O*- glycosidic bond (Durán et al., 2021). Nevertheless, the purification of these compounds is still a challenge due that they appear as complex mixtures with similar structurally related forms. Thereby, a dereplication strategies by combination of NMR spectra with UPLC-QTOF/MS^E and HMAI method it is proposed and it allowed the identification of the main saponins of *Agave* enriched fractions (Simonet et al., 2021). This strategy has allowed the dereplication of four fractions of *Agave* plants to led the identification of a total of 26 saponins, 14 of which have already been described and 12 of which are proposed as new saponins. Some of this saponins have structural isomers in their aglycone as well as in their sugar chain and these differences, not observed by UPLC/MS analysis, could be identified by NMR data.

Keywords: steroidal saponins, *Agave*, dereplication, HMAI

References

Durán, A.G.; Benito, J.; Macías, F.A.; Simonet, A.M. 2021. *Agave* Steroidal Saponins as Potential Bioherbicides. *Agronomy*, 11, 2404.

Simonet, A. M.; Durán, A. G.; Pérez, A. J.; Macías, F. A. 2021. Features in the NMR spectra of the aglycones of *Agave* spp. saponins. HMBC method for aglycone identification (HMAI). *Phytochem. Anal.*; 32, 38–61.

* Presenting author. E-mail address: ana.simonet@uca.es (Ana M. Simonet)

S1.P146 Understanding the chemistry of the genus *Amylostereum* in its multilateral symbioses

Esther Schulzke^a, Marc Stadler^a, Bernard Slippers^b, Yaming Liu^c, Leonie Berger^a, Christine Beemelmaans^c, Miroslav Kolarik^d, Birthe Sandargo^{a,*}

^a Microbial Drugs Department, Helmholtz Centre for Infection Research, Inhoffenstr. 7, 38124 Braunschweig, Germany

^b Department of Biochemistry Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute, University of Pretoria, South Africa

^c Anti-Infectives from Microbiota, Helmholtz Institute for Pharmaceutical Research Saarland, Campus E8.1, 66123, Saarbrücken, Germany

^d Laboratory of Fungal Genetics and Metabolism, Institute of Microbiology of the Czech Academy of Sciences, Vídeňská 1083, CZ-14220 Prague, Czech Republic

The genus *Amylostereum*, with many species tragically known as forest pathogens, especially in the southern hemisphere, live in multilateral symbioses with their corresponding, exclusive siricid woodwasps infesting coniferous trees causing white rot and subsequent pine tree diebacks (Slippers et al., 2012).




Fig 1. *Amylostereum* and its symbiotic lifestyle (created with biorender.com; based on Ryan & Hurley, in: Slippers et al. 2012).

To unravel the chemistry of this multilateral symbiosis we deployed, among other, an ecomimetic cultivation and OSMAC (one strain – many compounds) approach to activate silent gene clusters in multiple species of the genus *Amylostereum*, combined with untargeted metabolomics. The effect of isolated secondary metabolites on corresponding woodwasp parasites was investigated. Results showed a variety of decorated (nor-)sesquiterpenoids and lanostane-type triterpenoids in *A. chailletii*, *A. areolatum* and *A. laevigatum*, hinting to a closer relation to species of the Echinodontiaceae.

Keywords: *Amylostereum*, siricid woodwasp, sesquiterpenoids, white rot

References

Slippers, B., de Groot, P., Wingfield, M.J. (Eds.). 2012. The Sirex Woodwasp and its Fungal Symbiont: Research and Management of a Worldwide Invasive Pest. Springer, Dordrecht, Netherlands.

* Presenting author. E-mail address: birthe.sandargo@helmholtz-hzi.de (Birthe Sandargo)

S1.P147 *In vitro* evaluation of proliferative activity of Turkish endemic *Prangos hulusii* S.G. Şenol, H. Yıldırım & Ö. Seçmen and its major compounds

Gokay Albayrak^{a,*}, Ali Erguc^b

^a Department of Pharmaceutical Botany, Faculty of Pharmacy, Izmir Katip Celebi University, 35620 Izmir, Türkiye

^b Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Izmir Katip Celebi University, 35620 Izmir, Türkiye

Prangos sp. roots have been used as an external wound-healing agent in Anatolian traditional medicine (Kaval et al., 2014). Therefore, we obtained *n*-hexane(PH), chloroform(PC), *n*-butanol(PB), aqueous fractions(PW), and total extract(80% ethanol,PT) of the *P. hulusii* roots. We isolated and identified oxypeucedanin, isoimperatorin, and osthole as three major compounds using chromatographic(column chromatography, HPLC) and spectroscopic(¹H-NMR, LC-MS) methods.

We evaluated the potential wound-healing effects of these extract/fractions(50 and 100 µg/mL,24 and 48h.) and three major compounds(50 and 100 µM,24 and 48h.) on CCD-34Lu cells through cell proliferation assay(MTT). The activity results are shown in Table 1.

Table 1. The proliferative effects of *P. hulusii* root extracts and its major compounds

Extracts	50 µg/mL(Final%)	100 µg/mL(Final%)
<i>n</i> -hexane	218±8%*	17±4%*
Chloroform	198±5%*	141±24%*
<i>n</i> -butanol	115±7%	118±3%
Aqueous	119±2%	115±4%
Total	147±14%*	177±24%*
Compounds	50 µM(Final%)	100 µM(Final%)
Oxypeucedanin	117±6%	114±8%
Isoimperatorin	106±11%	111±6%
Osthole	126±20%	164±31%*
Control(DMSO,0.5%)	100±8%	
Doxorubicin(20 µg/mL)	76±7%*	

CCD-34Lu cell proliferation activities of tested materials after 24 h of incubation.

Values are the mean±SD from three independent experiments (n:3).

(*) significantly different(p<0.05) than the solvent control(0.5%DMSO).

n-hexane, chloroform fractions, and total extract showed higher proliferative effects than other fractions. *n*-hexane fraction demonstrated a hormetic effect, evidenced by an increase(218%) in proliferation(50 µg/mL) and a decrease(17%) in proliferation(100 µg/mL). Osthole showed the highest proliferative activity among the tested compounds. These results suggest that apolar fractions of *P. hulusii* roots and osthole have the potential to be used as wound healing agents.

Keywords: *Prangos hulusii*, Apiaceae, cell proliferation, MTT, furanocoumarins

References

Kaval, I., Behcet, L., Cakilcioglu, U., 2014. Ethnobotanical study on medicinal plants in Geçitli and its surrounding (Hakkari-Turkey). J. Ethnopharmacol., 155(1), 171-184.

* Presenting author. E-mail address: albayrakgokay@gmail.com (Gokay Albayrak)

S1.P148 Spectral data mining of museum-sourced soft corals collected in the 19th century for natural product discovery


Jennifer Netjes^{a,*}, Sam Afoullouss^a, Dale Chaput^b, Andrea Quattrini^c, Bill Baker^a

^a Department of Chemistry, University of South Florida, Tampa, FL USA

^b Proteomics and Mass Spectrometry Core Facility, University of South Florida, Tampa, FL USA

^c National Museum of Natural History, Smithsonian Institution, Washington, DC USA

Conventional marine natural product workflows rely on collection trips to the sea for the retrieval of sponges, corals, and other invertebrates. However, an alternative source of samples was newly realized: museums of natural history. Within museums are holdings of tens of thousands of invertebrates preserved in ethanol – valuable extracts for natural product chemists. In this study, museum-sourced marine invertebrate extracts were subjected to high-resolution accurate mass (HRAM) mass spectrometry, NMR, and spectral data mining techniques to profile the metabolome of the preservative extracts, providing a novel, targeted pipeline for the discovery of new natural products from an unconventional source. This newly established spectral data mining workflow is a combination of selective NMR experiments, HRAM mass spectrometry, and SIRIUS (MS² annotation tool). This method allowed for ‘turnkey chemical profiling’, by which the compound class composition of marine invertebrates is described from the crude extract (**Figure 1**), allowing for streamlined isolations of natural products through targeting both reported and unreported ion masses belonging to compound classes of interest.



Paragorgia arborea extracts, originally collected from the 1800's and supplied by the Smithsonian's National Museum of Natural History, were among the various invertebrates subjected to this described workflow. The presence of xenicane diterpenoids was detected by SIRIUS via *de novo* predictions based on fragmentation pattern and binary molecular fingerprinting, leading to the isolation and purification of unreported natural products. This technique demonstrates the utility of museum-sourced samples as well as the benefit of spectral data mining of crude extracts for targeted natural product discovery.

Keywords: marine natural products, data mining, mass spectrometry

* Presenting author. E-mail address: jenniferwilliams@usf.edu (Jennifer Netjes)

S1.P149 Influence of flavonoid 3',8'' dimerization on antioxidant, neuroprotective, anti- hyperpigmentation and anti-diabetic activities: comparative *in vitro* analysis of acacetin and isoginkgetin

Iva Jurčević Šangut*, Dunja Šamec

University North, Department of Food Technology, Trg dr. Žarka Dolinara 1, 48000 Koprivnica, Croatia

Flavonoids are the most extensively researched group of specialized plant metabolites that play a crucial role in various plant-environment interactions, with their biological activity greatly influenced by their molecular structure. They may occur as dimers, known as biflavonoids. Although over 600 biflavonoids have been described so far, their function and biosynthetic pathways in plants are not fully understood (He et al., 2021). As well it is unknown how dimerization influences the biological activity of flavonoids. We conducted a direct comparison of the biological activities of acacetin and its 3',8'' dimer, isoginkgetin (Fig. 1). We assessed their radical scavenging activity utilizing DPPH, as well as their inhibitory effects on the enzymes acetylcholinesterase, tyrosinase, α -amylase, and α -glucosidase.



Fig. 1. Chemical structure of a) isoginkgetin and b) acacetin.

Both compounds exhibited weak antioxidant activity when measured by DPPH, up to a concentration of 1 mg/mL. Notably, isoginkgetin demonstrated significantly stronger inhibition of acetylcholinesterase compared to acacetin, registering at $41.99 \pm 3.68\%$ and $19.86 \pm 0.78\%$, respectively, at a concentration of 100 μM . Additionally, the dimer isoginkgetin displayed nearly double the inhibition potency against α -amylase and α -glucosidase compared to acacetin, suggesting potential antidiabetic properties. However, acacetin exhibited increased tyrosinase inhibition activity compared to the dimer isoginkgetin. Our findings suggest that 3',8''-dimerization could potentially enhance the neuroprotective properties and antidiabetic activities of flavonoids. Conversely, monomers may be better suited for addressing hyperpigmentation disorders.

Keywords: acacetin, 3',8''-biflavones, isoginkgetin

References

He, X., Yang, F., Huang, X., 2021. Proceedings of Chemistry, Pharmacology, Pharmacokinetics and Synthesis of Biflavonoids. *Molecules* 26, 6088.

* E-mail address: ijurcevic@unin.hr (Iva Jurčević Šangut)

S1.P150 Development of quality criteria for *Vitex pinnata* Leaf as an herbal medicine

Zohorul Islam^{a,*}, Maryam Malmir^a, Manuela Caniça^b, Nurul Islam^c, Olga Silva^a

^a Research Institute for medicines (iMed.Ulisboa), Faculty of Pharmacy, Universidade de Lisboa, 1649-003 Lisbon, Portugal

^b National Reference Laboratory of Antibiotic Resistance and Healthcare-Associated Infections, Department of Infectious Diseases, National Institute of Health Dr. Ricardo Jorge, 1649-016 Lisbon, Portugal

^c Department of Zoology, Faculty of Biological Sciences, University of Rajshahi, Rajshahi 6250, Bangladesh

Vitex pinnata L. a shrub tree from the Lamiaceae family, is widely used in traditional medicine in Asian countries and Western Europe (Kamal *et al.*, 2022), with its leaf used to treat hypertension, fever, and gastrointestinal diseases (Islam *et al.*, 2024). The present study aims to establish the standard parameters for proper identification and quality control of *V. pinnata* dried leaf through botanical (macroscopic and microscopic) identification and phytochemical (TLC, colorimetry, and UV-visible spectroscopy) assessments of the leading chemical classes of secondary metabolites. Macroscopically, dried leaves showed pale grey to yellowish color, 3/5 foliate with acuminate apex and ovate/ elliptic shape, leaflet with 6-16 cm long and 2-8 cm wide, petiolule 2-3 mm long and petiole 3-10 cm long. Microscopically, the leaf exhibited the presence of capitate glandular trichomes, non-glandular trichomes, spheroidal trichomes, and thick and prominent midribs. Stomata were found to be of anomocytic type and densely distributed. Chemical characterization revealed the presence of phenolic compounds (phenolic acids, flavonoids, condensed and hydrolysable tannins) and terpenoids (iridoids) among the main classes of secondary metabolites of the leaf 70% hydroethanolic extract. Total phenol and flavonoid contents were calculated as 206.5±14.2 mg gallic acid equivalent/g dried extract and 92.2±5.2 mg catechin equivalent/g dried extract, respectively, and luteolin derivatives were detected among the main marker compounds. The data obtained will contribute to establishing botanical and chemical quality criteria for identifying and authenticating *V. pinnata* leaf as a possible herbal substance.

Keywords: *Vitex pinnata*, leaf, herbal medicine, botanical identification, chemical characterization

References

Islam, Z., Caldeira, G.I., Caniça, M., Islam, N., Silva, O., 2024. *Vitex Genus* as a Source of Antimicrobial Agents. *Plants* 13, 401.

Kamal, N., Mio Asni, N.S., Rozlan, I.N.A., Mohd Azmi, M.A.H., Mazlan, N.W., Mediani, A., Baharum, S.N., Latip, J., Assaw, S., Edrada-Ebel, R.A., 2022. Traditional Medicinal Uses, Phytochemistry, Biological Properties, and Health Applications of *Vitex* sp. *Plants (Basel)* 11, 1944.

* Presenting author. E-mail address: zislam@edu.ulisboa.pt (Zohorul Islam)

S1.P151 Monographic characterization of *Vitex peduncularis* Leaf as an Herbal Medicine

Zohorul Islam^{a,*}, Maryam Malmir^a, Manuela Caniça^b, Nurul Islam^c, Olga Silva^a

^a Research Institute for medicines (iMed.Ulisboa), Faculty of Pharmacy, Universidade de Lisboa, 1649-003 Lisbon, Portugal

^b National Reference Laboratory of Antibiotic Resistance and Healthcare-Associated Infections, Department of Infectious Diseases, National Institute of Health Dr. Ricardo Jorge, 1649-016 Lisbon, Portugal

^c Department of Zoology, Faculty of Biological Sciences, University of Rajshahi, Rajshahi 6250, Bangladesh

Vitex peduncularis Wall. ex Schauer, a deciduous shrub, typically reaches heights of 6-12 meters and is noted for its slender branches and aromatic foliage. This species has been recognized for its medicinal properties, particularly in treating joint pain, diabetes, fever, and skin disorders, as emphasized in Islam et al.'s 2024 study. The current research focuses on establishing monographic parameters for accurately identifying and quantifying the dried leaf of *V. peduncularis*. This involves botanical (macroscopic and microscopic) identification and phytochemical evaluation (including Thin Layer Chromatography, Colorimetry, and UV-visible spectroscopy) of the principal chemical classes of secondary metabolites. The dried leaf is macroscopically characterized by its green to yellowish color, being 3-foliate with an acute apex, and a lanceolate/elliptic shape. Each leaflet measures 5-20 cm in length and 2-5 cm in width, with a petiolule 1-2 cm long and a petiole extending 3-11 cm. Microscopically, the leaf demonstrates the presence of capitate glandular trichomes and non-glandular trichomes. The stomata are of the anomocytic type and are densely distributed. Chemical profiling of the leaf's 70% hydroethanolic extract revealed the presence of flavonoids, condensed tannins, and terpenoids (notably iridoids) as the main classes of secondary metabolites. The total phenolic and flavonoid contents were quantified as 90.9 ± 4.7 mg of gallic acid equivalent per gram of dried extract and 42.9 ± 1.4 mg of catechin equivalent per gram of dried extract. Additionally, apigenin derivatives were identified among the main marker compounds. This study contributes valuable data towards establishing botanical and chemical quality standards for the authentication of *V. peduncularis* leaf as herbal medicine.

Keywords: *Vitex peduncularis*, leaf, herbal medicine, botanical identification, chemical characterization

Reference:

Islam, Z., Caldeira, G.I., Caniça, M., Islam, N., Silva, O., 2024. *Vitex* Genus as a Source of Antimicrobial Agents. *Plants* 13, 401.

Kamal, N., Mio Asni, N.S., Rozlan, I.N.A., Mohd Azmi, M.A.H., Mazlan, N.W., Mediani, A., Baharum, S.N., Latip, J., Assaw, S., Edrada-Ebel, R.A., 2022. Traditional Medicinal Uses, Phytochemistry, Biological Properties, and Health Applications of *Vitex* sp. *Plants (Basel)* 11, 1944.

* Presenting author. E-mail address: zislam@edu.ulisboa.pt (Zohorul Islam)

S1.P152 Identification of ergot alkaloids and lysergic acid derivatives in Kykeon, a traditional concoction of the Eleusinian Mysteries

Romanos K. Antonopoulos^a, Evangelos Dadiotis^b, Antigoni Cheilari^{b,*}, Vangelis Mitsis^c, Eleni Melliou^b, Prokopis Magiatis^b, Nikolas Fokialakis^b, Zacharoula Gonou-Zagou^a, Nektarios Aligiannis^b

^a Department of Ecology and Systematics, Faculty of Biology, National and Kapodistrian University of Athens, Panepistimiopolis Zografou 17551, Greece

^b Department of Pharmacognosy and Natural Products Chemistry, Faculty of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou 17551, Greece

^c Ekati Alchemy Lab SL, Carretera Barcelona 11, 08180 Moia, Spain

To explore traditional preparations of ancient Greece, we revisited the theory regarding the ceremonial use of kykeon, a sacred concoction used during the Eleusinian Mysteries (Wasson et. al, 1978). Beside the hallucinogenic properties attributed to the drink, kykeon is mentioned by Homer to be offered as a tonic to soldiers in battle, as well as for wounds' treatment (Homer, Iliad, 635-641). Kykeon was a mixture consisting mainly of barley flour, water, and herbs. *Claviceps purpurea* is a fungus known for infecting cereals while producing sclerotia containing ergot alkaloids and probably is the psychoactive agent responsible for kykeon's properties. Our aim was to support the hypothesis that ancient Greeks had discovered a way of hydrolysing ergot alkaloids into lysergic acid amide and lysergic acid. Following a traditional formulation, a concoction utilizing lye was prepared and monitored by NMR. *C. purpurea* sclerotia were powdered and boiled in a lye solution. After 30 min, ¹H-NMR spectra revealed that the peak at 9.6 ppm of the amide linkage of ergocryptine connecting the lysergic acid moiety and the peptide disappeared, and the occurrence of the characteristic peaks at 8.03 and 6.94 ppm indicating the presence of ergine and lysergic acid (Fig. 1).




Fig. 1. Conversion of ergocryptine in ergine and lysergic acid with lye solution treatment

UPLC/Q-TOF-MS analysis in positive ionization mode confirmed the presence of ergine (m/z 268.14), lysergic acid (m/z 269.12) and ergocryptine (m/z 576.31). The transformation of ergot alkaloids was confirmed into the traditional concoction prepared with lye and support the hypothesis that the hallucinogenic properties of kykeon can be attributed to the use of ergot.

Keywords: Eleusinian Mysteries, kykeon, *Claviceps purpurea*, ergocryptine, lysergic acid

References

Wasson, R.G., Hofmann, A., Ruck, C.A.P. 1978. The Road to Eleusis. Unveiling the Secret of the Mysteries. Harcourt Brace Jovanovich, New York.

Homer's Iliad, Rhapsody XI, verses 635–641.

* Presenting author. E-mail address: cheilarianti@pharm.uoa.gr (Antigoni Cheilari)

S1.P153 Molecular networking-based dereplication and structural elucidation of compounds derived from liquid culture supernatants of marine-sourced *Streptomyces* strains

Nikola Milic^a, Christina Koufali^{a,*}, Christina Stamataki^a, Nikolaos Tsafantakis^a, Fernando de la Calle Verdu^b, Nikolas Fokialakis^a

^a Department of Pharmacognosy and Chemistry of Natural Products, Faculty of Pharmacy, National and Kapodistrian University of Athens, Athens 15771, Greece

^b PharmaMar SA, Avda. De los Reyes, 1, 28770 - Colmenar Viejo, Madrid, Spain

Marine environments are a treasure trove of microbial diversity teeming with actinomycetes, renowned as producers of chemically diverse bioactive metabolites. Marine actinobacterial compounds—including amphiphilic biosurfactants and siderophores, studied within the EU's Horizon2020 SECRETed[‡] project—offer a promising avenue for potentially novel therapeutic and/or industrial applications. In this work, a liquid-liquid extraction of culture supernatants of ten marine- derived *Streptomyces* strains was conducted with ethyl acetate, followed by UPLC-ESIHRMS analyses of the yielded extracts. The acquired spectral data were subjected to a molecular networking- based dereplication workflow to: (a) cluster compounds based on their spectral and structural similarities, (b) generate annotations through matching against curated spectral libraries, and (c) perform a chemical classification of the annotated natural products. A multi-step data processing, parametrisation, and visualisation in several chemoinformatic/computational environments (MZMine3 (Schmid et al., 2023), GNPS (Wang et al., 2016), R (Posit team, 2023), and Cytoscape (Shannon et al., 2003)) were involved in this process, including manual workflow validations. The extracts underwent Sephadex® LH-20-based fractionation and the fractions were further purified and analysed via UPLC-ESIHRMS and NMR to structurally elucidate the compounds of interest, pinpointed as potential targets through the dereplication. A diverse range of N-containing compounds were identified, including carboxylic and amino acid derivatives, cyclic and linear peptides, various alkaloids, alongside compounds with both hydrophobic and hydrophilic moieties, suggesting an amphiphilic nature thereof and, hence, their potential function as biosurfactants and/or siderophores.

Acknowledgments: European Union's Horizon2020 Research and Innovation Programme (Grant Agreement No. 101000794).

Keywords: marine *Streptomyces*, UPLC-HRMS, molecular networking, NMR structural elucidation

References

- Posit team (2023). RStudio. Posit Software, PBC, Boston, MA. URL <http://www.posit.co/>.
- Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., Ideker, T. 2003. Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res.* 13, 2498–2504.
- Schmid, R., Heuckeroth, S., Korf, A., Smirnov, A., Myers, O., Dylund, T.S., Bushuiev, R., Murray, K.J., Hoffmann, N., Lu, M., Sarvepalli, A., Pluskal, T., 2023. Integrative analysis of multimodal mass spectrometry data in MZmine 3. *Nat. Biotechnol.* 41, 447–449.
- Wang, M., Carver, J.J., Phelan, V.V., Sanchez, L.M., Garg, N., Peng, Y., Nguyen, D.D., Watrous, J., Kapono, C. A., Luzzatto-Knaan, T., Bandeira, N., 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat. Biotechnol.* 34, 828–837.

* Presenting author. E-mail address: chrikouf@pharm.uoa.gr (Christina Koufali)

S1.P154 Src tyrosine kinase inhibitory activity of selected flavonoids

Sanda Vladimir-Knežević^{a,*}, Dubravko Jelić^b, Maja Bival Štefan^a, Biljana Blažeković^a

^a University of Zagreb, Faculty of Pharmacy and Biochemistry, Department of Pharmacognosy, Trg Marka Marulića 20, 10000 Zagreb, Croatia

^b Selvita Ltd., Prilaz Baruna Filipovića 29, 10000 Zagreb, Croatia

The aim of the present study was to investigate various flavonoids for their inhibitory effect on Src tyrosine kinase, as this enzyme plays an important role in multiple cellular processes and is over-activated in both cancer and inflammatory cells (Martellucci et al., 2020; Brian and Freedman, 2021). Inhibiting the activity of Src tyrosine kinase is an important strategy to prevent chronic inflammation and thus the development of cancer and other inflammation-related diseases. Twenty flavonoids in the form of aglycones were analysed using a time-resolved fluorescence resonance energy transfer (TR-FRET) assay, with staurosporine serving as a positive control (Vladimir-Knežević et al., 2023). The tested flavonoids belonged to the structurally diverse types of flavonoids, including flavones, flavonols, flavanones, isoflavones, flavan-3-ols and chalcones. Among them, luteolin was found to be the strongest Src tyrosine kinase inhibitor ($IC_{50} = 9$ mM), followed by baicalein, myricetin, fisetin and diosmetin ($IC_{50} = 27-38$ mM). The IC_{50} values obtained for the other flavonoids were in the range of 104-708 mM or the inhibitory effect was not detected at a concentration of 2000 mM. In this study, the relationship between the structure and activity of the tested flavonoids was discussed and the most effective candidates were selected for further investigation.

Keywords: Src tyrosine kinase inhibitory activity, flavonoids, luteolin

References

- Martellucci, S., Clementi, L., Sabetta, S., Mattei, V., Botta, L., Angelucci, A. 2020. Src family kinases as therapeutic targets in advanced solid tumors: What we have learned so far. *Cancers*. 12, 1448.
- Brian, B.F., Freedman, T.S. 2021. The Src-family kinase Lyn in immunoreceptor signaling. *Endocrinology*. 162, 152.
- Vladimir-Knežević, S., Bival Štefan, M.; Blažeković, B., Jelić, D., Petković, T., Mandić, M., Šprajc, E., Lovković, S. 2023. Src tyrosine kinase inhibitory and antioxidant activity of black chokeberry and bilberry fruit extracts rich in chlorogenic acid. *Int. J. Mol. Sci.* 24, 15512.

Presenting author. E-mail address: sanda.vladimir@pharma.unizg.hr (Sanda Vladimir-Knežević)

S1.P155 Green extraction of bioactive volatile terpenes from Croatian *Salvia* species

Mateja Mervić^a, Marija Kindl^a, Maja Bival Štefan^a, Biljana Blažeković^a, Matija Cvetnić^b, Sanda Vladimir-Knežević^{a,*}

^a University of Zagreb, Faculty of Pharmacy and Biochemistry, Department of Pharmacognosy, Trg Marka Marulića 20, 10000 Zagreb, Croatia

^b University of Zagreb, Faculty of Chemical Engineering and Technology, Department of Analytical Chemistry, Trg Marka Marulića 20, 10000 Zagreb, Croatia

Salvia L. is the largest genus within the Lamiaceae family and includes many medicinal plants used in traditional and modern medicine. In addition to *S. officinalis* L., the most important representatives of this genus growing wild in Croatia are *S. fruticosa* Mill., *S. glutinosa* L., *S. sclarea* L. and *S. verticillata* L., which have not yet been sufficiently studied (Mervić et al., 2022). The aim of this study was to isolate volatile terpenes from selected *Salvia* species using environmentally friendly techniques such as hydrodistillation and supercritical carbon dioxide (sCO₂) extraction and to determine their composition and biological activity. The leaves of *S. officinalis* and *S. fruticosa* proved to be very rich sources of essential oil, containing 22 mL/kg and 28 mL/kg, respectively, while 0.3-0.8 mL/kg of essential oil was isolated from the other *Salvia* species by hydrodistillation. A total of 143 compounds were identified by GC/MS analysis, accounting for 87.34-99.62% of the oil content. The main constituents of the tested oils were α -thujone, 1,8-cineole, camphor, γ -gurjunene, caryophyllene oxide and germacrene D. The essential oils of *S. fruticosa* and *S. officinalis* showed antioxidant and acetylcholinesterase inhibitory activity, with α/β -thujone and 1,8-cineole being most responsible for the effects. To obtain the highest yield of the extract with the highest content of the most bioactive volatile terpenes, the sCO₂ extraction process was optimised. The extraction yield varied between 0.60% and 4.51%, depending on the applied pressure (100-220 bar) and temperature (40-60°C). A pressure of 220 bar and a temperature of 40°C were optimal parameters for the extraction of α/β -thujone and 1,8-cineole.

Keywords: *Salvia* species, essential oils, sCO₂ extracts, neuroprotective effect

References

Mervić, M., Bival Štefan, M., Kindl, M., Blažeković, B., Marijan, M., Vladimir-Knežević, S. 2022. Comparative antioxidant, anti-acetylcholinesterase and anti- α -glucosidase activities of Mediterranean *Salvia* species. *Plants*. 11, 625.

* Presenting author. E-mail address: sanda.vladimir@pharma.unizg.hr (Sanda Vladimir-Knežević)

S1.P156 *Anti-Fusarium* activities of microemulsions of essential oils of *Artemisia annua* L. (Asteraceae) and *Salvia fruticosa* L. (Lamiaceae) to be used as pre-harvest treatment of fruits

Lucia Grifoni^{a,*}, Cristiana Sacco^b, Rosa Donato^b, Tziakas Spyros^c, Tomou Ekaterina-Michaela^c, Skaltsa Helen^c, Giulia Vanti^a, Anna Rita Bilia^a

^a Department of Chemistry Ugo Schiff, via Ugo Schiff 6, 50019 Sesto Fiorentino, Florence, Italy

^b Department of Health Sciences, University of Florence, Florence, Italy Viale G.B. Morgagni, 48 – 50134 Florence, Italy

^c Department of Pharmacognosy and Chemistry of Natural Products, Faculty of Pharmacy, National & Kapodistrian University of Athens, Panepistimiopolis, Zografou, 15771, Athens, Greece

The study covers the analysis, formulation and biological evaluation of two essential oils (EOs), *Artemisia annua* L. (Asteraceae) and *Salvia fruticosa* L. (Lamiaceae). Both EOs have been reported for their bactericidal and fungicidal activities (Santomauro et al, 2016; id., 2018; Jaradat et al., 2022). Their activity could be optimized if nanoencapsulated in drug delivery systems to decrease their volatility, improving the stability, water solubility, and efficacy. Composition of essential oils was obtained by GC-MS analysis. The main constituents characterized the *Salvia* EO were the oxygenated monoterpenes (65.1%) with 1,8-cineole (38.5%) and camphor (7.8%). Main constituents of *Artemisia* EO were monoterpenes (ca. 88%) with camphor (25.2%), 1,8-cineole (20%), and artemisia ketone (12.5%). The microemulsions were developed using vitamin E acetate as oil (8.22%), Cremophor RH 40 and Labrasol ALF as surfactants (14.8%) in a 1:1 ratio. The formulation was selected using a pseudo-ternary diagram. The microemulsion was loaded with *S. fruticosa* and *A. annua* EOs at a concentration of 10 mg/mL with 100% recovery using HPLC. Antifungal activity was carried out using a microplate dilution method using amphotericin B[®] as a positive control, while sterile Sabouraud broth was the negative control. Twenty microliters of *Fusarium verticillioides* subcultured in Sabouraud agar to 1.0×10^5 spores/mL were added to the wells. The plate was sealed and incubated for 24 hours at 28°C. *Fusarium* ufc per mL resulted 2.4×10^7 . Microemulsions containing 1.8, 1.6, 1.4, 1.2 and 1.0 mg of EOs were added and incubated in the for 5 days. The results are the average of two tests repeated in triplicate. MIC values were determined when a logarithms reduction of 4 was found. The two microemulsions displayed a similar activity towards *Fusarium* spp., being the MIC of *S. fruticosa* 1.2 mg and that of *A. annua* 1.4 mg. The unloaded microemulsions had no activity. These two formulations could be very useful for the in-field application of the tested EOs in order to treat *Fusarium* infections.

Keywords: essential oils, microemulsions, antifungal activity, *Fusarium verticillioides*

Acknowledgments: This research was funded by the project EthnoHERBS (H2020-MSCA-RISE-2018, Grant Agreement No. 823973)

References

- Santomauro, F, Donato, R, Pini, G, Sacco, C, Ascrizzi, R, Bilia, AR., 2018. Liquid and Vapor-Phase Activity of *Artemisia annua* Essential Oil against Pathogenic *Malassezia* spp. *Planta Med.* 84(3):160-167
- Santomauro F, Donato R, Sacco C, Pini G, Flamini G, Bilia AR., 2016 Vapour and Liquid-Phase *Artemisia annua* Essential Oil Activities against Several Clinical Strains of *Candida*. *Planta Med.* 82(11-12):1016-20.
- Jaradat N, Abdallah S, Al-Maharik N, Altamimi M, Hawash M, Qneibi M, Abu Khair A, Zetawi A, Jabarin L., 2022. Constituents, Antibacterial Adhesion, Cytotoxic and in Vitro Metastasis Blocking Properties of *Salvia fruticosa* Essential Oils from Three Palestinian Localities. *Chem Biodivers.* 19(4)

* Presenting author. E-mail address: lucia.grifoni@unifi.it (Lucia Grifoni)

S1.P157 Antibacterial and cytotoxic activities of bark and branches extracts from *Lafoensia replicate* Pohl (Lythraceae)

Débora Machado de Lima^a, Ana Carolina Costa Santos^b, Carlos Ueira-Vieira^b, Matheus Reis Santos de Melo^c, Denise Crispim Tavares^c, Raquel Maria Ferreira Sousa^{a,*}

^a Institute of Chemistry, Federal University of Uberlândia, 38400-902, Uberlândia, MG Brazil

^b Institute of Biotechnology, Federal University of Uberlândia, 38405-319, Uberlândia, MG Brazil

^c University of Franca, 14404-600, Franca, São Paulo, Brazil

Lafoensia replicate Pohl (Lythraceae) is an endemic tree from Cerrado vegetation in central Brazil. The bark of this species is used in the treatment of liver and kidney diseases, general inflammations, wound healing, gastritis, high blood pressure, headache, and stomach pain (Sobrinho et al, 2005). In this study, the antibacterial (*Staphylococcus aureus* and *Escherichia coli*) and cytotoxicity (human non-tumor keratinocyte cell line) activities of bark and branches extracts from *L. replicata* were evaluated and the chemical composition was analyzed by chromatography coupled to mass spectrometry with an electrospray ionization system (LC/ESI-MS/MS-negative mode). The extraction was carried out by maceration at room temperature with hexane (HE), followed by ethanol extraction (EE). The activities were performed by the broth microdilution technique (Santos, 2022; Riss et al. 2013). The extract shows MIC against *S. aureus* of 6.25, 0.78, 0.39 and 12.5 µg/mL for EE-bark, EE-branches, HE-bark and HE-branches, respectively. The MIC against *E. coli* was 0.39, 6.25, >800 and 50 µg/mL, for EE-bark, EE-branches, HE-bark and HE-branches, respectively. In general, EE-bark and EE-branches showed better results against the bacteria evaluated. The cytotoxicity of these extracts was evaluated and IC₅₀ of the EE-bark and EE-branches were 407.90 and 376.33 µg/mL, respectively. The composition annotated by LC/ESI-MS/MS for these extracts showed variated hexahydroxydiphenols (HHDP), (*epi*)gallo catechin, variated punicalins, gallic acid, punicalagin, flavogallonic acid, terflavin A, isorhamnetin hexoside, quercetin hexoside, ellagic acid and hexoside kaempferol. This study showed that *L. replicate* is a natural source of compounds with antibacterial activity and low cytotoxicity.

Acknowledgments: CNPq, FAPEMIG and CAPES.

Keywords: *Lafoensia replicate*, cytotoxicity, antibacterial, tannin, HHDP

References

- Riss T.L., Moravec R.A., Niles A.L., et al. 2013. Cell Viability Assays. In: Assay Guidance Manual; Markossian S., Grossman A., Brimacombe K., et al., Eds.; Eli Lilly & Company and the National Center for Advancing Translational Sciences: Bethesda, MD.
- Santos, A.C.C., Malta, S.M., Dantas, R.C.C., Rocha, N.D.C., Azevedo, V.A.C., Ueira-Vieira, C., 2022. Antimicrobial activity of supernatants produced by bacteria isolated from Brazilian stingless bee's larval food. BMC Microbiol., 22, 127.
- Sobrinho, F.C.B., Almeida, A.L.S., Monteiro, J.M., 2016. Ethno-pharmacological study about *Lafoensia replicate* Pohl. in east Maranhão, Brazil: a promising species for bioprospection. Desenvol. Meio Ambiente, 39, 207-216.

* Presenting author. E-mail address: rsousa@ufu.br (Raquel Maria Ferreira Sousa)

S1.P158 Phytochemical composition and biological activities of *Pleroma granulosum* (Melastomataceae) leaf extract

Thaywane Azevedo Marquesa^a, Débora Machado de Lima^a, Anna Livia Oliveira Santos^b, Carlos Henrique Gomes Martins^b, Carla de Moura^c, Alberto de Oliveira^a, Raquel Maria Ferreira Sousa^{a,*}

^a Institute of Chemistry, Federal University of Uberlândia, 38400-902, Uberlândia, Minas Gerais, Brazil

^b Institute of Biomedical Sciences, Federal University of Uberlândia, 38405-319, Uberlândia, Minas Gerais, Brazil

^c Federal Institute Goiano, 75650-000, Morrinhos, Goiás, Brazil

Pleroma granulosum is a widely distributed tree and widely used in square decoration due to its exuberant flowering. This species has been little explored scientifically. Therefore, this study investigated the antioxidant and antifungal activities and the chemical composition of *P. granulosum* leaf extracts. Extracts were obtained by maceration in hexane (HE) followed by ethanol (EE). Antifungal activity was evaluated by the broth microdilution method (CLSI, 2012). The antioxidant activity was analyzed by DPPH, ABTS and FRAP methods (Quaresma et al., 2020, Malta, Liu, 20014). The composition was carried out by liquid chromatography coupled to electrospray ionization source mass spectrometry (LC/ESI-MS/MS). The antioxidant activity by the DPPH method showed EC₅₀ of 6.07±0.28 and 206.08±25.50 µg mL⁻¹ for EE and HE, respectively. The ABTS method presented 3068.00±123.60 and 24.10±6.50 µmol ET (Equivalente Trolox) g⁻¹ for EE and HE, respectively. The FRAP method showed 736.09±7.97 and 180.41±2.21 µmol ET g⁻¹ for EE and HE, respectively. In all methods, greater activity was observed for EE. The antifungal activity showed MIC of 2.93, 1.46, 11.72 and 1.46 µg mL⁻¹ for the EE against *C. albicans*, *C. glabrata*, *C. tropicalis* and *C. parapsilosis*, respectively. HE presented MIC>3000 µg mL⁻¹ against all fungi. Therefore, EE presented better activities compared to HE. Therefore, the composition of EE showed several phenolic compounds, including flavonoids, phenolic acids and tannins. The results demonstrate that the extract of *P. granulosum* is a source rich in phenolic compounds with high antioxidant and antifungal activity. Acknowledgments: CNPq, FAPEMIG and CAPES.

Keywords: *Pleroma granulosum*, antioxidant, antifungal, tannin, phenolic

References

- CLSI Clinical and Laboratory Standards Institute. 2012. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically. Approved standard M7-A9. Wayne, PA: CLSI.
- Malta, L.G., Liu, R.H., 2014. Analyses of Total Phenolics, Total Flavonoids, and Total Antioxidant Activities in Foods and Dietary Supplements, 2014. In: Van Alfen, N.K. (Ed.), Encyclopedia of Agriculture and Food Systems. Academic Press, Oxford, pp. 305-314.
- Quaresma, D.M.O., Justino, A.B., Sousa, R.M.F., Munoz, R.A.A., Aquino, F.J.T., Martins, M.M., Goulart, L.R., Pivatto, M., Espindola, F.S., De Oliveira, A. Antioxidant compounds from *Banisteriopsis argyrophylla* leaves as α-amylase, α-glucosidase, lipase, and glycation inhibitors. 2020. Bioorg. Chem. 105, 104335, 2020.

* Presenting author. E-mail address: rsousa@ufu.br (Raquel Maria Ferreira Sousa)

S1.P159 Combining a ‘multi-bioassay screening’ approach with metabolomics for in depth exploration of four Andean-Patagonian fungi

Bastien Petit^{a,b,*}, Ines Michrafy^{a,b}, Jonhatan Gómez Espinoza^c, Jaime Roberto Cabrera-Pardo^c, Sandra Loesgen^{a,b}

^a Whitney Laboratory for Marine Bioscience, St. Augustine, FL, USA

^b Department of Chemistry, University of Florida, Gainesville, FL, USA

^c Departamento de Química, Universidad del Bío-Bío, Concepción, Chile

Finding compounds that are both bioactive and structurally novel is arguably the most desired and challenging goal in natural products research. Bioguided fractionation is considered the gold standard for isolating bioactive compounds, despite some drawbacks. These drawbacks include time- consuming and costly lab work as well as the substantial production of noxious chemical waste, among others. Additionally, bioactivity often takes precedence over structural novelty in bioguided fractionation, resulting in the isolation of bioactive but already-known compounds. Recent advancements in mass spectrometry-based bioinformatics have facilitated the organization and visualization of large datasets, such as specialized metabolites from natural extracts. These tools, relying on spectroscopic data, are particularly suitable for profiling and prioritizing compounds to identify structural novelty. Furthermore, metabolomics platforms are generally flexible enough to incorporate various layers of additional information into their processes. In theory, this flexibility enables the identification of compounds with desired characteristics based on analytical, taxonomical, experimental conditions, or bioactivity data. Based on these principles, we present a multilateral approach that combines standard bioactivity screening with metabolomics guidance. This methodology is applied for the first time to four Basidiomycetes fungi from the Andean-Patagonian region of Chile: *Stereum hirsutum*, *Stereum greslebinii*, *Aleurodiscus vitellinus*, and *Anthracoephyllum discolor*. These strains were grown and extracted under various conditions to promote chemical diversity through experimental variations. The resulting panel of extracts underwent a ‘multi-bioassay screening’ against *Staphylococcus aureus*, *Escherichia coli*, *Candida albicans*, HCT-116 cells, and zebrafish. Additionally, a feature-based molecular network integrating taxonomy, experimental variations, and bioactivity data was generated to identify compounds with unique characteristics within the dataset.

Keywords: fungi, Basidiomycetes, bioactivity, metabolomics, antimicrobials

* Presenting author. E-mail address: petitbastien@ufl.edu (Bastien Petit)

S1.P160 Diketopiperazines from the marine sediment-derived bacterium *Streptomyces hoynatensis*

Ozlem Demirkiran^{a,*}, Elif Coskun^a, Fatma Kaynak Onurdag^b, Suzan Okten^b

^a Department of Pharmacognosy, Faculty of Pharmacy, Trakya University, Edirne, Turkey

^b Department of Pharmaceutical Biology, Faculty of Pharmacy, Trakya University, Edirne, Turkey

Cyclic dipeptides also known as diketopiperazines (2,5-DKPs), or dipeptide anhydrides, are among the most common peptide derivatives found in nature (Prasad 1995). They have been obtained from a variety of natural resources, including marine organisms (Huang et al. 2010). The interest in DKPs is high because they exhibit not only antimicrobial, antitumor, and antiviral activities (Huang et al. 2014) but also activities such as a relatively rare treatment of ischemic brain injury (Guan 2008), anti- Alzheimer's (Turkez et al. 2020) and inhibition of microtubule polymerization (Ding et al. 2020). *Streptomyces hoynatensis* isolated from Black Sea sediments, produced a new diketopiperazine, Hoynatenamide (**1**) along with six known diketopiperazines (**2-7**) (Fig. 1). The structures of the compounds were determined based on 1D- and 2D-NMR spectroscopic analysis, and mass spectrometry data. The absolute configuration of compound **1** was deduced by traditional hydrolysis, derivative formation, and chromatographic analyses compared with standards. Among the compounds tested, compound **2** showed the best activity against HeLa cancer cell lines and **3** against MCF-7 cell lines with an IC₅₀ value of 19.4 and 32.4 μ M, respectively. Except for compound **1**, all of the compounds showed mild antifungal activity against *C.albicans* with an IC₅₀ value of 64 μ g/mL.




Fig. 1. Structure of the isolated diketopiperazines.

Keywords: diketopiperazines, *Streptomyces*, microbial secondary metabolites, antimicrobial activity, antiproliferative activity

References

- Ding Z, Feifei L, Changjiang Z, Feng L, Yuqian L, Shixiao W, Jianchun Z, Wenbao L. 2020. Structure-based design and synthesis of novel furan-diketopiperazine-type derivatives as potent microtubule inhibitors for treating cancer. *Bioorg Med Chem*. 28, 115435.
- Guan J. 2008. Insulin-like growth factor-1 and its derivatives: potential pharmaceutical application for ischemic brain injury. *Recent Pat CNS Drug Discov*. 3, 112-27.
- Huang, RM, Xiang-Xi Y, Yuying Z, Xiangdong S, Yan P, Cheng-Hai G. 2014. An Update on 2,5 Diketopiperazines from Marine Organisms. *Marine Drugs*, 12: 6213-35.
- Huang, RM, Xuefeng Z, Tunhai X, Xianwen Y, Yonghong L. 2010. Diketopiperazines from Marine Organisms. *Chem Biodivers*. 7: 2809-29.
- Prasad C. 1995. Bioactive Cyclic Dipeptides. *Peptides*. 16, 151-64.
- Turkez H, Cacciatore I, Arslan ME, Fornasari E, Marinelli L, Di Stefano A, Mardinoglu A. 2020. Histidyl-Proline Diketopiperazine Isomers as Multipotent Anti-Alzheimer Drug Candidates, *Biomolecules*, 10, 737.

* Presenting author. E-mail address: ozlemdemirkiran@trakya.edu.tr (Ozlem Demirkiran)

S1.P161 The phenolic acids production in stationary and agitated shoot cultures of *Aronia melanocarpa*, *A. arbutifolia* and *A. × prunifolia*

Paweł Kubica, Halina Ekiert, Agnieszka Szopa*

Department of Pharmaceutical Botany, Medical College, Jagiellonian University, Medyczna 9 str, 30-688 Kraków, Poland

Aronia melanocarpa (Michx.) Elliott (black chokeberry), *A. arbutifolia* (L.) Pers. (red aronia) and *A. × prunifolia* (Marsh.) Rehd. (purple aronia) are known crop and medicinal plant species of North American origin (Kokotkiewicz et al. 2010). Different plant *in vitro* culture systems (e.g., stationary and agitated cultures) have a great influence on the accumulation of bioactive metabolites in the cultivated biomasses. The aim of this study was to compare the productivity of *Aronia* sp. shoots cultivated in stationary and agitated cultures. The object of interest were phenolic acids - the metabolites with anticancer, anti-inflammatory and other activities important in phytotherapy and cosmetology. The cultures were cultivated on three variants of Murashige-Skoog medium (MS) supplemented with cytokinin, 6-benzyladenine (BA) and auxin, 1-naphthaleneacetic acid (NAA) in the following concentrations [mg/L]: variant I - 0.5 and 2.0; variant II - 2.0 and 2.0; variant III - 3.0 and 1.0. In the methanolic extracts from biomasses, 19 compounds (derivatives of cinnamic acid, benzoic acid and depsides – chlorogenic, neochlorogenic and rosmarinic acids) were estimated by an LC-DAD method (Ellnain and Wojtaszek, 1999). The studied *in vitro* cultures produced as the main metabolites three depsides: neochlorogenic acid, chlorogenic acid and rosmarinic acid. The maximal amounts of these compounds were higher in shoots cultivated as agitated cultures in comparison with stationary cultures and reached respectively 82.00, 105.45 and 134.24 mg/100g DW (*A. melanocarpa*), 25.48, 175.94 and 132.49 mg/100g DW (*A. arbutifolia*), and 70.91, 260.34 and 225.26 mg/100g DW (*A. × prunifolia*). The studied *Aronia in vitro* cultures could be proposed as an innovative source of the chosen phenolic compounds.

Keywords: plant biotechnology, *in vitro* cultures, phenolic acids, *Aronia* species

References

- Kokotkiewicz, A., Jaremicz, Z., Łuczkiwicz, M., 2010. *Aronia* plants: a review of traditional use, biological activities, and perspectives for modern medicine. J. Med. Food 13, 255-269.
- Ellnain-Wojtaszek, M., Zgórk, G., 1999. High-performance liquid chromatography and thin-layer chromatography of phenolic acids from *Ginkgo biloba* L. leaves collected within vegetative period. J. Liq. Chrom. Rel. Tech. 22, 1457-1471.

* Presenting author. E-mail address: a.szopa@uj.edu.pl (Agnieszka Szopa)

S1.P162 Meroterpenoids from *Centrapalus pauciflorus* leaves with antiproliferative activity

Muhammad Bello Saidu^a, Gordana Krstić^a, Anita Barta^a, Hazhmat Ali^b, István Zupkó^b, Umar Shehu Gallah^c, Dóra Rédei^a, Judit Hohmann^{a,d,*}

^a Department of Pharmacognosy, University of Szeged, Eötvös u. 6, 6720 Szeged, Hungary

^b Institute of Pharmacodynamics and Biopharmacy, University of Szeged, Eötvös u. 6, 6720 Szeged, Hungary

^c Bioresource Department, National Research Institute for Chemical Technology (NARICT), Zaria, Nigeria

^d HUN-REN-USZ Biologically Active Natural Products Research Group, University of Szeged, Eötvös u. 6, H-6720 Szeged, Hungary

Centrapalus pauciflorus (Willd.) H. Rob (Asteraceae) is an annual or sometimes short-lived perennial herb with tropical African origin. In the African traditional medicine, its leaves are used as tea or cooked into porridge to ease chest pain and stomach-ache (Chhabra et al., 1989). Previous phytochemical studies of *C. pauciflorus* revealed the presence of fatty acids, coumarins, flavonoids, phenolic acids, sesquiterpenes, and triterpenoids (Maroyi, 2000). In continuation of our ongoing study on *C. pauciflorus* (Saidu et al., 2023; Krstić et al., 2023), the present paper reports the isolation, and structure determination of twelve previously unreported meroterpenoids from the chloroform extract of the leaves besides the known ethuliaconyzophenone. Solvent-solvent partition, OCC, VLC, and HPLC were used for isolation of the compounds. The structures were determined using extensive NMR spectroscopy and HRESIMS. The compounds proved to be hybrid molecules of monoterpenes with 5-methylcoumarin, 5-methylchromone, or acetophenone. New carbon skeletons, dimeric meroterpenoids, and the first meroterpenoid containing a monoterpene unit fused with both a coumarin and an acetophenone part are among the structural novelties. Nine isolated compounds were tested for antiproliferative activity against human adherent malignant cell lines of gynecological origin using the MTT assay, and six of them exhibited moderate activity (IC₅₀ 10 – 25 µM) against MCF-7, HeLa, and A2780 cells.

Keywords: *Centrapalus pauciflorus*, meroterpenoids, centrapalus coumarin, pauciflorin, X-ray

References

- Chhabra, S. C., Mahunnah, R. L. A., Mshiu, E. N. 1989. Plants used in traditional medicine in Eastern Tanzania. II. Angiosperms (Capparidaceae to Ebenaceae). J. Ethnopharmacol. 25, 339–359.
- Maroyi, A., 2020. *Centrapalus pauciflorus*: Review of its medicinal uses, phytochemistry and pharmacological properties. J. Pharm. Nutr. Sci. 10, 287–294.
- Saidu, M. B., Krstić, G., Todorović, N., Berkecz, R., Ali, H., Zupkó, I., Hohmann, J., Rédei, D. 2023. Monoterpenoid 5- methylcoumarins from *Centrapalus pauciflorus* with antiproliferative activity. Arab. J. Chem. 16, 104777.
- Krstić, G., Saidu, M. B., Bombicz, P., De, S., Ali, H., Zupkó, I., Berkecz, R., Gallah, U. S., Rédei, D., Hohmann, J. 2023. Pauciflorins A–E, unexpected chromone–monoterpene-derived meroterpenoids from *Centrapalus pauciflorus*. J. Nat. Prod. 86, 891–896.

* Presenting author. E-mail address: hohmann.judit@szte.hu (Judit Hohmann)

S1.P163 Volatiles derived from Philippine plant species exhibited antimicrobial activity, antioxidative and cytotoxicity using a newly developed microplate-based assay

Genesis Albarico^{a,f,*}, Marketa Houdkova^a, Ivo Daskocil^b, Jan Tauchen^c, Klara Urbanova^d, Edgardo E. Tulin^e, Ladislav Kokoska^a

^a Department of Crop Sciences and Agroforestry, Faculty of Tropical AgriSciences, Czech University of Life Sciences Prague, 165 00 Prague, Czech Republic

^b Department of Microbiology, Nutrition and Dietetics, Faculty of Agrobiolgy, Food and Natural Resources, Czech University of Life Sciences Prague, 165 00 Prague, Czech Republic

^c Department of Food Science, Faculty of Agrobiolgy, Food and Natural Resources, Czech University of Life Sciences Prague, 165 00 Prague, Czech Republic

^d Department of Sustainable Technologies, Faculty of Tropical AgriSciences, Czech University of Life Sciences Prague, 165 00 Prague, Czech Republic

^e Philrootcrops, Visayas State University, Baybay City 6521, Philippines

^f Department of Pure and Applied Chemistry, Visayas State University, Baybay City 6521, Philippines

Essential oils derived from Philippine plant species were evaluated for bioactivity using a newly developed microplate-based bioassay. It has been known that several volatile plant-derived products exhibit considerable bioactivities, however, results of conventional microplate-based assays used to evaluate such highly shifting vapor products can sometimes yield erratic and unusual results. With the aim to demonstrate such phenomenon, antimicrobial, antioxidant, and cytotoxic activities of three Philippine essential oils (*Alpinia elegans*, *Cinnamomum iners*, and *Xanthostemon verdugonianus*) were evaluated in a series of experiments including both ethylene vinyl acetate (EVA) Capmat sealed and nonsealed microplates. The results clearly illustrated that vapor transition to adjoining wells causes false-positive results of bioassays performed in nonsealed microtiter plates. The microplate layout and a duration of the assay were demonstrated as the key aspects defining level of the results affected by the vapors of volatile agents. Additionally, we reported biological activities and chemical composition of essential oils from *A. elegans* seeds and *X. verdugonianus* leaves, which were, according to our best knowledge, were analyzed for the first time. Considering our findings, certain modifications of conventional microplate-based assays are necessary (e.g., using EVA Capmat as vapor barrier) to obtain reliable results when biological properties of volatile agents are evaluated.

Keywords: broth microdilution, essential oil, microtiter plate, supercritical CO₂ extract, volatilization

References

Houdkova M, Albarico G, Daskocil I, Tauchen J, Urbanova K, Tulin EE, Kokoska L. 2020. Vapors of Volatile Plant-Derived Products Significantly Affect the Results of Antimicrobial, Antioxidative and Cytotoxicity Microplate-Based Assays. *Molecules*. 25, 6004.

* Presenting author. E-mail address: genesis.albarico@vsu.edu.ph (Genesis C. Albarico)

S1.P164 Metabolomics-driven discovery of new ophiobolin-type sesterterpenes from marine-derived fungus *Pleosporales* sp. and their antimicrobial activities

Sailesh Maharjan^{a,*}, Johan Mattias Isaksson^b, Monika Krupova^c, Teppo Rämä^a, Kine Østnes Hansen^a, Jeanette Hammer Andersen^a, Espen Holst Hansen^a

^a Marbio, Norwegian College of Fishery Science (NFH), Faculty for Biosciences, Fisheries, and Economics, UiT-The Arctic University of Norway, Tromsø, Norway

^b Department of Pharmacy (IFA), Faculty of Health Sciences, UiT-The Arctic University of Norway, Tromsø, Norway

^c Hylleraas Centre for Molecular Sciences, Department of Chemistry (IK), Faculty of Science and Technology, UiT-The Arctic University of Norway, Tromsø, Norway

In the quest to discover new bioactive chemical entities from nature, a bottleneck is natural product isolation, which can be overcome by integrating several layers of data into multi-informative approaches (Martínez-Aldino et al., 2023). A combination of taxonomy, nontargeted metabolomics, and bioactivity information has led to the selection of a marine-derived *Pleosporales* fungus i.e. *Uzbekistanica* sp. to isolate biologically active new natural products. The strain was prioritized based on the hypothesis that investigating poorly studied and highly metabolic taxa could lead to the isolation of new natural products. A non-targeted metabolomics investigation of the fungal extract using molecular networking indicated the presence of a terpenoid cluster with several unidentified compounds within (Wang et al., 2016). The isolation of the targeted compounds then led to new Ophiobolin-type sesterterpenes (**1-8**). Their structures were determined by extensive spectroscopic analysis and the absolute configurations were further determined by ECD assessment. The antimicrobial effects of these compounds were tested against several pathogenic bacteria and a fungal pathogen. Compounds **5**, **6**, and **8** exhibited inhibitory activity against *Streptococcus agalactiae* with MIC values of 85, 70, and 70 µM, respectively. In addition, **2** showed moderate activity against *Candida albicans* at 100 µM concentration.

Keywords: pleosporales, *Uzbekistanica*, sesterterpenoids, bipolarolides, ophiobolin

References

Martínez-Aldino, I.Y., Rivera-Chávez, J., & Morales-Jiménez, J., 2023. Integrating Taxonomic and Chemical Diversity of Mangrove-Associated Ascomycetes to Discover or Repurpose Bioactive Natural Products. *J. Nat. Prod.* 86, 2423-2434. Wang, M., Carver, J.J., Phelan, V.V., Sanchez, L.M., Garg, N., Peng, Y., Nguyen, D.D., Watrous, J., Kapon, C.A., Luzzatto-Knaan, T., et al., 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat. Biotechnol.* 34,828-837.

* Presenting author. E-mail address: sailesh.maharjan@uit.no (Sailesh Maharjan)

S1.P165 Molecular networking of actinobacteria sourced from thai environments reveals the presence of metabolites with antimicrobial properties

Yuwathida Sunghanghwa^a, Samson Cheruiyot Koech^{a,b}, Amit Jaisi^{a,b,*}

^a College of Graduate Studies, Walailak University, Nakhon Si Thammarat, Thailand

^b School of Pharmacy, Walailak University, Nakhon Si Thammarat, Thailand

^c Biomass Excellence Center, Walailak University, Nakhon Si Thammarat, Thailand

Actinomycetes, particularly from mangrove and marine sediments in Thailand, are relatively unexplored for their potential antimicrobial properties. The aim of this study was to isolate actinobacteria using different media and evaluate the antimicrobial potential of the isolated actinobacteria against methicillin resistant *Staphylococcus aureus*, *Acinetobacter baumannii*, *Klebsiella pneumoniae* and the chloroquine-resistant *Plasmodium falciparum* strain K1 (Septama et al 2022; Phuwajaroanpong et al 2022). Methanolic crude extracts of isolates demonstrated significant in vitro inhibitory activities against chloroquine-resistant *P. falciparum* lactate dehydrogenase (pLDH), agar-overlay and microdilution bacterial assays. The active isolates were further subjected for metabolite profiling using LCMS aided by a global molecular networking platform. We identified numerous active and known compounds, such as Geldanamycin and elaiophyllin. Additionally, we also performed the identification of the selected actinobacterial isolates using 16S rRNA sequencing. Further studies are needed to determine the complete chemical structures and explore the biosynthetic gene clusters. Lastly, our research aimed to provide sustainable solutions for antimicrobial drug resistance, highlighting the potential of actinobacteria-derived compounds in global health initiatives.

Keywords: Microbial natural products, antibiotics, actinobacteria, antimicrobial resistance, drug discovery

References

- Septama, A.W., Tasfiyati, A.N., Kristiana, R. and Jaisi, A., 2022. Chemical profiles of essential oil from Javanese turmeric (*Curcuma xanthorrhiza* Roxb.), evaluation of its antibacterial and antibiofilm activities against selected clinical isolates. *S Afr J Bot.* 146. 728-734.
- Phuwajaroanpong, A., Chaniad, P., Plirat, W., Phoopha, S., Septama, A.W., Chukaew, A. and Punsawad, C., 2022. Antiplasmodial properties of aqueous and ethanolic extracts of ten herbal traditional recipes used in Thailand against *Plasmodium falciparum*. *Infect Dis Trop Med.* 7. 417.
- Wang, M., Carver, J.J., Phelan, V.V., Sanchez, L.M., Garg, N., Peng, Y., Nguyen, D.D., Watrous, J., Kaponov, C.A., Luzzatto-Knaan, T. and Porto, C., 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat Biotechnol.* 34. 828-837.

* Presenting author. E-mail address: amit.ja@mail.wu.ac.th (Amit Jaisi)

S1.P166 MSMS-based molecular networking and fragmentation patterns reveal new insights into the astins chemodiversity from the fungal endophyte *Cyanodermella asteris*

Valdet Uka*, Shen Tang, Tobias A.M. Gulder

Chair of Technical Biochemistry, TU Dresden, Bergstr. 66, 01069 Dresden, Germany

Among the endless number of biologically active cyclopeptides encountered in nature, the plant-derived family of astins represents an interesting case of cyclic peptides with promising anti-tumor and immunomodulating bioactivity. Astins were first isolated from the root extracts of the flowering plant *Aster tataricus* L.f. (Asteraceae) which has a long history of use in traditional Chinese medicine (TCM) as antitussive and expectorant remedy (Yu et al., 2015). Chemically, astins are macrocyclic 16-membered pentapeptides which may contain unusual non-proteinogenic (L-Abu, L-*allo*-Thr, b- Phe, L-Ava, and mono- or dichlorinated L-Pro) and proteinogenic (L-Ser and L-Pro) amino acids. Up to the present, sixteen derivatives (Astin A-I and K-P) from the astin series of cyclopentapeptides have been characterized as natural-occurring peptides (Xu et al., 2013). Recently it was shown that three members of astin family can be produced by the newly discovered fungal endophyte *Cyanodermella asteris* (Schafhauser et al., 2019). In this study, mass spectrometry-based molecular networking and detailed inspection of the fragmentation trees uncovered the idea that *C. asteris* can produce many more astins including proline desaturated derivatives, aminovaleric acid-containing analogues, formylated and acetylated derivatives and some lipopeptide astins. These findings give new information on astin biosynthesis in fungi.

Keywords: astins, fungi, molecular networking, high-resolution MS

References

- Yu P., Cheng S., Xiang J., Yu B., Zhang M., Zhang C., Xu X., 2015. Expectorant, antitussive, anti-inflammatory activities and compositional analysis of *Aster tataricus*. *J. Ethnopharmacol.* 164, 328-33.
- Xu H.M., Zeng G.Z., Zhou W.B., He W.J., Tan N.H., 2013. Astins K-P, six new chlorinated cyclopentapeptides from *Aster tataricus*. *Tetrahedron.* 69, 7964-9.
- Schafhauser, T., Jahn, L., Kirchner, N., Kulik, A., Flor, L., Lang, A., Ludwig-Müller, J., 2019. Antitumor astins originate from the fungal endophyte *Cyanodermella asteris* living within the medicinal plant *Aster tataricus*. *Proc. Nat. Acad. Sci.*, 116, 26909-26917.

*Presenting author. E-mail address: valdet.uka@tu-dresden.de (Valdet Uka)

S1.P167 Phytochemicals and Enzyme inhibition potential of *Rydingia* and *Moluccella* Species in Iran

Amir Reza Jassbi*, Mina Doorandishan, Najmeh Khalighian, Somayeh Pirhadi

Medicinal and Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

Three plants of the genus *Otostegia* Benth (Lamiaceae): *O. persica* (Burm.) Boiss., *O. aucheri* Boiss.; and *O. michauxii* Briq (Mozaffarian, 2003) are renamed to: *Rydingia persica* (Burm.f.) A.-C. Scheen & V. A. Albert and *Rydingia michauxii* (Briq.) A.-C. Scheen & V. A. Albert and *Moluccella aucheri* (Boiss.) A.-C. Scheen, based on their molecular and morphological studies (Scheen and Albert, 2009). We have isolated and identified several diterpenoids and flavonoids with α -glucosidase and acetylcholine esterase activities. *ent*-Labda-8(17),13-dien-18-oic acid-15,16-olide, its glycoside ester, and a trimethoxy flavone: 5, 4'-dihydroxy-6,7,3'-trimethoxy flavone (Eupatorin) were isolated from

R. michauxii. 6 β -acetoxy-9 α -hydroxy-7-oxa-labd-13-ene-15,16-olide, 5-hydroxy-7,4-dimethoxy- flavone and genkwanin were reported from *M. aucheri* (Doorandishan et al., 2021). Finally, four labdane diterpenoids: 15,16-epoxy-3 α , 7 β , 9 α -trihydroxylabdan-13-(16), 14-dien-6-one; 15,16- epoxy-3 α , 7 α , 9 α -trihydroxylabdan-13-(16),14-dien-6-one; 9,13,15,16-diepoxy-3 α ,7 β ,15 α (β)- trihydroxy-labdan-6-one were characterized in an ethyl acetate fraction of an ethanol extract of the aerial parts of *R. persica*. The structures of the compounds were elucidated by spectral data analyses of their ^1H and ^{13}C NMR and ESIMS spectroscopy. The *in vitro* and *in silico* α -glucosidase and acetylcholine esterase activity of the isolated compounds are reported in this presentation.

Acknowledgments: The study was financially supported by the National Institutes for Medical Research Development (NIMAD), Tehran, Iran (grant number: 988197).

Keywords: *Otostegia*, *Rydingia persica*, *Rydingia michauxii*, *Moluccella aucheri*, labdane diterpenoids

References:

- Mozaffarian V., 2003. A dictionary of Iranian plant names. Tehran: Farhang Moaser. 387.
- Scheen AC, Albert VA., 2009. Molecular phylogenetics of the Leucas group (Lamioideae; Lamiaceae). Syst Bot. 34, 173- 181.
- Doorandishan M, Pirhadi S, Swilam M. M, Gholami M, Ebrahimi P, El-Seedi H. R, Jassbi A. R., 2021. Molecular docking and simulation studies of a novel labdane type- diterpene from *Moluccella aucheri* Scheen (Syn. *Otostegia aucheri*) as human- AChE inhibitor. J. Mol. Struct, 1245, 131034.

* Presenting author. E-mail address: jassbiar@sums.ac.ir (Amir Reza Jassbi)

S1.P168 Phenanthrenes from the aerial parts of *Juncus tenuis*

Anita Barta^a, Petra Petz^a, Dragica Purger^b, Andrea Vasas^{a,c}, Judit Hohmann^{a,c,*}

^a Department of Pharmacognosy, University of Szeged, Eötvös u. 6, 6720 Szeged, Hungary

^b Department of Pharmacognosy, University of Pécs, Rókus u. 2, 7624 Pécs, Hungary

^c HUN-REN-USZ Biologically Active Natural Products Research Group, University of Szeged, Eötvös u. 6, 6720 Szeged, Hungary

Phenanthrenes are a promising group of natural small molecules, with great structural variety and noteworthy pharmacological (e.g., antiproliferative, antibacterial, and anti-inflammatory) activities (Bús et al., 2018; Tóth et al., 2018). Although, Juncaceae is a relatively small plant family with approximately 500 plant species worldwide, almost 20% of currently known natural phenanthrenes were described from species belonging to this family. Especially members of the *Juncus* genus are rich sources of phenanthrenes. Various *Juncus* species are used mainly in traditional Chinese medicine for the treatment of numerous disorders (e.g. fidgetiness, insomnia, painful urination, pharyngitis, and aphtha). Medulla Junci, the dried stem pith of *J. effusus*, is official in the Pharmacopoeia of the People's Republic of China. In continuation of our search for phenanthrenes from plants belonging to the family Juncaceae, *Juncus tenuis* was investigated. The dried and ground plant material was extracted with methanol. After evaporation, it was subjected to solvent-solvent partition using *n*-hexane, chloroform, and finally ethyl acetate. Phenanthrenes were enriched in the chloroform phases; therefore, it was fractionated by vacuum liquid chromatography, gel filtration on Sephadex LH-20 gel, and in the last step by HPLC to obtain pure compounds. The structure elucidation of the compounds was carried out by extensive NMR spectroscopic analysis, and HRMS experiments as well as by comparison of spectroscopic data with literature values. The results allowed the identification of 17 phenanthrenes, among them 14 monomers and three dimers from *J. tenuis*. Four compounds are previously undescribed natural products.

Keywords: *Juncus tenuis*, Juncaceae, phenanthrenes

References

- Bús, C., Tóth, B., Stefkó, D., Hohmann, J., Vasas, A., 2018. Family Juncaceae: promising source of biologically active natural phenanthrenes. *Phytochem. Rev.* 17, 833-851.
- Tóth, B., Hohmann, J., Vasas, A., 2018. Phenanthrenes: a promising group of plant secondary metabolites. *J. Nat. Prod.* 81, 661-678.

* Presenting author. E-mail address: hohmann.judit@szte.hu (Judit Hohmann)

S1.P169 Discovery of anti-inflammatory cyanobacterial secondary metabolites targeting the Keap1/Nrf2 pathway

Fatma H. Al-Awadhi^{a,b,*}, Ranjala Ratnayake^b, Valerie J. Paul^c, Hendrik Luesch^b

^a Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Kuwait University, P.O. Box 24923, Safat 13110, Kuwait

^b Department of Medicinal Chemistry and Center for Natural Products, Drug Discovery and Development (CNPDP3), University of Florida, Gainesville, FL 32610, USA

^c Smithsonian Marine Station, Fort Pierce, Florida, FL 34949, USA

Chemical investigation of marine cyanobacterial samples collected from Florida Keys resulted in the discovery of two secondary metabolites belonging to two different structural classes. The planar structures of both compounds were elucidated using a combination of 1D and 2D NMR spectroscopy and mass spectrometry. The structures feature an α,β -unsaturated carbonyl system, a key motif required for the activation of the Keap1/Nrf2–ARE pathway, a cytoprotective pathway that controls the activation of antioxidant genes and phase II detoxification enzymes. The compounds were screened in ARE-luciferase reporter gene assay using stably transfected HEK293 cells and found to induce Nrf2 activity up to 10 μ M. Due to the established crosstalk between Nrf2 and NF- κ B pathways, the anti-inflammatory potential of both compounds was investigated in LPS-induced mouse macrophages (RAW264.7 cells), a commonly used model for inflammation. The compounds significantly upregulated *Nqo1* (Nrf2 target gene) and downregulated *iNos* (NF- κ B target gene) resulting in a significant reduction of nitric oxide (NO) levels. Furthermore, we performed RNA-sequencing and demonstrated the transcriptional activity of both compounds on a global level and identified canonical pathways and upstream regulators involved in inflammation, immune response, and certain oxidative stress-underlying diseases such as multiple sclerosis.

Keywords: marine cyanobacteria, inflammation, RNA sequencing, immune response, multiple sclerosis

* Presenting author. E-mail address: fatma.h@ku.edu.kw (Fatma H. Al-Awadhi)

S1.P170 A terpenoids-enriched fraction of *Inula viscosa*: its chemical composition and anti-cancerous effects against human lung cancer cells

Fatiha Seglab^a, Mazen Mounir Abu Asali^b, Thoraya Al-Yafei^b, Hassan Hassan^b, Abdullah A. Shaito^{a,*}

^a Biomedical Research Center, Qatar University, Doha P.O. Box 2713, Doha, Qatar

^b Environmental Science Center, Qatar University, Doha P.O. Box 2713, Doha, Qatar

Inula viscosa has been a heavily used plant in traditional Mediterranean and Middle Eastern medicine to handle various illnesses. *I. viscosa* has been shown to have anticancerous effects against a variety of cancers, but its effects against lung cancer have been under limited investigation. The aim of the present study is to examine the potential anticancerous properties of methanolic or aqueous extracts of stems and leaves of *I. viscosa* against human lung cancer A549 cells. *I. viscosa* leaves methanolic extracts (IVLM) showed the highest reduction in viability of A549 cells among all the extracts. IVLM also reduced the viability of a panel of human cancer cell lines. Fractionation of IVLM by liquid-liquid separation using hexane, ethyl acetate and dichloromethane (DCM) solvents revealed that *I. viscosa* DCM fraction (IVL DCM) displayed the most notable reduction in viability of A549 cells ($IC_{50} = 28.68 \pm 3.83 \mu\text{g/mL}$ at 48 h), and was not cytotoxic to normal human embryonic fibroblasts (HDFn). This led to a detailed assessment of IVL DCM phytochemical constituents using GC-MS analysis which revealed 22 metabolites, highlighting an enrichment in terpenoids like 9,19-cyclolanostan-3-ol acetate, betulin, longifolene and lupeol, known for their pro-apoptotic, pro-autophagic and anti-metastatic functions. Furthermore, analysis of proliferation, autophagy, and apoptosis markers by Western blotting indicated that IVL DCM reduced proliferation (reduction of proliferation marker Ki67 and induction of proliferation inhibitor proteins P21 and P27) and induced intrinsic apoptotic pathway (P38/P53/BCL-2/BAX/ Caspase 3) in A549 cells. IVL DCM also reduced the migration of A549 cells, potentially through reduction of FAK activation. Future identification of anticancer metabolites of IVL DCM may emphasize its terpenoids as anti-lung cancer agents, and confirm *I. viscosa* as a resource of herbal anticancer agents.

Keywords: *Inula viscosa*, lung cancer, A549 cells, phytochemical analysis, cytotoxicity

* Presenting author. E-mail address: abdshaito@qu.edu.qa (Abdullah A. Shaito)

S1.P171 Molecular identification and metabolite profiling of endophytes isolated from *Bambusa vulgaris* var. *striata* shoot

Nusirat Aderinsola Sadiku*, Nurah Eniola Shuaib

Department of Forest Resources Management, University of Ilorin, Ilorin, Nigeria

Endophytic fungi have been reported to produce many plant-derived biochemicals. In this study, endophytic fungi were isolated from *Bambusa vulgaris* var. *striata* shoot and the biocompounds in the ethanolic extract of the endophytes were investigated. Three endophytes comprising of 2 strains of *Candida albicans* and 1 strain of *Fusarium oxysporum* were characterized by 18S rRNA gene and internal transcriber spacer region sequence analysis. GC-MS analysis revealed 18, 30 and 36 biocompounds in the ethanolic extracts of *C. albicans* strain KY101911.1, MK805514.1 and *F. oxysporum* respectively, with variations in the concentration as well as the type of the phytoconstituents in these endophytes. However, the main biocompounds in each of the endophytes were the 9-octadecenoic acid, methyl ester (E)- (18.58%), 11,14-octadecadienoic acid, methyl ester (18.77%), hexadecanoic acid ethyl ester (21.66%) and butyl 9,12-octadecadienoate (12.15%) for *C. albicans* strain MK805514.1, KY101911.1, and *F. oxysporum* respectively. Generally, some of the compounds occurred in fairly large amounts while others occurred in trace amounts. However, the phytoconstituent with the largest concentration was hexadecanoic acid ethyl ester in the extract of *C. albicans* strain MK805514.1. Seven compounds (hexadecanoic acid, ethyl ester; hexadecanoic acid, methyl ester; ethyl stearate; 11,14-octadecadienoic acid, methyl ester; 9-octadecenoic acid, methyl ester (E)-; n-propyl 9,12-octadecadienoate; ctadecanoic acid, 17-methyl-, methyl ester) were similalry revealed in each of the endophytes but in varying quantities. These findings indicate that *B. vulgaris* var. *striata* shoot is a huge reservoir of micro-organisms that could be useful for the development of novel drugs with therapeutic properties (Khiralla et al., 2015).

Keywords: bamboo shoot, endophytes, *C. albicans*, *F. oxysporum*, phytochemicals

References

Khiralla, A., Mohamed, I., Thomas, J., Mignard, B., Spina, R., Yagi, S., Laurain-Mattar, D., 2015. A pilot study of antioxidant potential of endophytic fungi from some Sudanese medicinal plants. *Asian Pac J Trop Med.* 8, 701-4.

* Presenting author. E-mail address: sadiku.na@unilorin.edu.ng (Nusirat Aderinsola Sadiku)

S1.P172 *Annulohypoxylon stygium*, an endophyte from the marine red alga *Asparagopsis taxiformis* is a source of antiviral binaphthalenes

Dulce Silva^{a,*}, Givaldo S. da Silva^{a,b}, Iatã Mendonça^a, Victor Rufino^a, Yan Fernandes^a, Lucas Moura^a, Edelson Dias^a

^a São Paulo State University - UNESP, Institute of Chemistry, Araraquara, SP, Brazil

^b Federal Institute of Education, Science and Technology of Acre – IFAC, AC, Brazil

Naphthalene derivatives with unique structures are widely found in *Annulohypoxylon* species. Truncatones A-D have been isolated from *A. leptascum* and *A. truncatum* and showed cytotoxic activity against mouse fibroblast cells (Sudarman, 2016). A strain of *A. stygium* was isolated from the red alga *Asparagopsis taxiformis* and grown on red rice solid medium for 20 days. After extraction with MeOH and solvent removal, the MeOH extract was dissolved into MeOH:H₂O (4:1) and defatted with *n*-hexane. The hydromethanol fraction was evaporated to yield 1.2 g of extract, which was subjected to subsequent MPLC (C18) and Sephadex LH-20 fractionation steps to yield verrucosapyrone B and diketopiperazines along with asperlone B (**1**), obtained as an amorphous red powder. Asperlone B showed a molecular formula C₂₀H₁₂O₈, determined by ESI-QqTOF-MS in positive mode, and was isolated for the first time as a fungal metabolite from the genus *Annulohypoxylon*. Its structure was established by NMR and ESI-QqTOF-MS data, and comparison with the literature (Xiao, 2015).




Fig. 1. Structure of asperlone B.

Molecular docking analysis of asperlone B evidenced effectiveness in modulating the biomacromolecular target of the Mayaro virus, confirmed by the release of -7.0 kcal/mol, mainly due to the resonance effect inherent to its aromatic rings, which allowed PI-type interactions with the amino acids lysine-138 and alanine-256, as well as Van der Waals interactions with further amino acids in the reaction site. Once this ligand is complexed in the hydrophobic site, the virus capsid bonds are no longer effective, thus interrupting its cycle. Such results confirmed marine endophytic fungi as a relevant source for potential antiviral metabolites.

Keywords: marine fungi, asperlones, endophytic fungus, *Annulohypoxylon*, molecular docking

References

- Sudarman, E., et al., 2016. Truncatones A-D, benzo[j]fluoranthenes from *Annulohypoxylon* species (Xylariaceae, Ascomycota). *Tetrahedron* 72, 6450-6454.
- Xiao Z., et al., 2015. Asperlones A and B, dinaphthalenone derivatives from a mangrove endophytic fungus *Aspergillus* sp. 16-5C. *Marine Drugs* 13, 366-378.

* Corresponding author. E-mail address: dulce.silva@unesp.br (Dulce Silva)

S1.P173 Antibacterial anthraquinone produced by the fungal endophyte *Botryosphaeria rhodina*, isolated from the mangrove *Xylocarpus granatum* J. Koenig

Rudi Hendra^{a,*}, Tetty Marta Linda^b, Yuli Haryani^a

^a Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Riau, Pekanbaru, Indonesia

^b Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Riau, Pekanbaru, Indonesia

The exploration of the secondary metabolites of fungus originating from mangroves, known for their notable biological activity and promise in pharmaceuticals, has attracted substantial attention (Deshmukh et al., 2020). Our previous investigation focused on isolating endophytic fungi from mangrove vegetation with antimicrobial characteristics. Among these fungi, *Botryosphaeria rhodina* was identified from the mangrove *Xylocarpus granatum* J. Koenig. (Linda et al., 2024).

In the continuum of our research focused on deriving antibacterial agents from endophytic fungi associated with mangroves, this study aims to isolate these agents from the fungi. To this end, solid-state fermentation was employed to yield an extract rich in antibacterial compounds, which was then subjected to vacuum liquid chromatography using a solvent system of n-hexane, ethyl acetate, and methanol, resulting in ten distinct fractions. From fraction 6, an orange precipitate was obtained, which was further purified through re-crystallization to produce orange crystals. Subsequent elucidation and comparison of spectroscopic data confirmed the compound as macrosporin, an anthraquinone derivative (**1**), as depicted in Figure 1.




Fig. 1. Structure of macrosporin 1

Compound **1** demonstrated remarkable activity against a range of pathogenic bacteria, with Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) values between 25 and 100 µg/mL. This data underscores the potency of macrosporin as an antibacterial agent, instilling confidence in the potential of our findings. Furthermore, *in silico* analyses of **1**'s antibacterial mechanism revealed a strong affinity for the bacterial receptor's active and allosteric sites, enabling it to disrupt cell wall integrity and inhibit protein synthesis.

Keywords: antibacterial, anthraquinone, *Botryosphaeria rhodina*, *in vitro*, *in silico*

References

- Deshmukh, S. K., Agrawal, S., Prakash, V., Gupta, M. K., Reddy, M. S. 2020. Anti-infectives from mangrove endophytic fungi. South African Journal of Botany, 134, 237-263.
- Linda, T. M., Hendra, R., Farniga, A., Delfira, N., Ismed, F., Haryani, Y. 2024. Potential antibacterial activity of *Botryosphaeria rhodina*: Mangrove *Xylocarpus granatum* J. Koenig. derived fungal endophyte. Journal of Applied Pharmaceutical Science, 14, 136-143.

* Presenting author. E-mail address: rudi.hendra@lecturer.unri.ac.id (Rudi Hendra)

S1.P174 Elucidating phytochemical profiles and antibacterial potential of *Helminthostachys zeylanica* (L.) through untargeted LC-HRMS metabolomics analysis

Hilwan Yuda Teruna, Monika Kerry Army, Yuana Nurulita, Rudi Hendra*

Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Riau, Pekanbaru, Indonesia

Helminthostachys zeylanica (L.) Hook. is renowned for its traditional therapeutic uses in managing diabetes mellitus and other illnesses in China, Malaysia, and Indonesia (Ridhasya et al., 2019). During our previous investigation into the antibacterial properties of several extracts derived from the roots of this species, we discovered that dichloromethane and ethyl acetate extracts showed inhibitory effects against various pathogenic bacteria (Army et al., 2023). This study employs LC-HRMS for untargeted metabolomic analysis of *H. zeylanica*'s various parts (root, stem, leaf) to elucidate its chemical profile and antibacterial potential. We identified diverse secondary metabolites, highlighting the plant's complex chemistry. The antibacterial properties were assessed using the Kirby-Bauer disc diffusion method and further quantified by determining the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC). Our research findings suggest that extracts from *H. zeylanica* demonstrate moderate antibacterial activity, particularly against Gram-positive bacteria. However, their effectiveness is lower compared to chloramphenicol. Advanced metabolomic profiling techniques were used, including Principal Component Analysis (PCA) and Partial Least Squares Discriminant Analysis (PLS-DA). This analysis revealed distinct metabolic patterns in different sections of plants, highlighting the diverse phytochemical contents. These investigations helped identify promising antibacterial metabolites. The study affirms the existence of distinct bioactive chemicals in *H. zeylanica*. However, their effectiveness against Gram-negative bacteria is limited compared to conventional antibiotics. This comprehensive metabolomic method improves our comprehension of the pharmacological properties of *H. zeylanica*.

Keywords: LC-HRMS, metabolomics, *Helminthostachys zeylanica*, antibacterial activity, secondary metabolites

References

- Army, M.K., Khodijah, R., Haryani, Y., Teruna, H. Y., Hendra, R., 2023. Antibacterial *in vitro* screening of *Helminthostachys zeylanica* (L.) Hook. root extracts. J. Pharm. Pharmacogn. Res. 11, 291-296.
- Ridhasya, F. E., Teruna, H. Y., Hendra, R., Almurdani, M., 2019. Natural Antidiabetic of Tunjuk Langit (*Helminthostachys zeylanica*) Rhizome Extracts. Pharmacol. Clin. Pharm. Res. 4, 67-70.

* Presenting author. E-mail address: rudi.hendra@lecturer.unri.ac.id (Rudi Hendra)

S1.P175 Sesquiterpenoids and hexanorcucurbitacin from *Aquilaria malaccensis* agarwood with anti- inflammatory effects by inhibiting STAT1/AKT/MAPK/NLRP3 pathway

Chi Thanh Ma^a, Tianqi Huang^b, Hyun Ok Yang^b, Jae Sik Yu^{b,*}

^a Faculty of Pharmacy, University of Medicine and Pharmacy at Ho Chi Minh City, 41-43 Dinh Tien Hoang st., dist. 1, Ho Chi Minh city 700000, Vietnam

^b Department of Integrative Biological Sciences and Industry and Convergence Research Center for Natural Products, Sejong University, 209, Neungdong-ro, Gwangjin-gu, Seoul 05006, Korea

Seven undescribed compounds **1-7**, including four sesquiterpenoids, one azulene-type, one indene-type, and one rare hexanorcucurbitacin, together with eleven known ones (**8-16**), were isolated from the agarwood chips of *Aquilaria malaccensis* (Fig. 1.). The structures of isolated compounds were elucidated by extensive spectroscopic methods such as mass spectrometry, UV, IR, NMR spectroscopy. The precise stereo-chemical configurations of new compounds were determined by calculated ECD spectra data, as well as a single-crystal X-ray diffraction analysis. This study demonstrates the phytochemical investigation of the extract of *A. malaccensis* and the pharmacological activities of seven previously undescribed compounds. The isolated compounds **1-7** were evaluated by estimating the levels of nitric oxide (NO), TNF- α , and the expression of enzyme iNOS, and COX-2. Among them, a rare hexanortriterpenoid (**7**) derived from a cucurbitane-type triterpenoid showed the significantly attenuated neuro-inflammatory effects via STAT1/AKT/MAPK/NLRP3 signaling pathway on the mechanistic studies.




Fig. 1. Chemical structure of compounds **1-16**.

Keywords *Aquilaria malaccensis*, agarwood, hexanorcucurbitacin, isodaucane-type sesquiterpenoid, NLRP3 signaling

References

M. A. Eissa, Y. Z. H. Y. Hashim, S. S. S. Abdul Azziz, H. M. Salleh, M. L. M. Isa, N. M. Abd Warif, F. Abdullah, E. Ramadan and D. M. El-Kersh, Phytochemical Constituents of *Aquilaria malaccensis* Leaf Extract and Their Anti- Inflammatory Activity against LPS/IFN- γ -Stimulated RAW 264.7 Cell Line, ACS Omega, 2022, 7, 15637-15646.

* Presenting author. E-mail address: jsyu@sejong.ac.kr (Jae Sik Yu)

S1.P176 Isolation of *Salvia apiana* Jeps. (Lamiaceae) essential oil—the influence of distillation technique and process parameters on the yield and composition of volatile fraction

Adam Kokotkiewicz^{a,*}, Pourya Hesami^a, Agata Król^a, Bożena Zabiegała^b, Maria Łuczkiwicz^a

^a Department of Pharmacognosy, Faculty of Pharmacy, Medical University of Gdańsk, al. gen. J. Hallera 107, 80-416 Gdańsk, Poland

^b Department of Analytical Chemistry, Faculty of Chemistry, Gdańsk University of Technology, ul. Narutowicza 11/12, 80-233 Gdańsk, Poland

Salvia apiana (white sage) is an aromatic plant of mint family, endemic to California Floristic Province. It has long been known to native North American Chumash people who employed white sage as a medicinal and ritual plant. Modern studies confirmed antioxidant, antimicrobial, anti-inflammatory, analgesic and anti-cancer activities of *S. apiana*, which can be largely attributed to its high essential oil content (Krol et al., 2022). In the current work, the influence of distillation technique (conventional hydrodistillation [HD] vs simultaneous distillation-extraction [SDE]), as well as different parameters of the process (sample size, water volume, distillation time and condensate flow) on the yield and composition of essential oil, isolated from dried white sage leaves, have been examined. Of the techniques employed, SDE proved to be superior in terms of essential oil yield (83 ml/kg dry wt, ca. 1.5-2-times higher as compared to HD, depending on the apparatus type used). Regardless of the technique used, the composition of the volatile fraction remained largely the same, with 1,8-cineole, alpha-pinene, beta-pinene and bornanone being the major constituents, as revealed by GC-MS and GC-FID analysis. HD experiments demonstrated that sample to water ratio had negligible effect on essential oil yield and composition. However, the study showed that by extending HD time from 1.5 to 3 h, the yield of volatiles can be increased from ca. 40 to 46 ml/kg dry wt. Higher amounts of essential oil were also obtained at higher condensate flows which enables to shorten the HD process, while maintaining its effectiveness.

Keywords: *Salvia apiana*, essential oil, hydrodistillation, simultaneous distillation-extraction, process optimization

References

Krol, A., Kokotkiewicz, A., Luczkiewicz, M., 2022. White Sage (*Salvia apiana*)—a Ritual and Medicinal Plant of the Chaparral: Plant Characteristics in Comparison with Other *Salvia* Species. *Planta Med.* 88, 604-627.

* Presenting author. E-mail address: adamkokot@gumed.edu.pl (Adam Kokotkiewicz)

S1.P177 Biomasses of *Phyllanthus* spp. (Euphorbiaceae) as a source of sulfonic derivatives of flavonoids

Barbara Sparzak-Stefanowska*, Mirosława Krauze-Baranowska

Department of Pharmacognosy with Medicinal Plant Garden, Faculty of Pharmacy, Medical University of Gdańsk, Gen. J. Hallera 107, 80-416 Gdańsk, Poland

Naturally occurring sulfonic derivatives of flavonoids are an extremely rare group of compounds. Due to their limited distribution, little is known about their biological activity, but also about their biosynthesis levels in plants. Moreover, there is no information on the accumulation of these compounds in *in vitro* conditions. In nature, some species of the genus *Phyllanthus* are unique sources of these secondary metabolites (Than et al., 2006; Thanh et al., 2014). The aim of the study was to evaluate *Phyllanthus* spp. biomasses obtained *in vitro* for the presence of sulfonic derivatives of flavonoids. The study included methanol extracts from biomasses (shoot, callus and adventitious root cultures) obtained from 5 species – *P. grandifolius*, *P. juglandifolius*, *P. glaucus*, *P. amarus* and *P. multiflorus*. Conditions for screening analysis using HPTLC and HPLC methods were developed and the presence of the tested compounds was demonstrated in *P. amarus* (shoot and adventitious root culture) and *P. multiflorus* (shoot culture and callus culture). Shoot cultures of both species were characterized by the presence of one compound, while the callus of *P. multiflorus* and adventitious roots of *P. amarus* were the source of a more complex set of compounds. Preliminary studies indicate that niruriflavone is the dominant compound in all tested biomasses (Fig. 1).




Fig. 1. Chemical structure of niruriflavone.

Keywords: *Phyllanthus*, sulfonic derivatives of flavonoids, niruriflavone, plan *in vitro* cultures

References

- Than, N. N., Fotso, S., Poeggeler, B., Hardeland, R., Laatsch, H., 2006. Niruriflavone, a new antioxidant flavone sulfonic acid from *Phyllanthus niruri*. Zeitschrift fur Naturforschung - Section B Journal of Chemical Sciences 61, 57- 60.
- Thanh, N. V., Huong, P. T. T., Nam, N. H., Cuong, N. X., Thao, N. P., Dejaegher, B., Gordien, A., Heyden, Y. V., Quetin-Leclercq, J., Minh, C. V., 2014. A new flavone sulfonic acid from *Phyllanthus urinaria*. Phytochemistry Letters 7, 182-185.

* Presenting author. E-mail address: bsparzak@gumed.edu.pl (Barbara Sparzak-Stefanowska)

S1.P178 Antibiotic potentiating effects of *Salix* twig extracts against *Bacillus cereus*

Pia Fyhrqvist^{a,*}, Eunice Ego Mgbeahuruike^a, Enass Y.A. Salih^a, Brigita Novak^b

^a Division of Pharmaceutical Biosciences, Faculty of Pharmacy, 00014-University of Helsinki, Finland

^b Faculty of Pharmacy and Biochemistry, HR-1000, University of Zagreb, Croatia

Plant extracts and compounds are known to possess antibiotic potentiating effects (Abreau et al., 2017). The use of plant extracts and compounds as antibiotic adjuvants could be a way to repurpose antibiotics that have lost their effects against resistant bacteria. Moreover, in the combinations it is possible to use the antibiotics in extremely small concentrations, thus reducing their toxic effects. In addition, the risk of development of resistance is reduced for a combination. Only a few *Salix* species, such as the African *S. ledermannii*, have so far been evaluated for their combination effects with antibiotics (Demgne et al., 2022). Thus, we have chosen to study *Salix starkeana*, *S. myrsinifolia* and *S. aurita* that grow naturally in Finland. In our studies, methanol extracts of the winter twigs of these *Salix* species gave growth inhibitory effects against *B. cereus* with MIC values of 625, 1250 and 2500 µg/ml, respectively. In addition, two antibiotics, tetracycline with MIC of 0.488 µg/ml and ciprofloxacin hydrochloride with MIC of 0.06 µg/ml against *B. cereus* were chosen for the combination studies. A 96 well microplate checkerboard method was used to test the combination effects (Mordmuang et al., 2019). To evaluate the results of the combinations, we calculated the FIC indexes (SFIC or FICI) for each combination producing 90% or more of growth inhibition. The FICI was calculated as $\Sigma FIC = FICA + FICB = (CA/MICA) + (CB/MICB)$, where MICA and MICB are the MICs of drugs A and B alone, respectively, and CA and CB are the concentrations of the drugs in combination, respectively. Several combinations of all tested *Salix* extracts with tetracycline hydrochloride resulted in synergistic effects. Moreover, our combination tests showed that even at concentrations of 0.03 ' MIC (0.061 µg/ml), tetracycline produced synergistic effects when combined with a *Salix aurita* methanol twig extract at 625 µg/ml (Fig. 1).

			MIC	0,5 MIC	0,25 MIC	0,125 MIC	0,0625 MIC	0,0312 MIC	0,0156 MIC	0,0078 MIC	
Inh%		Extr in well	2500	1250	625	313	156,5	78,25	39,125	19,56	µg/ml
	Atb in well	Eppendorf	10000	5000	2500	1250	625	313	156,5	78,25	µg/ml
MIC	0,488	1,952	106,000497	100,80752	104,323339	99,105516	100,409972	99,7515322	100,086964	99,8757661	
0,5 MIC	0,244	0,976	132,077191	118,312076	109,690243	101,279609	98,8322014	100,03727	100,273315	100,472089	
0,25 MIC	0,122	0,488	92,2229584	92,272652	107,044062	101,006295	100,633593	100,03727	99,8384959	100,049694	
0,125 MIC	0,061	0,244	99,043399	94,4094749	104,459997	111,479212	98,6831208	91,1297002	84,4459168	93,105019	
0,0625 MIC	0,0305	0,122	124,958589	102,447408	100,496936	88,8686434	57,3629286	46,4676164	45,2128541	41,0510187	
0,0312 MIC	0,01525	0,061	126,113964	86,3591188	90,8936558	51,7351333	46,2688421	41,8958092	31,2862349	5,8555574	
	µg/ml	µg/ml									

Fig. 1. Checkerboard test results as FICI of combinations of a methanol twig extract of *S. aurita* (Extr) with tetracycline (Atb) against *Bacillus cereus*. The y-axis shows tetracycline and the x-axis the plant extract at their MIC related concentrations in Eppendorf tubes and in the microplate wells. The green colour indicates synergistic, the blue additive and the orange intermediate effects.

Keywords: *Salix myrsinifolia*, *S. aurita*, *S. starkeana*, antibiotic potentiating

References

- Abreau, A.C., Coqueiro, A., Sultan, A.R., Lemmens, N., Kim, H.K., Verpoorte, R., van Wamel, W.J.B., Simões, M., Choi, Y.H., 2017. Looking to nature for a new concept in antimicrobial treatments: isoflavonoids from *Cytisus striatus* as antibiotic adjuvants against MRSA. *Sci. Rep.* 7, 1-16.
- Demgne, O.M.F., Tchinda, C.F., Mbaweng, A.T., Beng, V.P., Kuete, V., 2022. Antibacterial and antibiotic-potentiating activities of nine Cameroonian medicinal plants against multidrug-resistant bacteria expressing active efflux pumps. *Inv. Med. Chem. and Pharm.* 5, 1-11.
- Mordmuang, A., Brouillette, E., Voravuthikunchai, S.P., Malouin, F., 2019. Evaluation of a *Rhodomyrtus tomentosa* ethanolic extract for its therapeutic potential on *Staphylococcus aureus* infections using in vitro and in vivo models of mastitis. *Phytotaxa*. 72, 1-111.

* Presenting author. E-mail address: pia.fyhrqvist@helsinki.fi (Pia Fyhrqvist)

S1.P179 HPLC-MS and 2D-TLC analysis of secondary metabolites from the leaves of selected species and hybrids from the genus *Populus* with the estimation of their antioxidant properties

Loretta Pobłocka-Olech*, Sylwia Godlewska, Katarzyna Kimel, Mirosława Krauze-Baranowska

Department of Pharmacognosy with Medicinal Plants Garden, Medical University of Gdańsk, 107 Hallera Str., 80-416 Gdańsk, Poland

Rheumatic diseases require long-term therapy, and usually steroid and non-steroidal anti-inflammatory drugs are used, which cause many side effects. Therefore, there is a growing interest among patients in medicines of plant origin, which effectively alleviate the symptoms associated with rheumatic diseases and are characterized by high safety of use (Uehleke et al., 2011). One of the herbal medicinal products used to treat rheumatism are poplar leaves (*Populi folium*). Currently, the pharmaceutical industry has problems with obtaining plant raw material that meets the requirements contained in the *Populi folium* monograph in the Polish Pharmacopoeia due to the increasingly rare occurrence of the *Populus nigra* species, which is the main source of plant raw material. Therefore, research is being conducted on other *Populus* species regarding compounds with high antioxidant and anti-inflammatory activity. This study presents the HPLC-MS analysis of leaves from three taxa from *Populus* genus - *P. alba*, *P. × candicans* and *P. nigra*, selected in our previous studies (Pobłocka-Olech et al., 2021). Moreover, the separation of secondary metabolites occurring in analyzed poplar leaves by 2D-TLC method were optimized and used for bioautography of their antioxidant and anti-inflammatory properties. The antioxidant activity was estimated by DPPH, ABTS and FRAP assays. In the results, it was found that all analyzed leaves were characterized by the similar contents of phenolics (81.75-85.29 mg/g d.w.), flavonoids (6.23-8.69 mg/g d.w.) and differential content of salicylic derivatives (4.42-36.16 mg/g d.w.). The highest antioxidant activity was assessed for the leaves of *P. nigra*, and similar for *P. × candicans*.

Keywords: *Populus*, phytochemical analysis, antioxidant activity

References

- Uehleke, B., Brignoli, R., Rostock, M., Saller, R., Melzer, J., 2011. Phytodolor (R) in musculoskeletal disorders: re-analysis and meta-analysis. *Forsch. Komplementarmed.*, 18, 249–256.
- Pobłocka-Olech, L., Głód, D., Jesionek, A., Łuczkiwicz M., Krauze-Baranowska, M., 2021. Studies on the polyphenolic composition and the antioxidant properties of the leaves of poplar (*Populus* spp.) various species and hybrids. *Chem. Biodiversity*, 18, e2100227.

* Presenting author. E-mail address: loretta.poblocka-olech@gumed.edu.pl (Loretta Pobłocka-Olech)

S1.P180 Antibacterial and phytochemical study of Bornean Myristicaceae and Clusiaceae species

M. Valmiki^{a,*}, A. Ganesan^a, P. de Resende^b, S. Gibbons^c, J. Wright^a, A. Morritt^a

^a School of Pharmacy, University of East Anglia, Norwich Research Park, NR4 7TJ, UK

^b John Innes Centre, Norwich Research Park, NR4 7UH, UK

^c Centre for Natural Products Discovery, Liverpool John Moores University, L3 3AF, UK

Antimicrobial resistance is a leading cause of morbidity and mortality worldwide. It is defined as the resistance of a microorganism to an antimicrobial agent, to which it was originally susceptible (de Sousa Oliveira, K. et al. 2016). In 2019, 1.3 million deaths were directly caused by antibiotic resistant infections, with estimations that by 2050, such infections could cause 10 million annual premature deaths worldwide, as well as a large economic burden (Murray et al. 2022; O'Neill, J. 2018). The island of Borneo has remarkable biodiversity, including a variety of unexplored plant species, some of which belong to families known for their pharmacological activities, such as the Myristicaceae and Clusiaceae. This project investigates the bioactivity and phytochemistry of the previously unexplored plant species *Horsfieldia splendida*, *H. polyspherula*, *Knema elmerii*, *K. latifolia*, *K. membranifolia*, *Gymnacranthera contracta*, *G. forbesii* (Myristicaceae), *Garcinia caudiculata*, *Calophyllum pulcherimum* and *Mesua calciphila* (Clusiaceae). Here, biological assays including broth microdilution and AlamarBlue, as well as chromatographic techniques (TLC, VLC and HPLC), spectroscopic techniques (LC-MS, NMR) and x-ray crystallography are used to isolate plant metabolites and assess their antibacterial and anticancer potentials. To date, 7 Myristicaceae and 3 Clusiaceae species have been extracted and assessed for antibacterial activities (MIC $\leq$ 64 μ g/mL against *Staphylococcus aureus*). Following this, two known (example, Figure 1) and two novel anacardic acids, as well as one resorcinol and two acetophenones have been isolated from *K. membranifolia*. In addition, one known vitamin E derivative, alpha-tocopherol quinone has been isolated from *G. contracta*. Lastly, two lactone compounds, including one novel, have been isolated from *G. caudiculata*. Furthermore, potent antibacterial activities have been demonstrated by these compounds for the first time. Spectroscopic analysis of further isolated compounds is being undertaken.




Fig. 1. 2-Hydroxy-6-(12-phenyl dodecyl)benzoic acid isolated for the first time from *Knema membranifolia* (Myristicaceae).

Keywords: ethnopharmacology, antimicrobial resistance, secondary metabolites, x-ray crystallography, NMR

References

- de Sousa Oliveira, K. et al. 2016. Mechanisms of Antibacterial Resistance: Shedding Some Light on These Obscure Processes, Antibiotic Resistance: Mechanisms and New Antimicrobial Approaches, pp. 19–35.
- Murray, C.J. et al. 2022. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis, The Lancet, 399, 10325, 629–655.
- O'Neill, J. 2018. Tackling drug-resistant infections globally: Final report and recommendations,” Straits Times [Preprint].

*Presenting author. Email address: M.Valmiki@uea.ac.uk (M. Valmiki)

S1.P181 Screening for inhibition of β -haematin formation in *Rhaphiostylis beninensis* (Hook. f. ex Planch) Planch ex Benth. (Icacinaeae)

Abiodun Adeyem^{a,*}, Maryam Malmir^b, Kamila Gromek^b, Rita Serrano^b, Jones Moody^a, Olga Silva^b

^a Department of Pharmacognosy, Faculty of Pharmacognosy, Faculty of Pharmacy, University of Ibadan, Ibadan, Nigeria

^b Research Institute for Medicines and Pharmaceutical Sciences (iMed.UL) Pharmacological Sciences Group, Faculty of Pharmacy, University of Lisbon, Av. Prof. Gama Pinto, 1649-019 Lisbon, Portugal

Rhaphiostylis beninensis leaf has demonstrated antimalarial activity in a murine antimalarial model (Oluseje, 2016), but the mode of antimalarial action (important in rational antiplasmodial drug discovery) appears lacking in literature. Therefore, the screening of leaves and roots of *R. beninensis* for inhibition of β -haematin formation is hereby presented. Methanol and 70 % ethanol extracts (25 mg/mL) of each of air-dried, powdered leaves and roots as well as root acidified-water extract (25 mg/mL) of the plant were screened for inhibition of β -haematin formation by modified method of Vargas *et al.*, 2011. Quinine hemi-succinate (10 mg/mL) was used as reference antiplasmodial drug. Activity (Ianalysis, Absorbance Unit, AU) was detected by formation of pink colour and a positive net absorbance value; maximal inhibition was equivalent to absorbance of initial haematin in reaction mixture. The chemical profiles of major groups of secondary metabolites present were also obtained by chromatography (TLC, LC/DAD) followed by quantitative estimation. The hydroethanol extracts of both plant parts had higher Ianalysis than their methanol extracts: 0.173 $\pm$ 0.127/ 0.115 $\pm$ 0.036 AU and 0.121 $\pm$ 0.051/0.091 $\pm$ 0.051 AU for hydroethanol and methanol extracts of leaf/roots, respectively. The acidified water extract had Ianalysis of 0.113 $\pm$ 0.070 AU. The reference drug had Ianalysis, 0.971 $\pm$ 0.141 AU. Terpenoids, phenolics and alkaloids were detected in the parts especially in the hydroethanol extracts. Higher levels of terpenoids and phenolics occurred in the hydroethanol extracts of leaves.

R. beninensis showed inhibition of β -haematin formation in leaves and roots possibly due to the detected secondary metabolites suggesting potential for discovery of inhibitors.

Keywords: *Rhaphiostylis beninensis*, inhibition of β -haematin formation, antimalarial, LC/DAD

References

- Oluseje, O.A. 2016. *In vivo* activity of the leaf extract of *Rhaphiostylis beninensis* (Hook.f.) Planch (Icacinaeae) on *Plasmodium berghei*. B. Pharm. dissertation, Department of Pharmacognosy, Faculty of Pharmacy, University of Ibadan, p. 46.
- Vargas, S., Ioset, K.N., Hay, A.E., Ioset, J.R., Wittlin, S., Hostettmann, K. 2011. Screening medicinal plants for the detection of novel antimalarial products applying the inhibition of β -hematin formation. *Journal of Pharmaceutical and Biomedical Analysis* 56, 880-886.

*Presenting author. Email address: gokeadeyemi11@gmail.com (Abiodun Adeyemi)

S1.P182 New bryostatin derivatives from the North Atlantic Nudibranch *Janolus cristatus*

Laurence K. Jennings^{a,*}, Filippo Spriano^b, Francesco Bertoni^b, Olivier P. Thomas^a

^a School of Biological and Chemical Sciences, University of Galway, University Road, Galway H91 TK33, Ireland

^b Institute of Oncology Research, Faculty of Biomedical Sciences, Bellinzona, Switzerland

Bryostatin-1 is a macrocyclic lactone first isolated from the bryozoan *Bugula neritina* in 1960 (Pettit et al., 1982). It was identified as a potent modulator of protein kinase C (PKC) and in the 1990's quickly progressed through phase I and II clinical trials for cancer treatment (Mohammad et al., 1993). Subsequently, bryostatin-1 has been under investigation for its potential in treating Alzheimer's disease and HIV. However, a significant challenge from the trials of Bryostatin-1 was its limited supply due to low yield within the bryozoan and complexity of its synthesis (Kollár et al., 2014). This underscores the importance of exploring alternative sources of Bryostatins. As part of our ongoing project to isolate bioactive compounds from Irish marine invertebrate, we conducted extensive screening against lymphoma cell lines and identified the nudibranch *Janolus cristatus* exhibiting potent and selective activity. The purification from 1g of *J. cristatus* led to the isolation of the known bryostatin-13 alongside a new C-18 demethylated bryostatin. Further analysis of unpure fractions also revealed the presence of the known bryostatin-11, and two other new C-20 deoxybryostatin derivatives. This study marks the first isolation of bryostatin derivatives from a mollusk. It is widely recognized that nudibranchs typically acquire chemical defence metabolites from their diet, and *J. cristatus* is reported to feed on *B. neritina* (Trindade-Silva et al., 2010). *J. cristatus* was found to exclusively contain C-20 deoxybryostatin derivatives potentially indicating a selective uptake of these specific analogues. Furthermore, *J. cristatus* was found to significantly bioaccumulate and concentrate the bryostatins, as evidenced by the quantity of material isolated.

Keywords: bryostatin, lymphoma, Nudibranch, Ireland

References

- Kollár, P., Rajchard, J., Balounová, Z., Pazourek, J., 2014. Marine natural products: bryostatins in preclinical and clinical studies. *Pharm. Biol.* 52, 237-242.
- Mohammad, R.M., Al-Katib, A., Pettit, G.R., Sensenbrenner, L.L., 1993. Differential effects of bryostatin 1 on human non-hodgkin's B-lymphoma cell lines. *Leuk. Res.* 17, 1-8.
- Pettit, G.R., Herald, C.L., Doubek, D.L., Herald, D.L., Arnold, E., Clardy, J., 1982. Isolation and structure of bryostatin 1. *J. Am. Chem. Soc.* 104, 6846-6848.
- Trindade-Silva, A.E., Lim-Fong, G.E., Sharp, K.H., Haygood, M.G., 2010. Bryostatins: biological context and biotechnological prospects. *Curr. Opin. Biotechnol.* 21, 834-842.

* Presenting author. E-mail address: laurence.jennings@universityofgalway.ie (Laurence Kane Jennings)

S1.P183 Study of the metabolic diversity of Antarctic endophytic fungus *Aspergillus unguis* in relation to different macroalgal hosts and environmental stressors

Márcia G. O. Lopes^{a,c,*}, Pio Colepicolo^b, Paula Carolina Pires Bueno^c, Hosana Maria Deboni^a, Anna Fricke^c

^a Department of Pharmaceutical Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

^b Department of Biochemistry, Chemistry Institute, University of São Paulo, São Paulo, SP, Brazil

^c Department Plant Quality and Food Security, Leibniz Institute of Vegetable and Ornamental Crops, Theodor-Echtermeyer-Weg 1, 14979 Großbeeren, Germany

The extreme environmental conditions of the Antarctic continent formed organisms with unique adaptive abilities and intricate molecular relations. For that reason, the interaction between endophytic fungi and their macroalgal hosts are characterized by the production of unique metabolites (Araújo et al. 2019). To study the diversity of these commercially interesting compounds, the present study focused on the endophytic fungus *Aspergillus unguis*. In a multifactorial approach strain from two different macroalgal hosts, the Rhodophyte *Palmaria decipiens* and the Phaeophyceae *Ascoseira mirabilis*, were subjected to different environmental key factors (e.g., light regime and cultivation media) and subsequent metabolic characterizations were performed by UHPLC-UV-DAD-MS/M. This research sheds light on the diversity and dynamics of metabolites produced by *A. unguis* and might support further targeted metabolite extractions.

Keywords: marine natural products, *Aspergillus unguis*, macroalgal hosts, UVB

References

Araújo, S. D., Silva, L. J., Sartori, A. L. B., Rosa, L. H., Parma, M. M., Queiroz, S. C. N., Barreto-Bergter, E. (2019). Endophytic fungi associated with Antarctic macroalgae: a potential source of bioactivity and new natural products. *Marine drugs*, 17(8), 463.

* Presenting author. E-mail address: marcialopes.qoam@usp.br (Márcia Gracielle de Oliveira Lopes)

S1.P184 Oxidation studies on selected betacyanins from the fruits of *Hylocereus polyrhizus* under influence of ABTS cation radicals

Łukasz Koziol*, Sławomir Wybraniec, Katarzyna Sutor-Świeży

Department C-1, Faculty of Chemical Engineering and Technology, Cracow University of Technology, st. Warszawska 24, 31-155 Krakow, Poland

Betalains, derived from tyrosine, include red-violet betacyanins and yellow-orange betaxanthins. In industry, these pigments can be valued for their potent coloring capabilities owing to their high extinction coefficients. Moreover, they exhibit antioxidant properties due to their ability to scavenge free radicals (Chander et al., 2008). In addition to their industrial utility, betalains demonstrate a spectrum of health benefits, including the induction of phase II enzymes, implicated in cancer protection (Lee et al., 2005), and the inhibition of Intercellular Adhesion Molecule-1 expression (Gentile et al., 2005). Our studies explored dehydrogenation process of selected betacyanins alongside with identifying newly formed derivatives through oxidation. The main objective was to decipher the mechanistic complexities underlying the interaction of cation radicals with betacyanins, thus increasing our understanding of the factors limiting betalain utilization in both food and pharmaceutical sectors. A dehydrogenation study was performed on three pigments present in *Hylocereus polyrhizus* fruits: betanin, as well as its acylated derivatives, phyllocactin and hylocerenin. The isolation of the pigments was conducted using preparative chromatography in a reversed-phase system. The composition of fractions obtained from preparative separations was examined using a mass spectrometer coupled with an HPLC system. The influence of the effect of the degree of the pigment extract purification on the stability of the pigments was performed. The results indicate that degradation of the pigments through reaction with the ABTS cation radicals not only catalyzes the dehydrogenation reactions but also leads to the formation of decarboxylated derivatives.

Acknowledgements: This research was funded by the National Science Centre (NCN, Poland), project no: 2021/41/N/NZ9/03046.

Keywords: *betalains, betacyanins, Hylocereus polyrhizus, dehydrogenation*

References

- Chander, S., Guo, Y. Q., Yang, X. H., Zhang, J., Lu, X. Q., Yan, J. B., Song, T. M., Rocheford, T. R., & Li, J. S. 2008. Using molecular markers to identify two major loci controlling carotenoid contents in maize grain. *Theoretical and Applied Genetics*, 116(2), 223–233.
- Gentile, C., Tesoriere, L., Allegra, M., Livrea, M. A., D'Alessio, P. Antioxidant betalains from cactus pear (*Opuntia ficus-indica*) inhibit endothelial ICAM-1 expression. *Ann. N. Y. Acad. Sci.* 2004, 1028, 481– 486
- Lee, C.-H., Wettasinghe, M., Bolling, B. W., Ji, L.-L., Parkin, K. L. 2005. Betalains, phase II enzyme-inducing components from red beetroot (*Beta vulgaris* L.) extracts. *Nutrition and Cancer*, 53(1), 91–103

* Presenting author. E-mail address: lukasz.koziol@doktorant.pk.edu.pl (Łukasz Koziol)

S1.P185 Identification of novel betaxanthin pigments in jewel-like fruits of *Talinum paniculatum* (Jact.) Gaertn

Renata Górská*, Katarzyna Sutor-Świeży, Sławomir Wybraniec

Department of Chemical Technology and Environmental Analysis, Faculty of Chemical Engineering and Technology, Cracow University of Technology, Warszawska 24, 31-155 Cracow, Poland

Betaxanthins belong to the unique pigments occurring in Caryophyllales order plants providing them yellowish color. Due to their health-promoting properties, natural sources of these pigments are continually sought (Kumorkiewicz-Jamro et al. 2021). Unlike betacyanins, betaxanthin pigments can be readily synthesized by coupling selected amines or amino acids to betalamic acid (Schliemann, et al., 1999). Previous studies have demonstrated that a plant of the *Talinum* genus – *Talinum triangulare* (Jacq.) Willd – produces dopamine-betaxanthin and tyrosine-betaxanthin (Swarna et al., 2013). However, *Talinum paniculatum* has not yet been studied for its betaxanthin profile. The plant belongs to a group of herbs used in folk medicine. It is distinguished by its succulent leaves, purplish flowers and yellowish fruits located on long stems (Tolouei et al., 2019). This study determined betaxanthin profile for fruits of *T. paniculatum* using liquid chromatography coupled with mass spectrometry. The results of the analyses indicated the presence of two novel, previously unknown betaxanthin derivatives with m/z ratio of 361. These compounds were identified as synephrine- betaxanthin and phenylephrine-betaxanthin. Their presence in the extract was confirmed based on the synthesized standards formed by coupling respective molecules with betalamic acid produced during betanin hydrolysis. These pigments were purified using flash chromatography and preparative high-performance liquid chromatography. The structures were confirmed by high-resolution mass spectrometry (LC-Q-Orbitrap-MS) and NMR spectroscopy. The finding of novel betaxanthin pigments in herbaceous plants suggests a promising direction for future research, including investigating their potential bioactive properties. Furthermore, this represents promising prospect for using betaxanthins for food fortification.

Keywords: *Talinum paniculatum*, betaxanthins, mass spectrometry, chromatography

References

- Kumorkiewicz-Jamro, A., Świergosz, T., Sutor, K., Spórna-Kucab, A., Wybraniec, S., 2021. Multi-colored shades of betalains: Recent advances in betacyanin chemistry. *Nat. Prod. Rep.* 38, 2315–2346.
- Schliemann, W., Kobayashi N., Strack D., 1999. The Decisive Step in Betaxanthin Biosynthesis Is a Spontaneous Reaction. *Plant Physiol.* 119, 1217–1232.
- Swarna, J., Lokeswari, T. S., Smita, M., Ravindhran, R., 2013. Characterisation and Determination of in Vitro Antioxidant Potential of Betalains from *Talinum Triangulare* (Jacq.) Willd. *Food Chem.* 141, 4382–4390.
- Tolouei, S. E. L., Palozi, R. A. C., Tirloni, C. A. S., Marques, A. A. M., Schaedler, M. I., Guarnier, L. P., Silva, A. O., de Almeida, V. P., Manfron Budel, J., Souza, R. I. C., dos Santos, A. C., Silva, D. B., Lourenço, E. L. B., Dalsenter, P. R., Gasparotto Junior, A., 2019. Ethnopharmacological Approaches to *Talinum Paniculatum* (Jacq.) Gaertn. – Exploring Cardiorenal Effects from the Brazilian Cerrado. *J. Ethnopharmacol.* 238, 111873.

* Presenting author. *E-mail address:* renata.gorska@doktorant.pk.edu.pl (Renata Górská)

S1.P187 Evaluation of a standardized *Melissa officinalis* extract anti-obesity effects in high fat high sugar diet-induced obesity in rat

Anne-Sophie Roy^a, Thaïs Vorgers^{b,c}, Sekhou Cisse^c, Amine Benarbiab^{b,c}, Yassine Mallema^a

^a Nutrition, PathoPhysiology and Pharmacology (NP3) Unit, Oniris, 101 route de Gachet, Nantes 44300, France

^b Nor-Feed SAS, 3 rue Amedeo Avogadro, Beaucouzé 49070, France

^c FeedInTech Labcom, 42 rue Georges Morel, Beaucouzé 49070, France

Many studies suggest the use of *Melissa officinalis* extracts (MOE) to decrease anxiety. However, their anti-obesity effects remain insufficiently documented. In this context, the anti-obesity effect of a standardized MOE was evaluated in an induced obesity rat model, and compared to one of its main active compound: Rosmarinic acid (RA). Thirty-two 9-week-old Wistar rats were divided into 4 groups. A control group fed with a standard diet (ctSTD) and 3 experimental groups fed with High Fat High Sugar Diet (HFHSD) and respectively supplemented with placebo (ctHFHSD, 200 mg/kg), MOE (MOE200, 200 mg/kg) and RA (RA10.8, 10.8 mg/kg). The trial duration was 12 weeks and several measures linked to obesity such as body weight, adiposity index, abdominal circumference, glucose tolerance, and Total Antioxidant Status (TAS) from rats' livers were monitored at the end of the trial. Results showed that HFHSD significantly increased Body weight, adiposity Index, abdominal Circumference, and reduced glucose tolerance compared to ctSTD group ($p < 0.05$ to $p < 0.005$). These effects were attenuated by MOE and RA supplementations. As expected, the TAS of the ctHFHSD group was also significantly lower than the TAS of the ctSTD group. Nevertheless, surprisingly, TAS of rats supplemented with MOE and RA was also lower than the TAS of ctHFHSD group. These results highlight the advantage of MOE and RA supplementation to reduce visceral obesity. However, their anti-obesity effect was not correlated with TAS increase, suggesting that their metabolic benefit involves antioxidant-independent mechanisms. Further studies are underway to confirm this hypothesis.

Keywords: *Melissa officinalis*, rosmarinic acid, obesity, rats

* Presenting author. E-mail address: yassine.mallem@oniris-nantes.fr (Yassine Mallem)

S1.P188 Biotransformation and antimicrobial studies of flavonoids with bromine and chlorine atoms

Martyna Perz^{a,*}, Daria Szymanowska^{b,c}, Agnieszka Krawczyk-Łebek^a, Monika Dymarska^a, Tomasz Janeczko^a, Edyta Kostrzewa-Susłow^a

^a Department of Food Chemistry and Biocatalysis, Faculty of Biotechnology and Food Science, Wrocław University of Environmental and Life Sciences, 50-375 Wrocław, Poland

^b Department of Biotechnology and Food Microbiology, Faculty of Food Science and Nutrition, Poznań University of Life Sciences

^c Department of Pharmacognosy and Biomaterials, Faculty of Pharmacy, Poznań University of Medical Sciences, 60-806 Poznań, Poland

Flavonoids are known for their anti-inflammatory properties, antioxidant, and antimicrobial ones. However, due to their low water solubility, orally administered flavonoids are poorly absorbed, and therefore, their therapeutic potential is limited (Karak et al., 2019; Lewin et al., 2013). It can be improved using glycosylation via biotransformation, which has been proven to be an effective way to increase water solubility and bioavailability of flavonoids (Yang et al., 2016; Hofer, 2016). In our study, we carried out the biotransformation of chalcone, flavanone, and flavone, which contained bromine and chlorine atoms, using the strains *Beauveria bassiana* KCH J1.5, *Isaria fumosorosea* KCH J2 and *Isaria farinose* KCH J2.6. Then, we checked the antimicrobial activities of aglycones and one glycoside 8-bromo-6-chloroflavone 4'-O-β-D-(4"-O-methyl)-glucopyranoside on four pathogenic and four probiotic bacteria. We measured the effect of flavonoids on bacteria growth over a 72-hour period at hourly intervals using a microplate reader. The chalcone exhibited highest antibacterial properties, followed by flavones and then flavanones. We also noticed that the glycoside derivative had enhanced antimicrobial effect against pathogenic bacteria, while being less harmful to probiotic bacteria than the corresponding aglycone. Our research suggests that bromine and chlorine atoms in flavonoids increase their antimicrobial properties, and that attachment of a glycoside unit to a flavone selectively enhances its bactericidal properties and bioavailability of the compound. This study adds to the knowledge about compounds not previously described in the literature with their antibacterial properties and further potential for their use in industry (Perz et al., 2023).

Keywords: flavonoids, biotransformations, entomopathogenic filamentous fungi, antimicrobial activity

References

- Karak, P., 2019. Biological Activities of Flavonoids: An Overview. *Int. J. Pharm. Sci. Res.* 10, 1567–1574.
- Lewin, G.; MacLuk, A.; Moncomble, A.; Cornard, J. P., 2013. Enhancement of the Water Solubility of Flavone Glycosides by Disruption of Molecular Planarity of the Aglycone Moiety. *J. Nat. Prod.* 76, 8–12.
- Yang, B.; Liu, H.; Yang, J.; Gupta, V. K.; Jiang, Y., 2018. New Insights on Bioactivities and Biosynthesis of Flavonoid Glycosides. *Trends Food Sci. Technol.* 79, 116–124.
- Hofer, B., 2016. Recent Developments in the Enzymatic O-Glycosylation of Flavonoids. *Appl. Microbiol. Biotechnol.* 2016, 100, 4269–4281.
- Perz, M.; Krawczyk-Łebek, A.; Kostrzewa-Susłow, E.; Dymarska, M.; Janeczko, T., 2023. Biotransformation of Flavonoids with -NO₂, -CH₃ Groups and -Br, -Cl Atoms by Entomopathogenic Filamentous Fungi. *Int. J. Mol. Sci.* 24, 9500.

* Presenting author. E-mail address: martyna.perz@upwr.edu.pl (Martyna Perz)

S1.P189 *In situ* approaches to marine and terrestrial metabolomics

Daniel Jordan Otto^{a,*}, Morgane Mauduit^b, Scott A. Jarmusch^b, Ling Ding^b, Aaron John Christian Andersen^b

^a CeMiSt – Center for Microbial Secondary Metabolites, Section for Microbial and Chemical Ecology, Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kgs Lyngby, Denmark

^b Section for Microbial and Chemical Ecology, Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kgs Lyngby, Denmark

This poster will cover *in situ* chemical detection of secondary metabolites in marine environments. *In situ* metabolomics allows for more unbiased studies of complex microbial systems, natural metabolite expression and interactions between microbial communities than what is afforded by traditional laboratory culturing of microbes (Tomm et al., 2019). Historically *in situ* detection has been difficult, but with advances in both instruments, capture methods, and metabolomic approaches it is a promising avenue for answering basic research questions about the microbial world, as well as identifying novel and useful compounds for real world applications. Marine-based *in situ* systems have been studied utilizing custom designed devices for *in situ* transcriptomic and meta-genomic studies (Bech et al., 2024). These devices allow for the colonization of marine bacteria and formation of biofilms after placement in a natural harbor environment. Combining exo-metabolite capture via pumping of surrounding seawater through HP-20 resin, metabolomics of resulting biofilms and captured exo-metabolites, mass spectrometry imaging, transcriptomics and meta-genomics, key secondary metabolites can be identified and linked to the different life stages and organisms present during microbial colonization in marine environments. Additionally, these key metabolites are likely to be of great interest due to wide varieties of possible bioactivity associated with their ecological roles as marine microorganisms (Li et al., 2023).

Keywords: *in situ*, metabolomics, marine microbes, multi-omics, chemical ecology

References

- Bech, P.K., Jarmusch, S.A., Rasmussen JA, Limborg MT, Gram L, Henriksen NNSE. 2024. Succession of microbial community composition and secondary metabolism during marine biofilm development. ISME Commun. 2024 Jan 20;4(1):ycae006.
- Li X, Xu H, Li Y, Liao S, Liu Y. 2023. Exploring Diverse Bioactive Secondary Metabolites from Marine Microorganisms Using Co-Culture Strategy. Molecules. 2023; 28(17):6371.
- Tomm HA, Ucciferri L, Ross AC. 2019. Advances in microbial culturing conditions to activate silent biosynthetic gene clusters for novel metabolite production. J Ind Microbiol Biotechnol. 2019 Oct;46(9-10):1381-1400.

* Presenting author. E-mail address: dajoot@dtu.dk (Daniel Jordan Otto)

S1.P190 Correlation between saponin content and biological activities of *Beta vulgaris* L.

Anna Tekieli^{a,*}, Aneta Spórna-Kucab^a, Agnieszka Grzegorzczak^b, Łukasz Świątek^c

^a Department of Chemical Technology and Environmental Analytics, Faculty of Chemical Engineering and Technology, Cracow University of Technology, Warszawska 24, 31-155 Krakow, Poland

^b Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Medical University of Lublin, Chodźki 1, 20-093 Lublin, Poland

^c Department of Virology with SARS Laboratory, Faculty of Medicine, Medical University of Lublin, Chodźki 1, 20-093 Lublin, Poland

Beta vulgaris L. represents functional food due to the presence of diverse bioactive compounds such as saponins, betalains, polyphenols, and minerals (Thiruvengadam et al., 2024). Saponins, which are natural glycosides, have exhibited varied pharmacological activities in clinical trials, effectively treating conditions such as type II diabetes, pancytopenia, and aplastic anemia (Galas et al., 2023). Triterpenoid saponins, notably prevalent in beetroot, are complex molecules composed of varied glycone and aglycone structures (Mikołajczyk-Bator et al., 2016). The diversity in structures and the proven pharmacological activity of certain saponins underscore significance of beetroot as a promising source of these compounds. The aim of the study is to investigate the correlation between saponin content in both the flesh and peel of five *B. vulgaris* varieties (Ceryl, Chrobry, Forono, Tytus, and Boldor) and their respective anticancer and antifungal activities. Using HPLC-ESI-MS/MS, 24 saponins with oleanolic acid, hederagenin, akebonoic acid, and gypsogenin as aglycones were identified. The Boldor cultivar (peel) exhibited the highest saponin content at 20.8 g/kg fresh weight. Notably, Tytus (peel) displayed the strongest antifungal activity against the majority of tested fungi (MIC = 0.125–0.5 mg/mL), and showed low cytotoxicity towards non-cancerous cells (CC50 = 405 µg/mL). Conversely, Tytus (flesh) exhibited high cytotoxicity against human cervical adenocarcinoma (HeLa) cells, with a CC50 of 282 µg/mL. Most saponins influenced the anticancer activity of beets against colon cancer cell line (RKO) and HeLa. Furthermore, betavulgarosides V and VIII, as well as compounds with *m/z* = 779 and 631, demonstrated effects on all tested fungi strains.

This research was funded in whole by National Science Centre, Poland for the years 2024–2027 (Project No. UMO-2023/49/N/NZ9/01706).

Keywords: *Beta vulgaris* L., saponins, anticancer, antifungal, LC-MS/MS

References

- Thiruvengadam, M., Chung, I. M., Samynathan, R., Chandar, S. H., Venkidasamy, B., Sarkar, T., Rebezov, M., Gorelik, O., Shariati, M., Simal-Gandara, J., 2024. A comprehensive review of beetroot (*Beta vulgaris* L.) bioactive components in the food and pharmaceutical industries. Crit. Rev. Food Sci. Nutr. 64, 708-739.
- Galas, A., Niemiec, R., Matysek, M., Olszanicka, A., Kołodziejczyk, K., Hop, I., Branewska, B., Imioło, J., Maciąg, A., 2023. Multidirectional pharmacological effects of saponins in light of clinical trials: a systematic review of literature. J. Educ. Health Sport. 39, 83-97
- Mikołajczyk-Bator, K., Błaszczak, A., Czyżniewski, M., Kachlicki, P., 2016. Characterisation and identification of triterpene saponins in the roots of red beets (*Beta vulgaris* L.) using two HPLC-MS systems. Food Chem. 192, 979-990.

* Presenting author. E-mail address: anna.tekieli@doktorant.pk.edu.pl (Anna Tekieli)

Natural products in the era of emerging infectious diseases

S1.P191 A metabolomic approach in assessing the anti-biofilm potentials of *F. proliferatum*-a fungal endophyte isolated from an indigenous Nigerian medicinal plant

Elizabeth Nwagwu*, Dinesh Raganathan, Tabassum Ara, Rothwell Tate, Louise Young, RuAngelie Edrad-Ebel

Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, The John Arbuthnott Building, 161 Cathedral Street, Glasgow G4 0RE, UK

Our research is aimed at isolating fungal endophytes from indigenous Nigerian medicinal plants and bio-prospecting their potential for the treatment of biofilm-forming methicillin-resistant *Staphylococcus aureus*. The fungal endophyte BLS1 was isolated from the stem of *Vernonia amygdalina*, an indigenous Nigerian medicinal plant used in the treatment of various ailments (Asante et al., 2016, Asante et al., 2019). DNA sequencing identified BLS1 as *Fusarium proliferatum* (Altschul, 1990).

Crude extract of BLS1, initially grown on malt agar, exhibited 96.3% antimicrobial activity, 99.8% biofilm prevention, and 67.3% biofilm eradication at concentration of 100 µg/ml. Culture optimization on rice, oat, malt extract and potato dextrose broth incubated at 7, 15, and 30 days determined the most suitable conditions to scale-up BLS1 on rice media for 15 days at highest extract yield with 89.5% antimicrobial activity while preventing biofilm formation at 100%. The scaled-up extract was fractionated by normal-phase flash chromatography using hexane and ethyl acetate. The polar fractions exhibited almost 100% antimicrobial and antibiofilm activity, while multivariate OPLS-DA of their NMR spectral data showed chemical shifts of 0-7 ppm representing fatty acids, acetylated compounds, hydroxylated aliphatics, and phenolics. Dereplication of the LC-HRMS data indicated that the antimicrobial- and antibiofilm-active fractions afforded known *Fusarium* chemotaxonomic markers, fusarin A and fusaric acid (Proctor et al., 2009).

Keywords: *Fusarium proliferatum*, antimicrobial, antibiofilm, fungal endophyte, MRSA

References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. *Mol. Biology*. 215 (3), 403–410.
- Asante, D.B., Effah-Yeboah, E., Barnes, P., Abban, H.A., Ameyaw, E.O., Boampong, J.N., Ofori, E.G., & Dadzie, J.B., 2016. Antidiabetic Effect of Young and Old Ethanolic Leaf Extracts of *Vernonia amygdalina*: A Comparative Study. *J. Diabetes Res*.
- Asante, D.B., Henneh, I.T., Acheampong, D.O., Kyei, F., Adokoh, C.K., Ofori, E.G., Domey, N.K., Adakudugu, E., Tangella, L.P., & Ameyaw, E.O., 2019. Anti-inflammatory, anti-nociceptive and antipyretic activity of young and old leaves of *Vernonia amygdalina*. *Biomed. Pharmacother*. 111, 1187–1203.
- Proctor, R.H., McCormick, S.P., Alexander, N.J., & Desjardins, A.E., 2009. Evidence that a secondary metabolic biosynthetic gene cluster has grown by gene relocation during evolution of the filamentous fungus *Fusarium*. *Mol. Microbiol*. 74(5), 1128–1142.

* Presenting author. E-mail address: elizabeth.nwagwu@strath.ac.uk (Elizabeth Nwagwu)

S1.P192 Flavonoids content, anti-inflammatory effect and antioxidant activities methanol leaf extract of *Funtumia africana* (Benth)

Bamidele Martin Amos-Tautua^{*}, Precious Ezougha Oyaseiye

Department of Chemical Sciences, Niger Delta University, Wilberforce Island, Amassoma, Nigeria

Funtumia africana (Apocynacea) a versatile plant in the Niger Delta region of Nigeria, have been used to treat various diseases, such as inflammation, diabetes and bacterial infections. Likewise flavonoids have received great attention due to the health benefits they offer varying from lowering blood pressure, inhibiting pro-inflammatory cytokines and protecting against oxidative stress (Al- Khayri et al., 2022).

This study was conducted to appraise the antioxidants activities and anti-inflammatory effect of aqueous methanol leaf extract of *F. africana*. The leaf extract was assessed for total phenol, flavonoids, total antioxidant capacity and free radical-scavenging activities. HPLC profiling of the plant sample was carried out for quantification of flavonoids. Acute anti-inflammatory activity was determined using the formalin induced paw edema method.

The antioxidant results show that *F. africana* has strong antioxidant activities (Table 1). HPLC fingerprinting gave kaempferol (154mg/100g), (+) – catechin (36.55 mg/100 g), naringin (35.90 mg/100 g), quercetin (35.67 mg/100 g), luteolin (31.26 mg/100 g) and apigenin (16.46 mg/100g) as the major flavonoids present in the sample. *F. africana* extract demonstrated significant anti- inflammatory effects in a dose-dependent manner. It showed significant reduction of the paw edema size in Wistar rats at 500 mg/kg dose compared to the standard drug (aspirin). The antioxidant and anti-inflammatory activities could be attributed to the high content of flavonoids and phenolics in *F. africana* leaves.

The presence of these phytochemicals might justify the use of the plant in the treatment of inflammation and other degenerate diseases by traditional practitioners in Niger Delta region of Nigeria.

Keywords: *Funtumia africana*, anti-inflammation, antioxidant, flavonoids

References

Al-Khayri, J.M., Sahana, G.R., Nagella, P., Joseph, B.V., Alessa, F.M., Al-Mssallem, M.Q. 2022. Flavonoids as Potential Anti-Inflammatory Molecules. A Review. *Molecules*, 27, 2901.

* Presenting author. *E-mail address:* bamideleamos-tautua@ndu.edu.ng (Bamidele Martin Amos-Tautua)

S1.P193 Demystifying Bangladeshi ethnomedicines with modern approaches: *Congea tomentosa* as a source of potential anti-microbial and anti-cancer agents

Mohammad Omar Faruque^{*a}, Shaikh Bokhtear Uddin^a, Hu Xuebo^b

^a Department of Botany, University of Chittagong, Chittagong-4331, Bangladesh

^b College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, China

The traditional healers of indigenous communities of Bangladesh use several plant parts, with often unknown ingredients, for the treatment of cancers and microbial diseases. To investigate such treatments, we documented the traditional use of 276 ethnomedicinal plant species, utilising 292 informants of Bangladesh. Notably, 28 species were reported with new therapeutic uses and 13 species described have never been studied. Of these, *Congea tomentosa* was selected for *in-vitro* studies. Bioassay-guided screening led us to identify 10 compounds, and 5 compounds were isolated and identified. Among the isolated compounds, compound-2 (Stigmasterol) displayed promising antimicrobial activity against tested microorganisms, while compound-4 (5,6-Dimethoxy-2-methyl- 1H-benzoimidazole) significantly decreased the ratio of Bcl-2/Bax and increased the expression levels of cleaved caspases-9 and -3, suggested that compound-4 had the potential to induce apoptosis in U-251 cells, through activation of the intrinsic/mitochondrial pathway which might be triggered by the inhibition of Stat3 and Akt expression. Despite the ethnobotanical importance of *Congea tomentosa*, no chemical or biological studies have been published to date which support its traditional applications in Bangladesh. However, our study demonstrates the possible use of this plant in the treatment of cancers/microbial diseases, and support its traditional use.

Keywords: ethnobotany, *Congea tomentosa*, cytotoxicity, antimicrobial activity, apoptosis

* Presenting author. E-mail address: omf@cu.ac.bd (Dr. Mohammad Omar Faruque)

S1.P194 In silico testing of flavonoids of the genus *Mentha* (Lamiaceae) as potential inhibitors of NS3- protease domains of dengue viruses

Marianna Ines Lopez

International School Manila, University Parkway, Fort Bonifacio Global City, Taguig City 1634, Philippines

Dengue fever is a vector-borne disease with 4 serotypes. From 2000-2019, the WHO documented a ten-fold surge in reported cases worldwide, as cases skyrocketed from 500,000 to 5.2 million. In 2023, 80 countries/territories cumulatively reported over 5 million dengue cases and over 5,000 dengue-related deaths. Clinical symptoms of the virus include severe headache, muscle and joint pain, nausea, vomiting, swollen glands, rash, and high fever. Currently, only one vaccine exists for the virus as a preventative measure, but according to the CDC, no antivirals have been identified to be used for treatment. In the present work, the flavonoids of medicinal plants (Eftekhar et al., 2021) known to block the virus infection in vitro were explored against the DENV NS3-protease of 4 serotypes. UCSF Chimera and AutodockVina were utilized for docking and optimal binding visualization to obtain the binding affinity scores for each ligand and DENV protein. Finally, the best ligands were evaluated by molecular dynamic simulations through GROMACS software.




Fig 1. Structure of luteolin-7-*O*-rutinoside (LNR26)




Fig 2. Structure of eriocitrin (ETN13)

Molecular docking, binding affinity scores suggested that the luteolin-7-*O*-rutinoside (**Fig 1.**) and eriocitrin (**Fig 2.**) compounds are able to tightly bind to NS3-pro. These molecules were then subjected to molecular dynamics to determine and visualize how they would bind to the DENV proteases.

Results indicate that LNR26 and ETN13 could block NS3-pro activity in DENV. For further research, follow-up in vitro experiments can be conducted to ensure the integrity of results, which can be later utilized for drug development.

Keywords: Dengue virus, NS3-protease, Molecular docking, *Mentha* sp., flavonoids

References

- Aziz Eftekhar, Ameer Khusro, Elham Ahmadian, Solmaz Maleki Dizaj, Amir Hasanzadeh, Magali Cucchiari, Phytochemical and nutraceutical attributes of *Mentha* spp.: A comprehensive review, *Arabian Journal of Chemistry*, Volume 14, Issue 5, 2021, 103106, ISSN 1878-5352,
- Cruz-Arreola O, Orduña-Díaz A, Domínguez F, Reyes-Leyva J, Vallejo-Ruiz V, Domínguez-Ramírez L, Santos-López
- G. *In silico* testing of flavonoids as potential inhibitors of protease and helicase domains of dengue and Zika viruses. *PeerJ*. 2022 Aug 4;10:e13650. doi: 10.7717/peerj.13650. PMID: 35945938; PMCID: PMC9357371.
- "Dengue - Global Situation" World Health Organization. 21 December 2023, <https://www.who.int/emergencies/disease-outbreak-news/item/2023>

*Presenting author. E-mail address: lopezmar@ismanila.org (Marianna Ines Lopez)

S1.P195 Application of feature based molecular network analysis for the investigation of the Covid-19 remedy Hanshiyi

Stefanie Tiefenbacher^{a,*}, Eva-Maria Pferschy-Wenzig^a, Yanyan Zhou^b, Min Li^c, Xiaolin Tong^c, Rudolf Bauer^a

^a Institute of Pharmaceutical Sciences, Division of Pharmacognosy, University of Graz, 8010 Graz, Austria

^b Institute of Chinese Materia Medica, China Academy of Chinese Medical Sciences, Dongcheng District, 100010 Beijing, China

^c China Academy of Chinese Medical Sciences, Guang'anmen Hospital, Department of Endocrinology, Xicheng District, 100053 Beijing, China

In the nascent phase of the Covid-19 pandemic, the Traditional Chinese Medicine (TCM) formula known as Hanshiyi (HSYF) was developed with the aim of mitigating the risk of severe disease progression among patients (Tian et al., 2020). Comprising primarily botanical ingredients, HSYF consists of 20 components. Previous investigations have detailed the chemical composition through the application of analytical techniques such as High-Performance Thin-Layer Chromatography (HPTLC) and Liquid Chromatography-Mass Spectrometry (LC-MS) (Tiefenbacher et al., 2022), along with the exploration of its pharmacological properties (Tiefenbacher et al., 2023).

This study introduces the utilization of feature-based molecular network analysis to delve into the constituents of HSYF. Different decoctions, were prepared and subjected to LC-MS analysis using an Ultimate 3000 RS system coupled with a Q Exactive Orbitrap Mass Spectrometer. Subsequently, the acquired data underwent conversion; alignment and were then exported to the Global Natural Products Social Molecular Networking (GNPS) platform (Aron et al., 2020). Employing Cytoscape, the resulting visualizations revealed 1536 nodes and 2533 edges in the positive mode, and 2456 nodes and 3897 edges in the negative mode. For the validation of the proposed compounds, 108 reference substances were utilized, with approximately 60 confirmed to be constituents of the mixture. Consequently, feature-based molecular network analysis emerges as a contemporary and valuable approach, amalgamating traditional analytical methods with innovative database-driven methodologies.

Keywords: Hanshiyi, TCM, Covid-19, feature based molecular network, LC-MS

References

- Aron AT, Gentry E, McPhail KL, Nothias LF, Nothias-Esposito M, Bouslimani A, Petras D, Gauglitz JM, Sikora N, Vargas F, van der J. Hoof J, Ernst M, Kang KB, Aceves CM, Caraballo-Rodríguez AM, Koester I, Weldon KC, Bertrand S, Roullier C, Sun K, Tehan RM, Boya CA, Martin H. C, Gutiérrez M, Ulloa AM, Mora JAT, Mojica-Flores R, Lakey- Beitia J, Vásquez-Chaves V, I. Calderón A, Tayler N, Keyzers RA, Tugizimana F, Ndlovu N, Aksenov AA, Jarmusch AK, Schmid R, Truman AW, Bandeira N, Wang M, Dorrestein P, 2020. Reproducible Molecular Networking of Untargeted Mass Spectrometry Data Using GNPS. *Nat Protoc* 15
- Tiefenbacher S, Schwarz P, Perhal A, Pirker T, Pferschy-Wenzig EM, Yao C, Tian C, Wang W, Zhou Y, Liu H, Li M, Dirsch V, Tong X, Bauer R, 2023. Pharmacological profiling of Hanshiyi formula, a herbal mixture used for the treatment of COVID-19 in China. *Planta Med* 89: 1422.
- Tian J, Yan S, Wang H, Zhang Y, Zheng Y, Wu H, Li X, Gao Z, Ai Y, Gou X, Zhang L, He L, Lian F, Liu B, Tong X, 2020. Hanshiyi Formula, a medicine for Sars-CoV2 infection in China, reduced the proportion of mild and moderate COVID-19 patients turning to severe status: A cohort study. *Pharmacological research* 161: 105127
- Tiefenbacher S, Zhou Y, Whang W, Chao Y, Li M, Tong X, Bauer R, 2022. Development of HPTLC and UPLC methods for quality control of Hanshiyi formula. *Planta Med* 88(15): 1459.

*Presenting author. E-mail address: stefanie.tiefenbacher@uni-graz.at (Stefanie Tiefenbacher)

S1.P196 Pathophysiological assessment of trypanocidal activities of ethanolic leaf extract and fractions of *Azadiractha indica* on infected experimental model

Nnodim, Nnedinma Nancy^{a*}, Okafor, Fabian^b, Ebiziem, Chukwuma Joseph^a, Samuel, Monday Dadi^a

^a Parasitology Unit, Department of Biological Science, Federal University of Technology, Owerri, Imo State, Nigeria

^b Department of Zoology and Environmental Biology, University of Nigeria, Nsukka, Enugu State, Nigeria

Trypanosomiasis is a vector borne parasitic infection caused by genus Trypanosome. This infection precipitates increased red blood cell destruction that predispose the host to anemia and tissue damage (Sirak, *et al.*, 2024;). The pathophysiological evaluation of the trypanocidal activity of ethanolic leaf extract and fractions of *Azadiractha indica* was studied. Phytochemical and toxicity studies were done using standard procedures (Mikail, 2009). A total number of 25 rats (5 groups (A - E) of 5 rats) were used for the study. All the groups were induced with 0.1ml of Trypanosome inoculum containing 10^6 of Trypanosome except the positive control. Groups A and B were treated with 400mg/kg and 600mg/kg (b.wt.) respectively for 5 days after the establishment of parasitemia. Groups C, D and E served as positive and negative controls. At the end of therapeutic study, the organs (liver and kidney) were harvested for histopathological studies.

The phytochemical screening revealed the presence of flavonoids, terpenoids, tannis, alkaloids, reducing sugar, phenols and saponins. The oral medial lethal dose (LD₅₀) of the extract was 3162mg/kg. Treatment with 600 mg/kg b.wt. cleared the parasite from circulating blood on day 9 post treatment (pt) similar to the positive control group. Out of the four fractions tested, Ethyl acetate fraction showed the highest activity compared to other test groups. This study established that the crude extract and fractions of *A. indica* were active against *T. brucei brucei* sub specie which has not been reported. Histopathological studies confirmed no infiltration of the intracellular cells of the organs examined. Therefore treatment with *A. indica* may not cause any serious toxic effects on the biological system of the host if utilized as a potential treatment for trypanosomiasis. Characterization of the most active fraction is ongoing.

Keywords: *Trypanosoma brucei brucei*, *Azadiractha indica*, phytotherapeutic, histopathology and albino rats

References:

Mikail, H. G., 2009. In vitro trypanocidal effect of methanolic extract of *Sclerocarya birrea*, *Commiphora kerstingii* and *Khaya senegalensis*. *African journal of Biotechnology*, 8:2047-2049

Sirak, B., Bizuneh, G. K., Imming, 2024. In vitro and In vivo antitrypanosomal activity of the fresh leaves of *Ranunculus Multifidus forsk* and its major compound anemonin against *Trypanosoma congolense* field isolate. *BMC veterinary research* 20: 32.

*Presenting author. E-mail address: ihechukwu03@gmail.com (Nnodim, Nnedinma Nancy)

S1.P197 Antioxidant activity and Structural-Activity Relationship (SAR) study of stilbenoids isolated from the stem bark of *Anisoptera laevis* (Dipterocarpaceae)

Noorazlina Adnan^{a,b,c}, Aisyah Salihah Kamarozaman^{a,b,*}, Siti Zakirah Azahar^{b,c}, Nik Fatini Nik Azmin^c, Muhammad Sulaiman Mohd Johari^{b,c}, Norizan Ahmat^a

^a Centre of Foundation Studies, Universiti Teknologi MARA, Cawangan Selangor, Kampus Dengkil, 43800 Dengkil, Selangor, Malaysia

^b Atta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA, Cawangan Selangor, Kampus Puncak Alam, 42300 Bandar Puncak Alam, Selangor, Malaysia

^c Faculty of Applied Sciences, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

Anisoptera laevis, a genus in the family of Dipterocarpaceae, is well-known to contain abundant sources of stilbenoids, exhibited a wide range of biological activities (Duta-Bratu et al., 2023). This study revealed antioxidant activities and structural-activity relationships of stilbenoids isolated from *A. laevis*. The stem bark of *A. laevis* (900g) was collected from Pahang, Malaysia and macerated in acetone for 24 hours, concentrated in vacuo to yield crude extract (65g). Fractionation using VLC, *n*-hexane:ethyl acetate as eluent gave eight fractions (AL1-8). Piceid (**1**) (4.0mg) and resveratrol-12-C- β -D-glucopyranoside (**2**) (1.9mg), were obtained by purifying AL5 (1.73g) via column chromatography (CC). Vaticanol B (**3**) (204mg) was obtained from AL4 (1g) using CC, solvent- system chloroform:methanol. The antioxidant activities of isolated stilbenoids were assessed using DPPH, ABTS and hydroxyl radicals scavenging.




Fig. 1. Structures of compounds 1-3

Compounds (**1-3**) (Fig.1) demonstrated significant efficacy in scavenging DPPH radicals, as evidenced by their respective IC_{50} values of 69.00, 50.58 and 66.34 μ g/mL. Compounds **2** and **3** exhibited very strong activity against ABTS radicals (IC_{50} =16.44, 33.43 μ g/mL, respectively), while compound **1** demonstrated strong activity at 63.57 μ g/mL. Compounds **1** and **2** exhibited potent antioxidant activity in scavenging hydroxyl radicals (IC_{50} =20.06, 17.08 μ g/mL, respectively), while compound **3** demonstrated weak antioxidant effect (IC_{50} =248.5 μ g/mL). The most potent antioxidant activity was observed in Compound **2**. This results indicated that the position of a sugar moiety (Xie et al., 2022) and the degree of polymerisation are the factors contributing the antioxidant activity. We would like to express our appreciation to UiTM, Malaysia for the financial support [600-UITMSEL (PI. 5/4) (127/2022)]

Keywords: *Anisoptera laevis*, stilbenoids, antioxidant activity, structural-activity relationship

References

- Duta-Bratu, C.-G., Nitulescu, G.M., Mihai, D.P., Olaru, O.T., 2023. Resveratrol and Other Natural Oligomeric Stilbenoid Compounds and Their Therapeutic Applications. *Plants*, 12, 2935.
- Xie, L., Deng, Z., Zhang, J., Dong, H., Wang, W., Xing, B., Liu, X., 2022. Comparison of Flavonoid O-Glycoside, C- Glycoside and Their Aglycones on Antioxidant Capacity and Metabolism during In Vitro Digestion and In Vivo. *Foods*, 11, 882.

* Presenting author. E-mail address: nooraz2753@uitm.edu.my (Noorazlina Adnan)

S1.P198 Resveratrol dimer and tetramers from the isolation of *Shorea pauciflora* King. (Dipterocarpaceae)

Nik Fatini Nik Azmin^a, Noorazlina Adnan^{a,b,c}, Siti Zakirah Azahar^{a,c}, Norizan Ahmat^b, Aisyah Salihah Kamarozaman^{b,c,*}

^a Faculty of Applied Sciences, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

^b Centre of Foundation Studies, Universiti Teknologi MARA, Cawangan Selangor, Kampus Dengkil, 43800 Dengkil, Selangor, Malaysia

^c Atta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA, Cawangan Selangor, Kampus Puncak Alam, 42300 Bandar Puncak Alam, Selangor, Malaysia

Polyphenols are a broad chemical class of phytochemicals, with the stilbenoid (resveratrol) subclass being extensively studied. Both individual compounds and plant extracts enriched with stilbenoids have been widely examined (Kasiotis et al., 2013, Shen et al., 2009) due to their diverse biological effects (Duta-Bratu et al., 2023). Stilbenoids have garnered attention for their isolation and investigation as monomers, oligomers and glycosylated derivatives. In continuation to the previous research, the phytochemical study from the stem bark of *Shorea pauciflora* has successfully afforded three resveratrol oligomers. Fractionation of the crude acetone extract (148 g) was executed using vacuum liquid chromatography with the increasing polarity of *n*-hexane:ethyl acetate and ethyl acetate:methanol to give ten semi-purified fractions. Compounds **1** and **2** were obtained from fraction 9 (1 g) using column chromatography twice, with *n*-hexane: ethyl acetate followed by chloroform: methanol as a solvent system, respectively. Fraction 4 (1 g) was also subjected to column chromatography followed by purification using preparative thin layer chromatography with chloroform: methanol as eluent to obtain compound **3**. Based on the spectroscopic analyses and comparison with reported data, the compounds were elucidated as resveratrol tetramers, namely isohopeaphenol (**1**) (Kang et al., 2020) and hemsleyanol D (**2**) (Lim et al., 2023) as well as a new resveratrol dimer, compound Z (**3**) (Fig. 1). This research was supported by the Ministry of Higher Education, Malaysia through the Fundamental Research Grant Scheme (FRGS/1/2022/STG04/UITM/02/24).

Keywords: *Shorea pauciflora*, resveratrols, isohopeaphenol, hemsleyanol D, new compound Z




Fig 1. Structure of resveratrol oligomers from *Shorea pauciflora*

References

- Kasiotis, K.M., Pratsinis, H., Kletsas, D., Haroutounian, S.A., 2013. Resveratrol and related stilbenes: their anti-aging and anti- angiogenic properties. *Food and Chemical Toxicology*, 61, 112-120.
- Shen, T., Wang, X.N., Lou, H.X., 2009. Natural stilbenes: an overview. *Natural product reports*, 26(7), 916-935.
- Duta-Bratu, C.G., Nitulescu, G.M., Mihai, D.P., Olaru, O.T., 2023. Resveratrol and other natural oligomeric stilbenoid compounds and their therapeutic applications. *Plants*, 12(16), 2935.
- Kang, J.E., Jeon, B.J., Park, M.Y., Yang, H.J., Kwon, J., Lee, D.H., Kim, B.S., 2020. Inhibition of the type III secretion system of *Pseudomonas syringae* pv. tomato DC3000 by resveratrol oligomers identified in *Vitis vinifera* L. *Pest management science*, 76(7), 2294-2303.
- Lim, P.C., Ramli, R., Manshoor, N., 2023. Miyabenol C isomers and other oligostilbenes from the stem of *Dipterocarpus semivestitus* Sloot. and their chemotaxonomic significance. *Biochemical Systematics and Ecology*, 110, 104685.

* Presenting author. E-mail address: aisyah2910@uitm.edu.my (Aisyah Salihah Kamarozaman)

S1.P199 In-silico and *in-vitro* evaluation of natural product scaffolds to overcome the scourge of drug resistant tuberculosis by virtual based drug design approach

Vilas Jagatap*, Harun Patel

R C Patel Institute of Pharmaceutical Education and Research, Shirpur, Dhule, Maharashtra, India- 425405

In pursuit of new anti-tubercular agents, we here report the anti-mycobacterial (H37Rv) and DNA gyrase inhibitory potential of daidzein and khellin natural products (NPs). We procured a total of 16 NPs based on their pharmacophoric similarities with known antimycobacterial compounds (Coronado-Aceves et al., 2017, Krishna et al., 2019). The H37Rv strain of *M. tuberculosis* was found to be susceptible to only two out of the 16 NPs procured; specifically, daidzein and khellin each exhibited an MIC of 25 µg/mL. Moreover, daidzein and khellin inhibited the DNA gyrase enzyme with IC₅₀ values of 0.042 and 0.822 µg/mL, respectively, compared to ciprofloxacin with an IC₅₀ value of 0.018 µg/mL. Daidzein and khellin were found to have lower toxicity toward the vero cell line, with IC₅₀ values of 160.81 and 300.23 µg/mL, respectively. Further, molecular docking study and MD simulation of daidzein (**Fig. 1**) indicated that it remained stable inside the cavity of DNA GyrB domain for 100 ns.




Fig. 1. Structure of daidzein natural product

Keywords: Tuberculosis, anti-mycobacterials, natural products, daidzein, DNA gyrase

References

- E.W. Coronado-Aceves, G. Gigliarelli, A. Garibay-Escobar, R. E. R. Zepeda, M. Curini, J. López Cervantes, C. I. Inés Espitia Pinzón, S. Superchi, S. Vergura, M. C. Marcotullio, New Isoflavonoids from the extract of *Rhynchosia precatória* (Humb. & Bonpl. ex Willd.) DC. and their anti-mycobacterial activity, *J. Ethnopharmacol.* 206 (2017) 92– 100.
- V. S. Krishna, S. Zheng, E. M. Rekha, L. W. Guddat, D. Sriram, Discovery and evaluation of novel *Mycobacterium tuberculosis* ketol-acid reductoisomerase inhibitors as therapeutic drug leads, *J. Comput. -Aided Mol. Des.* 33 (2019) 357– 366.

* Presenting author. E-mail address: jagtapv390@gmail.com (Vilas Jagatap)

S1.P200 Exploring Malaysian biodiversity in the fight against antibiotic resistance

Isabel Ramón Martínez^{a,*}, Saoussen Oueslati^{b,c}, Cécile Apel^a, Thierry Naas^{b,c}, Marc Litaudon^a, Sandy Desrat^a, Fanny Roussi^a

^a Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, UPR 2301, 1 Avenue de la Terrasse, 91198 Gif-sur-Yvette, France

^b Team ReSIST, INSERM U1184, School of Medicine, Université Paris-Saclay, 94270 Le Kremlin-Bicêtre, France

^c Bacteriology-Hygiene Unit, Bicêtre Hospital, APHP Paris-Saclay, 94270 Le Kremlin-Bicêtre, France

Carbapenemases are enzymes produced as a resistance mechanism by various bacteria to hydrolyse carbapenem antibiotics. These enzymes propagate rapidly, contributing to the alarming ineffectiveness of these kind of large spectrum antibiotics all over the world (Taconelli *et al.*, 2018). A series of 824 Rutaceae and Annonaceae plant extracts from the extract library of the Institut de Chimie de Substances Naturelles (Gif-sur-Yvette, France) was screened for inhibition of three of the most extended varieties of carbapenemase enzymes: KPC-2, OXA-48 and NDM-1 (Bush *et al.*, 2020).

The bark extract of *Fissistigma litseaefolium* (King) Merr, a Malaysian endemic climber Annonaceae species, was found active and was thus selected for a bioassay-guided isolation. The fact that this species has not been explored before from a metabolomic standpoint added an extra dimension to its study. A comparison of the metabolome of this species with others of the genera by molecular networking using MetGem software (Olivon *et al.*, 2018) and annotation tools revealed differences and guided to the isolation of novel structures.

Two new molecules derived from an original double cyclisation of a prenylated-chalcone have been isolated. Their absolute configuration has been determined thanks to X-Ray spectroscopy. All these results (**1**), as well as the proposed biosynthetic pathway conducting to these structures and hemisynthesis will be presented.




Fig. 1. Toward carbapenemase inhibitors isolation from ICSN's extract library

Keywords: *Fissistigma* sp., carbapenemases, molecular networking, isolation, hemisynthesis

References

- Bush K, Bradford PA. 2020. Epidemiology of β -Lactamase-Producing Pathogens. Clin Microbiol Rev. 33(2).
- F. Olivon, N. Elie, G. Grelier, F. Roussi, M. Litaudon, and D. Touboul, 2018. MetGem software for the generation of molecular networks based on t-SNE algorithm, Anal. Chem. 90, 23, 13900-13908.
- Taconelli E. *et al.* 2018. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis. 18(3):318-327.

* Presenting author. E-mail address: isabel.ramon-martinez@cnrs.fr (Isabel Ramón Martínez)

S1.P201 Effect of seasonal and geographical location on the secondary metabolic contents of *Artemisia afra* and *Artemisia annua*. Anti-plasmodial properties

Lahngong Methodius Shinyuy^{a,*}, Ledoux Allison^a, Olivia Jansen^a, Frédéric Michel^a, Palmaerts Benjamin^b, Hallot Eric^b, E. Loe Gisèle^c, Ghogomu Stephen^d, Souopgui Jacob^e

^a Pharmacognosy Laboratory, Center of Interdisciplinary Research on Medicine (CIRM), Uliège, 4000 Liège, Belgium

^b Remote Sensing and Geodata Unit, Institut Scientifique de Service Public (ISSeP), 4000 Liège, Belgium

^c Laboratory of Pharmacochemical and natural pharmaceutical substances, Doctoral Training Unit in Health Sciences. Faculty of Medicine and Pharmaceutical Sciences, University of Douala, Cameroon

^d Molecular and Cell Biology Laboratory (MCBL), Department of Biochemistry and Molecular Biology, Faculty of Science, University of Buea, Cameroon

^e Embryology and Biotechnology Laboratory, Université Libre de Bruxelles, Belgium

Malaria is a disease of public health concern, and natural products have been investigated as potential treatments. Due to the emergence of resistance to Artemisinin, traditional medicinal preparation using *A. afra* and *A. annua* is practiced in Africa where malaria is endemic. This study explores the phytochemical diversity and anti-plasmodial potential of *A. annua* and *A. afra* sourced from distinct geographical locations within Cameroon aiming to discern potential variations in phytochemical profiles and correlated to observed activity. Crude extracts were prepared from plants collected from diverse regions in Cameroon during the rainy/dry seasons and metabolic contents analyzed by TLC, HPLC, and GC. Additionally, anti-plasmodial potential was assessed on 3D7 *Plasmodium falciparum* strain. The activity profiles of the samples were correlated with their environment, with distinct phytochemical compositions observed for each sample based on its geographical origin and season of collection. Traces of artemisinin were detected in some *A. afra* samples but present in all *A. annua* samples with a high concentration in the rainy season samples (better anti-plasmodial activity). The Adamawa region had the highest artemisinin contents (8.9% m/m artemisinin in dry extract) with better anti-plasmodial activity. These findings suggest that the selection of a suitable *Artemisia* sample for use as a potential antimalarial treatment should take into consideration its geographical origin and season. By understanding how location and seasonal changes influence the phytochemical composition of *Artemisia* samples, this study may provide an insight into how natural products can be effectively used as a preventative or curative measure against malaria.

Keywords: malaria, *Artemisia annua*, *Artemisia afra*, resistance, *Plasmodium*

References

[1]. W. Trager and J. B. Jensen. Human Malaria Parasites in Continuous Culture. *Science* (1979). doi: 10.1126/SCIENCE.781840.

* Presenting author. E-mail address: lahngongmethodius@gmail.com (Methodius Shinyuy Lahngong)

S1.P202 Ecology, metabolome and occurrence of *Pseudallescheria* sp. associated with termites

Tomas Herrera-Durigneux^a, Jérôme Muriienne^b, David Touboul^c, Véronique Eparvier^{a,*}

^a Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, UPR 2301, 91198, Gif-sur-Yvette, France

^b Laboratoire Évolution & Diversité Biologique UMR 5300 Université Toulouse III Paul Sabatier–Bâtiment 4R1 118, route de Narbonne 31062 Toulouse cedex 9, France

^c LCM, CNRS, Ecole polytechnique, Institut Polytechnique de Paris, Route de Saclay, 91120 Palaiseau, France

Termites are eusocial insects that live in homeostatic environments, confined spaces and with high population density (Liu *et al.*, 2019). This life style leads to an intimate association with microscopic life forms. These microorganisms can be parasites, mutualists, opportunists or pathogens (Douglas *et al.*, 2016). Until now, studies on the termite holobiont have focused on trophobiosis-related symbioses that allows the degradation of lignocellulose in the termite gut. However, exploration of other symbiotic relationships may show their ecological role in termite survival whilst environmental conditions may promote the transmission of infectious agents. It can also lead to the discovery of new chemical entities with therapeutic interest (Beemelmans *et al.*, 2016).

During previous investigations carried out by ICSN, fungal strains of the genus *Pseudallescheria boydii*, from different termite species of French Guiana were identified (frequently associated with *Nasutitermes* sp.). The fungi's metabolome was analyzed. When co-inoculated with the entomopathogen *Beauveria* sp., antimicrobial compounds like Tyroscherin and *N*-methyltyroscherin (Fig. 1) were overproduced by the symbiont (Sorres *et al.*, 2017, 2022), indicating a probable host defense role under biotic stress when in association with the termite species. Current studies aim to determine if this termite-*Pseudallescheria* sp. relationship is widespread focusing on the occurrence of this interaction in anthropophized and non-anthropophized environments. Today, 14 new nests from French Guiana are being analyzed by LC-HRMS/MS, using Tyroscherin as an indicator of the presence of fungi in the various parts of the macro-holobiont „termite nest” (including termites, nest and tunnels).




Fig. 1. Structure of Tyroscherin 1 and *N*-methyltyroscherin 2 from *Pseudallescheria* sp. strains.

Keywords: *Nasutitermes* sp., *Pseudallescheria* sp., Tyroscherin, Symbiosis, Chemical ecology

References

- Beemelmans, C.; Guo, H.; Rischer, M.; Poulsen, M. 2016. Natural Products from Microbes Associated with Insects. *Beilstein J. Org. Chem.* 12, 314–327.
- Douglas, A. E. 2015. Multiorganismal Insects: Diversity and Function of Resident Microorganisms. *Annu. Rev. Entomol.* 60, 17–34.
- Sorres, J.; Nirma, C.; Eparvier, V.; Stien, D. 2017. Pseudallicins A-D, four complex Ovalicin derivatives from *Pseudallescheria boydii* SNB-CN85. *Organic Letters*, 9, 3978–3981.
- Sorres, J.; Hebra, T.; Elie, N.; Leman-Loubière, C.; Grayfer, T.; Grellier, P.; Touboul, D.; Stien, D.; Eparvier, V. 2022. Bioactive fatty-acid derivated ovalicin from *Pseudallescheria boydii*. *Molecules* 27, 1182.
- Liu, L.; Zhao, X.-Y.; Tang, Q.-B.; Lei, C.-L.; Huang, Q.-Y. 2019. The Mechanisms of Social Immunity Against Fungal Infections in Eusocial Insects. *Toxins* 11 (5), 244.

* Presenting author. E-mail address: veronique.eparvier@cnrs.fr (Véronique Eparvier)

S1.P203 *Tetracera alnifolia* (Wild) Drake popular in Guinean traditional medicines possesses some pharmacological activities

Elhadj Saidou Baldé^{a,b}, Aboubacar Kadiatou Camara^{a,c,*}, Mamadou Saliou Telly Diallo^b, Mohamed Kerfalla Camara^b, Tanou Valdez Bah^{a,b}, Mamoudou Condé^c, Aboubacar Soumah^c, Ian Tietjen^d, Aliou Mamadou Baldé^{a,b}

^a Faculty of Health Sciences and Techniques, Gamal Abdel Nasser University of Conakry, BP 1147, Guinea

^b Institute for Research and Development of Medicinal and Food Plants of Guinea-Dubréka, BP 6411, Guinea

^c National Institute of Public Health, Ministry of Health and Public Hygiene, Conakry, BP 6623, Guinea

^d The Wistar Institute, Philadelphia, Pennsylvania, USA

Several parts of the woody plant *T. alnifolia* are traditionally used for treating infectious diseases, sexually transmitted infections, skin diseases, and malaria in Guinean traditional medicine (Baldé et al., 2020; Magassouba et al., 2007; Traore et al., 2013). In this context, based on the ethnomedical investigations conducted by IRDPMAG, *T. alnifolia* was selected to confirm some of its traditional uses in Guinea as well as to survey its biological properties. We conducted an ethnomedical survey across several markets of the city of Conakry to identify and interview healers. Chloroform, methanol, dichloromethane, and aqueous extracts of *T. alnifolia* were tested for activities against protozoa, bacteria, fungi, HIV, and SARS-CoV-2.

39 traditional healers indicated that *T. alnifolia* is used in the treatment of more than 15 pathologies including *Fassa* (marasmus/malnutrition), *Soukhoulou kouyé* (white discharge in women), and *Tèmou bankhi* (sexual weakness in men). Leaves were the most used part. Modes of preparation included decoction and powder. Data from biological activities identified good activities of the methanolic extract against *Leishmania infantum* (MIC = 8.1 µg/ml) and a moderate activity against *Trypanosoma brucei* (MIC = 28.2 µg/ml) and *Staphylococcus aureus* (MIC = 29.9 µg/ml). Dichloromethane extracts inhibited up to 53.4% of SARS-CoV-2 replication at 50 µg/mL.

These results explain at least in part the traditional use of *T. alnifolia*. Aiming to rationalise the use of *T. alnifolia*, bioassay-guided fractionations are in progress to determine useful fractions or molecules for clinical investigations.

Keywords: *Tetracera alnifolia*, ethnomedical investigations, biological activity, SARS-CoV-2, protozoa

References

- Baldé, M.A., Tuenter, E., Traoré, M.S., Matheeuissen, A., Cos, P., Maes, L., Camara, A., Haba, N.L., Gomou, K., Diallo, M.S.T., Baldé, E.S., Pieters, L., Balde, A.M., Foubert, K., 2020. Antimicrobial investigation of ethnobotanically selected guinean plant species. *J. Ethnopharmacol.* 263, 113232.
- Magassouba, F.B., Diallo, A., Kouyaté, M., Mara, F., Mara, O., Bangoura, O., Camara, A., Traoré, S., Diallo, A.K., Zaoro, M., Lamah, K., Diallo, S., Camara, G., Traoré, S., Kéita, A., Camara, M.K., Barry, R., Kéita, S., Oularé, K., Barry, M.S., Donzo, M., Camara, K., Toté, K., Berghe, D.V., Totté, J., Pieters, L., Vlietinck, A.J., Baldé, A.M., 2007. Ethnobotanical survey and antibacterial activity of some plants used in Guinean traditional medicine. *J. Ethnopharmacol.* 114, 44–53.
- Traore, M.S., Baldé, M.A., Diallo, M.S.T., Baldé, E.S., Diané, S., Camara, A., Diallo, A., Balde, A., Kéita, A., Keita, S.M., Oularé, K., Magassouba, F.B., Diakité, I., Diallo, A., Pieters, L., Baldé, A.M., 2013. Ethnobotanical survey on medicinal plants used by Guinean traditional healers in the treatment of malaria. *J. Ethnopharmacol.* 150, 1145–1153.

* Presenting author. E-mail address: esbalde@irdpmag.edu.gn (Elhadj Saidou Baldé)

S1.P204 Specialized metabolites with antiviral properties from a Vietnamese plant: isolation and synthesis

Phan Linh^{a,*}, Petit Bastien^a, Haddad Juliano^b, El Kalamouni Chaker^b, Cachet Xavier^c, Gaslonde Thomas^c, Litaudon Marc^a, Apel Cécile^a, Desrat Sandy^a, Roussi Fanny^a

^a Université Paris-Saclay, CNRS, Institute of Chemistry of Natural Substances (ICSN), UPR 2301, 91198, Gif-sur-Yvette, France

^b Université de Réunion, Mixed Unit for Infectious Processes in Tropical Island Environments (PIMIT), CYROI Technological Platform, INSERM U1187, CNRS UMR 9192, IRD UMR 249, 94791 Sainte Clotilde, Réunion, France

^c Université Paris Cité, Faculty of Pharmacy of Paris, CNRS UMR 8038, 4 Avenue de l'Observatoire, 75270 Paris Cedex 06, France

The plant metabolites group at the Institute of Chemistry of Natural Substances (ICSN) aims to study and valorize specialized metabolites isolated from plants. For that, the plant extract library named Extractothèque ICSN, containing more than 15,000 extracts, is regularly screened by biologist partners.

After the COVID-19 pandemic, as part of this project, a selection of 824 plant extracts from Extractothèque ICSN was evaluated on an inhibition cell-based assay on human Coronavirus (HCoV- 229E and SARS-CoV-2). The EtOAc and MeOH extracts of the leaves of the Vietnamese species *Melodorum fruticosum* were selected for their good activity on HCoV-229E. Bio-guided fractionation led to the isolation of 11 pure products, including 3 novel compounds. Their antiviral evaluation confirmed the interest of some metabolites with IC₅₀ around 1 M for toussaintine C. To validate their activity and improve our knowledge of the structure-activity relationships, we first performed the total synthesis of racemic toussaintine C (Fig. 1) and developed a divergent synthesis of analogs. As one of the enantiomers of toussaintine C (Fig. 1) has been found to exhibit greater activity against HCoV-229E, we developed an asymmetric total synthesis leading to both enantiomers.




Fig. 1. Total synthesis pathway of toussaintine C and synthesis analogs *Keywords:* *Melodorum fruticosum*, HCoV-229E, SARS-CoV-2, antiviral, toussaintine C

References

Zhou, B., Gong, Q., Fu, Y., Zhou, J.S., Zhang, H.Y., Yue, J.M., 2023. Sarglamides A–E, indolidinoid – monoterpenoid hybrids with anti-neuroinflammatory activity from a *Sarcandra* species. *Org. Lett.* 25 (9), 1464–1469.

* Presenting author. E-mail address: linh.phan@cnrs.fr (Phan Hoang Linh)

S1.P205 From signaling molecules to secondary metabolites: harnessing c-di-GMP for enhanced natural product biosynthesis in *Streptomyces*

Olaf Latta^a, Robin Teufel^b, Markus Häner^c, Henning J. Jessen^c, Andreas Bechthold^a

^a Institute for Pharmaceutical Biology and Biotechnology, University of Freiburg, Germany

^b Division of Pharmaceutical Biology, Department of Pharmaceutical Sciences, University of Basel, Switzerland

^c Institute for Organic Chemistry, University of Freiburg, Germany

The regulatory mechanisms governing natural product biosynthesis in *Streptomyces* species have long intrigued researchers due to their potential in drug discovery and biotechnology. Among these mechanisms, cyclic di-GMP (c-di-GMP) signaling has emerged as a key player, orchestrating various cellular processes (1). In this study, we delve into the intricate world of c-di-GMP signaling in *Streptomyces* and its implications for natural products research. In addition to unraveling the interplay between c-di-GMP signaling and natural product biosynthesis, our study highlights *Streptomyces*' crucial role in antibiotic discovery, especially amidst rising resistance, tapping into their potential to produce new antibiotics.

Through capture compound experiments, utilizing a modified c-di-GMP molecule with a UV-reactive group for crosslinking and biotin-residue for affinity purification, we elucidated novel c-di-GMP effectors (2). This molecule was synthesized in-house. Investigating the strain *Streptomyces* sp. Tu6071, we are searching for potential targets involved in transcriptional regulation, that are crucial for activating new biosynthetic gene clusters.

These findings not only deepen our understanding of c-di-GMP signaling dynamics, but also offer promising avenues for manipulating natural product biosynthesis. In the future, these potential targets will be confirmed, and their direct connection to the biosynthesis of natural products will be elucidated.

Moreover, our study opens doors to the activation of silent gene clusters, harboring untapped reservoirs of bioactive compounds. Unraveling the intricate interplay between c-di-GMP signaling and natural product biosynthesis holds tremendous potential for accelerating drug discovery efforts and harnessing the vast chemical diversity of *Streptomyces*.

Keywords: c-di-GMP signaling, *Streptomyces*, natural products, capture compound experiments, gene regulation

References

1. Makitrynskyy R, Tsyplik O, Nuzzo D, Paululat T, Zechel DL, Bechthold A. Secondary nucleotide messenger c-di-GMP exerts a global control on natural product biosynthesis in streptomycetes. *Nucleic Acids Res.* 2020;48(3):1583–98.
2. Nesper J, Reinders A, Glatter T, Schmidt A, Jenal U. A novel capture compound for the identification and analysis of cyclic di-GMP binding proteins. *J Proteomics.* 2012 Aug 3;75(15):4874–8.

* Presenting author. E-mail address: Olaf.latta@pharmazie.uni-freiburg.de (Olaf Latta)

S1.P206 A novel natural product from the leaves of a *Solanum spec.* (Solanaceae) shows antimicrobial activity against multiple multi-resistant pathogens

Alexander Sonntag^a, Karin Gorzolka^b, Myong-Su Lenz^b, Fabian Herrmann^c, Matthias F. Melzig^a, Alexander Weng^a

^a Freie Universität Berlin, Institute of Pharmacy, Department of Pharmaceutical Biology, Königin-Luise-Str. 2+4, 14195 Berlin, Germany

^b Julius-Kühn-Institut, Institute of Ecological Chemistry, Plant Analysis and Stock Protection, Königin-Luise-Str. 19, 14195 Berlin, Germany

^c University of Münster, Institute of Pharmaceutical Biology and Phytochemistry, Corrensstr. 48, 48149 Münster, Germany

Due to the extensive use of antibiotics in both human medicine and factory farming, there is an alarming increase of infections with antibiotic-resistant bacteria that do not respond to standard therapy. These strains develop resistance to reserve antibiotics, posing a threat to public health (Frieri et al., 2017). Between 1981 and 2019 approximately 55 % of new antibiotics are natural products or their derivatives (Newman et al. 2020), and consequently, substances from natural sources are promising in drug development.

A new natural product was isolated and identified from the leaves of a wild potato species (Solanaceae) and was shown to be bioactive against the Colorado potato beetle (*Leptinotarsa decemlineata*) and plant pathogenic fungi (not published). Here we investigated this product for its activity against human pathogens.

Hence, seven different bacteria from the WHO-Pathogen-Priority-List (*Acinetobacter baumannii*, *Campylobacter jejuni*, *Enterococcus faecium*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*) and *Candida albicans* were tested against the natural product and the MIC-values were determined using the broth microdilution method of EUCAST. The substance showed activity against every pathogen (MIC between 1024 µg/mL and 32 µg/mL).

Atomic force microscope imaging indicated a short-term membrane toxic effect on the bacteria.

To assess the cytotoxicity, an MTT assay using the immortalized epithelial kidney cell line Vero E6 (*Cercopithecus aethiops*) and human primary fibroblasts was performed. These data indicated high toxicity (IC₅₀ = 2.2 µg/mL and 1.2 µg/mL) making the natural product most likely to be unsuitable for use in human medicine to treat infections.

Keywords: *Solanum*, Solanaceae, anti-infective, cytotoxicity, bioactive

References

Frieri, M., Kumar, K., Boutin, A., 2017. Antibiotic resistance. J Infect Public Health. 10-4, 369-378.

Newman, D.J., Cragg G.M., 2020. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. J. Nat. Prod. 83-3, 770-803.

* Presenting author. E-mail address: alexander.sonntag@fu-berlin.de (Alexander Sonntag)

S1.P207 Prenylated phenolics from Fabaceae as inhibitors of the multidrug efflux pump NorA


Marina Ika Irianti^a, Jean-Paul Vincken^a, Sarah van Dinteren^a, Ellen ter Beest^a, Klaas Martinus Pos^b, Carla Araya-Cloutier^{a,*}

^a Laboratory of Food Chemistry, Wageningen University and Research, Bornse Weiland 9, 6708 WG, Wageningen, The Netherlands

^b Institute of Biochemistry, Goethe-University Frankfurt, 60438, Frankfurt am Main, Germany

Considering their key role in the emerge of antimicrobial resistance (AMR), efflux pumps are an interesting target for the development of novel resistance-modifying agents, such as efflux pump inhibitors (EPIs). Among different efflux pumps, members of the Major Facilitator Superfamily (MFS) are the most prevalent efflux pumps in gram-positive bacteria, including *Staphylococcus aureus*, for which the overexpression of the MFS efflux pump NorA is observed in 43% of strains, particularly in MRSA strains (Patel et al., 2010). EPIs are a promising solution as they can re-sensitize resistant bacteria to common antibiotics and reduce the emerge of AMR.

Plants are an excellent source of bioactive compounds, including EPIs. Some specific plant EPIs identified contain prenyl groups (isoprene moieties) in their backbone (Prasch & Bucar, 2015, Ika Irianti et al., 2023). These prenyl groups increase hydrophobicity of the molecule and their affinity towards biological targets. The Fabaceae plant family produces prenylated phenolics as main defence mechanism against microbial stress.



To understand the structure-function relationship as potential NorA EPIs, an extended collection of prenylated phenolics was investigated, belonging to different flavonoid and isoflavonoid subclasses, substituted with different number and configuration of prenyl groups. Characterization was done in terms of NorA inhibition using ethidium accumulation and antibiotic potentiation assays, using both *norA* overexpressing and knockout *S. aureus* strains. Cytotoxicity of promising candidates was assessed using Caco-2 cell lines. Given the chemical diversity of the collection investigated, an in- silico analysis was performed to understand key molecular properties for NorA inhibition.

Keywords: Prenylated isoflavonoids, Fabaceae, *Staphylococcus aureus*, NorA, efflux pump

References

- Patel, D., Kosmidis, C., Seo, S. M. & Kaatz, G. W. 2010. Ethidium bromide MIC screening for enhanced efflux pump gene expression or efflux activity in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 54, 5070–5073.
- Prasch, S., Bucar, F. 2015. Plant derived inhibitors of bacterial efflux pumps: an update. *Phytochem Rev* 14, 961–974. Ika Irianti, M., Vincken, J.P., van Dinteren, S. ter Beest, E., Pos, K.M., Araya-Cloutier, C. 2023. Prenylated isoflavonoids from Fabaceae against the NorA efflux pump in *Staphylococcus aureus*. *Sci Rep* 13, 22548

* Presenting author. E-mail address: carla.arayacloutier@wur.nl (Carla Araya-Cloutiera)

S1.P208 Improving the activity of pleuromutilin antibiotics

Kerrin Hainsworth^{a,*}, Melissa M. Cadelis^{a,b}, Florent Rouvier^c, Marie-Lise Bourguet-Kondracki^d, Jean-Michel Brunel^d, Brent R. Copp^a

^a School of Chemical Sciences, The University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

^b School of Medical Sciences, The University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

^c Membranes et Cibles Thérapeutiques, SSA, INSERM, Aix-Marseille Université, 27 bd Jean Moulin, 13385 Marseille, France

^d Laboratoire Molécules de Communication et Adaptation des Micro-organismes, UMR 7245 CNRS, Muséum National d'Histoire Naturelle, 57 rue Cuvier (C.P. 54), 75005 Paris, France

With rising levels of antimicrobial resistance and the slow development of new antibiotics, there is increasing interest in re-vitalizing existing agents. Pleuromutilin is a diterpenoid natural product antibiotic that was first isolated in the 1950s from *Pleurotus mutilis* (now *Clitopilus scyphoides*) (Kavanagh, et al., 1952). The ease of modification of the C-14 glycolate side chain has led to the discovery of more active variants, however, they remain predominantly active against Gram-positive and mycoplasma bacteria (Egger et al., 1976, Goethe et al., 2019).

In our ongoing efforts to develop antibiotics against Gram-negative bacteria, we synthesised a series of pleuromutilins with modified sidechains that explored several strategies to turn Gram-positive- only antibiotics into Gram-negative agents. The first is increasing the presence of ionizable amines which has been suggested to increase the uptake and accumulation of small molecule antibiotics in Gram-negative bacteria (Richter et al., 2017). We have also extended the sidechain with a variety of end groups through peptide coupling to facilitate targeted delivery (Fig. 1). The synthesis and antimicrobial activities of these derivatives, along with their cytotoxic and hemolytic activity, will be presented.




Fig. 1. General structure of pleuromutilin derivatives for targeted delivery. X is S or NH.

Keywords: pleuromutilin, antibiotics, antifungal activity, targeted delivery

References

- Egger, H., Reinshagen, H., 1976. New pleuromutilin derivatives with enhanced antimicrobial activity II. Structure activity correlations. J. Antibiot. 29, 923-927.
- Goethe, O., Heuer, A. Ma, X. Wang, Z. Herzon, S.B., 2019. Antibacterial properties and clinical potential of pleuromutilins. Nat. Prod. Rep. 36, 220-247.
- Kavanagh, F., Hervey, A., Robbins, W.J., 1952. Antibiotic substances from basidiomycetes. Proc Natl Acad Sci U S A. 38, 555-560.
- Richter, M.F., Drown, B.S., Riley, A.P., Garcia, A., Shirai, T., Svec, R.L., Hergenrother, P.J., 2017. Predictive compound accumulation rules yield a broad-spectrum antibiotic. Nature. 545, 299-304.

* Presenting author. E-mail address: khai892@aucklanduni.ac.nz (Kerrin Hainsworth)

S1.P209 Bioactivity-guided investigations on biofilm inhibition with *Rubus chamaemorus* leaf extracts

Hildegard Kieninger*, Sebastian Günther, Christian Schulze

Pharmaceutical Biology, Institute of Pharmacy, University of Greifswald, Friedrich-Ludwig-Jan- Straße 17, 17489 Greifswald, Germany

Cloudberry (*Rubus chamaemorus* L., Rosaceae) is known for its health-promoting fruits rich in antioxidants and historical use against scurvy (Luca et al., 2011; Kähkönen et al., 2001). While the fruits are well studied, cloudberry leaves remain less investigated (Fig.1). Traditional use as antidiarrheal and for wound treatment are reported. (Rocabado et al., 2008). Recent investigations show biofilm inhibition against multidrug-resistant Gram-negative (MRGN) *Escherichia coli* (Neumann et al., 2022). Further investigation of this plant is highly interesting, considering its potential for dual utilization of leaves and fruits, cultivation in CO₂-storing bogs, and therapeutic application against multidrug-resistant pathogens. This study examines cloudberry leaf extracts, fractions, and sub-fractions using a bioactivity-guided approach.



Fig. 1. Cloudberry leaves.

Five different methods were used subsequent to methanol extraction: solid phase extraction utilizing a reversed-phase column for extract fractionation, High-Performance Liquid Chromatography for a fingerprint of the extracts, size exclusion chromatography for further fractionation, and mass spectrometry to identify active compounds. In parallel to all extraction and separation steps, bioactivity-guided tests were performed using a macrocolony biofilm assay with *E. coli*.

One fraction exhibited biofilm inhibition against MRGN *E. coli* at 10 µg/mL, outperforming the crude extract at 30 µg/mL. Mass spectrometric analysis suggested sanguin-H6 and lambertianin C, ellagitannins, as potential biofilm inhibition contributors. The fraction containing ellagic acid, quercetin-glucuronide, quercetin-galloylglucuronide, and kaempferol-glucuronide showed no biofilm inhibition. Ongoing assays aim to elucidate the potential involvement of quorum sensing in this biological activity. Understanding cloudberry's biofilm inhibition holds promise for phytopharmaceutical development and peatland preservation through cultivation.

Keywords: *Rubus chamaemorus*, anti-biofilm, MRGN *E. coli*, bioactivity-guided, tannins

References

- Kähkönen, M.P., Hopia, A.I., Heinonen, M., 2001. Berry phenolics and their antioxidant activity. J. Agric. Food Chem. 49 (8), 4076-4082.
- Luca, L.M. de, Norum, K.R., 2011. Scurvy and cloudberrries: a chapter in the history of nutritional sciences. J. Nutr. 141 (12), 2101–2105.
- Neumann, N., Honke, M., Povydysh, M., Guenther, S., Schulze, C., 2022. Evaluating Tannins and Flavonoids from Traditionally Used Medicinal Plants with Biofilm Inhibitory Effects against MRGN *E. coli*. Molecules 27 (7), 2284.
- Rocabado, G.O., Bedoya, L.M., Abad, M.J., Bermejo, P., 2008. *Rubus* - A Review of its Phytochemical and Pharmacological Profile. Nat. Prod. Commun. 3 (3), 423-436.

* Presenting author. E-mail address: hildegard.kieninger@uni-greifswald.de (Hildegard Kieninger)

S1.P210 Insights into the nature of the microalgal toxins from the *Chrysochromulina leadbeateri* blooms in Northern Norwegian fjords

Xinhui Wang^{a,*}, Mathias Fon^{b,c}, Aaron J.C. Andersen^a, Anita Solhaug^b, Richard A. Ingebrigtsen^{d,e}, Ingunn A. Samdal^b, Silvio Uhlig^b, Christopher O. Miles^{b,f}, Bente Edvardsen^c, Thomas O. Larsen^a

^a Technical University of Denmark, DTU Bioengineering, Kgs. Lyngby, Denmark

^b Norwegian Veterinary Institute, Ås, Norway

^c University of Oslo, Department of Biosciences, Oslo, Norway

^d Cawthron Institute, Nelson, New Zealand

^e UiT - The Arctic University of Norway, Norwegian College of Fishery Science, Tromsø, Norway

^f National Research Council Canada, Halifax, Canada

This presentation will cover the discovery and analysis of two novel polyhydroxylated polyketides, leadbeaterin-1 (1) and leadbeaterin-2 (2), from *Chrysochromulina leadbeateri*, identified as potential fish-killing toxins. In May–June 2019, the microalga *C. leadbeateri* caused a massive fish-killing event in several fjords in Northern Norway, resulting in the largest direct impact ever on aquaculture in northern Europe due to toxic algae (Samdal and Edvardsen, 2020; John et al., 2022). Motivated by the fact that no algal toxins have previously been described from *C. leadbeateri*, we set out to investigate the chemical nature and toxicity of secondary metabolites in extracts of two strains (UIO 393, UIO 394) isolated from the 2019 bloom, as well as one older strain (UIO 035) isolated during a bloom in Northern Norway in 1991. Bioassay-guided fractionation using the RTgill-W1 cell line (Bols et al., 1994; Dorantes-Aranda et al., 2011) and metabolomics analysis pointed to a major compound affording $[M+H]^+$ ions at m/z 1399.8333 as a possible toxin. Moreover, our study unveiled a series of minor analogues exhibiting distinct patterns of chlorination and sulfation, together defining a new family of compounds, which we propose to name leadbeaterins. Leadbeaterin-1 (1) and leadbeaterin-2 (2) were isolated and purified from UIO 394 cultures. Their structures were determined by 1D and 2D NMR experiments and tandem mass spectrometry. Remarkably, these suspected toxins were detected *in situ* in samples collected during the 2019 bloom close to Tromsø, thereby substantiating their likely role in fish kills.

Keywords: *Chrysochromulina leadbeateri*, algal bloom, ichthyotoxins, bioassay-guided fractionation, RTgill-W1

References

- Samdal, I., Edvardsen, B., 2020. Massive salmon mortalities during a *Chrysochromulina leadbeateri* bloom in Northern Norway. *Harmful Algae News*. 64, 4–6.
- John, U., Šupraha, L., Gran-Stadniczeňko, S., Bunse, C., Cembella, A., Eikrem, W., Janoušek, J., Klemm, K., Kühne, N., Naustvoll, L., Voss, D., Wohlrab, S., Edvardsen, B., 2022. Spatial and biological oceanographic insights into the massive fish-killing bloom of the haptophyte *Chrysochromulina leadbeateri* in northern Norway. *Harmful Algae* 118, 102287.
- Bols, N.C., Barlian, A., Chirinp-Trejo, M., Caldwell, S.J., Goegan, P., Lee, L.E.J., 1994. Development of a cell line from primary cultures of rainbow trout, *Oncorhynchus mykiss* (Walbaum), gills. *J. Fish Dis.* 17, 601–611.
- Dorantes-Aranda, J.J., Waite, T.D., Godrant, A., Rose, A.L., Tovar, C.D., Woods, G.M., Hallegraeff, G.M., 2011. Novel application of a fish gill cell line assay to assess ichthyotoxicity of harmful marine microalgae. *Harmful Algae* 10, 366–373.

* Presenting author. E-mail address: xinwang@dtu.dk (Xinhui Wang)

S1.P211 Revisiting F.P. Porcher's Resources of Southern Fields and Forests (pub. 1863): Insights into the Forgotten Medicinal Botany of the Southeastern United States

George S. Hanna*

Department of Public Health, Medical University of South Carolina, Charleston, SC 29425

Following the American Civil War, the Southeast United States was under blockade with limited the access to medical supplies. Dr. Francis Pyre Porcher, a physician and medicinal botanist at the Medical University of South Carolina was contracted to create a “repertory of scientific and popular knowledge as regards the medicinal, economical, and useful properties of the trees, plants, and shrubs found within the limits of the Confederate States.” In this seminal work, Porcher reported on the uses of nearly 800 species of plants with a focus on those native to his home region. Discussed here, with a focus on infectious disease, is an overview of Porcher's reported medicinal plant uses. Additionally, a bioassay guided isolation of anti-mycobacterial compounds from *Baccharis halimifolia* is presented. Porcher describes *B. halimifolia* thus, “This plant is of undoubted value and of very general use in popular practice in South Carolina as a palliative demulcent in consumption and cough. I have frequently seen it used with advantage and have often heard those employing it confess the benefit derived from it... Like many other of our indigenous plants possessed of unequivocal utility, is unnoticed in the dispensaries and other works.”

Keywords: *Mycobacterium*, *Baccharis*, antimicrobial, ethnobotany, medicinal plants

* Presenting author. E-mail address: hannag@musc.edu (George S. Hanna)

S1.P212 Bioassay-guided interpretation of antimicrobial compounds from *Limonia acidissima*

Ajmal Khan^{a,b,*}, Sujogya Kumar Panda^{a,c}, Walter Luyten^a

^a Department of Biology, Animal Physiology and Neurobiology Section, KU Leuven, Leuven, Belgium

^b Center for Animal Sciences and Fisheries, University of Swat, Swat, Khyber Pakhtunkhwa, Pakistan

^c Center of Environment, Climate Change and Public Health, Utkal University, Vani Vihar, Bhubaneswar, Odisha, India

Medicinal plants are a primary source of natural products which are used for the treatment of various infections throughout the world (Abdallah et al., 2023). Among, many medicinal plants, *Limonia acidissima* L. (Wood apple), Rutaceae, emerged out to be one of the most valuable plants, owing multiple medicinal properties (Gopinath et al., 2023). The present study was conducted to investigate the antimicrobial components of *L. acidissima* through bioassay-guided purification against *Staphylococcus aureus*, *Micrococcus luteus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Candida albicans* and *Saccharomyces cerevisiae*. The active peaks were collected and identified via LC- MS/MS. For identification, the COCONUT Database and a manually built LA database were used, in combination with prediction and calculation of mass fragmentation and retention time. A total of 10 active compounds were identified, including 3 alkaloids, 4 flavonoids, 1 anthraquinone, and 1 oxidoreductase against *S. aureus*; 1 alkaloid, 1 flavonoid, and 1 anthraquinone against *E. coli*; and only 1 alkaloid was active against *C. albicans*. All of them exhibited strong antimicrobial activities with good potential to be developed as antibiotics.

Keywords: plants, microbes, infections, *Limonia acidissima*, chromatography

References

- Abdallah, E. M., Alhatlani, B. Y., de Paula Menezes, R., and Martins, C. H. G., 2023. Back to Nature: Medicinal Plants as Promising Sources for Antibacterial Drugs in the Post-Antibiotic Era. *Plants*, 12, 3077.
- Gopinath, S. M., Sinchana, Anupama, G., Shravani, Megeri, T., and Danapur, V., 2023. "Pharmacognostic Studies and Biological Activities of *Limonia Acidissima* ", *Int. J. Pharma. Bio-Med. Sci.*, 3, 107-114.

* Presenting author. Email adress: ajmalkhan399@hotmail.com (Ajmal Khan)

S1.P213 Targeted isolation of potent antiviral cyclopeptides from tropical plant extracts

Cécile Apel^{a,*}, Juliano Haddad^b, Clément Grisel^a, Stéphanie Boutet^c, Fanny Roussi^a, Sandy Desrat^a, Chaker El Kalamouni^b, Marc Litaudon^a

^a Institut de Chimie des Substances Naturelles, CNRS, UPR 2301, Université Paris-Saclay, 91198 Gif-sur-Yvette, France

^b Unité Mixte Processus Infectieux en Milieu Insulaire Tropical, Université de la Réunion, INSERM U1187, CNRS UMR 9192, IRD UMR 249, Plateforme Technologique CYROI, 94791 Sainte Clotilde, France

^c Institut Jean-Pierre Bourgin, INRAE, AgroParisTech, Université Paris-Saclay, 78000 Versailles, France

Viral diseases represent a significant public health concern. The emergence of novel viruses, such as coronaviruses, and the resurgence of known pathogens such as dengue and Zika viruses, underscores the necessity for effective treatment options. The rich chemical diversity of plant metabolites is of great interest in the search for antiviral agents and requires innovative approaches to be explored.

In this context, biological assays were conducted on 824 plant extracts from the ICSN's extract library. Among them, eight extracts from phylogenetically related species exhibited significant inhibitory activities against Zika virus replication. The 824 samples were analyzed by LC- HRMS/MS, and the resulting data were processed using MZmine3 (Schmid et al., 2023). The metabolite content of the extracts was then organized and visualized as molecular networks using MetGem (Wang et al., 2016; Olivon et al., 2018). Overlaying the biological activity results onto the networks subsequently highlighted a cluster of ions highly specific to the eight active extracts (Fig. 1) (Olivon et al., 2017). Targeted isolation of the corresponding analogues afforded 8 undescribed cyclopeptides, which were further evaluated against a broader panel of viruses including dengue, chikungunya, Ross River and SARS-CoV-2 viruses. The most active compounds demonstrated antiviral activity in the nanomolar range, with low cytotoxicity. A medicinal chemistry project is underway for the development of an antiviral lead compound. Two patent applications are currently pending.




Fig. 1. Bioactivity-based molecular networking revealing a cluster of antiviral cyclopeptides

Keywords: Tropical plants, emerging viruses, antiviral agents, molecular networking

References

- Schmid, R., Heuckeroth, S., Korf, A. et al., 2023. Integrative analysis of multimodal mass spectrometry data in MZmine 3. *Nat. Biotechnol.* 41, 447-449.
- Wang, M., Carver, J., Phelan, V. et al., 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat. Biotechnol.* 34, 828-837
- Olivon, F., Elie, N., Grelier, G. et al., 2018. MetGem software for the generation of molecular networks based on t-SNE algorithm. *Anal. Chem.* 90, 23, 13900-13908.
- Olivon, F., Allard, PM., Koval, A. et al., 2017. Bioactive Natural Products Prioritization Using Massive Multi- informational Molecular Networks. *ACS Chem. Biol.* 12, 2644-2651.

* Presenting author. E-mail address: cecile.apel@cnrs.fr (Cécile Apel)

S1.P214 Anti-*Helicobacter pylori* compounds isolated from *Sambucus williamsii*

Dong-Min Kang^a, Woo-Jin Jeong^a, Atif Ali Khan Khalil^b, Mi-Jeong Ahn^{a,*}

^a College of Pharmacy and Research Institute of Pharmaceutical Sciences, Gyeongsang National University, Jinju 52828, Republic of Korea

^b Department of Pharmacology, School of Medicine, Institute of Health Sciences, Gyeongsang National University, Jinju 52727, Republic of Korea

Sambucus williamsii Hance (Viburnaceae) has been widely used in herbal medicine and food industry. It is known to have various pharmacological effects including antitumor, antioxidant, anti-inflammatory, and antimicrobial activities (Xiao et al., 2016). During our search for anti-*Helicobacter pylori* compounds from natural resources, the methanol extract of the *S. williamsii* branch significantly inhibited the growth of *H. pylori*. Four lignans and two phenolic compounds were isolated from the methylene chloride fraction that showed the most potent anti-*H. pylori* activity among the five fractions of hexane, methylene chloride, ethyl acetate, butanol, and water fractions. The chemical structures were identified to be boehmenan (**1**), pinnatifidanin B VII (**2**), guaiacylglycerol β -coniferyl ether (**3**), guaiacylglycerol- β -O-6'-(2-methoxy) cinnamyl alcohol ether (**4**), guaiacylglycerol (**5**) and 3-hydroxy-5-methoxy benzenemethanol (**6**) by UV, ¹H-NMR, ¹³C-NMR and mass spectrometry (Fig. 1) (Huang et al., 2013). Among the isolates, compounds **1** and **5** exhibited significant anti-*H. pylori* activity against strains 26695 and 51. Compound **5** displayed more potent anti-bacterial activity with MIC values of 3.13 and 6.25 μ M, and MIC₅₀ values of 28.5 and 56.8 μ M against the two strains, respectively. Their inhibitory activities were higher than those of a positive control, quercetin. Furthermore, these two compounds showed moderate urease inhibitory activity. A molecular docking study revealed the high binding ability of **1** and **5** to the active site of *H. pylori* urease. These results will provide further insights into the design of more potent natural products for eradicating *H. pylori*.




Fig. 1. Two anti-*H. pylori* compounds isolated from *S. williamsii*

Keywords: *Sambucus williamsii*, Viburnaceae, anti-*Helicobacter pylori*, boehmenan, guaiacylglycerol

References

- Xiao, H. H., Zhang, Y., Cooper, R., Yao, X. S., Wong, M. S. 2016. Phytochemicals and potential health effects of *Sambucus williamsii* Hance (Jiegumu). Chin. Med. 11, 1-16.
- Huang, X. -X., Zhou, C. -C., Li, L. Z., Li, F. -F., Lou, L. L., Li, D. -M., Ikejima, T., Peng, Y., Song, S. -J. 2013. The cytotoxicity of 8-O-4' neolignans from the seeds of *Crataegus pinnatifida*. Bioorg. Med. Chem. Lett. 23, 5599-5604.

* Presenting author. E-mail address: amj5812@gnu.ac.kr (Mi-Jeong Ahn)

S1.P215 Lignans from autochthonous Argentinians *Larrea* species: potential use in the control of microorganisms in agriculture

Sarah Tanguy Guillo^{a,*}, Mariana Sanchez Matías^b, Marisol Cristina Jiménez^b, Diego Sampietro^b, Raphaël Grougnet^a, Marina Kritsanida^a

^a PNAS team, CiTCoM laboratory, UMR 8038, 4, avenue de l'observatoire, Université Paris Cité, Paris, France

^b University of Biochemistry, Chemistry and Pharmacy of the National University of Tucuman, Batalla de Ayacucho, Tucuman, Argentina

For decades, the growing global food demand has driven agricultural intensification, necessitating increased output in both crop and livestock production (Tudi *et al.*). Despite advancements in global food safety, crops and livestock remain susceptible to infectious diseases. Ensuring this safety requires the constant control of such illnesses. Hence, the ongoing search for new molecules is crucial to combat pathogens that are increasingly resistant to chemical treatments (Qiu *et al.*). This study aims to discover new antifungal and antibacterial compounds through bioguided isolation from two active plant extracts native to Argentina, *Larrea cuneifolia* and *Larrea divaricata* (Vogt *et al.*). Twenty molecules were isolated, especially lignans, flavonoids and original esters. Seven of them were tested against two *Salmonella* serovars affecting chicken farms and a phytopathogenic fungus, *Fusarium graminearum*. Nordihydroguaiaretic acid (NDGA) demonstrated significant activity against the tested *Salmonella* strains, with a minimum inhibitory dose (MIC) of 1.6 µg, and the (*R-R*) ent of Parakmerin A showed notable activity against the *Fusarium* strain with a MID of 3.1 µg. Isobolographic analysis revealed the synergistic effect of these two molecules with commonly used agricultural controls such as chloramphenicol and tebuconazole. Finally, the content of Ent (*R-R*) of Parakmerin A in *Larrea divaricata* was measured by HPTLC representing 0.90% of the total ethanolic extract.

Keywords: *Larrea*, lignans, antimicrobial activity, isobologram, agricultural control

References

- Qiu, D., Mingjing K., Qi Z., Fan Z., Tao L., Liwei S., et Haifeng Q., 2022. Response of microbial antibiotic resistance to pesticides: An emerging health threat. *Science of The Total Environment* 850, 15805.
- Tudi, M., Huada D. R., Li W., Jia L., Ross S., Des C., Cordia C., et Dung T.P., 2021 Agriculture Development, Pesticide Application and Its Impact on the Environment. *International Journal of Environmental Research and Public Health* 18, n° 3, 1112.
- Vogt, V., Diego C., Carlos T., Liliana S., et Susana R., 2013. Antifungal activity *in vitro* and *in vivo* of extracts and lignans isolated from *Larrea divaricata* Cav. against phytopathogenic fungus. *Industrial Crops and Products* 42, 583- 86.

* Presenting author. E-mail address: sarah.tanguy-guillo@etu.u-paris.fr (Sarah Tanguy Guillo)

S1.P216 Traditional Middle Eastern spice blends (Baharat): Antimicrobial activity, metabolomic profile, and trace element analysis

Mauro Di Stasi^{a,*}, Ata Kaboudari^b, Mancini Simone^c, Veronique Vacchina^d, Alessandra Braca^{a,e,f}, Marinella De Leo^{a,e,f}, Filippo Fratini^{c,e}

^a Department of Pharmacy, University of Pisa, Via Bonanno 33, 56126 Pisa, Italy

^b Department of Food Hygiene and Quality Control, Urmia University, Urmia P.O. Box 1177, Iran

^c Department of Veterinary Science, University of Pisa, Viale delle Piagge 2, 56124 Pisa, Italy

^d UT2A, Ultra-Traces Analyses Aquitaine, 2 avenue du president Angot, 64000 Pau, France

^e Interdepartmental Center NUTRAFOOD “Nutraceuticals and Food for Health”, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

^f CISUP, Centre for Instrumentation Sharing, University of Pisa, Lungarno Pacinotti 43/44, 56126 Pisa, Italy

Spices and aromatic herbs have traditionally served culinary, preservative, and health purposes (Liu et al., 2017). This study assessed Middle Eastern spice blends (Baharat) for antimicrobial activity, comprising black and red pepper, cinnamon, clove, allspice, garlic, mint, black dry lime, turmeric, cumin, and ginger. EtOH:H₂O (3:7) was used to extract the spices individually and in four mixtures (blends 1-4). Blends 2 and 4 exhibited good antibacterial activity through the plate microdilution method (Wiegand et al., 2008) against Gram-positive bacteria (MIC = 0.390 and 3.125 mg/mL against *Staphylococcus aureus* and *Listeria monocytogenes*, respectively) and inhibition against *Enterococcus faecalis* (MIC = 12.5 and 6.25 mg/mL for blends 2 and 4, respectively). A reduced inhibitory activity was found against Gram-negative bacteria, except blend 4 against *Salmonella typhimurium* (MIC = 6.25 mg/mL). A mild bactericidal activity was shown against Gram-positive bacteria (50-200 mg/mL). The MIC values of positive control (gentamycin) ranged 7-28 µL/mL. Blends 2 and 4 underwent metabolomic analysis using UHPLC-HR-Orbitrap/ESI-MS. Altogether, 123 compounds were identified, including phenolic and hydroxycinnamic acids, flavonoids, piperamides, and organosulfur compounds (Ali et al., 2021). Blend 4 also contained nine limonoids exclusive to *Citrus* L. fruits due to black dry lime inclusion. Finally, ICP- QQQ-MS analyzed 12 trace elements in the spices, comparing their concentrations to FAO/WHO limits (FAO/WHO, 2023): Al ≤ 49, Cu ≤ 9.9, Fe ≤ 588, Mn ≤ 484, Zn ≤ 17.4 mg/L; As ≤ 38, Cd ≤ 207, Cr ≤ 636, Hg ≤ 10, Ni ≤ 1266, Pb ≤ 1548, Sn ≤ 60 µL/L.

Keywords: spices, metabolomics, antimicrobial, UHPLC-HR-Orbitrap/ESI-MS, ICP-QQQ-MS

References

- Ali, A., Wu, H., Ponnampalam, E.N., Cottrell, J.J., Dunshea, F.R., Suleria, H.A.R., 2021. Comprehensive profiling of most widely used spices for their phenolic compounds through LC-ESI-QTOF-MS² and their antioxidant potential. *Antioxidants* 10(5), 721.
- FAO/WHO, 2023. Joint FAO/WHO Food Standards Programme Codex Committee on Contaminants in Foods. Working Document for Information and use in discussions related to contaminants and toxins in the GSCTF.
- Liu, Q., Meng, X., Li, Y., Zhao, C.N., Tang, G.Y., Li, H.B., 2017. Antibacterial and antifungal activities of spices. *J. Mol. Sci.* 18(6), 1283.
- Wiegand, I., Hilpert, K., Hancock, R.E., 2008. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat Protoc.* 3(2), 163–175.

* Presenting author. E-mail address: mauro.distasi@phd.unipi.it (Mauro Di Stasi)

S1.P217 Ethnopharmacological survey of medicinal plants traditionally used in Tanzania for the management of sickle cell disease

Godfrey Sambayi^{a,b,*}, Rogers Mwakalukwa^b, Ramadhani Adamu^b, Luc Pieters^a, Emmy Tuenter^a

^a Natural Products & Food Research and Analysis-Pharmaceutical Technology (NatuRAPT), Department of Pharmaceutical Sciences, University of Antwerp, Universiteitsplein 1, B-2610, Antwerp, Belgium

^b Department of Pharmacognosy-School of Pharmacy, Muhimbili University of Health and Allied Sciences, P.O. Box 65013, Dar es Salaam, Tanzania

Introduction: Previous ethnopharmacological and bioactivity studies reported the use of medicinal plants in the management of sickle cell disease (SCD)^{1,2}. *In vitro* and *in vivo* experiments showed anti-sickling, anti-inflammatory, antioxidant, analgesic, anti-anemic and antibacterial activities for these plants³⁻⁵. The use of medicinal plants against infectious and non-infectious diseases was reported, but little is known about medicinal plants used to manage SCD in Tanzania⁶⁻⁸.

Methods: In-person interviews were conducted with traditional health practitioners (THPs) using semi-structured, open-ended questions and medicinal plants traditionally used to manage SCD were documented. The interview was followed by a field visit to collect plant specimens. A comprehensive literature review was performed for each of the medicinal plants mentioned by THPs and little studied plants were selected for further investigation.

Results: In total, 37 participants were interviewed. Most participants had primary education (81%) and mainly gained their knowledge on medicinal plants via revelations (54%). A total of 95 medicinal plants were collected belonging to 35 families. Roots (38%) and leaves (33%) were most frequently used, and decoction was the most common preparation method (56%). The vast majority of herbal treatments was recommended to be taken orally (89%).

Conclusion: This is the first ethnopharmacological survey on medicinal plants used in Tanzania for the management of SCD. The survey reported 95 medicinal plants claimed to have anti-sickling potential. The results provide a starting point in the search for new bioactive phytochemicals for the management of SCD.

Keywords: ethnopharmacology, sickle cell disease, Tanzania

References

- Yembeu, N.L., Biapa Nya, P.C., Pieme, C.A., Tchouane, K.D., Kengne Fotsing, C.B., Nya Nkwikeu, P.J., Feudjio, A.F., Telefo, P.B., 2022. Ethnopharmacological study of the medicinal plants used in the treatment of sickle cell anemia in the West region of Cameroon. *Evid-Based Compl. Alt.* 2022:5098428.
- Amujoyegbe, O.O., Idu, M., Agbedahunsi, J.M., Erhabor, J.O., 2016. Ethnomedicinal survey of medicinal plants used in the management of sickle cell disorder in Southern Nigeria. *J. Ethnopharmacol.* 185, 347–360.
- Azubuike, C.P., Uzoeto, C.A., Igbokwe, N.H., Igwilo, C.I., 2016. In vitro antisickling, antimicrobial and antioxidant potentials of extracts of *Sorghum bicolor* (L) Moench seeds and *Mangifera indica* (L) Anacardiaceae leaves and their formulations. *Journal of Science and Practice of Pharmacy.* 3, 135–144.
- Mishra, S., Sonter, S., Dwivedi, M.K., Singh, P.K., 2021. Anti-sickling, anti-inflammatory and anti-oxidant potential of *Rubia cordifolia* L.: A traditionally used medicinal plant of Amarkantak, Madhya Pradesh (India). *Medicinal Plants – International Journal of Phytomedicines and Related Industries.* 13, 499–514.
- Fulata, A.M., Usman, H., Tijjani, M.A., Daja, A., Bukar, A., 2022. Phytochemical, anti-sickling and polymerization inhibition studies of the crude methanol leaf recipe extracts of three medicinal plants. *Int. J. Adv. Chem. Res.* 4, 42–47
- Peter, E.L., Rumisha, S.F., Mashoto, K.O., Malebo, H.M., 2014. Ethno-medicinal knowledge and plants traditionally used to treat anemia in Tanzania: A cross sectional survey. *J. Ethnopharmacol.* 154, 767–773.
- Marealle, A.I., Innocent, E., Marobela, K.A., Qwarse, M., Moshi, M., 2020. Ethnopharmacological survey of medicinal plants used in the management of HIV and AIDS-related conditions in Makete district, Southern highlands of Tanzania. *European J. Med. Plants.* 31, 56–83.
- Maregesi, S.M., Messo, C.W., Mathias, J., 2017. Ethnomedicinal survey and safety evaluation of traditional eye medicines used in Misungwi district, Tanzania. *J. Intercult. Ethnopharmacol.* 6, 75–83.

* Presenting author. E-mail address: godfrey.sambayi@student.uantwerpen.be (Godfrey Sambayi)

S1.P218 Bio-elicitation of isoquinoline alkaloids in *Chelidonium majus* in vivo & in vitro

Weronika Kozłowska^a, Anna Sroka-Bartnicka^{b,c}, Mikołaj Krysa^c, Malwina Brożyna^d, Adam Junka^d, Magdalena Dziągwa-Becker^e, Adam Matkowski^{f,g}, Sylwia Zielińska^{a,*}

^a Department of Pharmaceutical Biology and Biotechnology, Division of Pharmaceutical Biotechnology, Wrocław Medical University, Borowska 211, 50-556 Wrocław, Poland

^b Department of Genetics and Microbiology, Institute of Biological Sciences, Maria Curie-Skłodowska University, Akademicka 19, 20-033 Lublin, Poland

^c Independent Unit of Spectroscopy and Chemical Imaging, Medical University of Lublin, Chodzki 4a, 20-093 Lublin, Poland

^d Platform for Unique Models Application (PUMA), Department of Pharmaceutical Microbiology and Parasitology, Wrocław Medical University, Borowska 211 50-556 Wrocław, Poland

^e Department of Weed Science and Tillage Systems, Institute of Soil Science and Plant Cultivation State Research Institute, Orzechowa 61, 50-540 Wrocław, Poland

^f Department of Pharmaceutical Biology and Biotechnology, Division of Pharmaceutical Biology and Botany, Wrocław Medical University, Borowska 211A, 50-556 Wrocław, Poland

^g Botanical Garden of Medicinal Plants at the Department of Pharmaceutical Biology and Biotechnology, Kochanowskiego 10-12-14, 51-601 Wrocław, Poland

Cell culture technology and hydroponic systems were used for the examination of *Chelidonium majus* metabolic response to biotic stimuli. In *in vitro* experiments, *C. majus* cells were cultured on *Komagataeibacter xylinus* derived non-purified nanocellulose matrices (BNC) (Zielińska et al., 2022). Three microbial strains (*Candida albicans*, *Staphylococcus aureus* and *Pseudomonas aeruginosa*) were used as bio-elicitors and were inoculated at the top or the bottom of cellulose discs. In turn, in the hydroponic plant cultivation *in vivo* system, *P. aeruginosa* at three different concentrations (1, 10, 50 ml/L) as well as methyl jasmonate (50 ml/L) were used. After the biotic elicitors treatment the isoquinoline alkaloid profile in plant material was examined using LC-MS/MS and MALDI-MS imaging. MALDI MSI chemical maps showed higher content of coptisine, sanguinarine, berberine, chelerythrine, chelidonine and allocryptopine in BNC containing *S. aureus* compared to *C. albicans* and *P. aeruginosa* or elicitor-free cellulose carriers. In hydroponic experiments, the concentration of 10 ml/L of lyophilized *P. aeruginosa* was the most effective in terms of protoberberine and benzophenanthridine derivatives' production in the *C. majus* plants. Further investigation on the individual microbial strains effect including the concentration and the exposure time, both for plants grown *in vitro* and *in vivo* is required to understand their possible mechanism of action and specialized metabolites production control.

Keywords: isoquinoline alkaloids, *Chelidonium majus*, biotic elicitors

References

- [1] Zielińska S, Matkowski A, Dydak K, Czerwińska ME, Dziągwa-Becker M, Wójciak M, Sowa I, Fijałkowski K, Ciecholewska-Juśko D, Broda M, Gorczyca D, Junka A. Bacterial nanocellulose fortified with antimicrobial and anti-inflammatory natural products from *Chelidonium majus*. *Materials*, 2022; 15,16.

Acknowledgments

The study was funded by the National Science Centre (NCN) - SONATA 15, 2019/35/D/NZ7/00266.

* Presenting author. E-mail address: sylwia.zielinska@umw.edu.pl (Sylwia Zielińska)

S1.P219 *In vitro* anti-*Helicobacter pylori* activity of extracts and essential oil of *Thymus pannonicus* All. (Lamiaceae)

Jelena Arsenijević^{a,*}, Francesca Sisto^b, Milica Drobac^a, Ivan Šoštarić^c, Zoran Maksimović^a, Nada Kovačević^a

^a University of Belgrade – Faculty of Pharmacy, Department of Pharmacognosy, Vojvode Stepe 450, 11221 Belgrade, Serbia

^b Department of Biomedical, Surgical and Dental Sciences, University of Milan, Via Pascal 36, 20133 Milan, Italy

^c University of Belgrade – Faculty of Agriculture, Department of Crop and Vegetable Science, Nemanjina 6, Zemun, 11080 Belgrade, Serbia

Thymus pannonicus All. is an aromatic herbaceous plant distributed in Central and Eastern Europe (Jalas, 1972). The aerial parts of the lemon-scented chemotype from Serbia are considered a traditional remedy for various gastrointestinal complaints (Maksimović et al., 2008). In this work, we tested its antibacterial activity against *Helicobacter pylori*, a bacterium associated with the risk of developing complicated chronic diseases of the gastrointestinal tract.

T. pannonicus hexane (HE), dichloromethane (DE), and methanol extracts (ME), the fraction of diethyl ether extract soluble in 80% aqueous methanol (FDM), and essential oil (EO), were tested by broth micro-dilution method against *H. pylori* ATCC 43504 (reference strain) and 10 clinical isolates, including strains resistant to metronidazole and/or clarithromycin.

ME showed the best activity, with MIC₅₀ (minimum concentration that inhibits the growth of 50% of the tested strains) and MIC₉₀ of 4 and 16 µg/mL, respectively. FDM exhibited MIC₅₀ 16 µg/mL and MIC₉₀ 32 µg/mL, while EO showed MIC₅₀ 2 µg/mL and MIC₉₀ 64 µg/mL. HE and DE were less active (MIC₅₀ and MIC₉₀ were 32 and 128 µg/mL, respectively).

Liquid chromatography with ultraviolet and mass spectrometric detection revealed that ME was characterized by a high content of rosmarinic acid (64 mg/g) and salvianolic acid H, while FDM was rich in methylated flavonoid aglycons. The dominant compounds in EO, analyzed by gas chromatography with mass spectrometry and flame-ionization detectors, were geranial and neral (36.7% and 27.2%, respectively).

The results indicate that preparations of *T. pannonicus* may be potential therapeutic agents against *H. pylori* infections.

Keywords: *Thymus*, *Helicobacter pylori*, rosmarinic acid, geranial, neral

References

Jalas, J., 1972. *Thymus* L. In: Tutin, T.G., Heywood, V.H., Burges, N.A., Moore, D.M., Valentine, D.H., Walters, S.M. et al. (Eds.), *Flora Europaea*, Volume 3. Cambridge University Press, Cambridge, pp. 172–182.

Maksimović, Z., Milenković, M., Vučićević, D., Ristić, M., 2008. Chemical composition and antimicrobial activity of *Thymus pannonicus* All. (Lamiaceae) essential oil. *Cent. Eur. J. Biol.* 3, 149–154.

* Presenting author. E-mail address: jelena.arsenijevic@pharmacy.bg.ac.rs (Jelena Arsenijević)

S1.P220 Assessing the structural interaction profile differences between natural products and synthetics, a data-driven approach

Yojana Gadiya*, António José Preto, Christoph A. Krettlér, David Healey, Daniel Domingo-Fernández, Viswa Colluru

Enveda Biosciences, 5700 Flatiron Parkway, Boulder, CO, 80301, USA

The potential of natural products (NPs) is a largely untapped source when comparing its prevalence in the drug market with that of its synthetic counterparts. By leveraging structure-based drug design (SBDD), we explore the coupling of conventional computational tools and the impact of state-of-the-art artificial intelligence (AI)-based tools. Moreover, NPs have in the past years, demonstrated increased efficacy and potency. In our work, we aimed to elucidate the target promiscuity profiles of synthetic and NP compounds to shed light on the rationale behind NPs' selection as primary candidates for drug discovery. To achieve this, we conducted a comprehensive comparative structure analysis of the two compound groups (i.e. NPs and synthetics) targeting similar biological targets. By systematically analyzing the interactions between synthetic and NP compounds and their respective targets, we sought to uncover insights into the unique properties and mechanisms underlying NP efficacy and potency. Our findings contribute to a deeper understanding of the molecular interactions and functional characteristics of synthetic and NP compounds, providing valuable insights for optimizing drug discovery strategies and harnessing the therapeutic potential of NPs. Further research in this area holds promise for the development of novel therapeutics and the advancement of drug discovery efforts.

Keywords: structure-based drug design (SBDD), in-silico computation, natural products, small molecules, protein-ligand interaction (PLI)

* Presenting author. E-mail address: yojana.gadiya@envedabio.com (Yojana Gadiya)

S1.P221 Guttiferones-targeted dereplication integrating LC-HRMS/MS and ^{13}C -NMR data: A comparative metabolomic study of *Symphonia globulifera* latex

David Hozain^{a,*}, Raimundo Goncalves De Oliveira Junior^a, Annabelle Dugay^a, Florence Souquet^a, Gilles Pakora^b, Séverine Derbre^c, Leandro Rocha^d, Lucas Silva Abreu^d, Massuo Jorge Kato^e, Didier Buisson^f, Marie-Christine Lallemant^a

^a UMR CNRS 8038 CiTCoM, Paris Cité University, 75006 Paris, France

^b Biology and Health Laboratory, Félix Houphouët-Boigny University, 22 BP 582, Abidjan 22, Ivory Coast

^c SONAS, EA921, Faculty of Health Sciences, Department of Pharmacy, 49045 Angers, France

^d Fluminense Federal University, Niterói, RJ, 24241-000, Brazil

^e Institute of Chemistry, University of São Paulo, São Paulo, 05508-000, Brazil

^f UMR CNRS 7245 MCAM, National Museum of Natural History, 75005 Paris, France

Symphonia globulifera is a Clusiaceae species, mostly distributed in South America and Africa. This medicinal plant has been the focus of phytochemical studies leading to the identification of several PPAPs. Guttiferones are the main PPAPs isolated from *S. globulifera* latex, exhibiting various pharmacological properties such as antiparasitic, antimicrobial, antiviral, and cytotoxic activities (Conceição et al., 2023). To gain access to this natural resource, our team has established a collaboration with French Guiana (FG), Ivory Coast (IC) and Brazil (BR), to conduct a comparative metabolomics study. Latex was collected from these three regions, and guttiferones were extracted by maceration (48h), using methanol. Extracts were then analyzed by LC-HRMS/MS and a molecular network (MN) was constructed using GNPS and Cytoscape[®] software. The MN revealed a main cluster for guttiferones with m/z 602 or 670, corresponding to guttiferones A and C and their isomers, respectively. Due to the presence of numerous stereoisomers, a very similar MS/MS fragmentation pathway is often observed, limiting their putative identification. We then proceed with ^{13}C -NMR dereplication experiments using MixONat[®] software (Bruguère et al., 2020). Latex spectral data were compared with predicted NMR data from a Clusiaceae database (DB) including 1992 natural compounds described on LOTUS. This MS/MS-NMR integrated approach allows the annotation of guttiferones C and D (m/z 670) in all samples, suggesting these are important chemical markers for *S. globulifera*. Guttiferone A (m/z 602) was found in FG latex, while guttiferone E/xanthochymol were identified in BR latex. Phytochemical investigations are ongoing in order to isolate the unknown guttiferones.

Keywords: Clusiaceae, dereplication, guttiferones, PPAPs (PolycyclicPolyprenylated Acylphloroglucinols)

References

- Bruguère, A.; Derbré, S.; Dietsch, J.; Leguy, J.; Rahier, V.; Pottier, Q.; Bréard, D.; Suor-Cherer, S.; Viault, G.; Le Ray, A.-M.; Saubion, F.; Richomme, P. MixONat, a Software for the Dereplication of Mixtures Based on ^{13}C NMR Spectroscopy. *Anal. Chem.* 2020, 92 (13), 8793–8801.
- Conceição, M.; Beserra, F. P.; Aldana Mejia, J. A.; Caldas, G. R.; Tanimoto, M. H.; Luzenti, A. M.; Gaspari, P. D. M.; Evans, N. D.; Bastos, J. K.; Pellizzon, C. H. Guttiferones: An Insight into Occurrence, Biosynthesis, and Their Broad Spectrum of Pharmacological Activities. *Chem. Biol. Interact.* 2023, 370, 110313.

* Presenting author. E-mail address: david.hozain@u-paris.fr (David Hozain)

S1.P222 QSAR-derived opportunities of prenylated phenolic compounds from Fabaceae as antimicrobial phytochemicals

Sylvia Kalli^{a,b}, Sarah van Dinteren^a, Jos Hageman^c, Jean-Paul Vincken^a, Carla Araya-Cloutier^{a,*}

^a Laboratory of Food Chemistry, Wageningen University and Research, Wageningen, The Netherlands

^b Wageningen Food Safety Research, Wageningen, The Netherlands

^c Biometris, Applied Statistics, Wageningen University & Research, Wageningen, The Netherlands

Prenylated phenolic compounds (Fabaceae), bearing C5-isoprene moieties, are antimicrobial phytochemicals known for their wide structural diversity (Al-Maharik, 2019). Evidence shows that even minor structural variations can greatly influence their antimicrobial properties (Araya-Cloutier et al., 2018).

This presentation aims to provide an overview of opportunities identified in recent structure-activity relationships (SARs) studies of prenylated phenolics as antimicrobial agents. Using a diverse collection of prenylated phenolics, tested against various microorganisms, including healthcare-associated pathogen methicillin-resistant *Staphylococcus aureus* (MRSA) and oral pathogen *Streptococcus mutans*, SARs were defined and in-silico quantitative SARs (QSAR) models developed and validated. The models were compared to unravel the most important molecular properties for antimicrobial activity of prenylated phenolics.

SARs showed that the number of prenyl groups attached to the backbone influences the spectrum of activity of these phytochemicals, i.e., diprenylated phenolics were among the most active antibacterials against Gram positive bacteria, whereas monoprenylated phenolics showed superior activity against yeast or Gram negative bacteria *Escherichia coli* (for the latter one in the presence of an efflux pump inhibitor). The backbone of the compounds was also shown to play a major role in the SARs; e.g. C-ring opening of prenylated flavanones, resulting in prenylated chalcones, significantly restored antimicrobial activity.

Based on the QSAR models, extent of hydrophobicity, shape, flexibility, charge and hydrogen-bonding were identified as critical contributors of antimicrobial activity of prenylated phenolics. Overall, prenylated phenolics are potent natural antimicrobials and the developed QSAR models are valuable tools for gaining insights into the complex SARs governing their activity.

Keywords: Prenylated phenolic compounds, Fabaceae, Structure-Activity Relationships, QSAR modelling

References

Al-Maharik, N. (2019) Isolation of naturally occurring novel isoflavonoids: an update. Nat. Prod. Reports 36:8, p. 1156- 1195.

Araya-Cloutier, C., Vincken, J.-P., Schans, M.G.M. van de, Hageman, J., Schaftenaar, G., Besten, H.M.W. den, and Gruppen, H. (2018) QSAR-based molecular signatures of prenylated (iso)flavonoids underlying antimicrobial potency against and membrane-disruption in Gram positive and Gram negative bacteria. Sci. Rep. 8, 9267.

Kalli, S., Araya-Cloutier, C., Hageman, J., and Vincken, J.-P. (2021) Insights into the molecular properties underlying antibacterial activity of prenylated (iso)flavonoids against MRSA. Sci. Rep. 11, 14180.

* Presenting author. E-mail address: carla.arayacloutier@wur.nl (Carla Araya-Cloutier)

S1.P223 Phytochemistry and biological activities of *Ostericum palustre* (*Angelica palustris*) (Apiaceae)

Grażyna Łaska^{a,*}, Elwira Sieniawska^b, Magdalena Maciejewska-Turska^c, Łukasz Świątek^d, Marcin Stocki^e, Mirosław Angielczyk^f

^a Department of Agri-Food Engineering and Environmental Management, Białystok University of Technology, 15-351 Białystok, Poland

^b Department of Natural Products Chemistry, Medical University of Lublin, 20-093 Lublin, Poland

^c Department of Pharmacognosy with Medicinal Plant Unit, Medical University of Lublin, 20-093 Lublin, Poland

^d Department of Virology with Viral Diagnostics Laboratory, Medical University of Lublin, 20-093 Lublin, Poland

^e Department of Silviculture and Forest Utilization, Białystok University of Technology, 15-351 Białystok, Poland

^f Podlaski Botanical Garden, Koryciny 73b, 17-315 Grodzisk, Poland

The genus *Angelica* has a Euroasiatic-continental range that includes several Central European countries (Dittbrenner et al. 2005). *Ostericum palustre* Besser (= *Angelica palustris* (Besser) Hoffm.) is a representative of genus *Angelica*. In Poland, this species has been subject to strict and active species protection since 2001 (Journal of Laws 2014, item 1409). The species is a rhizomatous perennial (hemicryptophyte) herb 140-160 cm high that prefers humid and eutrophic habitats on mineral substrates. The genus *Angelica* is indicated as a pharmaceutical raw material due to the richness of chemical compounds with anti-allergic, anti-cancer, and anti-inflammatory properties and is used in the treatment of diabetes, rheumatism, injuries, burns, and blood deficiencies (Patents 2012, 2014, 2015, 2016). This presentation will cover the phytochemical analysis and biological evaluation of the extracts prepared from aerial and underground parts of this species.

The antimalarial activity was determined against the chloroquine-sensitive (D6) strain of *Plasmodium falciparum* by measuring plasmodial LDH activity according to the procedure of Makler and Hinrichs (1993). Laboratory tests showed high biological activity of the methanol extract from the aerial part (stems, leaves, and flowers) of this species, which inhibited the growth of the malaria-causing strain D6 of *Plasmodium falciparum* in 41%. In comparison, chloroquine (positive control) showed 94–98% inhibition. Evaluation of cytotoxicity was conducted using selected normal and cancer cell lines.

Among the secondary metabolites of *O. palustre*, coumarins, organic acids (angelic, valeric, succinic, malonic, oxalic, citric, fumaric), phenolics (coumaric, feruloylquinic, caffeoylquinic and di-caffeoylquinic acids, luteolin rutoside) and terpenes (cymene, camphene, limonene) were identified.

Keywords: *Ostericum palustre*, Apiaceae, antimalarial activity, *Plasmodium falciparum*, cytotoxicity

References

- Dittbrenner, A., Hensen, I., Wesche, K., 2005. Genetic structure and random amplified polymorphic DNA diversity of the rapidly declining *Angelica palustris* (Apiaceae) in Eastern Germany in relation to population size and seed production. *Plant Species Biol* 20:191–200.
- Makler, M.T., Hinrichs, A.D., 1993. Measurement of the Lactate Dehydrogenase activity of *Plasmodium falciparum* as an assessment of parasitemia. *Am J Trop Med Hyg* 48(2): 205–210.
- Patent No CN 102755500 A 20121031; Patent No CN 104056084 A 20140924; Patent No CN 104623155 A 20150520; Patent No CN 105213835 A 20160106.
- Rozporządzenie Ministra Środowiska z dnia 9 października 2014 roku w sprawie ochrony gatunkowej roślin. *Dz.U.* 2014, poz. 1409.

* Presenting author. E-mail address: g.laska@pb.edu.pl (Grażyna Łaska)

S1.P224 Discovery of two new bicyclic macrolides from *Kitasatospora acidiphila*

Yegang Hur^{a,*}, Joon Soo An^a, Seung Bum Kim^b, Dong-Chan Oh^a

^a Natural Products Research Institute, College of Pharmacy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

^b Department of Microbiology and Molecular Biology, College of Bioscience and Biotechnology, Chungnam University, 99 Daehak-ro, Daejeon, 34134, Republic of Korea

Forests are an under-explored habitat for chemically prolific actinomycetes (Srivastava et al., 2019). Because rare actinomycete phylogenetic groups, besides commonly studied *Streptomyces*, are recognized as important sources of novel bioactive, we prioritized rare actinomycetes strains from forest soil for deeper chemical analysis (Tiwari et al., 2012).

The MMS16-CNU292 strain was isolated from a soil sample collected from the surface of a pine forest in Daejeon City, Republic of Korea. It has been identified as a new species of the genus *Kitasatospora*, and named *Kitasatospora acidiphila* based on its phylogenetic, chemotaxonomic, and phenotypic properties (Kim et al., 2020).

Cultivation and LC/MS-based chemical analysis of *K. acidiphila* MMS16-CNU292 enabled the detection of two previously unreported compounds. Chromatographic purification and spectroscopic analysis based on 1D/2D NMR, HR-ESI-MS, and UV data elucidated the planar structures of the new natural products as unique bicyclic macrolides bearing 14- and 6-membered rings. Further analysis of ³J_{H,H} and ROESY correlations determined the geometry of two *trans-trans* dienes in the new macrolides.

Based on *J*-based configuration analysis, ROESY correlations, and acetonide derivatization of an 1,3-diol (Rychnovsky et al., 1997), the relative stereochemistry of the new polyketides was established. The determination of the absolute configuration of the compounds is in progress based on multi-step chemical derivatizations, including Mosher's esterification.

Keywords: *Kitasatospora acidiphila*, rare actinomycetes, macrolide, structure elucidation

References

- Srivastava, N.; Nandi, I.; Ibeyama, A.; Gupta, S.; Sarethy, I. P., 2019. Microbial diversity of a Himalayan forest and characterization of rare actinomycetes for antimicrobial compounds. 3 Biotech. 9, 27.
- Tiwari, K.; Gupta, R. K., 2012. Rare actinomycetes: a potential storehouse for novel antibiotics. Crit. Rev. Biotechnol. 32, 108-132.
- Kim, M. J.; Roh, S. G.; Kim, M.-K.; Park, C.; Kim, S. B., 2020. *Kitasatospora acidiphila* sp. nov., isolated from pine grove soil, exhibiting antimicrobial potential. Int. J. Syst. Evol. Microbiol. 70, 5567-5575.
- Rychnovsky, S. D.; Richardson, T. I.; Rogers, B. N., 1997. Two-Dimensional NMR Analysis of Acetonide Derivatives in the Stereochemical Assignment of Polyol Chains: The Absolute Configurations of Dermostatins A and B. J. Org. Chem. 62, 2925-2934.

* Presenting author. E-mail address: whuyakang@snu.ac.kr (Yegang Hur)

S1.P225 The potential of curry leaf extract in controlling SARS-CoV-2 infectious syndromes

Mohamed Abbas Umare Faruk, Michal Korinek, Hui-Chun Wang*

Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung, 80708, Taiwan

Coronavirus disease 2019 (COVID-19) is caused by infection of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), a worldwide pandemic with asymptomatic, mild or severe symptoms. The severe symptoms linked to a massive viral load may induce a severe immune response that causes tissue damage (Purbey et al., 2023). Evidence supports that several eatable spices, including curry leaves, could potentially control the cytokine storm of lung infectious diseases (Kunnumakkara et al., 2021). This study aims to identify the active components of a spice curry leaf (*Murraya koenigii*) that can prevent SARS-CoV-2 infection and its related allergy response. The assays of cell-free spike RBD-ACE2 binding and calcium ionophore A23187-induced mast cell degranulation guided the plant extraction. Curry leaves were initially extracted using 95% ethanol and partitioned by ethyl acetate and water with 1% HCl. The n-hexane and methanol subjects partitioned the ethyl acetate (non-alkaloids) fraction, which showed potent inhibition of spike-ACE2 binding. The most bioactive active hexane extract was fractionated using MPLC column chromatography. The acid water fraction was partitioned by dichloromethane (DCM) and water with ammonium hydroxide, and the DCM fraction with anti-allergy activity was further fractionated. The current evidence supports that leaf extract fractions of non-alkaloid have potential activities to inhibit spike-ACE2 protein binding by a concentration range of ng/mL, and alkaloid fractions have the potential to inhibit cellular allergy response by a concentration range of µg/mL.

Keywords: Curry leaf (*Murraya koenigii*), SARS-CoV-2

References

- Purbey P.K., Roy K., Gupta S., Paul M.K., 2023 Mechanistic insight into the protective and pathogenic immune-responses against SARS-CoV-2. *Mol Immunol.* 156:111-126.
- Kunnumakkara A.B., Rana V., Parama D, Banik K., Girisa S., Henamayee S., Thakur K.K., Dutta U., Garodia P., Gupta S.C., Aggarwal B.B., 2021 COVID-19, cytokines, inflammation, and spices: How are they related? *Life Sci.* 284:119201.

* Presenting author. E-mail address: wanghc@kmu.edu.tw (Hui-Chun Wang)

S1.P226 Antiviral effect of *Hoveniae Semen Seu Fructus* against Influenza A viral infection

Won-Kyung Cho*, Min-Ho Cha, Nam Hui Yim, Hee-Jeong Choi, Jin Yeul Ma

Korean Medicine Application Center, Korea Institute of Oriental Medicine, Daegu, South Korea, 41062

Hoveniae Semen Seu Fructus is the fruit and seed of *Hovenia dulcis* Thunb. Here, we first demonstrate that *Hoveniae Semen Seu Fructus* water extract contains a potent anti-influenza A virus effect via inhibition of neuraminidase and hemagglutinin and a direct virucidal effect. We investigated the effect of *Hoveniae Semen Seu Fructus* on Influenza A virus infection using the GFP- tagged Influenza A virus. FACS analysis and fluorescent microscopy showed a dose-dependent inhibitory effect of *Hoveniae Semen Seu Fructus* against Influenza A virus infection. *Hoveniae Semen Seu Fructus* also suppressed the cytopathic effects of H1N1 and H3N2 Influenza A virus. Immunofluorescence and qPCR analyses confirmed that *Hoveniae Semen Seu Fructus* significantly represses Influenza A virus protein and RNA expressions. Time of addition and plaque assays showed that *Hoveniae Semen Seu Fructus* changes hemagglutinin and causes virucidal effects early on in the viral replication process. Furthermore, *Hoveniae Semen Seu Fructus* inhibited the neuraminidase activity of H1N1 and H3N2 Influenza A virus. Quercetin, not taxifolin in *Hoveniae Semen Seu Fructus*, inhibited Influenza A virus infection. Our results suggest that *Hoveniae Semen Seu Fructus* could be developed as a natural antiviral agent to prevent Influenza A virus infection.

Keywords: *Hoveniae Semen Seu Fructus*, Influenza A virus, hemagglutinin, neuraminidase, quercetin

* Presenting author. E-mail address: wkcho@kiom.re.kr (Won-Kyung Cho)

S1.P227 *Thuja orientalis Folium* contains antiviral effect against Influenza A viral infection via modulating hemagglutinin and neuraminidase

Myong-Min Lee, Won-Kyung Cho, Min-Ho Cha, Nam Hui Yim, Jin Yeul Ma*

Korean Medicine Application Center, Korea Institute of Oriental Medicine, Daegu, South Korea, 41062

Thuja orientalis Folium (TOF) has been traditionally used as an expectorant for inflammatory airway disease. In the present study, we examined the antiviral activity of TOF using GFP-tagged influenza A virus (IAV). The fluorescence microscopy and fluorescence-activated cell sorting analysis showed that TOF potently inhibits IAV infection, dose-dependently. Consistent with that, immunofluorescence and Q-PCR analysis results confirmed TOF significantly reduces IAV protein and RNA expression. TOF blocked IAV infection at the binding and entry step upon viral infection and interfered with HA protein. Further, TOF exhibited a direct virus-killing effect and inhibited the neuraminidase activity of IAV. Additionally, TOF prevented the cytopathic effect caused by H1N1 and H3N2 IAV infection. Amentoflavone among the constituents in TOF exerted the strongest anti- IAV effect. Myricetin, quercetin, and quercitrin in TOF also inhibited IAV infection. However, the potent anti-IAV effect of TOF may be related to the synergistic effect of constituents, not by a single specific compound. Our results suggest TOF exhibits a significant inhibitory effect against IAV infection at multi-stages via the blockage of viral attachment and entry, inhibition of neuraminidase, and induction of virucidal effects.

Keywords: *Thuja orientalis Folium*, influenza A virus, virucidal effect, hemagglutinin, neuraminidase

* Presenting author. E-mail address: jyma@kiom.re.kr (Jin Yeul Ma)

S1.P228 Exploring the phytochemistry, antibacterial and anti-inflammatory activities of three Ericaceous species native to Ireland

Zeeshan Anwar^a, Gaia Scalabrino^{a,*}, Ismael Obaidi^a, Mohammed A. Selo^a, Kavita Gadar^b, Ronan McCarthy^b, Jandirk Sendker^c, Helen Sheridan^a

^a NatPro Centre, School of Pharmacy and Pharmaceutical Sciences, Trinity College Dublin, Dublin 2, Ireland

^b College of Health and Life Sciences, Brunel University London, UK

^c University of Münster, Institute of Pharmaceutical Biology and Phytochemistry, Münster, Germany

Plant derived natural products have promising therapeutic candidates with multifaceted bioactivities. As part of the 'Unlocking Nature's Pharmacy from Bogland Species' (UNPBS) project, we have investigated the chemistry and biology of a range of bogland species. The present study investigates three Ericaceous species (Heathers) native to Ireland which grow abundantly across raised bogs and mountain heath. We report the antibacterial and anti-inflammatory activities of *Erica cinerea* L., *Erica tetralix* L., and *Calluna vulgaris* (L.) Hull. toward the highly virulent and antibiotic resistant bacterial ESKAPE pathogens *E. faecalis*, *S. aureus*, *A. baumannii* and *E. cloacae*. *C. vulgaris* demonstrated the most potent antibiofilm activity reducing *A. baumannii* biofilm formation by 38%, at a concentration of 0.05 mg/ml. The effects of extracts on Interleukin 6 (IL6), RANTES (CCL5), TNF α and IFN β were evaluated in the THP-1 cell line, after stimulation with LPS. All three extracts significantly reduced the LPS induced cytokine release, most notably of IL6 at a subtoxic concentration range of 10-100 μ g/mL.

Chemical fingerprints of the extracts were generated by LCMS, GCMS, HPTLC (normal and reverse phase) and NMR. Bioactivity guided fractionation of *E. cinerea* was carried out and the most bioactive fraction 'S5' was shown to contain high concentrations of scopoletin, protocatechuic acid, p-coumaric acid along with isofraxidin, ferulic acid, caffeic acid and ursolic acid.

This research highlights the significance of three Irish heathers. From the findings it can be concluded that the three plants have antibacterial, antibiofilm and anti-inflammatory potential and serve as promising candidates for further development of novel therapeutic or cosmetic agents.

Keywords: Ericaceous species, antibacterial, anti-inflammatory, IL6, phytochemical profile

* Presenting author. E-mail address: scalabrg@tcd.ie (Gaia Scalabrino)

S1.P229 Funicone-like compounds: potential antiviral agents towards canine coronavirus

Maria Michela Salvatore^a, Luca Del Sorbo^b, Rosario Nicoletti^a, Marina DellaGreca^a, Valentina Iovane^c, Filomena Fiorito^b, Anna Andolfi^a

^a Department of Chemical Sciences, University of Naples Federico II, via Cinthia 4, 80126 Naples, Italy

^b Department of Veterinary Medicine and Animal Production, University of Naples Federico II, 80137 Naples, Italy

^c Department of Agricultural Sciences, University of Naples Federico II, Portici, via Università 100, 80055 Portici, Italy

The exploitation of bioactive properties of microbial products is of great pharmaceutical interest. Fungi are a promising source of novel drugs with a broad spectrum of activities and functions, including antiviral properties. Funicone-like compounds, as well as related compounds, constitute a homogeneous group of fungal polyketides possessing remarkable biological activities which have promoted their consideration as drug possibilities (Salvatore et al. 2022). This group of secondary metabolites is essentially produced by species in the genus *Talaromyces* (Eurotiales: *Trichocomaceae*). This communication provides interesting findings concerning the antiviral properties of three well-known members of this group (i.e., 3-O-methylfunicone (1), penisimplicissin (2) and vermistatin (3) (Fig.1) obtained from an isolate of *Talaromyces pinophilus*, previously recovered from the rhizosphere of tobacco (*Nicotiana tabacum*) (Salvatore et al. 2018). In particular, the antiviral effects of 1-3 were investigated on canine coronavirus (CCoV) infection. The virus, an alphacoronavirus, causes self-limiting enteric disease in dogs, especially in puppies but, due to noteworthy plasticity of CoVs, mutation and recombination processes occur generating new dangerous variants. In this context, developing antiviral compounds certainly may have potential applications in the treatment of CoVs infection. Thus, during CCoV infection in canine fibrosarcoma cells (A72), a fibrosarcoma cell line suitable for investigating CCoV, a non-toxic dose of these funicone-like compounds markedly increased features of cell viability and reduced virus yield. In addition, we observed that these compounds caused a strong inhibition in the expression of the aryl hydrocarbon receptor (AhR), a ligand-activated transcription factor which is activated during CCoV infection.




Fig 1. Structures of 3-O-methylfunicone, penisimplicissin and vermistatin (1-3).

Keywords: coronavirus, *Talaromyces*, fungal secondary metabolites, metabolomics

References

- Salvatore, M.M., DellaGreca, M., Andolfi, A. and Nicoletti, R., 2022. New insights into chemical and biological properties of funicone-like compounds. *Toxins*, 14, 466.
- Salvatore, M.M., DellaGreca, M., Nicoletti, R., Salvatore, F., Vinale, F., Naviglio, D. and Andolfi, A., 2018. Talarodioidide, a new 12-membered macrodiolide, and GC/MS investigation of culture filtrate and mycelial extracts of *Talaromyces pinophilus*. *Molecules*, 23,950.

* Presenting author. E-mail address: mariamichela.salvatore@unina.it (Maria Michela Salvatore)

S1.P230 Efficacy of food waste as a potential source of antibacterial agents and antibiotic adjuvants against MRSA and MSSA isolates

James Blee, Peter O'Hara, Thomas J. Smyth, Owen Kenny

Department of Health & Nutritional Sciences, Atlantic Technological University, Sligo Campus, Sligo, Ireland

Staphylococcus aureus is a leading cause of nosocomial and community acquired infections, with up to 50% of isolates exhibiting methicillin resistance (European Centre for Disease Prevention and Control, 2014). Methicillin-resistant *Staphylococcus aureus* (MRSA) infection is associated with increased healthcare costs and higher mortality rates, with over fifty thousand MRSA associated deaths reported in the WHO European region for 2019 (O'Neill, 2014). In this study, the antibacterial properties of extracts, derived from thirteen fruit and vegetable waste streams, were investigated against clinical MRSA and MSSA isolates, using a broth microdilution assay. Of the 65 crude extracts tested, 11 were shown to have activity against both strains at minimum inhibitory concentrations (MICs) ≤ 1 mg/mL. Crude extracts from onion skins (0.25 mg/mL), kiwi skins (0.8 mg/mL) and broccoli stalks (0.8 mg/mL) demonstrated the greatest antibacterial activity. Extracts were also evaluated for their efficacy as synergistic agents with several beta-lactam antibiotics against clinical MRSA isolates. The strongest antibacterial effect was shown from the combination of the onion skin extract with amoxicillin, resulting in a twenty-five-fold reduction of its MIC (0.31 μ g/mL) and a Fractional Inhibitory Concentration Index of

0.29. Total phenolic content analysis of the onion skin extract ($262.39 \pm$

0.05 mg GAE/g DW suggested that these compounds may play a key role in its outlined bioactivity. Overall, the results of this study demonstrate the potential of food waste revalorisation as a viable strategy for combatting the rise of AMR. Studies regarding the antibacterial mechanism of action and identification of bioactive compound(s) are currently ongoing.

Keywords: antimicrobial resistance, methicillin-resistant *Staphylococcus aureus*, food waste, antibiotic modulation

References

European Centre for Disease Prevention and Control., 2014. Antimicrobial resistance surveillance in Europe 2013. Annual Report of the European Antimicrobial Resistance Surveillance Network (EARS-Net). Stockholm: ECDC. O'Neill J. I. M., 2014. Review on Antimicrobial Resistance: Tackling a crisis for the health and wealth of nations. London: Review on Antimicrobial Resistance.

* Presenting author. E-mail address: s00171423@atu.ie (James Blee)

S1.P231 Coumarins of *Angelica archangelica* roots inhibit efflux pumps in *Escherichia coli*

Vanessa Baumgartner*, Franz Bucar

University of Graz, Institute of Pharmaceutical Sciences, Department of Pharmacognosy, Beethovenstraße 8, 8010 Graz, Austria

Angelica archangelica L. (Apiaceae), native and widespread in Europe, has been cultivated as a medical plant with antibacterial, antispasmodic and antianxiety properties. *A. archangelica* has a complex chemical composition of essential oils and coumarins, contributing to the plant's biological activities (Lunz et al., 2021).

Former studies unveiled, that essential oils obtained from the roots show antibacterial activity against Gram-positive bacteria (Fraternali et al., 2014). Additionally, *A. archangelica* arrested attention as coumarins were identified as being responsible for antibacterial activity against *Staphylococcus aureus*, *Escherichia coli* (*E.coli*) and *Pseudomonas aeruginosa* (Nemeth et al., 2015; Alloush et al., 2022).

So far, with increasing prevalence of untreatable infections induced by antibiotic resistance of bacteria, coumarins were identified as inhibitors of bacterial efflux pump systems (Roy et al., 2013). While many former studies focused on efflux pumps in Gram-positive bacteria, the effects of coumarins on Gram-negative bacteria like *E. coli* are still poorly investigated. Hence, combined analytical systems like thin-layer-chromatography (TLC) and high-pressure-liquid-chromatography coupled with mass spectrometry (HPLC-MS) were used, to get a deeper insight into coumarin pattern of *A. archangelica* and to identify coumarins with efflux pump modifying effects.

To this, the antimicrobial and resistance-modifying profile of plant-derived coumarins were evaluated, and *E. coli* AG100 and the *acrB* gene knockout mutant *E. coli* AG100A were treated with isolated fractions. It could be shown, that fractions containing archangelicin and 2'-angeloyl-3'-isovaleryl vaginate (Wszelaki et al., 2011) induced the most potent modulatory effects on ciprofloxacin (CIP) against *E. coli* AG100 in comparison to *E. coli* AG100A.

Keywords: *Escherichia coli*, *Angelica archangelica*, coumarins, antibacterial activity, efflux

References

- Alloush, M., Mallion, C., Sarker, S. D., Rahman, M. M. 2022. Coumarins from the roots of *Angelica archangelica* and antibacterial activity against methicillin resistant *Staphylococcus aureus*. Dhaka Univ. J. Pharm. Sci. 20, 275–281.
- Fraternali, D., Flamini, G., Ricci, D., 2014. Essential oil composition and antimicrobial activity of *Angelica archangelica* L. (Apiaceae) roots. J. Med. Food 17, 1043–1047.
- Lunz, K., Stappen, I., 2021. Back to the roots—an overview of the chemical composition and bioactivity of selected root- essential oils. Molecules 26, 3155.
- Nemeth, S., Paşca, B., Teodorescu, A., Coita, I., Teaha, D., 2015. Coumarins isolated from the dry roots of *Angelica archangelica* L. and their antibacterial activity. Analele Universităţii din Oradea, Fascicula: Ecotoxicologie, Zootehnie şi Tehnologii de Industrie Alimentară 14, 355–362.
- Roy, S.K., Kumari, N., Pahwa, S., Agrahari, U.C., Bhutani, K.K., Jachak, S.M., Nandanwar, H., 2013. NorA efflux pump inhibitory activity of coumarins from *Mesua ferrea*. Fitoterapia, 90, 140 – 150.
- Wszelaki, N., Paradowska, K., Jamróz, M. K., Granica, S., & Kiss, A. K., 2011. Bioactivity-guided fractionation for the butyrylcholinesterase inhibitory activity of furanocoumarins from *Angelica archangelica* L. roots and fruits. J. Agric. Food Chem. 59, 9186–9193.

* Presenting author. E-mail address: vanessa.baumgartner@uni-graz.at (Vanessa Baumgartner)

S1.P232 Thymoquinone potentiate activity of antibiotics against clinical isolates of *Mycobacterium tuberculosis*

Anna Zabost^a, Ewa Augustynowicz-Kopec^a, Rafał Sawicki^b, Elwira Sieniawska^{c,*}

^a Department of Microbiology, National Tuberculosis and Lung Diseases Research Institute, 01-138 Warsaw, Poland

^b Chair and Department of Biochemistry and Biotechnology, Medical University of Lublin, 20-093 Lublin, Poland

^c Department of Natural Products Chemistry, Medical University of Lublin, 20-093 Lublin, Poland

A numerous plant species are being screened in order to find molecules inhibiting the growth of *Mycobacterium tuberculosis* (Mtb). These include in vitro determination of minimal inhibitory concentration against sensitive and multidrug resistant (MDR)-Mtb strains, inhibition of intracellular bacilli in vitro and ex vivo, and killing of replicating and dormant bacteria (Gupta et al., 2017). However, because molecules active in nano- or micromolar concentrations are found rarely, the plant secondary metabolites are evaluated for their ability to enhance activity of antimycobacterial antibiotics. The synergistic action of plant extracts or isolated compounds against Mtb was already described (Gupta et al., 2017; Polak and Kapka-Skrzypczak, 2017). Interestingly, the synergy was more frequently found when natural compounds were combined with rifampicin rather than with isoniazid or ethambutol (Sieniawska et al., 2018). These combinations showed variable activity against sensitive and drug resistant Mtb clinical strains (Sieniawska et al., 2018, 2020) suggesting the possible differences in their mode of action.

To better understand the differences in the sensitivity of Mtb clinical strains we evaluated the possible synergistic action of first line antibiotics in a combination with thymoquinone, a natural antimicrobial and anti-inflammatory agent. The studies of tuberculostatic activity of antibiotics in the presence of subinhibitory concentrations of thymoquinone showed enhanced activity of rifampicin against all tested strains. This activity was further evaluated in order to investigate the antimycobacterial mechanism of action of thymoquinone.

Keywords: antimicrobials, *Nigella sativa*, tuberculosis, MIC, FICI

References

- Gupta, V.K., Kumar, M.M., Bisht, D., Kaushik, A. 2017. Plants in our combating strategies against *Mycobacterium tuberculosis*: progress made and obstacles met. *Pharm Biol.* 55, 1536-1544.
- Polak, Z., Kapka-Skrzypczak, L. 2017. Anti-tubercular substances produced by plants. *J Pre-Clin Clin Res.* 11, 147-152.
- Sieniawska E, Sawicki R, Golus J, Georgiev, M.I. 2020. Untargeted metabolomic exploration of the *Mycobacterium tuberculosis* stress response to cinnamon essential oil *Biomolecules*, 2020, 10, 357, 1-13.
- Sieniawska E, Sawicki R, Swatko-Ossor M, Napiorkowska, A., Przekora, A., Ginalska, A., Augustynowicz-Kopec, E. The effect of combining natural terpenes and antituberculous agents against reference and clinical *Mycobacterium tuberculosis* strains. *Molecules*, 2018, 23, 176.

Acknowledgements

This work was funded by Polish National Science Centre within the project 2022/45/B/NZ7/00412.

* Presenting author. E-mail address: esieniawska@pharmacognosy.org (Elwira Sieniawska)

S1.P233 Exploring olikomycin a: a novel lipopeptide antibiotic - insights into its mechanism of action

Luisa Munz^{a,*}, Roman Makitrynsky^a, Olga Tsypik^a, Sarah Crocoll^a, Sinja Götz^a, Lena Keller^{b,c}, Amninder Kaur^b, Rolf Müller^b, Heiko Heerklotz^a, Andreas Bechthold^a

^a Institute of Pharmaceutical Sciences, University of Freiburg, Germany

^b Helmholtz Institute for Pharmaceutical Research Saarland (HIPS), Saarland University, Germany

^c Weincampus Neustadt, University of Applied Science Kaiserslautern, Germany

In the world of antibiotics, *Streptomyces* stand out for their large number of biosynthetic gene clusters (BGCs), which enable them to produce a diverse range of natural products. Typically, only a limited number of these BGCs are abundantly expressed under laboratory conditions, while the rest are either poorly expressed or completely silent for unknown reasons.

By specifically deleting the regulatory gene *wblA* in a *Streptomyces* strain, we successfully awakened a silent BGC, leading to the discovery of the natural product olikomycin A with a strong antibiotic activity. Olikomycin A is a cyclic lipopeptide synthesized via NRPS.

Olikomycin A was extracted by liquid-liquid extraction and purified by reversed-phase HPLC. In order to investigate its antibiotic efficacy, a comprehensive analysis was carried out to assess its effects on both peptidoglycan biosynthesis and the structural integrity of the cell membrane. In particular, isothermal titration calorimetry (ITC) was used to analyse interactions with a model membrane.

Olikomycin A has been found to show a similar mechanism of action to daptomycin, which is based on dependence on calcium and phosphoglycerol. Interestingly, however, it exhibits unique pattern in binding behaviour.

This study underlines the potential of *Streptomyces* in producing diverse natural compounds through its wealth of BGCs. The discovery of Olikomycin A, coupled with insights into its purification and antibiotic mechanisms, provides valuable contributions to the field of antibiotics and opens doors for further exploration.

Keywords: *Streptomyces*, novel antibiotics, cyclic lipopeptide, calcium-dependent, non-ribosomal peptide synthetases (NRPS)

*Presenting author. E-mail address: luisa.munz@pharmazie.uni-freiburg.de (Luisa Munz)

S1.P234 GC-MS analysis of essential sandalwood oils with in vitro activity against *Madurella mycetomatis*

Stephan Seegers^{a,*}, Shereen O. Abd Algaffar^b, Prabodh Satyal^c, William N. Setzer^d, Sami A. Khalid^b, Thomas J. Schmidt^a

^a University of Münster, Institute of Pharmaceutical Biology and Phytochemistry (IPBP), Pharma-Campus – Corrensstrasse 48, D-48149 Münster, Germany

^b Faculty of Pharmacy, University of Science and Technology, Omdurman, Sudan

^c Essential Oil Science, dōTERRA International, 1248 W 700 S, Pleasant Grove, UT 84062, USA

^d Department of Chemistry, University of Alabama in Huntsville, Huntsville, AL 35899, USA

Mycetoma is a chronic subcutaneous infection occurring in tropical and subtropical regions. Since 2016, Mycetoma is included in the WHO's list of neglected tropical diseases. Depending of the causative organism, mycetoma is classified as actinomycetoma, caused by bacteria, and eumycetoma, caused by fungi (e.g. *Madurella mycetomatis*). With limited treatment options due to high costs, low availability and poor efficacy of antimycotic drugs, the urge to find new active compounds against eumycetoma is high (Zijlstra et al., 2016).

In an in vitro screening of essential oils against *M. mycetomatis* (Abd Algaffar et al., 2021), Indian sandalwood oil (SWO) the essential oil of *Santalum album* (Santalaceae), with MICs $\leq 0.0039\%$ (v/v) was the most active among 27 tested oils. Therefore, 15 SWOs from 5 different *Santalum* species were tested and thoroughly analyzed by GC-QTOF-MS on two columns (DB-5 and DB-HeavyWax). A biotechnological SWO-surrogate and two oils from other plants marketed as "SWOs" were also tested and analyzed. Over 270 constituents were detected, and

>100 of them identified, many for the first time in the investigated oils. All tested oils showed considerable activity with MICs ranging from 0.0078-0.0019% (v/v). Column-chromatographic separation of SWO on silica yielded fractions of sesquiterpene-hydrocarbons and -alcohols. Only the latter showed activity against *M. mycetomatis*. The main compounds of SWO (Fig. 1), (Z)- α -santalol (**1**) and (Z)- β -santalol (**2**) were obtained by CC on AgNO₃-impregnated silica (Daramwar et al., 2012). The former was twice as active as the latter (MIC = 27.5 vs. 55 $\mu\text{g/mL}$).




Fig. 1. Structures of (Z)- α -santalol (**1**) and (Z)- β -santalol (**2**)

Keywords: *Santalum*, Santalaceae, essential oil, GC-MS, Mycetoma, *Madurella mycetomatis*

References

1. Zijlstra, E.E.; van de Sande, W.W.J.; Welsh, O.; Mahgoub, E.S.; Goodfellow, M.; Fahal, A.H. Mycetoma: a unique neglected tropical disease. *Lancet Infect. Dis.* 2016; 16, 100–112.
2. Abd Algaffar, S.O.; Verbon, A.; van de Sande, W.W.J.; Khalid, S.A. Development and validation of an in vitro resazurin-based susceptibility assay against *Madurella mycetomatis*. *Antimicrob Agents Chemother.* 2021, 65, e01338-20.
3. Daramwar, P. P.; Srivastava, P. L.; Priyadarshini, B.; Thulasiram, H. V. Preparative separation of α - and β -santalenes and (Z)- α - and (Z)- β -santalols using silver nitrate-impregnated silica gel medium pressure liquid chromatography and analysis of sandalwood oil. *The Analyst*, 2012, 19, 4564-4570.

*Presenting author. Email-address: s.seegers@uni-muenster.de (Stephan Seegers)

S1.P235 Bio-guided approach as a valuable tool in the phytochemical investigation of three underexplored Myrtaceae plants

Francesca Guzzo^{a,*}, Elisabetta Buommino^b, Leslie Landrum^c, Antonio Fiorentino^a, Brigida D'Abrosca^a

^a Department of Environmental, Biological and Pharmaceutical Sciences and Technologies- DiSTABiF, University of Campania "Luigi Vanvitelli", via Vivaldi 43, I-81100 Caserta, Italy

^b Department of Pharmacy, University of Naples "Federico II", Via Domenico Montesano 49, 80131 Naples, Italy

^c School of Life Sciences, Arizona State University

Myrtaceae Juss is a large family of flowering plants with privileged chemical and biological profiles (Celaj et al., 2020). In particular, it is a very rich source of phloroglucinol derivatives, which have become attractive targets for organic chemists thanks to their structural features and biological activities, including antimicrobial activity (Nicoletti et al., 2018). Nevertheless, some species in this family, have been scarcely investigated. Thus, this work explores the chemical complexity of the underexplored species, *Myrcianthes cisplatensis*, *Psidium friedrichsthalianum* and *Psidium oligospermum* as a potential source of antimicrobial phytochemicals, through the combination of antimicrobial assay and NMR analysis. Three extracts at increasing polarity, hexane, chloroform and methanol, obtained from leaves of each plant, grown and collected in Arizona, were evaluated for their antimicrobial potential against two strains of *Staphylococcus aureus*: ATCC 29213 and 43300 (a methicillin-resistant *Staphylococcus aureus* strain, MRSA). Hexane and methanol of *M. cisplatensis* (MIC of 16 µg/mL and 64 µg/mL, respectively) and polar extracts of *P. friedrichsthalianum* (MIC of 64 µg/mL and 128 µg/mL, respectively) were the most promising against both the strains. Starting from the most polar ones, a liquid-liquid separation was useful to concentrate the flavonoidic components in ethyl acetate fraction, giving back a more active dichloromethane fraction that reported an increasing antimicrobial potential against both the strains (MIC of 16 µg/mL for *M. cisplatensis*; MIC of 128 µg/mL for *P. friedrichsthalianum*). The fractionation, using different chromatographic techniques, has let to identify the component responsible of the highlighted antimicrobial properties. Four new cinnamoylated alkylphloroglucinol glucosides (Guzzo et al., 2023), along with tricyclic sesquiterpenes, acylated pentacyclic triterpenes, were isolated from bioactive fractions and characterized by extensive 2D NMR analysis (HSQC, H2BC CIGAR-HMBC, COSY, HSQCTOCSY) and HR-MS analyses. The antimicrobial potential of the isolated compounds was evaluated, showing how cinnamoylated alkylphloroglucinol glucosides derivatives were the most active respect to the others and how the bioguided approach is an excellent tool for the characterization of new natural bioactive compounds.

Keywords: bio-guided approach, *Myrcianthes cisplatensis*, *Psidium friedrichsthalianum*, *Psidium oligospermum*, *Staphylococcus aureus*, NMR analysis

References

- Celaj, O., Garcia Duràn, A., Cennamo, P., Scognamiglio, M., Fiorentino, A., Esposito, A., D'Abrosca, B.- 2020. Phloroglucinols from Myrtaceae: attractive targets for structural characterization, biological properties and synthetic procedures. *Phytochemistry Reviews*, 20, 259-299.
- Nicoletti, R., Salvatore, M.M., Ferranti, P.- 2018. Structures and bioactive properties of Myrtucommulones and related acylphloroglucinols from Myrtaceae. *Molecules*, 23:3370.
- Guzzo, F., Buommino, E., Landrum, L., Russo, R., Lembo, F., Fiorentino, A., D'Abrosca, B.- 2023. Phytochemical investigation of *Myrcianthes cisplatensis*: structural characterization of new *p*-coumaroyl alkylphloroglucinols and antimicrobial evaluation against *Staphylococcus aureus*. *Plants*, 12(5), 1046.

* Presenting author. E-mail address: francesca.guzzo@unicampania.it (Francesca Guzzo)

S1.P236 From poison to potential: Nb-methyl usambarensine reveals antiviral activity against SARS- CoV-2

Kristi Leka^{a,b,*}, Olivier Bonnet^a, Sophie Vroomen^b, Michel Frédérich^a, Allison Ledoux^a, Mutien-Marie Garigliany^b

^a University of Liege Faculty of medicine, CIRM, Building B36 Quartier Hôpital 4000, Liege, Belgium

^b University of Liege Faculty of veterinary medicine, FARA, Building B42 Quartier Vallée 2 Avenue de Cureghem 6 4000, Liege, Belgium

Nb-methyl usambarensine is a quaternary alkaloid originally isolated from the roots of *Strychnos usambarensis* (Loganiaceae) (Angelot. L *et al.*, 1975). This plant was historically recognized for its curarizing properties, traditionally employed as an arrow poison in Africa. In response to the SARS- CoV-2 pandemic in 2020, an investigation for potential antiviral agents from the enriched pharmacognosy library was undertaken. Although Nb-methyl usambarensine exhibited no significant activity against *Plasmodium falciparum* in previous studies (Wright *et al.*, 1991), it emerged as a promising candidate during screening against SARS-CoV-2. Additionally, a structurally similar compound, 10-hydroxyusambarensine, was identified as a hit-compound through an in-silico approach, suggesting its potential as an inhibitor of the coronavirus 3-chymotrypsin-like protease (Gyebi *et al.*, 2021).

The antiviral activity of Nb-methyl usambarensine was evaluated in BSL-3 facilities using VERO- E6 cells. The virus was introduced with a multiplicity of infection ratio or MOI of 0.01, alongside varying concentrations of the molecule (1.56 µg/ml- 50 µg/ml). Sampling occurred at 24-hour intervals over 3 days to capture the viral kinetics. The RT-qPCR results at 72h demonstrated a reduction of more than 4 log₁₀ in RNA copies/ml at the concentration of 50 µg/ml of the Nb-methyl usambarensine, while no signs of cytotoxicity were observed. This study highlights the previously unrecognized antiviral potential of Nb-methyl usambarensine against SARS-CoV-2. Further research is needed to elucidate the molecular mechanisms underlying this antiviral activity and to explore its potential as a therapeutic agent against COVID-19.




Fig. 1. Structure of Nb-methyl usambarensine

Keywords: Nb-methyl usambarensine, SARS-CoV-2, antiviral-activity, RT-qPCR

References

Angelot. L *et al.* (1975). *Chemical Structure and Pharmacological (Curarizing) Properties*. 215(2).

Wright, C. *et al.* (1991). Antiamoebic and antiparasmodial activities of alkaloids isolated from *Strychnos usambarensis*. *Planta Medica*, 57(4), 337–340.

Gyebi, G. *et al.* (2021). Potential inhibitors of coronavirus 3-chymotrypsin-like protease (3CLpro): An in silico screening of alkaloids and terpenoids from African medicinal plants. *Journal of Biomolecular Structure and Dynamics*, 39(9), 3396– 3408.

* Presenting author. E-mail address: Kristi.Leka@uliege.be (Kristi Leka)

S1.P237 Semi-synthetic ecdysteroids as promising new antichagasic agents

Márton B. Háznagy^{a,*}, Máté Vágvölgyi^a, Sandhya R. Krishnan^b, Kaushavi Cholke^b, Jürg Gertsch^b, Attila Hunyadi^a

^a Institute of Pharmacognosy, Faculty of Pharmacy, University of Szeged, Eötvös utca 6., 6720 Szeged, Hungary

^b Institute of Biochemistry and Molecular Medicine, Faculty of Medicine, University of Bern, Bülhlstrasse 28., CH-3012 Bern, Switzerland

The parasite *Trypanosoma cruzi* is the causative agent of a severe infectious disease, the so-called Chagas' disease, which is one of the neglected tropical diseases, according to WHO. As therapy is limited to two available drugs, (WHO; Rassi et al 2010) seeking new effective agents is crucial.

In this study, the screening of fifty-two ecdysteroids for their antichagasic effects revealed that cinnamic ester and *tert*-butyl oxime ether moieties might have a key role in the antiparasitic effect. Based on the screening results, new compounds were designed and prepared, incorporating the two potential pharmacophores. 20-Hydroxyecdysone (20E) was used as starting material, which is the most widespread natural ecdysteroid. Six compounds were semi-synthesised, among which two cinnamic mono-esters of 20E 6-*tert*-butyl oxime ether derivatives exerted the most potent activity. (Háznagy et al., 2022) IC₅₀ measurements (MTT and XTT assays) (Salm A, et al, 2021) were carried out with compound (1) (Fig. 1). The results demonstrated that compound (1) has six times selective toxicity on *T. cruzi* epimastigotes as compared to cytotoxicity on the native cell line. The FACS- based quantitation of released parasites from infected cells indicated that (1) exerted an IC₅₀ value of 2.7 µm.




Fig. 1. Structure of the most active ecdysteroid derivative 1

Semi-synthetic preparation of new derivatives is currently ongoing. A further aim is to investigate the role of the substitution pattern of the cinnamic acid moiety.

Keywords: *Trypanosoma cruzi*, ecdysteroids, semi-synthesis, cinnamic esters

Acknowledgements: Grants NKFIH K-134704 and TKP2021-EGA-32 are acknowledged.

References

- Háznagy, M., Vágvölgyi, M., Krishnan, S.R., Gertsch, J., Hunyadi, A., 2022. Antitrypanosomal activity of semisynthetic enone-type derivatives *Planta Medica*. 88(15): 1534
- Rassi, A.Jr, Rassi, A., Marin-Neto, J.A., 2010. Chagas disease. *The Lancet*. 375 (9723): 1388-402
- Salm, A., Krishnan, S.R., Collu, M., Danton, O., Hamburger M., Leonto M., Almanza G., Gertsch, J., 2021. Phylobioactive hotspots in plant resources used to treat Chagas disease *iScience*. 24(4):102310.
- World Health Organization. "Chagas disease (American trypanosomiasis)" Retrieved from <https://www.who.int/health-topics/chagas-disease>

* Presenting author. E-mail address: haznagy.marton.benedek@szte.hu (Márton Benedek Háznagy)

S1.P238 Exploring the potential biopesticide application of nemertide alpha-1 on parasitic and disease vector mosquitoes

Quentin Laborde^{a,*}, Melika Hajkazemian^b, Justine Bonnassieux^b, Håkan S. Andersson^{a,c}, Ulf Göransson^a, S. Noushin Emami^b

^a Division of Pharmacognosy, Department of Pharmaceutical Bioscience, Faculty of Pharmacy, Uppsala University, Sweden

^b Department of Molecular Biosciences, Wenner-Gren Institute, Stockholm University, Stockholm, Sweden

^c Department of Medical Biochemistry and Biophysics, Karolinska Institute, Sweden

This poster presents the initial exploration of nemertide alpha-1 as a potential biopesticide targeting pest insects. Nemertide alpha-1, a neurotoxic peptide consisting of 31 amino acids, has shown potent biopesticidal activity, with an EC₅₀ of 8.6 nM on BgNav1 (*Blatella germanica*) (Jacobsson et al., 2018). Additionally, it demonstrated complete inhibition of inactivation in Nav channels of *Drosophila melanogaster* and *Varroa destructor* at 1 µM (Jacobsson et al., 2018). Bell et al. (2021) also reported the insecticidal efficacy of a recombinant nemertide alpha-1 propeptide on cabbage moth larvae (*Mamestra brassicae*) and proboscis feeding insects such as adult aphids (*Acyrtosiphon pisum* and *Myzus persicae*) and adult bees (*Apis mellifera*). Building upon these promising findings, our study aims to explore the insecticidal potency of alpha-1 while considering its implications for human health and the emergence or re-emergence of infectious diseases. Here, we present a preliminary investigation into the acute toxicity effects of nemertide alpha-1 on *Anopheles* mosquitoes (vectors of malaria: *Plasmodium falciparum*) and *Aedes* mosquitoes (vectors of viral diseases: dengue fever, Zika fever, yellow fever, and Chikungunya).

Keywords: nemertide, marine, peptide toxin, biopesticide, mosquito

References

- Bell, J., Sukiran, N. A., Walsh, S., & Fitches, E. C., 2021. The insecticidal activity of recombinant nemertide toxin α-1 from *Lineus longissimus* towards pests and beneficial species. *Toxicon*, 197, 79-86.
- Jacobsson, E., Andersson, H. S., Strand, M., Peigneur, S., Eriksson, C., Lodén, H., ... & Göransson, U., 2018. Peptide ion channel toxins from the bootlace worm, the longest animal on Earth. *Scientific Reports*, 8(1), 4596.

* Presenting author. E-mail address: quentin.laborde@uu.se (Quentin Laborde)

S1.P239 Chemical constituents of *Sambucus nigra* fruit (European Elderberry)

Zulfiqar Ali^{a,*}, Fazila Zulfiqar^a, Avula Bharathi^a, Ikhlas A. Khan^{a,b}

^a National Center for Natural Products Research, School of Pharmacy, University of Mississippi, University, MS 38677, USA

^b Division of Pharmacognosy, Department of BioMolecular Sciences, School of Pharmacy, University of Mississippi, University, MS 38677, USA

Sambucus spp. are broadly dispersed throughout the tropical/subtropical and temperate areas. *Sambucus nigra* L. (syn. *S. nigra* subsp. *nigra*) is one of the five best-known among 30 species in the genus *Sambucus* L. (elder) (Viburnaceae). *S. nigra*, a shrub commonly known as elderberry, elder, European elderberry, European elder, black elder, and European black elderberry, is prevalent in Europe, North Africa, and West and Central Asia. Various biological activities such as anti-inflammatory, diuretic, and antioxidative have been associated with *S. nigra* (Ahmadiani et al., 1998; Ebrahimzadeh et al., 2006). *S. nigra* fruit is a cultivated as well as a wild-harvested species and is one of the richest sources of flavonoids and anthocyanins (Duymuş et al., 2014).

Elderberry products famed over the past couple of years due to their use in the COVID-19 pandemic, though, there is no robust evidence to support the prevention or treatment of this disease. Two iridoids including one previously undescribed, an indole alkaloid, and various flavonoid glycosides (Fig.1) were isolated and characterized from the dried fruit of *S. nigra*. The compounds were purified from the hydroalcoholic extract of the dried berries after various chromatographic procedures. Structure elucidation was accomplished from NMR and mass data analysis.




Fig. 1. Structure of iridoid, alkaloid, and flavonoids

Keywords: *Sambucus nigra*, Viburnaceae, iridoid, alkaloid, flavonoid

References

- Ahmadiani, A., Fereidoni, M., Semnani, S., Kamalinejad, M., Saremi, S., 1998. Antinociceptive and anti-inflammatory effects of *Sambucus ebulus* rhizome extract in rats. *J. Ethnopharmacol.* 61, 229-235.
- Duymuş, H.G., Göger, F., Başer, K.H.C., 2014. In vitro antioxidant properties and anthocyanin compositions of elderberry extracts. *Food Chem.* 155, 112-119.
- Ebrahimzadeh, M.A., Mahmoudi, M., Salimi, E., 2006. Antiinflammatory activity of *Sambucus ebulus* hexane extracts. *Fitoterapia* 77, 146-148.

* Presenting author. E-mail address: zulfiqar@olemiss.edu (Zulfiqar Ali)

S1.P240 Invasive plants as promising solutions against biofilms

Guillaume Hamion^a, Sufi Desrini^a, Willy Aucher^a, Caroline Rouger^b, Jérôme Labanowsky^c, Mustofa^{d,f}, Titik Nuryastuti^{e,f}, Christine Imbert^a, Marion Girardot^{a,*}

^a Laboratoire EBI, UMR CNRS 7267, Université de Poitiers, F-86000 Poitiers, France

^b Université de Bordeaux, UMR 1366 INRAE, Bordeaux INP, OENO, ISVV, F-33140 Villenave d'Ornon, France

^c IC2MP, UMR CNRS 7285, Université de Poitiers, F-86000 Poitiers, France

^d Department of Pharmacology and Therapy, Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

^e Department of Microbiology, Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

^f Indonesian Biofilm Research Collaboration Center UGM-BRIN, Yogyakarta, Indonesia

Biofilm is a widespread way of life among human pathogenic microorganisms and conventional drugs are poorly effective against biofilm-associated infections. Invasive plants, due to their often higher chemical diversity than native species, constitute interesting sources of active molecules.

Among 73 extracts obtained from 10 invasive plants collected in France, the essential oil (EO) from the leaves of *Baccharis halimifolia* was the most active against *Candida albicans* (Ca) biofilms, acting against the maturation phase and the 24h biofilm (IC₅₀=4 and 74 µg/mL respectively). The GC-MS analysis of this EO showed the presence of 48 compounds mainly identified as oxygenated sesquiterpenes (62%). The most abundant compound was β-caryophyllene oxide (37%) which showed, as well as aromadendrene oxide-(2) and (±)-β-pinene also present, significant anti- maturation and antibiofilm effects of Ca (IC₅₀=9-630 µmol/L) (Desrini et al., 2023). Also, the 2- methyltetrahydrofuran stem extract of *Ludwigia grandiflora* was the most active against the bi- species biofilm *Staphylococcus aureus* (Sa)-Ca, inhibiting the mature biofilm by more than 50% at 50 µg/mL. An approach using bioguided molecular networks correlating data from UHPLC-MS/MS analyzes of extracts and fractions with the results of antibiofilm tests revealed seven compounds of interest in *L. grandiflora*, belonging to pentacyclic triterpenes and lipids including acylglycerols and hydroxylated and epoxidized derivatives of fatty acids. The most correlated compound, betulinic acid, significantly reduced Sa-Ca preformed biofilms by several pairs of strains (inhibition>40% at 25 µg/mL) (Hamion et al., 2022).

This study clearly confirms the potential of invasive plants as promising solutions against biofilms.

Keywords: invasive plants, biofilm, *Baccharis halimifolia*, *Ludwigia grandiflora*, antibiofilm compounds

References

- Desrini, S., Ducloux, J., Hamion, G., Bodet, C., Labanowski, J., Mustofa, M., Nuryastuti, T., Imbert, C., Girardot M., 2023. Antibiofilm activity of invasive plants against *Candida albicans*: Focus on *Baccharis halimifolia* essential oil and its compounds. Chem Biodivers. 20 :e202300130.
- Hamion, G., Aucher, W., Tardif, C., Miranda, J., Rouger, C., Imbert, C., Girardot, M., 2022. Valorization of invasive plant extracts against the bispecies biofilm *Staphylococcus aureus*-*Candida albicans* by a bioguided molecular networking screening. Antibiotics. 11, 1595.

* Presenting author. E-mail address: marion.girardot@univ-poitiers.fr (Marion Girardot)

S1.P241 Isolation and characterization of antimicrobial compounds from *Picralima nitida* (Apocynaceae)

Makafui Adzo Gbemu^a, Susana Dufie Boatey^a, Denick Amon^a, Prince Okyere Dagadu^b, Priscilla Kolibea Mante^b, Yaw Duah Boakye^c, Edmund Ekuadzi^a

^a Department of Pharmacognosy, FPPS, KNUST, Kumasi, Ghana

^b Department of Pharmacology, FPPS, KNUST, Kumasi, Ghana

^c Department of Pharmaceutics, FPPS, KNUST, Kumasi, Ghana

Our current work is focused on re-isolating akuammidine (**1**), and akuammine (**2**), two major monoterpene indole alkaloids from *Picralima nitida* Stapf. Th & H. Durand (Apocynaceae) and their biological evaluation. The plant is used traditionally for treating infectious diseases (Menzies et al., 1998). The global quest to curb antimicrobial resistance elicits the need to harness antimicrobial and resistant modifying compounds from nature. Seeds were collected from Kpetoe in the Volta region of Ghana. The alkaloidal extract showed significant resistance-modifying properties in efflux pump inhibition and biofilm formation assays against resistant microbial strains.




Fig 1. Chemical structures of isolated compounds, akuammidine (**1**) and akuammine (**2**)

Structure elucidation of the compounds was achieved using spectroscopic techniques. Biofilm inhibition assay revealed the compounds to be ineffective against *E. coli* and *C. albicans*. However, they demonstrated significant inhibition of biofilm formation in *P. aeruginosa*, *S. aureus*, and *B. subtilis* for akuammidine by 60%, 67%, 53%, respectively and akuammine by 61%, 27%, and 61% respectively. Additionally, they exhibited superior efflux pump inhibition activity on the organisms except *S. aureus*. These findings suggest the potential application of the compounds as promising leads in the development of antimicrobial agents to combat antimicrobial resistance.

Keywords: *Picralima nitida*, resistant strains, isolation, biofilm formation, efflux pump inhibition

References

Menzies, J.R., Paterson, S.J., Duwiejua, M., Corbett, A.D., 1998. Opioid activity of alkaloids extracted from *Picralima nitida* (fam. Apocynaceae). Eur. J. Pharmacol. 350, 101–108.

* Presenting author. E-mail address: makafuiadzogbemu100@gmail.com (Makafui Adzo Gbemu)

S1.P242 Identification of a Potent Antibiotic Compound from *Geodermatophilus* Bacteria

Sunmin Woo^a, Brendan Lin^e, Paul M. Dunman^e, Cassandra L. Quave^{a,b,c,d}

¹ Center for the Study of Human Health, Emory University, Atlanta, GA 30322, USA

² Department of Dermatology, Emory University, Atlanta, GA 30322, USA

³ Molecular and Systems Pharmacology Graduate Program, Emory University, Atlanta, GA 30322, USA

⁴ Emory University Herbarium, Emory University, Atlanta, GA 30322, USA

⁵ Department of Microbiology and Immunology, University of Rochester Medical Center, Rochester, NY 14642, USA

The majority of antibiotics in clinical use today are natural products derived from microorganisms in the environment. One such example is gentamicin, which Dr. Marvin Weinstein discovered in soil bacteria and produced through fermentation of *Micromonospora echinospora*. One of Dr. Weinstein's gentamicin project members, Dr. Luedemann (Weinstein et al., 1963) continued to collect microbial samples from arid desert environments over twenty years and left a large collection of source organisms when he passed away. Herein, we screened over 300 *Geodermatophilus* bacteria from the over 2000 isolate Luedemann collection to discover potent antibiotics candidates from uncharacterized microbial sources. We identified five species with potent antibacterial activity. Among them, lyophilized 9005BA culture supernatant exhibited a relative strong MIC of 100 µg/mL against *Acinetobacter baumannii* and weaker activity against *S. aureus*. The 9005BA was selected for further isolation and structural elucidation studies on the active compound. Bioactivity-guided isolation was performed following a microfractionation technique using HPLC-DAD to target the active compound. In parallel to HPLC-based fractionation, HPLC-ESI-Q-TOF MS in positive ion mode was used for the early detection and prediction of the active compound presented in the extract and fraction. Compound isolation was achieved by solid phase extraction and further semi-preparative HPLC and its structure was established by carrying out 1D and 2D NMR experiments along with HRMS analyses. Here we present our findings on a novel potent antibiotic alkaloid for the gram- negative pathogen, *A. baumannii*.

Keywords: *Geodermatophilus* bacteria, antibiotic, alkaloid, microfractionation, *Acinetobacter baumannii*

References

Weinstein, M. J., Luedemann, G. M., Oden, E. M., Wagman, G. H., 1963. Gentamicin, a New Broad-Spectrum Antibiotic Complex. *Antimicrob Agents Chemother* (Bethesda). 161, 1–7.

* Presenting author. E-mail address: sun-min.woo@emory.edu (Sunmin Woo)

S1.P243 Ethnobotany as a tool for targeted discovery of bioactive natural products for emerging infectious diseases

Cassandra L. Quave*

Department of Dermatology, Emory University School of Medicine, Atlanta, Georgia, USA

Out of the estimated 374,000 species of plants on Earth, 9% have been documented as being used in medicine. Billions of people across the globe currently rely on plants to meet their primary healthcare needs, including for the treatment of infection. Many of our life-saving therapies for treating cancer, pain, heart disease, and infection were developed based on chemical scaffolds first discovered in plants. Yet, most medicinal plants have never been examined through the lens of modern science. This presentation will cover the historical importance of plants in the evolution of current therapeutics and explore emerging technologies, such as microelectron diffraction (Delgadillo et al., 2023), enabling scientists to look deeper into the pharmacological properties of plants. Major classes of antimicrobial natural products will be discussed, and insights on research bottlenecks and challenges in scientific rigor in the field will be addressed (Porras et al., 2021). Examples of a novel antibiotic potentiator **1** (Dettweiler, et al. 2020) from *Callicarpa americana* L., Lamiaceae and antivirulence **2** (Tang et al., 2020) and antifungal **3** (Marquez et al., 2023) agents from *Schinus terebinthifolia* Raddi, Anacardiaceae will be used to illustrate the ethnobotanical approach to drug discovery (Fig. 1).




Fig. 1. Structure of a clerodane diterpene **1**, triterpenoid acid **2**, and gallotannin **3**.

Keywords: ethnobotany, antimicrobial, drug discovery, antimicrobial resistance

References

- Delgadillo, D.A., Burch, J.E., Kim, L.J., Moraes, L.S., Niwa, K., Williams, J., Tang, M.J., Lavallo, V.G., Chhetri, B.K., Jones, C.G., Rodriguez, I.H., Signore, J.A., Marquez, L., Bhanushali, R., Woo, S., Kubanek, J., Quave, C., Tang, Y., Nelson, H., 2023. High-throughput identification of crystalline natural products from crude extracts enabled by microarray technology and microED. *ACS Cent Sci.* 10(1), 176-183.
- Dettweiler, M., Melander, R.J., Porras, G., Risener, C., Marquez, L., Samarakoon, T., Melander, C., Quave, C.L., 2020. A clerodane diterpene from *Callicarpa americana* resensitizes Methicillin-resistant *Staphylococcus aureus* to β -Lactam antibiotics. *ACS Infect Dis.* 6(7), 1667-1673.
- Marquez, L., Lee, Y., Duncan, D., Whitesell, L., Cowen, L.E., Quave, C., 2023. Potent antifungal activity of Penta- O- galloyl- β -d-Glucose against drug-resistant *Candida albicans*, *Candida auris*, and other non-albicans *Candida* species. *ACS Infect. Dis.*, 9(9), 1685-1694.
- Porras, G., Chassagne, F., Lyles, J.T., Marquez, L., Dettweiler, M., Salam, A.M., Samarakoon, T., Shabih, S., Farrokhi, D.R., Quave, C.L., 2021. Ethnobotany and the role of plant natural products in antibiotic drug discovery. *Chem Rev.* 121(6), 3495-3560.
- Tang, H., Porras, G., Brown, M.M., Chassagne, F., Lyles, J.T., Bacsa, J., Horswill, A.R., Quave, C.L., 2020. Triterpenoid acids isolated from *Schinus terebinthifolia* fruits reduce *Staphylococcus aureus* virulence and abate dermonecrosis. *Sci Rep.* 10(1), 8046.

* Presenting author. E-mail address: cquave@emory.edu (Cassandra L. Quavea)

S1.P244 Amaryllidaceae alkaloids as promising anti-dengue phytochemicals

Ngoc-Thao-Hien Le^{a,b}, Suzanne Kaptein^c, Elke Maas^c, Johan Neyts^c, Luc Pieters^a, Emmy Tuenter^{a,*}

^a Natural Products & Food Research and Analysis - Pharmaceutical Technology (NatuRAPT), Department of Pharmaceutical Sciences, University of Antwerp, Antwerp, Belgium

^b Université Paris Cité - INSERM Unit 1284, Paris, France

^c Department of Microbiology, Immunology and Transplantation, Rega Institute for Medical Research, Laboratory of Virology and Chemotherapy, KU Leuven, Leuven, Belgium

Amaryllidaceae alkaloids (AAs) are compounds belonging to the large group of plant-derived isoquinoline alkaloids, which exhibit various pharmacological activities such as anticancer, antiparasitic, anti-inflammatory, antiviral, etc (Ding et al, 2017). This presentation will cover the anti-dengue serotype-2 (DENV-2) evaluation of AAs isolated from *Hymenocallis littoralis* (Le et al., 2022; 2023a), *Scadoxus multiflorus* (Le et al., 2023b) and *Pancratium maritimum* (Le et al., 2023c). In total, a library of 37 Amaryllidaceae alkaloids was built from the three plant species, belonging to seven skeletal subclasses: 14 lycorine-type, 3 lycorenine-type, 3 narciclasine-type, 8 crinine/haemanthamine-type, 6 galanthamine-type, 1 montanine-type and 2 norbelladine-type compounds. The library was screened *in vitro* against the DENV2 virus and cytotoxicity was assessed on Vero- E6 cells. As a result, 12 compounds displayed potent inhibition of DENV2 replication ($EC_{50} = 0.023 - 3.22 \mu M$); 4 compounds exhibited weak inhibition ($EC_{50} = 12.9 - 29.4 \mu M$); and 21 compounds were devoid of activity and cytotoxicity. All potent anti-DENV-2 AAs described in this study showed a certain level of cytotoxicity, but did possess selective antiviral activity ($SI = 15 - 127$). Preliminary evaluation of structure-activity relationships revealed four skeleton-types (lycorine, homolycorine, narciclasine, and haemanthamine) as promising scaffolds for further research. Synergy was assessed for dual combinations of the four most potent AAs and moderate synergistic effects were also observed.

Keywords: Amaryllidaceae alkaloids, anti-DENV2, cytotoxicity, synergy

References

- Ding, Y., Qu, D., Zhang, K.-M., Cang, X.-X., Kou, Z.-N., Xiao, W., Zhu, J.-B., 2017. Phytochemical and biological investigations of Amaryllidaceae alkaloids: a review. *Journal of Asian Natural Products Research*, 19(1), 53–100.
- Le, N. T. H., Vermeyen, T., Aerts, R., Herrebout, W. A., Pieters, L., & Tuenter, E., 2022. Epimeric Mixture Analysis and Absolute Configuration Determination Using an Integrated Spectroscopic and Computational Approach - A Case Study of Two Epimers of 6-Hydroxyhippeastidine. *Molecules*, 28(1), 214.
- Le, N. T. H., De Jonghe, S., Erven, K., Neyts, J., Pannecouque, C., Vermeyen, T., Herrebout, W. A., Pieters, L., Tuenter, E., 2023a. Anti-SARS-CoV-2 activity and cytotoxicity of Amaryllidaceae alkaloids from *Hymenocallis littoralis*. *Molecules*, 28(7), 3222.
- Le, N. T. H., De Jonghe, S., Erven, K., Neyts, J., Pannecouque, C., Vermeyen, T., Herrebout, W. A., Pieters, L., Tuenter, E., 2023b. Comprehensive study of alkaloids from *Scadoxus multiflorus* by HPLC-PDA-SPE-NMR and evaluation of their anti-SARS-CoV-2 activity. *Phytochemistry Letters*, 57(August), 156–162.
- Le, N. T. H., De Jonghe, S., Erven, K., Neyts, J., Pannecouque, C., Vermeyen, T., Herrebout, W. A., Pieters, L., Tuenter, E., 2023c. A new alkaloid from *Pancratium maritimum* - Structure elucidation using computer-assisted structure elucidation (CASE) and evaluation of cytotoxicity and anti-SARS-CoV-2 activity. *Phytochemistry Letters*, 58(September), 1–7.

* Presenting author. E-mail address: emmy.tuenter@uantwerpen.be (Emmy Tuenter)

S1.P245 Multi-target action of volatile compounds from essential oils in the treatment of periodontitis

Ruxandra Ștefănescu^{a,*}, Bianca Eugenia Ösz^b, Amelia Tero-Vescan^c, Dan Dragoș Sita^d

^a Pharmacognosy and Phytotherapy Department, Faculty of Pharmacy, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Târgu Mureș, 540139 Târgu Mureș, Romania

^b Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Targu Mures, 540139 Targu Mures, Romania

^c Medical Chemistry and Biochemistry Department, Faculty of Medicine in English, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Târgu Mureș, 540139 Târgu Mureș, Romania

^d Department of Odontology and Oral Pathology, Faculty of Dental Medicine, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Târgu Mureș, 540139 Târgu Mureș, Romania

Periodontal disease has a multifactorial etiology including genetic predisposition, environmental factors and microbial dysbiosis, where Gram-negative anaerobic bacteria play a crucial role. Essential oils have long been used to treat a variety of disorders affecting the oral cavity, and they are mostly known for their antibacterial effect. However, emerging evidence suggests that the predominant bioactive compounds within these oils confer distinct specificity in targeting various bacterial strains. The main mechanism of action seems to be the disruption of the bacterial cell membrane and the potency is higher for the alcohols and phenols (e.g. thymol and carvacrol). Other volatile compounds prevent bacterial replication and transcription by inhibiting the expression of some genes linked to bacterial energy metabolism, and DNA metabolism (Álvarez-Martínez et al, 2021). Evidence from the clinical studies suggests that the antibacterial and anti-inflammatory effects of essential oils are maintained *in vivo*, by inhibiting the growth and biofilm formation of these pathogenic microorganisms (Azad et al, 2016; Takallu et al, 2024). Recent studies suggest that some volatile compounds target pro-inflammatory cytokines, such as interleukin-1 β (IL-1 β) and tumor necrosis factor- α (TNF- α), as well as inflammatory enzymes, including cyclooxygenase-2 (COX- 2) and matrix metalloproteinases (MMPs) (Tazehjani, 2021). Future research should focus on elucidating the molecular mechanisms underlying compound-specific antibacterial and anti- inflammatory effects in order to develop targeted strategies that can enhance the treatment of periodontal disease.

Keywords: periodontal disease, essential oils, inflammation modulators, antibacterial activity

References

1. Álvarez-Martínez, F.J.; Barrajón-Catalán, E.; Herranz-López, M.; Micol, V., 2021. Antibacterial Plant Compounds, Extracts and Essential Oils: An Updated Review on Their Effects and Putative Mechanisms of Action. *Phytomed.* 90, 153626.
2. Azad, M.F.; Schwiertz, A.; Jentsch, H.F.R., 2016. Adjunctive Use of Essential Oils Following Scaling and Root Planing –a Randomized Clinical Trial. *BMC Complement. Altern. Med.* 16, 171.
3. Takallu, S.; Mirzaei, E.; Zakeri Bazmandeh, A.; Ghaderi Jafarbeigloo, H.R.; Khorshidi, H., 2024. Addressing Antimicrobial Properties in Guided Tissue/Bone Regeneration Membrane: Enhancing Effectiveness in Periodontitis Treatment. *ACS Infect. Dis.* 4.Tazehjani, D.A.J.; Farahpour, M.R.; Hamishehkar, H., 2021. Effectiveness of Topical Caraway Essential Oil Loaded into Nanostructured Lipid Carrier as a Promising Platform for the Treatment of Infected Wounds. *Colloids Surf. A: Physicochem. Eng. Asp.* 610, 125748.

Acknowledgments

This work was supported by the University of Medicine, Pharmacy, Science and Technology „George Emil Palade“ of Târgu Mureș, Research Grant number 163 /7/ 10.01.2023.

*Presenting author: E-mail address: ruxandra.stefanescu@umfst.ro (Ruxandra Ștefănescu)

S1.P246 Metabolomic-guided phytochemical investigation and antimicrobial activity of *Morus alba* L. twigs extract

Ernesto Gargiulo*, Elisabetta Buommino, Francesca Lembo, Giuseppina Chianese, Orazio Tagliatela-Scafati

Department of Pharmacy, School of Medicine and Surgery, University of Naples "Federico II", 80131 Naples, Italy

In the realm of public health, a pressing concern is antimicrobial resistance, a significant global threat (De Oliveira et al., 2020). Plant natural products offer promise in combating multi-drug resistance by acting as antibacterial agents through various mechanisms (Woo et al., 2023). Widespread natural product classes, such as terpenes, flavonoids, alkaloids, and phenolics have demonstrated antimicrobial activity (Drasar, 2020).

Morus alba L. (Moraceae) twigs have historical use in traditional medicine for muscle-related symptoms, yet their chemical constituents and clinical potential remain underexplored (Kwak et al., 2023). Here, we conducted a metabolomic-guided phytochemical investigation, based on a LC- HRMS/MS and Molecular Networking combined approach, of an industrial extract of *M. alba* L. twigs, obtained in the field of our scientific partnership with Indena Spa (Milan, Italy). Our analysis allowed us to swiftly profile and isolate 17 secondary metabolites including stilbenoids, flavonoids, flavanones, and chalcones. Molecular networking uncovered a cluster of prenylated polyphenols, with several nodes having a similar parent mass. Manual annotation based on LC-HRMS combined with isolation and NMR-based structure elucidation allowed the identification of a new chromenochalcone derivative.

Isolated metabolites were tested for their antimicrobial activity against *Staphylococcus spp.* The most active compound resulted to be kuwanon C, exhibiting a MIC of 8 mg/mL against *S. aureus* ATCC 43300 (methicillin resistant, MRSA) and *S. epidermidis* ATCC 3598 (a biofilm producers' strain). The same concentration resulted bactericidal. We also observed an additive interaction between 4 mg/mL kuwanon C in combination with low oxacillin dosage against the MRSA.

Keywords: *Morus alba*, metabolomics, phytochemical investigation, antimicrobial, MRSA

References

- De Oliveira D. M. P., Forde B.M., Kidd T.J., Harris P.N., Schembri M.A., Beatson S.A., Paterson D.L., Walker M.J., 2020. Antimicrobial Resistance in ESKAPE Pathogens. *Clinical Microbiology Reviews* 33 (3).
- Drasar P. D., 2020. Growing importance of natural products research. *Molecules* 25(1), 6.
- Kwak, H.J., Kim, J., Kim, S., Park S., Choi J., Kim S.H., 2023. Moracin E and M isolated from *Morus alba* Linné induced the skeletal muscle cell proliferation via PI3K-Akt-mTOR signaling pathway. *Sci Rep* 13, 20570.
- Woo S., Marquez L., Crandall W.J., Risener C.J., Quave C.L., 2023. Recent advances in the discovery of plant-derived antimicrobial natural products to combat antimicrobial resistant pathogens: insights from 2018–2022. *Natural products report* 40, 1271–1290.

* Presenting author. E-mail address: ernesto.gargiulo@unina.it (Ernesto Gargiulo)

S1.P247 A new approach to study the mechanisms of action of antimicrobial plant-derived molecules in prokaryotic cells

Raffaella Nocera^{a,*}, Valentina Parisi^a, Emanuele Rosa^a, Fabrizio Dal Piaz^b, Giuliana Donadio^a, Nunziatina De Tommasi^a

^a Department of Pharmacy, University of Salerno, Via Giovanni Paolo II, 132, 84084 Fisciano (SA), Italy

^b Department of Medicine and Surgery, University of Salerno, Via Salvador Allende, 43, 84081 Baronissi (SA), Italy

The identification of molecular targets of small-molecule drugs is a major challenge of pharmaceutical research (Lomenick et al., 2009). In this context, proteomic-based approaches appear to be very promising. In particular, Drug Affinity Responsive Target Stability (DARTS) coupled with mass spectrometry emerged as an efficient, quick and easy method (Dal Piaz et al., 2016). Although the DARTS approach is widely used to investigate protein interactions in eukaryotic cells and protein extracts, it is still not exploited in bacteria. Therefore, the use of this approach to study antibacterial compounds is limited. The main problem with bacteria is their very fast duplication times and stress- response. To overcome such a limitation, we developed an experimental protocol based on the use of minimal medium, thus slowing down bacterial proliferation and metabolism. We used this approach to identify the molecular target of a plant diterpene with antimicrobial activity. As a bacterium, we chose *Streptococcus mutans*, which recently emerged as a novel Gram-positive model organism (Lemos et al., 2013). The experiment was performed by incubating *S. mutans* with the diterpene while cultured in minimal media, and then performing the typical DARTS protocol followed by LC-MS/MS analysis. The DARTS approach was also carried out using *S. mutans* protein lysate, and the results obtained in the two experiments were compared. Remarkably, most of the proteins identified in the two procedures were the same and their inhibition by the diterpene could actually underlie its antimicrobial activity. These results confirmed that our DARTS experimental model can also be applied to prokaryotic cells.

Keywords: prokaryotic cells, drug affinity responsive target stability assay, plant-derived molecules, bacteria, molecular target(s)

References

- Dal Piaz, F., Vera Saltos, M.B., Franceschelli, S., Forte, G., Marzocco, S., Tuccinardi, T., Poli, G., Ebrahimi, S. N., Hamburger, M., De Tommasi, N., Braca, A., 2016. Drug affinity responsive target stability (DARTS) identifies laurifolioside as a new clathrin heavy chain modulator. *J. Nat. Prod.* 79, 2681-2692.
- Lemos, J. A., Quivey Jr, R.G., Koo, H., Abranches, J., 2013. *Streptococcus mutans*: a new Gram-positive paradigm? *Microbiol.* 159, 436-445.
- Lomenick, B., Hao, R., Jonai, N., Chin, R.M., Aghajan, M., Warburton, S., Wang, J., Wu, R.P., Gomez, F., Loo, J.A., Wohlschlegel, J.A., Vondriska, T.M., Pelletier, J., Herschman, H.R., Clardy, J., Clarke, C.F., Huang, J., 2009. Target identification using drug affinity responsive target stability (DARTS). *Proc. Natl. Acad. Sci. U S A* 106, 21984-21989.

* Presenting author. *E-mail address:* ranocera@unisa.it (Raffaella Nocera)

S1.P248 Harnessing the constrained macrocyclic peptide frameworks for the development of antimicrobial peptides to treat infections

Taj Muhammad^{a,*}, Sunithi Gunasekera^a, John Kerr White^b, Adam A. Strömstedt^a, Soumitra Mohanty^b, Anna Blasi-Romero^c, Carlos Palo-Nieto^c, Natalia Ferraz^c, Annelie Brauner^b, Ulf Göransson^a

^a Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden

^b Department of Microbiology, Tumor and Cell Biology, Division of Clinical Microbiology, Karolinska Institute, Stockholm, Sweden

^c Nanotechnology and Functional Materials, Department of Materials Science and Engineering, Uppsala University, Uppsala, Sweden

The need for new innovative treatments for common infections like urinary tract infections (UTI) and skin infections is critical, and among the top 10 reasons to visit outpatient clinics. This is also of particular concern because of increasing antibiotic resistance. Despite widespread acknowledgment of this huge and demanding problem, progress in developing alternative therapeutics has been slow, and current efforts are insufficient. Where do we find novel chemical entities with therapeutic potential against pathogens? Nature has been a tremendous inspiration for novel chemistry throughout history and continues to be a rich reservoir of new chemical entities. One such source of inspiration is antimicrobial peptides (AMPs) produced by all living organisms as important and indispensable components of their innate immune system (Moretta et al., 2021).

Several AMPs have reached clinical trials. However, their development is hampered by a lack of stability because of enzymatic degradation, and a lack of activity when tested in biological contexts. The constrained macrocyclic peptides have experienced a renaissance in medicinal chemistry in recent years (Torres et al., 2019).

Over the years, we have investigated several macrocyclization methodologies, which involve dimerization, backbone cyclization, and the creation of cross-links via a disulfide bond. Our research has utilized KR-12, a truncated peptide derived from the human host defense peptide LL-37, for peptide engineering purposes (Muhammad et al., 2023). Furthermore, we have refined a lead compound for its application in treating bacterial infections (White et al., 2022).

Keywords: antimicrobial peptides, infections, LL-37, KR-12, macrocyclic peptides

References

- Moretta, A., Scieuzo, C., Petrone, M. A., Salvia, R., Manniello, M. D., Franco, A., Lucchetti, D., Vassallo, A., Vogel, H., Sgambato, A., Falabella, P., 2021. Antimicrobial Peptides: A new hope in biomedical and pharmaceutical fields. *Front. Cell Infect. Microbiol.* 11, 668632.
- Muhammad, T., Strömstedt, A. A., Gunasekera, S., Göransson, U., 2023. Transforming cross-linked cyclic dimers of KR- 12 into stable and potent antimicrobial drug leads. *Biomed.* 11, 504.
- Torres, M. D. T., Sothiselvam, S., Lu, T. K., Fuente-Nunez, C. de la., 2019. Peptide design principles for antimicrobial applications. *J. Mol. Biol.* 431, 3547-3567.
- White, J. K., Muhammad, T., Alsheim, E., Mohanty, S., Blasi-Romero, A., Gunasekera, S., Strömstedt, A. A., Ferraz, N., Göransson, U., Brauner, A., 2022. A stable cyclized antimicrobial peptide derived from LL-37 with host immunomodulatory effects and activity against uropathogens. *Cell. Mol. Life Sci.* 79, 411.

* Presenting author. *E-mail address:* taj.muhammad@fa.uu.se (Taj Muhammad)

S1.P249 A steroid-rich fraction from parasitic climber *Cocculus hirsutus* (L.) W.Theob inhibits coronavirus disease and neutrophilic inflammation

Vidya Febrasca Tenderly^{a*}, Thiyagarajan Raviraj^a, Yen Chi Loo^a, Olha Mykhailenko^b, Marcela Safratova^c, Stefan Kosturko^c, Mahendra Jadhao^d, Yu-Li Chen^e, Tsong-Long Hwang^f, Fang-Rong Chang^{a*}, Michal Korinek^a

^a Graduate Institute of Natural Products, College of Pharmacy, Kaohsiung Medical University, Kaohsiung, Taiwan

^b Department of Pharmaceutical Chemistry, National University of Pharmacy of Ministry of Health of Ukraine, Kharkiv, Ukraine

^c Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmacy, Charles University, Hradec Kralove, Czech Republic

^d Department of Cancer Biology, University of Cincinnati College of Medicine, Cincinnati, OH, USA

^e Research Center for Chinese Herbal Medicine, Graduate Institute of Health Industry Technology, College of Human Ecology, Chang Gung University of Science and Technology, Taoyuan, Taiwan

^f Graduate Institute of Natural Products, College of Medicine, Chang Gung University, Taoyuan, Taiwan

Infections from coronavirus diseases and their complications pose significant threats to human health, economies, and societies. Excessive inflammatory responses in human neutrophils are linked to high mortality rates (Chiang et al., 2020). There is a pressing need for affordable and safe treatments, prompting interest in herbal remedies.

Cocculus hirsutus (L.) W.Theob (CH, Menispermaceae) is a perennial climbing plant found widely across Asia and Africa. We have obtained aerial parts of CH from farm in India where it grows as weed and parasite, damaging crops. Notably, various parts of CH are used in folk medicine to treat fever, skin issues, stomach ailments, and urinary disorders. Alkaloid compounds are among major constituents identified in this plant (Logesh et al., 2020). However, other CH secondary metabolites are unexplored.

This study evaluated crude extracts, partitioned layers, and subfractions using SARS-CoV-2 spike/ACE2 pseudovirus neutralization assay (crude CH extract 27.16% and CHEA6 subfraction 54.85% inhibition at 10 µg/mL). Moreover, the anti-inflammatory assays including superoxide anion generation and elastase release in fMLF/CB-induced human neutrophils showed notable activity (crude CH extract 53.95% and 80.39%, respectively; CHEA layer and CHEA6 subfraction >90% inhibition for both assays at 10 µg/mL). Importantly, the bioactivity-guided fractionation of non-alkaloid subfraction CHEA6 (ethyl acetate fraction 6) yielded major steroid triterpenes, including ecdysone type, including 20-hydroecdysone (**1**), makisterone A (**2**), and phytosterols, such as β -sitosterol glucoside (**3**). Additionally, three compounds, including (+)-*proto*-quercitol (**4**), lutein (**5**), and formononetin (**6**) were isolated. The structure of the compounds was confirmed by 1D, 2D NMR, and mass spectroscopic analysis.




Fig. 1. Steroid-rich fraction isolation scheme, HPLC, and NMR chemical profile

Keywords: *Cocculus*, SARS-CoV-2, human neutrophils, triterpene, phytochemistry

References

- Chiang, C.-C., Korinek, M., Cheng, W.-J., & Hwang, T.-L. 2020. Targeting Neutrophils to Treat Acute Respiratory Distress Syndrome in Coronavirus Disease. *Frontiers in Pharmacology*, 11.
- Logesh, R., Das, N., Adhikari-Devkota, A., & Devkota, H. P. 2020. *Cocculus hirsutus* (L.) W.Theob. (Menispermaceae): A Review on Traditional Uses, Phytochemistry and Pharmacological Activities. *Medicines*, 7(11), 69.

*Presenting author. E-mail address: vfebrasca@gmail.com (Vidya Febrasca)

S1.P250 *Sargassum fusiforme* and its components inhibits Respiratory Syncytial Virus replication *in vitro* and *in vivo*

Kiramage Chathuranga, Jong-Soo Lee*

College of Veterinary Medicine, Chungnam National University, Daejeon 34134, Republic of Korea

Here, we demonstrate that *S. fusiforme* extract (SFE) has antiviral effects against the respiratory syncytial virus (RSV) *in vitro* and *in vivo* mouse models. *Sargassum fusiforme* is an edible brown alga widely distributed around the coastlines of China, Korea, and Japan, and this nutritious marine vegetable has been applied as a therapeutic in traditional Chinese medicine for thousands of years. It is a popular functional seaweed that can prolong life expectancy. (Chen et al., 2018) Treatment of HEp2 cells with non-cytotoxic concentrations (30, 50, or 100 µg/mL) of SFE significantly reduced RSV replication, RSV-induced cell death, RSV gene transcription, RSV protein synthesis, and syncytium formation. Moreover, oral inoculation of SFE significantly improved RSV clearance from the lungs of BALB/c mice.




Fig. 1. Anti-RSV effect of eicosane, docosane and tetracosane in Hep2 cells.

Previous reports identified several active components of SFE. Among these, eicosane, dotriacontane, tritetracontane, docosane, heptacosane, and tetracosane are major components of SFE and have been proposed to have antimicrobial activity (El Shafay et al., 2016). Interestingly, the phenolic compounds eicosane, docosane, and tetracosane were identified as active components of SFE. Treatment with a non-cytotoxic concentration of these three components elicited similar antiviral effects against RSV infection as SFE *in vitro*. Collectively, these results suggest that SFE and its potential components are a promising natural antiviral agent candidate against RSV infection, and consumption of SFE would have beneficial effects in the prevention or therapy of RSV infection.

Keywords: *Sargassum fusiforme*, therapeutic effects, RSV, eicosane, docosane, tetracosane

References

- Chen, P., Yang, S., Hu, C.; Zhao, Z.; Liu, J., Cheng, Y.; Wang, S., Chen, Q., Yu, P., Zhang, X, 2018. *Sargassum fusiforme* Polysaccharide Rejuvenates the Small Intestine in Mice Through Altering its Physiology and Gut Microbiota Composition. Curr. Mol. Med. 17, 1.
- El Shafay, S.M., Ali, S.S., El-Sheekh, M.M. 2016. Antimicrobial activity of some seaweeds species from Red sea, against multidrug resistant bacteria. Egypt. J. Aquat. Res. 42, 65–74.

Acknowledgments

The National Research Foundation of Korea (2021R1A6A1A03045495).

* Presenting author. E-mail address: jongsool@cnu.ac.kr (Jong-Soo Lee)

S1.P251 Inhibition of Respiratory Syncytial Virus *in vitro* and *in vivo* by *Apostichopus japonicus* (Selenka) extract

Kiramage Chathuranga, Jong-Soo Lee*

College of Veterinary Medicine, Chungnam National University, Daejeon 34134, Republic of Korea

Apostichopus japonicus (Selenka) a sea cucumber, is an invertebrate animal inhabiting the coastal sea around Korea, Japan, China, and Russia. It is one of the highest commercially valuable species of seafood, and has been commonly used for centuries in indigenous and folk medicine. *A. japonicus* is used in Chinese medicine to replenish the kidney, and supplement the essence, nourish the blood, and moisten dryness (Lu et al., 2023, Wang et al., 2021).

Although it has many therapeutic effects, its antiviral activity against Respiratory Syncytial Virus (RSV) has not been reported in detail. In this study, we show that extracts from *A. japonicus* (AJSE) have antiviral effects against RSV *in vitro* cell cultures and an *in vivo* mouse model. Treatment of a human respiratory tract cell line (HEp2) with a non-cytotoxic concentration of AJSE significantly reduced RSV replication, RSV-induced cell death, RSV gene transcription, and RSV protein synthesis.




Fig.1. AJSE abrogates RSV-mediated syncytial formation (A), and protein translation (B)

Additionally, the treatments significantly diminished syncytial formation after RSV infection in HEp2 cells. Time-dependent treatment of AJSE after RSV infection in HEp2 cells showed that treatment with two-hour post-infection of virus infections can provided a better result by demolishing further replication of the RSV virus in the Hep2 cell line. Interestingly, oral inoculation with AJSE significantly improved viral clearance in the lungs of BALB/c mice. Together, our results suggested that extracts of *A. japonicus* could be used as a potent natural anti-RSV candidate.

Keywords: *Apostichopus japonicus* (Selenka), RSV, therapeutic effects, virus replication

References

- Lu, C., Wang, X., Ma, J., Wang, M., Liu, W., Wang, G., Li, Y., 2023. Chemical substances and their activities in sea cucumber *Apostichopus japonicus*: A review. Arch. Pharm. 357, 2300427.
- Wang, C., Yao, L., Wang, W., Sang, S., Hao, J., Li, C., Zhang, Q., 2021. First report on natural infection of Nodavirus in an Echinodermata, sea cucumber (*Apostichopus japonicus*). Viruses, 13(4)-636.

Acknowledgments

The National Research Foundation of Korea (2021R1A6A1A03045495)

* Presenting author. E-mail address: jongsool@cnu.ac.kr (Jong-Soo Lee)

S1.P252 *Aster tataricus* and its components display a broad spectrum of antiviral effects *in vitro* and *in vivo*

Nuwan Gamage, Jong-Soo Lee*

College of Veterinary Medicine, Chungnam National University, Daejeon, Republic of Korea

This presentation will cover the antiviral activities and mechanisms of action associated with *Aster tataricus* both *in vitro* and *in vivo*. *Aster tataricus*, a member of the family Asteraceae, is a perennial terrestrial herb with a rich history in traditional medicine, native to Siberia, Korea, Japan, and eastern Asia (Shen et al., 2018). It is characterized by violet-blue flowers with a conspicuous yellow center (Chen et al., 2022). The investigation comprised cytotoxicity assessment, antiviral tests, mRNA analysis, and assessment of protein phosphorylation. In the context of *in vivo*, mRNA induction, viral challenge in mice, histopathological examination, and cytokine analysis followed by HPLC chemical characterization and compound antiviral testing.

Preliminary assessments indicate that *Aster tataricus* Extract (ATE) exhibits low toxicity with a CC_{50} of 134.0 ± 1.08 $\mu\text{g/ml}$, while an effective concentration of 10 $\mu\text{g/ml}$ is demonstrated marked inhibition of Influenza A virus (PR8), Newcastle disease virus (NDV), and Herpes simplex virus (HSV) replication in immune cells (RAW264.7). This inhibition was attributed to the induction of an antiviral state through the upregulation of interferon (IFN)-related genes, coupled with the secretion of IFNs and pro-inflammatory cytokines. *In vivo*, experiments with ATE-treated BALB/c mice revealed enhanced survivability and reduced lung viral titers when challenged with lethal doses of the highly pathogenic influenza A subtypes (H1N1, H5N2, and H9N2). The observed prophylactic effects correlated with increased secretion of IL-6, IFN- λ , and IFN- β in bronchoalveolar lavage fluid (BALF) of treated mice.




Fig. 1. Antiviral effect of *Aster tataricus* in RAW264.7 cells and structures of HPLC analysis identified active compounds in the aqueous fraction

High-performance liquid Chromatography (HPLC) analysis identified three main active compounds in the aqueous fraction. Subsequent evaluation highlighted the antiviral properties of Quercetin, Kaempferol, and Ferulic acid. This study confirms *Aster tataricus* and its components as potent broad-spectrum anti-viral and anti-influenza agents.

Keywords: *Aster tataricus*, antiviral effect, anti-influenza effect, Active compounds, Herbal medicine

References

- Shen X, Guo S, Yin Y, Zhang J, Yin X, Liang C, Wang Z, Huang B, Liu Y, Xiao S. 2018. Complete chloroplast genome sequence and phylogenetic analysis of *Aster tataricus*. *Molecules* 23:2426.
- Chen T, Yang P, Chen H-j, Huang B. 2022. A new biflavonoids from *Aster tataricus* induced non-apoptotic cell death in A549 cells. *Natural Product Research* 36:1409-1415.

Acknowledgments

The National Research Foundation of Korea (2021R1A6A1A03045495)

* Presenting author. E-mail address: jongsool@cnu.ac.kr (Jong-Soo Lee)

S1.P253 Extracts of *Costus speciosus* suppresses influenza A H1N1 activity *in vitro* and *in vivo*

Amal Senevirathne, Jong-Soo Lee*

College of Veterinary Medicine, Chungnam National University, Daejeon, Republic of Korea

This study demonstrates the antiviral activity of aqueous leaf extract of *Costus speciosus* (J. Koenig) Sm, (TB100) against influenza viruses. *Costus speciosus* (J. Koenig) Sm is a well-known medicinal herb used in traditional medicine in South Asian countries such as Sri Lanka, India, and other tropical regions (Waisundara et al., 2015). TB100, a perennial herb, contains many medicinal phytochemicals with potent antimicrobial, antioxidant, insecticidal, anticancer, and antidiabetic properties (Wang et al., 2015).

Initially, antiviral activity against green fluorescence-expressing influenza A (H1N1) PR8-GFP virus was assessed. EC₅₀ and CC₅₀ values for RAW264.7 cells were determined as 15.19 ± 0.61 and 117.12 ± 18.31 µg/mL, respectively. TB100 reduced GFP expression and viral copy number, confirming its inhibition of viral replication in murine RAW264.7 and human A549 and HEp2 cells. In vitro pre-treatment with TB100 induced phosphorylation of transcriptional activators TBK1, IRF3, STAT1, IKB-α, and p65, associated with interferon pathways, indicating activation of antiviral defences. The antiviral effects of TB100 against multiple influenza A strains (H1N1, H3N2, and H9N2) were investigated in BALB/c mice. TB100 demonstrated the highest protective response against H1N1, followed by H3N2 and H9N2.




Fig. 1. Chemical compounds in TB100 were analysed by the reversed-phase HPLC

High-performance liquid chromatography (HPLC) analysis of aqueous extracts led to the identification of cinnamic, caffeic, and chlorogenic acids as potential chemicals for antiviral responses. Further confirmatory studies using these acids revealed that each of them confers significant antiviral effects against influenza when used as pre-treatment and enhances the antiviral response in a time-dependent manner. These findings suggest that TB100 has the potential to be developed into an antiviral agent that is effective against seasonal influenza.

Keywords: *Costus speciosus*, leaf extract, TB100, influenza A, antiviral effect, transcriptional activator

References

- Waisundara, V.Y.; Watawana, M.I.; Jayawardena, N. *Costus Speciosus* and *Coccinia Grandis*: Traditional Medicinal Remedies for Diabetes. *South Afr. J. Bot.* 2015, 98, 1–5.
- Wang, Y.J.; Pan, K.L.; Hsieh, T.C.; Chang, T.Y.; Lin, W.H.; Hsu, J.T.A. Diosgenin, a Plant-Derived Sapogenin, Exhibits Antiviral Activity In Vitro against Hepatitis C Virus. *J. Nat. Prod.* 2011, 74, 580–584.

Acknowledgments

The National Research Foundation of Korea (2021R1A6A1A03045495)

* Presenting author. E-mail address: jongsool@cnu.ac.kr (Jong-Soo Lee)

S1.P254 Exploring faba bean pods as a potential nutritional supplement for piglet gut health

Inna Vlasova^{a,*}, Philip Krüsselmann^b, Yuliia Kostenko^a, Marcin Równicki^a, Jürgen Zentek^b, Jakub P. Piwowarski^a

^a Microbiota Lab, Department of Pharmaceutical Biology, Faculty of Pharmacy, Medical University of Warsaw

^b Institute of Animal Nutrition, Freie Universität Berlin, Berlin, Germany

One of the significant health concerns in the farm industry is post-weaning diarrhea (PWD) in piglets (Fairbrother et al., 2005). Due to a reduction in the use of antibiotics, it is necessary to find a source to uphold the gut health of piglets (Cheng et al., 2014). A potential source could derive from a by-product, such as faba bean pods (FBP, *Vicia faba* L.), which are rich in bioactive compounds (Valente et al., 2018). Our research focuses on nutritional profile, phytochemical analysis and biological activity of FBP.

Nutritional analysis revealed notable concentrations of crude protein (144 g/kg), potassium (27.8 g/kg), and iron (126 mg/kg) in dry FBP. Phytochemical analysis of methanolic extract (ME) showed significant levels of total polyphenols (38.43±0.59 mg gallic acid equivalents (GAE)/ g dry FBP), tannins (30.67±1.03 mg GAE/ g dry FBP), and procyanidins (0.66±0.13 mg procyanidin B1/g dry FBP). Utilizing a UHPLC-DAD-MSn method, 34 compounds were found in the ME; 16 were identified and classified as derivatives of quercetin and kaempferol, alongside levodopa and piscidic acid. The extract was also tested for inhibiting two common gastrointestinal pathogens. A dose-dependent growth inhibition of *Escherichia coli* ATTC25922 and a total elimination of visible growth of *Salmonella enterica* when exposed to the ME concentrations of 1–8 mg/mL was obtained. Additionally, the ME showed insignificant activity against porcine digestive enzymes (α -amylase, lipase, and trypsin) (IC₅₀>4mg/mL).

Our research demonstrates that FBP is a rich reservoir of varied polyphenols with antimicrobial properties, including the influence of FBP on porcine digestive enzymes. It is a promising feed additive for mitigating PWD in piglets.

Keywords: faba bean pods, food by-product, antimicrobials, post-weaning diarrhea, phytochemical analyses

References

- Fairbrother, J. M., Nadeau, É., Gyles, C. L., 2005. *Escherichia Coli* in Postweaning Diarrhea in Pigs: An Update on Bacterial Types, Pathogenesis, and Prevention Strategies. *Anim. Health Res. Rev.* 6(1), 17-39.
- Cheng, G., Hao, H., Xie, S., Wang, X., Dai, M., Huang, L., Yuan, Z., 2014. Antibiotic Alternatives: The Substitution of Antibiotics in Animal Husbandry? *Front. Microbiol.* 5.
- Valente, I. M.; Maia, M. R. G.; Malushi, N.; Oliveira, H. M.; Papa, L.; Rodrigues, J. A.; Fonseca, A. J. M.; Cabrita, A. R. J., 2018. Profiling of Phenolic Compounds and Antioxidant Properties of European Varieties and Cultivars of *Vicia Faba* L. Pods. *Phytochemistry*, 152, 223-229.

Acknowledgments

Project financially supported by Polish National Science Centre research grant OPUS LAP UMO-2020/39/I/NZ7/02547.

* Presenting author. E-mail address: inna.vlasova@wum.edu.pl (Inna Vlasova)

S1.P255 Anti-inflammatory properties of twelve Norwegian medicinal plants and their protective properties against impairments in the intestinal epithelial barrier

Andrea Angelov^{1,*}, Helle Wa-ngensteen¹, Marit Inngjerdengen²

¹ Section for Pharmaceutical Chemistry, Department of Pharmacy, University of Oslo, Oslo, Norway

² Department of Pharmacology, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

Chronic inflammation is linked to various diseases like cardiovascular diseases, cancer, and diabetes. There is a lack of satisfying treatments for these diseases and medicinal plants may represent sustainable and effective sources for new drugs or drug leads. Polyphenols are known to act on several immune signaling pathways and can modulate the production and release of cytokines and affect proliferation and differentiation of leukocytes. The aim of this study was to investigate the *in vitro* anti-inflammatory properties and the ability to counteract impairments in the intestinal epithelial barrier of polyphenol rich extracts from Norwegian medicinal plants. Based on the traditional use of plants towards various inflammatory diseases, twelve Norwegian plants were identified. The plant material was extracted with 80% ethanol and dichloromethane, and the extracts were analyzed by ¹H NMR spectroscopy. By stimulating a dendritic cell line to produce NO, the extracts' ability to inhibit NO secretion was determined. Furthermore, inhibition of TNF- α secretion was performed using ELISA with peripheral blood mononuclear cells. The ability to counteract impairments in the intestinal epithelial barrier was investigated by indomethacin treated Caco-2 cells. The results show that the most promising ethanol extracts were *Alnus incana* (59% reduction in NO, 100 μ g/mL; 79% reduction in TNF- α , 10 μ g/mL), *Antennaria dioica* (38% reduction in NO, 100 μ g/mL; 42% reduction in TNF- α , 10 μ g/mL), and *Geranium sylvaticum* (36% reduction in NO, 100 μ g/mL; 59% reduction in TNF- α , 10 μ g/mL). Among the dichloromethane extracts, *Lepidothea suaveolens* (100% reduction in NO, 100 μ g/mL; 81% reduction in TNF- α , 10 μ g/mL), *Malva moschata* (96% reduction in NO, 100 μ g/mL; 77% reduction in TNF- α , 10 μ g/mL), and *Tanacetum vulgare* (100% reduction in NO, 100 μ g/mL; 99% reduction in TNF- α , 10 μ g/mL) showed strongest effects. Results from Caco-2 barrier protection experiments will be presented. The NMR spectra indicate a high phenolic content in the active ethanolic extracts with a diverse content in structure type.

Keywords: Medicinal plants, polyphenols, anti-inflammatory, TNF- α , nitric oxide

* Presenting author. E-mail address: andrang@uio.no (Andrea Angelov)

S1.P256 Could Bistort rhizome serve as a potential alternative to antibiotics in preventing diarrhea in weaned pigs? *In vitro* study

Marcin Równicki^{a,*}, Inna Vlasova^a, Philip Krüsselmann^b, Yuliia Kostenko^a, Karolina A. Pawłowska^a, Sebastian Granica^a, Jürgen Zentek^b, Jakub P. Piwowarski^a

^a Microbiota Lab, Department of Pharmaceutical Biology, Medical University of Warsaw, 1 Banacha St., 02-097, Warsaw, Poland

^b Institute of Animal Nutrition, Department of Veterinary Medicine, Freie Universität Berlin, Berlin, Germany

Post-weaning diarrhea is a major threat in the swine industry, which results in the high use of antibiotics for its treatment. There is a global agenda to decrease the use of antibiotics in order to reduce the risk of antimicrobial resistance. Therefore, alternative methods to prevent the development of post-weaning diarrhea and maintain piglets' gut health are needed.¹

Bistort (*Bistorta officinalis*) rhizome (BR) has long been esteemed worldwide for its therapeutic properties, particularly in alleviating diarrhea. Its efficacy is attributed to its high tannin content, making it a popular choice in traditional medicine. However, in piglet nutrition, tannins are widely regarded as anti-nutritional factors.²

This study investigated the phytochemical and nutritional content of BR. Considering the attributed anti-nutritional properties and the potential application of BR as feed additives, the *in vitro* biological activity was evaluated, including the influence of BR on porcine digestive enzymes and antimicrobial activity against porcine gastrointestinal pathogens.

The UHPLC analysis of the aqueous extract of BR revealed 28 major compounds, mainly procyanidins and their galloylated derivatives. Extract from BR inhibited porcine amylase (IC₅₀= 0.05 mg/ml) and lipase (IC₅₀= 0.55 mg/ml) and had no impact on porcine trypsin (IC₅₀= 17.04 mg/ml). Antibacterial activity of the extract against major porcine intestinal pathogens *Salmonella enterica* and *Escherichia coli* in concentrations 0.25 – 8 mg/ml was also tested. Additionally, BR extract exhibited low cytotoxicity (IC₅₀= 330 µg/ml) on the intestinal porcine epithelial cell line (IPEC-J2). The obtained results provide basic evidence of the usage of BR as a novel feed additive in the prevention of post-weaning diarrhea in piglets.

Keywords: *Bistort rhizome*, *Bistorta officinalis*, antibacterials, cytotoxicity, gut health

References:

1. Xiong, X. *et al.* Nutritional Intervention for the Intestinal Development and Health of Weaned Pigs. *Front. Vet. Sci.* 6, 46 (2019).
2. Pawłowska, K. A. *et al.* Antibacterial and anti-inflammatory activity of bistort (*Bistorta officinalis*) aqueous extract and its major components. Justification of the usage of the medicinal plant material as a traditional topical agent. *J. Ethnopharmacol.* 260, 113077 (2020).

Acknowledgments

Project financially supported by Polish National Science Centre research grant OPUS LAP UMO-2020/39/I/NZ7/02547.

* Presenting author. E-mail address: marcin.rownicki@wum.edu.pl (Marcin Równicki)

S1.P257 Detection and insight into structural requirements of SARS_CoV2 antibodies by peptide- oligonucleotides derived from sunflower trypsin inhibitor 1 (SFTI-1) peptide

Sunithi Gunasekera^{a,*}, Hongxing Zhao^b, Camilla Eriksson^a, Mingshu Zhang^a, Taj Muhammad^a, Ulf Landegren^b, Ulf Göransson^a

^a Department of Pharmacognosy, Department of Pharmaceutical Biosciences, Box 591, Uppsala University, SE 75124, Uppsala, Sweden

^b Department of Immunology, Genetics and Pathology, Science for Life Laboratory, Uppsala University, Sweden

SARS_CoV2 virus represents the most devastating pandemic of modern times. As part of epidemiology studies, it has been crucial to understand the prevalence of antibodies following COVID-19 infection and vaccination, providing insight into protection against the virus. Recently Antibody proximity extension assay (AbPEA) was successfully used in SARS_CoV2 antibody detection (Zhao et al. 2022), offering high sensitivity with minimum samples volumes. However, the AbPEA typically utilizes complete viral antigenic protein domains (eg. spike protein subunit S and nucleocapsid protein), examining total antibody reactivity.

To enhance antibody detection specificity, we designed peptide-oligonucleotides containing short epitopes (10-15 residues), originating from immunodominant regions of the spike protein. These constructs were then employed in AbPEA to detect antibodies in convalescent sera and dry blood spots of vaccinated individuals. Two different counterpart epitopes were compared, in their linear form and incorporated within a cyclic scaffold derived from sunflower trypsin inhibitor 1 (SFTI-1) (Korsinczky et al. 2001; Eriksson et al. 2023). Peptides were synthesized by Fmoc-SPPS and their secondary structural characteristics were determined by ¹H NMR.

Several epitopes from S1 and spike protein subunit S2 regions were able to detect antibodies in seropositive samples. Notably, peptide conformation appeared to influence antibody binding for some epitopes derived from the S1 and S2 regions, but not for several other epitopes. Certain epitopes reacted with antibodies regardless of whether they were linear or cyclic, indicating that the conformation does not affect binding. Overall, we gained insight into heterogeneity in antibody profiles across patient samples as well as structural requirements of antibodies for antigen binding. This study highlights the potential for improving current SARS-CoV2 antibody detection methods to achieve greater specificity in antibody profiling. If these fine specificity profiles in patient samples can be correlated with clinical symptoms and disease severity, AbPEA will become an invaluable tool for better understanding of seroprevalence.

Keywords: SARS_CoV2, antibody proximity extension, seroprevalence, peptide, SFTI-1

References

- Eriksson C., Gunasekera S., Muhammad T., Zhang M., Lauren I., Mangsbo S.M., Lord M. & Göransson U. (2023) Epitopes Displayed in a Cyclic Peptide Scaffold Bind SARS-COV-2 Antibodies. *Chembiochem* 24, e202300103.
- Korsinczky M.L., Schirra H.J., Rosengren K.J., West J., Condie B.A., Otvos L., Anderson M.A. & Craik D.J. (2001) Solution structures by 1H NMR of the novel cyclic trypsin inhibitor SFTI-1 from sunflower seeds and an acyclic permutant. *J Mol Biol* 311, 579-91.
- Zhao H., Wang M., Muthelo P., Lof L., Sterky F., Gallini R., Kumar N.V., Monsen T., Nilsson K., Aberg M., Kamali- Moghaddam M., Mei Y.F. & Landegren U. (2022) Detection of SARS-CoV-2 antibodies in serum and dried blood spot samples of vaccinated individuals using a sensitive homogeneous proximity extension assay. *N Biotechnol* 72, 139-48.

* Presenting author. *E-mail address:* sunithi.gunasekera@farmbio.uu.se (Sunithi Gunasekera)

S1.P258 Chemical analyses and *in vitro* biological properties of *Lythrum salicaria* L. for potential use in piglet nutrition

Yuliia Kostenko^{a,*}, Alicja Zadarko^a, Inna Vlasova^a, Marcin Równicki^a, Philip Krüsselmann^b, Jürgen Zentek^b, Jakub P. Piwowski^a

^a Microbiota Lab, Department of Pharmaceutical Biology, Faculty of Pharmacy, Medical University of Warsaw

^b Institute of Animal Nutrition, Freie Universität Berlin, Berlin, Germany

Post-weaning diarrhoea (PWD) is a significant gastrointestinal disease affecting piglet production. Traditionally, PWD was treated by antibiotics, however this has contributed to the spread of antimicrobial resistance, making new therapeutic strategies urgently needed (Fairbrother et al., 2005). One possible approach to reduce the risk of PWD is the application of hydrolysable tannins, which were known for anti-diarrhoeal properties in the pre-antibiotic era. (Girard et al., 2020).

The present work aims to establish the phytochemical composition and biological activity of Purple loosestrife (*Lythrum salicaria* L.) aqueous extract (LSH) to assess the potential beneficial effects on gastrointestinal health of post-weaning piglets (Piwowski et al., 2013). Firstly, we found a high content of tannins (TTC=268.06 µg/mg) as dominating compounds among polyphenols (TPC=274.75 µg/mg) in LSH. Then, we verified the inhibitory activity of LSH against *Salmonella typhimurium* LT2, determining the minimal inhibitory concentration at 0.5 mg/mL and the dose-dependent effect on *Escherichia coli* ATCC 25922 in concentrations 0.5-2 mg/mL. We also determined cytotoxicity of LSH for the intestinal porcine epithelial cell line J2 (IC₅₀>100µg/mL). Additionally, we studied LSH interaction with porcine digestive enzymes. LSH had no effect on porcine trypsin with IC₅₀=27.45 mg/mL, and minimal effect on amylase with IC₅₀=2.65 mg/mL. However, we found significant activity against lipase with IC₅₀=0.33 mg/mL. To further investigate the bioactivity of the compounds found in LSH, we fractionated the LSH using column chromatography and spectroscopic methods. We isolated several hydrolysable tannins including vescalagin, castalagin, and salicarinin A, B, and C. Our results support the usage of LSH as novel feed additive in reducing the incidence of PWD in piglets.

Keywords: anti-microbial properties, hydrolysable tannins, *Lythrum salicaria* L., phytochemical composition, post-weaning diarrhoea

References

- Fairbrother, J. M., Nadeau, E. & Gyles, C. L., 2005. *Escherichia coli* in post-weaning diarrhea in pigs: an update on bacterial types, pathogenesis, and prevention strategies. *Anim Health Res Rev* 6, 17-39.
- Girard, M. & Bee, 2020. Invited review: Tannins as a potential alternative to antibiotics to prevent coliform diarrhea in weaned pigs. *G. Animal* 14, 95-107.
- Piwowski, J. P. & Kiss, A. K., 2013. C-glucosidic Ellagitannins from *Lythri herba* (European Pharmacopoeia): Chromatographic Profile and Structure Determination. *Phytochem Analysis* 24, 336-348.

Acknowledgments

Project financially supported by Polish National Science Centre research grant OPUS LAP UMO-2020/39/I/NZ7/02547.

* Presenting author. E-mail address: yuliia.kostenko@wum.edu.pl (Yuliia Kostenko)

S1.P259 Therapeutic potential of stilbene/ β -diketone/gingerol allied naturally occurring non- flavonoids in triple-positive breast cancer: tracing the path from extraction to efficacy

Lairikyengbam Deepti Roy*, Jyotsna Kumar

Department of Chemistry, Faculty of Mathematical and Physical Sciences, M.S. Ramaiah University of Applied Sciences, Bangalore-560058, Karnataka, India

Triple-positive breast cancer (TPBC) is a less aggressive type of breast cancer often treated with long-term oral medications, but resistance can occur (Wang et al., 2019). To alleviate this issue, researchers are exploring natural and therapeutic interventions, including polyphenols, which have excellent anticancer properties (Schmidt et al., 2020). However, only a few studies have explored the role of non-flavonoid polyphenols in TPBC, while the major focus was on ER+ or triple-negative BC.

The present investigation aimed to evaluate the antiproliferative activity of three distinct non- flavonoids derived from different plant sources – resveratrol with a stilbene moiety from *Vitis vinifera*, curcumin with a β -diketone moiety from *Curcuma longa*, and 6-gingerol with a gingerol moiety from *Zingiber officinale* against two breast cancer cells: TPBC-BT-474 and TNBC-MDA- MB-231 via MTT assay (Mosmann 1983). Plant extracts of dried+sieved grapes, turmeric, and ginger powder were prepared via reflux extraction using ethanol and water mixture. Isolation and purification of respective polyphenols were carried out by Flash chromatography and HPLC/MS. Structures were elucidated by ^1H -NMR, ^{13}C -NMR, FT-IR, and HPLC/MS.

Among the three natural polyphenols, resveratrol showed superior antiproliferative activity against TPBC-BT-474 compared to TNBC-MDA-MB-231 cells. The other two polyphenols isolated from the rhizome family – curcumin and 6-gingerol showed better inhibiting properties against MDA-MB-231 cells. The chemopreventive effect of polyphenols is attributed to their excellent anti- inflammatory properties, especially through their antioxidant activity. The study highlights the potential of non-flavonoid polyphenols, particularly resveratrol (comprising stilbene moiety) as a scaffold for developing novel therapeutic derivatives for TPBC suggesting that their synergistic effect with anticancer drugs could enhance future treatments.

Keywords: non-flavonoids, reflux extraction, MTT, BT-474 cells, anti-breast cancer

References

- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* 65, 55-63.
- Schmidt, B., Ferreira, C., Passos, C.L.A., Silva, J.L., Fialho, E., 2020. Resveratrol, curcumin and piperine alter human glyoxalase 1 in MCF-7 breast cancer cells. *Int. J. Mol. Sci.* 21, 5244.
- Wang, J., Xu, B., 2019. Targeted therapeutic options and future perspectives for HER2-positive breast cancer. *Sig. Transduct. Target Ther.* 4, 34.

* Presenting author. E-mail address: deepti07roy@gmail.com (Lairikyengbam Deepti Roy)

S1.P260 Drug discovery based on essential oils concurrently active against eumycetoma and actinomycetoma with *Pimpinella anisum* emerged as a forerunner

Shereen Abd Algaffar^{a,b,*}, Prabodh Satyal^c, Naglaa Ashmawy^{d,e}, Annelies Verbon^b, Wendy van de Sande^b, Sami Khalid^a

^a Faculty of Pharmacy, University of Science and Technology, Omdurman, Sudan

^b Erasmus MC, University Medical Center Rotterdam, Department of Medical Microbiology & Infectious Diseases, Dr. Molewaterplein 40, 3015 GD, Rotterdam, The Netherlands

^c Aromatic Plant Research Center, 230N 1200E, Lehi, UT 84043 USA

^d Department of Pharmaceutical Sciences, College of Pharmacy, Gulf Medical University, Ajman P.O. Box 4184, United Arab Emirates

^e Department of Pharmacognosy, Faculty of Pharmacy, Ain Shams University, Abbassia, 11566, Cairo, Egypt

Mycetoma is a neglected infection of the subcutaneous tissues. It is caused by fungal (eumycetoma) and bacterial (actinomycetoma) pathogens (Relhan et al., 2017). There is an urgent need to develop novel, effective, broad-spectrum antimycetomal agents. Treatment differs for both entities, and a diagnostic tool differentiating eumycetoma from actinomycetoma is limited and unaffordable. Therefore, discovering drugs concurrently acting on both types would circumvent the diagnosis to initiate treatment promptly. Eight essential oils (EOs) from diverse taxonomical origins belonging to five botanical families were screened for antimicrobial activity against *Madurella mycetomatis* and *Actinomyces madurae* strains. EOs were prepared by steam distillation, tested *in vitro* against mycetoma strains (Abd Algaffar et al., 2021, 2022), and subsequently subjected to gas chromatography, mass spectrometry, and radical scavenging activity (RSA). The most promising EOs of *Myristica fragrans* L. (Myristicaceae) and *Pimpinella anisum* L. (Apiaceae) demonstrated dual *in vitro* activity against all tested strains with MICs of 0.004 to 0.125% v/v, and DPPH inhibition of 90.30±0.008 to 98.00±0.003 %RSA±SD, respectively. Both EOs were further tested for *in vivo* toxicity and efficacy in *Galleria mellonella* larvae models (Konings et al., 2021). *P. anisum* EO enhanced the survival of *M. mycetomatis*-infected *Galleria* larvae, with no signs of toxicity to uninfected larvae. A tendency towards enhancing survival was noted for *A. madurae*-infected larvae. The use of these EOs as broad-spectrum dual antimycetomal agents, could cutting diagnosis cost and time and subsequently resulting in an early treatment. This approach could possibly pave the way for the next generation of antimycetomal agents.

Keywords: *Madurella mycetomatis*, *Actinomyces madurae*, essential oils, *Galleria mellonella*, metabolomics

References

- Abd Algaffar SO, Verbon A, vande Sande WWJ, Khalid SA. 2021. Development and validation of an in vitro resazurin- based susceptibility assay against *Madurella mycetomatis*. Antimicrob Agents Chemother. 65: e01338-20.
- Abd Algaffar SO, Verbon A, Khalid SA, van de Sande WWJ. 2022. Development and validation of a resazurin assay for in vitro susceptibility testing of *Actinomyces madurae*: a common causative agent of actinomycetoma. J Antimicrob Chemother. 78,155-160.
- Konings M, Eadie K, Lim W, Fahal AH, Mouton J, Tesse N, van de Sande W. 2021. The synthetic synergistic cinnamon oil CIN-102 is active against *Madurella mycetomatis*, the most common causative agent of mycetoma. PLoS Negl Trop Dis. 15(6).
- Relhan V, Mahajan K, Agarwal P, Garg VK. 2017. Mycetoma: an update. Indian J Dermatol. 62(4):332.

* Presenting author. E-mail address: phd_sh086@hotmail.com (Shereen Abd Algaffar)

S1.P261 Synthesis of autoinducer-2 prodrugs and analogues as a new antimicrobial strategy

Peter Kis^{a,b}, Miguel V. Rodrigues^a, Karina B. Xavier^c, M. Rita Ventura^{a,*}

^a Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Av. da República, 2780-157 Oeiras, Portugal

^b Institute of Chemistry, Slovak Academy of Sciences, 84538 Bratislava, Slovakia

^c Instituto Gulbenkian de Ciência, 2780-156 Oeiras, Portugal

Many species of bacteria control gene expression on a community-wide scale by producing, secreting, detecting and responding to extracellular signaling molecules that accumulate in the environment in proportion to cell density. This process is called quorum-sensing and enables these organisms to coordinate their behaviours which are more beneficial when cells are working in unison.^[1,2] Autoinducer-2 (AI-2, Fig.1) is one of the extracellular signal molecules and is unique as it is produced and detected by many phylogenetically distinct bacteria.^[1,2]

In past years we have developed the synthesis of (S)-4,5-dihydroxypentane-2,3-dione (DPD 1, Fig.1),^[3] which is the AI-2 precursor, and analogues in optically pure form.^[2,4-6] Based on previous results, we hypothesise that AI-2 plays an important role in controlling colonisation and homeostasis of mammalian gut microbiota contributing to the protective properties of these poly-species against pathogens.^[1] This community of bacteria is crucial for production of nutrients, maturation of the immune system and host protection against pathogens. Therefore, the ability to drive this community from disease to healthy states by manipulating the native signals and interactions occurring between its members offers great potential for therapeutic benefit. AI-2 analogues have high potential as a complementary therapy that can avoid the use of antibiotics, contributing to the fight against the development of antibiotic resistance, a worldwide problem.




Fig. 1. Structures of linear and two cyclic forms of Autoinducer-2 (AI-2)

Here we present the synthetic strategies to obtain new analogues and prodrugs of AI-2 and the results of their biological evaluation.

Keywords: autoinducer 2, organic synthesis, quorum sensing, antibiotic resistance

References

- [1] Pereira, C.S.; Thompson, J.A.; Xavier, K.B. *FEMS Microbiology Reviews* 2013, 37(2), 156.
- [2] Rodrigues, M. V.; Kis, P.; Xavier, K. B.; Ventura, M. R. *Is. J. Chem.* 2023, e202200091.
- [3] Ascenso, O. S.; Marques, J. C.; Santos, A. R.; Xavier, K. B.; Ventura, M. R.; Maycock, C. D. *Bioorg. Med. Chem.* 2011, 19, 1236.
- [4] Ascenso, O. S.; Carrau, G.; Xavier, K. B.; Ventura, M. R.; Maycock, C. D. *Molecules* 2021, 26, 369.
- [5] Miranda, V.; Torcato, I. M.; Xavier, K. B.; Ventura, M. R. *Bioorganic Chemistry* 2019, 92, 103200.
- [6] Ascenso, A. S.; Torcato, I. M.; Miguel, A. S.; Marques, J. C.; Xavier, K. B.; Ventura, M. R.; Maycock, C. D. *Bioorganic Chemistry* 2019, 85, 75.

* Presenting author. E-mail address: rventura@itqb.unl.pt (M. Rita Ventura)

S1.P262 Bioactive compounds, antibacterial and antioxidant activities of methanol extract of *Axonopus compressus* (Sw.) P. Beauv

Olowe Olaoluwa Ayomikun*, Olajuyigbe Olufunmiso Olusola, Adefisoye Martins Ajibade, Adeoye-Iisjola Morenike

Department of Microbiology, School of Science Technology, Babcock University Ilisan-Remo, Ogun State, Nigeria

This presentation will cover the various phytochemical constituents, antibacterial, antioxidant activities present in the methanol extract of *Axonopus compressus* responsible for its pharmacological activities. *Axonopus compressus* is a perennial, terrestrial stem compressed grass with a bearded or hairy nodes belonging to the family Poaceae [Cabi, 2019] commonly known as carpet grass. *Axonopus compressus* is widely used in the treatment of a wide range of infectious diseases by a large number of locals [Tow et al., 2018].

It has become very crucial to explore natural compounds due to their therapeutic potential in the treatment of infections [Vaou et al., 2021]. The extract was obtained through standardized maceration methods and subjected to various phytochemical screening tests, gas chromatography-mass spectrometry analysis to determine the chemical compounds present and the antibacterial activity was tested against some pathogens invitro.

The phytochemical screening tests revealed the presence of tannins, saponins, flavonoids, phenols in varying concentrations. The result of GC-MS analysis revealed the presence of 61 bioactive compounds in the extract with peak area such 1,2, 3 – Benzentriol (20.69%), gamma. -Sitosterol (10.89%), 9,12-Octadecadienoic acid (Z, Z) (6.27%), 5-Hydroxymethylfurfural (6.62%) with different activities such as antibacterial, anti-inflammatory, anticancer. Reports on antioxidant activity indicated a notable rise with increasing plant concentration. The extracts demonstrated inhibitory potential against all tested pathogen.

GC-MS analysis revealed the presence of the bioactive constituents of the plants with various biological activities and this justifies the folkloric use of this plant as a therapeutic remedy by a large number of locals.

Keywords: *Axonopus compressus*, phytochemical constituents, antioxidant activity

References

- CABI. *Axonopus compressus* (carpet grass) <https://www.cabi.org/isc/datasheet/8094> (2019)
- Tow SWT, Eng ZX, Wong SP, Ge L, Tan SN, Yong JWH. 2018. *Axonopus compressus* (Sw.) Beauv.: a potential biomonitor for molybdenum in soil pollution. *International Journal of Phytoremediation* 20:1363–1368.
- Vaou N, Stavropoulou E, Voidarou C, Tsigalou C, Bezirtzoglou E. Towards Advances in Medicinal Plant Antimicrobial Activity: A Review Study on Challenges and Future Perspectives. *Microorganisms*. 2021 Sep 27;9(10):2041. doi: 10.3390/microorganisms9102041.

* Presenting author. E-mail address: olowe0376@pg.babcock.edu.ng (Olowe Olaoluwa Ayomikun)

S1.P263 The anti-inflammatory and pro-resolving role of selected South African medicinal plants used traditionally as immunomodulatory remedies

Gugulethu P. Khumalo^{a,d,*}, Ben-Erik Van Wyk^b, Yunjiang Feng^c, Ian E. Cock^a, Caroline Maake^d

^a School of Environment and Science, Nathan Campus, Griffith University, 170 Kessels Rd, Nathan, Queensland 4111, Australia

^b Department of Botany and Plant Biotechnology, University of Johannesburg, P.O. Box 524, Auckland Park 2006, Johannesburg, South Africa

^c Griffith Institute for Drug Discovery, Griffith University, 46 Don Young Rd, Nathan, Queensland 4111, Australia

^d Institute of Anatomy, University of Zürich, Winterthurerstr. 190, 8057 Zürich, Switzerland

There is a growing interest in the use of natural products as alternative and/or combination therapy against inflammatory conditions. The use of medicinal plants to treat auto-immune disorders is a common practice in southern Africa and a primary source of healthcare for many individuals. However, the pharmacodynamic and pharmacokinetic properties related to the traditional use of herbal remedies as immunomodulatory agents remain relatively unknown. This study investigates the anti-inflammatory properties of extracts prepared from the leaves of *Warburgia salutaris* (G. Bertol.) Chiov., and the bark of *Pterocelastrus rostratus* (Thunb.) Walp., widely used traditionally to treat inflammation and pain. The inhibitory effect of aqueous and ethanol extracts as well as fractions on the release of pro-inflammatory cytokines was determined in lipopolysaccharide (LPS) stimulated and unstimulated RAW 264.7 murine macrophage cells. The levels of interleukin (IL)-1 β , IL-6, tumour necrosis factor- α (TNF- α), and monocyte chemoattractant protein 1 (MCP-1) release were determined using cytokine multiplex-bead assays. The inhibitory properties of extracts and fractions were also evaluated on the release of prostaglandin E2 (PGE2), leukotriene-B4 (LTB4), cyclooxygenase-2 (COX-2), and nuclear factor-kappa B (NF- κ B) in LPS-stimulated RAW 264.7 cells. The ethanol extracts and fractions of *W. salutaris* and *P. rostratus* demonstrated the greatest inhibitory activity, with over 50-fold inhibition of IL-1 β , IL-6 and TNF- α levels in LPS-stimulated RAW 264.7 macrophages without toxicities in cell viability assays. These findings give an insight into understanding the mechanism of action of widely used southern African medicinal plants against chronic pain and rheumatological diseases.

Keywords: medicinal plants, inflammation, cytokines, cyclooxygenases, southern Africa

References

- Van Wyk, B.-E., Gericke, N., 2018. *People's Plants: A Guide to Useful Plants of Southern Africa*, 2nd edition. Briza Publications, Pretoria.
- Elgorashi, E.E., McGaw, L.J., 2019. African plants with in vitro anti-inflammatory activities: A review. *South Afr. J. Bot.* 126, 142–169.
- Khumalo, G.P., Van Wyk, B.E., Feng, Y. and Cock, I.E., 2022. A review of the traditional use of southern African medicinal plants for the treatment of inflammation and inflammatory pain. *J. Ethnopharmacol.* 283, p.114436.

* Presenting author. E-mail address: khumalo.gugulethu@uzh.ch (Gugulethu Khumalo)

S1.P264 Phytochemical analysis and antiplasmodial activity of *Pseudolachnostylis maprouneifolia* fruits extracts

Patricia Mbombo Mungitshi^{a,b,*}, Carla Hamann^a, Allison Ledoux^a, Paulin Mutwale Kapepula^b, Alembert Tchinda^c, Nadege Ngombe Kabamba^b, Michel Frederich^a

^a Laboratory of Pharmacognosy, Centre Interdisciplinaire de Recherches sur les Médicaments(CIRM), University of Liege, Liege , Belgium

^b Centre d'Etude des Substances Naturelles d'Origine Végétale, Faculty of Pharmacy, University of Kinshasa, Department of Medicinal Chemistry and Pharmacognosy, Kinshasa, Democratic Republic of the Congo

^c Laboratory of Phytochemistry, Institute Medicinal Research and Medicinals Plant Studies (IMPM), Yaounde, Cameroon

Pseudolachnostylis maprouneifolia, a member of the Phyllanthaceae family, is recognized for its distinctive botanical features and potential medicinal properties. In this study, we conducted a comprehensive phytochemical screening and evaluation of the antiplasmodial activity of a methanolic extracts derived from the fruits of *Pseudolachnostylis maprouneifolia*, sourced from Katanga in the Democratic Republic of the Congo. The extracts exhibited significant antiplasmodial activity against the chloroquine-sensitive *Plasmodium falciparum* strain (3D7), with an IC₅₀ value of 4.56 ± 0.19 µg/mg, underscoring their potential as a source of antimalarial compounds (Jansen et al., 2010).

Using a multidisciplinary analytical approach, we employed thin-layer chromatography (TLC) and high-performance liquid chromatography (HPLC) for component identification, followed by fractionation using column chromatography (CC). While gallic acid and methyl gallate were identified as key bioactive compounds through nuclear magnetic resonance (NMR) spectroscopy (Arsianti et al., 2018), ongoing fractionation efforts acknowledge the potential contribution of additional constituents to the observed antiplasmodial activity (Degotte et al., 2021, Arsianti et al., 2018). The goal of this study is to fully understand the chemical makeup of fruit extracts from *Pseudolachnostylis maprouneifolia* and find new antimalarial drugs that show in different mechanisms of action.

Our findings highlight the therapeutic potential of *Pseudolachnostylis maprouneifolia* as a valuable botanical resource for the development of antimalarial agents and underscore the importance of continued exploration to harness its full pharmacological potential.

Keywords: *Pseudolachnostylis maprouneifolia*, Gallic acid, Methyl gallate, Malaria, NMR

References

- Arsianti A, Astuti H, Fadilah F, Simadibrata D. M, Adyasa Z. M, Amartya D, Bahtiar A, Tanimoto H, Kakiuchi K. Synthesis and in Vitro Antimalarial Activity of Alkyl Esters Gallate as a Growth Inhibitors of Plasmodium Falciparum. Orient J Chem 2018;34(2).
- Degotte G, et al. Potential of Caffeic Acid Derivatives as Antimalarial Leads. Letters in Drug Design and Discovery, 2021, 18.
- Jansen O, Angenot L, Tits M, Nicolas JP, De Mol P, Nikiéma JB, Frédérick M. Evaluation of 13 selected medicinal plants from Burkina Faso for their antiplasmodial properties. J Ethnopharmacol. 2010 Jul 6;130(1):143-50.

* Presenting author. E-mail address: pmbombo@uliege.be (Patricia Mbombo Mungitshi)

S1.P265 Reconstruction of an ancient treatment of infections by incense

Selina Grollitsch^{a,*}, Johannes-Paul Fladerer^{a,b}, Franz Bucar^a

^a University of Graz, Institute of Pharmaceutical Sciences, Department of Pharmacognosy, Beethovenstraße 8, 8010 Graz, Austria

^b Apomedica Pharmazeutische Produkte GmbH, Roseggerkai 3, 8010 Graz, Austria

The increasing number of antimicrobial resistances (AMRs) is one of the major health challenges nowadays. Effective strategies for management of AMRs include revitalizing already existing antimicrobials and combination therapies. In ancient drug books written in the 17th century in Austria herbal combinations have been used as incense to treat infectious diseases. Within of that treatment, incense was burned and smoked through sheets. Afterwards, the sheets were put on wounds or wombs in case of urinary infections. The aim of this work was to reconstruct and proof the described treatment. Therefore, an apparatus containing Erlenmeyer flasks, heater and hoses was created to fumarate herbal mixtures and draw the smoke through a solvent by vacuum. In short, herbs were burned in an Erlenmeyer flask on a heater and the fume was suck via vacuum through a hose in a second flask containing a solvent (10% DMSO in water). The hose transporting the smoke ended in the solvent. A second hose reaching into the flask was connected to the vacuum and drew the smoke through the solvent. To contain 1 ml of extract 2.56 mg dried material were burned. The resulting extracts were tested against MRSA by microdilution assays using a time-kill methodology (Fladerer et al., 2023). One formulation containing eggshell, coriander, silver birch and myrrh significantly reduced the growth of MRSA ($p < 0.05$) pointing towards the effectivity of this 300-year-old treatment against bacterial infections. As a result, we were able to develop a new method to obtain extracts from incense formulations and to prove their antimicrobial activity against MRSA.

Keywords: Antimicrobial resistance, MRSA, incense, method development

References

Fladerer, J.P., Grollitsch, G., Bucar, F., 2023. SMS: A new screening method for synergism to test multi-herbal drug interactions for their antimicrobial activity. *Phytochem. Lett.* 57, 195-199.

* Presenting author. E-mail address: selina.grollitsch@uni-graz.at (Selina Grollitsch)

S1.P266 Chelation of metal ions by tannins: from protolytic equilibrium to antibacterial activity

Franjo Frešer^{a,*}, Matja Zalar^a, Urban Bren^{a,b,c}, Gregor Hostnik^a

^a Faculty of Chemistry and Chemical Technology, University of Maribor, Smetanova 17, SI-2000 Maribor, Slovenia

^b Faculty of Mathematics, Natural Sciences and Information Technologies, University of Primorska, Glagoljaška 8, SI-6000 Koper, Slovenia

^c Institute of Environmental Protection and Sensors, Beloruska 7, SI-2000 Maribor, Slovenia

Ellagitannins are a subgroup of tannins, ubiquitous secondary polyphenolic plant metabolites. They are technologically important compounds with several health-enhancing properties like antibacterial activity. Essential metal ions depletion is one of the major mechanisms of antibacterial action (Serrano et al., 2009). We studied the interactions of six ellagitannins (vescalin, castalin, vescalagin, castalagin, roburin A and D), isolated from chestnut wood extract, with Fe²⁺ and Al³⁺ ions. The main goal of the present study was to determine, how structural motifs of ellagitannins (NHTP and HHDP groups; green and orange in Fig. 1) affect the stoichiometry of complexes formed. This was performed using UV-Vis spectroscopy in conjunction with Job's method. Moreover, the effect of pH on tannins affinity for metal ions was examined. Since the tannin-metal ions interactions strongly depend on pH (Frešer et al., 2021; Frešer et al., 2024), the detailed study of ellagitannins microscopic protolytic equilibria was performed using a combination of ¹³C NMR and DFT methods. The results provided us with information on exact deprotonation positions, as well as pK_a constants. This enabled us to develop a model that explains the observed ellagitannin-metal ions interactions and can predict the stoichiometry and metal affinity of new ellagitannins, consisting of the NHTP and HHDP groups. The significance of the study does not end here because insight into ellagitannins' microscopic protolytic equilibrium is crucial for understanding their other properties, such as their bioavailability, reactivity, interactions with proteins and pH-dependent oxidative stability.



Fig. 1. Chemical structure of vescalagin (R₁=OH, R₂=H) and castalagin (R₁=H, R₂=OH)

Keywords: ellagitannins, metal ions chelation, proteolytic equilibrium, Job's method, NMR

References

- Frešer, F., Hostnik, G., Tošović, J., Bren, U., 2021. Dependence of the Fe(II)-Gallic Acid Coordination Compound Formation Constant on the pH. *Foods*, 10, 2689.
- Frešer, F., Bren, U., Hostnik, G., 2024. Chelation of iron(II) ions by ellagitannins—Effects of hexahydroxydiphenoyl and nonahydroxytriphenoyl groups. *Spectrochim. Acta A*, 313, 124079.
- Serrano, J., Puupponen-Pimiä, R., Dauer, A., Aura, A.-M., Saura-Calixto, F., 2009. Tannins: Current knowledge of food sources, intake, bioavailability and biological effects. *Mol. Nutr. Food Res.*, 53, S310–S329.

* Presenting author. E-mail address: franjo.freser@um.si (Franjo Frešer)

S1.P267 Effect of pH on UV/Vis and CD spectrum of cohumulone and colupulone

Gregor Hostnik^{a,*}, Anja Petek^a, Urban Bren^{a,b,c}

^a Faculty of Chemistry and Chemical Engineering, University of Maribor, Smetanova ulica 17, SI- 2000 Maribor, Slovenia

^b Faculty of Mathematics, Natural Sciences and Information Technologies, University of Primorska, Glagoljaška 8, SI- 6000 Koper, Slovenia

^c Institute for Environmental protection and Sensors, d.o.o., Beloruska ulica 7, SI-2000 Maribor, Slovenia

This presentation will cover the isolation and characterization of cohumulone and colupulone from hop (*Humulus lupulus* L.) cones. The isolated compounds are of utmost technological importance because especially cohumulone is the source of the bitter beer taste, while they are researched also for other health-enhancing effects, such as antimicrobial (Kolenc et al., 2023), diuretic, and progestogenic properties. (Arruda et al. 2021; Zanolini et al., 2008)




Fig. 1. Comparison of experimental and calculated spectrum of cohumulone and visualisation of orbitals taking place in HOMO→LUMO transition.

The compounds were dissolved in aqueous buffers of various pH values. The corresponding dependence of cohumulone and colupulone UV/Vis spectrum was measured. Additionally, the CD spectrum of cohumulone various species was examined. The UV/Vis spectrum of all possible species of cohumulone and colupulone species was calculated using TD-DFT on B3LYP level of theory and 6—311++G(d,p) basis set in the implicit CPCM water model. Additionally, CD spectra of all possible rotamers of cohumulone were calculated and compared to the experimental spectra. Next to observing which spectral transitions are responsible for the experimental absorption spectrum, the used method also enabled us to examine, which species are most likely present in the solution. What is more, the stereoconfiguration of cohumulone was examined and confirmed to be in good agreement with the stereoconfiguration already determined (Urban et al. 2013) – demonstrating that combination of CD spectroscopy and TD-DFT can relatively easily provide information on absolute stereoconfiguration.

Keywords: cohumulone, colupulone, stereoconfiguration, CD-spectroscopy, TD-DFT

References

- Arruda, T.R., Pinheiro, P.F., Silva, P.I., Bernardes, P.C., 2021. A New perspective of a well-recognized raw material: Phenolic content, antioxidant and antimicrobial activities and a- and b-acids profile of Brazilian hop (*Humulus lupulus* L.) extracts. *LWT*. 141, 110905.
- Kolenc, Z., Langerholc, T., Hostnik, G., Ocvirk, M., Štumpf, S., Pintarič, M., Košir, I.J., Černak, A., Garmut, A., Bren, U. 2023. Antimicrobial Properties of Different Hop (*Humulus lupulus*) Genotypes. *Plants*. 12, 120.
- Urban, J., Dahlberg, C.J., Carroll, B.J., 2013. Absolute Configuration of Beer's Bitter Compounds. *Angew. Chem.* 52, 1553-1555.
- Zanolini, P., Zavetti, M. 2008. Pharmacognostic and Pharmacological Profile of *Humulus lupulus*. *J. Ethnopharmacol.* 116, 383-396.

* Presenting author. E-mail address: gregor.hostnik@um.si (Gregor Hostnik)

S1.P268 Inhibitory activity of *Vitex negundo* (Lagundi) leaf extracts against biofilm formation and quorum sensing of *Vibrio cholerae*

Franco Miguel Nodado^a, Adrienne Marrie Bugayong-Janagap^b, Yoo Ki Bastol^a, Leopoldo Bucquial^a, Zeb Jose Joshua Cabalfin^a, Kathleen Mae Donoso^a, Danielle Elevado^a, Nove Kris Herida^a, Jemma Dawn Leonardo^a, Angel Mae Malones^a, Maio Vendell Sabidong^a

^a West Visayas State University, College of Medicine, La Paz, Iloilo City, Philippines

^b University of the Philippines Visayas, Division of Biological Sciences, Miagao, Iloilo, Philippines

Cholera, endemic in 69 countries including the Philippines, affects 1.3 billion individuals, with antibiotic use threatening antibiotic-resistant strains due to the biofilm formation of *Vibrio cholerae* (Almagro-Moreno et al., 2015; Yoon et al., 2019). The bacterium's virulence is regulated by quorum sensing (QS), controlled by LuxO, which is regulated by *hapR* and *qrr4*, quantified in this study (Watve et al., 2020; Hema et al., 2017). Lagundi (*Vitex negundo*), a traditional Philippine cholera treatment, is investigated for anti-QS and anti-biofilm properties (Basri et al., 2014; WIPO, 2015; Haq et al., 2012; Dwivedi et al., 2021).

This study determined the efficacy of *V. negundo* leaf methanolic extracts against *V. cholerae* by measuring the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC), assessing antibiofilm activity against both *V. cholerae* and *Chromobacterium violaceum*, and evaluating anti-QS activities against *C. violaceum*. Quantifying gene expression of QS-regulating *qrr4* and *hapR* was also conducted. Positive control used is gentamicin.

Lagundi leaf extracts showed promise against cholera-causing *V. cholerae*. The minimum concentration needed to inhibit (MIC) and kill (MBC) *V. cholerae* was 7.81 and 125 mg/mL, respectively. Lagundi extracts, especially at 500 mg/mL, significantly reduced biofilm formation in both *V. cholerae* and a standard bacterium (*C. violaceum*). Interestingly, this concentration even seemed more effective than gentamicin against biofilms in *C. violaceum*. Lagundi extracts also inhibited *C. violaceum*'s cell-to-cell communication (quorum sensing) in a dose-dependent manner, with an inhibition zone similar to gentamicin (0.8 cm vs 1.5 cm). However, Lagundi appears to work through a different mechanism than directly suppressing a key QS gene (*qrr4*).

These findings underscore the therapeutic potential of *V. negundo* against *V. cholerae*. Mechanistic studies are necessary to identify bioactive compounds, optimize dosing regimens, and assess pharmacokinetics. Exploring synergistic combinations with existing antimicrobials are essential.

Keywords: *Vitex negundo*, *Vibrio cholerae*, quorum sensing, biofilm formation, gene expression

References

- Almagro-Moreno S, et al. Intestinal Colonization Dynamics of *Vibrio cholerae*. PLoS Pathog. 2015 May 21;11(5):e1004787.
- Basri F, et al. A Review Of Ethnomedicinal Plant- *Vitex negundo* Linn. International Journal of Advanced Research (IJAR). 2014;2
- Dwivedi MK, et al. Evaluation of ethnopharmacologically selected *Vitex negundo* L. for In vitro antimalarial activity and secondary metabolite profiling. J Ethnopharmacol. 2021 Jul 15;275:114076.
- Haq RU, et al. Antitussive and toxicological evaluation of *Vitex negundo*. Nat Prod Res. 2012;26(5):484-8.
- Hema M, et al. Modulating the Global Response Regulator, LuxO of *V. cholerae* Quorum Sensing System Using a Pyrazine Dicarboxylic Acid Derivative (PDCApy): An Antivirulence Approach. Front Cell Infect Microbiol. 2017 Oct 12;7:441.
- Watve S, et al. Parallel quorum-sensing system in *Vibrio cholerae* prevents signal interference inside the host. PLoS Pathog. 2020 Feb 14;16(2):e1008313.
- World Intellectual Property Organization. From Herbal Folklore to Modern Medicine [Internet]. Philippines: National Integrated Research Program on Medicinal Plants; 2013
- Yoon SH, Waters CM. *Vibrio cholerae*. Trends Microbiol. 2019 Sep;27(9):806-807.

* Presenting author. E-mail address: francomiguel.nodado@wvsu.edu.ph (Franco Miguel Nodado)

S1.P269 Exploring the role of tannin-metal ion complex formation in antimicrobial activity

Sara Štumpf^a, Marija Djurič^b, Sara Seršen^b, Miha Fermšek^a, Urban Bren^{a,c,d}, Gregor Hostnik^{a,*}

^a Faculty of Chemistry and Chemical Engineering, University of Maribor, Smetanova 17, 2000 Maribor, Slovenia

^b Slovenian National Building and Civil Engineering Institute, Dimičeva 12, 1000 Ljubljana, Slovenia

^c Faculty of Mathematics, Natural Sciences and Information Technologies, Glagoljaška 8, University of Primorska, 6000 Koper, Slovenia

^d Institute of Environmental Protection And Sensors, Beloruska 7, 2000 Maribor, Slovenia

The focus of the study was the antibacterial effect of ellagitannins, which are polyphenolic compounds known for their wide range of biological activities [1]. We specifically examined the Minimum Inhibitory Concentration (MIC) values of chestnut extract components, including a) gallic acid, b) vescalal and castalal as well as c) vescalagin and castalagin, against the Gram-positive bacterium *Staphylococcus aureus* and the Gram-negative bacterium *Escherichia coli* using the broth microdilution method [2]. Our findings reveal that all tested samples displayed significant antibacterial activity. It was also observed that the concentration of the growth medium impacts MIC values by affecting the availability of nutrients to the bacteria. In the case of *E. coli*, MIC values increased proportionally to the medium concentration, whereas for tannin concentrations below the MIC, the duration of the lag phase extended exponentially with rising tannin concentration [3]. This effect was less pronounced for *Staphylococcus aureus* [4]. Additionally, when essential metal ions were added to the Mueller-Hinton Broth (MHB), an elevation in MIC values was observed for all tannins, suggesting that tannins' chelation of metal ions inhibits *E. coli* growth. Inductively Coupled Plasma Mass Spectrometry (ICP-MS) was employed to track metal ion concentrations in MHB, revealing that *E. coli* accumulates specific essential metal ions while others are unaffected. Notably, gallic acid was found to reduce the accumulation of these ions significantly.




Fig. 1. Structural formula of a) gallic acid, b) vescalal ($R_1=H$, $R_2=OH$) and castalal ($R_1=OH$, $R_2=H$) and c) vescalagin ($R_1=H$, $R_2=OH$) and castalagin ($R_1=OH$, $R_2=H$).

Keywords: metal ions, tannins, *Escherichia coli*, antimicrobial activity, ICP-MS, MIC

References

- [1] Taguri T, Tanaka T, Kouno I. Antimicrobial Activity of 10 Different Plant Polyphenols against Bacteria Causing Food-Borne Disease, *Biol. Pharm. Bull.* 2004; 27: 1965–1969
- [2] Štumpf S, Hostnik G, Primožič M, Leitgeb M, Salminen J.-P, Bren U. The Effect of Growth Medium Strength on Minimum Inhibitory Concentrations of Tannins and Tannin Extracts against *E. coli*, *Molecules* 2020; 25: 2947
- [3] Štumpf S, Hostnik G, Primožič M, Leitgeb M, Bren U. Generation Times of *E. Coli* Prolong with Increasing Tannin Concentration while the Lag Phase Extends Exponentially, *Plants* 2020; 9: 1680
- [4] Štumpf S, Hostnik G, Langerholc T, Pintarič M, Kolenc Z, Bren U. The Influence of Chestnut Extract and Its Components on Antibacterial Activity against *Staphylococcus aureus*, *Plants* 2023; 12: 2043

* Presenting author. E-mail address: sara.stumpf@um.si (Sara Štumpf)

S1.P270 Isolation of flavonoids from the stem bark of *Cynometra malaccensis*

Noorazlina Adnan^{a,b,c}, Aisyah Salihah Kamarozaman^{a,b,*}, Nurul Shahira Zakaria^c, Siti Zakirah Azahar^{b,c}, Nik Fatini Nik Azmin^c, Norizan Ahmat^a, Nurulfazlina Edayah Rasol^{b,c}

^a Centre of Foundation Studies, Universiti Teknologi MARA, Cawangan Selangor, Kampus Dengkil, 43800 Dengkil, Selangor, Malaysia

^b Atta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA, Cawangan Selangor, Kampus Puncak Alam, 42300 Bandar Puncak Alam, Selangor, Malaysia

^c Faculty of Applied Sciences, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia

Cynometra malaccensis, belongs to the Fabaceae family (Leguminosae). Even though there are 122 species of *cynometra* (GBIF, 2024), less than 10% of the species has been studied (Sabiha, 2022). *Cynometra* species were reported to contain flavonoids and phenolic compounds which are secondary metabolites known for their potent antioxidant and antibacterial activities (Ado, 2019 & Yuan, 2021). In this study, *C. malaccensis* was selected to be phytochemically investigated. The powder from the stem bark of *C. malaccensis* (750 g) was macerated in acetone at room temperature for 24 hours and repeated three times. The initial crude acetone extract (15 g) was subjected to treatment with diethyl ether in order to reduce tannin, resulting in a tannin-reduced extract (7.9 g). The process of fractionation was carried out using vacuum liquid chromatography (VLC) with the eluent *n*-hexane:ethyl acetate, which was gradually increased in polarity. This resulted in the separation of the mixture into eight semi-purified fractions (CMS 1-8). Fraction CMS 5 (425 mg) underwent purification using column chromatography (CC) with the solvent system of chloroform:ethyl acetate. This process resulted in the isolation of a pure compound **1** (0.9 mg). Compound **2** (28.4 mg) was obtained from CMS 6 (195 mg) using CC (chloroform:ethyl acetate). Isolation of compound **3** (0.9 mg) from CMS 7 (438 mg) was conducted via CC (chloroform:ethyl acetate) and further purified using preparative thin layer chromatography (pTLC). Based on spectroscopic techniques and comparison with existing literature data, the isolates were identified as apigenin (**1**), luteolin (**2**), and 7,4'-dihydroxyflavone (**3**).

Keywords: *Cynometra malaccensis*, apigenin, luteolin, 7,4'-dihydroxyflavone

References

- Ado, M.A., Mediani, A., Maulidiani, Ismail, I. S., Ghazali, H. M. and Abas, F., 2019. Flavonoids from *Cynometra cauliflora* and their antioxidant, α -glucosidase, and cholinesterase inhibitory activities. *Chemistry of Natural Compound*. 55(1). 112-114.
- Global Biodiversity Information Facility (GBIF) (<https://www.gbif.org/>; accessed on 8 February 2024).
- Sabiha, S., Serrano, R., Hasan, K., Moreira da Silva, I.B., Rocha, J., Islam, N., Silva, O., 2022. The Genus *Cynometra*: A Review of Ethnomedicine, Chemical, and Biological Data. *Plants*. 11. 3504.
- Yuan, G., Guan, Y., Yi, H. *et al.*, 2021. Antibacterial activity and mechanism of plant flavonoids to gram-positive bacteria predicted from their lipophilicities. *Sci Rep*. 11, 10471.

Acknowledgments

We would like to express our greatest appreciation to the Ministry of Higher Education, Malaysia through the Fundamental Research Grant Scheme (FRGS/1/2022/STG04/UITM/02/24).

* Presenting author. E-mail address: nooraz2753@uitm.edu.my (Noorazlina Adnan)

S1.P271 Chemical constituents from the leaves of *Dipterocarpus costulatus* (Dipterocarpaceae)

Siti Zakirah Azahar^a, Aisyah Salihah Kamarozaman^{b,c,*}, Noorazlina Adnan^{a,b,c}, Nik Fatini Nik Azmin^a, Nurunajah Ab. Ghani^{a,c}

^a Faculty of Applied Sciences, Universiti Teknologi MARA, 40450 Shah Alam, Selangor, Malaysia ^b Centre of Foundation Studies, Universiti Teknologi MARA, Cawangan Selangor, Kampus Dengkil, 43800 Dengkil, Selangor, Malaysia

^c Atta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA, Cawangan Selangor, Kampus Puncak Alam, 42300 Bandar Puncak Alam, Selangor, Malaysia

Dipterocarpus is the third largest genus in the group of Dipterocarpaceae family and is particularly prominent in the tropical rainforests of Southeast Asia (Appanah & Turnbull, 1998). This genus is reported to possess a broad spectrum of potential biological activities due to the presence of their resveratrol oligomers, flavonoids and triterpenoids (Xue et al., 2014) showing anti-bacterial, anti-fungal, anti-cancer, anti-diabetic and hepatoprotective effects (Fernandes & Maharani, 2019). In this study, *D. costulatus* was selected to be phytochemically studied. The granules from the leaves of *D. costulatus* (1040 g) were macerated in methanol to give crude methanol extract (765 g). Six fractions (DC 1-6) were obtained from the fractionation of the crude via vacuum liquid chromatography. Purification of fraction DC 4 (1 g) by column chromatography (CC) with the solvent system chloroform:methanol yielded two pure compounds **1** and **2**. Meanwhile, compound **3** was obtained from the purification of DC 3 (500 mg) with the same chromatographic method with the eluent chloroform:ethyl acetate. The structural elucidation of the pure compounds was accomplished based on the spectroscopic analyses and comparison with the literature data. The pure compounds were determined as suffruticosol A (**1**) (7 mg), vaticanol B (**2**) (14 mg) and scopoletin (**3**) (4 mg) (**Fig. 1**). We would like to express our greatest appreciation to the Ministry of Higher Education, Malaysia through Fundamental Research Grant Scheme (FRGS/1/2022/STG04/UITM/02/24).




Fig. 1. Structures of compounds **1** – **3**

Keywords: *Dipterocarpus costulatus*, resveratrol oligomers, vaticanol B, suffruticosol A, scopoletin

References

- Appanah, S., & Turnbull, J. M. (Eds.). (1998). A review of dipterocarps: taxonomy, ecology, and silviculture. Fernandes, A., & Maharani, R. (2019). Phytochemical and GC-MS analysis of oleoresin of *Dipterocarpus gracilis* Blume: As a basic consideration for human remedy. Intl J Pharm Sci Res, 10(5), 2224-2229.
- Xue, Y. Q., Di, J. M., Luo, Y., Cheng, K. J., Wei, X., & Shi, Z. (2014). Resveratrol oligomers for the prevention and treatment of cancers. Oxidative Medicine and Cellular Longevity, 2014, 1-9.

* Presenting author. E-mail address: aisyah2910@uitm.edu.my (Aisyah Salihah Kamarozaman)

S1.P272 16-Angeloyloxy tiglane diterpenoid esters from *Euphorbia cooperi* with anti-HIV activity

Sbonelo S. Hlengwa^{a,*}, Fanie R. van Heerden^a, Nompumelelo P. Mkkwanazi^b

^a School of Chemistry and Physics, University of KwaZulu-Natal, Pietermaritzburg, South Africa

^b HIV Pathogenesis Programme, Doris Duke Medical Research Institute, School of Laboratory Medicine and Medical Sciences, College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa

Tiglane diterpenoids constitute a significant class of anti-HIV diterpenoids in the *Euphorbia* genus (Wang et al., 2015, Otsuki et al., 2023). In an investigation to identify anti-HIV diterpenoids from South African *Euphorbia* species, an extract of *Euphorbia cooperi* N.E.Br. ex A.Berger, with 100% viral inhibition and IC₅₀ = 0.14 mg/mL, was investigated by a luciferase-based antiviral assay. The extract was analysed using LC-TOF-MS-MS for the presence of diterpenoids. The purification of the extract afforded four 16-O-tiglane diterpenoids, including two new compounds (**1**) (Fig. 1) and a 7-oxo derivative of **1**. Two known 12-deoxy phorbol esters were also identified. Tiglane phorbol esters fragment well in the ESI positive mode of the MS due to the ester bonds that are prone to MS fragmentation. The MS fragmentation patterns of the minor peaks in the LC chromatogram enabled us to propose structures for these compounds.

Other isolated compounds included a new norsesquiterpenoid and its known glycoside, arachiside A. Three phorbol esters demonstrated anti-HIV activity, with compounds **1** and one of the known phorbol esters exhibiting potent viral inhibition concentrations when compared with AZT (zidovudine). The enhanced activities of the two active diterpenoids may be attributed to structural features on the seven-membered ring of the tiglane diterpenoids.




Fig. 1. Structure of 12-deoxyphorbol ester **1**.

Keywords: *Euphorbia cooperi*, 16-O-tiglane diterpenoids, anti-HIV, LC-TOF-MS-MS

References

- Wang, H-B., Wang, X-Y., Liu, L-P., Qin G-W. and Kang, T-G., 2015. Tiglane diterpenoids from the Euphorbiaceae and Thymelaeaceae families. Chem. Rev. 115, 2975-3011.
- Otsuki, K. and Li, W., 2023. Tiglane and daphnane diterpenoids from Thymelaeaceae family: Chemistry, biological activity, and potential in drug discovery. J. Nat. Med. 77, 1-19.

* Presenting author. E-mail address: sbonelosanehlengwa@gmail.com (Sbonelo S. Hlengwa)

S1.P273 Anti-inflammatory and anti-oxidative effects of STW 42 and root extract of *Althaea officinalis* L. on endothelial cells, fibroblasts and macrophages *in vitro*

Laura Much^a, Heba Aziz-Kalbhenn^b, Marlon Palm^{b,*}, Olaf Kelber^b, Gabriel A. Bonaterra^a, Ralf Kinscherf^a

^a Philipps University of Marburg, Dept. of Medical Cell Biology, Marburg, Germany

^b Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt, Germany

Introduction: The root extract of marshmallow (*Althaea officinalis* L., REAo), known to cure mild irritation of pharyngeal mucosa as well as stomach/gut mucosa¹, has a physical action and also induces regeneration of irritated mucosa². Damage is associated with mucosal infiltration of activated leukocytes, such as MΦ, which produce excessive ROS and pro-inflammatory cytokines.

Objectives: We aimed to investigate whether REAo has anti-inflammatory/-oxidative properties in human dermal fibroblasts (NHDF), human umbilical vein endothelial cells (HUVEC), and human acute monocytic leukemia (THP-1) differentiated MΦ, critical cellular components of the gastrointestinal and oral mucosa.

Materials and methods: The cells were pre-treated with REAo or its formulation STW 42, containing REAo, and selected excipients, followed by incubation with H₂O₂ (pro-oxidative) or lipopolysaccharide (LPS, pro-inflammatory). Intracellular ROS were quantified with DCFDA. The releases of IL-6 and TNF-α were determined by ELISA. THP-1 cells were differentiated into MΦ with 0.1 µg/ml phorbol-12-myristate-13-acetate for 3–5 days.

Results: Pre-treatment (24 h) of HUVEC, NHDF or THP-1-MΦ with STW 42 or REAo (100-1000 µg/ml) significantly inhibited the H₂O₂ -induced intracellular ROS production by 30.0% to 58.7% and the LPS-activated IL6 release by 25.0-67.0%. The observed effects of STW 42 or REAo were similar to 25-100 µM diclofenac, which was used as an anti-inflammatory control.

Conclusion: These properties may support the benefit of STW 42 and its constituent REAo in patients with mucosal irritation with subsequent irritative cough or gastric complaints.

Keywords: *Althaea officinalis* L, STW 42, anti-inflammatory, anti-oxidative, mucosa

References

1. Fink C et al. Complement Med Res 2018;25:299-305; 2. Bonaterra GA et al. Front Pharmacol. 2022;13:948248.

Acknowledgements

The product was supported by the Bayer LSC project on GlycoScience.

* Presenting author. E-mail address: marlon.palm@bayer.com (Marlon Palm)

S1.P274 Ancient remedies for modern challenges: novel anti-infectives from Egyptian mummy embalming

Raphael Reher^{a,*}, Amira Naimi^a, Tim Berger^a, Kornelia Hardes^b, Tanja Pommerening^c

^a Philipps-University Marburg, Institute for Pharmaceutical Biology and Biotechnology, Department of Pharmacy, Robert-Koch-Straße 4, 35037 Marburg, Germany

^b Philipps-University Marburg, Institute for Pharmaceutical Biology and Biotechnology, Department of Pharmacy, Robert-Koch-Straße 4, 35037 Marburg, Germany

^c Fraunhofer Institute for Molecular Biology and Applied Ecology, Department of Bioresources, Ohlebergsweg 12, 35392 Giessen, Germany

The escalating global threat of antibiotic resistance and the resurgence of neglected tropical diseases (NTDs) underscore the urgent need for new anti-infective therapies.³ Of all anti-infective drugs approved from 1941-2014, 73% are natural products (NPs) or derivatives, underscoring the importance of NPs in drug discovery and development for this therapeutic area.^{1,2} Leveraging the rich pharmacopoeia of Traditional Egyptian Medicine from the Pharaonic Era (ca. 3000-323 B.C.), we embarked on a collaboration to unearth potential anti-infective agents from ancient remedies.

Partnering with Egyptologists, we decoded and analyzed ancient medical recipes, focusing on identifying compounds with anti-infective properties. Concurrently, we formed alliances with virologists to employ native metabolomics, a powerful technique integrating non-targeted liquid chromatography tandem mass spectrometry and native mass spectrometry, to discover inhibitors of viral proteins and other anti-infective targets.⁴

This interdisciplinary approach led us to isolate bioactive natural products from embalming residues of Egyptian mummies, confirmed through ¹⁴C analysis to originate from the 28th Pharaonic dynasty (407-386 B.C.). Utilizing AI-assisted NMR and LC-MS/MS methods, we elucidated their structures and identified promising compounds with selective antiviral properties against pandemic Influenza virus Type A (H1N1pdm)⁵⁻⁷.

Our findings represent a pioneering effort to bridge ancient wisdom with modern science in the quest for novel anti-infective treatments, offering hope in the face of emerging infectious threats.

Keywords: natural products discovery, Egyptian mummy, artificial intelligence, antiviral

References

- (1) *Neglected tropical diseases -- GLOBAL*. <https://www.who.int/health-topics/neglected-tropical-diseases> (accessed 2024-03-08).
- (2) Newman, D. J.; Cragg, G. M. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J. Nat. Prod.* 2020, 83 (3), 770–803.
- (3) Gerwick, W. H.; Moore, B. S. Lessons from the Past and Charting the Future of Marine Natural Products Drug Discovery and Chemical Biology. *Chemistry & Biology* 2012, 19 (1), 85–98.
- (4) Reher, R. et al. Native Metabolomics Identifies the Rivulariapeptolide Family of Protease Inhibitors. *Nat Commun* 2022, 13 (1), 4619.
- (5) Mullowney, M. W. et al. Artificial Intelligence for Natural Product Drug Discovery. *Nat Rev Drug Discov* 2023, 22 (11), 895–916.
- (6) Reher, R. et al. A Convolutional Neural Network-Based Approach for the Rapid Annotation of Molecularly Diverse Natural Products. *J. Am. Chem. Soc.* 2020, 142 (9), 4114–4120.
- (7) Dührkop, K. et al. Systematic Classification of Unknown Metabolites Using High-Resolution Fragmentation Mass Spectra. *Nat Biotechnol* 2021, 39 (4), 462–471.

* Presenting author. E-mail address: raphael.reher@pharmazie.uni-marburg.de (Prof. Dr. Raphael Reher)

S1.P275 Identification of natural compounds targeting the *Plasmodium falciparum* pyruvate kinases

Rym Attia^{a,*}, Lucille, Hervé^b, Edilson Beserra De Alencar Filho^c, Raphael Grougnet^a, Raimundo Gonçalves de Oliveira Junior^a, Boutefnouchet Sabrina^a, Anaïs Merckx^b

^a Team P.N.A.S, UMR-CNRS 8038 CiTCoM, Faculté de Pharmacie de Paris, Université de Paris Cité, France

^b UMR261 MERIT, IRD, Faculté de Pharmacie de Paris, Université de Paris Cité, France

^c Department of Pharmacy, Universidade Federal do Vale do São Francisco, Petrolina-PE, Brazil

Pyruvate kinases (PfPyrK1 and PfPyrK2) are essential for the survival of *Plasmodium falciparum* during its erythrocytic stage. These enzymes catalyze the phosphorylation of ADP to ATP, using phosphoenolpyruvate (PEP) that is converted to pyruvate. Therefore, they represent promising therapeutic targets (Chan, Tan, and Sim 2007; Swift et al. 2020). However, to date, no effective specific inhibitors for PfPyrK1 or PfPyrK2 have been identified.

In this context, 70 molecules were screened *in silico* by evaluating their binding energies to both the oxalate (a substrate analog inhibitor) and the ATP binding sites of PfPyrK1 and PfPyrK2. Among the compounds tested, including flavones, sesquiterpene lactones, and diterpenes obtained from plant extracts or hemisynthesis, a group of polymethoxyflavones (PMFs) showed the highest affinity for these sites. The binding energies ranged from -8.516 to -6.943 kcal/mol for the oxalate site and from -9.129 to -7.624 kcal/mol for the ATP site. Following the *in silico* screening, flavones available in sufficient quantities in our chemical library were tested *in vitro* on PfPyrK1 and PfPyrK2. Apigenin, with binding energies of -7.292 and -7.646 kcal/mol for the oxalate and ATP sites respectively, exhibited interesting inhibitory activity of 54% and 45% on PfPyrK1 and PfPyrK2, respectively, at 50 μ M (The IC₅₀ of suramin, a reference inhibitor, were 7 μ M and 3 μ M for PfPyrK1 and PfPyrK2, respectively).

Other flavones, such as viscosine (with binding energies of -7.403 and -8.339 kcal/mol for the oxalate and ATP sites respectively), are being isolated from *Gardenia oudiepe*, to evaluate their *in vitro* inhibition effects on PfPyrK.

Keywords: Malaria, *in silico* screening, pyruvate kinases, polymethoxyflavones

References

- Chan, Maurice, Doreen S. H. Tan, et T. S. Sim. 2007. *Plasmodium falciparum* pyruvate kinase as a novel target for antimalarial drug-screening. *Travel Medicine and Infectious Disease*, 1st International Conference of Travel Medicine and Infectious Disease. 5, 125-31.
- Swift, Russell P., Krithika Rajaram, Cyrianne Keutcha, Hans B. Liu, Bobby Kwan, Amanda Dziedzic, Anne E. Jedlicka, et Sean T. Prigge. 2020. The NTP Generating Activity of Pyruvate Kinase II Is Critical for Apicoplast Maintenance in *Plasmodium falciparum*. *eLife*. 9, e50807.

* Presenting author. E-mail address: rym.attia@u-paris.fr (Rym Attia)

S1.P276 The relationship between bioactivity and taxonomy of *Euphorbia* species in South Africa

Fanie R. van Heerden*, Sbonelo S. Hlengwa, Wonder Mpapane

School of Chemistry and Physics, University of KwaZulu-Natal, Private Bag X01, Scottsville, South Africa

The phytochemistry of Euphorbiaceae and Thymelaceae has recently received substantial attention due to the anticancer and anti-HIV activities of phorbol diterpenes in these families. Some of these diterpenes, such as prostratin, not only kill HIV at nanomolar concentrations but can reactivate the latent virus and have the potential to cure patients with HIV (Beans et al., 2013). Medication containing Euphorbiaceae diterpenes as the active ingredient has also reached the market for treating topical cancers in animals (Cullen et al., 2021).

Euphorbiaceae are one of the largest flowering plant families (6300 species), and as a result the selection of suitable species for phytochemical investigation may be problematic. In this project, we have focused on the identification of novel bioactive diterpenes from the genus *Euphorbia*. Southern Africa is home to 172 *Euphorbia* species, of which 74% are endemic to this area (Bruyns, 2023). To assist with selecting *Euphorbia* species for investigation, we have correlated the diterpene contents with the taxonomy of the different species. The diterpenes were identified by isolation of the compounds and/or identification by LC-MS-MS. The structures and bioactivity of the diterpenes will be discussed. The terpenes identified were phorbol esters, abietane, kaurane, and atisane diterpenoids, and these compounds were correlated with the taxonomic classification of the plants.

Keywords: Euphorbiaceae, Thymelaceae, *Euphorbia*, diterpenoid, phorbol ester

References

- Beans, E.J., Fournogerakis, D., Gauntlett, C., Heumann, L.V., Kramer, R., Marsden, M.D., Murray, D., Chun, T.-W., Zack, J.A., Wender, P.A., 2013. Highly potent, synthetically accessible prostratin analogs induce latent HIV expression in vitro and ex vivo. PNAS 110, 11698–11703.
- Cullen, J.K., Boyle, G.M., Yap, P.-Y., Elmlinger, S., Simmons, J.L., Broit, N. Johns, J., Ferguson, B., Maslovskaya, L.A., Andrei I. Savchenko, A.I., Mirzayans, P.M., Porzelle, A., Bernhardt, P.V., Gordon, V.A., Reddell, P.W. Pagani, A. Appendino, G., Parsons, P.G., Williams, C.M., 2021. Activation of PKC supports the anticancer activity of tigilanol tiglate and related epoxytiglanes. Sci. Rep. 11, 207.
- Bruyns, P.V., 2023. Euphorbias in the spotlight. Veld & Flora, Botanical Society of South Africa.

* Presenting author. E-mail address: vanheerdenf@ukzn.ac.za (Fanie van Heerden)

S1.P277 Structure activity studies of anti-parasitic norditerpenes from subterranean fungi

Christine Salomon^{a,*}, Roberta O'Connor^b, Yudi Rusman^a, Jessica Williams^a, Harrison VanKoten^a, Fernanda Fumuso^b, Alexis Cotto-Rosario^b

^a Center for Drug Design, University of Minnesota, Minnesota, USA 55455

^b Department of Veterinary and Biomedical Sciences, University of Minnesota, Minnesota, USA 55108

Cryptosporidiosis is a globally distributed water-borne gastrointestinal disease caused by protozoan parasites in the genus *Cryptosporidium*. *C. parvum* is one of the most important species that infects both humans and ruminant animals. The diarrheal disease is especially problematic for immunocompromised patients, and there are no vaccines or effective clinical drugs for these populations and new treatments are urgently needed.

We previously identified a suite of norditerpene lactone metabolites with anti-Cryptosporidial activity from the fungus *Oidiodendron truncatum*, isolated from the Soudan Iron Mine in northern Minnesota.¹ The two most active compounds with sub-micromolar EC₅₀s were oidiolactones A (1) and B (2), which differ in both anti-parasitic potency and cytotoxicity towards host cells. In order to expand structure activity relationship studies to better understand the structural features that might lead to higher activity and minimal off target effects, we identified a *Mucor* sp. of fungus from the same subterranean environment that also produced oidiolactones A and B as well as four additional new congeners, including a norditerpene lactam derivative. We also transformed the natural oidiolactone A into eight semi-synthetic analogs and tested these together with the natural compounds against *C. parvum* infected HCT- 8 host cells. None of the new analogs or semi- synthetic derivatives were more potent than 1 or 2, but the C-13 epimer of oidiolactone A displayed comparable activity and increased cytotoxicity. The structure activity relationships of the 21 norditerpenes will be discussed, as well as the implications for mechanism of action studies.




Fig. 1. Structures of compounds 1 and 2.

Keywords: *Cryptosporidium*, *Toxoplasma*, fungi, anti-parasitic, infectious disease

References

Rusman, Y, Wilson, MB; Williams, J, Held, BW, Blanchette, RA, Anderson, B, Lupfer, C, Salomon, CE. 2020. Antifungal norditerpene oidiolactones from the fungus *Oidiodendron truncatum*, a potential bio-control agent for white-nose syndrome in bats. *Journal of Natural Products*. 83, 344-353.

* Presenting author. E-mail address: csalomon@umn.edu (Christine Salomon)

S1.P278 Bioactive terpenoids from members of the Hericiaceae

Winnie Chemutai Sum^{a,b,*}, Sherif Elsayed Ebada^{a,c}, Nico Mitschke^{a,d}, Hedda Schrey^a, Harald Kellner^e, Josphat Matasyoh^f, Marc Stadler^{a,b}

^a Department Microbial Drugs, Helmholtz Centre for Infection Research GmbH (HZI), Inhoffenstraße 7, 38124 Braunschweig, Germany

^b Institute of Microbiology, Technische Universität Braunschweig, Spielmannstraße 7, 38106 Braunschweig, Germany

^c Department of Pharmacognosy, Faculty of Pharmacy, Ain Shams University, Abbasia, 11566 Cairo, Egypt

^d Research Group for Marine Geochemistry (ICBM-MPI Bridging Group), Institute for Chemistry and Biology of the Marine Environment (ICBM), Carl von Ossietzky Universität Oldenburg, Carl- von-Ossietzky-Str. 9-11, 26129 Oldenburg, Germany

^e Department of Bio- and Environmental Sciences, Technische Universität Dresden-International Institute Zittau, Markt 23, 02763 Zittau, Germany

^f Department of Chemistry, Egerton University, P.O. Box 536, Njoro 20115, Kenya

In our continued search for biologically active metabolites from cultures of rare Basidiomycota species, we identified twenty-two previously undescribed compounds from submerged and solid-state cultures of three European Basidiomycetes belonging to family *Hericiaceae* namely: *Dentipellis fragilis*, *Hericium flagellum* and *H. coralloides* (1–4). Their chemical structures were unambiguously determined based on extensive 1D and 2D NMR spectroscopy along with HR-ESI-MS analyses. The isolated compounds revealed different biological effects. Some exhibited significant activities in our standardized cell-based assays for the determination of cytotoxic and antimicrobial effects. In addition, most of the compounds demonstrated neurotrophic effects. These were determined using the rat pheochromocytoma cell (PC-12)-based assay for neurite outgrowth promotion and real-time quantitative reverse transcription polymerase chain reaction (RT-qPCR) assay, to assess the effects of the compounds on the expression of neurotrophins NGF and brain-derived neurotrophic factor (BDNF) on human astrocytoma cells.

Keywords: *Basidiomycetes*, NMR, HR-ESI-MS, cytotoxic, antimicrobial, neurotrophic

References

1. Sum WC, Gonkhom D, Ibrahim MAA, Stadler M, Ebada SS. 2024. New isoindolinone derivatives isolated from the fruiting bodies of the basidiomycete *Hericium coralloides*. *Mycol Prog* 23:1–7.
2. Mitschke N, Chemutai Sum W, Hassan K, Kirchenwitz M, Schrey H, Gerhards L, Kellner H, Stradal TEB, Matasyoh JC, Stadler M. 2023. Biologically active drimane derivatives isolated from submerged cultures of the wood- inhabiting basidiomycete *Dentipellis fragilis*. *RSC Adv* 13:25752–25761.
3. Sum WC, Ebada SS, Kirchenwitz M, Kellner H, Ibrahim MAA, Stradal TEB, Matasyoh JC, Stadler M. 2023. Hericioic Acids A-G and Hericiofuranoic Acid; Neurotrophic Agents from Cultures of the European Mushroom *Hericium flagellum*. *J Agric Food Chem* 71:11094–11103.
4. Sum WC, Mitschke N, Schrey H, Wittstein K, Kellner H, Stadler M, Matasyoh JC. 2022. Antimicrobial and Cytotoxic Cyathane-Xylosides from Cultures of the Basidiomycete *Dentipellis fragilis*. *Antibiotics* 11:1072.

* Presenting author. E-mail address: winnie.sumchemutai@helmholtz-hzi.de (Winnie Sum)

S1.P279 Isolation, structural characterization and antiprotozoal activity of steroidal alkaloids from *Pachysandra terminalis*

Lizanne Schäfer^{a,*}, Marcel Kaiser^b, Pascal Mäser^b, Thomas J. Schmidt^a

^a Institute of Pharmaceutical Biology and Phytochemistry (IPBP), University of Münster, PharmaCampus, Corrensstraße 48, D-48149 Münster, Germany

^b Swiss Tropical and Public Health Institute, Kreuzstraße 2, CH-4123 Allschwil, Switzerland

Trypanosoma brucei rhodesiense (*Tbr*) and *Plasmodium falciparum* (*Pf*) are protozoan parasites responsible for severe diseases, namely human African trypanosomiasis and malaria. With limited treatment options and emerging resistance, there is an urgent demand for new antiprotozoal compounds. Previous studies of our group have indicated that plants of the Buxaceae family contain aminosteroids with high activity against these parasites (Althaus et al., 2014), (Szabó et al., 2021). The alkaloid enriched fraction of *Pachysandra terminalis* Sieb. & Zucc. (Buxaceae) also showed prominent activity against *Tbr* (Flittner et al., 2021). Therefore, it is promising to further investigate the constituents of this plant and to systematically isolate and test individual compounds. An optimized extraction protocol was established to obtain an alkaloid-enriched fraction of *P. terminalis* leaves that was active against *Tbr* (IC₅₀ 1.84 µg/mL) and highly active against *Pf* (IC₅₀ 0.31 µg/mL). With centrifugal partition chromatography the alkaloid-enriched fraction was further divided into 18 subfractions with IC₅₀ values against *Tbr* ranging from 0.50 - 7.92 µg/mL and against *Pf* from 0.17 - 1.27 µg/mL. By preparative HPLC, eight aminosteroids could be isolated so far: 5,6-Dehydro-desacyl-epipachysamine A (**1**), desacyl-epipachysamine A (**2**), epipachysamine B (**3**), pactermine A (**4**), 3*R*-pachysanaximine A (**5**), pachysamine A (**6**), *N*-methyl-desacyl-epipachysamine A (**7**) and sarcodinine (**8**). The structures were established by UHPLC/+ESI-QqTOF-MS/MS and NMR spectroscopy. To the best of our knowledge, compounds **1** and **5** are described as natural products for the first time. The activity against *Tbr* and *Pf* is shown along with the structures in Fig. 1.




Fig. 1. Structures of the steroidal alkaloids (**1-8**) from the leaves of *Pachysandra terminalis*

Keywords: *Pachysandra terminalis*, Buxaceae, steroidal alkaloids, *Trypanosoma brucei rhodesiense*, *Plasmodium falciparum*

References

- Althaus, J.B., et al., 2014. Antiprotozoal Activity of *Buxus sempervirens* and Activity-Guided Isolation of O- tigloylcyclovirobuxeine-B as the Main Constituent Active against *Plasmodium falciparum*. *Molecules* 19, 6184-6201.
- Flittner, D., et al., 2021. The Alkaloid-Enriched Fraction of *Pachysandra terminalis* (Buxaceae) Shows Prominent Activity against *Trypanosoma brucei rhodesiense*. *Molecules* 26, 591.
- Szabó, L.U., et al., 2021. Antiprotozoal Nor-Triterpene Alkaloids from *Buxus sempervirens* L. *Antibiotics* 10, 696.

* Presenting author. E-mail address: l_scha57@uni-muenster.de (Lizanne Schäfer)

S1.P280 Antibacterial Activity of different extracts of *Plicosepalus acacia*

Omaish Alqahtani*

Department of Pharmacognosy, College of Pharmacy, Najran University, Najran, Saudi Arabia

A considerable global health crisis that threatens communities has been attributed to the emergence of resistant bacterial infections. As a result, the search for novel antibiotics from different natural sources has increased. *Plicosepalus acaciae* (Zuccarini) Wiens & Polhill) is used in Saudi Arabia for treating some infectious disease such as bronchitis. The aim of this study is to evaluate the antibacterial activity of different extracts of *Plicosepalus acacia*. MIC assay was used to assess the antibacterial activity for hexane, chloroform and methanol extracts of leaves. The findings of MIC assay illustrated that all extracts showed weak or no significant activities except for the methanol extract that possessed mild to moderate activities at 256 µg/mL in compared to a positive control: In conclusion, the methanol extract of *Plicosepalus acacia* demonstrated moderate activity against *Staphylococcus aureus* and *Bacillus subtilis*, weak activity against *Escherichia coli*, and no activity against *Pseudomonas aeruginosa* and *Klebsiella pneumonia*.

Keywords: *Plicosepalus acacia*., antibacterial agents, bacterial resistance

* Presenting author. E-mail address: osalqahtani@nu.edu.sa (Omaish Alqahtani)

S1.P281 *Epimedium koreanum* attenuates Influenza A viral infection via modulation of hemagglutinin and neuraminidase

Won-Kyung Cho, Jin Yeul Ma*

Korean Medicine Application Center, Korea Institute of Oriental Medicine, Daegu, South Korea, 41062

Epimedium koreanum also known as Nakai is a popular plant in Korean and Chinese medicine for treating various ailments. In this study, the water extract of *E. Koreanum* has a significant inhibitory impact on influenza A virus (IAV) infection by directly blocking viral attachment and having a virucidal effect. We examined the effect of the *E. koreanum* water extract on viral infection using fluorescent microscopy and fluorescence-activated cell sorting analysis with a green fluorescent protein-tagged Influenza A/PR/8/34 virus. At a concentration of 100 µg/mL, the Nakai water extract decreased GFP expression by up to 90% compared to the virus-infected control. Immunofluorescence and Western blot analyses against influenza viral proteins showed that it dose-dependently decreased the viral protein expression. The Nakai extract at 100 µg/mL inhibited the H1N1 influenza virus's hemagglutinin and neuraminidase, preventing viral binding to cells. Furthermore, it exhibited a virucidal impact and inhibited the cytopathic effects of H1N1, H3N2, and influenza B virus infection. Finally, our results showed that *E. koreanum* water extract could be developed as a natural viral inhibitor against influenza virus infection.

Keywords: *Epimedium koreanum*, antiviral effect, influenza virus, virucidal effect

* Presenting author. E-mail address: jyma@kiom.re.kr (Jin Yeul Ma)

S1.P282 Anti-biofilm and anti-quorum sensing activity of galloylquinic acids from *Copaifera lucens* leaves against clinical isolates of multidrug-resistant *Pseudomonas aeruginosa* in open wound infection: *in vitro* and *in vivo* study

Lamiaa Al-Maddboly^a, Mohamed Abd El-Salam^{b,c,*}, Jairo Bastos^d, Eiman Hasby^e, Ivan Kushkevych^f, Rasha El-Morsi^g

^a Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Tanta University, Tanta, Egypt

^b Department of Pharmacognosy, Faculty of Pharmacy, Delta University for Science and Technology, International Coastal Road, Gamasa, 11152, Egypt

^c School of Pharmacy and Biomolecular Sciences, Royal College of Surgeons in Ireland, Dublin D02 VN51, Ireland

^d Department of Pharmaceutical Sciences, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, SP, 14040-903, Brazil

^e Department of Pathology, Faculty of Medicine, Tanta University, Tanta, Egypt

^f Department of Experimental Biology, Faculty of Science, Masaryk University, 60200 Brno, Czech Republic ^gDepartment of Microbiology, Faculty of Pharmacy, Delta University for Science and Technology, International Coastal Road, Gamasa, 11152, Egypt

This presentation investigates the antibacterial, anti-biofilm, and anti-quorum sensing actions of galloylquinic acid compounds (GQAs) derived from the brazilian *Copaifera lucens* leaves against clinical isolates of multidrug-resistant (MDR) *Pseudomonas aeruginosa*. Our results showed significant inhibition zone diameters (25-40 mm) in the antibacterial activity assay, with minimal inhibitory concentrations (MIC) ranging from 1-4 µg/mL and minimal bactericidal concentrations (MBC) between 2-16 µg/mL. GQAs effectively interfered with planktonic *P. aeruginosa* isolates and inhibited their growth within pre-formed biofilms, with MBIC80 and MBEC80 values of 64 µg/mL and 128 µg/mL, respectively. Fluorescence staining and confocal microscopy revealed a substantial reduction in cell viability and biofilm thickness (62.5%) following exposure to 128 µg/mL of GQAs. Scanning electron micrographs further confirmed GQAs ability to disrupt biofilm and bacterial structures by interfering with biomass and exopolysaccharides (Fig. 1). Moreover, GQAs significantly ($p < 0.05$) reduced the production of virulence factors and bacterial motility, including rhamnolipid, pyocyanin, as well as swarming and swimming motility. This reduction is attributed to GQAs downregulation (up to 89%) of quorum-sensing genes (*lasI*, *lasR*, *pqsA*, and *pqsR*) involved in biofilm formation. In conclusion, GQAs derived from *C. lucens* represent promising antibiofilm agents with the potential to enhance wound healing in biofilm-associated infections caused by MDR *P. aeruginosa* clinical isolates.




Fig. 1. Scanning electron micrographs showing biofilms of *P. aeruginosa* after exposure to different concentrations of GQAs. (A) untreated preformed biofilm showing large number of the bacterial cells with intact cell walls, arranged in close contact with each other, covered with EPS matrix glycocalyx, (B) spaces were detected in the biofilm and a few distortions in the cell morphology were observed after treatment with 16 µg/mL of GQAs, (C) irregularities in cell morphology (arrow) are markedly noticed following treatment with 64 µg/mL of GQAs in addition to a significant reduction in the biomass and EPS matrix, (D) severe damage was detected in the biofilm in case of the treatment with 128 µg/mL of GQAs, with leakage of cellular content and death (arrows).

Keywords: Antibacterial, *Pseudomonas aeruginosa* clinical isolates, anti-quorum sensing, anti-biofilm, antivirulence, galloylquinic acid compounds

References

- Abd El-Salam M, El-Tanbouly G, Bastos J, Metwally H (2023) Suppression of VEGF and inflammatory cytokines, modulation of Annexin A1 and organ functions by galloylquinic acids in breast cancer model. *Scientific Reports* 13(1):12268 doi:10.1038/s41598-023-37654-4
- Abd El-Salam MA, El-Tanbouly GS, Bastos JK, Metwally HA (2022) Novel antitumor activity of the combined treatment of galloylquinic acid compounds with doxorubicin in solid Ehrlich carcinoma model via the Notch signaling pathway modulation. *Life Sci* 299:120497 doi:10.1016/j.lfs.2022.120497
- Al-Madbolay LA, Abd El-Salam MA, Bastos JK, El-Shorbagy SH, El-Morsi RM (2022) Novel Preclinical Study of Galloylquinic Acid Compounds from *Copaifera lucens* with Potent Antifungal Activity against Vaginal Candidiasis Induced in a Murine Model via Multitarget Modes of Action. *Microbiol Spectr* 10(5):e0272421 doi:10.1128/spectrum.02724-21

* Presenting author. E-mail address: mohamed.abdelsalam@deltauniv.edu.eg (M. Abd El-Salam)

S1.P283 Effects of raw and boiled ginger (*Zingiber officinale* Roscoe) on hyperlipidemia in high cholesterol diet-induced obese rats – A comparative therapeutic study

Akomolafe Seun Funmilola*, Atoyebi David Ayodeji

Department of Biochemistry, Ekiti State University, P.M.B. 5363, Ado Ekiti, Ekiti State, Nigeria

This presentation will cover the comparative therapeutic effects of raw and boiled ginger (*Zingiber officinale* Roscoe) on hyperlipidemia in high cholesterol diet-induced obese rats. Traditional herbal medicine has claimed to have successfully employed ginger rhizome to treat obesity in people all over the world (Pan et al., 2016; Gupta et al., 2015; Abdulrazaq et al., 2011).

As part of continuing search for new natural compounds and their bioactive components as potential novel drug development and illness treatment alternatives, we compared the therapeutic benefits of raw as well as boiled ginger at different percentages (2% and 4%) on hyperlipidemia in high cholesterol diet-induced obese rats. Feeding of high cholesterol diet (HCD)-induced obese rats with raw and boiled ginger for twenty-four weeks resulted in the levels of antioxidant status, lipid profile as well as 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMG Co-A reductase) activity to be significantly elevated ($p < 0.05$), with a concomitant reduction in activities of liver marker enzymes in the serum and liver of HCD-induced obese rats fed raw and boiled ginger in comparison with untreated obese rats. Levels Histology reports also depict the potential of the ginger to reduce abdominal fat deposits in HCD-induced obese rats fed raw and boiled ginger (Fig. 1).




Fig. 1. Histological changes of the adipose tissue (abdominal fat)

Both raw and boiled ginger had positive effects on the parameters studied, but boiled ginger at a higher percentage (4%) had the greatest significant result in comparison with raw ginger. Therefore, consuming boiled ginger will be more effective than eating raw ginger in preventing hyperlipidemia brought on by a high-cholesterol diet.

Keywords: obesity, ginger rhizome, thermal processing, lipid profile, lipogenesis, HMG-CoA reductase

References

- Abdul-Razak, S., Daher, A. M., Ramli, A. S., Ariffin, F., Mazapuspavina, M. Y., Ambigga, K. S., REDISCOVER investigators. (2016). Prevalence, awareness, treatment, control and socio demographic determinants of hypertension in Malaysian adults. BMC public health, 16, 1-10.
- Gupta, V., Mah, X. J., Garcia, M. C., Antonypillai, C., & van der Poorten, D. (2015). Oily fish, coffee and walnuts: Dietary treatment for nonalcoholic fatty liver disease. World Journal of Gastroenterology: WJG, 21(37), 10621.
- Pan, M. H., Tung, Y. C., Yang, G., Li, S., & Ho, C. T. (2016). Molecular mechanisms of the anti-obesity effect of bioactive compounds in tea and coffee. Food & function, 7(11), 4481-4491

* Presenting author. E-mail address: seun.akomolafe@eksu.edu.ng (Akomolafe Seun Funmilola)

S1.P284 Triterpenoids from the roots of *Anthocleista djalensis* and their antiplasmodial, antimicrobial, and antitrypanosomal properties

Ijeoma Solomon Okoro^{a,b,*}, Anita Kelechi Asekunowo^c, Sampson Dominic Umoh^a, John Ogbaji Igoli^a, Xavier SiweNoundou^b, Rui Werner Maçedo Krause^b

^a Department of Chemistry, Joseph Sarwuan Tarka University, Makurdi, Nigeria

^b Department of Chemistry, Rhodes University, P. O. Box 94, Grahamstown 6140, South Africa

^c Department of Chemistry, Faculty of Science, University of Lagos

This study explores the phytochemical composition and antibacterial, antiplasmodial, and antitrypanosomal properties of *Anthocleista djalensis* (A. Chev.) roots used in traditional medicine for treating malaria, and leprosy, among others. Chromatographic procedures were used to isolate the compounds from the species, and their structures were determined by 1D and 2D NMR data and compared with the literature. *In vitro*, antimicrobial properties, cell growth-inhibiting characteristics, and cytotoxicity were assessed. Bauerenone (**1**), bauerenol (**2**), and a mixture of stigmasterol and β - sitosterol (**3** & **4**) were isolated from this species. Antimicrobial effectiveness was demonstrated by the ethanol, acetone, and ethyl acetate extracts and all compounds against all tested microbes. MICs ranged from 16 to 100 $\mu\text{g/mL}$ in the crude extracts and 4 to 31 μM for the isolated compounds. Significant anti-malarial activity was recorded for hexane (IC_{50} : 0.0191 $\mu\text{g/mL}$), ethyl acetate (IC_{50} : 0.0139 $\mu\text{g/mL}$), acetone (IC_{50} : 0.0168 $\mu\text{g/mL}$), ethanol (IC_{50} : 0.0159 $\mu\text{g/mL}$), and the isolated compounds (bauerenone, IC_{50} : 0.0150 μM ; bauerenol, IC_{50} : 0.0138 $\mu\text{g/mL}$; a combination of β - sitosterol and stigmasterol, IC_{50} : 0.0145 $\mu\text{g/mL}$). The crude extracts (hex, IC_{50} : 0.0576; ethyl acetate, IC_{50} : 0.0318; acetone, IC_{50} : 0.0452; and ethanol, IC_{50} : 0.0313 $\mu\text{g/mL}$) with bauerenol (IC_{50} : 0.0554 μM) demonstrated substantial antitrypanosomal effects comparable to pentamidine (IC_{50} : 0.01252 μM). No toxicity was observed toward human cervical carcinoma (HeLa) cells. The study reveals that bauerenone, stigmasterol, and β -sitosterol, along with the root extracts of *Anthocleista djalensis*, are potential candidates for integration into the natural products drug discovery programs, highlighting the need for evidence-based investigation of the species.

Keywords: *Anthocleista djalensis*, cytotoxicity, HeLa cells, antimicrobial activity, antiparasitic activity

*Presenting author. E-mail address: okoro.ijeoma@uam.edu.ng (Ijeoma S. Okoro)

S1.P285 Enzymatic Synthesis of Tryptophan-Linked DKP Dimers: Expanding Biosynthetic Routes for Antifungal Drug Discovery

Natalia Harris^{a,*}, Danny Li^b, David Sherman^c

^a Department of Biological Chemistry, University of Michigan Medical School, Ann Arbor, Michigan

^b Department of Chemistry, University of Michigan, Ann Arbor, Michigan

^c Department of Medicinal Chemistry, University of Michigan College of Pharmacy, Ann Arbor Michigan

Dimeric diketopiperazine (DKP) alkaloids belong to a diverse class of natural products with an array of pharmacological applications such as anti-cancer and anti-microbial therapies. Synthesis of dimeric DKPs are of interest due to their varied biological activities and intricate structural features. Total chemical synthesis of tryptophan linked DKP dimers remains a challenging task which has motivated us to turn to a biocatalytic route. Currently, there are no catalyst-controlled approaches for accessing a variety of DKP dimerization modes from a single starting material. A previous in silico analysis of *Streptomyces* genomes revealed several bacterial DKP dimerases, which have been identified as cytochrome P450 (P450). Biochemical analysis revealed the diverse dimerization activities of the four cytochrome P450 DKP dimerases using Trp-Pro (brevianamide F) as the monomeric DKP substrate.

In this study, we explore the substrate scope of pivotal P450s, NzeB and AspB, which catalyze C–C and C–N bond formations in DKP dimer biosynthesis. Using 24 unnatural DKP substrates, which are analogs of the native substrate brevianamide F, we have expanded enzymatic control over C–H functionalization sites. This investigation not only enhances our understanding of selective dimer formation but also lead to the biosynthesis of novel small molecules with antifungal properties, crucial in combating rising fungal infections. Conducting this substrate scope study has led to the discovery of compounds effective against drug-resistant pathogen *Candida auris*.

Keywords: Biocatalysis, cytochrome P450, diketopiperazine, antifungals

References

- Borthwick AD. 2,5-diketopiperazines: Synthesis, reactions, medicinal chemistry, and bioactive natural products. *Chem Rev.* 2012;112(7):3641-3716.
- Martins MB, Carvalho I. Diketopiperazines: biological activity and synthesis. *Tetrahedron.* 2007;63(40):9923-9932. Tadano S, Ishikawa H. Synthesis of tryptophan-based dimeric diketopiperazine alkaloids using bioinspired reactions. *Synlett.* 2014;25(2):157-162.
- Shende V V., Khatri Y, Newmister SA, et al. Structure and Function of NzeB, a Versatile C–C and C–N Bond Forming Diketopiperazine Dimerase. *Journal of the American Chemical Society.* 2020; 142, 41, 17413–17424.
- Shende VV, Harris NR, et al. Molecular Dynamics Simulations Guide Chimeragenesis and Engineered Control of Chemoselectivity in Diketopiperazine Dimerases. *Angewandte Chemie.* 2023; Volume 62, Issue 20.

* Presenting author. E-mail address: hnatalia@umich.edu (Natalia Harris)

S1.P286 An investigation of *Origanum onites* and *Thymus fallax* in terms of the European Pharmacopoeia criteria

Ayten Acar^a, Pelin Taştan^{b,*}

^a Department of Pharmacognosy, Faculty of Pharmacy, Izmir Katip Çelebi University, Çiğli, Izmir, Turkey

^b Department of Pharmacognosy, Faculty of Pharmacy, Izmir Katip Çelebi University, Çiğli, Izmir, Turkey

The Lamiaceae family includes approximately 236 genera and 6900 to 7200 species (Tzima et al., 2018). In Turkey the Lamiaceae family is represented by 46 genera and 586 species, 260 of which are endemic, and is the third richest family in Turkey in terms of plant species and diversity, with an endemism of 44.2% (Davis, 1982). Medicinal plants belonging to this family have been used for many years in the treatment of various animal and human diseases based on their antibacterial, antioxidant, antitumoral, antifungal, analgesic and insecticidal effects (Hossain et al., 2010).

This presentation covers the determination of quality control methods and analytical studies for *Thymus fallax* (catri) and *Origanum onites* (kekik) which were collected from Şirnak and Izmir, respectively. Heavy metal analyses using the WDXRF method and the determination of ash content and loss on drying according to the European Pharmacopoeia 8.0 standards were carried out (EDQM). Furthermore, the total phenolic contents of the aqueous extracts of both plants were tested with the Folin-Ciocalteu colorimetric method to getting first step for determining their antioxidant activities (Kulisic et al., 2005).

The amounts of heavy metals determined for both species were within the limits of pharmacopoeia standards, according to WDXRF results. In addition, total ash and water amounts were determined to be 7%, 14% and 14%, 16% for *Origanum onites* and *Thymus fallax* respectively, and were determined to be in compliance with pharmacopoeia standards. Total phenolic contents (GAE µg/mL) were found to be 28.152 ± 0.389 and 41.009 ± 2.059 for *Origanum onites* and *Thymus fallax*, respectively. This is evidence that both species contain phenolic compounds that will cause them to display antioxidant activity. The data obtained within the scope of the study showed that if the plants are procured correctly and the water extracts are prepared with appropriate methods, the quality and effectiveness will increase.

Keywords: *Origanum onites*, *Thymus fallax*, Lamiaceae, phenolic content, WDXRF

References

- Council of Europe (EDQM). 2013. Oregano, Origani herba; Thyme, Thymi herba. European Pharmacopoeia 8.0. 1342- 1343, 1403-1405.
- Davis, P.H., (Edt.) 1982. Flora of Turkey and The East Aegean Island. Univ. Press, Edinburgh.
- Hossain, M.B., Rai, D.K., Brunton, N.P., Martin-Diana, A.B., Barry-Ryan, C., 2010. Characterization of phenolic composition in Lamiaceae spices by LC-ESIMS/MS. J Agr Food Chem. 58(19), 10576-10581.
- Kulisic, T., Radonic, A., Milos, M., 2005. Antioxidant properties of thyme (*Thymus vulgaris* L.) and wild thyme (*Thymus serpyllum* L.) essential oils. Italian Journal of Food Science, 17(3), 315-324.
- Tzima, K., Brunton, N.P., Rai, D.K., 2018. Qualitative and Quantitative Analysis of Polyphenols in Lamiaceae Plants- A Review. Plants (Basel). 7(25), 1-30.

* Presenting author. E-mail address: pelintastan35@gmail.com (Pelin Taştan)

S1.P287 Transformation of food by-products into high value added extracts with antimicrobial properties

Katarzyna Tyśkiewicz^{a,*}, Marcin Gruba^a, Marcin Konkol^a, Anita Wziątek^a, Zygmunt Fekner^a, Bernard Pawlak^a, Rafał Kowalski^a, Rafał Wiejak^a, Vicente Montes Jiménez^b, Joran Verspreet^c

^a *Lukasiewicz Research Network – New Chemical Syntheses Institute, Al. Tysiąclecia Państwa Polskiego 13A 24-110 Puławy, Poland*

^b *Department of Organic and Inorganic Chemistry, University of Extremadura, 06006 Badajoz, Spain*

^c *Flemish Institute for Technological Research (VITO), Boeretang 200, 2400 Mol, Belgium*

The transformation of food by-products into high-value added extracts with antimicrobial properties involves several steps and processes aimed at extracting bioactive compounds from these by-products and enhancing their antimicrobial activity. The first step involves selecting appropriate food by-products that are rich in bioactive compounds and have the potential for antimicrobial properties. Examples of such by-products include fruit peels, vegetable scraps, seed husks, and spent grains from brewing. Food by-products often require pre-treatment to facilitate the extraction process. This can involve washing, drying, grinding, or any other method to prepare the by-products for extraction.

Various extraction techniques can be employed to isolate bioactive compounds from the food by-products, including supercritical fluid extraction (SFE) as a green technique. After extraction, the crude extract may undergo fractionation and purification to isolate specific bioactive compounds responsible for antimicrobial activity. Techniques such as column chromatography, thin-layer chromatography, or high-performance liquid chromatography (HPLC) can be used for this purpose. Then the obtained are evaluated for their antimicrobial activity against a range of microorganisms such as bacteria, fungi, and viruses. This can be done using standard microbiological methods, including agar diffusion assays, broth microdilution assays, or other relevant techniques.

By transforming food by-products into high-value added extracts with antimicrobial properties, not only are valuable compounds recovered and utilized, but also environmentally sustainable practices are promoted by reducing waste and creating value from otherwise discarded materials.

Keywords: bioactive compounds, by-products, food, extracts, valorization

References

- Galanakis, C. M. 2012. Recovery of high added-value components from food wastes: conventional, emerging technologies and commercialized applications. *Trends in Food Science & Technology*, 26, 68–87.
- Teixeira, A., Baenas, N., Dominguez-Perles, R., Barros, A., Rosa, E., Moreno, D. A., & Garcia-Viguera, C. 2014. Natural bioactive compounds from winery by-products as health promoters: a review. *International Journal of Molecular Sciences*, 15, 15638–15678.
- Bampidis, V. A., & Robinson, P. H. 2006. Citrus by-products as ruminant feeds: A review. *Animal Feed Science and Technology*, 128, 175–217.
- Nazzaro, F., Fratianni, F., De Martino, L., Coppola, R., & De Feo, V. 2013. Effect of essential oils on pathogenic bacteria. *Pharmaceuticals*, 6, 1451–1474.

Acknowledgments

The research was financed by the European Commission as a part of the Circular Bio-based Europe Joint Undertaking project, entitled: *circular bioeconomy Business models owned by agroCOOPERatives* (project no. 101112407, project acronym: ROBOCOOP-EU).

* Presenting author. E-mail address: katarzyna.tyskiewicz@ins.lukasiewicz.gov.pl (Katarzyna Tyśkiewicz)

S1.P288 Secondary metabolites as chemotaxonomic markers in *Salix myrsinifolia* and *S. starkeana* (Salicaceae) twigs extracts

Enass Salih^{a,*}, Nina Sipari^b, Eunice Ego Mgbeahuruike^a, Pia Fyhrqvist^a

^a Division of Pharmaceutical Biosciences, Faculty of Pharmacy, 00014-University of Helsinki, Finland

^b Viikki Metabolomics Unit, Organismal and Evolutionary Biology research Programme, Faculty of Biological and Environmental Sciences, 00014-University of Helsinki, Finland

Chemotaxonomy is based on the special metabolite profiles in plants and is, together with the morphological characters, used as a tool in plant taxonomy and species identification (Julkunen- Tiitto, 1986). *Salix starkeana* and *S. myrsinifolia* (Salicaceae) are dioecious shrubs native to the temperate biome. Species identification within the genus *Salix*, and especially when it comes to the dormant winter twigs, can be difficult when based solely on morphological characters. Thus, we subjected twig extracts of *S. starkeana* and *S. myrsinifolia* to UPLC/QTOF-MS analysis, to elucidate the relations between these *Salix* spp. and to provide an additional guide to *Salix* taxonomy and identification.

We found that the *S. myrsinifolia* twig methanol extract exhibited various resemblances with the *S. starkeana* twigs methanol extract. For example, the condensed tannin, procyanidin B1 [*M-H*]⁻ 577,1348 and its isomers and the procyanidin monomers, epigallocatechin and catechin, were the main compounds in both species, although these compounds varied in quantity between the two species. Many compounds, and among them benzyl-β-primeveroside, caffeoylhexose, an isolariciresinol-pentoside isomer, salicortin, salicin-7-sulfate, naringenin and acetyl-*O*-salicortin were present in both species. Previously, salicin-7-sulfate, which is a novel salicylate in the genus *Salix*, was found only in *Salix caprea* (Lackus et al., 2020) and *Salix koryanagi* (Noleto-Dias et al., 2018).

However, some metabolites in *S. starkeana*, such as the unknown compounds at [*M-H*]⁻ 495,1511, (tR 2,97 min) and at [*M-H*]⁻ 453,1401 (tR 2,39 min) were found to be specific for *S. starkeana* but absent from *S. myrsinifolia*. Thus, these compounds may be used as chemotaxonomic markers to distinguish *S. starkeana* winter twigs.

It is worth to conclude that for pharmaceutical purposes, it is important to distinguish chemotypes with the most beneficial composition of therapeutic phytochemicals.

Keywords: *Salix myrsinifolia*, *S. starkeana*, Chemotaxonomy, UPLC-QTOF/MS, phenolic

References

Julkunen-Tiitto, R., 1986. A chemotaxonomic survey of phenolics in leaves of northern *Salicaceae* species. *Phytochemistry*. 25, 663-667.

Lackus, N.D., Müller, A., Kröber, T.D., Reichelt, M., Schmidt, A., Nakamura, Y., Paetz, C., Luck, K., Lindroth, R.L., Constabel, C.P., Unsicker, S.B., 2020. The occurrence of sulfated salicinoids in poplar and their formation by sulfotransferase1. *Plant physiol.* 183,137-151.

Noleto-Dias, C., Ward, J.L., Bellisai, A., Lomax, C., Beale, M.H., 2018. Salicin-7-sulfate: A new salicinoid from willow and implications for herbal medicine. *Fitoterapia*, 127, pp.166-172.

* Presenting author. E-mail address: enass.salih@helsinki.fi (Enass Salih)

S1.P289 Carvacrol solution and copper nanoparticles as an effective and non-phytotoxic treatments of carrot seeds against the pathogenic bacterium *Xanthomonas hortorum* pv. *carotae*

Dorota Tekielska^{a*}, Eliška Hakalová^a, Jana Čechová^a, Zuzana Bytešníková^b, Miroslav Baránek^a

^a Mendel University in Brno, Faculty of Horticulture, Mendeum–Department of Genetics, Valtická 334, 69144, Lednice, Czech Republic

^b Mendel University in Brno, Faculty of AgriSciences, Department of Chemistry and Biochemistry, Zemědělská 1, 61300 Brno, Czech Republic

The effectiveness of three phenolic compounds: carvacrol, eugenol and thymol, as well as three nanomaterials based on copper, silver and silver–selenium for elimination of bacterial pathogen from carrot seeds was evaluated in two presented studies. *Xanthomonas hortorum* pv. *carotae*, causing bacterial blight, leads to serious loss in carrot production worldwide, especially in seed production because of its seed-borne character. The treatment of bacterial diseases in plant production is challenging due to the restrictions in antibiotic use. Currently available methods for suppression of *Xanthomonas hortorum* pv. *carotae* in carrot seeds do not always provide a satisfactory effect, can reduce a germination and burden an environment. Therefore, the new alternative treatments have been tested *in vitro* and *in vivo*. The lowest minimum bactericidal concentration was observed for carvacrol and thymol (0.0098–0.0196%) and silver nanoparticles (10 mg/L⁻¹). The application of tested chemicals *in vivo* showed that the biological compound carvacrol at concentration 0.0196% completely eliminated pathogenic bacterium quantified in plants germinated from infected seeds. From three nanomaterials, only use of copper nanoparticles at concentration 993 mg/L⁻¹ resulted in a significant reduction of the bacterium quantity in carrot plants. The quantification was performed using real-time PCR assay targeting the hypersensitive response and pathogenicity-associated phosphatase (*hpaP*) gene. Both treatments did not reduce the seed germination percentage in comparison to non-treated controls and did not cause any visible morphological changes of plants, therefore can be considered as non-phytotoxic.

Keywords: bacterial blight, seed treatment, biological bactericide, phenolic compounds, nanomaterials

Acknowledgments

This article was created based on support from the project QK22010031 “Use of innovative potential of nanotechnology to enhance the rentability of selected areas of agricultural production” (funded by The Ministry of Agriculture of the Czech Republic).

*Presenting author. E-mail address: dorota.tekielska@mendelu.cz (Dorota Tekielska)

S1.P290 Characterization of polyphenolic compounds from an antibacterial twig methanol extract of *Salix aurita*

Eunice Ego Mgbeahuruike^{a,*}, Nina Sipari^b, Enass Salih^a, Pia Fyhrqvist^a

^a Division of Pharmaceutical Biosciences, Faculty of Pharmacy, University of Helsinki, Finland

^b Viikki Metabolomics Unit, Viikki Plant Science Centre, University of Helsinki, Finland

Polyphenols are natural and biologically active compounds found in plants, fruits and vegetables. They have numerous health benefits and serve as dietary antioxidants in promoting human health (Abdulkhaleq et al., 2017). Polyphenolic compounds that are common in *Salix* spp., such as the proanthocyanidin monomers; catechin and epicatechin, the procyanidins, and cinnamic acid and its derivatives, have been reported to be effective in the management and treatment of infectious diseases, diabetes, and cancer (Bouarab-Chibane et al., 2019; Adisakwattana, 2017). *Salix* species such as *S. aurita* L., are used traditionally in the treatment of inflammation and fever, and their efficacy has been attributed to the presence of its pharmacologically active compounds such as salicylates and polyphenols (Noleto-Dias et al., 2018).

The aim of this study was to conduct a phytochemical investigation and chemical profiling to identify the polyphenolic compounds present in a *Salix aurita* methanol twig extract using UPLC-PDA-QTOF/MS (ESI-mode). In our previous studies this extract showed good antibacterial activity. From the results, condensed tannins, flavonoids and salicylates were determined from a methanol twig extract of *S. aurita*. The chemical identification and molecular mass determination revealed procyanidin B1 (m/z 577.1346), naringenin (m/z 271.0606), salicin-7-sulfate (m/z 365.0542), salicin (m/z 285.0977), salicortin (m/z 423.1291), catechin (m/z 289.0712), luteolin (m/z 285.0399), and taxifolin (m/z 303.0505) as the major polyphenolic compounds present in the twigs of *S. aurita*. This study will pave way for our further research on the mechanism of actions of these identified polyphenols as antibacterial agents.

Keywords: *Salix aurita*, polyphenols, UPLC-PDA-QTOF/MS

References

- [1] Abdulkhaleq, L. A., Assi, M. A., Noor, M. H. M., Abdullah, R., Saad, M. Z., & Taufiq-Yap, Y. H. (2017). Therapeutic uses of epicatechin in diabetes and cancer. *Veterinary world*, 10(8), 869.
- [2] Adisakwattana, S. (2017). Cinnamic acid and its derivatives: mechanisms for prevention and management of diabetes and its complications. *Nutrients*, 9(2), 163.
- [3] Bouarab-Chibane, L., Forquet, V., Lantéri, P., Clément, Y., Léonard-Akkari, L., Oulahal, N., ... & Bordes, C. (2019). Antibacterial properties of polyphenols: characterization and QSAR (Quantitative structure–activity relationship) models. *Frontiers in microbiology*, 10, 440698.
- [4] Noleto-Dias, C., Ward, J. L., Bellisai, A., Lomax, C., & Beale, M. H. (2018). Salicin-7-sulfate: A new salicinoid from willow and implications for herbal medicine. *Fitoterapia*, 127, 166-172.

*Presenting author. E-mail address: Eunice.mgbeahuruike@helsinki.fi (Eunice Ego Mgbeahuruike)

S1.P291 Antimicrobial and anti-biofilm properties of geranylated flavonoids from *Paulownia tomentosa* Steud. fruit

Lenka Molčanová^{a,*}, Adam Tesárek^b, Karel Šmejkal^a, Jakub Trembl^b

^a Department of Natural Drugs, Faculty of Pharmacy, Masaryk University, Palackého tř. 1946/1, 612 00 Brno, Czech Republic

^b Department of Molecular Pharmacy, Faculty of Pharmacy, Masaryk University, Palackého tř. 1946/1, 612 00 Brno, Czech Republic

Paulownia tomentosa (Thunb.) Steud. (Paulowniaceae), a traditional Chinese medicinal plant, had been used for centuries as a component of remedies for various illnesses. It is a rich source of phenolic compounds, mainly geranylated flavonoids, which are currently studied for their promising biological activities.

An increasing resistance to conventional antibiotics is a threat to the global healthcare system, and therefore it is necessary to search for new alternatives in the treatment of bacterial infections. Also, production of biofilm by certain bacteria is connected to persistence of their infections and it is difficult to eradicate them. Thus, new treatments against such long lasting infections that inhibit biofilms are needed.

This experiment was carried out on a biofilm-producing gram-positive bacteria *Staphylococcus epidermidis*. Ten C-geranylated flavanones and dihydroflavonols (**1** – **10**) isolated from the fruit of *P. tomentosa* were tested for their antimicrobial and anti-biofilm properties. The evaluation of antimicrobial activity was performed using the microdilution method and the anti-biofilm activity was evaluated by crystal violet staining.

As a result of our study, seven of the tested compounds showed antimicrobial activity and six of them revealed anti-biofilm activity. Diplacone (**1**), diplacol (**4**), and 3'-O-methyl-5'-hydroxydiplacone (**6**) showed the most potent antimicrobial activity. Diplacone (**1**), mimulone (**2**), and mimulol (**10**) were the most effective in suppressing bacterial biofilm formation. These compounds need to be further investigated for their antimicrobial or anti-biofilm activity and safety, but there is a potential for their future use in the suppression of antibiotic resistance.

Keywords: anti-biofilm activity, antimicrobial activity, geranylated flavonoids, *Paulownia tomentosa*, *Staphylococcus epidermidis*

Acknowledgments

This work was supported by the Czech Science Foundation, grant number GA23-04655S: Role of prenylation and glycosylation patterns in anti-inflammatory activity and metabolism of natural phenolic compounds.

* Presenting author. E-mail address: molcanoval@pharm.muni.cz (Lenka Molčanová)

S1.P292 Navigating Experimental Metabolomics Knowledge Graphs for Targeted Isolation of Anti-Infective Natural Products

Olivier A. Kirchhoffer^{1,2*}, Jahn Nitschke³, Louis-Félix Nothias^{1,2,4}, Luis Quiros-Guerrero^{1,2}, Laurence Marcourt^{1,2}, Nabil Hanna³, Antonio Grondin⁵, Emerson F. Queiroz^{1,2}, Thierry Soldati³, Jean-Luc Wolfender^{1,2}

¹ Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, CMU, 1211 Geneva, Switzerland

² School of Pharmaceutical Sciences, University of Geneva, CMU, 1211 Geneva, Switzerland

³ Department of Biochemistry, Faculty of Sciences, University of Geneva, Quai Ernest-Ansermet 30, 1205 Geneva, Switzerland

⁴ Université Côte d'Azur, CNRS, ICN, France

⁵ Green Mission Department, Herbal Products Laboratory, Pierre Fabre Research Institute, Toulouse, France

The escalating threat posed by Multidrug-Resistant Tuberculosis (MDR-TB) caused by *Mycobacterium tuberculosis* (Mtb) has spurred the necessity to find new antibiotics to treat tuberculosis. In our study, we employed an innovative 3R infection model using *Mycobacterium marinum* alongside the amoeba and professional phagocyte *Dictyostelium discoideum* as stand-ins for Mtb and macrophages, respectively. We used this model in a high-throughput phenotypic assay which enables simultaneous assessment of a compound's impact on both the host and the pathogen (Hanna et al., 2020).

To find novel anti-infective compounds, a registered diverse collection of 1600 plant extract samples (Pierre Fabre Library, PFL) was screened using this assay. In parallel, this collection was analyzed by untargeted UHPLC-high-resolution tandem mass spectrometry (UHPLC-HRMS/MS) (Allard et al., 2023).

In a careful two-step validation process, we eventually reduced the collection to a list of 13 validated hit extracts. We then applied High-resolution HPLC microfractionation bioactivity profiling to swiftly pinpoint bioactive compounds within these selected hit extracts (Hell et al., 2022). Furthermore, by leveraging experimental metabolomic data organized within a knowledge graph (KG) (Gaudry et al., 2023), we could systematically explore the virtual chemical space generated. This approach facilitated the targeted isolation of new/bioactive structural analogues based on annotations, also in extracts not necessarily targeted in the initial bioactivity screening (Kirchhoffer et al., 2023). These were then fully characterized by NMR and electronic circular dichroism (ECD) to establish their absolute stereochemistry. The example chosen serves as a proof of concept for the use of a KG (integrating bioactivity and metabolomics) to efficiently identify bioactive NPs in collections of plant extracts.

Keywords: knowledge graph (KG), 3R infection model, novel anti-infective compounds, high-resolution tandem mass spectrometry (HRMS/MS), plant extracts library

References

- Allard P-M, Gaudry A, Quirós-Guerrero L-M, et al. Open and reusable annotated mass spectrometry dataset of a chemodiverse collection of 1,600 plant extracts. *GigaScience* 2023; 12: giac124.
- Gaudry A, Pagni M, Mehl F, et al. A Sample-Centric and Knowledge-Driven Computational Framework for Natural Products Drug Discovery. *ChemRxiv*. 2023.
- Hanna N, Kicka S, Chiriano G, et al. Identification of anti-mycobacterium and anti-legionella compounds with potential distinctive structural scaffolds from an HD-PBL using phenotypic screens in amoebae host models. *Frontiers in Microbiology* 2020; 11: 266.
- Hell T, Rutz A, Dürr L, et al. Combining Activity Profiling with Advanced Annotation to Accelerate the Discovery of Natural Products Targeting Oncogenic Signaling in Melanoma. *J Nat Prod* 2022; 85: 1540–1554.
- Kirchhoffer OA, Nitschke J, Allard P-M, et al. Targeted isolation of natural analogs of anti-mycobacterial hit compounds based on the metabolite profiling of a large collection of plant extracts. *Frontiers in Natural Products* 2023; 2.

* Presenting author. E-mail address: olivier.kirchhoffer@unige.ch (Olivier A. Kirchhoffer)

Biotransformation and biosynthesis of natural products

S1.P293 Discovery of novel terpene hydrocarbon scaffolds and their associated tailoring enzymes from *Streptomyces* bacteria

Diana Łomowska-Keehner*, Tyler Alsup, Melvin Osei Opoku, Xiuting Wei, Wenbo Ning, Caitlin McCadden, Jeffrey D. Rudolf

Department of Chemistry, University of Florida, 1645 W University Ave, Gainesville, FL, 32601, USA

Bacterial natural products (NPs) from the genus *Streptomyces* provide approximately 70% of antibiotics used to date (Kieser et al., 2000). Given the growing antimicrobial resistance crisis, it is critical to discover novel compounds with antibiotic activity. In this work, we look towards terpenoids, the largest class of NPs with over 100,000 characterized compounds (Rudolf et al., 2021). Terpenoids have broad bioactivities including antibacterial, cytotoxic, and anti-inflammatory properties. Historically, plants were considered the most prolific producers of terpenoids. However, genome mining reveals that bacteria, especially *Streptomyces*, have great terpenoid biosynthetic potential. We genome mined for uncharacterized bacterial terpene synthases (TSs) encoded in NP biosynthetic gene clusters (BGCs) also containing cytochrome P450 enzymes that may generate novel terpene scaffolds. These bacterial TSs of unknown function were heterologously expressed in *E. coli* containing upregulated precursor genes and a kinase system, allowing conversion of exogenously added isoprenol into terpene precursors *in vivo*. This system facilitates detection of terpenes at small scale to validate TS activity and supports reliable terpene production in large scale fermentations. *Streptomyces* normally produce complex NP mixtures; furthermore, BGCs with TSs of interest often are not expressed under laboratory conditions. Thus, expressing TSs in *E. coli* decreases complexity of fermentation extracts and increases the titer of terpenes produced. Once TS products are characterized, we can express tailoring enzymes including P450s clustered with TSs producing novel terpene scaffolds to discover highly functionalized terpenoid NPs for bioactivity testing. This poster will showcase novel terpenes isolated thus far from previously uncharacterized bacterial TSs, our heterologous expression workflow, and BGCs under study.

Keywords: biosynthesis, *Streptomyces*, heterologous expression, terpene synthases, cytochrome P450s

References

- Kieser, T., Bibb, M.J., Buttner, M.J., Chater, K.F., Hopwood, D.A., 2000. Practical *Streptomyces* Genetics (2nd ed.). Norwich, England: John Innes Foundation.
- Rudolf, J. D., Alsup, T. A., Xu, B., Li, Z. 2021. Bacterial terpenome. Nat. Prod. Rep. 38, 905-980.

* Presenting author. E-mail address: dstancic@chem.ufl.edu (Diana Łomowska-Keehner)

S1.P294 Effects of difference LED light colors on isoflavonoid production in callus cultures of *Pueraria candollei* Wall. ex Benth. (Fabaceae)

Worapol Sae-Foo^{a,*}, Witsarut Kraithong^a, Sukritta Singkham^a, Gorawit Yusakul^b, Waraporn Putalun^a

^a Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen, Thailand

^b School of Pharmacy, Walailak University, Nakhon Si Thammarat, Thailand

The tuberous root of *Pueraria candollei* Wall. ex Benth. (Fabaceae) is in high demand for dietary supplements and cosmetics, making it scarce. Callus culture is a source of secondary metabolites production (Udomsin et al. 2020). The primary factor for *in vitro* plant tissue culture is the light source. Light-emitting diode (LED) lights were introduced to plant tissue culture in recent years (Bantis et al., 2018). The current work used fluorescent, red, blue, white LED, and dark conditions to investigate the phytoestrogen compounds production in *P. candollei* callus for 14 and 28 days. Chalcone synthase (*CHS*), isoflavone synthase (*IFS*), isoflavone reductase (*IFR*), hydroxyisoflavanone dehydratase (*HID*), cytochrome P81E (*CYP81E*) and prenyltransferase-1 (*PT-1*) related biosynthesis pathway genes were determined expression level using quantitative reverse transcription-polymerase chain reaction (Suntichaikamolkul et al., 2019). The white LED condition showed the highest growth rate at day 28 increase of 1.89-fold compared to the initial culture day. White LED light stimulated a significant upregulation of *HID*, *CYP81E*, and *PT-1* genes by 1.6, 4.8, and 56-fold, respectively. Treated red LED callus extract showed daidzin, genistin, and deoxymiroestrol levels of 612.83 ± 42.80 , 549.39 ± 32.72 , and 1.46 ± 0.24 $\mu\text{g/g}$ dry weight, respectively. The total isoflavonoids concentration of callus extract increased 3 -fold and 2-fold under red and white LED, respectively compared to fluorescent light. The presence of red and white LED lights can stimulate the formation of phytoestrogens in *P. candollei* callus, particularly in the isoflavonoids and their derivatives. However, further studies should be performed to confirm the pharmacological activity of plant extracts.

Keywords: LED light, *Pueraria candollei*, Fabaceae, gene expression, isoflavonoids

References

- Bantis, F., Smirnakou, S., Ouzounis, T., Koukounaras, A., Ntagkas, N., Radoglou, K., 2018. Current status and recent achievements in the field of horticulture with the use of light-emitting diodes (LEDs). *Sci. hortic.* 235, 437-451.
- Suntichaikamolkul, N., Tantisuwannichkul, K., Prombutara, P., Kobtrakul, K., Zumsteg, J., Wannachart, S., Schaller, H., Yamazaki, M., Saito, K., De-eknamkul, W., Vimolmangkang, S., Sirikantaramas, S. (2019). Transcriptome analysis of *Pueraria candollei* var. *mirifica* for gene discovery in the biosyntheses of isoflavones and miroestrol. *BMC. Plant. Biol.* 19, 1-14.
- Udomsin, O., Yusakul, G., Kitisripanya, T., Juengwatanatrakul, T., Putalun, W., 2020. The deoxymiroestrol and isoflavonoid production and their elicitation of cell suspension cultures of *Pueraria candollei* var. *mirifica*: from shake flask to bioreactor. *Appl. Biochem. Biotech.* 190, 57-72.

* Presenting author. E-mail address: worapol.s@kkumail.com (Worapol Sae-Foo)

S1.P295 Elicitation of licorice callus for improving flavonoids production

Supaluk Krittanai, Rattanathorn Choonong, Khunkhang Butdapheng, Saponanai Vongwairat, Khomkrit Niamin, Pisitchai Masugarut, Saharat Patiwettham, Waraporn Putalun*

Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen 40002, Thailand

Licorice is an herbal plant that has been used as a medicinal source since the ancient era. The current issue with this plant is the depletion of raw materials in natural cultivation (Kwon et al., 2020). The callus culture is conducting research to resolve this problem. Ononin and its aglycone, formononetin, are bioactive licorice flavonoids with well-established anti-inflammatory activity (Krittanai et al., 2021). In this study, a combination of sample pre-treatment and elicitation using methyl jasmonate (MJ) or cellulase enzyme (CL) was implemented to increase the yield of these substances in *Glycyrrhiza inflata* callus culture. The largest ononin induction was seen on day 3, with a 3- and 5.93- fold induction in MJ and CL, respectively, compared to the control group. In the case of formononetin, the highest production was observed on day 7, with a 3.22- and 5.46-fold induction over the control. The mechanism of elicitation was revealed as stimulating the antioxidant defense mechanism and stimulating the flavonoid biosynthesis genes (*CHI*, *CHS*, and *IFS*). Therefore, the combination strategies of sample pre-treatment and elicitation present in the current study can be applied for enhancing the medicinal substance in licorice, especially in the form of callus culture, which has high potential to be the future source for the production of medicinal substances.

Keywords: elicitation, methyl jasmonate, cellulase enzyme, *Glycyrrhiza inflata*, callus

References

- Kwon, Y.J., Son, D.H., Chung, T.H., Lee, Y.J., 2020. A Review of the pharmacological efficacy and safety of licorice root from corroborative clinical trial findings. *J. Med. Food.* 23, 12–20.
- Krittanai, S., Buraphaka, H., Sakamoto, S., Putalun, W., 2021. Enhancement of nutraceutical properties of licorice callus cultures using sample pre-treatment strategy. *Int. J. Food Sci. Technol.* 56, 5843–5852.

* Presenting author. E-mail address: waraporn@kku.ac.th (Waraporn Putalun)

S1.P296 Chemopreventive properties of *Alchemilla vulgaris* L. extract biotransformed by the human gut microflora – preliminary study of a human cellular model of colon cancer

Katarzyna Jakimiuk^{a,*}, Aleksandra Kruk^b, Marta Kinga Lemieszek^c, Jakub W. Strawa^a, Sebastian Granica^d, Urszula Kuszel^e, Aleksandra Komendo^e, Adrian Wiater^e, Michał Tomczyk^a

^a Department of Pharmacognosy, Faculty of Pharmacy with the Division of Laboratory Medicine, Medical University of Białystok, ul. Mickiewicza 2a, 15-230 Białystok, Poland

^b Microbiota Lab, Center for Preclinical Studies, Faculty of Pharmacy, Medical University of Warsaw, ul. Banacha 1, 02-097 Warsaw, Poland

^c Department of Medical Biology, Institute of Rural Health, ul. Jaczewskiego 2, 20-090 Lublin, Poland

^d Department of Pharmaceutical Biology, Faculty of Pharmacy, Medical University of Warsaw, ul. Banacha 1, 02-097 Warsaw, Poland

^e Student's Scientific Association, Faculty of Pharmacy with the Division of Laboratory Medicine, Department of Pharmacognosy, Medical University of Białystok, ul. Mickiewicza 2a, 15-230 Białystok, Poland

^f Department of Industrial and Environmental Microbiology, Institute of Biological Sciences, Maria Curie-Skłodowska University, ul. Akademicka 19, 20-033 Lublin, Poland

Numerous scientific reports have demonstrated that substances introduced into the body are subject to metabolic changes caused by bacteria in the digestive tract (Kamada et al., 2013). Thus, we evaluated the chemical composition of lady's mantle herb extract subjected to gut biotransformation and assessed the possibility of using the obtained products for colon cancer chemoprevention. First, extraction was carried out using a eutectic mixture (urea/choline chloride). The obtained extract was exposed to the gut microflora from 3 donor faecal samples (D1-D3) in an anaerobic chamber. After incubation (24 h), the changes in the phytochemical composition of the biotransformed extract were assessed, followed by LC-MS characterization. In the next step, the impact of the extracts on human colon CCD841 CoN and HT-29 cells was examined. Raw-extract (non-biotransformed) did not affect either cancer or normal colon cells. Biotransformed extracts at concentrations of 25-250 µg/mL did not cause any changes in the viability or proliferation of colon epithelial cells, but did have a dose- dependent cytotoxic effect on HT-29 cells. LDH release from damaged HT-29 cells in response to D1, D2 and D3 at a concentration of 250 µg/mL increased by 12.6%, 25.3% and 30.0%, respectively. Moreover, biotransformed extracts at concentrations ranging from 50-250 µg/ml significantly decreased the proliferation of HT-29 cells. The strongest effect was caused by extract D3 (IC₅₀ = 471 µg/mL), while the weakest change was induced by extract D1 (IC₅₀ = 1440 µg/mL). The performed studies revealed the chemopreventive properties of the biotransformed extracts of *Alchemilla vulgaris* L. (Rosaceae).

Keywords: *Alchemilla vulgaris*, microbiota, colorectal cancer

References

Kamada, N., Seo, S.U., Chen, G.Y., Núñez, G., 2013. Role of the gut microbiota in immunity and inflammatory disease. *Nat. Rev. Immunol.* 13, 321–335.

* Presenting author. E-mail address: katarzyna.jakimiuk@umb.edu.pl (Katarzyna Jakimiuk)

S1.P297 Chemical characterization, prebiotic and anti-diabetic effects of galactomannans from two traditional Chinese medicines

Ching-Yi Lu^a, Hui-Kang Liu^{b,c}, Mei-Kuang Lu^b, Shoei-Sheng Lee^a, Chia-Chuan Chang^{a,*}

^a Department of Pharmacy, National Taiwan University, Taipei City, Taiwan

^b Natural Research Institute of Chinese Medicine, Taipei City, Taiwan

^c Graduate Institute of Pharmacognosy, Taipei Medical University, Taipei, Taiwan

The galactomannans of *Trigonella foenum-graecum* L. (fenugreek) and the fungus *Cordyceps militaris* L. are two TCMs that demonstrate anti-diabetic activities. Such activities might be caused by modulation of the homeostasis of gut microbiota. Among the gut microbes, *Bacteroides ovatus* is more abundant (about 1.2-fold) in healthy individuals than in diabetes patients. Another microbe, *Akkermansia muciniphila*, is known to degrade mucin and produce short-chain fatty acids (SCFAs), which aid in the growth of other bacteria and maintain healthy mucus. To chemically characterize and evaluate the prebiotic effects of the galactomannans from the two TCMs, they were column chromatographed to give 19 fractions. The major monosaccharide components of these were Man, Gal and Glc (in the ratio about 5:3:1 to 4:3:1). Among these, TFPS-1 (carbohydrates obtained from fenugreek), CMMPS-1a02 and CMFPS-1a03 (carbohydrates obtained from *C. militaris* L.), improved the growth of *B. ovatus* and *A. muciniphila* (1.3~1.5-fold) with the production of SCFAs (i.e., acetic acid, propionic acid and butyric acid, about 2~4-fold increase, respectively). To evaluate the anti-diabetic effects, these polysaccharides were evaluated by three *in vitro* assays. Among these, CMMPSs triggered the secretion of GLP-1 (STC-1 cell, 1.8-fold) and insulin (BRIN-BD11 cell, 1.5-fold); TFPS-5, CMMPS-1a02, CMFPS-1a03 showed significant inhibitory effect of DPP-4 enzyme; the combination of carbohydrates derived from *C. militaris* and cordycepin could enhance the DPP-4 inhibitory effect.

Keywords: fenugreek, *Cordyceps militaris*, galactomannans, gut microbiota, anti-diabetic effects

References

Lee CT, Huang KS, Shaw JF, Chen JR, Kuo WS, Shen GX, Grumezescu A.M., Holban A.M., Wang YT, Wang JS, Hsiang YP, Lin YM, Hsu HH, Yang CH, 2020. Trends in the Immunomodulatory Effects of *Cordyceps militaris*: Total Extracts, Polysaccharides and Cordycepin. *Frontiers in Pharmacology*. 11, 575704.

Kalyan C. Nagulapalli Venkata, Anand Swaroop, Debasis Bagchi, Anupam Bishayee, 2017. A small plant with big benefits: Fenugreek (*Trigonella foenum-graecum* Linn.) for disease prevention and health promotion. *Molecular Nutrition and Food Research*, 61, 1600950.

Jixian Zhang, Chaoting Wen, Yuqing Duan, Haihui Zhang, Haile Ma, 2019. Advance in *Cordyceps militaris* (Linn) Link polysaccharides: Isolation, structure, and bioactivities: A review. *International Journal of Biological Macromolecules*, 132, 906–914.

* Presenting author. E-mail address: d10423202@ntu.edu.tw (Ching-Yi Lu)

S1.P298 Phytochemical profiling of phenolic metabolites in flower buds of *Magnolia kobus* DC. and their antioxidant activity

Aldona Adamska-Szewczyk^{a,*}, Grażyna Zgórka^b

^a Bionorica Poland, Hrubieszowska 6B Str., 01-209 Warsaw, Poland

^b Department of Pharmacognosy with the Medicinal Plant Garden, Faculty of Pharmacy, Medical University of Lublin, 1 Chodzki Str., 20-093 Lublin, Poland

Some representatives of the genus *Magnolia* L. (Magnoliaceae) are popular in traditional Asian medicine systems due to their broad spectrum of bioactive specialised metabolites (Poivre et al., 2017). Following our previous research on *Magnolia × soulangeana* Soul.-Bod. var. 'Lennei' (Zgórka et al., 2023), dried flower buds of *M. kobus* DC. collected from two different locations in Poland, namely the Warsaw University Botanical Garden (MKW) and the Rogów Arboretum (MKR), were subjected to ultrasound-assisted extraction using 30, 50 and 70% (V/V) EtOH. Herbal preparations were analysed using coupled chromatographic (RP-LC), spectroscopic (PDA) and mass spectrometric (QTOF/ESI-MS/MS) techniques. The results showed similar quantitative profiles of the main phenolic compounds, including phenolic acids, phenylethanoids, flavonols and lignans. However, their total content, calculated as gallic acid equivalent (GAE), increased significantly with the elution strength of the extraction solvent and was highest in extracts obtained with 70% EtOH (60.59 and 56.76 mg GAE/g dry weight, for MKW and MKR, respectively). Interestingly, the antiradical capacity of the above-mentioned extracts, as assessed by the DPPH• assay, was strongly correlated with the total phenylethanoid content, which was about three times higher in MKR than MKW, so the former extract exhibited higher antiradical activity (IC₅₀ ~25.90 µg/mL) than the latter (IC₅₀ ~33.76 µg/mL). Our study showed a relationship between environmental factors (UV exposure) in the plant habitat and the content of phenolic products of biotransformation, especially phenylethanoids, which are synthesised in the flower buds of the same magnolia taxon.

Keywords: *Magnolia kobus* DC., phenolics, antioxidant properties

References

- Poivre, M., Duez P., 2017. Biological activity and toxicity of the Chinese herb *Magnolia officinalis* Rehder & E. Wilson (Houpo) and its constituents. J Zhejiang Univ Sci B. 18, 194-214.
- Zgórka, G., Adamska-Szewczyk, A., Baj, T., 2023. Response Surface Methodology in Optimising the Extraction of Polyphenolic Antioxidants from Flower Buds of *Magnolia × soulangeana* Soul.-Bod. var. 'Lennei' and Their Detailed Qualitative and Quantitative Profiling. Molecules 28, 6335.

* Presenting author. E-mail address: a.adamskaszewczyk@gmail.com (Aldona Adamska-Szewczyk)

S1.P299 A research of the optimization of recombinant protein expression based on plant system

Juho Lee*, Woo-Hyun Jeong, Seon-Kyeong Lee, Kyeong-Ryeol Lee

National Institute of Agricultural Sciences, Rural Development Administration, Jeonju, Korea

Plants are generally suitable for recombinant protein production. Recombinant proteins can be rapidly produced in 4-6 weeks using transient expression (Kapila et al., 1997). In addition, plant production systems cannot be infected by mammalian pathogens during protein production and are easy to scale up (Fischer et al., 1999). To effectively express recombinant proteins in plants, we introduce stable and transient expression systems in this study. In order to find the optimal stable expression combination for each promoter and terminator, plastid transformation was performed to confirm the level of GFP expression (Fig. 1). As a result, it was found that when the *rbcL* promoter-*psbA* terminator was used, the GFP expression level was 2 folds higher than when original combination (the *rrn* promoter-*psbA* terminator) was used.




Fig. 1. Accumulation of GFP in promoter and terminator combination.

In addition, several types of vectors were designed to increase the expression level of the antigen protein of foot-and-mouth disease virus (FMDV) using transient expression. Among various combinations, the expression level of the antigen protein was high when the *csVMV* promoter and *pinII*-*extensin* double terminator were combined. In the future, we will compare the expression level of the antigen protein through stable expression using the *rbcL* promoter and transient expression using a double terminator to identify a better method of producing the antigen protein and evaluate immunogenicity.

Keywords: recombinant protein, transient expression, transgenic plant

References

- Kapila J., De Rycke R., Van Montagu M., Angenon G., 1997. An *Agrobacterium*-mediated transient gene expression system for intact leaves. *Plant Sci.* 122(1). 101–108.
- Fischer R., Drossard J., Commandeur U., Schillberg S., Emans N., 1999. Towards molecular farming in the future: moving from diagnostic protein and antibody production in microbes to plants. *Biotechnol. Appl. Biochem.* 30,101–108.

* Presenting author. E-mail address: jhlee0102@korea.kr (Juho Lee)

S1.P300 Cyanobacterial natural products: bioactive cyclic and linear peptides

Chin-Soon Phan^{a,b,*}, Kenichi Matsuda^{c,d}, Jakia Jerin Mehjabin^b, Zhengyi Ling^b, Kenji Washio^b, Masaaki Morikawa^b, Taiki Umezawa^b, Toshiyuki Wakimoto^{c,d}, Tatsufumi Okino^{b,e}

^a Latvian Institute of Organic Synthesis, Aizkrauklesielā 21, 1006 Riga, Latvia

^b Faculty of Environmental Earth Science, Hokkaido University, Kita-ku, Sapporo 060-0810, Japan

^c Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan

^d Global Station for Biosurfaces and Drug Discovery, Hokkaido University, Kita 12, Nishi 6, Sapporo 060-0812, Japan

^e Graduate School of Environmental Science, Hokkaido University, Kita-ku, Sapporo 060-0810, Japan

Chemical diversity and biological activity of natural products from cyanobacteria is continuing to be a subject of interest to researchers. This presentation focus on the chemistry and biology of cyanobacterial cyclic and linear peptides. Research on cyanobacterial natural products has allowed us to isolate 1) argicyclamides, a new class of cyanobactin with unique mono- and bis-prenylation on guanidine moieties from *Microcystis aeruginosa* NIES-88. We performed biochemical characterization on the new prenyltransferase, AgcF a key enzyme involved in enhancing antibiotic activity on the cyclic peptide, argicyclamide A (Phan et al., 2021); 2) nostosin G and spiroidesin B, new analogue of three to four residues containing linear peptides from *Dolichospermum* sp. NIES- 1697. We showed that nostosin G exhibited strong activity in trypsin inhibition, and the biosynthetic gene clusters shared similarities to spumigin except for their genetic organization, where an unrelated NRPS module was positioned in between two NRPS modules involved in nostosin G biosynthesis (Phan et al., 2022); 3) five new microginins and two new anabaenopeptins from *M. aeruginosa*. We also discovered a unique residue of 2-amino-5-(4'-hydroxyphenyl) pentanoic acid (Ahppa) at the ureido linkage of both new anabaenopeptins, which prompted us to investigate the biosynthetic origin of this non-proteinogenic amino acid Ahppa. We performed heterologous expression in *E. coli* and biochemical assay to show MaHphABCDE involved in double homologation of Tyr residue. These works disclosed bioactive natural products from cyanobacteria and characterized enzymes which expand the biocatalysis toolbox for prenylations and homologations.

Keywords: *Cyanobacterium*, natural products, biosynthesis, bioactivity

References

Phan, C.-S., Matsuda, K., Balloo, N., Fujita, K., Wakimoto, T., Okino, T., 2021. Argicyclamides A-C unveil enzymatic basis for guanidine bis-prenylation. *J. Am. Chem. Soc.* 143, 10083-10087.

Phan, C.-S., Mehjabin, J. J., Anas, A. R. J., Hayasaka, M., Onoki, R., Wang, J., Umezawa, T., Washio, K., Morikawa, M., Okino, T., 2022. Nostosin G and spiroidesin B from the cyanobacterium *Dolichospermum* sp. NIES-1697. *J. Nat. Prod.* 85, 2000-2005.

* Presenting author. E-mail address: chinsoon@osi.lv (Chin-Soon Phan)

S1.P301 Investigation of the effect of solid state fermentation on the release of phenolic compounds from by-products of the agro-food industry

Ewa Szczepańska*, Magdalena Waloszczyk, Filip Boratyński

Wroclaw University of Environmental and Life Sciences, Department of Food Chemistry and Biocatalysis, Norwida 25 Str., 50-375 Wrocław, Poland

The main purpose of this research was to obtain valuable phenolic compounds (PC) from the by-products of the agricultural and food industry - linseed oil cake (LOC) and brewer's spent grain (BSG) - using solid state fermentation (SSF) and submerged fermentation (SmF) culture modes. The by-products used in the study are lignocellulosic materials with a proven content of PC, which are desirable in the industry due to their biological properties. Indeed, they exhibit antioxidant properties, which makes them widely used in medicine, cosmetology and food industry (Kyselka et al., 2017). Various biocatalysts (29 *Ascomycota* and *Basidiomycota* strains) were used in the screening experiment. A comparative analysis of the two culture systems was performed, as well as the selection of the most effective biocatalysts in terms of their ability to degrade lignocellulosic raw materials and release PC. Qualitative and quantitative analysis of the extracts obtained from the fermentation media allowed to identify several PC such as: ferulic acid, *p*-coumaric acid, caffeic acid, vanillic acid, veratric acid, and vanillin (Fig. 1).




Fig. 1. Structures of identified phenolic compounds.

This study indicates that the use of biotechnological methods to obtain biologically active compounds is an attractive alternative to value difficult to recycle by-products of the agro-food industry. Moreover, it is consistent with sustainable development and the business model of using agricultural waste in the bioeconomy (Šelo et al., 2021).

Acknowledgments: The work was financed as part of the LIDER XII project financed by the National Center for Research and Development (Poland) under the contract No. LIDER/44/0228/L-12/20/NCBR/2021.

Keywords: agro-food by-products, brewer's spent grains, linseed cake, solid state fermentation, phenolic compounds

References

- Kyselka, J., Rabiej-Kozioł, D., Dragoun, M., Kreps, F., Burčová, Z., Němečková, I., Smolová, J., Bjelková, M., Szydłowska-Czerniak, A., Schmidt, S., Šarman, L., Filip, V., 2017. Antioxidant and antimicrobial activity of linseed lignans and phenolic acids. *Eur. Food Res. Technol.* 243, 1633–1644.
- Šelo, G., Planinić, M., Tišma, M., Tomas, S., Koceva Komlenić, D., Bucić-Kojić, A., 2021. A comprehensive review on valorization of agro-food industrial residues by solid-state fermentation. *Foods* 10, 927.

* Presenting author. E-mail address: ewa.szczepanska@upwr.edu.pl (Ewa Szczepańska)

S1.P302 Biosynthesis of gramine in barley by a cryptic oxidative rearrangement

Sara Leite Dias^a, Ling Chuang^{b,c}, Shenyu Liu^{b,c,*}, Benedikt Seligmann^b, Fabian L. Brendel^a, Benjamin G. Chavez^a, Robert E. Hoffie^d, Iris Hoffie^d, Jochen Kumlehn^d, Arne Bültemeier^{b,c}, Johanna Wolf^b, Marco Herde^e, Claus-Peter Witte^e, John C. D'Auria^a, Jakob Franke^{b,c}

^a Department of Molecular Genetics, Leibniz Institute for Plant Genetics and Crop Plant Research (IPK), Gatersleben, Germany

^b Institute of Botany, Leibniz University Hannover, Hannover, Germany

^c Centre of Biomolecular Drug Research, Leibniz University Hannover, Hannover, Germany

^d Department of Physiology and Cell Biology, Leibniz Institute for Plant Genetics and Crop Plant Research (IPK), Gatersleben, Germany

^e Department of Molecular Nutrition and Biochemistry of Plants, Leibniz University Hannover, Hannover, Germany

The indole alkaloid gramine protects barley and other grasses from insects. Early studies demonstrated that gramine is derived from tryptophan via the key intermediate aminomethylindole (AMI) by an unknown enzyme that removes C-1 and C-2 of tryptophan while forming a new C-N bond (Dieter et al., 1974, Ishikawa et al., 2023). In this study, we elucidated the biosynthetic pathway of gramine in barley by identifying the missing AMI synthase (AMIS). Co-expression of AMIS with a previously described *N*-methyltransferase gene in *Nicotiana benthamiana*, *Arabidopsis thaliana* and *Saccharomyces cerevisiae* enabled the production of gramine in heterologous platforms (Fig. 1).




Fig. 1. Gramine biosynthetic pathway.

As a further part, we demonstrated that AMIS catalyzes an oxidative Stieglitz-type rearrangement of tryptophan by a combination of isotope labeling experiments, intermediate trapping and by-product investigation. In summary, our study reveals the genetic and biochemical basis of an unusual amino acid chain shortening in the biosynthesis of gramine. This enables further biotechnological applications utilizing this alkaloid for cereal protection.

Keywords: biosynthesis, gramine, alkaloid, oxidative rearrangement, barley

References

- Diete, G., Lehmann, H., Schütte, H. R., 1974. Zur Biosynthese des Gramins: Biosynthesis of Gramine. *Biochem. Physiol. Pflanz.* 166, 281-287.
- Ishikawa, E., Kanai, S., Sue, M., 2023. Detection of a novel intramolecular rearrangement during gramine biosynthesis in barley using stable isotope-labeled tryptophan. *Biochem. Biophys. Rep.* 34, 101439.

* Presenting author. E-mail address: shenyu.liu@oci.uni-hannover.de (Shenyu Liu)

S1.P303 Urolithin production in *ex vivo* incubations of gut microbiota – survey on different substrates and antibiotics' influence

Dominik Popowski^{a,b,c,*}, Paulina Średnicka^b, Jakub P. Piwowarski^c, Marek Roszko^b, Sebastian Granica^c, Emmy Tuenter^a, Nina Hermans^a, Luc Pieters^a

^a Natural Products and Food Research and Analysis – Pharmaceutical Technology (NatuRAPT), Department of Pharmaceutical Sciences, University of Antwerp, Wilrijk, Belgium

^b Wacław Dąbrowski Institute of Agricultural and Food Biotechnology-State Research Institute, Warsaw, Poland

^c Microbiota Lab, Department of Pharmaceutical Biology, Faculty of Pharmacy, Medical University of Warsaw, Warsaw, Poland

The most extensively explored of ellagitannins' postbiotic metabolites is urolithin A (Ryu et al., 2016). However, not every individual is able to have it produced in the gut, and certain urolithin metabolotypes were distinguished (García-Villalba et al., 2022). This might be associated with the occurrence of several already isolated species producing urolithins (Iglesias-Aguirre et al., 2023). Therefore, in subsequent studies, the metabolotypes need to be considered as the main variable. Nevertheless, other factors affecting gut microbiota, such as antibiotic treatment, might influence urolithin production *in vivo* as well. Thus, the authors performed a series of incubations with gut microbiota *ex vivo*, using fecal samples from donors of different metabolotypes. Substrates such as *Punica granatum* juice, corilagin, ellagic acid, and urolithins M7 and C were used. The influence of three commonly used antibiotics, i.e., amoxicillin-clavulanic acid, cefuroxime, and azithromycin, was tested. The samples collected during incubations were analyzed using LC-HRMS. Using an intermediate product - urolithin C, as substrate allowed detectable concentrations of bioavailable urolithins to be obtained more quickly than in incubations with other substrates. High variability of the biotransformation outputs (product ratio) was observed among metabolotype B donors. The addition of amoxicillin-clavulanic acid inhibited urolithin production in the batch, while azithromycin and cefuroxime showed less impact. The incubation of metabolotype B donor microbiota with azithromycin at 25 µg/mL resulted in a biotransformation output typical to metabolotype A (only urolithin A).

Acknowledgments: The presented research was financially supported by the Polish National Science Centre grant Miniatura 6 No. 2022/06/X/NZ7/01559 and by the European Union Horizon Europe MSCA-PF grant No. 101106272.

Keywords: ellagitannins, urolithins, biotransformation, antibiotics, metabolotypes

References

- García-Villalba, R., et al., 2022. Urolithins: a comprehensive update on their metabolism, bioactivity, and associated gut microbiota. *Mol. Nutr. Food Res.* 2101019.
- Iglesias-Aguirre, C.E., et al., 2023. Gut Bacteria Involved in Ellagic Acid Metabolism To Yield Human Urolithin Metabolotypes Revealed. *J. Agric. Food Chem.* 71, 4029–4035.
- Ryu, D., Mouchiroud, et al., 2016. Urolithin A induces mitophagy and prolongs lifespan in *C. elegans* and increases muscle function in rodents. *Nat. Med.* 22, 879–888.

* Presenting author. E-mail address: dominik.popowski@uantwerpen.be (Dominik Popowski)

S1.P304 Identification of gene cluster for biosynthesis of antiviral polyketide limocrocin

Sofia Melnyk^{a,b,*}, Marc Stierhof^c, Dmytro Bratiichuk^c, Yuriy Rebets^b, Andriy Luzhetskyy^c, Bohdan Ostash^a

^a Department of Genetics and Biotechnology, Ivan Franko National University of Lviv, Hrushevskoho st. 4, Lviv 79005, Ukraine

^b Explogen LLC, Zelena st. 20, Lviv 79005, Ukraine

^c Helmholtz Institute for Pharmaceutical Research Saarland (HIPS), Saarland University, University Campus Bldg C2.3, Saarbruecken 66123, Germany

Explosion of bacterial genome data has led to discovery of myriads of biosynthetic gene clusters (BGCs) for specialized metabolites of unknown chemical identity. There is urgent need for generic approach to elucidate such (often cryptic) BGCs. Over the last decade we focused on assessing various tools and approaches which ultimately would help us decipher the natural products hidden behind cryptic BGCs through heterologous expression. Here we report the use of *Streptomyces albidoflavus* J1074 as a platform to discover the BGC for polyketide limocrocin (LIM). LIM is an unusual linear polyene that interferes with viral reverse transcriptase that was discovered back in the 1950s (Brockmann et al., 1953; Hanajima et al., 1985). However, the genetic and biochemical logic of LIM biosynthesis remained unknown until this report. In course of the studies of *S. roseochromogenes* NRRL 3504 (Melnyk et al., 2022) our attention was attracted to exotic manumycin-like BGC within its genome. We generated NRRL 3504 cosmid library, and, through end sequencing, identified the one carrying aforementioned BGC. Its introduction into *S. albidoflavus* Del14 led to accumulation of yellow-pigmented compound. LC-MS and NMR results showed that the compound is LIM 1. Two oxidized derivatives of 1 were also identified for the first time. Detailed bioinformatic analysis of *lim* BGC as well as *lim* gene knockouts led to the initial hypothesis about biosynthetic pathway leading to 1. Further experiments are necessary to understand several obscure stages of LIM assembly to enable the overproduction and genetic engineering of LIM derivatives.

Keywords: *Streptomyces roseochromogenes* NRRL 3504, BGC, heterologous expression, limocrocin, reverse transcriptase inhibitor

References

- Brockmann, H., Grothe, G. 1953. Über Actinomycetenfarbstoffe, II. Mitteil: Limocrocin, ein gelber Actinomycetenfarbstoff. Chem. Berich. 86, 1110–1115.
- Hanajima, S., Ishimaru, K., Sakano, K., Roy, S.K., Inouye, Y., Nakamura, S. 1985. Inhibition of reverse transcriptase by limocrocin. J Antibiot. 38, 803–805.
- Melnyk, S., Stepanyshyn, A., Yushchuk, O., Mandler, M., Ostash, I., Koshla, O., Fedorenko, V., Kahne, D., Ostash B. 2022. Genetic approaches to improve clorobiocin production in *Streptomyces roseochromogenes* NRRL 3504. Appl Microbiol Biotechnol. 106, 1543–1556.

* Presenting author. E-mail address: Sofia.Melnyk@lnu.edu.ua (Sofia Melnyk)

S1.P305 Reconstitution of natural products biosynthetic pathways using cell-free synthetic biology

Deepa D Acharya^{a,*}, Krishna Madduri^a, Jeremy McFadden^a, Jade Bing^b, Paul Ketterer^b, Babu Raman^a

^a *Biologicals Discovery and Development, Corteva Agriscience, 9330 Zionsville Rd, Indianapolis, IN 46268*

^b *Crop Protection Discovery and Development, Corteva Agriscience, 9330 Zionsville Rd, Indianapolis, IN 46268*

We report the successful cell-free reconstitution of two natural product biosynthetic pathways of divergent complexity and structural classes – teleocidin and UK-2A. To enable the construction of the teleocidin biosynthetic pathway, we discovered a direct interaction between a known pathway enzyme TleA and an uncharacterized enzyme MbtH. We also illustrated the regulatory bottleneck of an intermediate step catalyzed by TleB. We then extended our methodology to successfully reconstitute UK-2 diol, the precursor to a commercially valuable secondary metabolite. The cell-free system enabled refactoring of a complex pathway of ten enzymes to build UK-2 diol from simple building blocks. We show that our cell-free biosynthesis platform is suitable for reconstructing pathways and identifying the functions of uncharacterized genes linked to biosynthetic gene clusters (BGCs) and rate-limiting biosynthetic steps.

Keywords: cell-free synthesis, biosynthesis, BGC, teleocidin, UK-2A

References

Awakawa, T. Enzymatic Reactions in Teleocidin B Biosynthesis. *Journal of Natural Medicines* 2021, 75 (3), 467–474.

Buntru, M.; Vogel, S.; Spiegel, H.; Schillberg, S. Tobacco BY-2 Cell-Free Lysate: An Alternative and Highly-Productive Plant-Based in Vitro Translation System. *BMC Biotechnology* 2014, 14 (1).

Madduri, K.; Armstrong, J.; Etter, A.; Buntru, M.; Vogel, S.; Spiegel, H.; Schillberg, S; Fischer, R. Eukaryotic cell-free protein expression system that does not require an artificial energy regeneration system. 2020 U.S. Patent No. 10,612,031.

* Presenting author. E-mail address: deepa.acharya@corteva.com

S1.P306 Biotransformation of phlorotannins from brown algae to enhance bioactivity

Lea Karlsberger^{a,b,*}, Nadiia Sadova^{a,b}, Stephanie Ladirat^c, Julian Weghuber^{a,b}

^a FFoQSI GmbH – Austrian Competence Centre for Feed and Food Quality, Safety and Innovation, Technopark 1D, 3430 Tulln, Austria

^b University of Applied Sciences Upper Austria, Centre of Excellence for Food Technology and Nutrition, Stelzhamerstraße 23, 4600 Wels, Austria

^c NUQO©, 13 Rue du Tertre, 35170 Bruz, France

The EU ban on antimicrobial growth promoters (AGPs) in animal production prompted a quest for plant-based alternatives (Yang et al., 2015). Several plant-derived compounds (“phytochemicals”) have been shown to support animal health. Marine plants like brown algae rich in phlorotannins possess e.g. antioxidative, and anti-inflammatory properties (Ford et al., 2019). However, the high molecular weight of phlorotannins limits their resorption in the intestine (Catarino et al., 2021), reducing their health-promoting effects. This poster will cover the biotransformation of brown algae by fermentation and chemical hydrolysis. The disruption of phlorotannin’s chemical bonds is hypothesized to yield smaller, more bioavailable oligomers, potentially increasing bioactivity. *In vitro* experiments using extracts from two brown algae species, *Ascophyllum nodosum* and *Fucus vesiculosus* provided by NUQO©, demonstrated their bioactive effects. Intracellular antioxidant activity was shown by reducing reactive oxygen species (ROS) in oxidative stressed Caco-2 cells. Treatment with the algae extracts led to significantly reduced nitric oxide (NO) production in LPS- stressed RAW264.7 cells, indicating anti-inflammatory effects. Moreover, treatment with *A. nodosum* and *F. vesiculosus* extracts led to reduced levels of pro-inflammatory cytokines IL-6, IL-8, IL-10, TNF- α in cell culture supernatants of LPS-activated THP-1 macrophages. Preliminary findings suggest that fermentation with lactic acid bacteria (LAB) and NaOH hydrolysis of *A. nodosum* enhance its antioxidant effects in oxidative-stressed Caco-2 cells. Further research is needed to elucidate the molecular mechanisms underlying this augmented bioactivity.

Keywords: phlorotannins, brown algae, bioactivity, biotransformation, fermentation

References

- Catarino, Marcelo D.; Silva, Artur M. S.; Cardoso, Susana M., 2017. Fucaceae: A Source of Bioactive Phlorotannins. In International Journal of Molecular Sciences 18 (6).
- Ford, Lauren; Theodoridou, Katerina; Sheldrake, Gary N.; Walsh, Pamela J., 2019. A critical review of analytical methods used for the chemical characterisation and quantification of phlorotannin compounds in brown seaweeds. In Phytochemical Analysis 30, 587–599.
- Yang, Chengbo; Chowdhury, M. A. Kabir; Huo, Yongqing; Gong, Joshua, 2015. Phytogetic compounds as alternatives to in-feed antibiotics: potentials and challenges in application. Pathogens (Basel, Switzerland) 4, 137–156.

* Presenting author. E-mail address: lea.karlsberger@fh-wels.at (Lea Karlsberger)

S1.P307 Enhancing polyphenol profiles and production: a novel approach through synergistic non symbiotic fungi and organic fertilization

Daniel Raus de Baviera^{a,*}, Enrique Barraón-Catalán^{b,c}, Antonio Ruiz-Canales^a

^a Department of Engineering, Area of Agroforestry, Miguel Hernández University, 03312 Orihuela, Spain

^b Institute for Research, Development and Innovation in Health Biotechnology, Miguel Hernández University, 03202 Elche, Spain

^c Department of Pharmacy, Elche University Hospital-FISABIO, 03203 Elche, Spain

Polyphenols, as naturally occurring high-value secondary metabolites, possess well-documented benefits for human health, catalyzing an augmented demand for these compounds. Attempts to economically synthesize polyphenols by alternative methods face industrial challenges, such as low yields and toxicity affecting cell growth (Ofosu et al., 2020). In this context, Mediterranean semideciduous shrubs, thriving under abiotic stress (thermal and solar radiation) without hydric stress, emerge as a potent source of pharmacologically relevant metabolites. *Cistus albidus* L. is such a species, traditionally recognized for its medicinal properties attributable to its prolific production of diverse natural metabolites under harsh Mediterranean climate conditions, with minimal resource requirements (Raus de Baviera et al., 2023). Through the implementation of a treatment strategy that combines fertilization with non-symbiotic inoculation with *Rhizophagus* sp., we have successfully augmented the production of several polyphenols of pharmacological significance. The annual average efficiency of leaves, quantified by the specific leaf area (SLA), exhibited a 57% enhancement relative to control plants, with a twofold increase observed in the spring. This enhancement was primarily manifested in the total phenolic content and antioxidant capacity, both of which saw increments exceeding 30%. Notably, the inoculation and organic fertilization regimen induced significant alterations in the compound profile, varying with the season, and predominantly affecting flavanols, flavonols, ellagitannins, and phenolic acids. It is proposed that the synergistic application of non-symbiotic fungi and adapted organic fertilization presents a novel strategy to modify the compound profile of *C. albidus*, thereby significantly elevating the production of targeted compounds.

Keywords: polyphenol production, *Cistus albidus*, non-symbiotic fungi, organic fertilization

References

Ofosu, F.K., Daliri, E.B.-M., Elahi, F., Chelliah, R., Lee, B.-H., & Oh, D.-H., 2020. New Insights on the Use of Polyphenols as Natural Preservatives and Their Emerging Safety Concerns. *Front. Sustain. Food Syst.* 4, 525810, 1-14. Raus de Baviera, D., Ruiz-Canales, A., & Barraón-Catalán, E., 2023. *Cistus albidus* L.-Review of a Traditional Mediterranean Medicinal Plant with Pharmacological Potential. *Plants*. 12, 2988, 1-43.

* Presenting author. E-mail address: daniel.raus@goumh.umh.es (Daniel Raus de Baviera)

S1.P308 The potential role of aqueous extract from fruits of *Chaenomeles japonica* and its post-digestive fractions in inflammation-related intercellular communication

Agata J. Olędzka^{a,b}, Aleksandra Sirak^c, Dariia Hovtvian^d, Maciej Matuszewski^c, Oleh Koshovyi^d, Monika E. Czerwińska^{a,b,*}

^a Department of Biochemistry and Pharmacogenomics, Faculty of Pharmacy, Medical University of Warsaw, 1 Banacha Str., 02-097 Warsaw, Poland

^b Centre for Preclinical Research, Medical University of Warsaw, 1B Banacha Str., 02-097 Warsaw, Poland

^c Student Scientific Association, Department of Biochemistry and Pharmacogenomics, Faculty of Pharmacy, Medical University of Warsaw, 1 Banacha Str., 02-097 Warsaw, Poland

^d Department of Pharmacognosy, National University of Pharmacy, 53 Pushkinska Str., 61002 Kharkiv, Ukraine

Chaenomeles japonica (Thunb.) Lindl. ex Spach (CJ), commonly cultivated in Poland, have been also used in traditional Chinese Medicine. The fruits of CJ are rich in polyphenol compounds, especially procyanidins (Du et al., 2013; Siegien et al., 2021). The study aimed to assess the role of phytochemicals present in CJ fruit extract in intercellular communication associated with inflammation. We evaluated the effect of CJ extract on cytokine secretion in peripheral blood mononuclear cells (PBMC) and human neutrophils (PMN). To assess potential bioavailable metabolites, we performed gastrointestinal digestion (GI) *in vitro* of said extract. We compared the activity of crude fruit extract and obtained GI fractions in the human adenocarcinoma cell line (Caco-2). Finally, we evaluated the effect of the GI fractions on the neutrophil extracellular traps (NETs) secretion in PMN. The aqueous extract from CJ fruit inhibited the secretion of all pro-inflammatory cytokines (TNF- α , interleukin(IL)-8, IL-6, IL-1 β) in PMN, PBMC, and Caco-2 cells and increased the concentration of anti-inflammatory IL-10 in PBMC. The release of TNF- α in PMN was 62.1% for 100 μ g/mL of CJ fruit extract, compared to samples treated with LPS. Each fraction obtained after digestion *in vitro* decreased significantly the release of IL-8 in Caco-2 cells. There are noticeable differences in the NETs secretion among the used GI fractions of CJ. In conclusion, both aqueous extract from the fruit of *C. japonica* and post-digestive fractions present inhibitory activity of cytokine secretion. The gastrointestinal digestion of the extract likely affects the composition and bioactivity of the fractions.

Keywords: *Chaenomeles japonica*, Rosaceae, cytokines, metabolites, procyanidins

References

Du, H., Wu, J., Li, H., Zhong, P.X., Xu, Y.J., Li, C.H., Ji, K.X., Wang, L.S., 2013. Polyphenols and triterpenes from *Chaenomeles* fruits: chemical analysis and antioxidant activities assessment. Food Chem 141, 4260-4268.

Siegien, J., Buchholz, T., Popowski, D., Granica, S., Osinska, E., Melzig, M.F., Czerwinska, M.E., 2021. Pancreatic lipase and alpha-amylase inhibitory activity of extracts from selected plant materials after gastrointestinal digestion *in vitro*. Food Chem 355, 129414.

* Presenting author. E-mail address: monika.czerwinska@wum.edu.pl (Monika Czerwińska)

S1.P309 *Phlegmariurus carinatus* *in vitro* culture and study on *Lycopodium* alkaloid biosynthesis

Amit Jaisi^{a,*}, Youli Xiao^b

^a School of Pharmacy, Walailak University, Nakhon Si Thammarat, Thailand

^b CAS Key Laboratory of Synthetic Biology, CAS Center for Excellence in Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Chinese Academy of Sciences, Shanghai, China

Phlegmariurus carinatus (Desv. ex Poir.) Ching plant of Lycopodiaceae family is a rich source of neuroactive 'huperzine A' and other Lycopodium alkaloids (LAs) and gained significant interest despite challenges such as low natural content, slow plant growth, and limited success in cultivation, *in vitro* propagation, and chemical synthesis (Xu et al., 2018). Recent biosynthetic studies have identified key genes involved (Fig. 1a; Nett et al., 2021 and 2023). As a part of our ongoing research on piperidine derived-alkaloid biosynthesis (Xu et al., 2018; Fan et al., 2021). We explored LAs biosynthesis in *in vitro* grown sporophytes of 9 plants in the Lycopodiaceae, including *P. carinatus*. It was observed that optimal sterilization techniques were essential for axenic culture establishment. Moreover, we saw slow growth, but sporophyte bifurcate within 4-6 months of *in vitro* culture. The LCMS/MS analysis of LAs showed varying concentration across sporophytes (Fig. 1b). Huperzine A was highest in the leaves. Furthermore, the transcriptome analysis showed that the early biosynthetic genes upregulated in the young tissues of sporophyte, while the late biosynthetic genes especially the modification step seems to be upregulated in specific tissues like roots and leaves (Fig. 1c). In conclusion, this study provides a new platform for neuroactive LA's production and biosynthetic studies.




Fig. 1. Huperzine A biosynthetic pathway (a), lycopodium production (b) and differentially expressed candidate genes of *in vitro* culture (c).

Keywords: *Phlegmariurus carinatus*, huperzine A, sporophyte culture, transcriptome

References

- Xu, B., Fan, Z., Lei, Y., Ping, Y., Jaisi, A. and Xiao, Y., 2018. Insights into pipecolic acid biosynthesis in *Huperzia serrata*. *Org. Lett.* 20, 2195-2198.
- Nett, R.S., Dho, Y., Low, Y.Y. and Sattely, E.S., 2021. A metabolic regulon reveals early and late acting enzymes in neuroactive *Lycopodium* alkaloid biosynthesis. *Proc. Natl. Acad. Sci. USA* 118, e2102949118.
- Nett, R.S., Dho, Y., Tsai, C., Passow, D., Martinez Grundman, J., Low, Y.Y. and Sattely, E.S., 2023. Plant carbonic anhydrase-like enzymes in neuroactive alkaloid biosynthesis. *Nature* 624, 182-191.
- Fan, Z., Jaisi, A., Chen, Y., Shen, L., Liu, Z., Wu, S., Liu, Y., Zhang, W. and Xiao, Y., 2021. Discovery and biosynthesis of ascorbylated *Securinega* alkaloids. *ACS Catal.* 11, 8818-8828.

* Presenting author. E-mail address: amit.ja@mail.wu.ac.th (Amit Jaisi)

S1.P310 Biosynthesis of α -aminopyrones produced by the marine-derived fungus *Aspergillus* sp. DLM38

Lamonielli F. Michaliski^a, Laura P. Ióca^a, Taicia P. Fill^b, Antonio G. Ferreira^c, Roberto G. S. Berlinck^{a,*}

^a Instituto de Química de São Carlos, Universidade de São Paulo, CP 780, CEP 13560-970, São Carlos, SP, Brazil

^b Instituto de Química-Departamento de Química Orgânica, Universidade Estadual de Campinas-UNICAMP, Barão Geraldo, Campinas, Brazil

^c Departamento de Química, Universidade Federal de São Carlos, São Carlos, SP, Brazil

α -Aminopyrones are metabolites produced in culture exclusively by *Aspergillus* spp. with cytotoxic and antiviral activity (Barnes et al., 1990; Varoglu et al., 2000; Zhang et al., 2010). Using a combination of stable-isotope feeding and gene knock-out experiments, we investigated the biosynthesis of the α -aminopyrones isopyrophen and naphthoquinoneimine, produced by *Aspergillus* sp. DLM38 (Ióca et al., 2016). ECD analysis confirmed the absolute configuration of naphthoquinoneimine. Feeding experiments with [1-¹³C]-D-glucose unveiled the biogenetic pathways of both compounds as organized by NRPS-PKS gene cluster, reinforced by genome mining analysis. To confirm the bioinformatics results, knock-out mutants were obtained by homologous recombination, followed by comparison of the chemical profiles produced by the wild-type and mutant strains. UPLC-HRMS/MS confirmed that α -aminopyrones production was abolished in the knock-out mutant strains.

Financial Support: LFM, LPI and RGSB thank the financial support by FAPESP procs. 2013/50228-8, 2016/05133-7, 2017/25100-9, 2019/07894-3, 2019/17721-9 and 2022/07831-4.

Keywords: aminopyrones, *Aspergillus*, genome mining, biosynthesis, gene knock-out

References

- Barnes, C. L. et al., 1990. Structure and absolute configuration of pyrophen, a novel pyrone derivative of L-phenylalanine from *Aspergillus niger*. Int. J. Pept. Protein Res. 36, 292.
- Varoglu, M. et al., 2000. Biosynthetically Diverse Compounds from a Saltwater Culture of Sponge-Derived *Aspergillus niger*. J. Nat. Prod., 63, 41
- Zhang, Y. et al., 2010. Phenethyl- α -pyrone derivatives and cyclodipeptides from a marine algous endophytic fungus *Aspergillus niger* EN-13. Nat. Prod. Res., 24, 1036.
- Ióca, L. P. et al., 2016. A strategy for the rapid identification of fungal metabolites and the discovery of the antiviral activity of pyrenocine a and harzianopyridone. Quím. Nova, 39, 720-731.

* Presenting author. E-mail address: rgsberlinck@iqsc.usp.br (Roberto G. S. Berlinck)

S1.P311 Plant tissue culture for production of bioactive metabolites in *Agrostemma githago* L.

Weronika Kozłowska^a, Adam Matkowski^{b,c,*}, Daria Zblewska^d, Maria Malicka^a, Weronika Korzeniowska^b, Fabian Bülow^e, Aleksander Weng^e, Sylwia Zielińska^a

^a Department of Pharmaceutical Biotechnology, Wrocław Medical University, Borowska 211, Wrocław, Poland

^b Department of Pharmaceutical Biology and Botany, Wrocław Medical University, Borowska 211A, Wrocław, Poland

^c Botanical Garden of Medicinal Plants, Wrocław Medical University, Al. Jana Kochanowskiego 14, 50367 Wrocław, Poland

^d Student Scientific Organization, Circle No. 76, Department of Pharmaceutical Biology and Biotechnology, Wrocław Medical University, Borowska 211, Wrocław, Poland

^e Department of Pharmaceutical Biology, Freie Universität Berlin, Königin-Luise-Straße, Berlin- Dahlem, Germany

Flavone-C-glycosides (FCG) are pharmacologically active plant natural compounds with vitexin and orientin as the most studied examples found in several botanical families. One of these, distinguished by a wide diversity of flavone-C-glycosides, is the Caryophyllaceae (pink or carnation family) which comprises numerous species but only a few have been used medicinally, mainly due to their high content of saponins (triterpenoid glycosides – TTG) as bioactive principle or as an industrial material with surfactant properties (Jakimiuk et al. 2022). However, the regulation of FCG biosynthesis and reasons of the phytochemical profiles diversity remain largely unresolved despite pertinently suggested role of environmental factors in various species/plant systems. Here, we used plant cell, tissue and organ cultures as models to find out if FCGs and TTGs can be biosynthesized in a highly controlled environment in cells at different levels of differentiation – callus culture, cell suspensions, regenerated organs during *in vitro* stimulated development. *Agrostemma githago* L. *in vitro* cultures were established in solid and liquid media. The content of specialized metabolites was analyzed using LC-qTOF-MS and UV-HPLC and compared to plants grown conventionally. As a result, the FCGs orientin, isoorientin and vitexin were detected with at least five derivatives with higher molecular weight supposedly with additional glycosylation. The content of TTG in all tested *in vitro* cultures was significantly lower and less complex than in field-grown plants. However, both content and HPLC profile of both compound classes differed between culture systems, such as callus culture on agar medium, cell suspension, organ cultures as well as between solid and liquid media.

Acknowledgements. This research is funded by National Science Center of Poland (NCN) # 2020/39/I/NZ7/01515 (RIPSAPO).

Keywords: *Agrostemma githago*, Caryophyllaceae, *in vitro* cultures, orientin

Reference

Jakimiuk, K., Wink, M., Tomczyk, M., 2022. Flavonoids of the Caryophyllaceae. *Phytochem. Rev.* 21, 179–218.

* Presenting author. E-mail address: pharmaceutical.biology@wp.eu (Adam Matkowski)

S1.P312 Analysis of host and gut microbial biotransformation products of kratom (*Mitragyna speciosa*) by mass spectrometry

William J. Crandall^{a,*}, Ken H. Liu^b, Choon M. Lee^c, Grant Singer^a, Jaclyn Weinberg^a, Edward T. Morgan^d, Dean P. Jones^c, Cassandra L. Quave^e

^a Molecular and Systems Pharmacology, Emory University, GA, USA

^b Department of Chemistry, Emory University, GA, USA

^c Division of Pulmonary, Allergy, Critical Care and Sleep Medicine, Emory University, GA, USA

^d Department of Pharmacology, Emory University, GA, USA

^e Department of Dermatology, Emory University, GA, USA

Despite the widespread use of botanical medicines and supplements, the metabolic fates of bioactive and other chemical constituents of these complex mixtures are without systematic characterization. This limits our understanding of the contributions of individual components to overall efficacy or toxicity. A main contributor to the biotransformation of xenobiotics in the body is the various drug- metabolizing enzymes present in the liver. Additionally, there has been significant focus on the role the gut microbiome plays in the pharmacokinetics and efficacy of drugs (Pant et al., 2023; Zimmermann et al., 2019). However, there is a severe lack of research regarding the poly- pharmacokinetics of botanical medicines in this regard. The vast metabolic repertoire of these microbes allows for entirely unique enzymatic reactions that contribute significantly to the metabolism of xenobiotics (Pant et al., 2023). In this study, pooled human liver S9 enzymatic fractions were used for *in vitro* biotransformation of pure compounds from Kratom (*Mitragyna speciosa* Korth., Rubiaceae) and used to test capabilities for identification in the Kratom extract. Additionally, fecal matter from healthy human donors was used for the *in vitro* culturing of gut microbiome communities under anaerobic conditions. Samples were analyzed by LC-MS/MS and data were analyzed with custom scripts in Python. An integrated workflow to generate and detect multiple phase-I and -II liver, and various gut microbial metabolites from a botanical mixture was established. Applications of molecular networking and various metabolomics tools will expand capabilities for the discovery of new metabolites of pharmacological interest from complex botanical mixtures.

Keywords: kratom, liver, gut microbiome, biotransformation, mass spectrometry

References

Pant, A., Maiti, T.K., Mahajan, D., Das, B., 2023. Human Gut Microbiota and Drug Metabolism. *Microb. Ecol.* 86, 97- 111.

Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R., Goodman, A.L. 2019. Mapping human microbiome drug metabolism by gut bacteria and their genes. *Nature* 570, 462-467.

* Presenting author. E-mail address: wjcrand@emory.edu (William J. Crandall)

S1.P313 Interactions with gut microbiota and bioavailability of natural products contained in lavender flowers

Diana Dolzhko^{a,*}, Aleksandra Kruk^a, Maciej Korczak^a, Dominik Popowski^{a,b}, Jakub Piwowarski^a, Sebastian Granica^a

^a Department of Pharmaceutical Biology, Medical University of Warsaw, 1 Banacha St., 02-097, Warsaw, Poland

^b Wacław Dąbrowski Institute of Agricultural and Food Biotechnology-State Research Institute, Warsaw, Poland

Lavandula angustifolia Mill., also called true/English lavender, is a source of important medicinal plant material - lavender flower (*Lavandulae flos*) (Ghadim, et al. 2020). It's essential oil fraction is rich in monoterpenoids, but also, the flowers are often used in the form of self-prepared infusions to treat mild symptoms of anxiety. However, the form of an infusion usually contains small amounts of essential oil and contains a variety of polar natural products, which can contribute to the beneficial effects *in vivo* (Bazrafshan, et al. 2020). Previous studies shown that various plant materials can affect the biodiversity of gut microbiota (GM) and the GM can modify the structure of natural products contained in plant extract. This lead to the production of novel bioactive metabolites (Morris, et al. 2018). The current proposal aims to provide scientific evidence on the interaction of lavender infusion (LOI) with GM in the context of the treatment and prevention of anxiety disorders in humans. Forty three compounds of LOI were detected by UHPLC-DAD-MS including flavonoid derivatives, caffeic acid derivatives, propanoic acid derivatives, ferulic acid derivatives, glucosyl hydroxycinnamic acid and methoxycinnamic acid derivatives. The chemical composition after incubation with GM was established by the UHPLC-DAD-MS analysis.




Fig. 1. Metabolites of GM and LOI.

The analysis of the GM metabolism of LOI showed at which timepoint the most diverse mixtures of metabolites are produced and which compounds undergo biotransformation (Fig.1). Up to know 9 potential metabolites were detected in analysed mixtures.

Keywords: *Lavandula officinalis*, water extract, gut microbiota, postbiotic metabolites

References

- Bazrafshan, M. R., Jokar, M. Shokrpour, N. Delam., 2020. The effect of lavender herbal tea on the anxiety and depression of the elderly: A randomized clinical trial. *Complementary Therapies in Medicine*, vol. 50.
- Ghadim, M. B. A., Neisy A., Sisakht M., Khoshdel Z., 2020. *Lavandula angustifolia* aqueous extract ameliorates anxiety and depressive-like behaviors in chronic mild stress-treated male rats. *Journal of Medicinal Plants Research* 14, 593-603.
- Morris, G., Fernandes B. S., Puri B. K., Walker A. J., Carvalho A. F. Berk M., 2018. Leaky brain in neurological and psychiatric disorders: Drivers and consequences. *Australian and New Zealand Journal of Psychiatry* 52, 924-948.

* Presenting author. E-mail address: dianadolzhko@gmail.com (Diana Dolzhko)

S1.P314 Using banana peels in fermentation: upstream bioprocesses

Mariana C. Pedrosa^{a,b,c}, Sandrina A. Heleno^{a,b}, Ana Susmozas^d, Paloma Manzanares^d, Ignacio Ballesteros^d, Manuela Pintado^c, Lillian Barros^{a,b}, Marcio Carcho^{a,b,*}

^a Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Bragança, Portugal

^b Laboratório Associado para a Sustentabilidade e Tecnologia em Regiões de Montanha (SusTEC), Instituto Politécnico de Bragança, Bragança, Portugal

^c Centro de Biotecnologia e Química Fina (CBQF), Universidade Católica Portuguesa, Portugal

^d Advanced Biofuels and Bioproducts Unit, Department of Energy, Research Centre for Energy, Environment and Technology (CIEMAT), 28040 Madrid, Spain

Banana (*Musa* spp., Musaceae family) is a fruit consumed worldwide (Hikal et al., 2022). Banana peels (BP), constituting 30–40% of the fruit's weight, are considered an industrial waste. This biomass presents an opportunity for more eco-friendly solutions, namely as source for biotechnological bioprocesses. BP can be considered as a lignocellulosic biomass and because of the inherent characteristics, pre-treatment becomes essential to enhance the effectiveness of enzymes on the hydrolysis of complex carbohydrates (Mahmoud et al., 2023; Mishra et al., 2020). Therefore, this study aims to determine the best pretreatment and enzymatic hydrolysis (EH) conditions to produce a fermentation substrate suitable for producing lactic acid. A diluted-acid pretreatment was used with H₂SO₄ (0.5% and 0.25%) in autoclave for 10 min at 121 °C. After autoclaving, the pretreated material was subjected to several EH tests in which two variables were evaluated for their influence on hydrolysis yield: solid load (% dw/v), and H₂SO₄ concentration in the pretreatment (% v/v). A commercial enzyme cellulase cocktail was used for the assays. Additionally, the following test were carried out; a control test in which the BP was subjected only to autoclave pretreatment (without the subsequent EH step), and another one in which only EH was performed in raw BP. The BP showed 11.42±0.09% of cellulose, 5.9±0.3% hemicellulose, 17.2±0.5% of total lignin and 44.8±0.2% of extractives. Pretreatment was essential, as shown by lower yields in the control and EH in raw BP (Figure 1). All yields surpassed 60% after pretreatment, with H₂SO₄ impacting viscosity and enhancing fluidity during hydrolysis. Acid concentration did not significantly affect yields, but 0.25% H₂SO₄ showed a slight improvement over 0.5% H₂SO₄ for 20% solid load. The concentration of total sugars during EH remained almost constant after 12 hours, indicating minimal further hydrolysis. Therefore, according to the results, the conditions chosen for the pretreatment and EH steps were 0.25% H₂SO₄ and 15% or 20% solid load.




Figure 1 -. Graphical representation of the glucose yield from BP hydrolysis with different solid loads. EH - enzymatic hydrolysis; TT - thermal treatment; HA - acid treatment; 0.5 and 0.25 - acid concentration (% v/v); and MFE - BP medium for fermentation.

Keywords: by-products, pre-treatment, tropical fruit, hydrolysis, carbon source

References

- Hikal, W. M., Said-Al Ahl, H. A. H., Bratovic, A., Tkachenko, K. G., Sharifi-Rad, J., Kačaniová, M., Elhourri, M., & Atanassova, M., 2022. Banana Peels: A Waste Treasure for Human Being. *Evid Based Complementary Altern Med*, 2022, 7616452.
- Mahmoud, K. F., Shedeed, N. A., & Hussein, A. M. S., 2023. Production and quality evaluation of corn crackers fortified with freeze-dried banana peel and pulp. *Food and Humanity*, 1, 1680–1690.
- Mishra, R. R., Samantaray, B., Chandra Behera, B., Pradhan, B. R., & Mohapatra, S., 2020. Process optimization for conversion of Waste Banana peels to biobutanol by A yeast Co-Culture fermentation system. *Renew. Energy*, 162, 478– 488.

* Presenting author. E-mail address: mcarcho@ipb.pt (Marcio Carcho)

S1.P315 Waste to wellness: hypocholesterolemic solutions from mushroom residues, boosting circular economy

Virgínie Xavier^{a,b,c}, Custódio M. Roriz^{a,b}, Márcio Carcho^{a,b,*}, Miguel Angel Lage^c, Rita Faustino^d, Naiara Fernández^d, Ana Teresa Serra^{d,e}, Maria do Rosário Bronze^{d,e,f}, Paula Oliveira^g, Ricardo C. Calhelha^{a,b}, Sandrina A. Heleno^{a,b}

^a Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia, Bragança, Portugal

^b Laboratório para a Sustentabilidade e Tecnologia em Regiões de Montanha, Instituto Politécnico de Bragança, Campus de Santa Apolónia, Bragança, Portugal

^c Grupo de Nutrición y Bromatología, Departamento de Química Analítica y Alimentaria, Facultad de Ciencias de Ourense, Universidad de Vigo-Ourense Campus, Spain

^d iBET- Instituto de Biologia Experimental e Tecnológica, Apartado, Oeiras, Portugal

^e Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Av. da República, 2780-157 Oeiras, Portugal

^f Faculdade de Farmácia, Universidade de Lisboa, 1649-019 Lisboa, Portugal

^g Centre for the Research and Technology of Agro-Environmental and Biological Sciences (CITAB), University of Trás-Os-Montes and Alto Douro (UTAD), Vila Real, Portugal

Although statins, the primary hypocholesterolemic agents in therapeutics, are widely utilized, they are also linked with various harmful side effects. Mushrooms are rich sources of mycosterols, that reduce cholesterol absorption, polysaccharides such as β -glucans and statin-like molecules, able to lower cholesterol levels by inhibiting the 3-hydroxy-3-methyl-glutaryl CoA reductase (HMGCoA- red), a key-enzyme in the cholesterol metabolism. The Mush4Chol project aims to transform mushroom industrial biowaste into a hypocholesterolemic formulation. By exploring unsalable mushrooms and broken stems, the project seeks to develop a validated and stabilized product capable of inhibiting cholesterol synthesis and absorption. Through a combination of conventional and emerging extraction techniques, including refinement, stabilization, and encapsulation, the project aims to produce extracts rich in mycosterols, β -glucans, and statins, either as fractions or isolated compounds. The project will employ a variety of *in vitro* and *in vivo* hypocholesterolemic methodologies to explore potential mechanisms of action comprehensively. Evaluation of the probiotic *Lactobacillus reuteri*'s ability to assimilate cholesterol, both alone and in synergy with the extracts, will also be conducted. Ultimately, the goal is to create a stabilized powder comprising natural hypocholesterolemic agents, offering versatile economic applications such as gelatin capsules or as an ingredient for developing functional foods. To embrace a zero-waste approach, the leftover extraction residues will be looped back into the beginning of the value chain, serving as nutrient-rich mushroom substrate, boosting circularity.

Acknowledgements: FCT for financial support to CIMO (UIDB/00690/2020 and UIDP/00690/2020) and SusTEC (LA/P/0007/2021); FCT through the institutional scientific employment for S.A. Heleno and R.C. Calhelha, and through the Mush4Chol project (<https://doi.org/10.54499/2022.08844.PTDC>).

Keywords: hypocholesterolemic agents, mushrooms, bioresidues, functional foods, circular economy

*Presenting author. E-mail address: mcarcho@ipb.pt (Márcio Carcho)

S1.P316 Bilateral interactions between human gut microbiota and xanthohumol

Maciej Korczak^{a,*}, Dominik Popowski^{a,b}, Aleksandra Kubiszewska^a, Sebastian Granica^c, Jakub Piwowarski^a

^a Microbiota Lab, Medical University of Warsaw, Warsaw, Poland

^b Department of Food Safety and Chemical Analysis, Prof. Waclaw Dąbrowski Institute of Agricultural and Food Biotechnology – State Research Institute, Warsaw, Poland

^c Department of Pharmaceutical Biology, Medical University of Warsaw, Warsaw, Poland

Xanthohumol (XH) is the main prenylated flavonoid found exclusively in the hop plant (*Humulus lupulus*). XH has gained attention as a potent leading molecule with promising results, positioning it as a potential agent for chemoprevention, antimicrobial, or anti-inflammatory purposes (Zanoli and Zavatti, 2008). Currently, the metabolism of gut microbiota emerges as a key regulator in the bioavailability of orally administered substances, leading to the formation of postbiotic metabolites (PMs). On the other hand, xenobiotics can substantially alter the composition of the gut microbiota. To evaluate these bilateral relationships in the context of the hop active constituent, isolated from hops XH (0.5 mg/mL) was incubated with human fecal microbiota sourced from six healthy donors under anaerobic conditions. During the 24-hour period, samples were collected and analyzed using high-resolution mass spectrometry. Subsequently, intestinal bacteria were collected and underwent 16S rRNA sequencing to evaluate changes in the α - and β -diversity indices. During the *ex vivo* incubation, the formation of certain reduced XH-derived PMs was observed. Additionally, bioinformatic analysis revealed significant changes in the abundance of a few intestinal bacteria; however, there was no negative effect on the overall diversity parameters of the microbiome. These findings highlight the critical role of gut microbiota in mediating XH's bioactivity through the production of XH-derived PMs and suggest that XH, despite its antibacterial attributes, does not lead to dysbiosis or reduce the diversity of gut bacteria at the moderate concentration.

Project financially supported by Medical University of Warsaw Grant for Young Scientists No. WF4L/1/F/MB/N/23.

Keywords: gut microbiota, xanthohumol, *Humulus lupulus*

References

Zanoli, P., Zavatti, M., 2008. Pharmacognostic and pharmacological profile of *Humulus lupulus* L. J Ethnopharmacol. 116, 383-396.

* Presenting author. E-mail address: maciej.korczak@wum.edu.pl (Maciej Korczak)

S1.P317 Engineering potential of the homology pathway of L-phenylalanine and L-tyrosine

Laura E. Stewart, Skyler L. Owens, Shopno R. Ahmed, Rebecca M. Lang, Shogo Mori*

Department of Chemistry and Biochemistry, Augusta University, Georgia, USA

Homologation of amino acid is a chemical transformation to add a methylene group to the amino acid side chain. Homologated amino acids (homoAAs) are not very common non-proteinogenic amino acids found in natural products (NPs), but they are seen in several NP, NP-derived, and synthetic peptide drugs. However, its potential has not been fully utilized because of a lack of knowledge of their biosynthesis and the cost of some homoAAs. This presentation will focus on the characterization and engineering of the first novel enzyme HphA in the homologation pathway for L-Phe and L-Tyr (Fig. 1). This pathway has been identified to produce homologated L-Phe (L-hPhe) and L-Tyr (L-hTyr) by one known and three novel enzymes (Koketsu et al., 2013) and studied for a mass-production of L-hPhe by metabolic engineering (Liu et al., 2020). We biochemically characterized one of the novel enzymes HphA by investigating its proposed reaction, substrate profile, and kinetics. It was also confirmed by site-directed mutagenesis studies on its metal binding site that HphA is a homologous enzyme to LeuA that is the first enzyme of L-Leu biosynthetic pathway.




Fig. 1. Homologation of L-Phe to L-hPhe. The enzyme studied herein is colored red.

These studies and follow-up mutagenesis studies showed the engineering potential of the enzyme to homologate other amino acids that are not homologated in Nature. This study is the first step towards developing the enzymatic homologation tool of amino acids as well as peptide NPs, which has never been reported as of today.

Keywords: amino acid, biosynthesis, engineering, homologation, nonribosomal peptide

References

- Koketsu, K., Mitsuhashi, S., Tabata, K., 2013. Identification of homophenylalanine biosynthetic genes from the cyanobacterium *Nostoc punctiforme* PCC73102 and application to its microbial production by *Escherichia coli*. *Appl. Environ. Microbiol.* 79, 2201-2208.
- Liu, Z., Lei, D., Qiao, B., Li, S., Qiao, J., Zhao, G.R., 2020. Integrative Biosynthetic Gene Cluster Mining to Optimize a Metabolic Pathway to Efficiently Produce L-Homophenylalanine in *Escherichia coli*. *ACS Synth. Biol.* 9, 2943-2954.

* Presenting author. E-mail address: smori@augusta.edu (Shogo Mori)

S1.P318 Unraveling the biosynthetic pathway of aorimycin: insights from gene deletion experiments

Elena Bialas^{a,*}, Andreas Dietl^b, Nikolas Eckert^c, Andriy Luzhetskyy^c, Andreas Bechthold^a

^a Department of Pharmaceutical Biology and Biotechnology, Institute of Pharmaceutical Sciences, Albert-Ludwigs University, 79104 Freiburg, Germany

^b Department of Pharmaceutical Sciences, University of Basel, 4056 Basel, Switzerland

^c Department of Pharmacy, Pharmaceutical Biotechnology, Saarland University, Campus C2.3, 66123 Saarbrücken, Germany

Aorimycin (**1**) is a novel natural product from *Streptomyces* sp. IB2014/011-12 classified as a type II polyketide (Eckert et al., 2022), containing a unique nine-ring structure as well as an epoxy functionality (Fig. 1). Understanding its biosynthetic pathway and the enzymes involved may help to expand the enzymatic tool box for combinatorial biosynthesis, increasing the chemical diversity of bioactive compounds and thus unlocking their therapeutic potential. In this study, we aimed to elucidate the role of specific oxidoreductases in the biosynthesis of aorimycin through gene deletion experiments.




Fig. 1. Structure of aorimycin 1.

Our investigation focused on two FMN-dependent two-component monooxygenase systems, hypothesized to be pivotal in generating the chemically intriguing structures observed in aorimycin. By deleting genes encoding for these monooxygenases and flavin reductase, we aimed to discern their individual contributions to the biosynthetic pathway. Additionally, putative intermediates and shunt products, which accumulated in the deletion mutant strains, were isolated and used for structure elucidation via NMR spectroscopy and HRESIMS as well as for *in vitro* assays. Our findings reveal the significant impact of these two-component monooxygenases on aorimycin biosynthesis, shedding light on the intricate enzymatic machinery responsible for its formation. Through detailed biochemical and structural analyses, we aimed to delineate the specific functions of these enzymes, providing valuable insights into the biosynthesis of complex natural products.

Keywords: *Streptomyces*, polyketide, knockout, oxidoreductase, two-component monooxygenase

References

Eckert, N.P., Luzhetskyy, A., 2022. Heterologous overproduction of new natural products in *Streptomyces* [Doctoral dissertation, Saarland University].

* Presenting author. E-mail address: elena.bialas@pharmazie.uni-freiburg.de (Elena Bialas)

S1.P319 Synthetic biology platform for production of novel anticancer agents

Rongbin Wang^{a,*}, Heli Tirkkonen^a, Khaled A. Shaaban^b, S. Eric Nybo^c, Mikko Metsä-Ketelä^a

^a Department of Life Technologies, University of Turku, Turku, FI-20014 Finland

^b Center for Pharmaceutical Research and Innovation and Department of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, Kentucky 40536, USA

^c Department of Pharmaceutical Sciences, College of Pharmacy, Ferris State University, Big Rapids, Michigan 49307, USA

Anthracyclines are important anti-cancer agents, which are in clinical use worldwide. However, anthracyclines can cause irreversible dose-dependent cardiotoxicity, which limits the amount of anthracyclines that can be administered in the lifetime of the patient. In addition, the stereochemical complexity of anthracyclines complicates structural modification by organic synthesis. This research exploits advances in synthetic biology and built up a platform for production of novel biosynthetic anthracycline agents on an unprecedented scale, where further modifications to anthracyclines were addressed for discovery of cardiotoxicity-free anthracyclines. We firstly assembled artificial anthracycline biosynthesis pathways in *Streptomyces* with a BioBricks approach. The biosynthesis routes were split to four modules, including polyketide aglycone pathways, TDP-carbohydrate pathways, glycosyltransferases, and tailoring enzymes. With the synthetic biology platform, we are able to *de novo* refactor 4 typical anthracycline pathways. In addition, we have made more than 10 new anthracycline compounds by combinatorial synthesis. The bioactivity test will be carried on these new compounds to discover cardiotoxicity-free anticancer compounds. This project also provides an example of rapid large-scale systematic exploration to allow the discovery of functional gene combinations for generation of novel chemistry.

Keywords: biobricks, anthracycline, *Streptomyces*, biosynthesis, *de novo*

References

- Hulst, M.B., Grocholski, T., Neefjes, J.J., van Wezel, G.P. and Metsä-Ketelä, M., 2022. Anthracyclines: Biosynthesis, engineering and clinical applications. *Natural Product Reports*, 39(4), pp.814-841.
- Gomez-Escribano, J.P. and Bibb, M.J., 2011. Engineering *Streptomyces coelicolor* for heterologous expression of secondary metabolite gene clusters. *Microbial biotechnology*, 4(2), pp.207-215.
- van Dissel, D., Claessen, D., Roth, M. and van Wezel, G.P., 2015. A novel locus for mycelial aggregation forms a gateway to improved *Streptomyces* cell factories. *Microbial cell factories*, 14, pp.1-10. *Robson, N.K., 2021.*
- Wang, R., Nguyen, J., Hecht, J., Schwartz, N., Brown, K.V., Ponomareva, L.V., Niemczura, M., van Dissel, D., Van Wezel, G.P., Thorson, J.S. and Metsä-Ketelä, M., 2022. A biobricks metabolic engineering platform for the biosynthesis of anthracyclines in *Streptomyces coelicolor*. *ACS Synthetic Biology*, 11(12), pp.4193-4209.

* Presenting author. E-mail address: rowang@utu.fi (Rongbin Wang)

S1.P320 Symbiotic relation between *Ammi visnaga* and *Rhizobium rhizogenes* in bioreactor conditions

Onur Sinan Türkmen

Çanakkale Onsekiz Mart University Department of Field Crops Çanakkale, Turkey Margeht Biotechnology Ltd. Çanakkale, Turkey

Ammi visnaga is a medicinal plant known by common names such as toothpick plant, toothpick herb, bisnaga, khella (Khalil, 2020). Visnagin is an important raw material in *A. visnaga* as it has a vasodilator effect that lowers blood pressure in rats (Lee et al., 2010). In the study, visnagin transformation was monitored during the bioreaction process in order to determine the visnagin level with *Rhizobium rhizogenes* inoculation. *A. visnaga* was propagated under tissue culture conditions with Murashige and Skoog (MS) medium containing $0.5\text{mg.L}^{-1} + 0.1\text{mg.L}^{-1}$ hormones. Internodes were used to transform friable callus with MS medium containing 2.0mg.L^{-1} 2,4-D and were infected with *R. rhizogenes*. Aplikon my-control bioreactor was used to carry out the transformation conditions. Khellin levels were determined by spectrophotometric method under 390 nm (Karawya et al. 1971). In the study, no positive effect of bacterial inoculation on visnagin yield was observed.

Keywords: bioreaction, biotransformation, callus, tissue culture, medicinal plants

References

- Lee J.K., Jung J.S., Park S.H., Park S.H., Sim Y.B., Kim S.M., Ha T.S., Suh H.W., 2010. Anti-inflammatory effect of visnagin in lipopolysaccharide-stimulated BV-2 microglial cells. Arch. of Pharm. Res. 33, 1843–1850.
- Karawya M.S., Nour G., Sina A., 1971. Estimation of The Non-Glycosidal Chromone and Coumarin Content in Fruits and Extracts of *Ammi visnaga*. Planta Medica 19, 368-77.
- Khalil N, Bishr M, Desouky S, Salama O., 2020. *Ammi visnaga* L., a potential medicinal plant: A Review. Molecules. 12, 301.

* Presenting author. E-mail address: osturkmen@icloud.com (Onur Sinan Türkmen)

S1.P321 Application of 1,2,4-triazolium in structural modification of natural products and their biological activity studies

Epole Ntungwe*, Jose P. L. Roque, Claudio M. Nunes, Rui Fausto

University of Coimbra, CQC-IMS, Department of Chemistry, 3004-535 Coimbra, Portugal

Nature products have been extensively used in the discovery and development of new drugs. 1,2,3- Triazoles are widely used in the fields of pesticides and medicines due to their broad-spectrum bioactive properties. The synthesis of a new class of triazole-containing natural product conjugates has gained interest in the past few years. However, these are associated with drug resistance, giving rise to the need to develop novel drugs with similar properties. In this regard, 1,2,4-triazolium appears significant due to its antibacterial, antifungal, anticancer, and antiprotozoal activities (Song et al., 2023). Some of the triazolium compounds, such as Biapenem and Cresemba, have been approved by the U. S. Food and Drug Administration (FDA) as marketed human drugs. The introduction of the cationic triazolium structure into a drug-candidate molecule can improve the aqueous solubility, efficacy, and safety of non-target organisms. In addition, triazolium salts have been employed as *N*- heterocyclic carbene catalysts (Jia et al., 2023). Therefore, we set out to synthesize different 1,2,4- triazolium compounds using 1,4-dimethyl-4*H*-1,2,4-triazolium iodide as the precursor. The synthesized triazolium molecules were evaluated for their antioxidant, antimicrobial, general toxicity, and cytotoxicity. We thus identified which of these 1,2,4-triazolium compounds have the best biological activities and suitable for the synthesis of new lead compounds. These triazolium molecules were then conjugated to royleanones previously isolated from *Plectranthus* species to improve their aqueous solubility and cytotoxicity. We also set to study the reactivity of the synthesized triazolium compounds proving the relationship between the basicity of the triazolium and the stability of the synthesised conjugate.

Acknowledgments: This work has run within the research project PTDC/QUI-QFI/1880/2020. The CQC-IMS is supported by the Fundação para a Ciência e a Tecnologia (FCT), through projects UI0313B/QUI/2020 (DOI: 10.54499/UIDB/00313/2020), UI0313P/QUI/2020 (DOI: 10.54499/UIDP/00313/2020) and LA/P/0056/2020.

Keywords: triazolium compounds, natural products, bioactivity, synthesis

References

- Jia, S., Juan, Z., Jiamiao, J., Jie, L., Chengli, M., and Yonggui, R. C., 2023. Bioactivities of Triazolium-Based *N*- Heterocyclic Carbene Salts. *Chem. Eur. J.* 29, e202203661.
- Song, J., Lv, J., Jin, J., Jin, Z., Li, T., Wu, J., 2023. Research Advances on the Bioactivity of 1,2,3-Triazolium Salts. *Int. J. Mol. Sci.*, 24, 10694.

* Presenting author. E-mail address: epole.ntungwe@uc.pt (Epole Ntungwe)

S1.P322 Biotransformation of bufadienolides from *Drimia maritima* and *in silico* biological studies

B. Tembeni*, S. Basiru, J. O. Olatunji

African Genome Center, Mohammed VI Polytechnic University, Benguerir, 43150, Morocco

This presentation will cover the isolation and structure elucidation and biotransformation of *Drimia maritima* extracts and *in silico* anti-influenza A properties of selected compounds. *D. maritima* (Asparagaceae) is a medicinal plant with bufadienolides as major components (Saadane et al., 2021). Bufadienolides are effective against a broad spectrum of diseases and have poor solubility deeming them unsuitable for pharmacological applications (Bedir et al., 2021; Manganyi et al., 2021). In this study, different extracts of *D. maritima* bulbs were selected for microbial biotransformation studies. Microbial biotransformation is a process that makes use of microbial enzymes from fungi or bacteria as biocatalysts in the conversion of compounds to obtain derivatives with improved solubility. Different extracts (ethanol, 80% ethanol, water, and DCM: MeOH) were separately incubated with *Fusarium sp.* in PDA at 25°C for 5 days. The DCM: MeOH extract was used for the purification of bufadienolides using flash chromatography and preparative HPLC. NMR spectroscopy and detection with UHPLC-MS/MS confirmed the isolated compounds as bufadienolides. The binding capacity of bioactive compounds to H5N1 neuraminidase was comprehensively evaluated using molecular docking analysis with zanamivir as positive control.




Fig. 1. Protein ligand interactions of selected bufadienolides.

The binding energies exhibited a range from -7.24 to -8.53 kcal/mol. Specifically, resibufogenin, cinobufagin, and bufalin demonstrated markedly higher binding affinities to the 3CKZ protein than zanamivir. The specific-strong interactions identified between these compounds and the protein suggest their potential effectiveness as inhibitors of influenza A neuraminidase. These insights provide a promising avenue for the development of novel antiviral therapeutic strategies.

Keywords: biotransformation, bufadienolides, *Drimia maritima*, biological activity

References

- Badalamenti, N., Bruno, M., Pavela, R., Maggi, F., Marinelli, O., Zeppa, L., Benelli, G., Canale, A., 2022. Acaricidal Activity of Bufadienolides Isolated from *Drimia panchratum* against *Tetranychus urticae*, and Structural Elucidation of arenobufagin-3-O- α -L-rhamnopyranoside. *Plants* 11.
- Bedir, E., Karakoyun, Ç., Doğan, G., Kuru, G., Küçüksolak, M., Yusufoglu, H., 2021. New cardenolides from biotransformation of gitoxigenin by the endophytic fungus *Alternaria eureka* 1e1b11: Characterization and cytotoxic activities. *Molecules* 26.
- Manganyi, M.C., Tlatsana, G.S., Mokoroane, G.T., Senna, K.P., Mohaswa, J.F., Ntsayagae, K., Fri, J., Ateba, C.N., 2021. Bulbous plants *Drimia*: “a thin line between poisonous and healing compounds” with biological activities. *Pharmaceutics* 13, 1–20.
- Saadane, F.Z., Habbachi, W., Habbachi, S., El Imene Boublata, N., Slimani, A., Tahraoui, A., 2021. Toxic effects of *Drimia maritima* (Asparagaceae) ethanolic extracts on the mortality, development, sexual behaviour and oviposition behaviour of *Drosophila melanogaster* (Diptera: Drosophilidae). *J. Anim. Behav. Biometeorol.* 9, 1–8.

* Presenting author. E-mail address: babalwa.tembeni@um6p.ma (Babalwa Tembeni)

S1.P323 Diverse ketoreductases involved in tailoring of atypical angucyclines

Aleksi Nuutila^{a,*}, Xiansha Xiao^b, Helga van der Heul^b, Gilles van Wezel^b, Pedro Dinis^a, Somayah Elsayed^b, Mikko Metsä-Ketelä^a

^a Department of Life Technologies, University of Turku, Turku/Finland

^b Molecular Biotechnology, Leiden University, Leiden/Netherlands

Soil dwelling bacteria of *Streptomyces* genus are the best producers of chemically diverse secondary metabolites, which have been sourced for the production of majority of therapeutic antibiotics. Angucyclines are the largest class of polyketide secondary metabolites produced by *Streptomyces*, and they display a tremendous diversity in their chemical structure. Typical angucyclines are characterized by a bent four-ring structure, while atypical angucyclines have undergone cleavage of the carbon scaffold leading to drastic structural rearrangements and chemical diversification (Mikhaylov et al., 2021). In this study we focused on SDR-family ketoreductases targeting positions 1, 6, 7 and 12 (Fig. 1), encoded by lugdunomycin biosynthetic gene cluster (*lug* BGC) from *Streptomyces* sp. QL37 (Wu et al., 2019), and thioangucycline BGC (*tac*) from *Streptomyces* sp. CB00072 (Cao et al., 2021), which encode for the production of C-ring cleaved angucyclinones.



Fig. 1. SDR-family ketoreductases targeting positions 1, 6, 7 and 12.

Angucycline precursor UWM6 was converted into 8-O-methyltetrangomycin using 12-hydroxylase PgaE, 6-ketoreductase LugOIred and O-methyltransferase LugN. We elucidated the function and substrate scope of LugG, TacO and TacA encoded by *lug* and *tac* BGCs, and characterized LugG and TacO as 7-ketoreductases through NMR-analysis of their reaction product. 8-O-methyltetrangomycin was converted into SM 196 A and hydranthomycin with LugG and TacA together with previously characterized 1/6-ketoreductase LugOIred (Xiao et al., 2020). We propose that competition of many tailoring enzymes for the same substrates, and differences in enzyme promiscuity lead to a branching biosynthetic network.

Keywords: *Streptomyces*, angucycline, ketoreductase

References

- Cao, M., Zheng, C., Yang, D., Kalkreuter, E., Adhikari, A., Liu, Y.C., Rateb, M.E., Shen, B., 2021. Cryptic Sulfur Incorporation in Thioangucycline Biosynthesis. *Angew. Chemie - Int. Ed.* 60, 7140–7147.
- Mikhaylov, A.A., Ikonnikova, V.A., Solyev, P.N., 2021. Disclosing Biosynthetic Connections and Functions of Atypical Angucyclinones with a Fragmented C-Ring. *Nat. Prod. Rep.*, 38, 1506–1517.
- Wu, C., van der Heul, H.U., Melnik, A.V., Lübber, J., Dorrestein, P.C., Minnaard, A.J., Choi, Y.H., van Wezel, G.P., 2019. Lugdunomycin, an Angucycline-Derived Molecule with Unprecedented Chemical Architecture. *Angew. Chemie.* 131, 2835–2840.
- Xiao, X., Elsayed, S.S., Wu, C., Van Der Heul, H.U., Metsä-Ketelä, M., Du, C., Protá, A.E., Chen, C.C., Liu, W., Guo, R.T., Abrahams, J.P., Van Wezel, G.P., 2020. Functional and Structural Insights into a Novel Promiscuous Ketoreductase of the Lugdunomycin Biosynthetic Pathway. *ACS Chem. Biol.* 15, 2529–2538.

* Presenting author. E-mail address: atvnuu@utu.fi (Aleksi Nuutila)

S1.P324 CBD biotransformation potential of endophytic fungi from *Cannabis sativa*

Eleni Georgoula^a, Christina Koufali^{b,*}, Nikolaos Tsfantakis^b, Zacharoula Gonou-Zagou^a, Nikolas Fokialakis^b

^a Department of Ecology and Systematics, Faculty of Biology, National and Kapodistrian University of Athens, Athens 15784, Greece

^b Department of Pharmacognosy and Chemistry of Natural Products, Faculty of Pharmacy, National and Kapodistrian University of Athens, Athens 15771, Greece

Endophytic fungi refer to a group of fungi that live inside plant tissues without causing any apparent harm to the host plant and establishing a symbiotic relationship with it. Endophytic fungi are known to produce a wide variety of secondary metabolites within the tissues of their host plants, which is also the case with cannabis (Kusari et al., 2012; Scott et al., 2018). Although numerous studies suggest the biotechnological potential of endophytic fungi (Zheng et al., 2015), the ability of *Cannabis sativa* mycobiota to biotransform cannabinoids is still an unexplored field. For this purpose, ten strains of endophytic fungi, isolated from cannabis plants of the dioecious variety Carmagnola, were used in biotransformation experiments of cannabidiol (CBD). The endophytic fungus that presented the most interesting chemical biotransformation profile of CBD was *Dichotomophilus erectus*, of the Chaetomiaceae family of the phylum Ascomycota, and was subjected to large-scale liquid cultivation to isolate its main biotransformation products. In total six cannabinoids were isolated by semi- preparative high performance liquid chromatography (HPLC). Their structure elucidation was performed by 1D&2D NMR spectroscopy and by UPLC-HRESIMS. Among the compounds, three new natural products were isolated: two belonging to cannabielsoin-type cannabinoids, and one to a hydroxylated metabolite of cannabidiolaldehyde.

Keywords: cannabidiol, biotransformation, *Cannabis sativa*, endophytic fungi

References

- Kusari, P., Kusari, S., Spiteller, M., Kayser, O., 2012. Endophytic fungi harbored in *Cannabis sativa* L.: diversity and potential as biocontrol agents against host plant-specific phytopathogens. *Fungal Diver.* 60, 137–151.
- Scott, M., Rani, M., Samsatly, J., Charron, J.-B., Jabaji, S., 2018. Endophytes of industrial hemp (*Cannabis sativa* L.) cultivars: identification of culturable bacteria and fungi in leaves, petioles, and seeds. *Can. J. Microbiol.* 1–17.
- Zheng, Y.K., Qiao, X.G., Miao, C.P., Liu, K., Chen, Y.W., Xu, L.H., Zhao, L.X., 2015. Diversity, distribution and biotechnological potential of endophytic fungi. *Ann. Microbiol.* 66, 529–542.

* Presenting author. E-mail address: chrikouf@pharm.uoa.gr (Christina Koufali)

S1.P325 Structure and biosynthesis of hyellamide, a glycosylated N-acyltyrosine derivative, from the cyanobacterium *Hyella patelloides* LEGE 07179

Ângela Brito^a, Teresa P. Martins^b, Adriana Rego^b, Pedro N. Leão^{b,*}

^a Instituto de Investigação e Inovação em Saúde, University of Porto, Porto, Portugal

^b IBMC - Instituto de Biologia Molecular e Celular, University of Porto, Porto, Portugal

^c CIIMAR - Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Matosinhos, Portugal

The chemical exploration of understudied taxa continues to be a valid strategy to reveal new natural products. Genome studies have indicated that the order Pleurocapsales is one of the most biosynthetically-rich among bacteria; however, so far, a single natural product has been reported from members of this order. Here, we report the discovery, isolation and NMR-based structure elucidation of hyellamide (**1**, Fig. 1), from the cultured cyanobacterium *Hyella patelloides* LEGE 07179. Compound **1** is an N-acylated tyrosine enamide with an acetylated sugar moiety. While a few non- glycosylated related metabolites had been discovered from the heterologous expression of eDNA- derived bacterial biosynthetic genes, this is the first reported example of a glycosylated version of such scaffolds. In addition, this class of metabolites had not been associated previously with cyanobacteria. Given the novelty of **1**, and having the genome of *Hyella patelloides* LEGE 07179 in hand, we sought to understand its biosynthesis. We encountered a single candidate biosynthetic gene cluster – *hye* – which contains the necessary functions to synthesize **1**, namely an acyltransferase, a glycosyltransferase, an O-acetyltransferase, and the key oxidative decarboxylase, which we propose decarboxylates the N-acylated tyrosine to generate an N-acylated enamide.




Fig. 1. Structure of hyellamide (**1**).

The discovery **1** will propel future discovery efforts in this natural-products rich order of cyanobacteria.

Keywords: cyanobacteria, biosynthesis, structure elucidation, *Hyella*

* Presenting author. E-mail address: pleao@ciimar.up.pt (Pedro Leão)

S1.P326 Transcriptome analysis of *Triphyophyllum peltatum*, a liana producing naphthylisoquinoline alkaloids

Darshana Joshi^{a,e}, Kiran K. Telukunta^{b,e}, Philipp Richter^c, Doris Feineis^d, Gerhard Bringmann^d, Karin Fester^{c,*}

^a Department of Bioinformatics, Guru Nanak Khasla College Matunga (East), 2VG4+7J8, Nathalal Parekh Marg (Matunga East), Mumbai, Maharashtra 400019, India

^b IOCB Prague - Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo nám. 542/2, 160 00 Praha 6-Dejvice, Czech Republic

^c Faculty of Natural and Environmental Sciences, Zittau/Görlitz University of Applied Sciences, Theodor-Körner-Allee 16, 02763 Germany

^d Institute of Organic Chemistry, Faculty of Chemistry and Pharmacy, University of Würzburg, Am Hubland, 97074 Würzburg, Germany

^e Tarunavadaanenasaha Muktbharatonnayana Samstha Foundation, Secunderabad, India

Triphyophyllum peltatum (Dioncophyllaceae) is a West-African liana, which produces acetogenic naphthoquinones and naphthylisoquinoline alkaloids (Feineis et al., 2023). Especially noteworthy are the naphthylisoquinoline alkaloids due to their unique structures and their promising efficacy against protozoan infections such as malaria, and against leukemia and cancer cells (Yücer et al., 2024). For further pharmacological studies, larger quantities of naphthylisoquinoline alkaloids are required, which cannot be provided from scarce natural sources or by demanding total chemical synthesis. Hence, we aim at elucidating the biosynthesis of these intriguing compounds in order to produce them in transgenic bacteria or yeast. To obtain further insight into the biosynthesis of naphthylisoquinoline alkaloids, we analyzed the transcriptome of *T. peltatum* callus cultures. *De novo* assembly of the sequences yielded 170,415 transcripts with an average size of 1494 bp. Functional annotation of the transcripts was performed with the BLAST2GO tool, and further gene ontology terms were assigned to them. In the KEGG pathway analysis, 64,608 transcripts were assigned to 22,495 KEGG pathway maps. According to this analysis, 4705 sequences are involved in the biosynthesis of secondary metabolites. In addition, a phylogenetic analysis of transcripts putatively involved in naphthoquinone and naphthylisoquinoline alkaloid biosynthesis was performed. The results of this study will increase the knowledge on the biosynthesis of these natural products and provide the base for their metabolic engineering.

Keywords: *Triphyophyllum peltatum*, naphthylisoquinoline alkaloids, antimalarial, anticancer, transcriptome analysis

References

- Feineis, D., Bringmann, G., 2023. Asian *Ancistrocladus* Lianas as Creative Producers of Naphthylisoquinoline Alkaloids. *Progr. Chem. Org. Nat. Prod.* (Eds.: Kinghorn, A.D., Falk, H., Gibbons, S., Asakawa, Y., Liu, J.-K., Dirsch, V.M.), Volume 119, Springer, Cham, Heidelberg, New York, Dordrecht, London.
- Yücer, R., Favez, S., Feineis, D., Klauck, S.M., Shan, L., Bringmann, G., Efferth, T., Dawood, M., 2024. Cytotoxicity of dioncophylline A and related naphthylisoquinolines in leukemia cells, mediated by NF-κB inhibition, angiogenesis suppression, G2/M cell cycle arrest, and autophagy induction. *Phytomedicine* 126, 155267.

* Presenting author. E-mail address: karin.fester@hszg.de (Karin Fester)

S1.P327 The roots of *Scorzonera kotschy* Boiss. a new source for unusual dihydroisocoumarin and saponin derivatives

Özlem Bahadır-Acıkara^{a,*}, Christian Zidorn^b

^a Department of Pharmacognosy, Faculty of Pharmacy, University of Ankara, 1 Degol Str., Number 4, 06560 Ankara, Turkey

^b Pharmazeutisches Institut, Abteilung Pharmazeutische Biologie, Christian-Albrechts-Universität zu Kiel, Gutenbergstraße 76, 24118, Kiel, Germany

Scorzonera L. is a genus of nearly 180 species that belongs to the Asteraceae family. It is naturally distributed in Asia, central and southern Europe, and northern Africa. The plants of this genus are traditionally used as vegetables or animal feed, as well as in folk medicine (Lendzion et al., 2021). In Turkey, the genus *Scorzonera* is represented by 52 species and among these, 31 species are endemic to Turkey (Coşkunçelebi et al., 2015). *Scorzonera kotschy* Boiss. is a perennial species native to Turkey (Chamberlain, 1975). Current research has investigated the plant's roots for their phytochemical structure. *S. kotschy* gave three saponins that have not yet been isolated from any *Scorzonera* species one of which has an undescribed chemical structure (2 α , 3 β , 19 α , 21 β , 23- pentahydroxy urs-12-en-21-O- β -glucopyranoside) has not been isolated from any natural sources. Furthermore, three undescribed dihydroisocoumarin derivative which were identified as 4-hydroxy hydrangenol-4-O- β -glucopyranoside, 4-hydroxy-hydrangenol-4-O- β -rhamnosyl-glucopyranoside and 4-hydroxy-hydrangenol-4-O- β -apiosyl-glucopyranoside as well as one new syringic acid derivative described as syringic acid-4-O-[(E)-6-O-feruloyl]- β -glucopyranoside have also been isolated. All structures were established using spectral data, including ¹H, ¹³C and 2D NMR techniques (HMBC, HSQC, COSY) and the HR-MS.

Keywords: *Scorzonera kotschy*, Asteraceae, dihydroisocoumarin, saponin

References

- Coşkunçelebi, K., Makbul, S., Gültepe, M., Okur, S., Güzel, M. E., 2015. A conspectus of *Scorzonera* s.l. in Turkey. Turkish J. Bot. 39, 76-87.
- Davis, P.H. (Ed.) 1975. Flora of Turkey and The East Aegean Islands, Volume 5. 632-657. University Press, Edinburgh.
- Lendzion, K., Gornowicz, A., Bielawski, K., Bielawska, A., 2021. Phytochemical composition and bioactivities of *Scorzonera* species. Inter. J. Mol. Sci. 22, 5128.

* Presenting author. E-mail address: obahadir@ankara.edu.tr (Özlem Bahadır-Acıkara)

S1.P328 Genome-wide identification and characterization of two potential antiviral plant protein families (Major Latex Protein and Glycine-Rich Protein) in *Chelidonium majus*

Joanna Gracz-Bernaciak, Jakub Barylski, Oliwia Mazur, Natalia Kielich, Oskar Musidlak, Robert Nawrot*

Faculty of Biology, Institute of Experimental Biology, Department of Molecular Virology, Adam Mickiewicz University, Poznań, Poland

Chelidonium majus L., a prominent latex-producing plant from the Papaveraceae family, boasts a wide array of medicinal properties, such as antiparasitic, insecticidal, anti-neoplastic, antibacterial, and antiviral (Zielińska *et al.*, 2018). Despite long-standing use of *C. majus* extracts in traditional medicine for treating visible symptoms of human papillomavirus (HPV) infection, the precise molecular mechanisms underlying its therapeutic action remains elusive. In recent years, two proteins—major latex protein (MLP) and glycine-rich protein (GRP)—have garnered our attention as potential contributors to the antiviral and anticancer activities of latex (Nawrot *et al.*, 2017). Moreover, their interaction with alkaloids could modulate therapeutic effects. The main goal of current studies is genome-wide characterization of *Chelidonium majus* GRP and MLP gene families and their antiviral and anticancer properties. To initiate the investigation, we performed hybrid *de novo* sequencing of *C. majus* genome by combining the Nanopore technology for long-read sequencing and Illumina technology for generating numerous short reads. After genome assembly and annotation, we managed to identify 72 genes encoding 91 MLP proteins and 18 genes encoding 21 GRP proteins. Using OrthoFinder, we found 13 ortholog gene clusters for MLP and 10 clusters for GRP genes in the Papaveraceae family. Moreover, we produced and purified CmGRP and CmMLP what was followed by cytotoxicity analysis of recombinant proteins against HPV-positive and HPV-negative cells. Through rigorous genome analysis and annotation approaches, we unravel the hidden intricacies of *Chelidonium majus*, providing valuable insights into its biological functions, potential therapeutic applications, and evolutionary relationships with other species within the plant kingdom.

Acknowledgements: This research was funded by the National Science Centre, Poland (grant number 2019/35/B/NZ9/03851).

Keywords: *Chelidonium majus*, latex, major latex protein, glycine-rich protein, genome-wide analysis, sequencing

References

- Nawrot, R. *et al.* 2017. Proteomic comparison of *Chelidonium majus* L. latex in different phases of plant development, Plant physiology and biochemistry: PPB/Societe francaise de physiologie vegetale, 112, 312–325.
- Zielińska, S. *et al.* 2018. Greater Celandine's Ups and Downs-21 Centuries of Medicinal Uses of *Chelidonium majus* From the Viewpoint of Today's Pharmacology, Frontiers in Pharmacology, 9, 299.

* Presenting author. E-mail address: rnawrot@amu.edu.pl (Robert Nawrot)

Medicinal natural products: from bench to bedside

S2.P1 Stelletin B inhibits VEGF-induced angiogenesis in human endothelial progenitor cells *in vitro* and *in vivo*

Shih-Wei Wang^{a,*}, Chi Chen^b, Jyh-Horng Sheu^c

^a Institute of Biomedical Science, MacKay Medical College, New Taipei City, Taiwan

^b Department of Medicine, MacKay Medical College, New Taipei City, Taiwan

^c Department of Marine Biotechnology and Resources, National Sun Yat sen University, Kaohsiung, Taiwan

Angiogenesis plays an important role during pathological processes, including various inflammatory diseases and tumor progression. Accumulating evidence reports that bone marrow derived endothelial progenitor cells (EPCs) regulate angiogenesis. It has been suggested that understanding the molecular targets and pharmacological functions of marine derived natural products is important for novel drug discovery. Stelletin B is a marine sponge derived triterpenoid and has been found to exert anti-cancer, anti-invasion, and anti-angiogenic activity. However, the function of stelletin B upon vascular endothelial growth factor (VEGF)-induced angiogenesis on human EPCs is undetermined. In this study, we found that stelletin B suppressed VEGF-induced EPCs tube formation in a concentration dependent manner. Furthermore, stelletin B markedly abrogated VEGF-induced microvessel formation in both CAM angiogenesis and murine Matrigel implant models. Mechanistic investigations showed stelletin B inhibited VEGF-induced phosphorylation of VEGF receptor-2 (VEGFR-2) and its downstream signals, including Akt, Erk, p38 and Src in EPCs. This is the first demonstration that stelletin B impeded VEGF-induced EPCs angiogenesis through VEGFR-2 dependent pathway on human EPCs. Stelletin B is a promising marine natural product worthy of further development for the treatment of angiogenesis related diseases.

Keywords: anti-angiogenesis, endothelial progenitor cells, marine natural products

* Presenting author. E-mail address: shihwei@mmc.edu.tw (Shih-Wei Wang)

S2.P2 Antidiarrheal coumarins from *Psydrax schimperianus* (Rubiaceae)

Getahun Damissie^a, Ketema Tolossa^b, Ariaya Hymete^c, Solomon Tadesse^d

^a Department of Pharmaceutical Chemistry and Pharmacognosy, School of Pharmacy, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia; Department of Pharmacy, College of Medicine and Health Sciences, Salale University, Fitcha, Ethiopia

^b Aklilu Lemma Institute of Pathobiology, Addis Ababa University, Addis Ababa, Ethiopia

^c Department of Pharmaceutical Chemistry and Pharmacognosy, School of Pharmacy, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia

^d Department of Pharmaceutical Chemistry and Pharmacognosy, School of Pharmacy, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia; Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, Kasiska Division of Health Sciences, Idaho State University, Pocatello, ID, 83209, USA

This presentation will cover the isolation, structure elucidation, and antidiarrheal evaluation of crude extracts and two coumarins, isoscopoletin, and scoparone, from *Psydrax schimperianus* (Rubiaceae). *Psydrax schimperianus* (A. Rich.) Bridson. In Ethiopia, the root of *P. schimperianus* is traditionally used to treat diarrhea (Dalle, 2019), and its leaves are used for the treatment of stomach aches and snake bites (Asfaw *et al.*, 2021). This study investigated the *in vivo* antidiarrheal activity of an 80% methanol crude extract and coumarins isolated from the roots of *P. schimperianus* to provide a pharmacological basis for its traditional use as an antidiarrheal agent (Dalle, 2019). Phytochemical investigation of this extract led to the isolation of two known coumarins, isoscopoletin and scoparone (Fig. 1).




Fig.1. Antidiarrheal activities of isoscopoletin and scoparone.

The crude root extract of *P. schimperianus*, at doses of 100, 200, and 400 mg/kg, inhibited defecation by 37.5%, 46.2%, and 61.2%, respectively. At a dose of 20 mg/kg, scoparone and isoscopoletin reduced defecation by 61.2% and 66.6%, respectively. The finding of this study indicated the strong antidiarrheal activities of 80% methanol crude extract and the isolates, this study warrants further investigation of isoscopoletin and scoparone towards development as a novel treatment for diarrheal diseases.

Keywords: Coumarins, diarrhea, isoscopoletin, *Psydrax schimperianus*, scoparone, traditional medicine

References

Dalle, M. 2019. Ethnobotanical study of medicinal plants in Nagelle Arsi district, West Arsi Zone of Oromia, Ethiopia.

J. Nat. Sci. 9(13), 1-19.

Asfaw, A., Lulekal, E., Bekele, T., Debella, A., Abebe, A. and Degu, S. 2021. Ethnobotanical investigation on medicinal plants traditionally used against human ailments in Ensaro district, north Shewa zone, Amhara regional state, Ethiopia., 1(1), 1-27.

* Presenting author. E-mail address: getchdr@slu.edu.et (Getahun Damissie)

S2.P3 Ethnopharmacological validation of antiproliferative and anti-inflammatory activities of four *Curcuma* species through chemical characterization and *in-vitro* studies

Swagat Mohanty*, Asit Ray, Ambika Sahoo, Sudipta Jena, Pratap Chandra Panda, Sanghamitra Nayak

Centre for Biotechnology, Siksha 'O' Anusandhan (Deemed to be University), Bhubaneswar, Odisha, India

The genus *Curcuma* (Zingiberaceae) is widely used in traditional folk medicine to treat diseases caused by inflammation. Several species have remained under-explored and lack scientific validation for their claimed ethnomedicinal properties (Tushar et al., 2010). Therefore, the current study aimed to investigate the chemical composition, antiproliferative and anti-inflammatory activities of essential oil of *C. alismatifolia*, *C. aromatica*, *C. raktakanta* and *C. xanthorrhiza* rhizomes. GC-MS analysis identified a total of 90 compounds from which curzerenone (2.5-60.2%), β -curcumene (1.6-31.83%), 1,8-cineole (1.1-16.78%) and xanthorrhizol (0.6-14.12%) were the major constituents. HepG2, PC3 and MCF7 cancer cell lines were subjected for MTT assay that revealed strong *in-vitro* antiproliferative activity where *C. alismatifolia* and *C. aromatica* showed the highest selective cytotoxicity against the MCF7 and PC3 cells exhibiting an IC₅₀ value of 52.86 μ g/mL and 69.29 μ g/mL, respectively. The toxicity of volatile oils was checked on normal 3T3-L1 cells and regarded as safe for normal human health. The *in-vitro* anti-inflammatory activity was initiated by the cytotoxicity of the essential oils (6.25–200 μ g/mL) to determine the non-toxic range of further assays (Hong et al., 2021). The levels of inflammatory mediators were significantly suppressed within 100 μ g/mL of essential oil in LPS-induced RAW 264.7 macrophages. *C. xanthorrhiza* showed the greatest anti-inflammation response by significantly lowering the production of NO (87%), PGE2 (91.2%) and pro-inflammatory cytokines viz., TNF- α (95.4%), IL-6 (92%) and IL-1 β (88%) compared to the LPS treated group. Thus, the current studies showed that the *Curcuma* species used in traditional folk medicine might be a potential source of bioactive constituents exhibiting therapeutic properties.

Keywords: *Curcuma*, GC-MS, MTT assay, antiproliferative, anti-inflammatory

References

- Tushar, Basak, S., Sarma, G.C., Rangan, L. (2010). Ethnomedicinal uses of Zingiberaceous plants of Northeast India. J. Ethnopharmacol. 132(1), 286-296.
- Hong, Y., Liu, X., Wang, H., Zhang, M., Tian, M., 2021. Chemical composition, antibacterial, enzyme-inhibitory, and anti-inflammatory activities of essential oil from Hedychium puerense rhizome. Agronomy 11 (12), 2506.
- Verma, P., Kundu, R., 2020. The ginger prophecy; a review of the underexplored genus, Hedychium against Cancer. Indian J. Pharm. Sci. 82(1), pp.11-20.

* Presenting author. E-mail address: swagatrude@gmail.com (Swagat Mohanty)

S2.P4 *In vitro* and in silico study of protein tyrosine phosphatase 1B (PTP1B) inhibitory compounds from the leaves of *Cleistocalyx operculatus* (Myrtaceae)

Byeol Ryu, Jorge-Eduardo Ponce-Zea, Won Keun Oh*

Research Institute of Pharmaceutical Sciences, College of Pharmacy, Seoul National University, Seoul 08826, Republic of Korea

In our continuing study of potential antidiabetic agents from traditional medicinal plants, we have identified two novel lupane-type triterpenoids (**1**, **2**) with dicarboxylic groups and a new nor-taraxastane-type triterpenoid (**3**), along with fourteen previously known compounds (**4**–**17**), from the leaves of *Cleistocalyx operculatus* (Roxb.) Merr. et Perry (Myrtaceae). The structures of these compounds were elucidated through comprehensive spectroscopic methods, including IR, HRESIMS, 1D, and 2D NMR, and the known compounds were identified by comparing them with existing data in the scientific literature. We assessed all isolated compounds (**1**–**17**) for their capacity to inhibit the protein tyrosine phosphatase 1B (PTP1B) enzyme (Rines et al., 2016). Of these, compounds **6**, **9**, and **17** demonstrated significant inhibition of PTP1B. The inhibition mechanism of PTP1B by these compounds was explored via enzyme kinetics, revealing a non-competitive inhibition mechanism as shown by Lineweaver-Burk plots. Dixon plots were also used to ascertain the inhibition constants, aligning well with the observed IC₅₀ values. Further understanding was developed through a structure-activity relationship study and molecular docking of the compounds with the PTP1B enzyme's crystal structure (Fig. 1).




Fig. 1. Inhibitory study on compound **6**, **9**, and **17** using Lineweaver-Burk plots and molecular docking simulation on the PTP1B enzyme

Additionally, the potential of all isolates (**1**–**17**) to stimulate the uptake of 2-deoxy-2-[(7-nitro-2,1,3-benzoxadiazol-4-yl)amino]-D-glucose (2-NBDG) was investigated in differentiated 3T3-L1 adipocyte cells. Here, compounds **6**, **13**, and **17** were found to notably enhance glucose uptake in a dose-dependent manner.

Keywords: *Cleistocalyx operculatus*, Myrtaceae, protein tyrosine phosphatase 1B (PTP1B), kinetic, molecular docking, glucose uptake

References

Rines, A.K., Sharabi, K., Tavares, C.D., Puigserver, P. 2016. Targeting hepatic glucose metabolism in the treatment of type 2 diabetes, Nat. Rev. Drug Discov. 15, 786-804.

* Presenting author. E-mail address: wkoh1@snu.ac.kr. (Won Keun Oh)

S2.P5 Revival of a traditional Austrian wound healing preparation

Elisabeth Eichenauer^{a,b,*}, Johannes Saukel^a, Sabine Glasl^a

^a Division of Pharmacognosy, Department of Pharmaceutical Sciences, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^b Vienna Doctoral School of Pharmaceutical, Nutritional and Sport Sciences, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

Herbal preparations represent a promising therapy option in the treatment of acute and chronic wounds. However, with the ongoing increase in chronic wounds further therapy alternatives than the already used herbal formulations are highly appreciated (Vitale et al., 2022). Traditional folk medicine might hold a plethora of old knowledge about already forgotten wound healing plants. The VOLKSMED database was created to compile and preserve Austrian folk medicine and consists of more than 43,000 records about the traditional use of plants and fungi, but also other materials of animal or mineral origin (Saukel et al., 1994). It can be used to identify promising folkloric traditions, such as the preparation of a wound healing ointment made of lard and Norway spruce balm (*Picea abies*, Pinaceae). This tradition has already been supported by identifying the main constituents and *in vitro* wound healing activities (Göls et al, 2020), as well as *in vivo* effects on castration wounds in piglets (Prokop et al., 2023) and first case studies in patients suffering from wounds. So far, the investigations led to a monograph of a lard-based Norway spruce balm ointment in the Austrian pharmacopeia and a lard-based medical device on the Austrian market. Alternatives to lard are already actively sought and investigated to provide a more animal-friendly, stable, and standardized vehicle, which might lead to higher patient compliance (Eichenauer et al., 2023). The revival of this folkloric tradition represents an excellent example of the hidden and forgotten treasures that still can be found in Austrian folk medicine.

Keywords: Austrian folk medicine, Norway spruce balm, *Picea abies*, traditional preparations, wound healing

References

- Eichenauer, E., Jozić, M., Glasl, S., Klang, V., 2023. Spruce balm-based semisolid vehicles for wound healing: effect of excipients on rheological properties and *ex vivo* skin permeation. *Pharmaceutics* 15, 1678.
- Goels, T., Eichenauer, E., Langeder, J., Hoeller, F., Sykora, C., Tahir, A., Urban, E., Heiss, E.H., Saukel, J., Glasl, S., 2020. Norway Spruce Balm: Phytochemical composition and ability to enhance re-epithelialization *in vitro*. *Planta Med.* 86, 1080-1088.
- Prokop, D., Spargser, J., Hagmüller, W., Tichy, A., Zitterl-Eglseer, K., 2021. Efficacy of Norway spruce ointments and bacterial and fungal alterations in the treatment of castration wounds in piglets. *Planta Med.* 88, 300-312.
- Saukel, J., Kubelka, W., 1994. VOLKSMED-Datenbank - Volksmedizinisch verwendete Arzneipflanzen in Österreich. *Sci. Pharm.* 62, 2.
- Vitale, S., Colanero, S., Placidi, M., Di Emidio, G., Tatone, C., Amicarelli, F., D'Alessandro, A.M., 2022. Phytochemistry and biological activity of medicinal plants in wound healing: an overview of current research. *Molecules* 27, 3566.

* Presenting author. E-mail address: elisabeth.eichenauer@univie.ac.at (Elisabeth Eichenauer)

S2.P6 Enhancing product stability by microencapsulation of a nanoemulsion of plant extracts

Ailane S. de Freitas^a, Ilza Maria de O. Sousa^a, Luan da S. D. Rabelo^a, Pamela Bendel^a, Lorena Ribeiro^a, Mayra Biccigo^a, Nubia C.A. Queiroz^a, Paulo Roberto Carvalho^b, Ana Ruiz^a, João E. Carvalho^a, Mary Ann Foglio^{a,*}

^a Faculty of Pharmaceutical Sciences, University of Campinas, Campinas, Brazil

^b Institute of Food Technology, Campinas, Campinas, Brazil

Fridericia chica (Bonpl.) L.G. Lohmann (Syn. *Arrabidaea chica* Verlot) (SisGen A5489C9), standardized extract proved wound healing activity in Phase I and II clinical trials of volunteers with head and neck cancer. The Anthocyanins (carajurin), involved in the species pharmacological activity are unstable under environmental conditions. Hence, the microencapsulation of a nanoemulsion containing the association of the standardized extract with 5% of an enriched tocotrienol fraction. *F. chica* leaves' extraction at -5°C with 70% ethanol with 0.3% citric acid provided the standardized crude extract. Further addition of 5% tocotrienol enriched fraction (70%) with high-energy using a tip sonicator gave the nanoemulsion (Rivera-Pérez et.al. 2023). Adding hydrolyzed collagen to the nanoemulsion provided a powder by spray drying. Carajurin, δ and γ tocotrienol contents were evaluated by high liquid pressure chromatography, at 470nm and 295nm, throughout 120 days, under three temperature conditions ($25^{\circ}\text{C} \pm 2^{\circ}\text{C}$), ($5^{\circ}\text{C} \pm 2^{\circ}\text{C}$), and ($40^{\circ}\text{C} \pm 2^{\circ}\text{C}$) (Sousa et al., 2022). The use of the nanoemulsion microencapsulation strategy demonstrated the ability to preserve carajurin, δ and γ tocotrienol content throughout 120 days at room temperature, providing an economically more suitable active plant pharmaceutical ingredient for use in topical formulations for wound healing purposes.

Keywords: *Fridericia chica*, *Arrabidaea chica*, carajurin, anthocyanins, stability

References

- Rivera-Pérez, E., Escobar-Ortiz, A., Pérez-Ramírez, I.F., Regalado-González, C., Zubieta-Otero, L.F., Rodríguez- García, M.E., Reynoso Camacho, R., 2023. J. Drug Deliv. Sci. Technol. 82, 104299.
- Sousa, I.M.O., Rivera, G.A., Pinto, D., Queiroz, N.C.A., Bastos, V.I.C., Braga, L.E., Palmeira, J.D., Amaral, H., Ferreira, H., Oliveira, H., Gonçalves, F.J., Ruiz, A.L., Ibanez, E., Oliveria, B., Foglio, M.A., 2022. Nat. Prod. Res. 36, 4793- 4797.

Acknowledgements: capes, CNPQ e FAPESP grant number 2021/01280-3.

* Presenting author. E-mail address: foglioma@gmail.com (Mary Ann Foglio)

S2.P7 Homoisoflavonoids from the Hyacinthaceae for use against ocular angiogenesis

Hannah Jefford^{a,*}, Dulcie Mulholland^a, Moses Langat^b, Timothy Corson^c

^a Natural Products Research Group, Department of Chemistry, Faculty of Engineering and Physical Science, University of Surrey, Guildford, Surrey, GU2 7XH, UK

^b Jodrell Laboratory, Department of Natural Capital and Plant Health, Royal Botanic Gardens, Kew, Richmond, TW9 3DS, UK

^c Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, Ontario, Canada M5S 3M2

Natural products chemistry involves the extraction of potentially active compounds from plants. A class of active compounds, called homoisoflavonoids, are commonly extracted from the Hyacinthaceae family. They are regarded as promising pharmacological candidates and are often used in traditional medicine (Mulholland et al., 2013). Homoisoflavonoids exhibit antiangiogenic activity, reducing excessive formation of blood vessels. Several homoisoflavonoids have been investigated as prospective treatments for wet age-related macular degeneration, characterised by abnormal blood vessel growth. Homoisoflavonoids have potential for delivery as eye drops. Homoisoflavonoids, extracted from the Hyacinthaceae and synthesised, have been screened for antiangiogenic activity (Schwikkard et al., 2019) and have shown important structure-activity relationships. Notably, the presence and position of methoxy groups, the presence of the 3,9-double bond and the configuration at the chiral centre (C-3) are important for the activity. (Fig. 1) shows a compound extracted from *Pseudoprospéro firmifolium*, showing strong antiangiogenic activity, with a GI₅₀ value of 0.13 µM against Human Retinal Endothelial Cells. From this screening, we have undertaken targeted synthesis and further extractions of *Scilla* and species from the genus *Eucomis*, to increase our library of compounds for screening. Furthermore, we are investigating the possibility of growing certain bulbs commercially to extract active compounds, and comparing this to the costs of synthesising the compounds, in partnership with a UK commercial bulb grower.

Keywords: Hyacinthaceae, homoisoflavonoids, structure activity relationships, antiangiogenic activity

References

- Mulholland D, Schwikkard S, Crouch N. 2013. The chemistry and biological activity of the Hyacinthaceae. *Nat. Prod. Rep.* 30, 1165-1210.
- Schwikkard S, Whitmore H, Sishtla K, Sulaiman R. S, Shetty T, Basavarajappa H. D, Waller C, Alqahtani A, Frankemoelle L, Chapman A, Crouch N, Wetschnig W, Knirsch W, Andriantiana J, Mas-Claret E, Langat M. K, Mulholland D, and Corson T.W., 2019. The Antiangiogenic activity of naturally occurring and synthetic homoisoflavonoids from the Hyacinthaceae. *J. Nat. Prod.* 82, 1227-1239.

* Presenting author. E-mail address: h.jefford@surrey.ac.uk (Hannah Jefford)

S2.P8 *Pistacia lentiscus*: a plant with multiple virtues for use in human medicine as palliative in the treatment of inflammatory-related disorders

Djebbar Atmani^{a,*}, Naima Saidene^a, Dina Kilani^a, Ghania Bouguellid^a, Tristan Richard^b

^a Laboratory of Applied Biochemistry, Faculty of Life and Nature Sciences, University of Bejaia, 06000 Algeria

^b UR CEnologie, MIB, EA 4577, USC 1366 INRAE, Bordeaux INP, ISVV, F-33140, Villenave d'Ornon, France

New research trend is focusing on medicinal plants due to their wealth in various substances which can be potentially used in medicine, food and cosmetics. *Pistacia lentiscus*, a plant growing in the Mediterranean basin, is widely used in traditional medicine by rural populations for the treatment of various inflammatory-related disorders. Therefore, the present study was designed to investigate the antioxidant, anti-inflammatory, anticancer, anti-hyperuricemic and neuroprotective potential, as well as identification of active compounds, using appropriate methodology. *Pistacia lentiscus* extracts and fractions exhibited an outstanding scavenging capacity against a number of deleterious radicals while significantly enhancing superoxide dismutase and catalase activities, as well as reducing malonaldehyde levels, thus counterbalancing the oxidant status. Administration of *P. lentiscus* extracts significantly decreased ATP- and H₂O₂-induced mice paw oedema, reduced interleukin levels in cell culture and inhibited the uric acid-producing enzyme, xanthine oxidase, thereby enhancing the gouty situation of mice, mainly through a substantial reduction of uric acid levels. Additionally, *P. lentiscus* extracts showed a promising anticancer activity on human cancer cell lines, such as ovarian and liver carcinoma cells. On the other side, *P. lentiscus* extracts showed good neuro-protection and restored cognitive functions in mice, partially through inhibition of acetylcholine esterase and amyloid plaque formation and reversion of the harmful effects caused by Aluminum. Several experiments, *in vitro* and *in vivo*, including Ames' test and chromosomal aberration test indicated that *P. lentiscus* extracts are free of any genotoxicity, even showing antimutagenic action. Phytochemical investigations using preparative HPLC, HPLC-MS and NMR allowed the identification and structural elucidation of several known phenolic compounds and new anthocyanins, known to be responsible for the described biological activities, thus shedding more light on the phytochemical constituents of this plant and their action against disease. In light of the obtained results in terms of phytochemistry and biological activities, we may conclude that *Pistacia lentiscus* can be considered as a serious and safe alternative in the treatment of inflammatory conditions and deterioration of cognitive functions.

Keywords: *Pistacia lentiscus*, anti-hyperuricemia, xanthine oxidase, anti-mutagenic, phenols, anti-genotoxic

References

- Azib, L., Debbache-Benaid, N., Da Costa, G., Atmani-Kilani, D., Saidene, N., Ayouni, K., Richard, T., Atmani D., 2019. *Pistacia lentiscus* L. leaves extract and its major phenolic compounds reverse Aluminium-induced neurotoxicity in mice. *Industrial Crops and Products*, 137, 576-584.
- Charid, I., Kessler, M., Darb-Esfahani, S., Zemojtel, T., Abobaker, S., Tyuarets, S., Schrauwen, S., Atmani-Kilani, D., Benaid-Debbache, N., Schafer, R., Castillo-Tong, D.C., Atmani, D., Cherbal, F., Amant, F., Sehouli, J., Kulbe, H., Braicu E.I., 2020. Pretreatment with methanolic extract of *Pistacia lentiscus* L. increases sensitivity to DNA damaging drugs in primary high-grade serous ovarian cancer cells. *Eur. J. Integr. Med.*
- Aissat, A.K., Chahe-Bazizi, N., Richard, T., Kilani-Atmani, Pedrot, E., Renouf, E., Atmani D., Fonayet J.V., 2021. Analysis of individual anthocyanins, flavanols, flavonols and other polyphenols in *Pistacia lentiscus* L. fruits during ripening. *J. Food Compos. Anal.* 106.
- Bouguellid, G., Debbache-Benaid, N., Atmani-Kilani, D., Russo, C., Lavorgna, M., Piscitelli, C., Ayouni, K., Berboucha-Rahmani, M., Isidori M., Atmani D., 2022. *Pistacia lentiscus* L. fruits showed promising antimutagenic and antigenotoxic activity using both *in-vitro* and *in-vivo* test systems. *J. Toxicol. Environ. Health Part A.* 85(15), 603-621.
- Medjahed, Z., Chahe-Bazizi, N., Atmani-Kilani, D., Ahmane, N., Ruiz-Larrea, M.B., Ignacio, J., Sanz, Z., Charid, I., Amant, F., Fonayet, J.V., Saidene, N., Atmani, D., Richard T., 2023. A novel flavonol glycoside and six derivatives of quercetin and kaempferol from *Clematis flammula* with antioxidant and anticancer potentials. *Fitoterapia* 170, 105642

* Presenting author. E-mail address: djebbar.atmani@univ-bejaia.dz (Djebbar Atmani)

S2.P9 Evaluation of the effective dose of the aqueous root extract of *Sphenocentrum jollyanum* Pierre: Role as an antioxidant and phosphodiesterase inhibitor for erectile function

Olorunfemi Molehin^{a,*}, Oluwakemi Adeleke^b, Tolulope Obasuyi^b

^a Department of Biochemistry, Faculty of Science, Ekiti State University, Ado Ekiti, P.M.B. 5363 Ado-Ekiti, Nigeria

^b Biochemistry Unit, bDepartment of Science Laboratory Technology, Faculty of Science, Ekiti State University, Ado Ekiti, P.M.B. 5363 Ado-Ekiti, Nigeria

Improved sexual function is important in human race and with the dominance of medicinal species in the environment used in folklore as an aphrodisiac, it is necessary to understand how this plant species work and provide evidence for their mode of action (Adeleke et al., 2022; Singh et al., 2013). This study evaluated the sexual stimulating effect of aqueous extract from the root of *Sphenocentrum jollyanum* Pierre. The dose considered are 50mg/kg, 100mg/kg, 500mg/kg, 1000mg/kg body weights of the aqueous extract of the plant. The effects were assessed on sexual behavioural pattern, phosphodiesterase activity, nitric oxide level and the antioxidant status in penile tissues of rats. The aqueous root extract of *S. jollyanum* improved sexual performance, decreased the activity of phosphodiesterase while increasing the level of nitric oxide and improving the antioxidant status of the rats. Generally, the aqueous root extract at 500 mg/kg b.w administration showed a better effect of *S. jollyanum*. *Sphenocentrum jollyanum* can be considered therapeutically as an aphrodisiac.

Keywords: *Sphenocentrum jollyanum*, aphrodisiac, erectile function, phosphodiesterase

References

- Adeleke, O.A., Oboh, G., Adefegha, S.A., Osesusi, A. 2022. Effect of aqueous extract from root and leaf of *Sphenocentrum jollyanum* Pierre on wounds of diabetic rats: influence on wound tissue cytokines, vascular endothelial growth factor and microbes. J Ethnopharmacol 293, 115266.
- Singh, R., Ali, A., Jeyabalan, G., Semwal, A., Jaikishan. 2013. An overview of the current methodologies used for evaluation of aphrodisiac agents. J. Acute Disease 2 (2), 85-91.

* Presenting author. E-mail address: olorunfemi.molehin@eksu.edu.ng (Olorunfemi Molehin)

S2.P10 Antiepileptic effects of some medicinal plant hydrosols

Mohamad Norani^{a,*}, Mohammad-Taghi Ebadi^a, Alexander Crawford^a, Yaser Tahamtani^{b,c}, Mahdi Ayyari^a

^a Department of Horticultural Science, Tarbiat Modares University, Tehran, Iran

^b Department of Stem Cells and Developmental Biology, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, IR 19395-4644 Iran

^c Department of Diabetes, Obesity and Metabolism, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, IR 19395-4644 Iran

Epilepsy is a common chronic neurological disorder that affects individuals across all age groups (Brillatz et al., 2020). Traditional European and Iranian folk medicines have utilized various medicinal plants to treat neurological conditions, including epilepsy (Zadali et al., 2022). In this particular study, dried *Syzygium aromaticum* L. flower buds, *Valeriana officinalis* L. roots, *Cinnamomum verum* barks, *Achillea millefolium* aerial parts, *Nigella sativa* L. fruits, *Ferula gummosa* roots, and *Heracleum persicum* seeds were subjected to hydrodistillation to extract hydrosols. The organic phase of hydrosol (OPH) was obtained by extracting the hydrosols with a solvent in a separating funnel. Gas chromatography techniques were employed to examine the content and compositions of the OPH. The anticonvulsant activity of the OPH from all the samples was tested using the PTZ epilepsy zebrafish model. The results indicated that *S. aromaticum* OPH exhibited the strongest anticonvulsant activity (Ali, 2019), reducing PTZ-induced locomotor activity at the tested concentration of 3 µg/mL ($p < 0.001$). *V. officinalis* and *H. persicum* OPH also decreased locomotor activity at 100 µg/mL ($p < 0.001$ for *V. officinalis* OPH and $p < 0.01$ for *H. persicum* OPH). Notably, *V. officinalis* OPH at 30 µg/mL showed a significant increase in locomotor activity. In summary, this study suggests that certain hydrosols derived from medicinal plants, including *S. aromaticum*, *V. officinalis*, and *H. persicum*, exhibit potential anticonvulsant activity. The presence of specific volatile compounds in these hydrosols, such as eugenol, anethole, pulegone, borneol, and limonene, may contribute to their therapeutic effects.

Keywords: epilepsy, hydrosol, zebrafish, anticonvulsant activity, gas chromatography

References

- Ali, J.S.H. (2019). *Syzygium aromaticum* L. and their effect on seized chickens. Basra. J. Vet. Res. 18(1).
- Brillatz, T., Jacmin, M., Queiroz, E. F., Marcourt, L., Slacanin, I., Petit, C., & Wolfender, J. L. (2020). Zebrafish bioassay-guided isolation of antiseizure compounds from the Cameroonian medicinal plant *Cyperus articulatus* L. Phytomedicine 70, 153175.
- Zadali, R., Bagheri, M., Abbasi, M., Yavari, N., Miran, M., & Ebrahimi, S. N. (2022). Anticonvulsant activity of Iranian medicinal plants and molecular docking studies of isolated phytochemicals. South African Journal of Botany 149, 646-657.

* Presenting author. E-mail address: Mohamadnorani1368@gmail.com, m.ayyari@modares.ac.ir (Mohamad Norani, Mahdi Ayyari)

S2.P11 Alisol B 23-acetate and Alisol A 24-acetate reverse MDR cancer via ATPase modulation and membrane fluidity enhancement

Yu-Ning Teng*

School of Medicine, College of Medicine, I-Shou University, 8 Yida Road, Kaohsiung 82445, Taiwan, R.O.C.

Alisol B 23-acetate and Alisol A 24-acetate (Fig. 1) are compounds isolated from *Alismatis* rhizome (Alismataceae) and belong to the triterpenoids (Feng et al., 2021). Their diverse biological effects, including oncology-related issues, were previously reported by several research groups. This study aims to investigate their modulating mechanisms on multi-drug resistant (MDR) cancer, especially the case associated with P-glycoprotein (P-gp). P-gp is an efflux pump largely expressed on the cell membrane of cancer cells, leading to the MDR problem in chemotherapeutic treatment.




Fig. 1. Structures of Alisol B 23-acetate and Alisol A 24-acetate (images from PubChem library)

The cytotoxic IC₅₀ range for Alisol B 23-acetate and Alisol A 24-acetate on the cell lines used in this study was 16.34 to 60.13 μ M. The results showed that both Alisol B 23-acetate and Alisol A 24-acetate exhibited significant reversing ability at 10 μ M when combined with chemotherapeutic drug doxorubicin on an extensive panel of multidrug-resistant cancer cell lines, including KB/VIN, MCF-7/DOX, and HepG2/VIN (reversal folds ranged from 3.81 to 8.14). Both compounds demonstrated prominent inhibitory effects on P-gp efflux function at 10 μ M. Alisol A 24-acetate exhibited higher and dose-dependent P-gp basal ATPase activity stimulation than Alisol B 23-acetate at the dose range 1 to 50 μ M. Meanwhile, Alisol A 24-acetate influenced the cell membrane fluidity more than the Alisol B 23-acetate at 10 μ M, in both P-gp overexpressing cell lines Flp-InTM-293/*ABCB1* and HepG2/VIN.

Keywords: *Alismatis* rhizome, Alismataceae, MDR cancer, P-gp ATPase activity, membrane fluidity

References

Feng L, Liu TT, Huo XK, Tian XG, Wang C, Lv X, Ning J, Zhao WY, Zhang BJ, Sun CP, Ma XC, 2021. Alisma genus: Phytochemical constituents, biosynthesis, and biological activities. *Phytother Res.* 35(4), 1872-1886.

* Presenting author. E-mail address: yuningteng@isu.edu.tw (Yu-Ning Teng)

S2.P12 The constituents and anti-inflammatory activity from *Murraya paniculata* var. *omphalocarpa* Hayata

Kun-Ching Cheng^a, Ping-Chung Kuo^b, Chia-Hung Wu^c, Chun Lee^c, Tian-Shung Wu^{b,*}

^a Taiwan Sugar Research Institute, Tainan 70176, Taiwan

^b School of Pharmacy, College of Medicine, National Cheng Kung University, Tainan 701, Taiwan

^c School of Post-Baccalaureate Chinese Medicine, China Medical University, Taichung 404, Taiwan

One undescribed coumarin, (2'S)-isomurralonginol 1'-O-(2''S)-methylbutyrate (**1**) (Fig. 1), together with fourteen known coumarins, including isomurralonginol isovalerate (**2**), isomurralonginol (**3**), osthol (**4**), omphamurin (**5**), (+)-coumurrayin (**7**), 5,7-dimethoxy-8-(3'-methyl-2'-oxobutyl)coumarin (**8**), murrayanone (**9**), seselinal (**10**), ($\pm$)-7-methoxy-8-(2'-hydroxy-1'-methoxy-3'-methyl-3'-butenyl)coumarin (**11**), murragatin (**12**), mupanidin (**13**), mexotycin (**14**), and murraculatin (**15**), were isolated successively from the dried leaves of *Murraya paniculata* var. *omphalocarpa* Hayata (Rutaceae). On the other hand, we also isolated two alkaloids, γ -fargarine (**16**) and murrapanine (**17**); two coumarin-naphthoquinones, 8-naphthoherniarin (**18**) and toddacoumaquinone (**19**); three coumarins, **7**, **11**, and scopoletin (**20**); and two alkyl phenylpropanoates, methyl sinapate (**21**) and feruloyl esters (**22**), from the stem bark of the same plant. The undescribed coumarin **1** was characterized by the spectroscopic and spectrometric analytical methods, and the known compounds were identified by comparison of their physical and spectral data with those reported (Ito et al., 1990; Zhou et al., 1991). Some of the isolated compounds were evaluated for their anti-inflammatory activity (Yang et al., 2013). The results showed that IC₅₀ values of **4**, **7**, **17**, and **19** (Fig. 1) for the inhibition of superoxide anion generation ranged from 2.11 to 5121 ng/mL.




Fig. 1. Structures of **1**, **4**, **7**, **17**, and **19**

Keywords: *Murraya paniculata* var. *omphalocarpa*, Rutaceae, anti-inflammatory, superoxide anion generation

References

- Ito, C., Furukawa, H., Ishii, H., Ishikawa, T., Haginiwa, J., 1990. The chemical composition of *Murraya paniculata*. The structure of five new coumarins and one new alkaloid and the stereochemistry of murrangatin and related coumarins. *J. Chem. Soc. Perkin Trans. 1*, 2047–2055.
- Yang, S. C., Chung, P. J., Ho, C. M., Kuo, C. Y., Hung, M. F., Huang, Y. T., Chang, W. Y., Chang, Y. W., Chan, K. H., Hwang, T. L., 2013. Propofol inhibits superoxide production, elastase release, and chemotaxis in formyl peptide-activated human neutrophils by blocking formyl peptide receptor 1. *J. Immunol.* 190, 6511–6519.
- Zhou, P., Berova, N., Nakanishi, K., Rohmer, M., 1991. Assignment of absolute stereochemistry of aminopolyols by the bichromophoric exciton chirality method. *J. Chem. Soc. Chem. Commun.* 256–258.

* Presenting author. E-mail address: tswu@mail.ncku.edu.tw (Tian-Shung Wu)

S2.P13 Exploring Chemical Composition and α -Glucosidase Inhibitory of *Nelumbo nucifera* Seedpods

Pai-Hsin Chen, Ching-Kuo Lee *

School of Pharmacy, Taipei Medical University, No. 250, Wuxing St., Xinyi Dist., Taipei City 110, Taiwan (R.O.C.)

Lotus (*Nelumbo nucifera* Gaertn.) is an aquatic plant cultivated throughout East Asia, serving various purposes such as food, as an ornamental plant, and in medicinal materials. The literature indicates that lotus exhibits a variety of biological activities, and lotus seedpods are often regarded as waste (Huang et al., 2022). More than one in 10 adults worldwide suffers from diabetes. It is a systemic metabolic disorder that is related to cardiac diseases such as heart disease and stroke. According to statistics, most of the current cases are type-2 diabetes (Tang et al., 2023). Alpha- glucosidase inhibitors are a widely used target for the treatment of type-2 diabetes. Therefore, using an alpha-glucosidase rapid screening platform to identify potential active substances from natural products is a common preliminary research method. In this study, a 95% ethanol lotus seedpod extract exhibited an alpha-glucosidase inhibitory activity of 99% at 500 $\mu\text{g/mL}$. The extract was partitioned sequentially with ethyl acetate, *n*-butanol, and water. Preliminary bioassays showed that the ethyl acetate layer had good activity against α -glucosidase, tyrosinase, and xanthine oxidase. Therefore, the ethyl acetate extract was further purified by MPLC and HPLC to obtain the pure compounds. Pure isolates were characterized as flavonoids and triterpenoids using NMR and mass spectrometry. The isolation is still ongoing and the resulting compounds will be evaluated for their alpha-glucosidase inhibitory activity. We are also continuing to use a Lab-developed UHPLC- MS/MS methods (Chen et al., 2010) employed to find additional α -glucosidase inhibitors in lotus seedpods.

Keywords: *Nelumbo nucifera* Gaertn, alpha-glucosidase, HPLC

References

- Jingxian Huang, Seo Young Yang, Wenting Li, Wei Zhang, Zhen Li, Yan Feng, Shuying Dong, Zhuoyuan Lv, Yuan Du, Yan Zhao, 2022. Valorization of the industrial waste of two uncommon parts from lotus (*Nelumbo nucifera* Gaertn.): Insight to phytochemicals and hypolipidemic potential. *Ind. Crops Prod.* 188, 115596.
- Cheng Tang, Tingting Bao, Qiqi Zhang, Hongyu Qi, Yishan Huang, Boxun Zhang, Linhua Zhao, Xiaolin Tong, 2023. Clinical potential and mechanistic insights of mulberry (*Morus alba* L.) leaves in managing type 2 diabetes mellitus: Focusing on gut microbiota, inflammation, and metabolism. *J. Ethnopharmacol.* 306, 116143.
- Chin-Hui Chen, Pi-Yu Chen, Kuang-Chuan Wang, Ching-Kuo Lee, 2010. Rapid Identification of the Antioxidant Constituent of *Koeleruteria henryi*. *J. Chin. Chem. Soc.* 57, 404-410.

* Presenting author. E-mail address: m301111015@tmu.edu.tw (Ching-Kuo Lee)

S2.P14 *Fragaria x ananassa* Duch aerial by-product activity against *Candida albicans*

Mayra Gabriella Biccigo^a, Larissa Pavanello^b, Ailane Souza de Freitas^a, Rogerio Jose Machado Jr.^c, João Ernesto de Carvalho^a, Ana Lúcia Tasca Gois Ruiz^a, Karina Cogo Müller^a, Mary Ann Foglio^{a,*}

^a Faculty of Pharmaceutical Sciences, University of Campinas, Campinas, Brazil

^b Faculty of Dentistry at Piracicaba, University of Campinas, Brazil

^c Faculty of Medical Sciences, University of Campinas, Brazil

A *Fragaria x ananassa* Duch. is a plant native to the temperate zones of the Americas and Europe. Due to the presence of polyphenols, this plant species has antioxidant activity, with antifungal potential which the mechanisms of action involve plasma membrane and inhibition of protein synthesis disruption (Aboody & Mickymaray, 2020). The pseudofruit, popularly known as strawberry, has high benefit, with a 30% increase in global production in the last 10 years (FAOSTAT, 2024). However, approximately 7 to 20% of the production is by-product (Villamil-Galindo et al., 2022). Seeking to reduce by-product disposal with benefit, prompted this work to evaluate *Fragaria*'s production residue for antifungal activity. An extraction of the aerial plant parts with 70% ethanol with 0.3% citric acid provided the sample tested against strains of *Candida albicans* (ATCC 90028 and CBS 562), to determine Minimum Inhibitory Concentration (MIC) and Minimum Fungicide Concentration (MFC). The extract gave MIC of 27.9 µg/mL and 55.76 µg/mL for the *C. albicans* strains ATCC 90028 and CBS 562, respectively, and MFC values of 55.8 µg/mL for ATCC 90028, showing a moderate and strong cytostatic and cytotoxic activity for this strain. The by-product's extract provided high antimicrobial activity with potential as an alternative for the treatment of candidiasis, a worldwide and recurrent disease in immunosuppressed patients. (Seleem et al., 2017).

Keywords: *Fragaria x ananassa* Duch., phenolic compounds, antifungal activity, *Candida albicans*, by-product

References

- Aboody M.S.A, Mickymaray S., 2020. Anti-fungal efficacy and mechanisms of flavonoids. *Antibiotics* (Basel). 2020 9(2),45.
- Food And Agriculture Organizations Of The United Nations – Faostat. Disponível em: <http://www.fao.org/faostat/en/#data>. Acesso realizado em 31 de janeiro de 2024.
- Seleem, D., Pardi, V., Murata, R.M., 2017. Review of flavonoids: A diverse group of natural compounds with anti-*Candida albicans* activity *in vitro*, *Archives of Oral Biology* 76, 76-83.
- Villamil-Galindo, E., Antunes-Ricardo, M., Piagentini, A.M., Jacobo-Velázquez, D.A., 2022. Adding value to strawberry agro-industrial by-products through ultraviolet A-induced biofortification of antioxidant and anti-inflammatory phenolic compounds. *Front. Nutr.* 9.

* Presenting author. E-mail address: foglioma@unicamp.br (Mary Ann Foglio)

S2.P15 Integrated *in vitro* and *in vivo* antilithiatic efficacy of a synergistic herbal formulation, URO-5 and exploration of *in silico* guided mechanism of action

Ankita Misra^{a,*}, Mridul kant Chaudhary^a, Poonam Rawat^a, Hafizurrahman Khan^b, Vikas Srivastava^b, Dharendra Singh^c, Saroj Kanta Barik^d, Sharad Srivastava^{a,*}

^a Pharmacognosy Division, CSIR-National Botanical Research Institute, Lucknow-226001, U.P. India

^b System Toxicology & Health Risk Assessment, CSIR-Indian Institute of Toxicology Research, Lucknow-226001, U.P. India

^c Regulatory Toxicology, CSIR-Indian Institute of Toxicology Research, Lucknow-226001, U.P. India

^d Department of Botany, North-Eastern Hill University, Shillong-793022, Meghalaya, India

Urolithiasis, a metabolic condition caused by deposition of calculi in urinary tract has high prevalence and recurrence worldwide (Rule, 2020). In more than 60% of the clinical cases, kidney stones are made up of calcium oxalates (CaOx) and the existing treatment has limitation of side effects due to long and repeated use (Chattaraj, 2023). Therefore, a clinically validated, safe and efficacious synergistic herbal formulation (URO-5) was developed, which can be used as an adjuvant therapy. URO-5 contains five Indian medicinal plants having folklore use in urinary disease. The antilithiatic efficacy of LC-MS standardized aqueous extract was analysed, along with *in silico* guided mechanism of action. *In vitro* studies showed, that URO-5 drastically reduces the number of crystals and alters their morphology for easy passage. The inhibition (IC₅₀) of crystal nucleation and aggregation was at low concentration of 4.51 and 11.17 mg/mL. The *in vivo* activity (500 mg/kg) reveals that efflux of calcium oxalate (CaOx) stones were significant, > 80%. The decrease in tissue calcium and increase in urine calcium was also significant ($p > 0.05$), at 15.14 and 74.08 µg/mg. Interestingly, URO-5 also repairs the tissue damage caused by CaOx stones. *In silico* molecular docking suggest that antilithiatic activity of URO-5 was mediated *via* calcium and oxalate dependent, and antioxidant dependent pathways (Chattaraj, 2023). The safety and toxicity profile were documented under GLP compliance and no abnormality was recorded. Double-blind, placebo-controlled, single centred clinical study advocates that URO-5 is an efficacious alternative and can be used as an adjuvant therapy in urolithiasis.

Keywords: Urolithiasis, calcium oxalate crystals, adjuvant therapy, LC-MS, molecular docking

References

- Rule, A. D., et al. 2020. Management of kidney stones in 2020. *Jama* 323(19), 1961-1962.
- Chattaraj, B., et al. 2023. Inhibitory activity of *Enhydra fluctuans* Lour. on calcium oxalate crystallisation through *in silico* and *in vitro* studies. *Front. Pharmacol.* 13, 982419.
- Chattaraj, B., et al. 2023. Network pharmacology and molecular modelling study of *Enhydra fluctuans* for the prediction of the molecular mechanisms involved in the amelioration of nephrolithiasis. *J. Biomol. Struct. Dyn.* 41(24), 15400-15410.

* Presenting author. E-mail address: anku.mis@gmail.com (Ankita Misra)

S2.P16 Ashwagandha improves cognition and attenuates depressive-like behavior in an Alzheimer's disease mouse model

Noah Gladen Kolarsky^{a,b}, Wyatt Hack^{a,b}, Olivia Monestime^{a,b}, Melissa Bollen^{a,b}, Amala Soumyanath^{a,b}, Nora E. Gray^{a,b,*}

^a BENFRA Botanical Dietary Supplements Research Center, Oregon Health & Science University (OHSU) Portland, OR, 97239, USA

^b Department of Neurology, OHSU, Portland OR 97239, USA

Withania somnifera (WS), also known as Ashwagandha, is a traditional Ayurvedic herb popularly used as a dietary supplement for effects on memory, stress, anxiety, depression and insomnia. Cognitive impairment and depression are common symptoms of Alzheimer's disease (AD), suggesting that WS may be of therapeutic value in that condition. Previous studies from our group with *Drosophila* have shown that an aqueous extract of WS root (WSAq) improves performance in a phototaxis assay and alleviates depressive symptoms. Here we sought to evaluate the effects of WSAq in a mammalian system using the 5xFAD mouse model of β -amyloid accumulation. Six- to seven-month-old male and female 5xFAD mice were treated with WSAq in their drinking water at either 0, 0.5 or 2.5 mg/mL for 4 weeks. Age-matched non-transgenic wild type littermates that received no WSAq were also included as a control group. In the fourth week of treatment mice underwent behavioral testing. The higher dose of WSAq (2.5 mg/mL) improved spatial memory and attenuated depressive-like symptoms in the 5xFAD mice. Analysis of plaque pathology as well as markers of neuroinflammation and oxidative stress is ongoing to gain mechanistic information on how WSAq could be eliciting these beneficial effects. Although future studies are needed to confirm these results in other models, these initial results suggest that WSAq does have potential as a therapeutic option for use in AD.

This work was supported by a pilot grant from the Oregon Partnership for Alzheimer's Research and the BENFRA center U19 grant U19AT010829 funded by NIH/NCCIH, USA.

Keywords: Ashwagandha, Alzheimer's disease, cognitive function, depression

* Presenting author. E-mail address: grayn@ohsue.edu (Nora Gray)

S2.P17 Chemical structure and bioactivities of a sulfated polysaccharide from precious medicinal fungi

Mei-Kuang Lu^{a,b}, Chi-Hsein Chao^a, Tsu-Yuan Chang^c, Ming-Che Cheng^c, Yu-Chi Hsu^a, Chia-Chuan Chang^{c,*}

^a National Research Institute of Chinese Medicine, Ministry of Health and Welfare, Taipei, Taiwan, R.O.C

^b Graduate Institute of Pharmacognosy, Taipei Medical University, Taipei, Taiwan, R.O.C

^c School of Pharmacy, College of Medicine, National Taiwan University, Taipei, Taiwan, R.O.C

Antrodia cinnamomea Chang & Chou, a valuable medicinal and edible fungi, demonstrated hepatoprotective, immunomodulative, anti-tumor, anti-inflammatory and anti-oxidative activities (Xie et al., 2022). Sulfated polysaccharides (SPSs) are polysaccharides with hemi-ester sulfate groups employed to maintain the integrity of the structural skeleton of fungi, marine invertebrate connective tissues, algae and bacteria. A SPS, **3-SS**, was isolated from *A. cinnamomea*. Its chemical structure was identified by utilizing 1D and 2D NMR techniques. The anti-inflammation effects of **3-SS** on RAW264.7 macrophage cells, such as IL-6 inhibition, restoration of LPS-induced IκB protein degradation, and inhibited LPS-induced TGFβII protein degradation, were confirmed to occur via AKT, ERK1/2, and p-38. In addition, **3-SS** impaired the proliferation of H1975 lung cancer cells through EGFR/ERK/slug signaling.

Keywords: medicinal fungi, sulfated polysaccharides, structural elucidation, anti-inflammation, anti-proliferation

References

Xie, L., Huang, Z., Qin, L., Yu, Q., Chen, Y., Zhu, H., Xie, J., 2022. Effects of sulfation and carboxymethylation on *Cyclocarya paliurus* polysaccharides: Physicochemical properties, antitumor activities and protection against cellular oxidative stress, Intl. J. Biol. Macromol. 204, 103-115.

* Presenting author. E-mail address: chiachang@ntu.edu.tw (Chia-Chuan Chang)

S2.P18 Unlocking the therapeutic potential: Comparative analysis of tormentic acid and its ester in combating inflammation and oxidative stress

Mara Heckmann^{a,*}, Gerald Klanert^{a,b}, Sylvain Legay^c, Julian Weghuber^{a,b}

^a Center of Excellence Food Technology and Nutrition, University of Applied Sciences Upper Austria, Stelzhamerstraße 23, 4600 Wels, Austria

^b FFoQSI GmbH–Austrian Competence Centre for Feed and Food Quality, Safety and Innovation, Technopark 1D, 3430 Tulln, Austria

^c Luxembourg Institute of Science and Technology (LIST), 5 Avenue Des Hauts-Fourneaux, L-4362, Esch-sur-Alzette, Luxembourg

This poster illustrates the anti-inflammatory and antioxidant properties of the pentacyclic triterpene tormentic acid (**1**) and its ester tormentic acid 3-O-p-coumaroyl ester (**2**). **1** has been shown to exhibit promising therapeutic potential, but **2** remains underexplored (Olech et al., 2021). By comparative analysis, we aim to elucidate how esterification affects the biological efficacy of such pentacyclic triterpenes. Using diverse *in-vitro* assays across various cell lines, we assessed **1** and **2**'s ability to mitigate oxidative stress by scavenging reactive oxygen and nitrogen species. Additionally, we investigated their capacity to modulate mRNA expression and protein production of pro-inflammatory cytokines upon lipopolysaccharide (LPS) challenge. Our findings revealed that **2** significantly surpassed **1** in effectively reducing levels of reactive oxygen species and nitric oxide. Moreover, we presented an extensive analysis of protein expression levels of over 100 cytokines and chemokines in THP-1 macrophages post-LPS stimulation. Addition of **1** or **2** to the cells resulted in different protein expression patterns, with **2** significantly reducing the levels of several pro-inflammatory cytokines involved in NF- κ B and Prostaglandin E2 signaling (e.g., TNF α , IL-8, IL-17A and IL-2), thereby counteracting cytokine storms. These results suggest that the esterified form **2** exerts superior biological activity compared to **1**, potentially through its impact on NF- κ B signaling. Future experiments will focus on analyzing their effects on TLR4 and downstream signaling pathways to further explore this hypothesis. In conclusion, elucidating the differential effects of esterification on the biological activity of pentacyclic triterpenes opens new avenues for developing effective and natural anti-inflammatory and antioxidant therapies.

Keywords: pentacyclic triterpene, esterification, tormentic acid, antioxidant, anti-inflammatory

References

Olech, M., Ziemichód, W., Nowacka-Jechalke, N., 2021. The occurrence and biological activity of tormentic acid - A review. *Molecules*. 26, 3797.

* Presenting author. E-mail address: mara.heckmann@fh-wels.at (Mara Heckmann)

S2.P19 Anti-diarrheal activity of *Psidium guajava* extract and partitions in *Drosophila melanogaster*

Chengmei Liu*, Valérie Jullian, François Chassagne

UMR 152 PharmaDev, Université Paul Sabatier, Institut de Recherche pour le Développement, Toulouse, France

Diarrheal diseases are among the most common causes of death, especially in developing countries (Ugboko et al., 2020). In traditional medicine from many tropical countries, *Psidium guajava* leaves represent one of the most frequently employed plant species to treat diarrhea, however no specific compounds have been described so far. Indeed, existing anti-diarrheal models are mainly murine models requiring large concentrations of plant materials and significant quantities of mice, thus hindering a bio-guided fractionation of the molecules involved. In this study, we used a genetically modified strain of *Drosophila melanogaster* for isolating antidiarrheal compounds from guava by bio-guided fractionation (Liu and Chassagne, 2023). Previously, it was shown that the *Drosophila* gene *Ion transport peptide* (*ITP*) is an important endocrine regulator of water intake and excretion. With *ITP* knockdown, the frequency of defecation events is increased and the transit through the digestive tract is faster, and so results in a phenotype resembling human diarrhea (Gáliková et al., 2018). In our experiments, antidiarrheal activity was assessed using this *D. melanogaster* strain by detecting various parameters of fecal deposits (number of fecal deposits, total area of deposits, total material excreted (total IOD)). The test groups, with 10 replicates and 6 flies per replicate, received 1g/100 mL ethanol extracts while the positive control group received racecadotril. Each ethanolic extract of guava collected in New Caledonia (NC), French Polynesia (FP) and Benin (B) significantly reduced the number of fecal deposits (reduction of 61.0% (NC), 60.9% (FP), 57.1% (B)), total area of deposits (reduction of 74.2% (NC), 67.0% (FP), 65.7% (B)), and total IOD (reduction of 77.3% (NC), 63.3% (FP), 71.1% (B)), compared to the group administrated with normal food. We are working on confirming the antidiarrheal activities of the ethanolic extract of *P. guajava* using *Drosophila* with other methods, as well as identifying the bioactive component.

Keywords: *Psidium guajava*, *Drosophila melanogaster*, antidiarrheal, fecal deposits

References

- Gáliková, M., Dirksen, H., Nässel, D.R., 2018. The thirsty fly: Ion transport peptide (ITP) is a novel endocrine regulator of water homeostasis in *Drosophila*. *PLOS Genetics* 14, e1007618.
- Liu, C., Chassagne, F., 2023. Assessment of The Effect of Antidiarrheal Drugs and Plant Extracts on *Drosophila melanogaster*. *Journal of Visualized Experiments* e65877. <https://doi.org/10.3791/65877>
- Ugboko, H.U., Nwinyi, O.C., Oranusi, S.U., Oyewale, J.O., 2020. Childhood diarrhoeal diseases in developing countries. *Heliyon* 6, e03690.

* Presenting author. E-mail address: liuchengmei1997@gmail.com (Chengmei Liu)

S2.P20 The influence of growth conditions on the amount of active substances in *Mentha x piperita* L.

Asta Klimienė, Erika Kaškelienė, Laura Normantė

Klaipėda University, H. Manto 84, 92294 Klaipėda, Lithuania

Mentha x piperita L. commonly known as peppermint, is a hybrid resulting from a cross between *Mentha spicata* L. (spearmint) and *Mentha aquatica* L. (watermint). Peppermint is an herbaceous rhizomatous perennial plant, native to Lithuania as well as in Europe and the Middle East which can be found wild with its parent species, typically in moist habitats. In this study was determine dependent of total phenolic, flavonoid and extractives content in extracts of *Mentha x piperita* L. on soil pH, humus, total nitrogen (N_{total}), plant – available/mobile phosphorus (P₂O₅) as well as potassium (K₂O), total calcium (Ca), magnesium (Mg) and sodium (Na) concentration. Experimental fields were formed in four locations in Lithuania. Soil and perennial peppermint samples were taken at three sites of each experimental field. A total 12 samples of soil and 12 samples of tested plant were collected and analysed. The concentration bioactive compounds in plants are significantly ($P < 0.05-0.01$) affected by some chemical properties of the soil: the total phenolic content reliably correlated with soil pH, N_{total}, Ca, Mg and P₂O₅; the flavonoid content – with soil P₂O₅, K₂O, Mg and Na; the extractives content – with soil humus, N_{total} and Na. The obtained results are useful in the development for commercial cultivation of peppermint

Keywords: *Mentha x piperita* L., total phenols, flavonoids, extractives, soil chemical properties

Presenting author. E-mail address: asta.klimiene@ku.lt (Asta Klimienė)

S2.P21 Inhalation - A promising administration route for anti-infectives from *Morus alba* root bark?

Sigrid Adelsberger^{a,b,*}, Jacqueline Schwarzwinger^{c,b}, Gabriela Hädrich^c, Judith M. Rollinger^a, Lea Ann Dailey^c, Ulrike Grienke^a

^a Department of Pharmaceutical Sciences, Division of Pharmacognosy, Faculty of Life Sciences, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^b Vienna Doctoral School of Pharmaceutical, Nutritional and Sport Sciences, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^c Department of Pharmaceutical Sciences, Division of Pharmaceutical Technology and Biopharmaceutics, Faculty of Life Sciences, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

Mulberry Diels-Alder type adducts (MDAAs) isolated from *Morus alba* root bark were identified as dual antiviral and antibacterial agents *in vitro* [Grienke et al., 2016]. However, no significant efficacy was observed for oral administration in an *in vivo* influenza A virus infection model [Langeder et al., 2023a]. Hence, the aim of this study was to assess the suitability of inhalation administration for this compound class. Two MDAA-containing extracts (MA21 and MA60, [Langeder et al., 2023b]) and their isolated MDAA stereoisomers, sanggenons C and D, were investigated for (i) cytotoxicity and (ii) permeability using the lung epithelium cell line, Calu-3, (iii) pH-dependent solubility, (iv) logP and pKa, and (v) plasma protein binding. Both extracts and pure compounds were well tolerated by Calu-3 cells up to ~800 µg/mL and ~170 µM. The apparent permeability (P_{app}) values were low ($0.07\text{--}0.26 \times 10^{-6} \text{ cm}\cdot\text{sec}^{-1}$) compared to the control compound, metoprolol ($10.3 \times 10^{-6} \text{ cm}\cdot\text{sec}^{-1}$ [Bosquillon et al., 2017]) which indicates the compounds could be retained in the lung following inhalation. Despite their structural similarity, sanggenon D showed a tenfold higher solubility compared to sanggenon C at a pH range of 1.2–7.4. Similarly, logP of sanggenon D was one log unit lower than sanggenon C (1.90 vs. 2.91). This study suggests that the purified sanggenons exhibit more favorable properties for pharmaceutical development compared to extracts. The low P_{app} values may indicate that membrane permeability is low, explaining the lack of efficacy in oral administration models. However, inhalation administration could be promising for treating respiratory infections.

Funding: FWF project P 35115

Keywords: *Morus alba*, Moraceae, inhalation, infections, permeability

References

- Bosquillon, C., Madlova, M., Patel, N., Clear, N., Forbes, B., 2017. A comparison of drug transport in pulmonary absorption models: isolated perfused rat lungs, respiratory epithelial cell lines and primary cell culture. *Pharm. Res.* 34(12), 2532–2540.
- Grienke, U., Richter, M., Walther, E., Hoffmann, A., Kirchmair, J., Makarov, V., Nietzsche, S., Schmidtke, M., Rollinger, J.M., 2016. Discovery of prenylated flavonoids with dual activity against influenza virus and *Streptococcus pneumoniae*. *Sci. Rep.* 6, 27156.
- Langeder, J., Koch, M., Schmietendorf, H., Tahir, A., Grienke, U., Rollinger, J.M., and Schmidtke, M., 2023a. Correlation of bioactive marker compounds of an orally applied *Morus alba* root bark extract with toxicity and efficacy in BALB/c mice. *Front. Pharmacol.* 14, 1193118.
- Langeder, J., Döring, K., Schmietendorf, H., Grienke, U., Schmidtke, M., Rollinger, J.M., 2023b. ¹H NMR-based chemometric analysis of *Morus alba* extracts toward a multipotent herbal anti-infective. *J. Nat. Prod.* 86(1), 8–17.

* Presenting author. E-mail address: sigrid.adelsberger@univie.ac.at (Sigrid Adelsberger)

S2.P22 Reporting of medical and health research - An assessment of studies on medicinal plant extracts and herbal medicines registered on PROSPERO

Banaz Jalil^{a,*}, Manael Baloch^a, Michael Heinrich^{a,b,*}

^a Pharmacognosy and Phytotherapy Group, UCL School of Pharmacy, London WC1N 1AX, UK;

^b Chinese Medicine Research Center, College of Chinese Medicine, China Medical University, Taichung, Taiwan

Studies evaluating the pharmacological effects, (clinical) efficacy or toxicity of medicinal plant extracts are constantly increasing, but the reporting quality and reproducibility of these studies remain unsatisfactory. The Consensus-based guidelines for the Phytochemical characterization of Medicinal Plant extracts (ConPhyMP), with the Open access ConPhyMP tool (Heinrich et al., 2022; Heinrich and Jalil, 2023), facilitate transparent reporting, reproducibility, and interpretations of studies. PROSPERO is an international prospective registry for systematic reviews with a health-related outcome, reducing duplication and reporting bias in studies. Using adapted ConPhyMP guidelines, we assess the reporting quality of 1727 registered studies involving medicinal plant extracts/herbal medicines on PROSPERO from inception to January 2024 (Fig. 1). Out of 65 studies included in our detailed analysis, only five report the minimum requirements (i.e., adhering to ConPhyMP guidelines), including reporting of starting materials, conducting, and reporting of phytochemical analysis of plant extracts/herbal medicines under investigation. Within 3-5 years of registration on PROSPERO, 83% of the studies remain under review, 13% are discontinued reviews, and only 4% reach publications. For specific medicinal plants, we found that *Cannabis sativa* L., *Curcuma longa* L., *Zingiber officinale* Roscoe, *Panax ginseng* C.A.Mey., and *Ginkgo biloba* L. have the highest registered studies on PROSPERO (n= 952, 362, 210, 176, 92, respectively), with n= 92, 32, 17, 2, 4, respectively reaching publications. There is a need for assessment and scrutiny for the registration of health-related research involving plant extracts/herbal medicines, especially for those with clinical relevance or generation of evidence-based information, which could enhance the study publications.




Fig. 1. Workflow chart for inclusion and exclusion of studies in the systematic review using PRISMA model (Page et al., 2021).

Keywords: PROSPERO, ConPhyMP, medicinal plant extracts, study registration, medical and health research

References

- Heinrich, M., Jalil, B., 2023. From the CONSORT to the ConPhyMP statement and beyond—how to ascertain best practice. *Front. Pharmacol.* 14, 1338710. Heinrich, M., Jalil, B., Abdel-Tawab, M., Echeverria, J., Kulić, Ž., McGaw, L.J., Pezzuto, J.M., Potterat, O., Wang, J.-B., 2022. Best practice in the chemical characterisation of extracts used in pharmacological and toxicological research—The ConPhyMP—Guidelines. *Front. Pharmacol.* 13, 953205.
- Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., Shamseer, L., Tetzlaff, J.M., Akl, E.A., Brennan, S.E., Chou, R., Glanville, J., Grimshaw, J.M., Hróbjartsson, A., Lalu, M.M., Li, T., Loder, E.W., Mayo-Wilson, E., McDonald, S., McGuinness, L.A., Stewart, L.A., Thomas, J., Tricco, A.C., Welch, V.A., Whiting, P., Moher, D., 2021. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 372, n71.

* Presenting author. E-mail address: b.jalil@ucl.ac.uk (Banaz Jalil)

S2.P23 Anti-inflammatory and antioxidant effects of *Marrubium vulgare* L. herb extracts in human immune cells and blood plasma

Anna Magiera^{a,*}, Małgorzata Kołtun-Jasion^b, Joanna Kołodziejczyk-Czepas^c, Oleksandra Liudvytska^c, Anna Kiss^b, Monika Anna Olszewska^a

^a Department of Pharmacognosy, Faculty of Pharmacy, Medical University of Lodz, 1 Muszynskiego St., 90-151 Lodz, Poland

^b Department of Pharmaceutical Biology, Medical University of Warsaw, 1 Banacha St., 02-097 Warsaw, Poland

^c Department of General Biochemistry, Faculty of Biology and Environmental Protection, University of Lodz, 141/143 Pomorska St., 90-236 Lodz, Poland

Marrubium vulgare (L.) herb has been valued in traditional medicine in many regions as a cardiogenic, vasoprotective and hypotensive remedy. The monograph of *Marrubii herba* is included in the European Pharmacopeia. However, the herb is rarely used in phytotherapy, mainly due to incomplete data on its pharmacological profile in the context of cardiovascular diseases (CVDs) (Teixidor-Toneu et al., 2016; Aćimović et al., 2020). This study focused on the impact of different *Marrubii herbae* extracts (characterized by LC-MS/MS) and their active components (marrubiin, verbascoside, apigenin glucoside) on the functions of neutrophils and PBMCs. Dysregulation of immune cells is largely associated with the development of CVDs. Hence, the effect on the release of reactive oxygen species (ROS), pro-inflammatory cytokines (IL-2, IL-8, TNF- α) and enzymes (ELA-2, MMP-9) in stimulated immune cells was measured. Moreover, the influence of extracts/active compounds on human fibrinogen and other plasma components under (ONOO⁻)-induced oxidative stress and on the expression of neutrophil adhesion molecules was evaluated. The biological activity tests were performed using immunoenzymatic, spectrophotometric/fluorometric, chemiluminescence, cytometric, SDS-PAGE, and western blot methods. As a result, the tested extracts/active compounds (5-100 μ g/mL, 5-50 μ M) were proved to significantly decrease the levels of ROS, TNF- α , IL-2, MMP-9, some biomarkers of lipid peroxidation and protein nitration, and to protect fibrinogen against oxidative/nitrative changes in its structure and function. The results might partly support the traditional use of the plant material in CVDs and encourage further mechanistic and in vivo studies in that matter.

Keywords: *Marrubium vulgare*, immune cells, fibrinogen, human plasma

References

- Teixidor-Toneu, I., Martin, G.J., Ouhammou, A., Puri, R.K., Hawkins, J.A., 2016. An ethnomedicinal survey of a Tashelhit-speaking community in the High Atlas, Morocco. *J. Ethnopharmacol.* 188, 96-110.
- Aćimović, M., Jeremić, K., Salaj, N., Gavarić, N., Kiprovski, B., Sikora, V., Zeremski T., 2020. *Marrubium vulgare* L.: a phytochemical and pharmacological overview. *Molecules.* 25,12, 2898.

* Presenting author. E-mail address: anna.magiera@umed.lodz.pl (Anna Magiera)

S2.P24 Antioxidant, anti-diabetic and anti-inflammatory potential of *Sorbus commixta* (Rosaceae) leaves in cellular, cell-free and human plasma models

Magdalena Rutkowska^a, Anna Magiera^{a,*}, Joanna Kolodziejczyk-Czepas^b, Martyna Pabich^a, Monika A. Olszewska^a

^a Department of Pharmacognosy, Faculty of Pharmacy, Medical University of Lodz, 1 Muszynskiego St., 90-151 Lodz, Poland

^b Department of General Biochemistry, Faculty of Biology and Environmental Protection, University of Lodz, 141/143 Pomorska St., 90-236 Lodz, Poland

Sorbus commixta Hedl. (Japanese rowan) has a long tradition of use as a herbal remedy (leaves, fruits, stem, bark) for atherosclerosis and liver or respiratory disorders (Sołtys et al., 2020). Considering the previous reports on the biological properties of *S. commixta* leaves (mainly antioxidant), as well as the available pharmacological data on anti-diabetic and anti-inflammatory potential of *S. commixta* bark and fruits, and the leaves of other *Sorbus* species (Matczak et al., 2018; Sołtys et al., 2020), we decided to evaluate these three activity directions for *S. commixta* leaves in cellular, cell-free and human plasma models (Fig. 1). As a result, we demonstrated for the first time the potent protective effects of the *S. commixta* leaf extracts against nitration and oxidation of plasma proteins and lipids under ONOO⁻-induced oxidative stress *ex vivo*; the ability to inhibit the formation of advanced glycation end products and α -amylase and α -glucosidase activity *in vitro*; and the potential to reduce the secretion of pro-inflammatory cytokines (TNF- α , IL-1 β , IL-2, IL-6) in the LPS-stimulated PBMC cells *in vitro*. Moreover, the phytochemical profiling of the tested extracts (qualitative and quantitative UHPLC-PDA-ESI-MS³ and HPLC-PDA analyses) revealed the presence of seventy-eight phenolic compounds (in comparison to only 13 phenolics reported previously). Furthermore, the contribution of individual polyphenols to the tested activity was verified with chemometric tools (PCA analysis).




Fig. 1. Study design

Keywords: *Sorbus commixta*, leaves, antioxidant, anti-diabetic, anti-inflammatory

References

- Matczak, M., Marchelak, A., Michel, P., Owczarek, A., Piszczan, A., Kolodziejczyk-Czepas, J., Nowak, P., Olszewska, M.A., 2018. *Sorbus domestica* L. leaf extracts as functional products: phytochemical profiling, cellular safety, pro-inflammatory enzymes inhibition and protective effects against oxidative stress *in vitro*. J. Funct. Foods, 40, 207-218.
- Sołtys, A., Galanty, A., Podolak, I., 2020. Ethnopharmacologically important but underestimated genus *Sorbus*: a comprehensive review. Phytochem. Rev. 19, 491-526.

*Presenting author. E-mail address: anna.magiera@umed.lodz.pl (Anna Magiera)

S2.P25 *In vitro* antioxidant capacity screening of *Piper auritum* extracts

Bertha Fenton Navarro^{a,*}, Adriana Aburto Ortiz^b, Aarón Vázquez Hernández^c

^a Laboratorio de Glicobiología y Farmacognosia, División de Estudios de Posgrado, Facultad de Ciencias Médicas y Biológicas "Dr. Ignacio Chávez", Universidad Michoacana de San Nicolás de Hidalgo (UMSNH), Morelia 58020, Michoacán, Mexico

^b Hospital de Especialidades, CMN SXXI, IMSS, Ciudad de México, México

Oxidative stress results from the production of unregulated reactive oxygen species (ROS). When low antioxidant levels and low regenerative capacity occur, tissue damage is produced. It has been reported that oxidative stress contributes to the progression of several diseases. Antioxidants are molecules that can donate electrons to free radicals without making them unstable; this causes the free radicals to stabilize and become less reactive, preventing, and repairing damage caused by ROS/RNS. *Piper auritum* is used to treat different ailments. To determine the antioxidant capacity *in vitro* of *Piper auritum* leaf extracts using different solvents. Aqueous extract (AQEPA), methanol extract (MEPA), ethanol extract (EEPA), acetone extract (AEPA), chloroform extract (CEPA), and hexane extract (HEPA) were obtained from *Piper auritum* leaves. The antioxidant activity *in vitro* was evaluated using the 1-1-dyphenyl-2-picrylhydrazyl (DPPH) method (Bonet and Brand- Williams, 1995), statistical ANOVA, and Tukey post hoc analysis was performed; results are presented as EC50, TEC50, and AE (IC50 x TEC50). The antiradical efficiency (AE) obtained was AQEPA:1051, MEPA:412.5, EEPA:19.74, AEPA:26.52, CEPA:33.6, and HEPA:3.71. The high AE obtained is due to the presence of anthocyanins, tannins, terpenoids, phenols, polyphenols, and flavonoids in aqueous ethanol and methanol. In acetone, phenol and flavonols are dissolved, while chloroform and hexane contain terpenoids, flavonoids, and coumarins. The presence of these secondary metabolites in the different solvents used explained the high antioxidant capacity obtained. *Piper auritum* extracts have high *in vitro* antioxidant capacity.

Keywords: *Piper auritum*, antioxidant capacity, solvents

* Presenting author. E-mail address: bertha.fenton@umich.mx (Bertha Fenton Navarro)

S2.P26 Chemical constituents and antioxidant properties of *Verbesina montanoifolia* leaf extracts

Bertha Fenton Navarro^{a,*}, Graciela Letechipía Vallejo^b, Aarón Vázquez Hernández^c

^a Laboratorio de Glicobiología y Farmacognosia, División de Estudios de Posgrado, Facultad de Ciencias Médicas y Biológicas “Dr. Ignacio Chávez”, Universidad Michoacana de San Nicolás de Hidalgo (UMSNH), Morelia 58020, Michoacán, Mexico

^b Laboratorio de Neurociencias. División de Estudios de Posgrado. Facultad de Ciencias Médicas y Biológicas “Dr. Ignacio Chávez”, Universidad Michoacana de San Nicolás de Hidalgo (UMSNH), Morelia 58020, Michoacán, Mexico

^c Hospital de Especialidades, CMN SXXI, IMSS, Ciudad de México, México

Reactive oxygen species (ROS) are products of normal cellular metabolism, they can damage lipids, DNA, RNA, and proteins, which, in theory, contribute to the physiology of aging. Oxidative stress may play a significant role in the development of different diseases such as cancer, diabetes, cardiovascular conditions like high blood pressure, atherosclerosis and stroke, inflammatory disorders, and chronic fatigue syndrome, among others. *Verbesina montanoifolia* (VM), a shrub native to Mexico, can measure up to 2 meters in height with elongated dark green leaves and yellow flowers. Is used in Mexican traditional medicine to treat different illnesses. However, there are no reports worldwide of the components or their antioxidant capacity, so this was the aim of the present work. An aqueous extract of *Verbesina montanoifolia* leaves was obtained. The total content of phenols, polyphenols, and flavonoids was quantified. The *in vitro* antioxidant capacity was determined using the compound DPPH (1,1-diphenyl-2-picrylhydrazyl) at different times. A gas chromatography-mass spectrometry on Agilent 7000C Triple Quad w/ 7890B GC/MS/MS system with an Agilent J&W GC Column was performed to identify the components. VM extracts contained high levels of flavonoids, phenols, and polyphenols that presented antioxidant effects *in vitro*. In an HPLC analysis, the main compounds were vanillic acid, hydroxybenzoic acid, and Gallic acid. These three compounds are known for their antioxidant properties. To the best of our knowledge, this is the first study to report the chemical constituents of VMAQE with antioxidant properties.

Keywords: reactive oxygen species, oxidative stress, antioxidants, flavonoids, polyphenols

* Presenting author E-mail address: bertha.fenton@umich.mx (Bertha Fenton Navarro)

S2.P27 Isolation and characterization of α -glucosidase inhibitors from *Ziziphora tenuior* using high- performance counter-current chromatography

Niloofer Zonoubi^{a,*}, Somayeh Mojtavai^b, Mohammad-Reza Delnavazi^a, Petri Turhanen^c, Mohammad Ali Faramarzi^b, Hamid-Reza Adhami^a

^a Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

^b Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

^c School of Pharmacy, Biocenter Kuopio, University of Eastern Finland, Kuopio, Finland

Diabetes Mellitus (DM) is a common chronic disease characterized by hyperglycemia, causing different complications. One successful approach to managing DM is to reduce postprandial hyperglycemia by inhibiting carbohydrate enzymes, such as α -glucosidase. Efforts have been made in recent years to find potent α -glucosidase inhibitors derived from medicinal plants (Phukhatmuen et al., 2020). *Ziziphora* spices from the Lamiaceae family are traditionally used for the treatment of DM (Mohammadhosseini, 2017). In this study, *Ziziphora tenuior* (L.) (Kakuti in Persian), with small leaves and pink-purple cluster flowers, was investigated for α -glucosidase inhibitory activity. The whole air-dried plant powder was extracted stepwise by hexane, EtOAc and MeOH and the extracts were evaluated for α -glucosidase inhibitory effects. The selected methanol extract was fractionated by vacuum liquid chromatography. The sub-fractions were subjected to high- performance counter-current chromatography (HPLCCC) obtaining five pure compounds. Their structures were elucidated as pinocembrin-7-O-rutinoside (1), rosmarinic acid (2), pinocembrin (3), narirutin (4) and chlorogenic acid (5) using NMR spectroscopy and mass spectrometry. All the pure compounds were examined for α -glucosidase inhibitory effect (*in-vitro* anti-type 2 diabetic assay). Compound 3 exhibited the best enzyme inhibition among the others with an IC₅₀ value of 3.8 ± 0.1 μ M. The findings indicate that *Ziziphora tenuior* has the potential for diabetic treatment.

Keywords: anti-diabetic, α -glucosidase inhibitory activity, *Ziziphora tenuior*, HPLCCC, pinocembrin

References

Phukhatmuen, P., Raksat, A., Laphookhieo, S., Charoensup, R., Duangyod, T. and Maneerat, W., 2020. Bioassay-guided isolation and identification of antidiabetic compounds from *Garcinia cowa* leaf extract. *Heliyon*, 6(4). Mohammadhosseini, M., 2017. The ethnobotanical, phytochemical and pharmacological properties and medicinal applications of essential oils and extracts of different *Ziziphora* species. *Industrial crops and products*, 105, pp.164-192.

* Presenting author. E-mail address: nzonoubi93@gmail.com (Niloofer Zonoubi)

S2.P28 Triterpenoids and flavonoids of Corncockle – a (‘n almost) forgotten weed and the (future) medicinal herb

Adam Matkowski^{a,b,*}, Izabela Nawrot-Hadzik^b, Reneta Gevrenova^c, Łukasz Pecio^d, Solomiya Pecio^d, Daria Zblewska^e, Marcin Surma^b, Marek Surma^{a,f}, Maria Malicka^f, Weronika Korzeniowska^b, Aleksander Smakosz^b, Weronika Kozłowska^f

^a Botanical Garden of Medicinal Plants, Wrocław Medical University, Al. Jana Kochanowskiego 14, 50367 Wrocław, Poland

^b Department of Pharmaceutical Biology and Botany, Wrocław Medical University, Borowska 211A, Wrocław, Poland

^c Department of Pharmacognosy, Sofia Medical University, ul. Dunav 2, Sofia, Bulgaria

^d Department of Crop Biochemistry, IUNG-Institute of Cultivation and Soil Sciences, ul. Krańcowa 8, 24-100 Puławy, Poland

^e Student Scientific Organization, Circle No. 76, Department of Pharmaceutical Biology and Biotechnology, Wrocław Medical University, Borowska 211, Wrocław, Poland

^f Department of Pharmaceutical Biotechnology, Wrocław Medical University, Borowska 211, Wrocław, Poland

The botanical family *Caryophyllaceae* (pink or carnation family) comprises herbaceous species with cosmopolitan distribution and characteristic morphology. With respect to phytochemistry, *Caryophyllaceae* typically contain triterpenoid saponins and several species have been used to obtain surfactants for washing, cosmetics, food industry (halva) and in pharmaceutical technology. Some other species have been considered medicinal and some as poisonous. *Agrostemma githago* L. was once a noxious weed in cereal fields but the agriculture modernization led to its eradication. Now, this plant is rare and survived mainly in fringe habitats and as ornamental. However, it also has a history as medicinal herb a few centuries ago but nowadays it is not used. Several saponins have been recently isolated with important pharmacokinetic properties (Clochard et al. 2020). A presence of orientin has been also reported in the past (Jakimiuk et al. 2022). Here, we used an LC- High-Resolution-Mass Spectrometry approach to contribute to *A. githago* phytochemistry and were able to detect 24 triterpenoid glycosides (mono- and bidesmosides with linear or branched glycosylation) where sapogenins were built of gypsogenin, quillajic acid, oleanolic acid and hydroxyoleanolic acid. In the aerial parts, the predominant compounds were orientin and isoorientin, along with 6 other flavone-C-glycosides, suggesting that this fast growing, high- biomass herb can provide a feasible, sustainable source of highly bioactive polyphenols, in addition to previously isolated saponins and ribosome-inactivating proteins.

Acknowledgements. This research is funded by National Science Center of Poland (NCN) # 2020/39/I/NZ7/01515 (RIPSAPO).

Keywords: *Agrostemma githago*, *Caryophyllaceae*, triterpene saponins, flavonoid C-glycosides

References

- Clochard, J., Jerz, G., Schmieder, P., Mitdank, H., Tröger, M., Sama, S., Weng, A., 2020. A new acetylated triterpene saponin from *Agrostemma githago* L. modulates gene delivery efficiently and shows a high cellular tolerance. *Intl. J. Pharmaceutics*. 589, 119822.
- Jakimiuk, K., Wink, M., Tomczyk, M., 2022. Flavonoids of the *Caryophyllaceae*. *Phytochem. Rev.* 21, 179–218.

* Presenting author. E-mail address: pharmaceutical.biology@wp.eu (Adam Matkowski)

S2.P29 Therapeutic effects of Menthacarin® and its individual components peppermint oil and caraway oil in an experimental colitis model

Sandra Christochowitz*, Silas F. Melcher, Sabine Tremmel, Sabrina Weisenburger, Martin D. Lehner

Preclinical R&D, Dr. Willmar Schwabe GmbH & Co. KG, Karlsruhe, Germany

Menthacarin® is a proprietary combination of peppermint oil (90 mg WS® 1340) and caraway oil (50 mg WS® 1520) of specified quality that is clinically used for the treatment of functional gastrointestinal disorders. Further, Menthacarin® has demonstrated to attenuate experimental colitis via anti-inflammatory effects (Alliger K. et al., 2020). However, its effect on the colonic mucus-producing cell loss, contributing to intestinal barrier dysfunction and thereby promoting intestinal inflammation, is unclear. Further, the relative contribution of the individual oils in the experimental colitis model is still not known. Therefore, mice with 3 % dextran sulfate sodium (DSS)-induced colitis were treated orally with Menthacarin® (30/60/90 mg/kg), peppermint oil (20/40/80 mg/kg), or caraway oil (10/20/40 mg/kg) for one week. Colonic tissue destruction and mucus area from Menthacarin®-treated mice were assessed in formalin-fixed paraffin-embedded Swiss rolls by Hematoxylin-Eosin and Alcian Blue periodic acid-Schiff (AB-PAS) staining, respectively. In line with a reduced body weight loss and colon length shortening, Menthacarin® (30 mg/kg) attenuated DSS-mediated mucosal destruction accompanied by an increased AB-PAS-positive area. Peppermint oil and caraway oil at the lowest dose had an ameliorative effect as determined by a reduction in body weight loss and colon length shortening. In conclusion, this data indicates that Menthacarin® promotes intestinal barrier function and shows for the first time that both, peppermint oil and caraway oil, contribute to the attenuation of experimental colitis.

Keywords: Menthacarin, peppermint, caraway, colitis, mucus

References

Alliger, K., Khalil, M., König, B., Weisenburger, S., Koch, E., Engel, M., 2020. Menthacarin attenuates experimental colitis. *Phytomedicine* 77, 153212.

* Presenting author. E-mail address: sandra.christochowitz@schwabe.de (Sandra Christochowitz)

S2.P30 *Bryophyllum pinnatum* inhibits oxytocin and vasopressin signalling in myometrial cells

Leonie Zurfluh^{a,*}, Lauriane Duvaud^b, Nejla İnçi^b, Olivier Potterat^c, Ana Paula Simões-Wuest^a, Johannes Mosbacher^b

^a Department of Obstetrics, University Hospital Zurich, University of Zurich, Zurich, Switzerland

^b Institute of Pharma Technology, School of Life Sciences, University of Applied Sciences and Arts Northwestern Switzerland FHNW, Muttens, Switzerland

^c Division of Pharmaceutical Biology, University of Basel, Basel, Switzerland

The medicinal plant *Bryophyllum pinnatum* (Lam.) Oken (Crassulaceae) is used in the treatment of preterm contractions and was previously shown to block oxytocin (OT)-induced signals in myometrial cells, consistent with its tocolytic effect observed in patients (Plangger et al., 2006; Santos et al., 2021). Since OT activates OT receptors and V1a receptors, both expressed in the myometrium (Åkerlund et al., 1999), we aimed to study the molecular pharmacology of *B. pinnatum* press juice (BPJ) using specific receptor ligands, the human myometrial cell line hTERT- C3, and cell lines expressing recombinant human OT and V1a receptors in a fluorescent calcium experiment. We found that BPJ inhibits both OT- and vasopressin (AVP)-induced intracellular calcium ($[Ca^{2+}]_i$) increase in hTERT-C3 myometrial cells. In the same functional assay, BPJ also inhibited signals mediated by recombinant human OT and V1a receptor with a similar potency (IC_{50} about 0.5 mg/mL). We further studied the details of endogenous OT- and AVP-sensitive receptors in hTERT-C3 cells and found that OT and AVP stimulated those receptors with similar potency (EC_{50} of ~1 nM), suggesting expression of both receptor subtypes. This interpretation was corroborated by the observed antagonist potencies of atosiban and relcovaptan. However, using qPCR, we almost exclusively found expression of OT receptors suggesting a pharmacological difference between recombinant OT receptors and those expressed in hTERT-C3 cells. In conclusion, *B. pinnatum* inhibits both, OT and AVP signalling, which may point beyond its tocolytic effects to other indications involving a disbalance in the vasopressinergic system.

Keywords: *Bryophyllum pinnatum*, oxytocin, vasopressin, myometrium

References

- Plangger, N., Rist, L., Zimmermann, R., von Mandach, U., 2006. Intravenous tocolysis with *Bryophyllum pinnatum* is better tolerated than β -agonist application. Eur. J. Obstet. Gynecol. Reprod. Biol. 124(2), 168-172.
- Santos, S., Zurfluh, L., Mennet, M., Potterat, O., Von Mandach, U., Hamburger, M., Simões-Wüst, A. P. 2021. *Bryophyllum pinnatum* compounds inhibit oxytocin-induced signaling pathways in human myometrial cells. Front Pharmacol. 12, 142.
- Åkerlund, M., Bossmar, T., Brouard, R., Kostrzewska, A., Laudanski, T., Lemancewicz, A., Secradeil-Le Gal, C., & Steinwall, M. 1999. Receptor binding of oxytocin and vasopressin antagonists and inhibitory effects on isolated myometrium from preterm and term pregnant women. BJOG 106(10), 1047-1053.

* Presenting author. E-mail address: leonie.zurfluh@usz.ch (Leonie Zurfluh)

S2.P31 Didesmethylocaglamide Cytotoxic Activity in High Grade Serous Ovarian Cancer in Preclinical *In Vitro* and *In Vivo* Models

Elizabeth N. Kaweesa^{a,*}, Tyler A. Wilson^b, Jaqueline M. Bazioli^a, Li Pan^b, A. Douglas Kinghorn^b, James R. Fuchs^b, Joanna E. Burdette^a

^a Department of Pharmaceutical Sciences, College of Pharmacy, University of Illinois at Chicago, Chicago, IL 60607, USA

^b Division of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, The Ohio State University, Columbus, OH 43210, USA

Didesmethylocaglamide (DDR) (**1**) is a naturally occurring derivative of rocaglamide with potent antitumor activity isolated from *Aglaia* plant species (Fig. 1) (Chang et al., 2020). DDR displays nanomolar cytotoxic activity in high grade serous ovarian cancer (HGSOC) cell lines including OVCAR8. HGSOC is the most lethal gynecological cause of death in women and requires new treatments to help tackle chemoresistance (Feng et al., 2021). Rocaglamides are known to function as translation inhibitors and trigger apoptosis in other types of solid tumors. We found that racemic **1** induced cytotoxicity in ovarian cancer cell lines as early as 24 hours and activated caspase-3 indicating pro-apoptotic activity. In addition, **1** was cytotoxic for the PEO4 and MCF7ADR cell lines that are resistant to cisplatin and paclitaxel, respectively. In addition, we evaluated **1** in OVCAR8 xenografts and observed a reduction in tumor burden. We hypothesized that combinatorial treatments of **1** with autophagy inhibitors like phyllanthusmin-34 would enhance cell death by blocking translation and the ability to recycle amino acids for protein production through autophagy (Salvi et al., 2022). The data obtained indicates that blocking autophagy or DNA repair in combination with **1** increased cell death in the sensitive and resistant models. Its effective cytotoxicity in drug sensitive and resistant cell models suggests **1** has potential as a new therapeutic strategy against HGSOC.




Fig. 1. Structure of (±)-didesmethylocaglamide (**1**)

Keywords: didesmethylrocaglamide, ovarian cancer, HGSOC, cytotoxicity, chemoresistance

References

- Chang, L.S., Oblinger, J.L., Burns, S.S., et al., 2020. Targeting protein translation by rocaglamide and didesmethylrocaglamide to treat MPNST and other sarcomas. *Mol. Cancer Ther.* 19(3),731-741.
- Feng, L.Y., Yan, B.B., Huang, Y.Z. et al., 2021. Abnormal methylation characteristics predict chemoresistance and poor prognosis in advanced high-grade serous ovarian cancer. *Clin. Epigenet.* 13, 141.
- Salvi A., Young A.N., Huntsman, A.C., et al., 2022. PHY34 inhibits autophagy through V-ATPase V0A2 subunit inhibition and CAS/CSE1L nuclear cargo trafficking in high grade serous ovarian cancer. *Cell Death Dis.* 13(1), 45.

* Presenting Author. E-mail address: ekaweesa@uic.edu (Elizabeth Kaweesa)

S2.P32 Isobavachalcone attenuates TNF- α -induced vascular inflammation in human umbilical vein endothelial cells

Sang Yoon Choi^{*}, Ae Sin Lee^a

Food Functionality Research Division, Korea Food Research Institute, 55365 Wanju, Republic of Korea

Isobavachalcone (Fig. 1) is an active molecule present in the medicinal plant *Psoralea corylifolia* Linn. and has been reported to have antioxidant, anti-inflammatory (Gao et al., 2019), antibacterial (Wang et al., 2021) and anticancer activities (He et al., 2021). However, to our knowledge, studies on the protective effects of isobavachalcone on vascular inflammation have not been reported. Vascular inflammation activated by pro-inflammatory cytokines is an inflammatory response that occurs in the early stages of atherosclerosis. Endothelial dysfunction in vascular inflammation begins with the expression of cell surface adhesion molecules by pro-inflammatory cytokines. In this study, we investigated the effects of isobavachalcone on inflammatory responses in vascular inflammation induced by the tumor necrosis factor- α (TNF- α) in human umbilical vein endothelial cells (HUVECs). TNF- α is critical for increasing the expression of adhesion molecules, such as intercellular adhesion molecule 1 (ICAM-1) and vascular cell adhesion molecule 1 (VCAM-1) in endothelial cells, and activates macrophages and neutrophils. Isobavachalcone decreased the expression of TNF- α -induced ICAM-1 and VCAM-1 in HUVECs, and the adhesion of monocytes to the vascular endothelium and the adhesion of monocytes to the vascular endothelium. In addition, isobavachalcone decreased the phosphorylation of the NF- κ B (necrosis factor κ B) p65 subunit. The result of this study demonstrate that isobavachalcone could prevents TNF- α -induced vascular inflammation and has the potential to protect against the early progression of atherosclerosis




Fig. 1. Structure of Isobavachalcone

Keywords: isobavachalcone, vascular inflammation, vascular cell adhesion molecule 1 (VCAM-1), induced intercellular adhesion molecule 1 (ICAM-1), NF- κ B

References

- Gao, D., Liu, F., Li, Z., Guan, Y., 2019. Isobavachalcone attenuates Sephadex-induced lung injury via activation of A20 and NRF2/HO-1 in rats. *Eur. J. Pharmacol.* 848, 49-54.
- He, H., Wang, C., Liu, G., Ma, H., Jiang, M., Li, P., Lu, Q., Li, L., Qi, H., 2021. Isobavachalcone inhibits acute myeloid leukemia: Potential role for ROS-dependent mitochondrial apoptosis and differentiation. *Phytother. Res.* 35, 3337-3350.
- Wang, M., Lin, L., Lu, J.J., Chen, X., 2021. Pharmacological review of isobavachalcone, a naturally occurring chalcone. *Pharmacol. Res.* 165, 105483.

* Presenting author. E-mail address: sychoi@kfri.re.kr (Sang Yoon Choi)

S2.P33 Synergistic therapeutic potential of quercetin, curcumin, and berberine combination in triple negative breast cancer: insights from *in vitro* models

Akanksha Kashyap, Sheikh Mohammad Umar, Arundhathi Dev JR, Chandra Prakash Prasad*

Department of Medical Oncology Lab, Dr BRA IRCH, AIIMS, New Delhi, 110029, India

In this study, we explored the therapeutic potential of QUE, CUR, and BBR in *in vitro* TNBC models. Our hypothesis posited that combining these polyphenols might enhance their efficacy in TNBCs compared to individual treatments. Using MDA-MB-468 and MDA-MB-231 as TNBC cell line models, we assessed the impact of individual phytochemicals and their combined treatment (COM^{QUE+CUR+BBR}) on Epithelial Mesenchymal Transition (EMT), migration potential, cancer stem cells (CSCs), clonogenic/3D matrigel growth, and drug resistance in TNBC cells. Our results revealed that the combination treatment (COM^{QUE+CUR+BBR}) exerted significant and synergistic effects on EMT proteins and cell migration, surpassing the impact of individual treatments. Furthermore, COM^{QUE+CUR+BBR} treatment significantly reduced the CD44⁺/CD24⁻ population, clonogenic potential, and inherent cisplatin resistance in TNBC cells. Notably, all combination experiments utilized 1/10th of the respective IC₅₀ concentrations, highlighting the robust efficacy of these phytochemicals in TNBC models. These findings represent the first evidence that the combination of QUE, CUR, and BBR holds promise for TNBC treatment, showcasing its anti-proliferative, EMT-inhibitory, and antagonistic cancer stemness functions.

Keywords: breast cancer, TNBC, quercetin, curcumin, berberine, EMT

References

Kashyap A., Umar S.M., Dev A.J.R., Prasad C.P., 2022. *In vitro* anticancer efficacy of a polyphenolic combination of Quercetin, Curcumin, and Berberine in triple negative breast cancer (TNBC) cells. *Phytomedicine Plus* 100265.

* Presenting author. E-mail address: researchchandra@gmail.com (Chandra Prakash Prasad)

S2.P34 Quercetin aims at the HuR-signaling axis to impede the migration of Triple Negative Breast Cancer (TNBC): Association with metabolic processes

Arundhathi Dev J.R., Akanksha Kashyap^a, Sheikh Mohammad Umar, Chandra Prakash Prasad*

Department of Medical Oncology Lab, Dr BRA IRCH, AIIMS, New Delhi, 110029, India

In our present research, our focus was on evaluating the influence of a natural compound, specifically Quercetin, a flavonoid, on the proliferation and migration of Triple Negative Breast Cancers (TNBCs). Our particular interest centered around examining the effects of quercetin on the RNA binding protein HuR, given previous indications that quercetin can disrupt HuR-mRNA interactions. Utilizing online meta-analysis, we delved into the role of HuR in TNBC datasets. Subsequently, we validated HuR protein expression in TNBC cell lines, specifically MDA-MB-231 and MDA-MB-468. We determined the IC₅₀ of quercetin in TNBC cell lines through MTT assays and assessed protein expression using western blotting. Additionally, we investigated cell migration and colony-forming potential in TNBC cells treated with quercetin. Our findings highlighted that HuR overexpression correlated with reduced overall survival in TNBC patients. Correspondingly, both TNBC cell lines exhibited heightened expression of HuR proteins. Quercetin treatment significantly suppressed HuR protein expression in TNBC cells. Further analysis revealed an active HuR-PFKP-LDHA signaling pathway in TNBC cells, and inhibiting this pathway led to reduced migration and clonogenic potential in TNBC cells. In summary, our research demonstrated that an elevated HuR-PFKP-LDHA axis is associated with TNBC progression and recurrence. The ability of quercetin to inhibit these proteins suggests its potential as an anticancer agent for treating aggressive TNBCs.

Keywords: breast cancer, TNBC, HuR, lactate, cell migration

References

- Umar S.M., Patra S., Kashyap A., Dev J. R. A., Kumar L., Prasad C.P., 2021. Quercetin impairs hur-driven progression and migration of Triple Negative Breast Cancer (TNBC) Cells. *Nutr Cancer* 19, 1-14.
- Umar S.M., Kashyap A., Kahol S., Mathur S., Gogia A., Deo S.V.S., Prasad C.P. 2020. Prognostic and therapeutic relevance of phosphofructokinase platelet-type (PFKP) in breast cancer. *Exp Cell Res.* 10, 112282.

* Presenting author. E-mail address: researchchandra@gmail.com (Chandra Prakash Prasad)

S2.P35 Biological activities of the polar and nonpolar extracts of Philippine Bamboo species

Chona D. Gelani*, Lina S. Lacandalo, Renzo Luis D. Reazo

Department of Chemistry, Mindanao State University-Iligan Institute of Technology, Iligan City, Philippines

Ethanol (E) and hexane (H) extracts of the shoots and outer covering (C) of three (3) bamboo species namely, *Dendrocalamus asper* (Da), *Bambusa blumeana* (Bb), and *Gigantochloa levis* (Gl) were subjected to an array of tests to determine their antibacterial, antidiabetic, anti-inflammatory, antioxidant, and angiotensin-converting enzyme (ACE) inhibition activity. Among the outer covering extracts, GlCE and BbCH exhibited the lowest MIC for both *Bacillus subtilis* and *Pseudomonas aeruginosa* at 0.780 µg/mL. At 800 µg/mL, DaE inhibited alpha amylase activity at

82.86 % followed by GlE and BbE with 79.31 and 75.95% inhibition, respectively. GlE exhibited the lowest IC₅₀ (i. e. 2.97 µg/mL), whereas IC₅₀ values for DaE and BbE are 3.03 and 3.13 µg/mL, respectively. The results show that these bamboo extracts have antidiabetic potential. Extracts from the bamboo shoots outer covering did not exhibit anti-inflammatory activity based on the standard 50% inhibition activity limit using 15-LOX inhibition assay. The highest % inhibition was that of DaCE (7.75%) followed by GlCE (6.19%) then BbCE (5.54%) at a concentration of 33.33 µg/mL. DPPH assay resulted to GlCH exhibiting the highest % Radical Scavenging Activity (RSA, 65.96%) followed by GlE at 58.13%. BbE, exhibited the lowest EC₅₀ at 92.53 µg/mL. GlCH, which exhibited the highest % RSA, is the most potent antioxidant among the samples with EC₅₀ value of 82.85 µg/mL. GlCE and BbCH show remarkable ACE inhibition activity with values 0.00044 and 0.00282 nmol/min, respectively.

Keywords: bamboo shoots, bamboo shoot covering, *Dendrocalamus asper*, *Bambusa blumeana*, *Gigantochloa levis*

References

- Nirmala, C., Bisht, M., Bajwa, H., Santosh, O., 2018. Bamboo: A rich source of natural antioxidants and its applications in the food and pharmaceutical industry. Trends in Food Science and technology. 77, 91-99.
- Park, E., Jhon, D., 2009. The antioxidant, angiotensin converting enzyme inhibition activity and phenolic compounds of bamboo shoot extracts. Journal of Food Science and Technology. 43(4), 655-659.
- Sarker, S.D., Nahar, L., Kumarasamy, Y., 2007. Microtitre plate-based antibacterial assay incorporating resazurin as an indicator of cell growth, and its application in the in vitro antibacterial screening of phytochemicals. Methods 42(4), 321-324.
- Tanaka, A., Zhu, Q., Tan, H., Horibo, H., Ohnuki, K., Mori, Y., Yamauchi, R., Ishikawa, H., Iwamoto, A., Kawahara, H., Shimizu, K., 2014. Biological Activities and Phytochemical Profiles of Extracts from Different Parts of Bamboo (*Phyllostachys pubescens*). Molecules 19(6), 8238–8260.

* Presenting author. E-mail address: chona.gelani@g.msuiit.edu.ph (Chona D. Gelani)

S2.P36 Terpenoids and polyphenols from *Carpesium cernuum* (Asteraceae) of European origin

Anna Stojakowska^{a,*}, Janusz Malarz^a, Klaudia Jakubowska^a, Danuta Jantas^a, Anna K. Kiss^b

^a Maj Institute of Pharmacology, Polish Academy of Sciences, 12 Smętna Str., 31-343 Kraków, Poland

^b Department of Pharmaceutical Biology, Medical University of Warsaw, 1 Banacha Str., 02-097, Warsaw, Poland

Carpesium cernuum L., the most widespread species of the genus *Carpesium* (Inuleae-Inulinae, Asteraceae), in the Far East and India found use as a folk remedy and as a food plant. To the best of our knowledge, the plant has not been medicinally used in Europe. Phytochemical examination of the plant material of European origin (Romania), cultivated in the Garden of Medicinal Plants, Maj Institute of Pharmacology PAS in Kraków, led to the isolation of a series of known monoterpenoid thymol derivatives and a new dihydrobenzofuran **1**, an analog of eupatobenzofuran (Chen et al., 2011), from the roots of plants. Aerial parts of the plants were rich in germacranolides with 4 β ,8 α - dihydroxy-5 β -isobutyryloxy-9 β -(3-methylbutyryloxy)-3-oxo-germacran-7 β ,12 α -olide, a sesquiterpene lactone of cytotoxic activity (Kłeczek et al., 2021), as one of the major constituents. An UHPLC-PAD MSⁿ analysis of hydroalcoholic extracts from the roots and aerial parts of *C. cernuum* revealed the presence of numerous caffeic acid conjugates (mainly caffeoylquinic and caffeoylhexaric acids) formerly unknown as constituents of *C. cernuum*.




Fig. 1. Structure of a new dihydrobenzofuran derivative **1**

One of the isolated thymol derivatives - 8-hydroxy-9,10-diisobutyryloxythymol - was assayed for cytotoxicity and potential neuroprotective activity using undifferentiated and differentiated SH- SY5Y neuroblastoma cells. The compound at concentrations higher than 25 μ M significantly reduced viability of both differentiated and undifferentiated cells. At a concentration range of 1-10 μ M the thymol derivative provided partial protection against H₂O₂-induced cell damage in undifferentiated neuroblastoma cells.

Keywords: *Carpesium cernuum*, Asteraceae, dihydrobenzofuran, hydroxycinnamates, thymol derivatives

References

- Chen, J.-J., Tsai, Y.-C., Hwang, T.-L., Wang, T.-C., 2011. Thymol, benzofuranoid, and phenylpropanoid derivatives: Anti-inflammatory constituents from *Eupatorium cannabinum*. *J. Nat. Prod.* 74, 1021-1027.
- Kłeczek, N., Malarz, J., Gierlikowska, B., Skalniak, Ł., Galanty, A., Kiss, A.K., Stojakowska A. Germacranolides from *Carpesium divaricatum*: Some new data on cytotoxic and anti-inflammatory activity. *Molecules* 26, 4644.

* Presenting author. E-mail address: stoja@if-pan.krakow.pl (Anna Stojakowska)

S2.P37 Diet-induced differences in gut microbiota determine the responder or non-responder of traditional herbal medicine

Kento Takayama^{a,b,*}, Masanori Iseki^b, Chiho Takahara^a, Ryoko Tomita^c, Katsuhiko Ishihara^d, Tadashi Hayama^c, Takashi Tanaka^c, Gen-ichiro Nonaka^f, Hirofumi Inoue^a

^a Faculty of Pharmacy and Pharmaceutical Sciences, Fukuyama University, 1 Sanzo, Gakuen-cho, Fukuyama, Hiroshima, Japan

^b Kawasaki Medical School, 577 Matsushima, Kurashiki, Okayama, Japan

^c Faculty of Pharmaceutical Sciences, Fukuoka University, 8-19-1 Nanakuma, Jyonan-ku, Fukuoka, Japan

^d Kawasaki University of Medical Welfare, 288 Matsushima, Kurashiki, Okayama, Japan

^e Institute of Biomedical Sciences, Nagasaki University, 1-12-4 Sakamoto, Nagasaki, Japan

^f Usaien Pharmaceutical Co., Ltd., 1-4-6 Zaimoku, Saga, Japan

Daiokanzoto (DKT), commonly used clinically to treat constipation, is a traditional herbal medicine comprising rhubarb and glycyrrhiza. The purgative action of DKT is primarily due to sennoside A, the main laxative constituent in rhubarb, which is metabolized by gut microbiota. It is widely acknowledged that dietary habits influence gut microbiota. Therefore, we hypothesize that these habits may influence the purgative activity of DKT, thereby categorizing individuals as either responders or non-responders to DKT treatment. In this study, we manipulated gut microbiota of mice by feeding them a high-carbohydrate or high-fiber diet. To examine the connection between laxatives and gut microbiota, we monitored the gut microbiome before and after administering laxatives. Mice on the high-carbohydrate diet exhibited significantly high purgative activity with DKT. This correlated with increased Enterobacteriaceae abundance due to reduced immunoglobulin A regulation. However, the high-fiber diet greatly reduced DKT efficacy. Additionally, rhein 8-O- β -D-glucopyranoside, gallate, and tannin, which are DKT constituents, contributed to the difference between responders and non-responders. These results demonstrate that diet-induced differences in gut microbiota determine the effectiveness of DKT. This is intriguing, given that Oriental medicines are typically tailored to specific functional states, or “patterns.” These results reveal that the usefulness of DKT depends on multiple constituents. Furthermore, dietary habits influence gut microbiota, impacting the success of this traditional medicine.

Keywords: daiokanzoto, gut microbiota, immunoglobulin A, short-chain fatty acids, diet

* Presenting author. E-mail address: takayama@fukuyama-u.ac.jp (Kento Takayama)

S2.P38 Antioxidant activity and cell viability assay on L6 myotubes of different parts of *Mitragyna speciosa*

Supattra Limsuwanchote^{a,*}, Rinnapat Boonsut^a, Charoenwong Premprasert^b, Juraithip Wungsintaweekul^b

^a Pharmacology Program, Division of Health and Applied Sciences, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla, 90110, Thailand

^b Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat-Yai, Songkhla, 90110, Thailand

Mitragyna speciosa Korth., commonly known as kratom (Rubiaceae), has been used in traditional medicine to alleviate several ailments such as diarrhea, cough, muscle pain, and some metabolic diseases like diabetes, hypertension, and dyslipidemia. Among its indole alkaloids, mitragynine is the principal component interacting with opioid receptors (Assanangkornchai et al., 2009; Nakaphan et al., 2016). In this study, different parts of kratom, including leaves extracted with methanol (LM) and water (LW), and branch (BM) and flower (FM) extracted solely with methanol were examined for their total phenolic and flavonoid contents and antioxidant activity. *In vitro* antioxidant activity was assessed including 2,2-diphenyl-1-picryl-hydrazil (DPPH) radical scavenging, ferric reduction (FRAP), and measurement of lipid peroxidation (TBARS). The results demonstrated that LM and LW contained high levels of total phenolic and flavonoid contents. In contrast, BM exhibited high total phenolic content but lower flavonoid content, which differs from the levels observed in FM. Among the different parts, leaves, whether LM or LW, exhibited the better antioxidant activity, followed by branch and flower, respectively. The impact of kratom on L6 myotube viability was evaluated using MTT assay. After 24 h of incubation with various concentrations (6.25-100 µg/mL), all extracts, except LW, did not induce cytotoxic effect within the range of 6.25-50 µg/mL. However, at 100 µg/mL, LM, BM, and FM decreased cell viability to less than 80%, whereas LW did not exhibit cytotoxicity. This study suggests that kratom leaves, particularly LW, could be a promising source of natural antioxidants deserving further research and development.

Keywords: *Mitragyna speciosa*, kratom, antioxidant, cell viability

References

- Assanangkornchai, S., Muekthong, A., Sam-Angsri, N., Pattanasattayawong, U., 2007. The use of *Mitragynine speciosa* ("Krathom"), an addictive plant, in Thailand. *Subst Use Misuse*. 42(14), 2145-2157.
- Nakaphan, T., Teerachaisakul, M., Puttum, S., Sompima, K., Nootim, P., 2016. Traditional uses of kratom (*Mitragyna speciosa* Korth) among folk healers in Southern Thailand. *J. Thai Trad. Alt. Med.* 14(3), 274-285.

* Presenting author. E-mail address: supattra.l@psu.ac.th (Supattra Limsuwanchote)

S2.P39 pH-dependent membrane permeabilisation through saponins and how it can enhance endosomal escape

Fabian E. Bülow*, Hardy Mitdank, Alexander Weng

Institute of Pharmacy, Department of Pharmaceutical Biology, Freie Universität Berlin, Königin- Luise-Straße 2+4, 14195 Berlin, Germany

Biologicals such as antibodies, enzymes and nucleic acids are arousing interest in pharmaceutical industry and academia. This rapidly growing field could be the key to cure diseases that were considered untreatable for a long time, with the addition that their therapeutic effect is limited through their poor cellular uptake. After endocytosis of these large molecules, they are subject to endosomal entrapment. The consequence: only a small amount of the applied biological reaches its cellular target. One approach for optimizing their cellular delivery is to lower the integrity of endosomal membranes through amphiphilic substances such as saponins. The saponin Agrostemmoside E (or rather AG1856), derived from the common corncockle (*Agrostemma githago* L.), displays such an enhancing effect on the endosomal escape. The main limitation of using saponins as transfection-enhancers is their cytotoxicity, caused by an unselective lysis of cytoplasmic membranes. This work investigated the influence of pH on the membrane- permeabilizing activity of triterpensaponins. The idea was to apply Agrostemmoside E in a concentration that would not affect cytoplasmic membranes at physiological pH, but only the endosomes as soon as they reach a specific acidity. Agrostemmoside E displayed such a pH- dependent behavior, showing enhanced membranolytic activity in acidic conditions. The data was generated via hemolysis- and LDH-assays. Saponin-concentrations and incubation-times for a selective endosomal escape were suggested as well as a possible biochemical explanation for the observed pH-sensitivity.

Keywords: *Agrostemma githago*, endosomal escape, saponin, pH-dependency, membranolysis

References

- Clochard, J. et al. 2020. A new acetylated triterpene saponin from a githago L. modulates gene delivery efficiently and shows a high cellular tolerance. *International Journal of Pharmaceutics* 589.
- Fuchs, H. et al. 2017. Glycosylated Triterpenoids as endosomal escape enhancers in targeted tumor therapies. *Biomedicines* 5(2), 14.

* Presenting author. E-mail address: fabian.buelow@fu-berlin.de (Fabian E. Bülow)

S2.P40 Cytotoxic potential of saponins and fruit extracts from two genotypes of *Sechium edule* (Cucurbitaceae)

Fátima Azucena Rasgado-Bonilla^{a,*}, Marcos Soto-Hernández^a, Jorge Cadena-Iñiguez^b, Ahtziri Socorro Carranza-Aranda^c, Sara Elisa Herrera-Rodríguez^c

^a Colegio de Postgraduados. Posgrado en Recursos Genéticos y Productividad, Fisiología Vegetal. Carretera Mexico- Texcoco Km. 36.5, Montecillo, Texcoco, Estado de Mexico 56230, Mexico

^b Colegio de Postgraduados. Posgrado en Inovación en Manejo de Recursos Naturales, Salinas de Hidalgo, San Luis Potosí 78620, Mexico

^c Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A. C., Subsede Yucatán, Tablaje Catastral 31264 Km 5.5 Carretera Sierra Papacal-Chuburna Puerto Parque Científico Tecnológico de Yucatán, Mérida, Yucatán 97302, México

Sechium edule (chayote) is a species with a wide distribution in the world and Mexico is the center of origin and domestication (Newstron, 1991). In Mexico, *S. edule* is known to have cytotoxic and anticancer potential (Soto-Hernández et al., 2016), even hybrids and wild relatives have this characteristic (Aguñiga-Sánchez et al., 2015). Within the genetic improvement of chayote, genotypes with bitter taste are associated with a high content of saponins, which are associated with antineoplastic processes (Koczurkiewicz et al., 2015). As part of the phytochemical characterization of the different *S. edule* genotypes obtained by hybridization, our research evaluated two segregating genotypes, 633-M11 and 387-M16. For this purpose, freeze-dried juice extracts and methanolic extracts of the pulp residue (bagasse) of fruits of both chayote genotypes were analyzed. The extraction of saponins from the lyophilized juice of 633-M11 was also carried out. Using HPLC-DAD, the presence of cucurbitacins (CuB, CuD, CuE, CuI, CuIIA), phenolic acids (β - resorcylic acid) and flavonoids (apigenin, catechin, rutin, naringenin and phloretin) were determined, and HPTLC was used to determine the content of total saponins. The juice and bagasse extracts of 633-M11 and 387-M16 showed significant cytotoxic activity on the prostate cancer cell line DU-145 in MTT assays. The lowest IC₅₀ of aqueous juice extract was of 633-M11 was 4.3 ± 1.2 $\mu\text{g/mL}$ while that 387-M16 was 0.47 ± 0.1 $\mu\text{g/mL}$ in the same type of extract. Bagasse extracts of both genotypes and that of 633-M11 total saponins showed cytotoxic effect on DU-145 cells, however, it was not as significant as that of juice extracts.

Keywords: *Sechium edule*, *Cucurbitaceae*, cancer, cytotoxicity, saponins

References

- Aguñiga-Sánchez, I., Soto-Hernández, R.M., Cadena-Iñiguez, J., Ruiz-Posadas, L.M., Cadena-Zamudio, J.D., González-Ugarte, A.K., Steider, B.W., Santiago-Osorio, E., 2015. Fruit extract from a *Sechium edule* hybrid induce apoptosis in leukaemic cell lines but not in Normal Cells. *Nutr Cancer*. 250-257.
- Koczurkiewicz, P., Czyz, J., Podolak, I., Wójcik K., Galanty, A., Janeczko, Z., Michalik, M., 2015. Multidirectional effects of triterpene saponins on cancer cells-minireview of *in vitro* studies. *Acta Biochim. Pol.* 62, 383-393.
- Newstron, L.E., 1991. Evidence for the origin of chayote *Sechium edule* (Cucurbitaceae). *Econ. Bot.*, 45, 410-428.
- Soto-Hernández, R.M., Ruiz-Posadas, L.M., Aguñiga-Sánchez, I., Santiago-Osorio, E., Arévalo-Galarza, M.L., Avendaño-Arrazate, C.H., Cisneros-Solano, V.M. y Cadena-Iñiguez, J. 2016. Uso de chayotes mexicanos para tratamiento de enfermedades de interés público. *Revista Agroproductividad*, suplemento. 51-53.

* Presenting author. E-mail address: msoto@colpos.mx (Marcos Soto-Hernández)

S2.P41 Studies *in vitro* and *in vivo* of anti-tumor activity of extracts derived from *Plukenetia volubilis* and *Moringa oleifera* in gastric tumor

Sara M. Robledo^{a,*}, Eyson Quiceno^b, Victoria Ospina^a, Natalia Arbelaez^b, Javier Murillo^b, Denny Cárdenas^c, Javier A. Soto^c

^a PECET-Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia

^b Grupo de Estudios Preclínicos, Corporación de Innovación CIDEPRO, Medellín, Colombia

^c Grupo BIOGEN, Universidad de Santander, Cúcuta, Colombia

Gastric cancer is the fifth most common cancer and the third most common cause of cancer death globally. Perioperative or adjuvant chemotherapy improves survival in patients with stage 1B or higher cancers. *M. oleifera* and *Plukenetia volubilis* are reported to enhance a broad range of biological functions including anti-tumor and antiproliferative activity (Anwar et al., 2007; Nascimento et al., 2013). With the aim of establishing a herbal extract for anti-tumor benefits, in this work we screened *in vitro* the effect of crude extracts (obtained with different solvents) from the leaves of the two plants and the seed oil from *P. volubilis* in the cytotoxicity, anti-proliferative activity, viability, cell cycle, apoptosis, membrane potential and cell migration of AGS (CRL-1739) tumor gastric cell line. The effect of the oil of *P. volubilis* in the progression of gastric tumor was also evaluated in a mouse model of gastric cancer in C57Bl/6 mice. All extracts evaluated were cytotoxic at concentrations (CC₅₀) between 90 and 160 µg/ml (Fig 1). The seed oil from *P. volubilis* showed 95% of mortality at 1% concentration (CC₅₀ = 0.86%). Cell proliferation was inhibited by 30%. At the cytotoxic dose, all extracts affected the cell cycle, observing an accumulation of AGS cells in the G0/G1 phase and induce DNA fragmentation as a mechanism of cell death. The ethanolic extract of *M. oleifera* leaf inhibits cell migration also.




Fig. 1. AGS cells exposed to cyclohexane extract of *P. volubilis* (a), cell culture medium (b) and dimethyl sulfoxide

The effect of the seed oil from *P. volubilis* in the progression of gastric tumors in C57Bl6 mice are under study.

Funding: Colombian Ministry of Science and Technology (Grant CT923-2019)

Keywords: Cell proliferation, cytotoxicity, apoptosis, DNA fragmentation, cell cycle

References

- Anwar F., Latif S., Ashraf M., Gilani A.H., 2007. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytother. Res.* 21, 17–25.
- Nascimento, A.K.L., Melo-Silveira, R.F., Dantas-Santos, N., Fernandes, J.M., Zucolotto, S.M., Rocha, H.A.O., Scortecci, K.C., 2013. Antioxidant and antiproliferative activities of leaf extracts from *Plukenetia volubilis* Linneo (Euphorbiaceae). *Evid. Based Complement. Alternative Med.* 2013, 1–10.

* Presenting author. E-mail address: sara.robledo@udea.edu.co (Sara M. Robledo)

S2.P42 Differential modulatory effects of *Populus balsamifera* (Salicaceae) and *Pinus Spp.* (Pinaceae) hydroethanolic extracts on IL-6 and TNF production by activated human peripheral blood mononuclear cells

Aaron Donny-Clark^{a,*}, Masa Sasagawa^d, Kaleb Lund^c, Anna Bernieri^a, Matthew Manniello^a, Mark R. Martzen^b, Cynthia A. Wenner^b

^a Doctorate of Natural Medicine Student, Bastyr University, Kenmore, Washington, USA

^b Bastyr University, Department of Basic Sciences, Kenmore, Washington, USA

^c Bastyr University, Department of Botanical Medicine, Kenmore, Washington, USA

^d Bastyr University, Bastyr University Research Institute, Kenmore, Washington, USA

Boswellia species are an example of plants whose harvest for western medicinal use is harmful to the species and has a high carbon footprint (Bongers, et al., 2019). This research contributes to the characterization of medicinal properties of local plants that can be sustainably harvested to reduce reliance on natural products requiring long-distance shipping. This study reports differential modulatory effects of commercially available extracts of *P. balsamifera* (Salicaceae) resinous buds and *Pinus Spp.* (Pinaceae) resin on human peripheral blood mononuclear cells (PBMC) stimulated without or with lipopolysaccharide (LPS). Both species are prolific in the United States, and harvest of their resins does not harm the trees. Both resins have documented use by indigenous peoples (Turner, 2014). PBMC were pre-treated with extracts at concentrations of 10-475 µg/mL for 2 hours. Cells were left unstimulated, or stimulated with 95 ng/mL LPS and incubated for 8-14 hours. Supernatants were harvested and IL-6 and TNF concentrations determined by ELISA. Cytotoxicity was determined by XTT assay.



Fig. 1. *Populus balsamifera*. K. Lund

Pinus spp. extract moderately decreased TNF and IL-6 release in a concentration-dependent manner. *P. balsamifera* extract induced significant TNF secretion beyond controls in a concentration-dependent manner and did not induce TNF in the absence of LPS. The TNF production enhancement is a novel finding, as previous studies have reported anti-inflammatory activity *in vitro* (e.g., Pannucci et al., 2022). Characterization of the extracts via HPLC, along with fractionation and identification of constituents responsible for the effects are warranted.

Keywords: *Populus balsamifera*, *Pinus Spp.*, cytokines, cytotoxicity, PBMC

References

- Bongers, F., Gorenendijk, P.L., et al. (2019). Frankincense in peril. *Nature Sustainability*, 2(7), 602-610.
- Pannucci, E., D'Eliseo, D., Ieri, F., Romani, A., Santi, L., Bernini, R., Sabatti, M., Velotti, F. (2022). Perspectives on *Populus spp.* (Salicaceae) bud extracts as antioxidant and anti-inflammatory agents. *Natural Product Research*, 36(6), 1648-1652.
- Turner, N. (2014). *Ancient pathways, ancestral knowledge: Ethnobotany and ecological wisdom of Indigenous peoples of northwestern North America*. McGill Queen's University Press.

* Presenting author. E-mail address: aaron.donny-clark@bastyr.edu (Aaron Donny-Clark)

S2.P43 Optimization of extraction conditions for salidroside isolated from *Acer tegmentosum*

Guijae Yoo*, In-Wook Choi, Sang Yoon Choi

Korea Food Research Institute, Wanju 55365, Republic of Korea

Salidroside is a major phenolic glycoside of *Acer tegmentosum* (Aceraceae) and known to be a hepatoprotective compound (Park et al., 2006; Kwon et al., 2011; Linlin et al., 2019). In this study, extraction conditions were optimized for maximum yield of salidroside from *A. tegmentosum*. For optimization, three extraction factors such as ethanol concentration (%), extraction temperature (°C), and solvent to material ratio (mg/mL) were tested and optimized for maximum yield of salidroside using response surface methodology (RSM). The optimal condition was obtained as an ethanol concentration of 53.4%, an extraction temperature at 67.11°C and a solvent to material ratio (mg/mL), 195.55 mg/mL. The salidroside yield under optimal conditions was found to be 1.59 mg/g dried samples, which were well-matched with the predicted value of 1.56 mg/g dried samples (Fig 1). These results will provide useful information about optimized extraction conditions for the development of salidroside as hepatoprotective therapeutics.




Fig. 1. HPLC chromatogram of *A. tegmentosum* extract prepared under optimal extraction conditions for salidroside.

Keywords: *Acer tegmentosum*, box–behnken design, response surface methodology, salidroside

References

- Park, K.M., Yang, M.C., Lee, K.H., Kim, K.R., Choi, S.U., Lee, K.R., 2006. Cytotoxic phenolic constituents of *Acer tegmentosum* Maxim. Arch. Pharm. Res. 29, 1086-1090.
- Kwon, D.J., Kim, J.K., Bae, Y.S., 2011. DPPH radical scavenging activity of phenolic compounds isolated from the stem wood of *Acer tegmentosum*. Mokchae Konghak 39, 104-112.
- Linlin, L., Shousheng, L., Quanjian, D., Yongning, X., 2019. Salidroside suppresses the metastasis of hepatocellular carcinoma cells by inhibiting the activation of the Notch1 signaling pathway. Mol. Med. Rep. 19, 4964-4972.

* Presenting author. E-mail address: gjyoo@kfri.re.kr (Guijae Yoo)

S2.P44 *Eleutherococcus senticosus* fruits intract - no toxicity in a balb/c mouse model

Filip Graczyk^a, Daniel Załuski^{a,*}, Rob Verpoorte^b

^a Department of Pharmaceutical Botany and Pharmacognosy, Nicolaus Copernicus University, Marie Skłodowska Curie 9, 85-094 Bydgoszcz, Poland

^b Natural Products Laboratory, IBL, Leiden University, Sylvius Laboratory, PO BOX 9505, 2300RA Leiden, the Netherlands

Eleutherococcus senticosus (Rupr. et Maxim.) Maxim. fruits and roots are used to treat immune- related diseases. Because of the overexploitation of the roots, the species is considered to be endangered therefore, the aerial parts of *E. senticosus* might be explored as a new sustainable source of compounds with an immunostimulative activity (Graczyk et al., 2021). This study was aimed to evaluate the safety profile (hepatotoxic, blood parameters) of the *E. senticosus* fruits intract to support the use of the fruits in folk medicine and in a modern phytotherapy in the EU's countries. Mice Balb/c have been treated for 5 days with water (group I) and the intract in doses 750 (group II) and 1500 (group III) mg/kg b.w. The body weight was measured every day, euthanasia on the eighth day and the activity of ALTL and ASTL was assayed. Additionally, the level of creatinine, urea and total protein as well as blood morphology were measured. There were no statistically significant differences in the body weight, ALT and AST activity between groups I, II and III, a slight decrease in urea level and total protein was observed. A decrease in total leukocytes count was observed (7.9 , 7.8 and 5.8×10^3 uL, resp.) with a percentage lymphocytes' increase (80.2, 81.8 and 82.6, resp.). Our observations justify the traditional use of the fruits and in the doses used, the intract is rather safe. On the basis of the results obtained, the dose 750 mg/kg b.w. is considered for the next research.

Keywords: *Eleutherococcus senticosus*, fruits, Balb/c mouse, toxicity

References

Graczyk, F., Orzechowska B., Franz D., Strzemski M., Verpoorte R., Załuski D., 2021. The intractum from the *Eleutherococcus senticosus* fruits affects the innate immunity in human leukocytes: From the ethnomedicinal use to contemporary evidence-based research. J. Ethnopharmacol. 268, 1-10.

* Presenting author. E-mail address: daniel.zaluski@cm.umk.pl (Daniel Załuski)

S2.P45 Potential molecular mechanisms of Silexan[®] identified by RNA-Sequencing in a mouse model of chronic social stress-induced excessive aversion sensitivity

Sabrina Weisenburger^{a,*}, Thomas Gantert^a, Hannes Sigrist^b, Christopher R. Pryce^b, Martin D. Lehner^a

^a Preclinical R&D, Dr. Willmar Schwabe GmbH & Co. KG, Willmar-Schwabe-Str.4, Karlsruhe, Germany

^b Preclinical Laboratory, Department of Psychiatry, Psychotherapy and Psychosomatics, University of Zurich, August Forel-Strasse 7, Zurich, Switzerland

Silexan[®] is a proprietary essential oil from *Lavandula angustifolia* approved for the treatment of temporary anxious mood. Silexan[®] has also shown antidepressant properties in a clinical study (Bartova et al., 2023). To further evaluate potential molecular mechanisms of Silexan[®] contributing to its activity we used a chronic social stress (CSS) mouse model inducing excessive Pavlovian aversion learning and memory, as a validated *in vivo* model for excessive aversion sensitivity. Silexan[®] (3 and 30 mg/kg per oral) dose-dependently reversed CSS-induced excessive Pavlovian aversion memory with similar efficacy as the SSRI escitalopram (3 mg/kg). Afterwards hippocampal RNA was isolated for RNA-Sequencing from the control (unstressed)-vehicle group, the CSS-vehicle group, the CSS-Silexan[®] 30 mg/kg group and the CSS-escitalopram group (n=5 per group). From 27,676 genes, 51 genes were both downregulated by CSS and upregulated by Silexan[®]. These included genes that, according to the literature, are altered in rodents following stress e.g. *Gucy1a2* (guanylate cyclase 1 soluble subunit alpha 2), *Ncam2* and *Ube3a*. Genes with a potential link to depression and anxiety in rodents and humans not affected by CSS but upregulated by Silexan[®] were e.g. *Hrtp1b*, *Pld5*, and *Scn5a*. For genes upregulated by CSS and downregulated by Silexan[®], no obvious association with depression and anxiety was identified in the literature. In summary, such RNA sequencing data might prove useful to establish first hypotheses on potential modes of action of Silexan[®] in the treatment of excessive aversion processing in stress-related psychiatric disorders such as anxiety and depressive disorders.

Keywords: Silexan[®], depression, hippocampal gene expression

References

Bartova, Lucie, et al., 2023. Beneficial effects of Silexan on co-occurring depressive symptoms in patients with subthreshold anxiety and anxiety disorders: randomized, placebo-controlled trials revisited. *European Archives of Psychiatry and Clinical Neuroscience* 273.1, 51-63.

* Presenting author. E-mail address: Sabrina.Weisenburger@schwabe.de (Sabrina Weisenburger)

S2.P46 The *Urtica dioica* root extract component in WS[®]1541 diminishes parameters of epithelial mesenchymal transition in human benign hyperplastic prostate epithelial cells

Carla Sens-Albert^a, Sabrina Weisenburger^{a,*}, Beatrix König^a, Martin Michel^b, Martin D. Lehner^a

^a Preclinical R&D, Dr. Willmar Schwabe GmbH & Co. KG, Willmar Schwabe-Str.4, 76227 Karlsruhe

^b Institut für Pharmakologie, Johannes Gutenberg Universität, Langenbeckstr. 1, 55131 Mainz

Epithelial mesenchymal transition (EMT) contributes to the pathogenesis of benign prostate enlargement (BPE). In a human hyperplastic prostate epithelial cell line (BPH-1), we evaluated the effect of the extract combination WS[®]1541 and its single components WS[®]1473 and WS[®]1031 on various EMT-parameters. WS[®]1541 is a proprietary combination of ethanolic extracts from *Sabal serrulata* fruits (WS[®]1473) and *Urtica dioica* roots (WS[®]1031) clinically used for treatment of lower urinary tract symptoms attributed to BPE. We treated BPH-1 cells with different concentrations of the three extracts and evaluated cell migration as a marker for EMT. The extract combination WS[®]1541 inhibited migration which was solely achieved by the presence of WS[®]1031, as WS[®]1473 had no effect on this parameter. Since TGF- β is a key driver of EMT, we examined the expression of the three TGF- β isoforms and its two receptors. WS[®]1541 diminished expression of all target genes. This effect was primarily mediated by the *Urtica dioica* root extract WS[®]1031, whereas the *Sabal serrulata* fruit extract WS[®]1473 only reduced expression of *TGFB2* and *TGFBR2*. Demonstrated by phalloidin staining, F-actin polymerization was not affected by either treatment. Inhibition of the migratory phenotype of BPH-1 cells by the combination product WS[®]1541 is conferred by the activity of the *Urtica dioica* root extract WS[®]1031. This effect may be due to a reduced expression of relevant TGF- β isoforms and their receptors. This mechanism could contribute to the clinical efficacy of WS[®]1541.

Keywords: WS[®]1473, benign prostate enlargement, *Urtica dioica*, *Sabal serrulata*, epithelial mesenchymal transition

* Presenting author. E-mail address: carla.sens-albert@schwabe.de (Carla Sens-Albert)

S2.P47 Antitumor activity of evodiamine analogs by targeting PKC β I in gemcitabine-resistant pancreatic cancer cells

Eun Seo Bae, Junhwa Hong, Yijae Lim, Woong Sub Byun, Simin Chun, Suckchang Hong, Sang Kook Lee*

College of Pharmacy, Seoul National University, Seoul 08826, Korea

Evodiamine, an indoloquinazolidine alkaloid, is a major component of the traditional herbal medicine *Evodia rutaecarpa* (Yu et al., 2013), which is widely used in Asian countries to alleviate vomiting, pain, and diarrhea. Evodiamine exerts various biological activities, including anti-inflammatory, anti-obesity, and antimicrobial effects. Accumulating evidences reported the antiproliferative and antitumor activities of evodiamine against various cancers through the metastasis inhibition, apoptosis induction, and cell cycle arrest. In cancer cells, protein kinase C β I (PKC β I) plays a crucial role in diverse cellular processes, including cell proliferation, invasion, and apoptotic pathways. In this present study, we created a scaffold to develop PKC β I inhibitors using evodiamine-based synthetic molecules.




Fig. 1. Structure of evodiamine

Based on various biological activities of evodiamine, the scaffold was synthesized and investigated for antiproliferative and antitumor activities against gemcitabine-resistant pancreatic cancer cell line PANC-1 (PANC-GR) cells. In addition, target validation experiment was employed to elucidate the molecular mechanism and verifying the target to overcoming gemcitabine resistance. Mechanistically, evodiamine analogs suppress the PKC β I expression, which subsequently inhibits the proliferation of PANC-GR cells *in vitro* and tumor growth in PANC-GR cells-implanted xenograft mouse model.

Keywords: *Evodia rutaecarpa*, synthetic analogs, antitumor activity, gemcitabine-resistant pancreatic cancer

References

- Yu, H., Jin, H., Gong, W., Wang, Z., Liang, H. 2013. Pharmacological actions of multi-target-directed evodiamine. *Molecules*, 18 (2), 1826-1843.
- Bae, E. S., Byun, W. S., Ock, C. W., Kim, W. K., Park, H. J., Lee, S. K. 2023. Periplocin exerts antitumor activity by regulating Nrf2-mediated signaling pathway in gemcitabine-resistant pancreatic cancer cells. *Biomed. Pharmacother.* 157, 114039.

* Presenting author. E-mail address: sklee61@snu.ac.kr (Sang Kook Lee)

S2.P48 Perceived changes in anxiety symptom burden during treatment with *Bryophyllum pinnatum*: a prospective, non-randomised, observational study

Tiffany Huber^{a,*}, Daniel Krüerke^a, Timotheus Haeck^a, Markus Schlemmer^a, Ana Paula Simões Wüst^{a,b}

^a Klinik Arlesheim, Arlesheim, Switzerland

^b Department of Obstetrics, University Hospital Zurich, University of Zurich, 8091 Zurich, Switzerland

Bryophyllum preparations are widely used in anthroposophic medicine, a form of integrative medicine. Prospective studies have revealed positive effects of treatment with *Bryophyllum* 50% tablets on sleep quality (Lambrigger-Steiner et al., 2014; Mirzayeva et al., 2023). These improvements are corroborated by preclinical studies showing that fractions from *B. pinnatum* leaves can prolong induced sleeping time in rodents, indicating a CNS depressant action (Pal et al., 2010). A recent study showed anxiolytic and psychoactive effects in young zebrafish (Martins Fernandes Pereira et al., 2022). We set to find out if patients perceive improvements of their anxiety symptoms during treatment with *Bryophyllum* 50% tablets. A total of 51 patients with anxiety symptoms were recruited among the waiting list from the Department of Psychiatry and Psychosomatics at the Klinik Arlesheim. During the three-week study, patients were treated with *Bryophyllum* 50 % tablets (3x2 per day) and filled out a questionnaire at baseline, after two and three weeks of tablet intake. The primary endpoint of the study was the change in anxiety symptoms measured with the Beck Anxiety Inventory (BAI). Depression, sleep quality, stress, health-related quality of life and internal coherence were assessed as well. Results show a statistically significant and clinically relevant decrease in the BAI-score between baseline and after two (and three) weeks of treatment. Additional improvements were observed in the secondary endpoints; tolerability and compliance were very good. We conclude that *Bryophyllum* 50% tablets have beneficial effects on anxiety-related symptoms and should be considered as a well-tolerated treatment for their management.

Keywords: *Bryophyllum pinnatum*, anxiety symptoms, Beck Anxiety Inventory, sleep quality

References

- Lambrigger-Steiner, C., Simões-Wüst, A.P., Kuck, A., Fürer, K., Hamburger, M., von Mandach, U. 2014. Sleep quality in pregnancy during treatment with *Bryophyllum pinnatum*: An observational study. *Phytomedicine* 21(5), 753–757.
- Martins Fernandes Pereira, K., Calheiros de Carvalho, A., André Moura Veiga, T., Melgoza, A., Bonne Hernández, R., Dos Santos Grecco, S., Uchiyama Nakamura, M., Guo, S. 2022. The psychoactive effects of *Bryophyllum pinnatum* (Lam.) Oken leaves in young zebrafish. *PloS One*, 17(3).
- Mirzayeva, N., Forst, S., Passweg, D., Geissbühler, V., Simões-Wüst, A.P., Betschart, C. 2023. *Bryophyllum pinnatum* and Improvement of Nocturia and Sleep Quality in Women: A Multicentre, Nonrandomised Prospective Trial. *Evidence-Based Complementary and Alternative Medicine*, 2023.
- Pal, S., Sen, T., Chaudhuri, A.K.N. 2010. Neuropsychopharmacological Profile of the Methanolic Fraction of *Bryophyllum pinnatum* Leaf Extract. *Journal of Pharmacy and Pharmacology* 51(3), 313–318.

* Presenting author. E-mail address: tiffany.huber@klinik-arlesheim.ch (Tiffany Huber)

S2.P49 Effects of orally applied sanggenons C and D in BALB/c mice

Julia Langeder^{a,b}, Mirijam Koch^c, Hannes Schmietendorf^d, Ammar Tahir^b, Judith M. Rollinger^b, Michaela Schmidtke^d, Ulrike Grienke^{b,*}

^a University of Vienna, Vienna Doctoral School of Pharmaceutical, Nutritional and Sport Sciences, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^b University of Vienna, Department of Pharmaceutical Sciences, Division of Pharmacognosy, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^c Jena University Hospital, Department of Medical Microbiology, Jena, Germany

^d Jena University Hospital, Department of Medical Microbiology, Section of Experimental Virology, Jena, Germany

As part of our continuing search for bioactive natural products, a standardized root bark extract (MA60) from *Morus alba* revealed as promising agent. *In vitro* activities against inflammation, influenza viruses, SARS-CoV-2, *S. aureus*, and *S. pneumoniae* were previously determined (Grienke et al., 2016; Langeder et al., 2023a; Wasilewicz et al., 2023). Major active constituents of MA60 are the sanggenons C and D. This presentation will cover a preclinical study conducted in female BALB/c mice to evaluate whether *in vitro* results are transferable to an *in vivo* setting. Oral application of MA60 was selected to determine i) the maximum tolerated dose (MTD) in an acute toxicity study and ii) its potential against influenza virus in an efficacy study. Moreover, inflammatory parameters were monitored. A further aim was to evaluate whether there is a correlation between the obtained results and the amount of sanggenons C and D in serum and tissues by quantitation using a validated UPLC-ESI-MS method. The MTD was reached at 100 mg/kg. In the efficacy study, the treatment effects were moderate. Dose-dependent quantities of sanggenon C in serum and sanggenon D in liver samples were detected. Only very low concentrations of sanggenons C and D were determined in lung samples and none of these compounds was found in spleen samples. There was no compound accumulation when MA60 was administered repeatedly (Langeder et al., 2023b). In conclusion, due to low serum concentrations after oral application once daily, alternative application routes, in particular inhalation or intranasal administration are currently pursued.

Funding: Wilhelm-Doerenkamp-Foundation, Switzerland (NATVANTAGE Grant 2018); Austrian Science Fund (FWF P34028 and P35115)

Keywords: *Morus alba*, sanggenon, influenza virus, *in vivo*, quantitation

References

- Grienke, U., Richter, M., Walther, E., Hoffmann, A., Kirchmair, J., Makarov, V., Nietzsche, S., Schmidtke, M., Rollinger, J.M., 2016. Discovery of prenylated flavonoids with dual activity against influenza virus and *Streptococcus pneumoniae*. *Sci. Rep.* 6, 27156.
- Langeder, J., Koch, M., Schmietendorf, H., Tahir, A., Grienke, U., Rollinger, J.M., Schmidtke, M., 2023b. Correlation of bioactive marker compounds of an orally applied *Morus alba* root bark extract with toxicity and efficacy in BALB/c mice. *Front. Pharmacol.* 14:1193118.
- Langeder, J., Döring, K., Schmietendorf, H., Grienke, U., Schmidtke, M., Rollinger, J.M., 2023a. ¹H NMR-based chemometric analysis of *Morus alba* extracts toward a multipotent herbal anti-infective. *J. Nat. Prod.* 86, 8-17.
- Wasilewicz, A., Kirchweiger, B., Bojkova, D., Jose Abi Saad, M., Langeder, J., Bütikofer, M., Adelsberger, S., Grienke, U., Cinatl Jr., J., Petermann, O., Scapozza, L., Orts, J., Kirchmair, J., Rabenau, H.F., Rollinger, J.M., 2023. Identification of natural products inhibiting SARS-CoV-2 by targeting viral proteases: A combined *in silico* and *in vitro* approach. *J. Nat. Prod.* 86, 264-275.

* Presenting author. E-mail address: ulrike.grienke@univie.ac.at (Ulrike Grienke)

S2.P50 Novel natural product inhibitors targeting oncogenic MAPK/ERK and PI3K/AKT signaling in melanoma: From large library screening to target identification

Maria Karpouchtsi^{a,*}, Lara Dürr^a, Maciej Dobrzynski^b, Silke Radetzki^c, Tanja Hell^a, Matthias Hamburger^a, von Jens-Peter von Kries^c, Olivier Pertz^b, Robin Teufel^a, Eliane Garo^a

^a Division of Pharmaceutical Biology, Department of Pharmaceutical Sciences, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

^b Institute of Cell Biology, University of Bern, Baltzerstrasse 4, 3012 Bern, Switzerland

^c Leibniz-Forschungsinstitut für Molekulare Pharmakologie, Robert-Rössle-Straße 10, 13125 Berlin, Germany

Malignant melanoma is the deadliest type of skin cancer with unmatched mutation rates arising in both MAPK/ERK and PI3K/AKT signaling pathways. Although specific inhibitors of these critical pathways show spectacular initial results, most patients relapse within just a few months. Combination therapy can improve overall survival rates, but the currently available options are limited (Tanda et al., 2020). Therefore, novel inhibitors targeting oncogenic ERK and AKT signaling in melanoma are urgently needed. To tackle this issue, our in-house library of crude plant extracts was combined with an innovative high-content screen (HCS) that quantifies downstream inhibitory activity at ERK and AKT level. HPLC-based activity profiling of the active hits and subsequent targeted isolation of the bioactive constituents was performed (Dürr et al., 2022). To further explore the coverage of chemical space for such inhibitors, we also screened large pure compound libraries by accessing high-throughput screening pipelines through an EU- OPENSREEN program. To this end, we screened 2,576 plant extracts and additional 25,696 pure natural and synthetic compounds. A total of 46 active compounds were confirmed as downstream inhibitors of ERK and/or AKT with IC₅₀ values in the low micromolar range. The current challenge aims towards target identification of the most promising hits. Several key kinases of the ERK-AKT network were produced through different cloning techniques and heterologous expression systems. Our strategy further includes the assessment of physical binding as well as enzymatic inhibition. Ultimately, we envisage to develop these newly discovered inhibitors into lead compounds for future drug development.

Keywords: melanoma, high-content screening, EU-OPENSREEN, MAPK/ERK and PI3K/AKT signaling, target identification

References

- Tanda, E.T., Vanni, I., Boutros, A., Andreotti, V., Bruno, W., Ghiorzo, P., Spagnolo, F., 2020. Current state of target treatment in BRAF mutated melanoma. *Front. Mol. Biosci.* 7, 154.
- Dürr, L., Hell, T., Dobrzynski, M., Mattei, A., John, A., Augsburger, N., Bradanini, G., Reinhardt, J.K., Rossberg, F., Drobnjakovic, M., Gupta, M.P., Hamburger, M., Pertz, O., Garo, E., 2022. High-content screening pipeline for natural products targeting oncogenic signaling in melanoma. *J. Nat. Prod.* 85, 4, 1006–1017.

* Presenting author. E-mail address: maria.karpouchtsi@unibas.ch (Maria Karpouchtsi)

S2.P51 Screening for natural products inhibiting PI3K/AKT pathways in melanoma: investigation of *Mammea americana* EtOAc extract

Ruzhe Han^{a,*}, Tamara Balsiger^a, Maciej Dobrzyński^b, Jakob K. Reinhardt^a, Lara Dürr^a, Tanja Hell^a, Pablo Solis^c, Matthias Hamburger^a, Olivier Pertz^b, Robin Teufel^a, Eliane Garo^a

^a Division of Pharmaceutical Biology, Department of Pharmaceutical Sciences, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

^b Institute of Cell Biology, University of Bern, Baltzerstrasse 4, 3012 Bern, Switzerland

^c Center for Pharmacognostic research and Panamanian Flora (CIFLORPAN), Faculty of Pharmacy, University of Panama, 0801 Panama City, Republic of Panama

Malignant melanoma is the deadliest skin cancer and its incidence has risen significantly over the past decades. Mutations are frequently observed in the PI3K/AKT and MAPK/ERK pathways, leading to abnormal cell proliferation. Targeted therapy showed spectacular initial results, but drug resistance appears after a few months only. Combination therapy has shown improvement in the progression-free and overall patient survival and additional compounds are urgently needed to complete the arsenal of available drugs. To address this issue, our natural product discovery platform was combined with an innovated high-content screening (HCS) assay to quantify AKT and ERK signaling in melanoma cells (Dürr et al., 2022). Here, the EtOAc extract from *Mammea americana* inhibited the PI3/AKT signaling pathway and was therefore selected for scaled-up isolation of the active constituents. To this end, an HPLC-based activity profiling approach was used, which led to the identification of novel betulinic acid derivatives (**1-10**) and known coumarins (**11-19**). Upon closer inspection, the coumarin theraphin B (**15**), was the only compound displaying a weak activity with an IC₅₀ of $37 \pm 5.9 \mu\text{M}$. In contrast, although betulinic acid has been reported to have anti-tumor activity in mice (Pisha et al., 1995), the analogues **1-10** did not show any inhibition of AKT and ERK, suggesting a different mechanism unrelated to the AKT signaling pathway to explain the in vivo activity.

Keywords: *Mammea americana*, melanoma, high-content screening, coumarin, ECD

References

- Dürr, L., Hell T., Dobrzyński, M., Mattei, A., John, A., Augsbürger, N., Bradanini, G., Reinhardt, J. K., Rossberg, F., Drobnjakovic, M., Gupta, M. P., Hamburger, M., Pertz, O., Garo, E., 2022. High-content screening pipeline for natural products targeting oncogenic signaling in melanoma. *Journal of Natural Products*. 185(4), 1006-1017.
- Pisha, E., Chai, H., Lee, I., Chagwedera, T.E., Farnsworth, N.R., Cordell, G.A., Beecher, C.W.W., Fong, H.H.S., Kinghorn, A.D., Brown, D.M., Wani, M.C., Wall, M.E., Hieken, T.J., Gupta, T.K.D., Pezzuto, J.M., 1995 Discovery of betulinic acid as a selective inhibitor of human melanoma that functions by induction of apoptosis. *Nature Medicine*. 1, 1046-1051

* Presenting author. E-mail address: ruzhe.han@unibas.ch (Ruzhe Han)

S2.P52 Exploring the anti-inflammatory potential of an herbal mixture preparation and individual herbal extracts on inflammatory bowel disease

Maria-Riera Piqué-Borràs^a, Johann Röhl^a, Mónica Mennet-von Eiff^b, Gerald Künstle^{a,*}

^a Weleda AG, Preclinical Research and Development, Dychweg 14, 4144 Arlesheim, Switzerland

^b Weleda AG, Research and Development Pharma, Dychweg 14, 4144 Arlesheim, Switzerland

Patients affected by inflammatory bowel disease (IBD) often suffer of compromised motility due to mucosal inflammation. The herbal mixture preparation found in Digestodoron[®] is traditionally used for treatment of gastrointestinal dysmotility disorders. This study aimed to investigate the anti-inflammatory properties of the herbal mixture and its four individual extracts—*Dryopteris filix-mas* (DF), *Phyllitis scolopendrium* (PS), *Polypodium vulgare* (PV) and *Salix mixture with alba/purpurea/viminalis* (SM)). Dry extracts were derived from Digestodoron[®] or from the individual ethanolic tinctures. Pro-inflammatory leukotrienes and prostaglandins, as well as the activation of the NF-κB pathway, are hallmark indicators of inflammation. NF-κB activation was evaluated *via* a reporter assay using the human Jurkat cell line. Inhibition analysis of the pain-related enzymes 5-lipoxygenase (5-LO) and cyclooxygenase-2 (COX-2) as part of the arachidonic acid pathway was performed using recombinant human enzymes. The whole extract, along with individual extracts PS and SM, notably suppressed 5-LO enzyme (IC₅₀: 12.6, 60.6, 16.6 µg/mL, respectively). In contrast, PV and DF showed comparatively lower inhibition effects (IC₅₀: 223.8 and 648.6 µg/mL, respectively). In terms of COX-2 enzyme suppression, the whole and all individual extracts exhibited effectiveness (IC₅₀: 12.6 (mixture), 25.5 (DF), 14.3 (PS), 7.9 (PV) and 2.7 (SM) µg/mL). Moreover, NF-κB activation was hindered by the whole extract and exclusively by the SM extract (IC₅₀: 223.1 and 79.6 µg/mL). These findings suggest that Digestodoron[®], with its potent anti-inflammatory and analgesic properties, could be a promising therapeutic option for individuals suffering from IBD. Ongoing research aims to elucidate further the potential therapeutic benefits of Digestodoron[®].

Keywords: herbal preparation, NF-κB, 5-lipoxygenase, cyclooxygenase-2, anti-inflammatory, inflammatory bowel disease

* Presenting author. E-mail address: preclinical-dev@weleda.ch (Gerald Künstle)

S2.P54 Extracts of *Sisymbrium officinalis* and *Plantago lanceolata* reduced inflammation induced by lipoteichoic acid from *Streptococcus pyogenes* in human tonsil epithelial cells

Olumayokun A. Olajide^{a,*}, Uzeme P. Aluta^a, Thomas Hallett^b

^a Department of Pharmacy, School of Applied Sciences, University of Huddersfield, Huddersfield, HD1 3DH, UK

^b Reckitt Benckiser Health LTD, Dansom lane, Hull, HU8 7DS, UK

Pharyngitis is an inflammation of the pharynx caused by viral, bacterial, or non-infectious triggers. *Sisymbrium officinalis* (L.) Scop (SO) and *Plantago lanceolata* L. (PL) leaf extracts are used in traditional medicine for the treatment of sore throat and inflammation of the upper respiratory tract. In this study, the anti-inflammatory effects of SO and PL dry extracts were evaluated in an *in vitro* model of streptococcal pharyngitis. Human primary tonsil epithelial cells were stimulated with lipoteichoic acid (LTA) from *Streptococcus pyogenes* (10 mg/mL) in the presence or absence of SO (12.5, 25 and 50 mg/mL). In similar experiments, tonsil cells were treated with PL (25, 50 and 100 mg/mL) prior to LTA stimulation. Culture supernatants were collected after 24 h and analysed for levels of PGE₂ using enzyme immunoassay (EIA) kit. Secretion of pro-inflammatory cytokines (TNF α and IL-6) were measured using ELISA. Results showed that pre-treatment of tonsil epithelial cells with SO produced significant ($p < 0.05$) and concentration-dependent reduction in LTA-induced production of PGE₂, TNF α and IL-6. The highest concentration of the extract reduced inflammatory mediator production by ~60%. Results also showed that PL significantly ($p < 0.05$) reduced LTA-induced increased production of PGE₂, TNF α and IL-6, with 100 mg/mL of the extract reducing elevated production of inflammatory mediators by ~70%. Cell viability experiments revealed no significant reduction in viability of tonsil epithelial cells at concentrations of extracts investigated. These results suggest that *Sisymbrium officinalis* and *Plantago lanceolata* are anti-inflammatory in pharyngitis caused by *Streptococcus pyogenes* and should be further investigated for healthcare applications.

Keywords: inflammation, pharyngitis, upper respiratory tract infections, *P. lanceolata*, *S. officinalis*

* Presenting author. E-mail address: O.A.Olajide@hud.ac.uk (Olumayokun A Olajide)

S2.P55 Protein kinase activity of chromatographic fractions of *Acacia auriculiformis* stem bark as potentials for anti-cancer drug discovery

Florence Tarfa^a, Aniefiok Udobre^b, Augustine A, Ahmaduc^{3,*}

^a Department of Medicinal Chemistry and Quality Control, National Institute for Pharmaceutical Research and Development, Idu, Abuja, Nigeria

^b Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, University of Uyo, Uyo, Nigeria

^c Department of Pharmaceutical and Medicinal Chemistry, Faculty of Pharmaceutical Sciences, Veritas University, Bwari Abuja, Nigeria

The incidence of deaths from cancer has been on the rise globally with reported cases of 10 million deaths in 2020 that is about one in six deaths and over one million new cases occur each year in Africa, with about 700,000 deaths reported. Data estimates showed a considerable increase in cancer mortality to nearly one million deaths per year by 2030. As a result of the negative side effects of the current treatment regimen plan which includes chemotherapy or a combination of radiotherapy and chemotherapy, this has justified the continuous search for new therapeutic agents from natural sources including medicinal plants. In this present work, we report the inhibitory activity of the chromatographic fractions of the ethyl acetate soluble of the stem bark extract of *Acacia auriculiformis* against some disease related protein kinases. The ethyl acetate fraction was subjected to chromatographic separation on silica gel and sephadex LH-20 to afford H1, H2 and H3 respectively. H1 was identified as Betulin using spectroscopic technique, while H2 and H3 gave positive test for flavonoid. H1-H3 were then subjected to kinase enzymatic assays performed using the ADP Glo assay kit in a 384- well plates. The result of the primary screening showed that H1 and H2 inhibited four and six of the kinases tested, while the best activity was exhibited by H3 against all the kinases tested. The result of the IC₅₀ revealed similar trend as H3 showed the best activity out of the three fractions with an IC₅₀ of 0.086 µg/mL against Haspin and 0.46 µg/mL against CDK 9 kinases, while H2 gave an IC₅₀ of 0.71 against CDK 9 but was inactive against the haspin kinase. The kinase activity of H2 and H3 were more than H1 (Betulin) previously reported as a multi target kinase inhibitor suggesting the potentials of H3 as a lead in the discovery of haspin and CDK 9 inhibitors which can be exploited in the search for anticancer drug.

Keywords: *Acacia auriculiformis*, chromatographic fractions, kinase activity, IC₅₀

References

- Calixto, J.B., 2000. Efficacy, safety, quality control and regulatory guidelines for herbal medicines (Pharmacotherapeutic agents). Braz. J. Med. Biol. 33, 179-189
- Rates, S.M., 2001. Plants as source of drugs. Toxicon. 39, 603-613 WHO (2020) Report, Geneva, Switzerland.

* Presenting author. E-mail address: e-mail: ahmadu2001@yahoo.com (Augustine A Ahmaduc)

S2.P56 Comparative protein kinase inhibitory activity of three *Acacia* species growing in Nigeria

Augustine A Ahmadu^{a,*}, Bilqis Abiola Lawal^b, Florence Tarfa^c, Patricia Odumosu^d

^a Department of Pharmaceutical and Medicinal Chemistry, Veritas University, Bwari-Abuja, Nigeria

^b Department of Pharmacognosy and drug development, University of Ilorin, Faculty of Pharm. Sciences Ilorin Nigeria

^c Department Medicinal Chemistry and quality control, National Institute for Pharmaceutical Research and development, Idu-Abuja, Nigeria

^d Department of Pharmaceutical Chemistry, Faculty of Pharm. Sciences, University of Jos, Nigeria

Significant improvements have been made in the field of drug discovery targets against cancer over the years, which have necessitated the search for novel chemical entities for the therapeutic treatment and management of cancer from natural sources. In this present work, we report herein the kinase inhibitory effect of the crude chloroform extract and the partitioned fractions of the hydroalcoholic stem bark extract of *Acacia ataxacantha* against a panel of disease related protein kinases and the comparison of the activity with that of *Acacia nilotica* and *Acacia auriculiformis*. The extract and the fractions of the stem bark of *Acacia ataxacantha* were subjected to Kinase enzymatic activities performed in 384-well plates using the ADP-Glo assay kit. The result showed that chloroform extract was inactive, while the ethyl acetate fraction gave the highest activity against haspin kinase with IC₅₀ of 0.1 µg/mL, followed by aurora B kinase with IC₅₀ of 0.45 µg/mL, while against CDK2, CDK5 and CDK9 kinases, the IC₅₀ were 0.6, 0.88 and 0.9 µg/mL respectively. The n-butanol fraction gave an IC₅₀ of 0.59 µg/mL and 0.7µg/mL against haspin and aurora B Kinases respectively. Comparison with *Acacia nilotica* and *Acacia auriculiformis* showed that the ethyl acetate fraction of *acacia ataxacantha* is the most active. The results showed the potentials of the ethyl acetate fraction of *Acacia ataxacantha* as source of inhibitors against Haspin, aurora B and CDK kinases which might serve as lead for antimitotic agent against cancer.

Keywords: *Acacia* species, protein kinases, ethyl acetate fraction, IC₅₀, anticancer agent

References

- Calixto, J.B., 2000. Efficacy, safety, quality control, marketing and regulatory guidelines for herbal medicines (phytotherapeutic agents). Braz. J. Med. Biol. 33, 179-189.
- Newmann, D.J., Cragg, G.M., 2016. Natural products as sources of new drugs from 1981-2014. J. Nat. Prod. 79, 629- 661.
- Siveen, K.S., Prabhu, K.S., Achkar, I. W., Kuttikrishnan, S., Shyam, S., et al., 2018. Role of non-receptor tyrosine kinases in hematological malignances and its targeting by natural products. Mol. Cancer 17(1), 1-21.

* Presenting author. E-mail address: ahmadu2001@yahoo.com (Augustine A Ahmadu)

S2.P57 Diversification of kratom alkaloids using biocatalysis for opioid use disorders

Sahar Amin^{a,b,*}, Hannah Boesger^{a,b}, Natalia R. Harris^a, Kayla Kroning^a, Harrison Snodgrass^c, Wenjing Wang^a, Jared Lewis^c, David H. Sherman^{a,b}

^a Life Sciences Institute, University of Michigan, Ann Arbor Michigan, USA

^b Department of Medicinal Chemistry, University of Michigan, Ann Arbor, Michigan 48109, USA

^c Department of Chemistry, Indiana University, Bloomington, Indiana, USA

This presentation will cover the diversification of the main indole alkaloids in the kratom plant (Fig. 1). Mitragynine, a plant-derived alkaloid natural product, has captured attention due to its intriguing analgesic properties, positioning it as a promising candidate for a safer alternative to existing mu- opioid receptor (MOR) agonists (Kruegel et al., 2016). We are exploring the creation of new analogs of mitragynine through late-stage hydroxylation and halogenation, which enables further interrogation of chemical space through additional functionalization. Two complementary strategies to identify and optimize biocatalysts capable of site-selectively modifying mitragynine have been employed. First, we are utilizing directed evolution on the 4V flavin dependent halogenase to enhance chlorination of mitragynine at variant positions on the indole ring. Directed evolution is being conducted to generate a diverse library of enzyme variants with improved activity and selectivity. Secondly, computational approaches, including density functional theory (DFT), inverse docking, and molecular dynamics (MD) simulations have been incorporated into our workflow to identify novel cytochrome P450s capable of late-stage C-H functionalization. Finally, the bioactivity of our analogs at the MOR will be assessed through M-SPOTIT and PathHunter β - galactosidase enzyme-complementation assays, providing detailed insights into their signaling and potential side effects (Kroning et al., 2021). We plan to investigate additional bioactive kratom alkaloids like speciogynine, paynantheine, and corynoxine for their synergistic potential. Through this comprehensive approach we strive to develop novel, improved mitragynine and related compound derivatives with improved analgesic properties.




Fig. 1. Overview of strategy for kratom alkaloid structural diversification

Keywords: biocatalysis, C-H functionalization, mitragynine, MOR, directed evolution

References:

- Kruegel, A.C., Gassaway, M.M., Kapoor, A., Váradi, A., Susruta Majumdar, S., Filizola, M., Javitch, J.A. Sames D., 2016. Synthetic and receptor signaling explorations of the mitragyna alkaloids: mitragynine as an atypical molecular framework for opioid receptor modulators J. Am. Chem. Soc. 138 (21), 6754-6764.
- Kroning, K.E., Wang, W., 2021. Designing a single protein-chain reporter for opioid detection at cellular resolution. Angew. Chem. 60, 13358-13365.

* Presenting authors E-mail address: saminn@umich.edu (Sahar Amin), hboesger@umich.edu (Hannah Boesger)

S2.P58 Study of *Carpesium abrotanoides* in dextran sulfate sodium-induced IBD and Caco-2 cells

So Yeon Kim*, Yun Mi Lee, Dong-Seon Kim

KM Science Research Division, Korea Institute of Oriental Medicine, 1672 Yuseongdae-ro, Yuseong-gu, Daejeon 34054, Republic of Korea

The genus *Carpesium* consists of approximately 21 species worldwide (Zhang et al., 2015). This species is particularly widespread in Asia and is used as a traditional treatment for various conditions such as colds, fever, insects, bruises, and inflammation (Zhang et al., 2015). Among them, research results have reported that *Carpesium abrotanoides* species has excellent anti-inflammatory, anti-cancer, and antioxidant effects (Lee et al., 2013). However, no studies have yet been reported on the effects of this natural product on inflammatory bowel disease. Therefore, in this study, we investigated the protective effects of *C. abrotanoids* on dextran sulfate sodium (DSS)-induced experimental colitis. Using an acute colitis model *in vivo*, we identified the possible mechanisms of anti-inflammatory activity and *C. abrotanoids* in C57BL/6 mice. *In vitro*, further studies were conducted to identify the molecular mechanisms of *C. abrotanoids* for the inflammatory response using Caco-2, a human intestinal epithelial cell. As a result, in C57BL/6 mice, *C. abrotanoids* alleviated disease activity index (DAI), colon length shortening, and inhibited pro-inflammatory cytokine production in blood and tissue. In addition, *C. abrotanoids* significantly reduced the secretion of pro-inflammatory cytokines such as TNF- α and IL-6 at LPS-induced Caco-2 cell levels. This study suggests that *C. abrotanoids* could be a potential agent for the treatment of inflammatory bowel disease.

Keywords: *Carpesium abrotanoides*, dextran sulfate sodium, inflammatory bowel disease, caco-2 cell, anti-inflammatory effect

References

- Xu, Y., Lu, Y., 2019. Noteworthy plants in Asteraceae from China. J. Zhejiang Univ. 46, 209–214.
- Zhang, J.P., Wang, G.W., Tian, X.H., Yang, Y.X., Liu, Q.X., Chen, L.P., Li, H.L., 2015. The genus *Carpesium*: A review of its ethnopharmacology, phytochemistry and pharmacology. J. Ethnopharmacol. 163, 173–191.
- Lee, J.H., Hwang, K.H., Kim, G.H., 2013. *In vitro* evaluation of anti-inflammatory activity for salad-food material *Carpesium abrotanoides*. J. Food Biochem. 37, 18–25.

* Presenting author. E-mail address: kimsy0913@kiom.re.kr (So Yeon Kim)

S2.P59 *In vitro* co-culture analysis of Caribbean medicinal plant extracts for bacterial vaginosis

Ella Vardeman^{a,b,c,*}, Ina Vandebroek^{b,c,d}, Edward J. Kennelly^{a,b}

^a Department of Biological Sciences, Lehman College, City University of New York, Bronx, NY 10468, USA

^b PhD Program in Biology, The Graduate Center, City University of New York, NY 10016, USA

^c The New York Botanical Garden, Bronx, NY 10458, USA

^d Department of Life Sciences and Caribbean Centre for Research in Bioscience (CCRIB), Faculty of Science and Technology, The University of the West Indies Mona, Kingston 7, Jamaica, West Indies

Bacterial vaginosis (BV) is the most common gynecological infection in reproductive-age women globally. BV is directly caused by dysbiosis of the vaginal microbiome – when beneficial *Lactobacillus* species are no longer dominant and are replaced by harmful anaerobic bacteria such as *Gardnerella vaginalis*. In Caribbean cultures, women use plants topically, such as *Argemone mexicana*, to treat BV and other vaginal infections. Until now, there has been very little research into how botanical extracts affect both beneficial and harmful bacteria in the vaginal microbiota, especially as these are often prepared in different ways for the same health condition. The aim of this study is to test the effect of botanical preparations using an *in vitro* co-culture assay with beneficial *Lactobacillus* species and BV-causing *G. vaginalis*. We hypothesized that chemical differences across five preparations would influence the ratio of beneficial and pathogenic bacteria. The different *A. mexicana* extractions (methanol, ethanol, water, acidified water, and heated water) were tested using an *in vitro* co-culture method with *G. vaginalis* and one of three vaginal *Lactobacillus* species. Additionally, UPLC-TQD-MS was used to quantify known antibacterial alkaloids, including berberine, protopine, and allocryptopine, across different extraction methods. Water extractions that had no antibacterial effect in monoculture suppressed the growth of *G. vaginalis* in co-culture with *Lactobacillus*. The results of this study provide practical information for women who use *A. mexicana* globally.

Keywords: vaginal microbiota, ethnobotany, women's health, caribbean medicinal plants

* Presenting author. E-mail address: evardeman@gradcenter.cuny.edu (Ella T. Vardeman)

S2.P60 Pharmacokinetic dynamics of the hybrid *Sechium* H387 07 and phytochemical characterization of its fruit

Marcos Soto-Hernández^{a,*}, Eugenia Elisa Delgado-Tiburcio^a, Itzen Aguiñiga-Sánchez^b, Jorge Cadena-Iñiguez^c, Lucero del Mar Ruiz-Posadas^a, Cecilia Beatriz Peña-Valdivia^a

^a Botany Department, Postgraduate College, Campus Montecillo, km 36.5 Carretera México-Texcoco, Texcoco 56230, Mexico

^b Hematopoiesis and Leukemia Laboratory, Research Unit on Cell Differentiation and Cancer, FES Zaragoza, National Autonomous University of Mexico, Mexico City 09230, Mexico

^c Innovation in Natural Resource Management, Postgraduate College, Campus San Luis Potosí, Salinas de Hidalgo, San Luis Potosí 78622, Mexico

Recently, a hybrid of *Sechium edule* named H387 07 has demonstrated potential as an antiproliferative, cytotoxic, and apoptotic agent in cancer cell lines, including leukemia and breast cancer. This effect is primarily attributed to the phytochemical content of phenols, flavonoids, and cucurbitacins present in the extract (Aguiñiga-Sánchez et al., 2015; Cadena-Iñiguez et al., 2022). This study aimed to assess the phytochemical content of the hybrid *Sechium* H387 07 and compare it with its segregants, 387M20 and 387M16. Additionally, the study sought to determine its antioxidant activity. On the other hand, the pharmacokinetic profile of cucurbitacins, phenolic acids, and flavonoids was investigated following the administration of the H387 07 hybrid extract at varying doses to mice. The obtained results indicated that segregant 387M16 showed a concentration of phytochemicals comparable to that of the hybrid *Sechium* H387 07. Consequently, 387M16 emerges as a promising candidate for subsequent bioassays. Furthermore, various pharmacokinetic parameters for apigenin, florentin, CuB, CuE, and CuI were determined. Notably, the study highlighted the presence of CuB and CuIIA due to their elevated concentration in the serum, with the latter being described for the first time in the extract of the *Sechium* hybrid.

Keywords: Pharmacokinetics, *Sechium*, hybrid, secondary metabolites

References

Aguiñiga-Sánchez, I., Soto-Hernández, M., Cadena-Iñiguez, J., Ruiz-Posadas, L.M., Cadena-Zamudio, J.D., González- Ugarte, A.K., Weiss Steider, B, Santiago-Osorio, E, 2015. Fruit extract from a *Sechium edule* hybrid induce apoptosis in leukaemic cell lines but not in normal cells. *Nutr Cancer*, 67, 250–257.

Cadena-Iñiguez, J., Aguiñiga-Sánchez, I., Uriostegui-Arias, M.T., Santiago-Osorio, E., Ruiz-Posadas, L.M., Soto- Hernández, M, 2022. Antiproliferative Effect of *Sechium edule* (Jacq.) Sw., cv. Madre Negra extracts on breast cancer *in Vitro*. *Separations*, 9.

* Presenting author. E-mail address: msoto@colpos.mx (Marcos Soto-Hernández)

S2.P61 Neolignans and diarylnonanoid derivatives from the seed of *Myristica fragrans* Houtt. and their cytotoxic activities against human ovarian cancer cell lines

Jeong Ah Kim^{a,b,*}, Hye Jin Kim^c, Byung Sun Min^d, Mi Jeong Kwon^{a,b}


^a Vessel-Organ Interaction Research Center, VOICE (MRC), College of Pharmacy, Kyungpook National University, Daegu 41566, Republic of Korea

^b BK21 FOUR Community-Based Intelligent Novel Drug Discovery Education Unit, College of Pharmacy and Research Institute of Pharmaceutical Sciences, Kyungpook National University, Daegu 41566, Republic of Korea

^c College of Pharmacy, Keimyung University, 1095 Dalgubeol-daero, Dalseo-Gu, Daegu 42601, Republic of Korea

^d College of Pharmacy, Drug Research and Development Center, Daegu Catholic University, Gyeongbuk 38430, Republic of Korea

Two new neolignans (myrifralignans F–G), four new diarylnonanoid derivatives (myrifragranones A–D), and 18 known compounds from the seed of *Myristica fragrans* Houtt. (nutmeg) were isolated and structurally elucidated. The absolute configurations of these secondary metabolites were accurately determined using the electronic circular dichroism method. The cytotoxic activities of these isolated compounds against cisplatin-sensitive and resistant human ovarian cancer cell lines were evaluated. In particular, the new compound myrifragranone C (**23**) exhibited cytotoxicity against all test cancer cell lines A2780, TOV-112D, and SK-OV3 with IC₅₀ values of 14.1, 16.9, and 33.4 μ M, respectively. Furthermore, compound **23** induced the death of A2780 and SK-OV3 cancer cells via apoptosis. Western blotting revealed that compound **23** significantly increased the expression of cleaved caspase-3 and poly-ADP ribose polymerase and promoted apoptosis via the mitogen-activated protein kinase signaling pathway.



Keywords: *Myristica fragrans*, nutmeg, myrifralignan, myrifragranone, ovarian cancer

References

Vu, T.O., Nguyen, V.P., Min, B.S., Yang, S.Y., Kim, J.A., 2023. Insights into the inhibitory activities of neolignans and diarylnonanoid derivatives from nutmeg (*Myristica fragrans* Houtt.) seeds on soluble epoxide hydrolase using *in vitro* and *in silico* approaches. J. Enzyme. Inhib. Med. Chem. 38, 225109.

* Presenting author. E-mail address: jkim6923@knu.ac.kr (Jeong Ah Kim)

S2.P62 The ameliorative effects of *Moschus* and its major compound L-muscone on traumatic brain injury mice model focusing on the mechanisms of action

Jang-Hoon Kim^{a,b,*}, Seogyong Lee^{a,b}, Jinhyun Bae^a, Hyejin Joo^a, Seok-Jae Ko^c, Beom-Joon Lee^c, Youngmin Bu^a

^a Department of Herbal Pharmacology, College of Korean Medicine, Kyung Hee University, Seoul 02447, Korea

^b Department of Science in Korean medicine, Graduate School, Kyung Hee University, Seoul 02447, Korea

^c Department of Internal Medicine, College of Korean Medicine, Kyung Hee University, Seoul 02447, Korea

Moschus, a dried secretion from musk cyst of male musk deer (*Moschus moschiferus* Linnaeus) has been used in Korean medicine to treat neurological and traumatic symptoms with the effect of resuscitation. It has been reported to reduce ischemic brain damage and chronic stress-induced functional deficits. L-muscone, a major component of *Moschus*, was also reported to have neuroprotective effects. However, there are few studies on its effects and mechanisms in traumatic brain injury (TBI). TBI was induced by controlled cortical impact (depth 2.5 mm; velocity 2 m/s; duration 300 ms) in mice. *Moschus* (10 and 100 mg/kg) and L-muscone (0.5 and 5 mg/kg) were suspended in corn oil and administered twice a day for 8 days after TBI. Brain damage, hemorrhage volume, Evans blue leakage, behavioral deficits were evaluated. In addition, blood brain barrier (BBB) compartments, pathological factors, and water channel were quantified. *Moschus* and L-muscone reduced brain damage, hemorrhage and BBB permeability along with functional improvements. Especially, they restored cognitive function to normal level. They also increased BBB compartments including zonula occludens-1, claudin, and occludin, decreased aquaporin-4, Heme oxygenase-1 and matrix-metalloproteinases -2 & -9 expressions. These results provide a scientific basis for the clinical use of *Moschus* in traditional medicine and suggest that L-muscone might be the main compound in the effects of *Moschus* on TBI, and that *Moschus* and L-muscone might be candidates for TBI therapeutic agents. This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (Ministry of Science and ICT) (No. 2020R1A2C1008603).

Keywords: *Moschus*, L-muscone, traumatic brain injury, blood brain barrier, cognitive function

* Presenting author. E-mail address: shfltm@khu.ac.kr (Jang-Hoon Kim)

S2.P63 Cycloartane-type triterpenoids from *Combretum quadrangulare* (Combretaceae) and their potential for the treatment of hypercholesterolemia

Chae-Yeong An*, Pisey Pel, Young-Won Chin

Natural Products Research Institute and Research Institute of Pharmaceutical Sciences, College of Pharmacy, Seoul National University, 1, Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

This presentation will cover the isolation, structure elucidation and biological evaluation of cycloartane-type triterpenoids including representative compound (**1**) (Fig. 1). *Combretum quadrangulare* Kurz. (Combretaceae) has been traditionally used in Southeast Asian countries as an herbal medicine with antipyretic, antidiysenteric and hepatoprotective properties (Banskota et al., 2003).

Previous studies (An et al., 2023; Won and Son et al., 2023) have reported acyclic and cyclic triterpenoid scaffolds as PCSK9 inhibitors. As part of our continuing search for PCSK9 inhibitory compounds from plants, we selected *C. quadrangulare*, known for its richness in cycloartane-type triterpenoids. Utilizing spectroscopic analysis, we successfully isolated 16 cycloartane-type triterpenoids, all of which underwent structural assignments and testing via ELISA bioassays.




Fig. 1. The structure of cycloartane-type triterpenoid **1**

Moreover, the exploration of the structure-activity relationship of cycloartane-type triterpenoids among the 16 isolates was conducted. Among these compounds, compound **1** was found to be the most potent in PCSK9 inhibitory activity, compared to berberine—a commonly utilized positive control in studies of PCSK9 inhibitory effects. These findings suggest that cycloartane-type triterpenoids could present a promising scaffold for the optimal treatment of hypercholesterolemia. Acknowledgments: This research was supported by grants from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR202207102) and National Research Foundation of Korea (NRF), funded by the Korean government (MSIT) (NRF-2022R1A2C1010084).

Keywords: *Combretum quadrangulare*, combretaceae, cycloartane-type triterpenoid, PCSK9, hypercholesterolemia

References

- Banskota, A.H., Tezuka, Y., Tran, Q., Kadota, S., 2003. Chemical constituents and biological activities of Vietnamese medicinal plants. *Curr. Top. Med. Chem.* 3(2), 227-248.
- An, C.Y., Son, M.G., Chin, Y.W., 2023. Acyclic Triterpenoids from *Alpinia katsumadai* Seeds with proprotein convertase subtilisin/kexin type 9 expression and secretion inhibitory activity. *ACS omega* 8(36), 32804-32816.
- Won, H., Son, M.G., Pel, P., Nhoek, P., An, C.Y., Kim, Y.M., Chae, H.S., Chin, Y.W., 2023. Chemical constituents from *Morus alba* with proprotein convertase subtilisin/kexin type 9 expression and secretion inhibitory activity. *Org. Biomol. Chem.* 21(13), 2801-2808.

* Presenting author. E-mail address: chaeyy02@snu.ac.kr (Chae-Yeong An)

S2.P64 Bitter-tasting plant extracts stimulate mechanisms of gastric acid secretion of immortalized human parietal cells via TAS2Rs depending on their polyphenol content

Phil Richter^{a,*}, Maria-Riera Piqué-Borràs^b, Gerald Künstle^b, V. Somoza^{a,c,d,#}

^a Leibniz Institute for Food Systems Biology at the Technical University of Munich, Lise-Meitner-Straße 34, 85354 Freising, Germany

^b Weleda AG, Preclinical Research and Development, Dychweg 14, 4144 Arlesheim, Switzerland

^c Chair of Nutritional Systems Biology, TUM School of Life Sciences, Technical University of Munich, Lise-Meitner-Straße 34, 85354 Freising, Germany

^d Department of Physiological Chemistry, Faculty of Chemistry, University of Vienna, Josef- Holaubek Platz 2 (UZA II), 1090 Wien, Austria

Bitter taste receptors (TAS2Rs) are not only involved in taste perception, but also in many extra-oral mechanisms. Previous work by our group has shown that bitter substances stimulate the proton secretion (Liszt et al., 2017) and also have anti-inflammatory effects (Tiroch et al., 2021 & 2023).

In this work, an herbal product and the nine different ethanolic plant extracts of which it is composed, namely tinctures of *Artemisia absinthium*, *Achillea millefolium*, *Centaureum erythraea*, *Cichorium intybus*, *Gentiana lutea*, *Juniperus communis*, *Peucedanum ostruthium*, *Salvia officinalis*, *Taraxacum officinalis*, were investigated for their effects on cellular proton secretion in immortalized human parietal cells (HGT-1). Quantitation of polyphenol and flavonoid contents were performed by means of Folin-Ciocalteu reagent and Al³⁺ complex formation. A TAS2R-dependent effect was studied by RT-qPCR experiments and validated by knock-out (ko) approaches. Seven of the investigated extracts as well as the mixture of all nine extracts led to a concentration-dependent stimulation of proton secretion, with a correlation between the respective polyphenol and flavonoid contents and the effect size ($p \leq 0.0001$). In accordance with the literature, the investigation of gene expression revealed regulation of *TAS2R4/5/39*, among others (Soares et al., 2013). Experiments with *TAS2R4ko* and *TAS2R39ko* cells showed a reduction of stimulatory effects on proton secretion ($p \leq 0.05$), validating an involvement of these TAS2Rs. In summary, depending on their polyphenol and flavonoid content, bitter plant extracts lead to a stimulation of proton secretion by the bitter taste receptors *TAS2R4* and *TAS2R39*, which are therefore involved in digestion-promoting mechanisms.

Keywords: bitter plant extracts, TAS2R, human parietal cells, proton secretion, polyphenols

References

- Liszt, K.I., Ley, J.P., Lieder, B., Behrens, M., Stöger, V., Reiner, A., Hochkogler, C.M., Köck, E., Marchiori, A., Hans, J., Widder, S., Krammer, G., Sanger, G.J., Somoza, M.M., Meyerhof, W., Somoza, V. 2017. Caffeine induces gastric acid secretion via bitter taste signaling in gastric parietal cells. In Proceedings of the National Academy of Sciences (Vol. 114, Issue 30). Proceedings of the National Academy of Sciences.
- Soares, S., Kohl, S., Thalmann, S., Mateus, N., Meyerhof, W., De Freitas, V. 2013. Different Phenolic Compounds Activate Distinct Human Bitter Taste Receptors. In Journal of Agricultural and Food Chemistry (Vol. 61, Issue 7, pp. 1525–1533). American Chemical Society (ACS).
- Tiroch, J., Sterneder, S., Di Pizio, A., Lieder, B., Hoelz, K., Holik, A.-K., Pignitter, M., Behrens, M., Somoza, M., Ley, J. P., Somoza, V. 2021. Bitter Sensing TAS2R50 Mediates the trans-Resveratrol-Induced Anti-inflammatory Effect on Interleukin 6 Release in HGF-1 Cells in Culture. In Journal of Agricultural and Food Chemistry (Vol. 69, Issue 45, pp. 13339–13349). American Chemical Society (ACS).
- Tiroch, J., Dunkel, A., Sterneder, S., Zehentner, S., Behrens, M., Di Pizio, A., Ley, J.P., Lieder, B., Somoza, V. 2023. Human Gingival Fibroblasts as a Novel Cell Model Describing the Association between Bitter Taste Thresholds and Interleukin-6 Release. In Journal of Agricultural and Food Chemistry (Vol. 71, Issue 13, pp. 5314–5325). American Chemical Society (ACS).

* Presenting author. E-mail address: p.richter.leibniz-lsb@tum.de (Phil Richter)

S2.P65 Anti-inflammatory testing of some isolated sesquiterpene lactones and methoxylated flavonoids from *Achillea millefolium* L. in an ICAM-1 *in vitro* assay

Carola Aumer*, Sabrina Zölch, Guido Jürgenliemk, Sebastian Schwindl, Jörg Heilmann

Department of Pharmaceutical Biology, University of Regensburg, Universitätsstraße 31, 93053 Regensburg, Germany

Achillea millefolium is a medicinal plant of the Asteraceae family used in both traditional and conventional medicine (Radušienė et al., 2005) mostly in form of alcoholic, hydroalcoholic, and aqueous extract, or infusion and decoction (Dias et al., 2013). Its main secondary metabolites are flavonoids, essential oil, phenolic acids, and sesquiterpene lactones including also proazulenes. The most important therapeutic indications for internal use are loss of appetite, dyspeptic disorders, and the treatment of cramps in dysmenorrhea, and for external use inflammations of the skin and mucous membranes (reviewed in Hutchins et al., 2021). Since the mechanism of action of the anti-inflammatory effect and the substances responsible for this are not known in detail, seven isolates from previous work on *Achillea millefolium* were investigated in an intracellular adhesion molecule-1 (ICAM-1) *in vitro* test (Freischmidt et al., 2012). Thereby, four sesquiterpene lactones of the guaianolide type, namely 8-desacetyl-8-angeloyl-4-epi-matricine, 8a-acetyl-2b,4a,10b-trihydroxy-6bH,7aH,11bH-1(5)-guaien-12,6a-olide, 8a-isopentyl-2b,4b,10b-trihydroxy-6bH,7aH,11bH-1(5)-guaien-12,6a-olide, and 8a-angeloxyl-2b,4b,10b-trihydroxy-6bH,7aH,11bH-1(5)-guaien-12,6a-olide, and two methoxylated flavonoids, pectolinarigenin and artemetin, reduced significantly the ICAM-1 expression in a human endothelial cell line (HMEC-1) in a concentration range from 25-75 µM after stimulation with TNFα. Pectolinarigenin proved to be the most effective substance with a reduction of ICAM-1 expression of 35.5%, while the most effective sesquiterpene lactone only showed an inhibition of 21.3%. These results indicate a slight influence of the tested isolates in the inflammatory process. However, to fully understand the anti-inflammatory effect of these compounds, further approaches regarding their effects in the ICAM-1/NF-kB related pathways are necessary.

Keywords: *Achillea millefolium*, Sesquiterpene lactones, methylated flavonoids, anti-inflammatory, ICAM-1

References

- Dias, M.I., Barros, L., Dueñas, M., Pereira, E., Carvalho, A.M., Alves, R.C., Oliveira, M.B.P.P., Santos-Buelga, C., Ferreira, I.C., 2013. Chemical composition of wild and commercial *Achillea millefolium* L. and bioactivity of the methanolic extract, infusion and decoction. Food Chemistry. 141(4), 4152-4160.
- Freischmidt, A., Jürgenliemk, G., Kraus, B., Okpanyi, S.N., Müller, J., Kelber, O., Weiser, D., Heilmann, J., 2012. Contribution of flavonoids and catechol to the reduction of ICAM-1 expression in endothelial cells by a standardised willow bark extract. Phytomedicine. 19:245–252.
- Hutchins, R., Mills, S., 2021. Millefolii herba (Yarrow), in: European Scientific Cooperative on Phytotherapy (ESCOP), ESCOP monographs The Scientific Foundation for Herbal Medicinal Products, Online series, Exeter, pp. 1-2.
- Radušienė, J., Gudaityte, O., 2005. Distribution of proazulenes in *Achillea millefolium* s.l. wild populations in relation to phytosociological dependence and morphological characters. Plant Genetic Resources. 3(2):136-143.

* Presenting author. E-mail address: carola.aumer@chemie.uni-regensburg.de (Carola Aumer)

S2.P66 Evaluation of aromatase enzyme inhibition by extracts of *Bergenia crassifolia* (Saxifragaceae)

Anna Nickl^{a,*}, Sanja Krstic^a, Jelena S. Katanić Stanković^b, Rudolf Bauer^a

^a University of Graz, Institute of Pharmaceutical Sciences, Department of Pharmacognosy, Beethovenstraße 8, 8010 Graz, Austria

^b University of Kragujevac, Institute for Information Technologies Kragujevac, Department of Science, Jovana Cvijića bb, 34000 Kragujevac, Serbia

Rhizomes and leaves of *Bergenia crassifolia* (L.) Fritsch (Saxifragaceae) are commonly used in the traditional medicine of several countries to treat a variety of diseases, such as gastrointestinal disorders, fevers and gynaecological conditions (Shikov et al., 2014; Koul et al., 2020). The plant contains several bioactive constituents like arbutin, bergenin, flavonoids and tannins. These compounds have been credited with some of the reported pharmacological activities of *B. crassifolia*, which include antimicrobial, antihypertensive, and anti-inflammatory effects (Shikov et al., 2010; Shikov et al., 2014). The aim of this ongoing study is to evaluate the ability of *B. crassifolia* extracts to inhibit aromatase (CYP19A1), a key enzyme in the biosynthesis of oestrogen, which is associated with different diseases, including breast cancer and endometriosis (Nelson et al., 2001). For this purpose, we established an aromatase enzyme inhibition assay, partially modelled on the basis of previously published methods (El-Kersh et al., 2021). Aromatase inhibition is assessed by measuring the fluorescence intensity of fluorescein benzyl ester, the aromatase metabolization product of dibenzylfluorescein (DBF). To optimise and streamline the assay, several different concentrations and combinations of substrate (DBF), enzyme (CypExpress™ 19A1), cofactors (NADPH and G6P) and positive control (ketoconazole) were evaluated before testing *Bergenia* extracts, which were obtained by accelerated solvent extraction (ASE) of flowers, leaves, and rhizomes with methanol. Initial results indicate that all three extracts exhibited a high degree of aromatase enzyme inhibition. Investigations to identify the constituents responsible for this activity are in progress.

Keywords: *Bergenia crassifolia*, endometriosis, aromatase, CYP19A1, inhibition

References

- El-Kersh, D.M., Ezzat, S.M., Salama, M.M., Mahrous, E.A., Attia, Y.M., Ahmed, M.S., Elmazar, M.M., 2021. Anti-estrogenic and anti-aromatase activities of citrus peels major compounds in breast cancer. *Sci.Rep.* 11, 7121.
- Koul, B., Kumar, A., Yadav, D., Jin, J.-O., 2020. *Bergenia* Genus: Traditional Uses, Phytochemistry and Pharmacology. *Molecules*. 25 (23), 5555.
- Nelson, L. R., Bulun, S. E., 2001. Estrogen production and action. *J. Am. Acad. Dermatol.* 45 (3 Suppl.), 116-24.
- Shikov, A.N., Pozharitskaya, O.N., Makarova M.N., Damien Dorman, H.J., Makarov, V.G., Hiltunen, R., Galambosi, B., 2010. Adaptogenic effect of black and fermented leaves of *Bergenia crassifolia* L. in mice. *J. Funct. Foods*. 2, 71–76.
- Shikov A.N., Pozharitskaya O.N., Makarova M.N., Makarov V.G., Wagner H., 2014. *Bergenia crassifolia* (L.) Fritsch- pharmacology and phytochemistry. *Phytomedicine*. 21, 1534–1542.

* Presenting author. E-mail address: anna.nickl@uni-graz.at (Anna Nickl)

S2.P67 Assessment of the anticonvulsant activity of *Myrica gale* L., a traditional irish plant in ethnoveterinary medicine, using a 4-AP induced seizure, electrophysiological model, *in vitro*

Sophie Whyms^a, Gaia Scalabrino^{a,*}, Kate Connor^b, Shipra Nagar^a, Maria Pigott^a, Helen Sheridan^a, Mark Cunningham^b

^a NatPro Centre, School of Pharmacy and Pharmaceutical Sciences, Trinity College Dublin, Dublin 2, Ireland

^b Discipline of Physiology, School of Medicine, Trinity College Dublin, Dublin 2, Ireland

The bogland species *Myrica gale* (MG) has ethnobotanical reports of its use for the treatment of ‘fits in dogs’ (Jackson, 2014) with ‘fits’ in Irish ethnography referring to ‘seizures’ (epilepsy). Other literature reports that many plant oils and their metabolites, especially terpenes, exhibit anticonvulsant activity. Seizures can be reduced or induced *via* inhalation of certain essential oils (EOs) (Tyagi et al., 2003; Yamada et al., 1994). This study assessed the acute anti-epileptic potential of two Irish MG EOs (EO1M, EO15C) and a commercial EO (COM) using *in vitro* electrophysiology. Extracellular local field potential (LFP) recordings measured ictal-like events (ILEs) in adult *Rattus norvegicus* brain slices (400µm). Application of 4-aminopyridine (26.7nM) elicited baseline ILE activity. Subsequently EOs were applied to the tissue *via* perfusate (0.5%v/v). LFP recordings of ILE activity were recorded from the superficial layers (II – III) of the medial entorhinal cortex. EO effects on ictal duration (ID), inter-ictal duration (IID), first spike amplitude (FSA), power spectrum density (PSD) and number of seizures (NoS), were analysed. Application of EO1M significantly reduced ID, IID, NoS, PSD (n=9, $P < 0.01$) and FSA ($P < 0.05$). EO15C significantly reduced ID, IID and NoS (n=9, $P < 0.01$). COM did not demonstrate significant effects on ID, IID, NoS, PSD or FSA (n=9, $P > 0.01$). MG extracted EOs can reduce ILEs in this model and our results support traditional ethnographic use. Qualitative and quantitative metabolomic differences between samples contribute to the variation in activity observed.

Keywords: *Myrica gale*, anticonvulsant, ILE, 4-aminopyridine, essential oils

References

Tyagi A., Delanty N., 2003. Herbal Remedies, Dietary Supplements, and Seizures, *Epilepsia* 44(2), 228 – 235.

Yamada K., Mimaki Y., Sashida Y., 1994. Anticonvulsant effects of inhaling lavender oil vapour. *Biol. Pharm. Bull.* 17, 359 – 360.

Jackson W.P., 2014. Ireland’s Generous Nature. *Missouri Bot. Press* 19 – 410.

* Presenting author. E-mail address: scalabrg@tcd.ie (Gaia Scalabrino)

S2.P68 Centrifugal partition chromatography as a tool for the isolation of polyphenols from ginger rhizomes

Wirginia Kukula-Koch^{a,*}, Dominik Tarabasz^a, Martyna Nizio^a, Apostolis Angelis^b, Lemonia Antoniadi^b, Maria Halabalaki^b, Alexios Leandros Skaltsounis^b

^a Department of Pharmacognosy with Medicinal Plant Unit, Medical University of Lublin, Poland

^b Department of Pharmacognosy and Natural Products Chemistry, School of Pharmacy, National and Kapodistrian University of Athens, Greece

Zingiber officinale Rosc. (Zingiberaceae) belongs to the most widely consumed spices around the world. Its vast application in traditional medicine as an antiemetic, antibacterial, anti-inflammatory or bile production enhancing agent still triggers numerous studies on its metabolites (Kukula-Koch et al., 2019). The study aimed to optimise the protocol for the recovery of ginger polyphenols using modern extraction techniques, like supercritical solvent extraction (SFE), accelerated solvent extraction (ASE), ultrasound-assisted extraction (UAE), but also to perform an efficient fractionation of the ginger oleoresin obtained by supercritical fluid extraction (SFE) using centrifugal partition chromatography (CPC). A successful separation of the two major groups of ginger secondary metabolites, namely phenolic components and terpenes remains a difficult task to achieve. In the study, the applied gradient of hexane, ethyl acetate, butanol and water enabled a fast recovery of the major phenolic components from the preparative CPC instrument equipped with a one-liter-volumed column, directly from the crude oleoresin. The performed optimisation of the biphasic solvent composition, rotation speed and solvent flow allowed for the selection of beneficial settings for the isolation of different polyphenols. The undertaken tasks allow for a large scale recovery of single polyphenols of ginger for food, cosmetic and pharmaceutical applications as both SFE and CPC can be operated in preparative or industrial scales.

Acknowledgements: The research was funded by the Polish National Science Centre project OPUS No 2021/41/B/NZ4/00337

References

Kukula-Koch, W., Czernicka, L. Gingerols and Shogaols from Food. In: Handbook of Dietary Phytochemicals; Xiao, J., Sarker, S.D., Asakawa, Y. (Eds.). Springer: Singapore, 2019, pp. 1–31.

* Presenting author. E-mail address: virginia.kukula@gmail.com (Wirginia Kukula-Koch)

S2.P69 Comparative assessment of the antioxidant capacity of Domat olive leaf and Sari Yaprak olive leaf

Fatma Gül Yoldaş*, Engin Celep

Faculty of Pharmacy, Department of Pharmacognosy, Yeditepe University, 34755 Istanbul, Atasehir, Turkey

Oxidative stress can cause several degenerative conditions, including inflammation, cardiovascular diseases, and cancer. Organisms have inherent mechanisms to reduce oxidative stress, but it is not entirely effective. Thus, they can benefit from additional antioxidants. Many fruits, vegetables, and plant leaves have antioxidant properties. Olive leaves are particularly interesting because they have been used in traditional treatments throughout the Mediterranean (Lawrendiadis, 1961), and their health benefits have been supported by scientific studies (Reece et al., 1991). On the other hand, different olive leaves cultivars showed different antioxidant activity (El et al., 2009).

This presentation will cover the antioxidant capacity of two different *Olea europaea* Domat olive cultivars and Sari Yaprak olive leaves collected from Izmir, Turkey. Domat olive leaves showed higher antioxidant capacity when compared with 21 other olive cultivars (Orak et al., 2012). The Sari Yaprak olive cultivar is exclusively grown in Turgutlu and Manisa, Turkey. In addition, how the antioxidant capacity of sari yaprak olive leaves was affected by geographical features was tested. Different cultivars of olive leaves were extracted with 80% methanol. Their antioxidant capacities were compared with DPPH radical scavenging activity, CUPRAC (cupric reducing antioxidant capacity), and TOAC (total antioxidant capacity) assays. Their phytochemical properties were compared with total phenolic content and total flavonoid content. Sari Yaprak olive leaves showed higher total flavonoid content than Domat olive leaves, whilst Domat olive leaves had higher antioxidant activity on CUPRAC than Sari Yaprak olive leaves. Furthermore, environmental conditions such as altitude, climate, and light affected the total phytochemical assays and antioxidant activities of Sari Yaprak olive leaves.

Keywords: *Olea europaea*, Domat, Sari Yaprak, Phenolic Profile, Antioxidant

References

- Lawrendiadis, G. (1961). Contribution to the knowledge of the medicinal plants of Greece. *Planta Medica* 9(02), 164– 169.
- Reece, R.J., Maxwell, A., 1991. DNA gyrase: structure and function. *Crit. Rev. Biochem. Mol. Biol.* 26, 335-375.
- El SN, Karakaya S, 2009. Olive tree (*Olea europaea*) leaves: potential beneficial effects on human health. *Nutr Rev.* Nov. 67(11), 632-8.
- Orak, H.H., Isbilir, S.S. Yagar, H., 2012. Determination of antioxidant properties of lyophilized olive leaf water extracts obtained from 21 different cultivars. *Food Sci. Biotechnol.* 21, 1065–1074.

* Presenting author. E-mail address: fgyoldas@gmail.com (Fatmagul Yoldas)

S2.P70 Syringin accelerated wound healing and reduced inflammation in human skin fibroblasts and keratinocytes *in vitro*

Andrzej Parzonko*, Anna K. Kiss

Chair and Department of Pharmaceutical Biology, Medical University of Warsaw, 1 Banacha Str., 02-097 Warsaw, Poland

Previous studies have shown that syringin, a trans-sinapyl alcohol glucoside isolated from common lilac (*Syringa vulgaris* L., Oleaceae) bark, stimulates the secretion of transforming growth factor (TGF β) by human monocytes/macrophages and inhibits tumor necrosis factor (TNF α) release (Filipek, 2019). TGF β is an important factor accelerating the regeneration process of damaged skin. It stimulates skin cell migration and proliferation, accelerating the process of restoring tissue continuity at the site of injury. On the other hand, a prolonged inflammatory process at the site of skin damage impedes the healing process. The aim of this study was to investigate the effects of syringin on the wound healing process *in vitro* using human dermal fibroblasts (NHDF) and keratinocytes (HaCaT) cell lines. In the presence of syringin (12.5 – 100 μ M), the cells showed an increase in the invasion potential in scratch assay. A significant increase of migration through a porous membrane in modified Boyden's chambers in skin fibroblasts stimulated with syringin was also observed. Syringin stimulated TGF β secretion by fibroblasts and keratinocytes and inhibited secretion of pro-inflammatory cytokines (IL-6, IL-8). Since the wound healing process can be impaired by prolonged inflammation, which hinders the transition to the proliferative phase, the presented action of syringin justifies its use as a potential agent for accelerating the healing of difficult wounds and ulcers.

Keywords: syringin, *Syringa vulgaris*, wound healing, migration, scratch assay

References

Filipek, A., Wyszomierka, J., Michalak, B., Kiss, A.K., 2019. *Syringa vulgaris* bark as a source of compounds affecting the release of inflammatory mediators from human neutrophils and monocytes/macrophages. *Phytochemistry Letters*. 30, 309-313.

* Presenting author. E-mail address: andrzej.parzonko@wum.edu.pl (Andrzej Parzonko)

S2.P71 Exploring the Impact of Different Heat Extraction Processes on the phytochemical profile of *Cichorium intybus* Extracts

Maria-Riera Piqué-Borràs^{a,*}, Friederike Tews^b, Johann Röhrle^a, Gerald Künstle^a, Mónica Mennet von Eiff^{b,c}

^a Weleda AG, Preclinical Research and Development, Dychweg 14, 4144 Arlesheim, Switzerland

^b Weleda AG, Anthroposophic Pharmacy Competence Center, Dychweg 14, 4144 Arlesheim, Switzerland

^c Weleda AG, Research and Development Pharma, Dychweg 14, 4144 Arlesheim, Switzerland

A wide array of extraction methods is described in pharmacopoeias for producing medicinal products from plant material, varying in extraction medium and temperature. Focusing on *Cichorium intybus*, we explored differences in extracts prepared at various heat levels, from room temperature to boiling point. *Cichorium intybus* extracts are used in Weleda AG's medicinal products for digestive complaints, atopic dermatitis, and allergic asthma, among others. Initially, we screened six *Cichorium intybus* planta total extracts, prepared from fresh, flowering, whole plants, using semi-quantitative analysis. In particular, four hydroethanolic and two aqueous extracts were prepared, including maceration, digestion, infusion, decoction, pressed juice, and rhythmic (Rh) extractions (Table 1). The extracts underwent filtration and lyophilization. Semi-quantitative analysis via UHPLC-HR-QTOF-MS/MS identified 32 secondary metabolites, such as flavonoids, polyphenols, acid esters, carboxylic and caffeic acids, and sesquiterpene lactones. Several metabolites are known for their anti-inflammatory, antioxidant, antimicrobial, and/or hepatoprotective properties, as well as more specific activities such as anti-allergenic, gastroprotective and lung protective effects. Notably, the infusion, followed by two other hydroethanolic extracts (decoction and digestion), showed the largest area under the curve (AUC) for over 60% of the detected metabolites, predominantly flavonoids, acid esters, and caffeic acids. In contrast, maceration, pressed juice, and Rh extraction exhibited smaller or undetectable AUCs for these metabolites, with higher AUC observed for carboxylic acids. These findings suggest that the therapeutic potential of *Cichorium intybus* may vary depending on the extraction methods and the extracted compounds. Subsequent *in vitro* cell models will assess the impact of these methods on pharmacological activities.

Table 1. Overview of the extraction methods for six *Cichorium intybus* extracts.

Sample	Extract medium	Temperature	Extraction time	Method
Maceration	Hydroethanolic	≤25°C	30 days	Ph.Eur. 1.1.7
Digestion	Hydroethanolic	37°C	1 hour	Ph.Eur. 1.2.5
Infusion	Hydroethanolic	Boiling point	5 minutes	Ph.Eur. 1.2.13
Decoction	Hydroethanolic	Boiling point (100°C)	30 minutes	Ph.Eur. 1.2.11
Pressed juice	Aqueous	≤25°C	Fresh plant	n.a.
Rhythmic	Aqueous	Alternating between 4°C (2-6°C) and 37°C (35- 39°C)	35 days	Ph Eur 1.5.1

Keywords: *Cichorium intybus*, plant extraction methods, metabolites, semi-quantitative analysis

*Presenting author: E-mail address: preclinical-dev@weleda.ch (Maria-Riera Piqué-Borràs)

S2.P72 Modulation of Gastric Motility by a Traditional Herbal Preparation in Functional Dyspepsia: Implications of Muscarinic Receptor Inhibition

Maria-Riera Piqué-Borràs^{a,*}, Johann Röhrl^a, Mónica Mennet-von Eiff^b, Gerald Künstle^a

^a Weleda AG, Preclinical Research and Development, Dychweg 14, 4144 Arlesheim, Switzerland

^b Weleda AG, Research and Development Pharma, Dychweg 14, 4144 Arlesheim, Switzerland

This study investigates the impact of Amara-Drops, a traditional herbal remedy for gastrointestinal dysmotility, on fundus accommodation in functional dyspepsia (FD). FD is characterized by impaired gastric accommodation after food intake, however, its etiology is poorly understood. Amara-Drops consist of a mixture of nine hydroethanolic herbal extracts, including *Juniperus communis* (JC), *Peucedanum ostruthium* (PO), and *Salvia officinalis* (SO). We examined the mechanisms of Amara-Drops' dry and individual extracts on gastric motility *ex vivo/in vitro* and their implication for muscarinic receptors, crucial in gastrointestinal disorders. Muscle strips from guinea-pig (GP) gastric fundus were exposed to the whole extract, resulting in significant relaxation at the highest concentrations tested ($p < 0.001$). Additionally, a significant relaxation was observed in Carbachol-induced contracted muscle strips ($p < 0.05$), indicating potential modulation of the M2 receptor. *In vitro* studies revealed binding of the whole extract to both M2 and M3 receptors (IC_{50} : 249 and 380 $\mu\text{g/mL}$, respectively), with selective impairment of M2 receptor activity (IC_{50} : 219 $\mu\text{g/mL}$) by inhibiting intracellular cAMP release. JC, SO, and PO extracts showed inhibitory effects on M2 receptor activity (IC_{50} : 32, 20.1, and 20.8 $\mu\text{g/mL}$, respectively). Additionally, PO extract was found to bind to M2 receptors (IC_{50} : 137 $\mu\text{g/mL}$) and to induce relaxation in Carbachol-induced contracted muscle strips ($p < 0.001$). These results suggest that the pharmaceutical product Amara-Drops, particularly through *Peucedanum ostruthium* extract, modulates gastric motility by inhibiting M2 receptor activity. This finding offers a molecular explanation for the therapeutic use of Amara-Drops to address impaired gastric accommodation in FD.

Keywords: Herbal preparation, gastrointestinal dysmotility, fundus accommodation, muscarinic receptors, Carbachol-induced contraction

*Presenting author. E-mail address: preclinical-dev@weleda.ch (Maria-Riera Piqué-Borràs)

S2.P73 *In vitro* liver metabolism of four biflavonoids and their corresponding monomers

Nicole Herl^{a,*}, Josef Kiermaier^b, Dejan Z. Orčić^c, Jörg Heilmann^a

^a Department of Pharmaceutical Biology, Faculty of Chemistry and Pharmacy, University of Regensburg, Universitätsstraße 31, 93053 Regensburg, Germany

^b Department of Central Analytical Services, Faculty of Chemistry and Pharmacy, University of Regensburg, Universitätsstraße 31, 93053 Regensburg, Germany

^c Department of Chemistry, Biochemistry and Environmental Protection, University of Novi Sad, Faculty of Sciences, Trg Dositeja Obradovića 3, Novi Sad, Serbia

Biflavonoids gained more attention during the last years. These secondary plant metabolites can be found in some well-known medical plants such as *Ginkgo biloba* L. (Šamec et al., 2022) or *Hypericum perforatum* L. (Carrubba et al., 2021) and may contribute to their pharmacological effects. The physiological role of biflavonoids in plants is still discussed but might be related to protection against UV radiation or predators as they are located in plant leaves (Šamec et al., 2022). Pharmacologically, these dimeric flavonoids are known for a wide range of effects such as their anti-inflammatory, antioxidant and antiviral properties (He et al., 2021). Recently, their application and therapeutic benefits in microbial diseases, Alzheimer's disease or metabolism-related diseases such as type2-diabetes was discussed (Menezes et al., 2021). Information on human liver metabolism of flavone dimers especially compared to their corresponding monomers are lacking. Thus, we investigated the metabolic stability of amentoflavone, I3,II8-biapigenin, bilobetin, hinokiflavone, apigenin and acacetin by incubation with human liver S9 fraction. Additional to the microsomal fraction of the liver, the S9 fraction contains the cytosolic fraction as well. Hence, it provides both phase I and phase II enzymes (Richardson et al., 2016). The received metabolites were identified using UHPLC-DAD-MS or HPLC-DAD-MS/MS revealing phase II glucuronidation products as main metabolites for all flavonoids. Interestingly glucuronidation pattern and number of metabolites varied for each compound based on the subunits and their linkage.

Keywords: biflavonoids, flavones, liver metabolism, S9 fraction

References

- Carrubba, A., Lazzara, S., Giovino, A., Ruberto, G., Napoli, E., 2021. Content variability of bioactive secondary metabolites in *Hypericum perforatum* L. *Phytochemistry Letters*. 46, 71-78.
- He, X., Yang, F., Huang, X., 2021. Proceedings of chemistry, pharmacology, pharmacokinetics and synthesis of biflavonoids. *Molecules* 26, 6088.
- Menezes, J.C.J.M.D.S., Diederich, M.F., 2021. Bioactivity of natural biflavonoids in metabolism-related disease and cancer therapies. *Pharmacological Research* 167, 105525.
- Richardson, S.J., Bai, A., Kulkarni, A.A., Moghaddam, M.F., 2016. Efficiency in drug discovery: Liver S9 fraction assay as a screen for metabolic stability. *Drug Metabolism Letters* 10, 83-90.
- Šamec, D., Karalija, E., Dahija, S., Hassan, S.T.S., 2022. Biflavonoids: Important contributions to the health benefits of Ginkgo (*Ginkgo biloba* L.). *Plants* 11, 1381.

* Presenting author. E-mail address: nicole.herl@ur.de (Nicole Herl)

S2.P74 Screening of flavonoids and phenolic compounds in *Betula litwinowii* Doluch. by LC-Q- TOF/MS and determination of their antioxidant potential

Buket Bozkurt^{a,*}, Zinar Pinar Gumus^b, Muzaffer Mukemre^c

^a Department of Pharmacognosy, Faculty of Pharmacy, Ege University, Bornova, 35100, Izmir, Türkiye

^b Central Research Test and Analysis Laboratory Application and Research Center (EGE-MATAL), Ege University, Bornova, 35100, Izmir, Türkiye

^c Yuksekova Vocational School, Department of Plant and Animal Production, Hakkari University, Hakkari, 30100, Türkiye

Birch tree (*Betula* L.) is a tree-shaped plant distributed in the Northern Hemisphere, and its leaves, buds, bark and essential oil are traditionally used for many purposes (Rastogi et al., 2015). To the best of our knowledge, the flavonoid and phenolic compounds of *Betula litwinowii* have not been investigated before. Therefore, the aim of this study is to reveal the flavonoids and phenolic compounds found in the buds and leaves of *B. litwinowii* collected from Van, Türkiye, to determine the total phenolic (TP) and flavonoid contents (TF) and to evaluate the antioxidant activity capacity. Qualification of compounds in aqueous alcohol extracts (80/20:v/v) was performed using a LC-Q- TOF/MS system equipped with an electrospray ionization interface operating in positive and negative ion mode (Sevimli-Gur et al., 2021). According to METLIN library results, extracts prepared from leaves and buds contain quinic acid, quercetin, quercetin 3-O-glucoside, quercetin 3-O-malonylglucoside, 5-desoxyquercetin, 3-O-methylquercetin, rutin, luteolin, diosmetin, genkwanin, gallic acid, epigallocatechin, epicatechin, catechin 7-β-D-xylopyranoside, chlorogenic acid, apigenin, sinapic acid, salidroside, naringenin, maslinic acid, kaempferol, kaempferol-3-glucuronide, 3-methylkaempferol, 8-hydroxykaempferol, caffeic acid, betulonic acid, betulinic acid, 4-O-methylgallate, 4-hydroxybenzaldehyde, 3,4-dihydroxybenzoic acid, procyanidin B2, and myricetin tetramethyl ether. The TP were $220,65 \pm 3.83$ mg gallic acid equivalent (GAE)/g extract and $243,65 \pm 4.27$ mg GAE/g extract for the leaves and buds respectively while the TF were $33,44 \pm 2.13$ and $49,6 \pm 2.75$ mg quercetin/g extract. The antioxidant capacity of the extracts was evaluated using the DPPH, ABTS, and FRAP techniques.

Keywords: *Betula litwinowii*, Betulaceae, LC-Q-TOF/MS

References

- Rastogi, S., Pandey, M.M., Rawat, A.K.S., 2015. Medicinal plants of the genus *Betula* - Traditional uses and a phytochemical-pharmacological review. J. Ethnopharmacol. 159, 62-83.
- Sevimli-Gur, C., Gezgin, Y., Oz, A., Al Sharqi, S., Gumus, Z.P., Dunford, N.T., 2021. Biological activity of the extracts from pecan shelling industry by products. Trans. ASABE. 64(3), 869-877.

* Presenting author. E-mail address: buket.bozkurt@ege.edu.tr (Buket Bozkurt)

S2.P75 The extracts of *Capparis cartilaginea* Decne attenuate inflammasome activation in LPS- stimulated THP-1 macrophages

Bashaer Alsharif^{1,2}, Nadhim Hante^{1,3}, María Jose Santos-Martinez^{1,4}, Fabio Boylan^{1,*}

¹ School of Pharmacy and Pharmaceutical Sciences, Trinity Biomedical Sciences Institute, Trinity College Dublin, Dublin, Ireland

² Department of Pharmacognosy, Faculty of Pharmacy, Umm Al-Qura University, Makkah, Saudi Arabia

³ Faculty of Pharmacy, University of Kufa, Al-Najaf, Iraq

⁴ School of Medicine, Trinity College Dublin, Dublin, Ireland

Capparis cartilaginea Decne. (CC), originating from arid regions in Asia and the Mediterranean basin, is recognized in traditional medicine for its efficacy in treating inflammatory conditions like rheumatoid arthritis, osteoarthritis, and gout through the common use of its leaf tea (Tounekti et al. 2019). However, the underlying anti-inflammatory mechanism of CC is still unclear. Recently, leucine-rich repeat protein-3 (NLRP3) inflammasome activation has been demonstrated to be essential in the pathogenesis of arthritis (Yin et al., 2022). In the present study, we aimed to investigate the effects of CC extract, tea, oil, and their isolated compounds (Alsharif et al., 2023) on NLRP3 inflammasome activation in LPS/ATP-stimulated THP-1 cells. The levels of inflammatory cytokines and NLRP3 inflammasome-related proteins were detected by ELISA and western blotting (WB), respectively. We found that CC extract significantly reduced NLRP3 expression, ASC speck formation, and caspase-1 cleavage, and therefore, IL-1 β and IL-18 secretion in THP-1 cells. The extract also inhibited NLRP3 and caspase-1 activation in LPS/ATP-stimulated THP-1 cells. Furthermore, mitogen-activated protein kinase (MAPK) activation in *P. aeruginosa* LPS/ATP stimulated THP-1 was also attenuated by the extract and some of the tested compounds. These results support the use of this medicinal plant to treat NLRP3 inflammasome-driven inflammatory diseases such as arthritis.

Keywords: *Capparis cartilaginea*, capparaceae, flavonoids, NLRP3 Inflammasome

References

- Alsharif, B., Hante, N., Govoni, B., Verli, H., Kukula-Koch, W. Jose Santos-Martinez, M., Boylan, F., 2023. *Capparis cartilaginea* Decne (Capparaceae): isolation of flavonoids by high-speed countercurrent chromatography and their anti-inflammatory evaluation. *Frontiers in Pharmacology*, 14, Article Number 1285243.
- Tounekti T, Mahdhi M, Khemira H., 2019. Ethnobotanical study of indigenous medicinal plants of Jazan region, Saudi Arabia. Evidence-based complementary and alternative medicine. Article Number 3190670.
- Yin, H., Liu, N., Sigdel, K.R., Duan, L., 2022. Role of NLRP3 Inflammasome in Rheumatoid Arthritis. *Frontiers in Immunology*, 13. Article Number 931690.

* Presenting author. E-mail address: fabio.boyland@tcd.ie (Fabio Boylan)

S2.P76 *Pistacia chinensis* Bunge subsp. *falcata* (Becc. ex Martelli) aqueous extract possesses acetylcholinesterase and monoamine oxidase inhibitory activities

Bashaer Alsharif^{a,b}, Matilde Biasiotti^c, Ammar Bader^b, Wirginia Kukula-Koch^d, Gavin P. Davey^e, Fabio Boylan^{a,*}

^a School of Pharmacy and Pharmaceutical Sciences and Trinity Biomedical Sciences Institute, Trinity College Dublin, Dublin 2, Ireland

^b Department of Pharmacognosy, Faculty of Pharmacy, Umm AlQura University, Makkah, Saudi Arabia

^c Department of Molecular medicine, University of Pavia, Pavia, Italy

^d Department of Pharmacognosy with Medicinal Plants Garden, Medical University of Lublin, Lublin, Poland

^e School of Biochemistry and Immunology, Trinity College, Dublin 2, Ireland

Plants in the genus *Pistacia* represent valuable natural resources for neuroprotection research based on ethnomedicinal knowledge (Al-Saghir and Porter 2005). Research findings have consistently affirmed the significance of bioactive compounds found in *Pistacia* for addressing various central nervous system disorders, including Alzheimer's disease, Parkinson's disease, multiple sclerosis, cerebral ischemia, depression, and anxiety (Moeini, Memariani et al. 2019). *Pistacia chinensis* Bunge subsp. *falcata* (Becc. ex Martelli) (*P. chinensis*) belongs to the Anacardiaceae family and has received limited study on phytochemistry and biological activities compared to other members (Crutcher, Schroeder et al. 2023). This study aimed 1- to characterise the phenolic compounds in *P. chinensis* extract using high-performance liquid chromatography coupled with electrospray ionisation mass spectrometry (HPLC-ESI-QTOF-MS) and 2- to evaluate the neuroprotective potential of the extract through the inhibition of CNS enzymes such as acetylcholinesterase (AChE), and monoamine oxidases (MAO-A and MAO-B). *P. chinensis* was found to be rich in phenolic acids and to a lesser extent in flavonoids, which explains its significant DPPH antioxidant activity (IC₅₀ of 23.37 ± 0.63 µg/mL). The extract was able to inhibit MAO-B and AChE. The anti- monoamine oxidase, anticholinesterase, and antioxidant properties exhibited by *P. chinensis* extract could contribute to its neuroprotective abilities. Thus, the plant can potentially serve as a source of raw material for the industry of functional foods and nutraceuticals targeting the management of neurodegenerative diseases.

Keywords: *Pistacia chinensis*, phenolic compounds, flavonoids, neuroprotective activity, neurodegenerative diseases

References

Al-Saghir, M.G., Porter, D.M., 2005. Stomatal distribution in *Pistacia* sp.(Anacardiaceae). Int. J. Bot. 1(2), 183-187. Crutcher, M., Schroeder, A., Wagner, M., Alsharif, B., Boylan, F., 2023. A review of the phytochemical properties and pharmacological uses of the genus *Pistacia* L.(Anacardiaceae). Infarma-Ciências Farmacêuticas 35(3), 285-331.

Moeini, R., Memariani, Z., Asadi, F., Bozorgi, M. and Gorji, N., 2019. *Pistacia* L. genus as a potential source of neuroprotective natural products. Planta Medica 85(17), 1326-1350.

* Presenting author. E-mail address: fabio.boylan@tcd.ie (Fabio Boylan)

S2.P77 Cyclotides from Bolivian plants

J. Ruiz^a, U. Göransson^b, S. Gunasekera^a

^a Division of Pharmacognosy, Faculty of Pharmacy, Uppsala University

^b Department of Pharmaceutical Biosciences, Uppsala University, Sweden

Cyclotides are a family of circular peptides derived from plants that possess a significant range of biological activities, such as uterotonic, antimicrobial, immunosuppressive, anti-HIV and cytotoxic (Niyomploy et al., 2018). They are cyclic compounds with around 28-37 amino acids (AA) where the N and C terminals are bound together by a peptide bond. They also contain six cysteine residues which form three disulfide bonds to form a Cyclic Cystine knot (CCK) (Burman et al., 2014). Currently there are many publications that give account of the properties of these compounds (Jennings et al., 2001; Svagnard et al., 2004) and there is an ongoing interest in finding new structures with different properties, which can be achieved either by synthetic modifications or by looking for cyclotides in different species that have not been studied before. In the current study, we aim for the latter option. Given the vast Bolivian biodiversity and because of the unique environmental characteristics of certain regions of the country (Ibisch et al., 2003), we believe that the plants that grow in these regions could develop compounds of interest. Therefore, different samples of species from the Violaceae and Rubiaceae families have been collected in Bolivia. These species have not been studied yet and they could represent a source of new structures of cyclotides.




Fig. 1. Sequence and representation of the cyclotide Kalata B1. (De Veer et al., 2014)

So far, from the eleven samples that have shown positive content of cyclotides, the compounds found in *Viola boliviana* have been isolated and processed for sequencing through mass spectrometry (MS). The cyclotides were reduced, alkylated and digested with enzymes for their structural elucidation. As for now, the structure of Kalata B1, Kalata S and Vpub A has been elucidated from one sample of *Viola boliviana*. Although these are known structures, we hope that further analysis will bring unknown structures, since there are more masses to be identified.

Keywords: cyclotide, *Viola boliviana*, kalata, Violaceae, Bolivia

References

- Niyomploy, P., Chan, I., Harvey, P., 2018. J. Nat. Prod. 81(11), 2512-2521.
- Burman, R., Gunasekera, S., Strömstedt, A., Göransson, U., 2014. J. Nat. Prod. 77(3):724-736, (2014). Jennings, C., West, J., Waine, C., 2001. PNAS 98(19), 10614-10619.
- Svagnard, E., Göransson, U., Hocaoglu Z., et al., 2004. J. Nat. Prod. 67(2), 144-147, (2004).
- Ibisch, P., Beck, S., Gerkmann, B., Ecoregiones y ecosistemas, in: Biodiversidad: Riqueza de Bolivia. Estado de Conocimiento y Conservación 47-88, (2003).
- De Veer, S., Kan, M., Craik, D., 2019. Chem. Rev. 119 (24), 12375-12421.

* Presenting author. E-mail address: jose.ruiz@uu.se (Jose Ruiz)

S2.P78 Cytotoxicity assessment of *Rosa pisiformis* (Christ) D. Sosn. (Rosaceae) endemic to Türkiye

Çiğdem Bilgi^a, Esma Sultan Baykara^{b,*}, Esma Yeniçeri^c, Hatice Demiray^d

^a Istanbul University-Cerrahpasa, Faculty of Pharmacy, Department of Pharmacognosy, 34500 Istanbul, Türkiye

^b Istanbul University-Cerrahpasa, Faculty of Pharmacy, 34500 Istanbul, Türkiye

^c Department of Chemistry, Institute of Science and Technology, Erzincan Binali Yıldırım University, 24030 Erzincan, Türkiye

^d Ege University, Faculty of Science Department of Biology, Section of Botany, Izmir, Türkiye

Interest in medicinal plants grows due to their low cost, availability, and minimal side effects. Among diverse medicinal plants, the genus *Rosa* L. is known to possess various species with promising anticancer activity (Fayaz et al., 2024). However, research exploring the bioactivity potential of *Rosa pisiformis* (Christ) D. Sosn. (Rosaceae), a Turkish endemic species (Kültür, 2003), remains limited. This study aims to evaluate the cytotoxic activity of *R. pisiformis* extracts against various cancer cell lines using the MTT assay (Mosmann, 1983). In this context, MCF-7 (breast cancer), HT-29 (colon cancer), and A-549 (lung cancer) cells were treated with ethanolic and hydroalcoholic (ethanol 80% v/v) extracts of leaves and fruits of the plant collected from Gumushane/Türkiye. Experiments were triplicated, with 5-Fluorouracil (5-FU) as the positive control. Cell viability was determined by comparing the formazan concentrations of the treated cells with those of the untreated control. The ethanolic fruit extract of *R. pisiformis* exhibited the most promising results among the tested extracts (Table 1). In the MCF-7 cell line, it exhibited an IC₅₀ value of $14.37 \pm 0.05 \mu\text{g/mL}$ compared to 5-FU's IC₅₀ value of $2.86 \pm 0.30 \mu\text{g/mL}$, indicating significant but less potent cytotoxicity compared to 5-FU. While most effective against MCF-7, it also displayed moderate activity against HT-29 and A549 cell lines, with IC₅₀ values of $36.22 \pm 0.13 \mu\text{g/mL}$ and $48.84 \pm 0.28 \mu\text{g/mL}$, respectively.

Table 1. Cytotoxic activity of ethanolic and hydroalcoholic extracts of *Rosa pisiformis*

Extracts	MCF-7 (IC ₅₀ , μM)	HT-29 (IC ₅₀ , μM)	A549 (IC ₅₀ , μM)
Ethanolic fruit extract (100%)	$14,37 \pm 0,05$	$36,22 \pm 0,13$	$48,84 \pm 0,28$
Hydroalcoholic fruit extract (80:20; v/v)	$33,07 \pm 0,20$	$84,82 \pm 0,24$	$78,44 \pm 0,23$
Ethanolic leaf extract (100%)	$54,84 \pm 0,25$	$89,23 \pm 0,23$	$110,34 \pm 0,23$
Hydroalcoholic leaf extract (80:20; v/v)	$40,99 \pm 0,32$	$90,15 \pm 0,28$	$52,99 \pm 0,37$
5-FU	$22 \pm 2,33$	$25,98 \pm 3,1$	$44,21 \pm 3,54$

This is the first report revealing the potent cytotoxicity of *R. pisiformis* against MCF-7 breast cancer cells, suggesting its potential for further investigation as a therapeutic candidate.

Keywords: *Rosa pisiformis*, MTT, cytotoxicity, cancer

References

- Fayaz, F., Singh, K., Gairola, S., Ahmed, Z., Shah, B.A., 2024. A Comprehensive review on phytochemistry and pharmacology of *Rosa* species (Rosaceae). *Curr. Top. Med. Chem.* 24(4), 364-378.
- Kültür, Ş., 2003. Morphological and palynological studies on *Rosa pisiformis* in Eastern Turkey. *Journal of Faculty of Pharmacy of Istanbul University.* 35, 111-119.
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods.* 65(1-2), 55-63.

* Presenting author. E-mail address: esmasultanbaykara@ogr.iuc.edu.tr (Esma Sultan Baykara)

S2.P79 Confirming (–)-epicatechin’s biological signature; universal modulation of NOS in chronic disease

Melissa M. Henckel^{a,b,*}, Desiree T. Reeves^a, Jane E.-B. Reusch^{a,b}, Amy C. Keller^{a,b}

^a RMR VA Medical Center, Research, 1700 N. Wheeling St., Aurora, CO 80045, USA

^b University of Colorado Anschutz Medical Campus, 12801 E. 17th St., Aurora, CO 80045, USA

Cardiovascular disease (CVD), common in those suffering from diabetes, is the leading cause of global mortality (WHO, 2021). Our work shows that botanical flavonoid (–)-epicatechin stimulates vasodilation and insulin secretion (Chun et al., 2022). As insulin regulates vessel reactivity, we hypothesized that (–)-epicatechin acts globally by modulating enzyme nitric oxide synthase (NOS), a regulator of vessel reactivity (eNOS) and pancreatic inflammation (iNOS), critical to insulin production. We administered 1 mg/kg/day of (–)-epicatechin to female Wistar rats for 7 days, then injected caerulein to induce pancreatitis. We measured glucose tolerance prior to caerulein treatment, and vascular reactivity, pancreatic iNOS and insulin production post-sacrifice. (–)-Epicatechin significantly lowered glucose area under the curve (AUC) as compared with control during a glucose tolerance test ($4,458 \pm 1167$ AUC vs. $7,973 \pm 1,177$ AUC, $p < 0.05$). Caerulein treatment resulted in diminished percentage of vasodilation response to acetylcholine (ACh, upstream of eNOS), restored by (–)-epicatechin ($55.3 \pm 8.2\%$ vs. $69.5 \pm 2.7\%$, $10 \mu\text{M}$ ACh, $p < 0.05$). (–)-Epicatechin treatment resulted in lower expression of iNOS in (–)-epicatechin-treated animals as compared with controls (0.0091 ± 0.0039 colorimetric intensity units [CIU] vs. 0.0542 ± 0.0232 CIU, $p < 0.05$), and elevated expression in animals treated with both (–)-epicatechin and caerulein as compared with caerulein alone (0.5353 ± 0.0649 CIU vs. 0.2710 ± 0.1068 CIU, $p < 0.05$). (–)-Epicatechin restores caerulein-damaged vasodilation and modulates pancreatic iNOS expression. These data support our hypothesis that (–)-epicatechin globally modifies NOS activity, suggesting its potential use in chronic disease.

Keywords: cardiovascular disease, diabetes, vasculature, flavonoid, nitric oxide synthase

References

World Health Organization. Cardiovascular diseases fact sheet. June 11, 2021; [www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](http://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)). Accessed February 14, 2024.

Chun, J., Henckel, M.M., Knaub, L.A., Hull, S.E., Pott, G.B., Ramirez, D.G., Reusch, J.E.-B., Keller, A.C. 2022. (–)-Epicatechin reverses glucose intolerance in rats housed at thermoneutrality. *Planta Med.* 88, 735-744.

* Presenting author. E-mail address: melissa.henckel@cuanschutz.edu (Melissa M. Henckel)

S2.P80 Analytical development on *Pulsatilla* species: comparison of two species

Frédéric Goncalves, Audrey Thorel, Khalil Taoubi, Stéphanie Chanut

Laboratoires Boiron, 2, avenue de l'Ouest Lyonnais, 69510 Messimy, France

This presentation will cover the analytical development and the validation of a method to assay the protoanemonin content in different species of *Pulsatilla*. Protoanemonin is one of the most characteristic compounds of *Pulsatilla* species and is considered as a toxic marker. In order to obtain a precise quantification of this compound in a hydro-alcoholic extract of the plant, an analytical method using liquid chromatography was set and a validation of the method was performed for two different species and protoanemonin contents were compared. The different results, particularly the accuracy profile of the methods is presented. Botanical characteristics and other constituents such as flavonoids and phenolic compounds are also given a complete comparison of *Pulsatilla* species. The methodology presented to set the botanical characterization and the analytical method can be considered as appropriate to elaborate pharmacopoeial monographs.

Keywords: *Pulsatilla pratensis* Mill., *Pulsatilla vulgaris* Mill., analytical method, development, validation, accuracy profile, protoanemonin, HPLC, flavonoids, phenolic compounds

* Presenting author. E-mail address: stephanie.chanut@boiron.fr (Stéphanie Chanut)

S2.P81 Wound healing effects of *Arnica montana* formulations administered by different routes in mice

Jean-François Bisson^a, Estelle Lager^a, Sandrine Lemoine^a, Julie Colin^a, Stéphanie Chanut^b, Pascal Trempat^{b,*}, Naoual Boujedaini^{b#}

^a Department of Dermatology, ETAP S.A., 13 rue du Bois de la Champelle, Technopôle Nancy-Brabois, 54500 Vandœuvre-lès-Nancy, France

^b Research Department, Laboratoires BOIRON, 2 avenue de l'Ouest Lyonnais, 69510 Messimy, France

[#] Lyon, France: NB is no longer belonging to Laboratoires BOIRON

This presentation will cover the use of different preparations of *Arnica* in wound healing. *Arnica montana* L. (*A. montana*) is a traditionally therapeutic plant, known for antalgic and anti-inflammatory properties (French Pharmacopeia 11th Edition, 2013; Iannitti et al., 2016). The plant is used to treat skin bruises, sprains, contusions, and hematomas but, to our knowledge was never tested so far in incisional wounds in mice. Blinded treatments of *A. montana* 5C or *A. montana* mother tincture gel at 7% (w/w) and their respective controls given by systemic or topical routes were performed daily during 20 days. An External Macroscopic Global score (EMG) taking into account the length, the width, the swelling and the visibility of the wound, were performed in order to evaluate the progression of wound healing. The inflammatory cell infiltration, necollagen, re-epithelialization and angiogenesis were quantified by histological and immunohistochemical analyses. At Day 2, both treatments significantly decreased EMG scores faster compared with their placebos. Moreover, *A. montana* 5C and *A. montana* gel significantly accelerated complete wound closure by 1.7 or 3.1 days and reduced inflammatory cell infiltrations by 40% or 55.6% respectively compared with placebos. Finally, *A. montana* 5C significantly improved the formation of neocollagen, re-epithelialization and angiogenesis while *A. montana* gel only favoured re-epithelialization. *A. montana* 5C does not impact cell viability in primary cells (Paumier et al., 2022). In conclusion, we showed that independent of the route of administration, *A. montana* acts in a similar way on the several overlapping phases of wound healing.

Keywords: *Arnica montana* L., inflammatory response, wound healing, mouse

References

French Pharmacopeia 11th Edition, 2013. *Arnica* (whole plant) for Homoeopathic Preparations. Agence Nationale de Sécurité des Médicaments (ANSM), Saint Denis, France.

Iannitti, T., Morales-Medina, J.C., Bellavite, P., Rottigni, V., Palmieri, B., 2016. Effectiveness and Safety of *Arnica montana* in Post-Surgical Setting, Pain and Inflammation. *Am. J. Ther.* 23, e184-197.

Paumier, A., Verre, J., Tribolo, S., Boujedaini, N., 2022. Anti-oxidant Effect of High Dilutions of *Arnica montana*, *Arsenicum Album*, and *Lachesis Mutus* in Microglial Cells *in Vitro*. Dose-Response. 20, 155932582211036.

* Presenting author. E-mail address: pascal.trempat@boiron.fr (Pascal Trempat)

S2.P82 Effect of *Tormentillae tinctura* and its postbiotic metabolites on human intestinal epithelium cellular model

Aleksandra Kruk^{a,*}, Jakub Piwowarski^a, Łukasz Grześkowiak^b, Jurgen Zentek^b, Sebastian Granica^c

^a Microbiota Lab, Department of Pharmaceutical Biology, Medical University of Warsaw, Poland

^b Institute of Animal Nutrition, Freie Universität Berlin, Germany

^c Department of Pharmaceutical Biology, Medical University of Warsaw, Poland

Tormentillae tinctura (TT) has been used for centuries to treat gastrointestinal tract ailments (Melzig et. al., 2020). Due to its tannin-rich composition, TT may have a beneficial effect on the intestinal epithelium. Thanks to these properties, TT could offer an opportunity for novel approaches in the therapy of leaky gut syndrome, which is a serious ailment that reduces the quality of life (Granica et. al., 2020). The presented research aimed to determine the effect of TT and its postbiotic metabolites (TTMs) on the integration and stability of the intestinal barrier. TTMs were obtained by incubation of TT with human fecal slurries from 3 healthy donors. After incubation, TTMs were analyzed (UHPLC-DAD-MSn) and isolated (solid phase extraction). As a cellular model, a Caco-2 monolayer with induced tight junction (TJ) disruption by bacterial toxins was used. The influence of TT and TTMs on cytotoxicity, transepithelial electrical resistance (TEER) value, and TJ proteins production (western blot) and expression (qPCR) on Caco-2 intestinal epithelium was determined. It was observed that TT and TTMs increased the TEER value and modulated the expression and production of selected TJ proteins, thereby beneficially influencing the stability and integrity of the intestinal epithelium.

Acknowledgments: The project has been funded by the Polish National Science Centre research grant Sonata 5 2021/40/C/NZ7/00231.

Keywords: intestinal epithelium, leaky gut, postbiotic metabolites, *Tormentillae tinctura*

References

- Granica, S., Vahjen, W., Zentek, J., Melzig M.F., Pawłowska, K.A., Piwowarski, J.P., 2020. *Lythrum salicaria* ellagitannins stimulate IPEC-J2 cells monolayer formation and inhibit enteropathogenic escherichia coli growth and adhesion. J. Nat. Prod. 83(12), 3614–3622.
- Melzig, M.F., Böttger, S., 2020. *Tormentillae rhizoma* - Review for an underestimated european herbal drug. Planta Med. 86(15), 1050–1057.

* Presenting author. E-mail address: akruk@wum.edu.pl (Aleksandra Kruk)

S2.P83 Effects of the Korean Herbal Prescription, Bojanggunbi-tang, on indomethacin--induced Small Bowel Injury focusing on mechanisms of action

Seogyong Lee^{a,b,*}, Jang-Hoon Kim^{a,b}, Jinhyun Bae^a, Hyejin Joo^a, Seok-Jae Ko^c, Beom-Joon Lee^c, Youngmin Bu^a

^a Department of Herbal Pharmacology, College of Korean Medicine, Kyung Hee University, Seoul 02447, Korea

^b Department of Science in Korean medicine, Graduate School, Kyung Hee University, Seoul 02447, Korea

^c Department of Internal Medicine, College of Korean Medicine, Kyung Hee University, Seoul 02447, Korea

Bojanggunbi-tang (BGT) is a well-known combination of ancient herbal medicines in Korea and has been widely used for treating gastrointestinal symptoms. The effects and mechanisms of action of BGT on non-steroidal anti-inflammatory drug-induced small intestinal injury (NSI) was evaluated by using a murine model of indomethacin-induced NSI. Indomethacin (15 mg/kg) was subcutaneously injected to induce NSI mice model. BGT was administered twice, 30 min before and 6 h after the induction of NSI, at doses of 50, 150, and 450 mg/kg, while the positive control received sodium alginate. Body weight, small intestine length, macroscopic damages, and histological damages were assessed 24 h after induction. Proinflammatory cytokines, tight junction proteins and gut commensal microbiota were investigated to reveal the mechanisms of action. BGT alleviated the shortening of small intestine, ulcer, and inflammation scores. In addition, BGT inhibited loss of the tight junction proteins including zonula occludens-1, occludin and claudin-1, the proinflammatory cytokines including interleukin (IL)-6 and IL-1b, and the gut commensal microbiota including *E. coli*. In conclusion, BGT ameliorated NSI via protective effects on tight junction, and anti-inflammatory and anti-ulcerative properties. The current study suggests that BGT could be a therapeutic option for NSI.

Acknowledgments: This study was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean Government (MSIT) (No. 2022R1C1C1004937).

Keywords: non-steroidal anti-inflammatory drugs, small intestinal injury, inflammation, tight junction, bojanggunbi-tang

* Presenting author. E-mail address: jewel9707@naver.com (Seogyong Lee)

S2.P84 Interaction of an herbal multi-component medicine for treatment of digestive disorders with the growth of selected intestinal lactic acid bacteria

Alina Henneck^a, Elisabeth Bauermann^a, Maria-Riera Piqué-Borràs^b, Sebastian Schmelzer^a, Johann Röhl^{b,*}, Lydia Nausch^a

^a Institute of Nutrition and Food Supply Management, University of Applied Sciences Weihenstephan-Triesdorf, Markgrafenstr. 16, 91746 Weidenbach, Germany

^b Weleda AG, Preclinical Research and Development, Dychweg 14, 4144 Arlesheim, Switzerland

The multi-component herbal medicine Amara-Drops is indicated for treatment of moderate digestive disorders like fullness, heartburn, nausea and disturbed gastrointestinal motility. It is composed of nine different hydroethanolic herbal extracts, *Artemisia absinthium* (AA), *Centaureum erythraea* (CE), *Cichorium intybus* (CI), *Gentiana lutea* (GL), *Juniperus communis* (JC), *Millefolii herba* (MH), *Peucedanum ostruthium* (PO), *Salvia officinalis* (SO) and *Taraxacum* (T). A well functioning digestion is inevitably linked to a healthy intestinal microbiota. Thus, medication should not interfere with intestinal commensal microbes. We aimed to evaluate the interaction of the herbal extracts with *Lactiplantibacillus plantarum* and *Ligilactobacillus salivarius* using an *in vitro* approach. Dry extracts were prepared from mother tinctures of extracts used for production of the final medicinal product Amara-Drops. The antibiotic tetracycline was used as reference compound (15 µg/mL). Bacteria were incubated in liquid culture under constant shaking at 37°C for up to 7 h together with test samples or controls. The growth of *L. plantarum* and *L. salivarius* was not affected at concentrations <5 mg/mL. Only JC and SO showed antibacterial effects towards *L. plantarum* and *L. salivarius* at very high concentrations between 5-10 mg/mL. JC and SO showed now effect at 1 mg/mL. The reference antibiotic completely inhibited the growth of tested bacterial strains (Tab. 1).

Tab. 1: Overview of differential effects of herbal extracts from Amara-Drops on growth of bacterial strains *in vitro*. ■ = growth promotion; □ = no effect; ■ = growth inhibition; ■ = no effect at low concentration (1 mg/ml), inhibition at high concentration (5-10 mg/ml).

	Amara extract mixture	<i>Artemisia absinthium</i>	<i>Centaureum erythraea</i>	<i>Cichorium intybus</i>	<i>Gentiana lutea</i>	<i>Juniperus communis</i>	<i>Millefolii herba</i>	<i>Peucedanum ostruthium</i>	<i>Salvia officinalis</i>	<i>Taraxacum</i>	Tetracycline (15 µg/ml)
<i>L. plantarum</i>	□	□	□	□	■	■	■	□	■	■	■
<i>L. salivarius</i>	□	□	□	□	□	■	□	□	■	■	■

These results indicate that Amara-Drops are compatible with the intestinal commensal bacterial strains *L. plantarum* and *L. salivarius* and might be considered as microbiome-friendly. Additional studies are required to further characterise the compatibility of Amara-Drops with intestinal microbiota.

Keywords: Amara, lactobacilli, intestinal microbiome, microbiome friendly

* Presenting author. E-mail address: preclinical-dev@weleda.ch (Johann Röhl)

S2.P85 *Vangueria infausta* Burch (wild medlar) extract improves semen quality by regulating testosterone levels and oxidative stress in testicular Leydig cells

Nelisiwe Masuku, Jeremiah Unuofin, Adewale Oladipo, Sogolo Lebelo

Department of Life and Consumer Sciences, College of Agriculture and Environmental Sciences, University of South Africa, Private Bag X06, Florida 1710, South Africa

Recently, men's sexual reproductive challenges have emerged as a public health problem. The utilization of medicinal plants for the treatment and management of male infertility could provide an alternative to available chemical drugs. This study aims to identify the bioactive compounds of *Vangueria infausta* Burch (VI) plant and investigate the antioxidant, cytotoxic, and androgenic activities of acetone (A), ethanol (E), methanol (M), and water (W) extracts against TM3 Leydig cells. Ultra-performance liquid chromatography-mass spectrometry (UPLC-MS) analysis revealed the presence of rutin, quercetin glucoside, nicotiflorin, quercitrin, quinic acid, caffeoylquinic acid (1, 2, and 3), 13S-Hydroxy-9Z,11E,15Z-octadecatrienoic acid, 12,13-Dihydroxy-9Z octadecenoic acid, and dicaffeoylquinic acid. Quantitative phytochemical analysis showed that the VI@A had the highest total flavonoid content (TFC) (221.95 mg QE/g) and proanthocyanidins content (PAC) (202.71 mg CE/g), while the VI@E had the highest total phenolic content (TPC) (83.76 mg 24 GAE/g). Also, the VI@M and VI@A extracts had the highest potential to scavenge DPPH and ABTS (482.44 and 0.36 $\mu\text{mol TE/g}$, respectively), while the VI@E induce strong ferric ion- reducing power (0.27 $\mu\text{mol TE/g}$). Besides, all four extracts of VI inhibited TM3 cell growth with increasing dosage while glutathione, lipid peroxidation, and DNA fragmentation pattern provided insights relating to the antioxidant effects on cells. Moreover, the enzyme-linked immunosorbent assay showed that VI@E and VI@W extracts increased the production of testosterone from TM3 cells in the presence of human chorionic gonadotropin (hCG). These findings provide the scientific basis for the use of *V. infausta* as a source of natural antioxidants with androgenic effects.

Keywords: *Vangueria infausta* Burch, antioxidant activity, cytotoxicity, testosterone, TM3 Leydig cells

* Presenting author. E-mail address: oladio@unisa.ac.za (Adewale Oladipo)

S2.P86 Phytochemical, antioxidant, and androgenic effects of multi-solvent extracts from *Strychnos spinosa* Lam on mouse testicular TM3 Leydig cells

Nelisiwe Masuku^a, Jeremiah Unuofin^{a*}, Adewale Oladipo^a, Sogolo Lebelo^a

^aDepartment of Life and Consumer Sciences, College of Agriculture and Environmental Sciences, University of South Africa, Private Bag X06, Florida 1710, South Africa

Men's sexual and reproductive health challenges including infertility, lack of libido, and erectile dysfunction are of concern in the last few decades. The utilization of medicinal plant extracts in their treatment and prevention could be beneficial. This study evaluated the antioxidant content, phytochemicals, and androgenic activity of *Strychnos spinosa* (SS) leaf extracts from different solvents (water W, methanol M, ethanol E, and acetone A) on TM3 Leydig cells. Ultra-performance liquid chromatography-mass spectrometry (UPLC-MS) analysis UPLC-Q-TOF-MS analysis of the plant material further revealed the presence of 13 bioactive compounds mainly polyphenols and other components such as caffeoylquinic acid (1, 2, and 3), dicaffeoylquinic acid (1, 2, and 3), 13S- Hydroxy-9Z,11E,15Z-octadecatrienoic acid, and 12,13-Dihydroxy-9Z-octadecenoic acid. Specifically, the SS-A extract contained the highest total phenolic content (TPC) (419.04 mg GAE/g), total flavonoid content (TFC) (145.46 mg QE/g), and proanthocyanidins content (PAC) (131.43 mg CE/g) compared to other extracts. Besides, the SS-M extract had the highest ABTS scavenging potential (0.3759 $\mu\text{mol TE/g}$) while the SS-W extract had the highest DPPH radical scavenging ability (542.78 $\mu\text{mol TE/g}$) and strong ferric-reducing power (FRAP) activity (0.2098 $\mu\text{mol TE/g}$). All four extracts of SS decreased the viability of TM3 cells in a dose-dependent manner while the mitochondrial membrane potential and DNA fragmentation pattern provided insights into the underlying mechanism of cell inhibition. Additionally, treatment with extracts enhanced testosterone production by TM3 cells in the presence of human chorionic gonadotropin (hCG). In summary, the fascinating androgenic properties displayed by SS extracts highlighted their potential natural remedy in sexual health treatment

Keywords: antioxidants, phytochemicals, *Strychnos spinosa* Lam., testosterone, TM3 Leydig cells

* Presenting author. E-mail address: unuofinjeremiah@gmail.com (Jeremiah Unuofin)

S2.P87 Unlocking longevity through the use of plant adaptogens: lifespan extension by Maral root (*Rhaponticum carthamoides* (Willd.) Iljin) extract in *Caenorhabditis elegans*

Monika N. Todorova^a, Velislava Todorova^c, Zhivka Balcheva-Sivenova^{a,*}, Martina S. Savova^{a,b}, Stanislava Ivanova^c, Kalin Ivanov^c, Milen I. Georgiev^{a,b}

^a Laboratory of Metabolomics, Institute of Microbiology, Bulgarian Academy of Sciences, 139 Ruski Blvd., 4000 Plovdiv, Bulgaria

^b Department of Plant Cell Biotechnology, Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

^c Department of Pharmacognosy and Pharmaceutical Chemistry, Faculty of Pharmacy, Medical University-Plovdiv, 4002 Plovdiv, Bulgaria

As human life expectancy continues to rise, the burden of age-related diseases and diminished healthspan in late-life years remains a significant challenge. Consequently, there is a growing need for effective strategies to enhance both the quality and duration of life (Guarente et al., 2024). Longevity-promoting interventions, particularly those derived from natural sources, are gaining attention due to their potential to address age-related health issues (Todorova et al., 2024). *Rhaponticum carthamoides* (Willd.) Iljin, commonly known as Maral root or Russian leuzea, is a renowned adaptogenic plant in Siberian traditional medicine (Todorova et al., 2023). The chemically diverse root extract of this plant represents a unique reservoir of bioactive compounds, including flavonoids and phytoecdysteroids. Notably, Maral root has been linked to fatigue reduction, improved strength, immunomodulatory and anti-aging properties (Todorova et al., 2023). Building upon this premise, our study explores the potential of Maral root to modulate the aging processes, utilizing *Caenorhabditis elegans* as a model system. To comprehensively evaluate the impact of Maral root on aging, we employed a range of approaches, including behavioral and stress assays. The results of our investigation demonstrate not only an extension of lifespan but also a notable enhancement in healthspan, accompanied by improved fitness in the nematode population. The observed effects underscore Maral root's potential as a promising natural adaptogen capable of mitigating the decline associated with aging. These findings contribute valuable insights to the growing body of research focused on natural interventions for promoting longevity and combating age-related health challenges.

Keywords: longevity, adaptogens, *Caenorhabditis elegans*, maral root, *Rhaponticum carthamoides*

References

- Guarente, L., Sinclair, D.A., Kroemer, G., 2024. Human trials exploring anti-aging medicines. *Cell Metab.* 36(2), 354- 376.
- Todorova, M.N., Savova, M.S., Mihaylova, L.V., Georgiev, M.I., 2024a. Nurturing longevity through natural compounds: Where do we stand, and where do we go? *Food Frontiers.* 5, 267-310.
- Todorova, V., Savova, M.S., Ivanova, S., Ivanov, K., Georgiev, M.I. 2023. Anti-adipogenic activity of *Rhaponticum carthamoides* and its secondary metabolites. *Nutrients*, 15(13), 3061.

* Presenting author. E-mail address: sivenova_jivka@abv.bg (Zhivka Balcheva-Sivenova)

S2.P88 Revealing the veil of barberry alkaloids: high-throughput non-targeted UHPLC-DIA-HRMS profiling enhanced by solid phase extraction based on mixed-mode sorbent

Štefan Kosturko^{a,c,*}, Anna Hošťálková^a, Michal Kořínek^b, Lucie Nováková^c, Marcela Šafratová^a

^a Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmacy in Hradec Králové, Charles University Akademika Heyrovského 1203, 500 03, Hradec Králové, Czech Republic

^b Graduate Institute of Natural Products, College of Pharmacy, Kaohsiung Medical University Shih-Chuan 1st Rd., 807 08, Kaohsiung, Taiwan

^c Department of Analytical Chemistry, Faculty of Pharmacy in Hradec Králové, Charles University Akademika Heyrovského 1203, 500 03, Hradec Králové, Czech Republic

Several isoquinoline alkaloids have been identified as antiviral agents with broad-spectrum activity against animal and human coronaviruses (Fielding et al., 2020). Based on our previous research, this study focuses on examining isoquinoline alkaloids within the genus *Berberis* L. (Berberidaceae). It's worth noting that a number of these alkaloids have shown promising efficacy against the SARS-CoV-2 spike protein in pseudovirus neutralization assays. Our study is focused on exploring and elucidating the phytochemical profiles of isoquinoline alkaloids, emphasizing the diversity of their structures and quantitative variations within this genus. Our research is focused on devising an analytical approach for the high-throughput comprehensive profiling of more than 20 isoquinoline alkaloids via UHPLC-HRMS with the qTOF platform. We evaluated multiple stationary phases during the method optimization process (BEH C18, CSH C18, Biphenyl, ACE C18-PFP) and experimented with different mobile phase compositions. This included variations in organic components (ACN, MeOH) and water components such as 10 mM ammonium formate at pH 3, 4, 5, and 9, along with concentrations of formic acid (0.02%, 0.1%, 0.5%) and ammonium hydroxide (0.02%, 0.1%, 0.5%). We designed a solid-phase extraction (SPE) procedure for sample preparation utilizing a mixed-mode polymeric sorbent with weak cation and reverse phase properties. This purification method effectively removes undesirable acidic and neutral secondary metabolites from the barberry crude extract, leading to improved performance of UHPLC-HRMS analysis. Our comprehensive methodology was effectively employed for non-targeted UHPLC-DIA-HRMS screening of various barberry species.

Keywords: *Berberis* spp., Berberidaceae, isoquinoline alkaloids, WCX-SPE, UHPLC-DIA-HRMS

References

Fielding, B.C., da Silva Maia Bezerra Filho, C., Ismail, N.S.M., Sousa, D.P.d., 2020 Alkaloids: therapeutic potential against human coronaviruses. *Molecules*. 25, 5496.

* Presenting author. E-mail address: kosturks@faf.cuni.cz (Štefan Kosturko)

S2.P89 *Viscum album* extracts in the perioperative oncological setting: evidence from bench to bedside

Florian Pelzer^{a,*}, Wilfried Tröger^b, Marcus Reif^a, Stephan Baumgartner^{a,b}

^a Iscador AG, Spitalstraße 22, 79539 Lörrach, Germany

^b Society for Cancer Research, Kirschweg 9, 4144 Arlesheim, Switzerland

This presentation will show preclinical and clinical evidence on the use of aqueous extracts from *Viscum album* (L.) to reduce postoperative immunosuppression in the oncological setting. Postoperative immunosuppression, especially a reduced cell-mediated immunity, is often seen after surgery and is associated with worse outcome, i.e. delayed wound healing, infections, sepsis, multiple organ failure and cancer recurrence (Lachmann et al., 2018). Aqueous extracts from the European mistletoe *Viscum album* (L.) contain mistletoe lectins, viscotoxins, flavonoids and polyphenols (Urech et Baumgartner, 2015). They stimulate the maturation of dendritic cells, activate natural killer cells and have anti-inflammatory properties by specifically inhibiting cyclooxygenase-2 *in vitro* (Oei et al., 2019). Given parenterally before tumour surgery, mistletoe extracts reduce natural killer cell and neutrophil suppression (Oei et al., 2019). A recent meta-analysis found no safety signals in 10 controlled trials using mistletoe extracts in the perioperative oncological setting (Cogo et al., 2023). The two included randomized controlled trials reporting on physical function showed an improvement in the mistletoe extract group. These results support the evidence from a randomized controlled trial performed in 220 postoperative pancreatic cancer patients, in which mistletoe extracts improved physical function in a clinically and statistically significant manner, together with seven of nine EORTC QLQ-C30 symptom scales (Tröger et al., 2014). Two follow-up randomized controlled trials have been performed in the same population with 290 participants each (EudraCT numbers: 2014-002386-30, 2014-004552-64) and are expected to be published soon.

Keywords: *Viscum album*, immunomodulation, anti-inflammation, cancer surgery, physical function

References

- Cogo, E., Elsayed, M., Bhardwaj, S., Cooley, K., Aycho, C., Liang, V., Papadogianis, P. Psihogios, A., Seely, D., 2023. Mistletoe extracts during the oncological perioperative period: a systematic review and meta-analysis of human randomized controlled trials. *Curr. Oncol.* 30, 8196-8219.
- Lachmann, G., von Haefen, C., Kurth, J., Yuerek, F., Spies, C., 2018. Innate immunity recovers earlier than acquired immunity during severe postoperative immunosuppression. *Int. J. Med. Sci.* 15(1), 1-9.
- Oei, S.L., Thronicke, A., Schad, F., 2019. Mistletoe and immunomodulation: Insights and implications for anticancer therapies. *Evid. Based Complement. Alternat. Med.* 2019.
- Tröger, W., Galun, D., Reif, M., Schumann, A., Stankovic, N., Milicevic, M., 2014. Quality of life of patients with advanced pancreatic cancer during treatment with mistletoe: A randomized controlled trial. *Dtsch. Arztebl. Int.* 111, 493-502.
- Urech, K., Baumgartner, S., 2015. Chemical constituents of *Viscum album* L.: Implications for the pharmaceutical preparation of mistletoe. In Chan S. H. H. (Ed) *Transl. Res. Biomed.* 4, 11-23.

* Presenting author. E-mail address: florian.pelzer@iscador.com (Florian Pelzer)

S2.P90 Potential benefits of *Malva sylvestris* in treating dry-eye pathology *in vitro*

Alexander Areesanan^{a,*}, Sven Nicolay^a, Morris Keller^b, Amy Marisa Zimmermann-Klemd^a, Olivier Potterat^b, Carsten Gründemann^a

^a Translational Complementary Medicine, Department of Pharmaceutical Sciences, University of Basel, Campus Rosental – Mattenstrasse 22, CH-4058 Basel, Switzerland

^b Division of Pharmaceutical Biology, University of Basel, Klingelbergstrasse 50, CH-4056 Basel, Switzerland

Given to the exposure to the pollution, digital lifestyle, and ultraviolet (UV) radiation, the worldwide prevalence of dry eye disease (DED) has increased (Patel et al., 2023). The grittiness and burning sensations of DED deteriorate one's quality of life immensely, making it a common cause of visits to ophthalmologists (Tovar et al., 2023). This study investigated the medical potential of *Malva sylvestris* flower extracts via *in vitro* cellular systems capable of simulating DED pathology.

M. sylvestris flower extract, its mucilage and polyphenol fractions facilitated 2D wound healing in human corneal epithelial cell line (HCE-T) and reduced intracellular reactive oxygen species when exposed to UVB at 30 µg/mL. Significant superoxide radical-scavenging properties were found to be prominent in flower extract and polyphenol fraction. The LPS-exposed monocytic (THP-1) cells when treated with polyphenol fraction lowered NF-κB expression at 50 µg/mL in a dose-dependent manner as well as significantly lowered pro-inflammatory cytokines (IL-6, IL-8, TNF-α) in the supernatants. The fraction was also demonstrated to diminish calcium influx in Jurkat cells. Moreover, one of the most crucial factors in DED development is a reduction of the tear film's lipid layer. Stimulation of lipid production was observable following the exposure to the flower extracts and mucilage fraction in an immortalized human meibomian gland epithelial cell (IHMGEC). Overall, the results indicate that the tested plant extracts interfere with the function of cells involved in DED pathology, thus providing a rationale for their potential therapeutic use. However, further investigation is needed to fully validate this hypothesis.

Keywords: *Malva sylvestris*, dry eye disease, reactive oxygen species, cytokines, lipid production

References

- Patel, S., Mittal, R., Kumar, N., Galor, A., 2023. The environment and dry eye—manifestations, mechanisms, and more. *Front Toxicol.* 5, 1173683.
- Tovar, A.A., Frankel, S.T., Galor, A., Sabater, A.L., 2023. Living with dry eye disease and its effects on quality of life: Patient, optometrist, and ophthalmologist perspectives. *Ophthalmol. Ther.* 12(5), 2219-2226.

* Presenting author. E-mail address: alexander.areesanan@unibas.ch (Alexander Areesanan)

S2.P91 The relationship among essential oil composition, morphological and climatic diversification - the case of *Teucrium montanum* from the Balkan Peninsula

Mirjana Marčetić^{a,*}, Miloš Zbiljić^b, Danilo Stojanović^b, Branislava Lakušić^b, Dmitar Lakušić^c

^a Department of Pharmacognosy, University of Belgrade-Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

^b Department of Botany, University of Belgrade-Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

^c Institute of Botany and Botanical Garden "Jevremovac", University of Belgrade-Faculty of Biology, Takovska 43, 11000 Belgrade, Serbia

Teucrium montanum L. (Lamiaceae) is a small, evergreen, semi-woody shrub that has a wide ecological preference and grows in various environments. The Balkan Peninsula is characterized by a complex geological history and very heterogeneous climatic conditions in a relatively small area. *Teucrium montanum* shows high morphological and anatomical variability in the Balkan Peninsula and seven morphological groups have been recognized (Zbiljić et al., 2023). The aim of our work was to determine whether the phytochemical variability of essential oils follows morphological diversification and to determine the influence of climatic conditions. Isolated essential oils from 46 populations were analysed by GC/FID/MS and multivariate statistical analysis. The main compounds in the investigated oils were: sesquiterpenes germacrene D-4-ol, shyobunol, *cis*- sesquiterpene hydrate, α -bisabolol, germacrene D and monoterpene limonene. Principal component analysis (PCA) showed significant chemical variability in the analysed samples, with the morphological group "montanum" showing the greatest changes, similar to the pronounced morphological variations of this group. In addition, there were specific compounds for the morphological groups that led to a clear separation in the discriminant analysis (CDA). Certain morphological groups ("parnassicum", "skadarensis" and "pannonicum") can be easily distinguished according to both phytochemical and morphological characteristics. The biosynthesis of various compounds was influenced by the climatic conditions. The lack of water caused the increase of limonene content in the population influenced by arid continental climate and of cryptomerione and (*E*)-nerolidol affected by arid submediterranean climate. However, it was found that phytochemical diversification followed morphological rather than climatic variability.

Keywords: *Teucrium montanum*, Lamiaceae, essential oil, morphological variability, climatic conditions

References

Zbiljić, M., Lakušić, B., Kuzmanović, N., Stojanović, D., Lakušić, D., 2023. Morphological diversification of *Teucrium montanum sensu lato* on the Balkan Peninsula. *Plant Biosystems*, 157(3), 670-687.

* Presenting author. E-mail address: mirjana.marcetic@pharmacy.bg.ac.rs (Mirjana Marčetić)

S2.P92 Salicylic alcohol content in the leaves of different *Salix* species and individuals: impact on pharmaceutical potential

Leonie Kayser^{a,*}, Thomas Olaf Gruber^a, Gregor Aas^b, Guido Jürgenliemk^a

^a Department of Pharmaceutical Biology, University of Regensburg, Universitaetsstraße 31, 93053 Regensburg, Germany

^b Ecological-Botanical Gardens Bayreuth, University of Bayreuth, Universitaetsstraße 30, 95447 Bayreuth, Germany

The genus *Salix* belongs to the family of Salicaceae and comprises around 500 different species (Meier et al., 1998). The bark is used traditionally in European countries due to its antiphlogistic, antipyretic, and analgesic efficacy (März et al., 2002). The Ph. Eur. does not define a species for the pharmaceutical use of Willow bark, but a minimum content of 1.5% total salicylic derivatives is required (Ph. Eur., 2023). The salicylic alcohols are metabolized *in vivo* to salicylic acid, the pharmacologically active form (März et al., 2002). Aim of the present study was whether *Salix* leaves could also be of pharmaceutical interest concerning their salicylic alcohol content. For that, the leaves of 12 different *Salix* species (*Salix aurita*, *S. bicolor*, *S. caesia*, *S. caprea*, *S. cinerea*, *S. daphnoides*, *S. fragilis*, *S. hastata*, *S. lapponum*, *S. purpurea*, *S. viminalis*, *S. x sepulcralis*) with 42 individuals were collected in the months of May-September from 2018-2019. The samples were analyzed for their phenolic profile and especially salicylic alcohol content using an RP18-UPLC[®] - PDA method (Wiesneth, 2017). Furthermore, inter-individual differences and differences between the sexes within a species were also of interest. *S. caesia*, *S. caprea*, *S. daphnoides*, *S. hastata*, *S. lapponum*, *S. viminalis*, and *S. x sepulcralis* contained hardly any relevant amounts of salicylic alcohol derivatives whereas *S. aurita*, *S. bicolor*, *S. cinerea*, *S. fragilis*, and *S. purpurea* contained more than 1.5%. Barely any differences were found between the sexes. This indicates that also the leaves of some *Salix* species could be of pharmaceutical interest.

Keywords: *Salix*, salicaceae, salicylic alcohol, Ph. Eur., European Pharmacopoeia

References

- Meier, B., Meier-Liebi, M., 1998. *Salix*, in: Blaschek, W., Hänsel, R., Keller, K., Reichling, J., Rimpler, H., Schneider, G., Hagers Handbuch der pharmazeutischen Praxis, Springer, Berlin, Heidelberg, New York, pp. 469-496.
- März, R. W., Kemper, F., 2002. Weidenrindenextrakt – Wirkungen und Wirksamkeit. Erkenntnisstand zu Pharmakologie, Toxikologie und Klinik. Wien. Med. Wochenschr. 15/16/2002, 354-359.
- Ph. Eur., 2023. Salicis cortex, in: European Pharmacopoeia 11.0, Deutscher Apotheker Verlag, Stuttgart, pp. 1583. Wiesneth, S., 2017. Phytochemische Untersuchung des phenolischen Inhaltsstoffspektrums in *Salix* Spezies unter besonderer Berücksichtigung der Flavan-3-ole (Doctoral Dissertation), University Regensburg, Germany, pp. 246-248.

* Presenting author. E-Mail address: Leonie.Kayser@chemie.uni-regensburg.de (Leonie Kayser)

S2.P93 Plant metabolites as glucose-induced insulin secretion modulators

Andrzej Patyra^{a,b,*}, Justine Vaillé^b, Jérémie Neasta^b, Anna Karolina Kiss^a, Catherine Oiry^b

^a Department of Pharmaceutical Biology, Medical University of Warsaw, 1 Banacha Str., 02-097 Warsaw, Poland

^b Institut des Biomolécules Max Mousseron, Université de Montpellier, CNRS, ENSCM, 1919 route de Mende, 34293 Montpellier, France

Type 2 diabetes mellitus (T2DM) is a disease that associates insulin resistance and pancreatic β -cell dysfunction. Based on the current understanding of the pathophysiology of this disease, multiple anti-diabetic therapies have been developed to improve glycemic control and slow disease progression. Nevertheless, they have limited effectiveness and numerous side effects. In this context, plant extracts and their active constituents are considered an important area of seeking new anti-diabetic treatments (Irondi et al., 2015). Research suggests that bioactive compounds (i.e., polyphenols) found in plants may help slow the aging process and reduce the risk of many chronic diseases, including type 2 diabetes mellitus. In this context, plant secondary metabolites are likely good candidates to protect β cells against dysfunction and regulate insulin secretion since several of these well-known antioxidant compounds are now widely regarded as pharmacological agents that regulate target proteins and signaling pathways in pancreatic β cell (Kasole et al., 2019). In this study flavonoids, lignans and coumarins (at 20 $\mu\text{mol/L}$) were screened for their ability to modulate β -cell function on INS-1 cell model in glucose-stimulating conditions. Insulin release was quantified by the homogeneous time-resolved fluorescence method. Some of the flavonoids and coumarins increased glucose-induced insulin secretion, while no such effect was observed for lignans. Modulation of the pancreatic β -cell function by plant secondary metabolites was highly dependent on their structure, suggesting that this activity is based on molecule-protein interaction rather than on antioxidative potential. Structure-activity relationship conclusions were drawn from this, leading to the identification of key substitution patterns.

Keywords: diabetes, insulin secretion, coumarins, flavonoids, lignans

References

- Irondi, E.A., Oboh, G., Akindahunsi, A.A., Boligon, A.A., Athayde, M.L., 2015. Phenolics composition and antidiabetic property of *Brachystegia eurycoma* seed flour in high-fat diet, low-dose streptozotocin-induced type 2 diabetes in rats. *Asian Pacific J. Trop. Dis.* 5, 59-165.
- Kasole, R., Martin, H.D., Kimiywe, J., 2019. Traditional Medicine and Its Role in the Management of Diabetes Mellitus: "Patients' and Herbalists' Perspectives". *Evid Based Complement. Alternat. Med.* 2019, 2835691.

* Presenting author. E-mail address: andrzej.patyra@wum.edu.pl (Andrzej Patyra)

S2.P94 Antioxidative capacity of European tree bark extracts

Sissy Häsler Gunnarsdottir^{a,*}, Gertie Janneke Oostingh^a, Lena Haretzmüller^b, Thomas Schnabel^c, Anja Schuster^a

^a Salzburg University of Applied Sciences, Health Sciences Department, Urstein Sued 1, 5412 Puch/Salzburg, Austria

^b Johann Kepler University, Analytical Chemistry Department, Altenberger Straße 69, 4040 Linz, Austria

^c Salzburg University of Applied Sciences, Green Engineering and Circular Design Department, Markt 136a, 5431 Kuchl, Austria

Tree bark was found to be a rich natural source for active antioxidant compounds (Dudonné et al. 2009) but there is little information on the effects of European tree bark. Within the presented study, we analyzed the potential of different tree bark extracts derived from European trees (oak, beech, birch, alder, pine and bird cherry) for their antioxidant capacity. Total phenolic content (TPC) and radical scavenging ability (DPPH) of the extracts was determined as well as their potential to reduce oxidative stress in a human keratinocyte cell line. For the latter, reactive oxygen species (ROS) production (using the DCFDA assay), lipid peroxidation (using the TBARS assay), GSH/GSSG ratio, and protein carbonylation were compared between cells treated with different concentrations of bark extracts and untreated cells. Within the chemical assays, oak bark extracts had the strongest anti-oxidant potential with a TPC of 366.5 µgGAE/mg and an DPPH inhibition of 76 %. For all wood extracts, a significantly lowered generation of ROS could be shown with the DCFDA method in the range of 50 to 600 µg/mL (except for beech at 50 µg/mL) compared to the untreated cells. The TBARS assay shows a strong tendency for alder bark extract to be the most promising agent, but this is currently being verified. GSH/GSSG ratio and protein carbonylation are still to be examined. Combined with their antimicrobial and wound-healing activity (Emrich et al. 2022), tree bark extracts could be promising compounds in the treatment of various skin diseases (Häsler Gunnarsdottir et al. 2023).

Keywords: European bark extracts, antioxidative capacity, skin applications

References

- Dudonné, S., Vitrac, X., Coutière, P., Woillez, M., Mérillon, J.-M., 2009. Comparative study of antioxidant properties and total phenolic content of 30 plant extracts of industrial interest using DPPH, ABTS, FRAP, SOD, and ORAC assays. *J. Agric. Food Chem.* 57(5), 1768–1774.
- Emrich, S., Schuster, A., Schnabel, T., Oostingh, G.J., 2022. Antimicrobial Activity and Wound-Healing Capacity of Birch, Beech and Larch Bark Extracts. *Molecules* 27(9).
- Häsler Gunnarsdottir, S., Sommerauer, L., Schnabel, T., Oostingh, G.J., Schuster, A. 2023. Antioxidative and antimicrobial evaluation of bark extracts from common european trees in light of dermal applications. *Antibiotics (Basel)* 12(1).

* Presenting author. E-mail address: sissy.haeslergunnarsdottir@fh-salzburg.ac.at (Sissy Häsler Gunnarsdottir)

S2.P95 Exploring the potential of ADAPT-232: plant adaptogens for healthy aging, and longevity in *Caenorhabditis elegans*

Monika N. Todorova^a, Biser K. Binev^{a,*}, Milen I. Georgiev^{a,b}

^a Laboratory of Metabolomics, Institute of Microbiology, Bulgarian Academy of Sciences, 139 Ruski Blvd., 4000 Plovdiv, Bulgaria

^b Department of Plant Cell Biotechnology, Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

In the ongoing scientific exploration of strategies to promote longevity and rejuvenation, there is a burgeoning interest in supplements aimed at enhancing healthspan and extending lifespan (Todorova et al., 2024). Plant adaptogens, revered for their multifaceted physiological benefits, have garnered attention as prospective contributors to this research trajectory (Vasileva et al., 2019). Among the most recognized adaptogens are the extracts from *Eleutherococcus senticosus* root, *Schisandra chinensis* berry, and *Rhodiola rosea* root, which constitute the herbal medicinal product ADAPT-323. This supplement has been investigated for its energy-boosting and stress-protective properties (Aslanyan et al., 2010; Panossian et al., 2021). We conducted an assessment of the potential of ADAPT-232 to modulate healthy aging and increase energy using *Caenorhabditis elegans*. This model system serves as a valuable model for studying age-related degradation at the tissue and molecular levels. Furthermore, nematodes exhibit diminished late-life fitness associated with muscle and neuronal degeneration. In our study, we focused on understanding the intricate interplay between ADAPT-232 and the physiological processes that influence aging. Through a comprehensive analysis employing both phenotypic and stress assays, we have determined that ADAPT-232 is a promising supplement for maintaining metabolism and homeostasis throughout lifespan. The combined effects of the three adaptogenic plants in ADAPT-232 contribute significantly to the extension of the healthspan of *C. elegans* through modulation of stress-response mechanisms. Our findings highlight the potential of plant adaptogens, particularly ADAPT-232, as a promising avenue in the pursuit of interventions that promote healthy aging and increased longevity.

Keywords: adaptogens, healthspan, *Caenorhabditis elegans*, ageing, longevity

References

- Aslanyan, G., Amroyan, E., Gabrielyan, E., Nylander, M., Wikman, G., Panossian, A., 2010. Double-blind, placebo- controlled, randomised study of single dose effects of ADAPT-232 on cognitive functions. *Phytomedicine*. 17, 494-499. Panossian, A.G., Efferth, T., Shikov, A.N., Pozharitskaya, O.N., Kuchta, K., Mukherjee, P.K., Banerjee, S., Heinrich, M., Wu, W., Guo, D.A., Wagner, H., 2021. Evolution of the adaptogenic concept from traditional use to medical systems: Pharmacology of stress- and aging-related diseases. *Med. Res. Rev.* 630-703.
- Todorova, M.N., Savova, M.S., Mihaylova, L.V., Georgiev, M.I., 2024a. Nurturing longevity through natural compounds: Where do we stand, and where do we go? *Food Frontiers*. 5, 267-310.
- Vasileva, L.V., Ivanovska, M.V., Murdjeva, M.A., Saracheva, K.E., Georgiev, M.I., 2019. Immunoregulatory natural compounds in stress-induced depression: An alternative or an adjunct to conventional antidepressant therapy? *Food Chem. Toxicol.* 127, 81-88.

* Presenting author. E-mail address: biser.binev@abv.bg (Biser K. Binev)

S2.P96 *Harpagophytum procumbens* DC. ex Meisn. hairy roots extract alleviates inflammatory response in psoriasis dermatitis

Ivanka K. Koycheva^{a,*}, Blagovesta D. Boneva^c, Emiliya Z. Stoyanova^c, Martina S. Savova^{a,b}, Iva D. Stoykova^{a,b}, Biser Binev^a, Nikolina M. Mihaylova^d, Milen I. Georgiev^{a,b}

^a Laboratory of Metabolomics, Department of Biotechnology, Institute of Microbiology, Bulgarian Academy of Sciences, 4000 Plovdiv, Bulgaria

^b Department of Plant Biotechnology, Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

^c Laboratory of Experimental Immunology, Department of Immunology, Institute of Microbiology, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria

^d Laboratory of Immunohistochemistry and Immunopathology, Department of Immunology, Institute of Microbiology, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria/

Psoriasis is an inflammatory skin pathology affecting 2-3% of the worlds' population. The disease manifests itself with characteristic skin lesions and is associated with multiple comorbidities and significant negative impact on patients' quality of life (Griffiths et al., 2021). Plant-derived bioactive compounds are promising as adjuncts to the conventional pharmacotherapy (Koycheva et al., 2021a, 2021b). We have investigated the anti-inflammatory and anti-psoriatic potential of the extract of transformed hairy root culture of *Harpagophytum procumbens* DC. ex Meisn. (HPE-HRs) and its secondary metabolite verbascoside on imiquimod-induced psoriasiform model in mice. The *in vitro* effects of HPE-HRs and verbascoside on the expression of the mouse chemokine receptors CXCR3, TGFβRI, as well as the production of the cytokines IFN-γ, IL-17F were analyzed by flow cytometry in skin- and spleen-derived T-lymphocytes, respectively. Concentration-dependent decrease was observed in the percentage of immune cells expressing CXCR3 and TGFβRI receptors and no induction of the inflammatory cytokines secretion upon HPE-HRs or VER treatment. Topical application of a cream containing HPE-HRs (1.25, 2.5 and 5%) on the shaved backs of C57BL/6 mice was done over the period of imiquimod-induced psoriasiform dermatitis in mice. The 5% HPE-HRs cream ameliorates psoriasis-like skin manifestations by an alleviation of epidermal hyperplasia, reduction of skin thickness and scale formation compared to the control groups. Further, gene expression analysis by RT-qPCR was performed to assess the changes in critical signalling pathways related to psoriasis development. The results suggest that HPE-HRs may serve as a promising therapeutic option in psoriasis management.

Keywords: psoriasis, *Harpagophytum procumbens*, CXCR3, TGFβRI, epidermal hyperplasia

References

Griffiths, C.E.M., Armstrong, A.W., Gudjonsson, J.E., Barker, J.N.W.N., 2021. Psoriasis. *Lancet*. 397, 1301-15. Koycheva, I.K., Vasileva, L.V., Amirova, K.A., Marchev, A.S., Balcheva-Sivenova, Z.P., Georgiev, M.I., 2021a. Biotechnologically produced *Lavandula angustifolia* Mill. extract rich in rosmarinic acid resolves psoriasis-related inflammation through Janus kinase/signal transducer and activator of transcription signaling. *Front. Pharmacol.* 12, 680168.

Koycheva, I.K., Vasileva, L.V., Amirova, K.A., Marchev, A.S., Balcheva-Sivenova, Z.P., Georgiev, M.I., 2021b. Leucosceptoside A from Devil's claw modulates psoriasis-like inflammation via suppression of the PI3K/AKT signaling pathway in keratinocytes. *Molecules*. 26(22), 7014.

* Presenting author. E-mail address: vkoy4eva@abv.bg (Ivanka Koycheva)

S2.P97 Caloric restriction mimicking through NRF-2-mediated mechanisms by plant-derived bioactive leads

Liliya V. Mihaylova^{a,b,*}, Martina S. Savova^{a,b}, Monika N. Todorova^b, Biser K. Binev^b, Milen I. Georgiev^{a,b}

^a Department of Plant Cell Biotechnology, Center of Plant Systems Biology and Biotechnology, 14 St. Knyaz Boris I Pokrastitel Street, Plovdiv, Bulgaria

^b Laboratory of Metabolomics, Institute of Microbiology, Bulgarian Academy of Sciences, 139 Ruski Boulevard, Plovdiv, Bulgaria

Obesity is a multifactorial disease associated with metabolic dysregulations, which could lead to multimorbidity affecting both the quality of life and the lifespan (Savova et al., 2023; Vasileva et al., 2020). Caloric restriction initiates the adaptive metabolic strategies that produce stress resilience and the nuclear factor erythroid 2-related factor 2 (NRF2) is known as a master regulator in the integration of these signaling pathways (Spadaro et al., 2022). Several natural products commonly used as dietary supplements (resveratrol, spermidine or rapamycin) have been defined as caloric restriction mimetics (CRM) due to the overlapping molecular mechanisms involved in their biological activity and caloric restriction-mediated healthspan (Madeo et al., 2019). For instance, curcumin is among the most potent natural NRF2 activators that has been proposed as a CRM. As an advancement in our continuing search for potent anti-obesogenic compounds from plants we have directed our focus toward the potential of natural NRF2 activators to promote improved metabolic homeostasis. We have evaluated the changes in *skn-1* (the orthologue of NRF2) transcriptional activity in *Caenorhabditis elegans* induced upon high glucose feeding and the potential of curcumin to modulate this process. The confocal GFP-based fluorescent detection revealed a 2-fold increase in *skn-1*/NRF2 upon curcumin (100 µM) supplementation that was further supported by the mRNA expression analysis. The obtained findings reinforced the idea that NRF2 serves as a hormetic mediator that not only provides integration of detoxification processes but also could account as a target to alleviate the severity of age-related illnesses including obesity and metabolic dysregulations.

Keywords: caloric restriction, obesity, aging, bioactive compounds, NRF-2

References

- Madeo F., Carmona-Gutierrez D., Hofer S.J., Kroemer G., 2019. Caloric restriction mimetics against age-associated disease: Targets, mechanisms, and therapeutic potential. *Cell Metab.* 29, 592-610.
- Savova M.S., Mihaylova L.V., Tews D., Wabitsch M., Georgiev M.I., 2023. Targeting PI3K/AKT signaling pathway in obesity. *Biomed. Pharmacother.* 159, 114244..
- Spadaro, O., Youm, Y., Shchukina, I., Artyomov, M.N., Dixit, V.D., 2022. Caloric restriction in humans reveals immunometabolic regulators of health span. *Science* 375, 671-677.
- Vasileva, L.V., Savova, M.S., Amirova, K.M., Dinkova-Kostova, A.T., Georgiev, M.I., 2020. Obesity and NRF2- mediated cytoprotection: Where is the missing link? *Pharmacol. Res.* 156, 104760.

* Presenting author. E-mail address: liliya.vl.mihaylova@gmail.com (Liliya V. Mihaylova)

S2.P98 *SKN1/NRF2*-dependent modulation of nutrient sensing network in *Caenorhabditis elegans* upon curcumin treatment

Martina S. Savova^{a,b,*}, Liliya V. Mihaylova^{a,b}, Monika N. Todorova^a, Milen I. Georgiev^{a,b}

^a Laboratory of Metabolomics, Institute of Microbiology, Bulgarian Academy of Sciences, 139 Ruski Blvd., 4000 Plovdiv, Bulgaria

^b Department of Plant Cell Biotechnology, Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

Metabolic dysregulations that root from obesity affect the nutrient sensing signaling pathways and mitochondrial homeostasis. The nuclear factor erythroid 2-related factor 2 (NRF-2) acts as a master regulator in cellular stress response and is suggested to be involved in obesity, diabetes, and metabolic syndrome (Vasileva et al. 2020). Curcumin and some of its derivatives are recognized as natural activators of NRF-2 and exhibit anti-inflammatory, antioxidant, chemopreventive, anti-nociceptive, anti-proliferative, and anti-parasitic properties (Shin et al., 2020). In the context of obesity and metabolic disorders, curcumin has been reported to inhibit adipogenesis, regulate lipid metabolism, stimulate energy expenditure, and benefit gut microbiota (Sampath et al., 2021). The current investigation provides further insights into the molecular mechanism of the beneficial effect of curcumin in modulating glucose-induced lipid accumulation in *Caenorhabditis elegans*. Moreover, we applied orlistat as an anti-obesogenic control, which is an approved drug for obesity therapy. Our preliminary experiments in *C. elegans* obesity model suggest that a combination of orlistat (12 μ M) and curcumin (100 μ M) significantly lowers accumulated lipids by 57.6% compared to vehicle group). Regarding the molecular mechanisms of curcumin as an NRF2- modulating caloric restriction mimetic agent, our results demonstrate that curcumin elevates the expression of *skn-1*, *sek-1* and *nhr-49*, which are important and evolutionary conserved genes involved in carbohydrate and lipid metabolism in *C. elegans*. Additionally, modulation of mitochondrial dynamics, including mitochondrial potential, was evaluated. Based on these results, we conclude that curcumin exerts its favorable biological activity through modulation of complex nutrient sensing signaling pathways.

Keywords: NRF-2, *Caenorhabditis elegans*, Curcumin, Caloric restriction mimetics, Obesity

References

- Sampath, C., Wilus, D., Tabatabai, M., Freeman, M.L., Gangula, P.R., 2021. Mechanistic role of antioxidants in rescuing delayed gastric emptying in high fat diet induced diabetic female mice. *Biomed. Pharmacother.* 137, 111370. Shin, J.W., Chun, K.-S., Kim, D.-H., Na, H.-K., Surh, Y.-J., 2020. Curcumin induces stabilization of Nrf2 protein through Keap1 cysteine modification. *Biochem. Pharmacol.* 173, 113820.
- Vasileva, L.V., Savova, M.S., Amirova, K.M., Dinkova-Kostova, A.T., Georgiev, M.I., 2020. Obesity and NRF2- mediated cytoprotection: Where is the missing link? *Pharmacol. Res.* 156, 104760.

* Presenting author. E-mail address: m.sav@abv.bg (Martina S. Savova)

S2.P99 The phenylethanoid glycoside forsythoside B as a promising candidate for photoprotective dermatological products

Iva D. Stoykova^{a,b,*}, Ivanka K. Koycheva^b, Liliya V. Mihaylova^{a,b}, Milen I. Georgiev^{a,b}

^a Laboratory of Metabolomics, Department of Biotechnology, Institute of Microbiology, Bulgarian Academy of Sciences, 4000 Plovdiv, Bulgaria

^b Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

Complex internal and extrinsic variables accelerate skin aging. The intrinsic aging leading to maturation processes is driven by epigenetic changes, circadian rhythm disruption, and senescence that impair skin texture. Premature skin aging caused by external factors, the main of which is chronic sun exposure, also known as photoaging, leads to irreversible photodamage and reduces skin capacity to self-renew (Stoykova et al., 2024; Zhang et al., 2023). Products that include sunscreen compounds and interfere with metabolism can be harmful and frequently cause serious cutaneous adverse effects such as irritations and allergy (Dhaliwal et al., 2019; Yamada et al., 2020). Over the past two decades, the scientific community has become increasingly interested in the “green pharmacy”, where natural ingredients are evaluated for potential use as cosmetic agents (Dhaliwal et al., 2019). The aim of our research is to determine the impact of ultraviolet radiation (UVA/UVB) on photoaged human keratinocytes and to prevent photodamage by application of plant-derived bioactive compounds. Forsythoside B (FORB), found in *Verbascum xanthophoeniceum* Griseb, was previously reported to have an anti-inflammatory effect in primary human epidermal keratinocytes (Georgiev et al., 2012). We determined the cytotoxic and photoprotective potential of FORB which is the primary aspect in personal care product. Identified as a secondary factor of photodamage, the high levels of reactive oxygen species (ROS) induced by the UV irradiation were measured by confocal-fluorescence microscopy. The obtained data exposed that FORB significantly inhibited the increased levels of ROS induced by UVA/UVB in HaCaT cells, which may be related to its photoprotective activity.

Keywords: Forsythoside B, keratinocytes, photoprotection, photoaging

References

- Dhaliwal, S., Rybak, I., Ellis, S.R., Notay, M., Trivedi, M., Burney, W., Vaughn A.R., Nguyen, M., Reiter, P., Bosanac, S., Yan, H., Foolad, N., 2019. Sivamani, R.K. Prospective, randomized, double-blind assessment of topical bakuchiol and retinol for facial photoageing. *Br. J. Dermatol.* 180, 289-296.
- Georgiev, M., Pastore, S., Potapovich, A., Panetta M., Korkina, L., 2012. *Verbascum xanthophoeniceum*-derived phenylethanoid glycosides are potent inhibitors of inflammatory chemokines in dormant and interferon-gamma- stimulated human keratinocytes. *J. Ethnopharmacol.* 144, 754-760.
- Stoykova, I.D., Koycheva, I.K., Binev, B.K., Mihaylova, L.V., Georgiev, M.I., 2024. Molecular approaches to prevent UV-induced premature skin aging: focus on phytochemicals as photo-protectants. *Phytochem. Rev.*, Submitted.
- Yamada, M., Mohammed, Y., Prow, W.T., 2020. Advances and controversies in studying sunscreen delivery and toxicity. *Adv. Drug Deliv. Rev.* 153, 72-86.
- Zhang, C., Gao, X., Li, M., Yu, X., Huang, F., Wang, Y., Yan, Y., Zhang, H., Shi, Y., He, X., 2023. The role of mitochondrial quality surveillance in skin aging: Focus on mitochondrial dynamics, biogenesis and mitophagy. *Ageing Res. Rev.* 87, 101917.

* Presenting author. E-mail address: ivastoykova_af@mail.bg (Iva D. Stoykova)

S2.P100 Exploring the therapeutic potential of *Punica granatum* leaves extract in enhancing healthspan and lifespan in *Caenorhabditis elegans*

Monika N. Todorova^{a,*}, Martina S. Savova^{a,b}, Biser K. Binev^a, Liliya V. Mihaylova^{a,b}, Milen I. Georgiev^{a,b}

^a Laboratory of Metabolomics, Institute of Microbiology, Bulgarian Academy of Sciences, 139 Ruski Blvd., 4000 Plovdiv, Bulgaria

^b Department of Plant Cell Biotechnology, Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

In recent decades, substantial efforts have been devoted to enhancing both healthspan and lifespan encompassing pharmacological and non-pharmacological interventions (Todorova et al., 2024a). Moreover, the exploration of natural products that modulate the molecular network associated with lifespan holds significant promise in the pursuit of longevity and disease-free years (Todorova et al., 2024b). *Punica granatum* L., commonly known as pomegranate, has served as a nutritional source and a traditional remedy for centuries. Abundant in bioactive compounds, predominantly flavonols and flavones, pomegranate leaves have been renowned for their anti-inflammatory and anti-diabetic properties (Abo-Saif et al., 2023; Nasir et al., 2024). However, there remains a gap in comprehending their role in modulating healthspan and longevity. Hence, our study has delved into the capacity of a crude extract from pomegranate leaves to mitigate the aging process using *Caenorhabditis elegans* model system. Our investigation revealed that the extract, in doses 50, 100 and 200 µg/mL, enhances resistance to stressors, including heat and oxidative stress. Supplementation with 200 µg/mL extract significantly increases survival upon heat stress to up to 85% and 78% on day 5 and day 10 of the nematodes' lifespan, respectively, compared to the control group. Moreover, the extract improves locomotion, as the mean bending rate upon 200 µg/mL was increased by up to 106% (day 5) and 110% (day 10), indicating a positive impact on neuronal and muscular function. Treatment of *C. elegans* with the highest concentration resulted in a mean lifespan of 24.8 days, which is 1.62-fold change in comparison to the control animals. In conclusion, our findings underscore the potential of *P. granatum* leaves extract in promoting healthspan and longevity, and its positive impact at both the molecular and functional levels.

Keywords: Longevity, Healthspan, *Caenorhabditis elegans*, *Punica granatum*, Pomegranate

References

- Abo-Saif, M.A., Ragab, A.E., Ibrahim, A.O., Saber-Ayad, M., El-Feky, O.A., 2023. Pomegranate peel extract protects against the development of diabetic cardiomyopathy in rats by inhibiting pyroptosis and downregulating LncRNA- MALAT1/DNA gyrase: structure and function. *Front. Pharmacol.* 14, 1166653.
- Nasir, A., Afridi, O.K., Ullah, S., Khan, H., Bai, Q., 2024. Mitigation of sciatica injury-induced neuropathic pain through active metabolites derived from medicinal plants. *Pharmacol. Res.* 200, 107076.
- Todorova, M.N., Savova, M.S., Mihaylova, L.V., Georgiev, M.I., 2024a. Nurturing longevity through natural compounds: Where do we stand, and where do we go? *Food Frontiers*. 5, 267-310.
- Todorova, M.N., Savova, M.S., Mihaylova, L.V., Georgiev, M.I., 2024b. Icariin improves stress resistance and extends lifespan in *Caenorhabditis elegans* through hsf-1 and daf-2-driven hormesis. *Int. J. Mol. Sci.* 25, 352.

* Presenting author. E-mail address: mntodorova@yahoo.com (Monika N. Todorova)

S2.P101 Baicalein, a major component of the root of *Oroxylum indicum* (Bignoniaceae) vasodilates isolated pulmonary arteries through endothelium-dependent and -independent mechanisms

Nattiya Chaichamnong^{a,*}, Céline Demougeot^a, Kornkanok Ingkaninan^b, Corine Girard^a, Perle Totoson^a

^a Université de Franche-Comté, EFS, INSERM, UMR RIGHT, F-25000 Besançon, France

^b Centre of Excellence in Cannabis Research, Department of Pharmaceutical Chemistry and Pharmacognosy, Faculty of Pharmaceutical Sciences, Naresuan University, Phitsanulok 65000, Thailand

Recently, we demonstrated that *Oroxylum indicum* (L.) Kurz (Bignoniaceae) root extract (OIRE) and its major constituent baicalein display phosphodiesterase-5 inhibitory activities (unpublished data), suggesting their potential use for treating pulmonary artery hypertension (PAH). The present study investigated the vasorelaxant effects of OIRE and baicalein on isolated pulmonary arteries (PA) compared to a conduit vessel from the systemic circulation (aorta). OIRE was prepared by maceration with 95% ethanol and baicalein was purchased from Sigma. Vascular reactivity was explored on isolated aorta and PA from Wistar rats. After pre-constriction with phenylephrine, vessels were exposed to cumulative concentrations of OIRE or baicalein. To unravel the role of endothelium-dependent and -independent pathways in PA relaxation, the same experiments were performed in the presence of specific inhibitors. OIRE and baicalein both induced a vasorelaxation on PA, greater than in the aorta ($E_{\max}=85\pm5\%$ vs $25\pm4\%$, $p<0.001$ for OIRE, and $E_{\max}=86\pm10\%$ vs $32\pm7\%$ for baicalein $p<0.001$). The EC_{50} values in PA and aorta were 0.035 and 0.018 $\mu\text{g/mL}$ for baicalein, and 7.13 and 0.59 $\mu\text{g/mL}$ for OIRE. Mechanistically, the effects of OIRE on PA relied on eNOS and COX activation, and the production of endothelium-derived hyperpolarizing factors. Baicalein-induced vasorelaxation involved the same endothelial pathways as OIRE but also endothelium-independent mechanisms such as α_1 -adrenergic receptor antagonism, K_v^+ channels blockade, and a decrease in intracellular Ca^{2+} release. The present study identified new pharmacological properties of OIRE and baicalein as PA vasodilators. In combination with their phosphodiesterase-5 inhibitory activities, our data encourage the investigation of their effects on PAH.

Keywords: *Oroxylum indicum*, baicalein, vasorelaxant effect, pathways

* Presenting author. E-mail address: nattiya.chaichamnong@univ-fcomte.fr (Nattiya Chaichamnong)

S2.P102 Chemical profile and cytotoxic activity of methanol extracts of eight *Geranium* L. species

Milan Ilić^a, Stevan Samardžić^a, Vesna Kojić^b, Dimitar Jakimov^b, Mirjana Marčetić^a, Nada Kovačević^a, Milica Drobac^{a,*}

^a Department of Pharmacognosy, University of Belgrade-Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

^b Oncology Institute of Vojvodina, Faculty of Medicine, University of Novi Sad, Put doktora Goldmana 4, 21204 Sremska Kamenica, Serbia

The aim of this study was to determine the chemical composition and cytotoxic activity of methanol extracts from underground and aerial parts of eight *Geranium* species (*G. macrorrhizum* L., *G. robertianum* L., *G. palustre* Torn, *G. sanguineum* L., *G. columbinum* L., *G. pyrenaicum* Burm. and *G. lucidum* L.). These polyphenol-rich extracts have previously been shown to have significant antioxidant and antimicrobial potential (Ilić et al., 2021). A total of seven phenolic acids (ellagic, gallic, galloylquinic, galloylshikimic, protocatechuic, chlorogenic and brevifolin carboxylic acid), 12 flavonoids (quercetin, quercetin-galloylhexoside, two pentosides of quercetin, quercetin-deoxyhexoside, hyperoside, isoquercitrin, kaempferol, astragalin, astragalin-2''- O-gallate, kaempferol-methylether and kaempferol-dimethylether) and 13 tannins (geraniin and its isomer, corilagin, two anomers of tellimagrandin I and three isomeric compounds, tellimagrandin II, hebulagic acid and 4 galloylglucoses) were identified and quantified by LC-MS analysis. The *in vitro* cytotoxic activity of extracts, geraniin and ellagic acid, was evaluated in the MTT assay against human prostate (PC-3), colon (HT-29), cervical (HeLa) and melanoma (Hs 294T) cancer cell lines. All tested extracts induced a reduction of viability against HeLa cancer cells, and the most effective were the extracts of the underground parts of *G. sanguineum*, *G. pyrenaicum*, *G. columbinum* and *G. macrorrhizum* (IC₅₀ = 2.52-8.94 µg/mL). The extracts showed no significant cytotoxicity against other cells, with the exception of the extract from the aerial parts of *G. lucidum* against HT-29 cells (IC₅₀ = 22.54 µg/mL). Geraniin was also most effective against HeLa cells (IC₅₀ = 12.88 µg/mL), with moderate activity against other cells (IC₅₀ = 44.41-80.94 µg/mL), while ellagic acid showed weaker activity (IC₅₀ = 65.62-221.51 µg/mL).

Keywords: *Geranium* species, LC-MS profile, cytotoxicity

References

Ilić, M., Samardžić, S., Kotur-Stevuljević, J., Ušjak, D., Milenković, M., Kovačević, N., Drobac, M., 2021 Polyphenol rich extracts of *Geranium* L. species as potential natural antioxidant and antimicrobial agents. Eur. Rev. Med. Pharmacol. Sci. 25(20), 6283-6294.

* Presenting author. E-mail address: milica.drobac@pharmacy.bg.ac.rs (Milica Drobac)

S2.P103 The composition of the essential oil of *Pimpinella serbica* and its relationship to other *Pimpinella* species from the central part of Balkan Peninsula

Milica Drobac^{a,*}, Mirjana Marčetić^a, Bojan Zlatković^b, Nada Kovačević^a, Violeta Slavkovska^c

^a Department of Pharmacognosy, University of Belgrade - Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

^b Department of Biology and Ecology, University of Niš, Faculty of Sciences and Mathematics, Višegradska 33, 18000 Niš, Serbia

^c Department of Botany, University of Belgrade - Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

Pimpinella serbica (Vis.) Drude (Apiaceae), is an endemic species of the Balkan Peninsula, formerly referred to as *Pancicia serbica* Vis. and considered a monotypic genus (Nikolić, 1973), while in the current taxonomic classification it is included in the genus *Pimpinella* L. (Euro+Med 2014). A number of plants in this genus are used for their medicinal properties, which are partly due to the presence of essential oils with various pharmacological/biological effects (Nasir and Yabalak, 2021). In this study we investigated the chemical composition of essential oils from different organs of *P. serbica* and their relationship with the previously analyzed essential oils of *P. tragiunum*, *P. saxifraga* (Slavkovska et al., 2022) and *P. alpina*, using cluster analysis. Phenylpropanoids dominated in the essential oils of *P. serbica*, dillapiole in the aerial parts (50.7-63.0%) and nothoapiole in the roots (79.0%) and fruits (73.4%). Although a high amount of nothoapiole (24.0%) was present in one sample of the aerial parts. Phenylpropanoids of pseudoisoeugenol type and trinosesquiterpenes, which are considered chemical markers of *Pimpinella* species (Tabanca et al., 2005), were not present in the *P. serbica* essential oils. Cluster analysis also revealed the separation of *P. serbica* essential oils from the essential oils of other *Pimpinella* species. These results indicate that the chemical profile of the essential oils of *P. serbica* does not support its classification in the genus *Pimpinella*, and suggest the possibility of using the essential oil composition as a chemotaxonomic marker in clarifying the status of this endemic species.

Keywords: *Pimpinella serbica*, essential oil composition, chemotaxonomy

References

- Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity. 2014. – Published at <http://www.europlusmed.org> [accessed February 2024].
- Nasir, A., Yabalak, E., 2021. Investigation of antioxidant, antibacterial, antiviral, chemical composition, and traditional medicinal properties of the extracts and essential oils of the *Pimpinella* species from a broad perspective: a review. *J. Essent. Oil Res.* 33, 411-426.
- Nikolić, V. *Pimpinella* L. In: Josifović, M. (Ed.). 1973. Flora of SR Serbia (In Serbian), Volume 5. SANU, Belgrade, 229-233.
- Slavkovska, V., Zlatković, B., Marčetić, M., Kovačević, N., Drobac, M. 2022. The anatomy and essential oil composition of the different organs of *Pimpinella tragiunum* and *P. saxifraga* (Apiaceae). *Bot. Serb.* 46, 269-284.
- Tabanca, N., Douglas, A.W., Bedir, E., Dayan, F.E., Kirimer, N., Baser, K.H.C., Aytac, Z., Khan, I.A., Scheffler, B.E., 2005. Patterns of essential oil relationships in *Pimpinella* (Umbelliferae) based on phylogenetic relationships using nuclear and chloroplast sequences. *Plant Genet. Resour.* 3, 149-169.

* Presenting author. E-mail address: milica.drobac@pharmacy.bg.ac.rs (Milica Drobac)

S2.P104 Different extracts of three *Veronica* species: the study of their antioxidant activity

Marija Nazlić^{a,*}, Ivana Vrca^a, Dario Kremer^b, Valerija Dunkić^a

^a Department of Biology, Faculty of Science, University of Split, Ruđera Boškovića 33, 21000 Split, Croatia

^b Department of Ornamental Plants, Landscape Architecture and Garden Art, Faculty of Agriculture, University of Zagreb, Svetošiminska cesta 25, 10000 Zagreb, Croatia

The genus *Veronica* L. is extremely variable in its morphology and its species are very well adapted to different living conditions (Barreira et al., 2014). There are a larger number of studies on various phenolic compounds of *Veronica* species and their antioxidant activities (Barreira et al., 2014; Živković et al., 2017; Sharifi-Rad et al., 2016). The *Veronica* species included in this study were *V. polita* Fr., *V. persica* Poir. and *V. anagallis-aquatica* L. For each selected species, three extracts with phenolic compounds were isolated from aerial parts of the plants: aqueous, methanolic and ethanolic. All extracts were tested for antioxidant activity using two methods: DPPH (2,2-diphenyl- 1-picrylhydrazyl) and ORAC (oxygen radical absorbance capacity). In addition, standards for compounds that were detected in the highest amount in all species were tested for their antioxidant activity: apigenin, *p*-hydroxybenzoic acid, caffeic acid and vanillic acid. Caffeic acid showed the highest antioxidant activity in both methods studied with an IC₅₀ value for DPPH activity of 1.99 µg/mL. Among the plant extracts, methanolic or ethanolic extracts generally showed higher activity than water extracts in both methods, which was to be expected, as organic solutions extract more phenolic compounds. Nevertheless, this was not the case for all species, so this will be discussed through relation of antioxidant activity and the amount of each compound in the phenolic composition. The methanolic extract of *V. polita* showed the highest antioxidant activity in the ORAC method. In the DPPH method, the methanolic extract of *V. anagallis-aquatica* and ethanolic extract of *V. persica* showed the highest activity.

Keywords: *Veronica* species, antioxidant activity, polyphenols, DPPH, ORAC

References:

- Barreira, J.C.M., Dias, M.I., Živković, J., Stojković, D., Soković, M., Santos-Buelga, C., Ferreira, I.C.F.R., 2014. Phenolic profiling of *Veronica* spp. grown in mountain, urban and sandy soil environments. *Food Chem.* 163, 275–283. Živković, J.C., Barreira, J.C.M., Savikin, K.P., Alimpic, A.Z., Stojković, D.S., Dias, M.I., Santos-Buelga, C., Duletić- Laušević, S.N., Ferreira, I.C.F.R., 2017 Chemical profiling and assessment of antineurodegenerative and antioxidant properties of *Veronica teucrium* L. and *Veronica jacquinii* Baumg. *Chem. Biodivers.* 14, e1700167
- Sharifi-Rad, M., Tayeboon, G.S., Sharifi-Rad, J., Iriti, M., Varoni, E.M., Razazi, S., 2016. Inhibitory activity on type 2 diabetes and hypertension key-enzymes, and antioxidant capacity of *Veronica persica* phenolic-rich extracts. *Cell. Mol. Biol.* 62, 80–85.

* Presenting author. E-mail address: mnazlic@pmfst.hr (Marija Nazlić)

S2.P105 Anti-inflammatory potential of chamomile flowers (*Matricariae flos*) extract in topical applications

Natalia Melnyk*, Weronika Skowrońska, Jakub Patryk Piwowarski, Sebastian Granica

Microbiota Lab, Department of Pharmaceutical Biology, Medical University of Warsaw, Banacha 1, Warsaw, Poland

Chamomile flowers (*Matricariae flos*) are one of the oldest and most well-documented plant materials in folk medicine from the Asteraceae family. It is a popular and widely used therapeutic plant traditionally used as an anti-inflammatory and wound-healing topical remedy for skin diseases (EMA, 2015; Srivastava, 2010). Moreover, chamomile flowers are an active ingredient in many popular ointments, creams, and cosmetics (Melnyk, 2022). The objective of the research was to conduct *in vitro* studies of 70% ethanolic chamomile flowers extract's effects on the inflammatory response of skin cells. Immortalized human keratinocytes (HaCat) and primary normal human fibroblasts (NHDF) were chosen as cell lines. The inflammation induced by tumor necrosis factor α (TNF- α)/interferon γ (IFN- γ), and lipoteichoic acid (LTA) accordingly. The secretions of interleukin 6 (IL-6) and 8 (IL-8) were determined using enzyme-linked immunosorbent assay. Additionally, the chemical composition of the extract was characterized by HPLC-DAD-MS. The experimental findings revealed a notable reduction in the release of IL-6 and IL-8 upon exposure to the extract in NHDF cells. Investigations conducted on HaCat cells indicated a significant decrease in the secretion of IL-6 following treatment with the extract. Meanwhile, no statistically significant variances were observed in the IL-8 secretion levels among the experimental groups. The obtained results confirmed the anti-inflammatory potential of chamomile flowers in topical therapy of skin inflammations.

The authors declare no conflict of interest. The presented research was financially supported by the NCN research grant Preludium Bis 2 No. 2020/39/O/NZ7/01109.

Keywords: *Matricariae flos*, Chamomile flowers, anti-inflammatory, skin diseases, interleukins

References

- European Medicines Agency. Assessment report on *Matricaria recutita* L., *flos* and *Matricaria recutita* L., *aetheroleum*. 2015; EMA/HMPC/55837/2011
- Srivastava J.K., Shankar E. and Gupta S., 2010. Chamomile: A herbal medicine of the past with a bright future (review). *Mol. Med. Rep.* 3, 895-901.
- Melnyk N., Vlasova I., Skowrońska W., Bazyłko A., Piwowarski J.P., Granica S., 2022. Current knowledge on interactions of plant materials traditionally used in skin diseases in Poland and Ukraine with human skin microbiota. *Int. J. Mol. Sci.* 23, 9644

* Presenting author. E-mail address: natalia.melnyk@wum.edu.pl (Natalia Melnyk)

S2.P106 A comparative study on biological activities of different extracts of *Scrophularia xanthoglossa* Boiss

Damla Kırıcı^{a,*}, Gökhan Zengin^b, Güneş Ak^b, Ayşe Esra Karadağ^c, Uğur Çakılciöğlu^d

^a İzmir Katip Çelebi University, Faculty of Pharmacy, Department of Pharmacognosy, İzmir, Türkiye

^b Selçuk University, Science Faculty, Physiology and Biochemistry Research Laboratory, Department of Biology, Konya, Türkiye

^c İstanbul Medipol University, Faculty of Pharmacy, Department of Pharmacognosy, İstanbul, Türkiye

^d Munzur University, Pertek Sakine Genç Vocational School, Tunceli, Türkiye

Scrophularia L. genus (Scrophulariaceae) is represented by 76 species in Türkiye, including 37 endemics. Some *Scrophularia* species were used in folk medicine for the treatment of various diseases including skin diseases, eczema and cancer (Lall and Mill, 1978; Özdemir and Alpınar, 2015; Chaibeddra et al., 2020). In the study, the extracts of *S. xanthoglossa* were prepared from ethyl acetate, ethanol, ethanol/water (70%), and water. We focused on the *in-vitro* enzyme inhibition (α -glucosidase, α -amylase, acetylcholinesterase (AChE)/butyrylcholinesterase (BChE/AChE), 5-lipoxygenase (5-LOX) and tyrosinase (TYR) and non-enzyme antioxidant activities (TPC, TFC, DPPH⁺ and ABTS⁺ radical scavenging assays, CUPRAC, FRAP and iron-chelating activity assays) of the extracts. The total contents of phenolics and flavonoids in the extracts varied from 18.68 to 24.22 mg GAE/g and from 2.12 to 15.43 mg RE/g, respectively. The total antioxidant capacity was also determined by the phosphomolybdenum assay (0.77-2.11 mmol TE/g). According to the results of enzyme inhibition activities of AChE (2.48 mg GALAE/g), BChE (9.27 mg GALAE/g), tyrosinase (46.73 $\pm$ 0.31 mg KAE/g) and α -amylase (0.67 mmol ACAE/g) enzyme inhibition activities, ethyl acetate extract was found to be the most effective extract. This study was the first to investigate the biological activities of *S. xanthoglossa* in Türkiye. The extracts obtained with organic solvents showed high biological effects. Furthermore, the results suggest that *S. xanthoglossa* could serve as a potential source of bioactive compounds.

Keywords: *Scrophularia xanthoglossa*, Scrophulariaceae, antioxidant activity, enzyme inhibition

References

- Chaibeddra, Z., Akkal, S., Ouled-Haddar, H., Silva, A.M., Zellagui, A., Sebti, M., Cardoso, S.M. 2020. *Scrophularia tenuipes* Coss. and Durieu: phytochemical composition and biological activities. *Molecules*. 25(7), 1647.
- Lall, S.S. Mill, R.R. 1978. *Scrophularia* L. flora of Turkey and the East Aegean Islands (ed. P.H. Davis) 6, 603- 647. Edinburgh University press., Edinburgh.
- Özdemir, E., Alpınar, K. 2015. An ethnobotanical survey of medicinal plants in western part of central Taurus Mountains: Aladaglar (Nigde-Turkey). *J. Ethnopharmacol.* 166, 53-65.

* Presenting author. E-mail address: damla.kirci@ikc.edu.tr (Damla Kırıcı).

S2.P107 Andrographolide as a potential anti-inflammatory molecule in prostate cancer

Gabriela Erazo^{a,b}, Mitzy Santiago-Rosario^{a,b}, María Sánchez-Vázquez^b, Magaly Martínez-Ferrer^{a,b,*}

^a Department of Pharmaceutical Sciences, University of Puerto Rico School of Pharmacy, San Juan, Puerto Rico

^b Division of Cancer Biology, University of Puerto Rico Comprehensive Cancer Center, San Juan, Puerto Rico

Prostate cancer is the most frequently diagnosed non-cutaneous cancer and the second cause of cancer-related deaths in American men (Siegel et al., 2020). An emerging area of cancer therapeutics is the combination of standard-of-care treatment with natural products (Chhabra et al., 2018). Andrographolide, a labdane diterpenoid and main bioactive component of *Andrographis paniculata*, has shown therapeutic potential. Our laboratory demonstrated that andrographolide inhibits cell growth, tumor growth and angiogenesis in prostate cancer. The objective of this study is to understand the role of andrographolide on inflammation in prostate cancer. Tumors from a mouse xenograft model injected with 22Rv1 cells and treated with andrographolide (10 mg/kg and 25 mg/kg), were used for microarray analysis. Ingenuity Pathway Analysis (IPA) revealed a total of 674 and 218 genes differentially expressed in tumors treated with andrographolide (10 mg/kg and 25 mg/kg), respectively, when compared to their control. Results showed alterations of inflammatory response genes such as ATM, DHFR, HIF-1- α , CDKAL1, NOX1, C5, CCL4, COG6, and TTC3. To further study the inflammatory response of andrographolide in prostate cancer, we used an array to measure changes in cytokine expression in PC3 and 22Rv1 prostate cancer cells. IGF-I, IL-1 β , IL-4, MCP-2, I-309, IL-1 α , IL-5, MDC, PDGF BB, SDF-1, and Angiogenin were differentially expressed in PC3 cells treated with andrographolide (25 mg/kg). Our findings reveal that andrographolide modulates the tumor microenvironment contributing to its possible mechanism of action and anticarcinogenic effect. Understanding these biological pathways is essential to determine pharmacological targets of andrographolide in prostate cancer.

Keywords: prostate cancer, inflammation, microarray analysis, andrographolide, mouse studies

References

- Siegel, D.A., O'Neil, M.E., Richards, T.B., Dowling, N.F., Weir H.K., 2020. Prostate cancer incidence and survival, by stage and race/ethnicity - United States, 2001-2017. *MMWR Morb. Mortal. Wkly. Rep.* 69(41), 1473-1480.
- Chhabra, G., Singh, C.K., Ndiaye, M.A., Fedorowicz, S., Molot, A., Ahmad, N., 2018. Prostate cancer chemoprevention by natural agents: Clinical evidence and potential implications. *Cancer Lett.* 422, 9-18.

* Presenting author. E-mail address: magaly.martinez1@upr.edu (Magaly Martínez-Ferrer)

S2.P108 The impact on proinflammatory cytokines secretion by human neutrophils and keratinocytes by extracts from *Arctii lappae fol*, *Serpylli hba* and *Millefolii hba* – raw materials traditionally used in Poland in skin disorders

Weronika Skowrońska^a, Marta Mainka^a, Monika Ewa Czerwińska^{b,c}, Agnieszka Bazyłko^{a,*}

^a Department of Pharmaceutical Biology, Medical University of Warsaw, 1 Banach Str., 02-97 Warsaw, Poland

^b Department of Biochemistry and Pharmacogenomics, Medical University of Warsaw, 1 Banach Str., 02-097 Warsaw, Poland

^c Centre for Preclinical Research, Medical University of Warsaw, 1B Banach Str., 02-097 Warsaw, Poland

In Poland, a wide range of plant materials are traditionally recognized as medicines for skin disorders (Kaczmarczyk-Sedlak et al., 2020; Kuźnicka et al., 1987). Among them, we have chosen thirteen to the first step of our project. All plant materials were harvested from a Warsaw University of Life Sciences cultivated field. We have determined the effect of aqueous and 70% (v/v) ethanolic extracts on the activity of enzymes (hyaluronidase and lipoxygenase) and their antioxidant activity in cell-free systems (Mainka et al., 2021). Based on the previous results to the second step of our research, six extracts from three plant materials were selected: *Arctii lappae fol*. (All_H₂O, All_EtOH), *Serpylli hba* (Th_H₂O, Th_EtOH) and *Millefolii hba* (K_H₂O, K_EtOH). The impact on ROS secretion by neutrophils stimulated with f-MLP, viability of cells (NRU test) and cytokines secretion by neutrophils and keratinocytes were analyzed. The studies were performed using chemiluminescence, colorimetry and ELISA methods, respectively. In the case of neutrophils stimulated with lipopolysaccharide TNF-α, IL-1β and IL-8 secretion, while on keratinocytes stimulated with UVB, IL-6 and IL-8 secretion were determined. As positive controls, dexamethasone and urolithin B were used. Tested extracts showed no or slight cytotoxicity on used cells. All extracts except All_H₂O inhibited ROS secretion. In neutrophil tests, all extracts showed stimulatory activity on TNF-α and IL-8 secretion, while moderate impact on IL-1β secretion. In tests on keratinocytes all extracts inhibited secretion of IL-6 and IL-8, in the concentration of 100 μg·mL⁻¹ the most active were All_H₂O (65.81±10.95%) and Th_H₂O (63.17±8.17%), respectively.

Keywords: *Achillea millefolium*, *Arctium lappa*, *Thymus serpyllum*, neutrophils, keratinocytes

References

Kaczmarczyk Sedlak, I., Ciolkowski, A. (Eds), 2020. Zioła w medycynie: choroby skóry, włosów i paznokci. Volume 2. PZWL, Warsaw.

Kuźnicka, B., Dziak, M. (Eds), 1987. Zioła i ich zastosowanie. PZWL, Warsaw.

Mainka, M., Czerwińska, M.E., Osińska, E., Ziąja, M., Bazyłko, A., 2021. Screening of antioxidative properties and inhibition of inflammation-linked enzymes by aqueous and ethanolic extracts of plants traditionally used in wound healing in Poland. *Antioxidants*. 10, 698.

* Presenting author. E-mail address: agnieszka.bazylko@wum.edu.pl (Agnieszka Bazyłko)

S2.P109 Enhancing the anticancer activity of oleandrin from *Nerium oleander* L. (Apocynaceae) through novel formulations

Esma Sultan Baykara^a, Çiğdem Bilgi^{b,c,*}

^a Istanbul University-Cerrahpasa, Faculty of Pharmacy, 34500 Istanbul, Türkiye

^b Istanbul University-Cerrahpasa, Faculty of Pharmacy, Department of Pharmacognosy, 34500 Istanbul, Türkiye

^c Health Biotechnology Joint Research and Application Center of Excellence, Istanbul, Türkiye

Nerium oleander Linn., also known as oleander, is a medicinal plant with potent bioactive compounds currently being explored for their potential therapeutic applications. Recent research focuses on its potential in cancer therapy due to the presence of a specific group of secondary metabolites called cardiotonic glycosides. Among these, oleandrin, the cardioactive compound, has emerged as a promising candidate due to its anticancer activity and ability to induce apoptosis in tumor cells. However, its narrow therapeutic window poses a significant challenge for clinical use. Therefore, developing new formulations with improved bioavailability and reduced systemic toxicity is crucial for advancing oleandrin's potential in cancer treatment. This project aims to address the bioavailability limitations of oleandrin and pave the way for its introduction into clinical therapy. In this context, *N. oleander* leaves were collected from Avcılar/Istanbul, dried at 50 °C, and ground into fine powder. Oleandrin was isolated using a modified method adapted from Ryer et al. (1948). Further purifications were performed via chromatographic methods, and the structure of the compound was elucidated by comparing our NMR data with the relevant literature (Karakoyun et al., 2021; Bedir et al., 2021). Oleandrin yield of the process was approximately 0.3% (w/w). Currently, we are developing and evaluating novel formulations designed to enhance the bioavailability and selectivity of oleandrin toward cancer cells, minimizing potential side effects. Developing highly bioavailable and selective formulations would allow for lower dosages and reduced side effects, improving patient outcomes and quality of life.

Acknowledgement: This study was funded by the Scientific Research Projects Coordination Unit of Istanbul University- Cerrahpasa. Project number:37113.

Keywords: oleandrin, cardiac glycoside, anticancer, secondary metabolite, *Nerium oleander*

References

- Ryer, A.I., Marie, F. Process for the Isolation of Oleandrin. U.S. Patent 2438418A, 23 March 1948. Bedir, E., Karakoyun, Ç., Doğan, G., Kuru, G., Küçüksoğak, M., Yusufoglu, H., 2021. New cardenolides from biotransformation of gitoxigenin by the endophytic fungus *Alternaria eureka* 1E1BL1: Characterization and cytotoxic activities. *Molecules*. 26(10), 3030.
- Karakoyun, Ç., Küçüksoğak, M., Bilgi, E., Doğan, G., Çömlekçi, Y.E., Bedir, E., 2021. Five new cardenolides transformed from oleandrin and nerigoside by *Alternaria eureka* 1E1BL1 and *Phaeosphaeria* sp. 1E4CS-1 and their cytotoxic activities. *Phytoch. Lett.* 41, 152-157.

* Presenting author. E-mail address: cigdem.karakoyun@iuc.edu.tr (Çiğdem Bilgi)

S2.P110 Oral acute toxicity and preliminary phytochemical characterization of *Hyptis albida*

Ávila-Villarreal Gabriela^{a,b,*}, Sarabia Dominguez Jazmin Abril^{a,b}, Aguilar Guadarrama Angelica Berenice^c, Irma Martha, Medina-Díaz^d

^a Centro Nayarita de Innovación y Transferencia de Tecnología, A.C. “Unidad Especializada en I+D+i en Calidad de Alimentos y Productos Naturales”, Universidad Autónoma de Nayarit.

^b Unidad Académica de Ciencias Químico Biológicas y Farmacéuticas, Universidad Autónoma de Nayarit

^c Centro de Investigaciones Químicas, IICBA, Universidad Autónoma del Estado de Morelos

^d Laboratorio de Contaminación y Toxicología Ambiental, Universidad Autónoma de Nayarit, Tepic, Mexico

Hyptis albida is a plant commonly used in remedies for the treatment of gastrointestinal disorders, skin infections, rheumatism, cramps, and muscle aches. (Sánchez *et al.* 2013) In the present work, different extraction methods were used with the purpose of carrying out the preliminary phytochemical characterization of *Hyptis albida*. A triterpene was purified: ursolic acid (Fig. 1). As a safety test we evaluate the acute oral toxicity, the LD50 of the hydroalcoholic extract of *Hyptis albida* was established following the OECD 423 guideline, starting with a dose of 300 mg/kg and continuing with a dose of 2000 mg/kg where no deaths or toxic symptoms of the animals were observed for both doses, which is why this extract belongs into category 5 or unclassified according to the Globally Harmonized System.




Fig. 1. Structure of ursolic acid

Keywords: oral toxicity, *Hyptis albida*, ursolic acid, OECD 423, Globally Harmonized System

References

Elizabeth Sánchez, M., Pérez, R., Fresán, O., Zavala, S. Pérez, G., 2013. Anti-inflammatory effects of *hyptis albida* chloroform extract on lipopolysaccharide-stimulated peritoneal macrophages. ISRN Pharmacol. 2013, 713060.

* Presenting author. E-mail address: gabyavila@uan.edu.mx (Ávila-Villarreal Gabriela)

S2.P111 *Ex vivo* assessment of potential *Eucalyptus cinerea* antihypertensive

Daniela Yusbizareth Rodriguez-Jimenez^{a,b,*}, Samuel Enoch Estrada-Soto^c, Luis Arias-Duran^d, Gabriela Maria Avila-Villarreal^{a,b}

^a Unidad académica de Ciencias Químico-Biológicas y Farmacéuticas, Universidad Autónoma de Nayarit. Ciudad de la cultura “Amado Nervo”, 63000 Nayarit, México

^b Unidad Especializada en Calidad de Alimentos y Productos Naturales, Tepic Nayarit, México

^c Facultad de Farmacia, Universidad Autónoma del Estado de Morelos, Cuernavaca, México

^d Laboratorio de farmacognosia Universidad Autónoma Metropolitana de Iztapalapa, México

Eucalyptus cinerea, a member of the Myrtaceae family, is primarily recognized for its ornamental and extensive traditional medicinal (MT) use due to its anti-inflammatory and antioxidant properties, particularly its essential oil and its applications in respiratory diseases such as relaxing activity (Surbhi et al., 2023). Despite its widespread use in MT there remains a lack of evidence regarding its potential as an antihypertensive agent linked to relaxing activity. In this study, our aim was to assess the broader pharmacological potential of leaves of *E. cinerea* hydroalcoholic extracts by evaluating their vasorelaxant activity using a standardized protocol in rat aortic tissue. Ethical considerations, following the principles of the 3Rs, were strictly observed throughout the experiment (Fig. 1).




Fig. 1. Concentration-response curves of the relaxing effect induced by hydroalcoholic (HE-Ec) and precipitate extract (HE-Ec P) on isolated rat aortic, the effect was partially endothelium-dependent.

A), HE-Ec P (Emax 70%, E+; Emax 81% E-) **B)** HE-Ec (Emax 80% E+; Emax 82% E-)

Our results reveal a partially endothelium-dependent vasorelaxant effect, indicating the presence of bioactive compounds in the extract that may interact with various endothelial pathways, including the prostacyclin pathway, endothelial M3-type muscarinic receptor agonists, nitric oxide synthase, or endothelium-derived factors (Sanchez et al., 2020).

Keywords: *Eucalyptus cinerea*, antihypertensive, relaxing effect

References

- Surbhi, Kumar, A., Singh, S., Kumari, P., Rasane, P., 2023. Eucalyptus: phytochemical composition, extraction methods and food and medicinal applications. *Adv. Tradit. Med.* 23(1). Springer.
- Sanchez-Recillas, A., Navarrete-Vazquez, G., Hidalgo-Figueroa, S., Bonilla-Hernandez, M., Ortiz-Andrade, R., Ibarra-Barajas, M., Sanchez-Salgado J.C., 2020. Pharmacological characterization of the cardiovascular effect of Nibethione: *ex vivo*, *in vivo* and *in silico* studies. *J. Pharm. Pharmacol.* 72, 1186-1198.

* Presenting author. E-mail address: Yusbi.jimenez@uan.edu.mx (Daniela Yusbizareth Rodriguez-Jimenez)

S2.P112 The anti-asthmatic effect of artichoke in BALB/c mouse model

Jae Pil Bae*, Venkata Prakash Annamneedi, Jun Woo Park, Tae Jin Kang

College of Pharmacy, Sahmyook University, Seoul 01795, Republic of Korea

Asthma is a chronic inflammatory disease of the respiratory tract, with symptoms such as wheezing, difficulty breathing, and coughing. Artichoke (*Cynara cardunculus* var. *scolymus* L) is a dicotyledonous plant native to the Mediterranean coast. Artichoke has traditionally been widely used for digestive diseases, hyperlipidemia, liver disease, and antioxidant research. However, its efficacy as a therapeutic agent for asthma is not well known. In this study, we investigated anti-asthmatic effect of artichoke using the animal model. Asthma was induced in BALB/C mice with a mixture of ovalbumin and aluminum hydroxide, and then artichoke was oral administered. The number of inflammatory cells was reduced in the bronchoalveolar lavage fluid (BALF) from mice treated with artichoke compared to non-treated mice. The level of serum IgE also reduced in mice treated with artichoke. Using Fluorescence Activated Cell Sorting (FACS) analysis, it was confirmed that the number of immune cells was reduced in BALF isolated from mice treated with artichoke. The level of pro-inflammatory cytokines such as TNF- α , IL-6, and IL-1 β was reduced in BALF from mice treated with artichoke and the level of IL-4, IL-5 and IL-13, a specific cytokine in asthma, was also reduced in BALF from mice treated with artichoke. Our findings suggest that artichoke has anti-asthmatic activity and can be applied to the development of treatments in atopic diseases.

Keywords: asthma, artichoke, BALF, animal model

References

- Agrawal, D.K., Shao, Z. 2010. Pathogenesis of allergic airway inflammation. *Curr. Allergy Asthma Rep.* 10, 39-48. Boo, K.-J., Lee, K., Park, I.-H., Kang, T.J. 2023. *Cynara scolymus* L. Inhibits the LPS-induced Inflammatory Reaction via Suppression of NF- κ B Activity in RAW 264.7 Cells. *Yakhak Hoeji*, 67, 45-50.
- Fritsche, J., Beindorff, C.M., Dachtler, M., Zhang, H., Lammers, J.G. 2002. Isolation, characterization and determination of minor artichoke (*Cynara scolymus* L.) leaf extract compounds. *Eur. Food Res. Technol.* 215, 149-157.

*Presenting authors. E-mail address: qowovlf08@naver.com (Jae Pil Bae)

S2.P113 Antimicrobial activity and chemical composition of *Taraxacum aleppicum* Dahlst. (Asteraceae)

Ebru Kuruldak^{a,b,*}, Fatıma Nur Yılmaz^c, Aynur Sari^d

^a Department of Pharmacognosy, Institute of Graduate Studies in Health Sciences, Istanbul University, Istanbul, Türkiye

^b Department of Pharmacognosy, Faculty of Pharmacy, Istanbul Health and Technology University, Istanbul, Türkiye

^c Department of Pharmaceutical Microbiology, Faculty of Pharmacy, Istanbul University, Istanbul, Türkiye

^d Department of Pharmacognosy, Faculty of Pharmacy, Istanbul University, Istanbul, Türkiye

Taraxacum aleppicum belongs to the genus *Taraxacum* which is a member of the Asteraceae family. The genus *Taraxacum* is found in warmer places of the Northern Hemisphere and it is represented in Türkiye as 54 taxa under nine sections (Uysal et al., 2016). *T. aleppicum* is a member of the Scariosa section which is characterized by a reddish, brown or straw-coloured achene and is used for eye diseases in animals (Gürdal, 2015, Selvi et al., 2023). As a part of our current study, *T. aleppicum* was collected from Sakarya in Marmara region, Türkiye. Air-dried roots and aerial parts were separated and extracted with ethanol. Ethanol extracts of roots and aerial parts were partitioned with petroleum ether, ethyl acetate and *n*-butanol, respectively. The chemical composition of root and aerial part fractions was elucidated using LC-HRMS. The main components of the root EtOAc fraction are chichoric acid, sinapinic acid and chlorogenic acid, respectively. The EtOAc fraction of the aerial part contain mainly chichoric acid, chlorogenic acid and luteolin-7-rutinoside. The main component of BuOH fractions of both root and aerial parts was found to be chichoric acid. The antimicrobial activity of both root and aerial parts fractions was examined against standard strains using the minimum inhibitory concentration (MIC) method (CLSI, 2000 & 2006).

Table 1. Antimicrobial activities of the fractions (µg/mL). PE: Petroleum ether, EtOAc: Ethyl acetate, BuOH:

Butanol, ap. fr.: aerial part fraction, ro. fr.: root fraction, S.a.: *Staphylococcus aureus*, S.e.: *Staphylococcus epidermidis*, K.p.: *Klebsiella pneumoniae*, E.c.: *Escherichia coli*, P.a.: *Pseudomonas aeruginosa*, C.a.: *Candida albicans*, C.t.: *Candida tropicalis*, C.p.: *Candida parapsilosis*, ATTC: American Type Culture Collection

Strains Samples	S.a. ATCC 29213	S.e. ATCC 12228	K.p. ATCC 4352	E.c. ATCC 25922	P.a. ATCC 27853	C.a. ATCC 10231	C.t. ATCC 750	C.p. ATCC 22019
PE ap. fr.	625	72.8	1250	625	625	312.5	625	625
EtOAc ap. fr.	625	156.25	1250	625	625	312.5	625	625
BuOH ap. fr.	625	312.5	625	1250	625	312.5	625	625
PE ro. fr.	625	625	625	1250	625	312.5	312.5	625
EtOAc ro. fr.	1250	625	625	625	625	312.5	625	625
BuOH ro. fr.	625	625	625	1250	625	625	312.5	625

According to the results (Table 1), while root fractions of *T. aleppicum* were found to be moderately effective against all standard strains, fractions of aerial parts were found to be very effective against *Staphylococcus epidermidis* ATCC 12228.

Keywords: *Taraxacum aleppicum*, antimicrobial, LC-HRMS

References

- Clinical and Laboratory Standards Institute (CLSI), 2000., Reference method for broth dilution antifungal susceptibility testing of yeasts; Approved Standard M27-A NCCLS, Wayne, Pennsylvania.
- Clinical and Laboratory Standards Institute (CLSI), 2006., Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: Approved Standard M7-A5. Wayne, PA: CLSI.
- Gürdal, B., 2015. Pharmaceutical botany investigations on taraxacum species growing in marmara region. Doctoral dissertation, Istanbul University.
- Selvi, S., Koç, F.A., Satıl, F. 2023. An ethnoveterinary study on plants used for in the treatment of livestock diseases in Ayvalık (Balıkesir, Turkey). Indian J. Nat. Prod. Resour. (IJNPR) [Formerly Natural Product Radiance (NPR)]. 14(2), 300-312.
- Uysal, A., Zengin, G., Gunes, E., Uysal, N., 2016. Natural two *Taraxacum* species effective against frame shift and base pair substitution mutations: mutagenic, antimutagenic and antioxidant evaluation. Marmara Pharm J. 20(3), 311-319.

* Presenting author. E-mail address: ebrukuruldak@gmail.com (Ebru Kuruldak)

S2.P114 The hydroethanolic extract from *Calendula officinalis* L. herba demonstrated potent anti- inflammatory properties *in vitro*

Johann Röhl^{a,*}, Maria-Riera Piqué-Borràs^a, Mónica Menett-von Eiff^b, Gerald Künstle^a

^a Weleda AG, Preclinical Research and Development, Dychweg 14, 4144 Arlesheim, Switzerland

^b Weleda AG, Research and Development Pharma, Dychweg 14, 4144 Arlesheim, Switzerland

Calendula officinalis L. is traditionally indicated for the symptomatic treatment of minor inflammations of the skin and as an aid in healing of minor wounds that are physiologically accompanied by local inflammation, including activation of the NF- κ B pathway and release of pro- inflammatory mediators such as leukotrienes and prostaglandins. Here we aimed to evaluate the anti-inflammatory efficacy of an hydroethanolic extract from *Calendula officinalis* L. herba. Dry extracts from *Calendula officinalis* L. were prepared using liquid extracts from fresh aerial parts of the flowering Calendula plant (mother tincture *Calendula officinalis* 2a Ø (GHP)). NF- κ B activation was analysed in a reporter assay with the human Jurkat cell line. Analysis of inhibition of the proinflammatory and pain-related enzymes 5-LO and COX-2 as part of the arachidonic acid pathway was performed using recombinant human enzymes. Calendula herba concentration- dependently inhibited NF- κ B activation (IC₅₀: 74.9 μ g/mL). In a cell-free enzyme inhibition assay, Calendula herba concentration-dependently inhibited both, 5-LO and COX-2 enzyme activity (IC₅₀: 39.6 μ g/mL and 29.4 μ g/mL). In this experimental *in vitro* approach the complex mixture of active compounds contained in the Weleda Calendula herba extract demonstrated potent anti- inflammatory and pain-related activity (Fig. 1). Additional studies are performed to further characterise therapeutic benefits of Calendula herba.




Fig.1. Summary of anti-inflammatory properties of the hydroethanolic Calendula herba extract.

Keywords: *Calendula officinalis*, inflammation, NF- κ B, cyclooxygenase-2, 5-lipoxygenase

* Presenting author. E-mail address: preclinical-dev@weleda.ch (Johann Röhl)

S2.P115 A study on the antiproliferative activity of the triterpenoid constituents in *Erica erigena*

Khadijah A. Jabal^{a,b,*}, John J. Walsh^a

^a School of Pharmacy and Pharmaceutical Sciences, Trinity College Dublin, Dublin 2, Ireland

^b Department of Pharmacognosy, Faculty of Pharmacy, Umm Al- Qura University, Makkah, Saudi Arabia

Erica erigena (Ericaceae) grows abundantly in specific regions around the west coast of Ireland, particularly in counties Galway and Mayo (Nelson, 2019). In our studies to date we have found that the aerial part of this plant is a rich source of numerous pentacyclic triterpenes based around the ursane, lupane and oleanane series. An ethyl acetate extract of the powdered leaves and flowers were prepared and loaded onto a flash column. Using a gradient mobile phase system, based around different ratios of hexane: EtOAc, specific bands were collected and analysed by GCMS to identify and quantify the triterpenoids present in the isolated fractions. Using the MTT assay, antiproliferative activity of each fraction was determined using the human prostate cell line, PC3.

Across the entire study, fractions were collected where various ratios of the following terpenoids were identified: α -amyrin, β -amyrin, lupeol, oleanolic acid, micromeric acid and ursolic acid. The minor constituents present included α -amyrenone, β -amyrenone, lupenone, erythrodiol, uvaol, betulin, ursolic aldehyde, β -sitosterol as well as coumaroyl triterpenes. The most active fractions were those that contained different ratios of ursolic:oleanolic:micromeric acids followed by a lupeol rich fraction. Further studies are planned for the evaluation of the active triterpenoid fractions against a panel of tumour cell lines.

Keywords: *Erica erigena*, Ericaceae, GC-MS, antiproliferative, human prostate cell line

Reference

Nelson, E.C., 2019. 905. *Erica erigena*. Curtis's Bot. Mag. 36(1), 32-45.

* Presenting author. E-mail address: Jabalk@tcd.ie (Khadijah A. Jabal)

S2.P116 Cyclic peptides derived from sunflower trypsin inhibitor 1 (SFTI-1) for autoantibodies neutralization in rheumatoid arthritis

Mingshu Zhang^{a,*}, Liyan Mei^{b,c}, Sunithi Gunasekera^a, Helena Idborg^b, Per-Johan Jakobsson^b, Ulf Göransson^a

^a Pharmacognosy group, Department of Pharmaceutical Biosciences, Faculty of Pharmacy, Biomedical Center, Uppsala University, Sweden

^b Rheumatology Unit, Department of Medicine, Karolinska Institutet & Karolinska University Hospital, Sweden

^c The Second Affiliated Hospital, Guangzhou University of Chinese Medicine (Guangdong Provincial Hospital of Chinese Medicine), Guangzhou, China

Bioactive peptides from plants can be used beyond their endogenous functions as defense and hormones. Sunflower trypsin inhibitor I (SFTI-1) is a 14 residues long peptide with a unique bicyclic structure, formed by the cyclic backbone and one disulfide bond. The native peptide is a potent trypsin inhibitor, but numerous examples have shown that this peptide can be modified to display an array of different pharmacological effects. (Luckett et al. 1999, Franke et al. 2018, De Veer et al. 2021) Drawing inspiration from the inherent stability of SFTI-1 found in sunflower seeds, we have employed SFTI-1 as a scaffold to detect and isolate autoantibodies. Previously, we successfully utilized SFTI-1 to display citrullinated peptide epitopes from autoantigens, resulting in grafted peptides with strong binding affinity for anti-citrullinated protein antibodies (ACPA), hallmark of rheumatoid Arthritis (RA). (Gunasekera et al. 2018). To further investigate grafted peptides as potential ACPA scavengers and their potential use in diagnosis and treatment of RA. Five citrullinated peptide antigen epitopes were chosen to replace the trypsin-inhibitory loop of SFTI-1 while preserving its secondary loop. Peptides synthesized via Fmoc-SPPS were then tested against ACPA-positive patient plasma in a ELISA-based ACPA neutralization assay, to evaluate the scavenging potential of grafted peptides. Additionally, the grafted peptides were employed in affinity purification to isolate sub-specific antibodies from ACPA positive individuals. The isolated ACPA subpopulation displayed partial specificity, meanwhile structural and functional studies are still ongoing.

Keywords: scaffold peptide, SFTI-1, rheumatic arthritis

References

- De Veer, S.J., White, A.M., Craik, D.J., 2021. Sunflower trypsin inhibitor-1 (SFTI-1): sowing seeds in the fields of chemistry and biology. *Angew. Chem. Int. Ed.* 60, 8050–8071.
- Franke, B., Mylne, J.S., Rosengren, K.J., 2018. Buried treasure: biosynthesis, structures and applications of cyclic peptides hidden in seed storage albumins. *Nat. Prod. Rep.* 35, 137–146.
- Gunasekera, S., Fernandes-Cerqueira, C., Wennmalm, S., Wähämaa, H., Sommarin, Y., Catrina, A.I., Jakobsson, P.-J., Göransson, U., 2018. Stabilized cyclic peptides as scavengers of autoantibodies: neutralization of anticitrullinated protein/peptide antibodies in rheumatoid arthritis. *ACS Chem. Biol.* 13, 1525–1535.
- Luckett, S., Garcia, R.S., Barker, J.J., Konarev, A.V., Shewry, P.R., Clarke, A.R., Brady, R.L., 1999. High-resolution structure of a potent, cyclic proteinase inhibitor from sunflower seeds. *J. Mol. Biol.* 290, 525–533.

* Presenting author. E-mail address: Mingshu.zhang@uu.se (Mingshu Zhang)

S2.P117 Detailed LC-MS analysis and evaluation of cytotoxic effects of various extracts obtained from aerial parts of *Glaucium acutidentatum* Hausskn. & Bornm

Magdalena Maciejewska-Turska^{a,*}, Elwira Sieniawska^b, Łukasz Świątek^c, Gokhan Zengin^d

^a Department of Pharmacognosy with the Medicinal Plant Garden, Medical University of Lublin, 20-093 Lublin, Poland

^b Department of Natural Products Chemistry, Medical University of Lublin, 20-093 Lublin, Poland

^c Department of Virology With SARS Laboratory, Medical University of Lublin, 20-093 Lublin, Poland

^d Department of Biology, Science Faculty, Selcuk University, Konya 42130, Turkey

Among *Glaucium* spp., *Glaucium acutidentatum* Hausskn. & Bornm. (GA) is one of the less investigated species endemic to Turkey. Only two papers report its significant anticancer activity (Kocanci et al., 2017; Hamamcioglu et al., 2018). The present study aims to evaluate the cytotoxicity of methanolic and aqueous extracts of *G. acutidentatum* obtained using different extraction techniques. Extracts prepared by maceration, homogenizer assisted extraction (HAE) and infusion were qualitatively analysed using LC/ESI-QToF/MS-MS technique to examine in detail their phytochemical profile. The cytotoxic effects of all GA extracts, were evaluated on non- cancerous (VERO) and cancer-derived (AGS - human gastric adenocarcinoma), (FaDu - human hypopharyngeal squamous cell carcinoma) and (RKO - human colon cancer) cell lines using MTT assay. According to the LC-MS results, three types of alkaloids were identified in all samples as the predominant compounds. Second class of polyphenols noted solely in methanolic GA extracts were glycosides of flavonoids. The *in vitro* studies showed that methanolic GA extracts showed lower cytotoxicity towards non-cancerous cells than aqueous extracts. Application of HAE method resulted in higher toxicity than maceration in terms of both extracts. All GA extracts showed selective cytotoxicity towards FaDu cancer cells (selectivity index between 1.62 and 9.04). The maceration-derived aqueous GA extract revealed the lowest CC50 value (24.98 µg/mL) against FaDu cells, which was significantly ($p < 0.0001$) lower than against non-cancerous cells. In conclusion, these results indicate that an appropriately selected extraction method provides GA extract with potential cytotoxic activity against certain types of cancer cells.

Keywords: *Glaucium acutidentatum*, cytotoxic effect, alkaloids

References

- Kocanci, F., Hamamcioglu, B., Aslim, B., 2017. The anti-AChE and anti-proliferative activities of *Glaucium acutidentatum* and *Glaucium corniculatum* alkaloid extracts. J. Appl. Pharm. Sci. 7(8), 191-200.
- Hamamcioglu, B., Kocanci, F., Aslim, B., 2018. Phytochemical screening and evaluation of neuroprotective, anti- mutagenic and anti-genotoxic effects of Turkish endemic *Glaucium acutidentatum*. S. Afr. J. Bot. 117, 232-239.
- Barakat, S., Feras, Q.A., Al Hammouri, M., 2016. Dereplication study on *Glaucium aleppicum* Boiss. in Jordan. Orient. J. Chem. 32(4), 1815-1822.

* Presenting author. E-mail address: magdalena.maciejewska@umlub.pl (Magdalena Maciejewska-Turska)

S2.P118 *In vivo* evaluation of the anti-inflammatory activity of the marine pigment echinochrome A on hairless mice skin

Stefanos Kikionis^{a,*}, Anastasia Panagou^b, Ioannis Svoliantopoulos^b, Andreas Vitsos^b, Efstathia Ioannou^a, Michael Rallis^b, Vassilios Roussis^a

^a Section of Pharmacognosy and Chemistry of Natural Products, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771 Athens, Greece

^b Section of Pharmaceutical Technology, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15784 Athens, Greece

Skin inflammation is the most common symptom observed in almost all dermatological diseases. Exposure to UV irradiation causes acute and chronic effects on skin, such as inflammation, aging and cancer. The UV-induced reaction depends on the intensity of solar radiation and the time of exposure, with acute exposure leading to skin damage, such as edema, erythema and itching. UV radiation generates reactive oxygen species resulting in oxidative stress and increase of inflammatory mediators. Thus, compounds with radical scavenging properties could prevent or attenuate the adverse effects of UV radiation on the skin. Inflamed skin is usually treated by topically applied products, such as creams, gels and lotions, with their main disadvantage being the non-controlled application. Topical patches could be a promising alternative since they can offer controlled release administration. Echinochrome A (7-ethyl-2,3,5,6,8-pentahydroxy-1,4-naphthoquinone, EchA) is the most abundant pigment found in various sea urchin species. In the form of its sodium salts it is the active ingredient of the antioxidant drug HistoChrome[®] used for the treatment of various cardiovascular diseases (Kim et al., 2021). In this study, in order to compare its anti-inflammatory activity in different pharmaceutical formulations, EchA was incorporated either in hydrophilic ointments or in electrospun micro/nanofibrous patches and applied on the UV- inflamed skin of SKH-1 hairless mice. Clinical evaluation, photo-documentation and histopathological analyses were performed and biophysical parameters, eg. erythema, hydration, transepidermal water loss and sebum production, were assessed. Both formulations exhibited significant efficacy but in different doses, ie. 0.1% for the ointment and 1% for the patches.

Keywords: marine natural products, echinochrome A, sea urchins, anti-inflammatory activity, SKH- 1 hairless mice

References

Kim, H.K., Vasileva, E.A., Mishchenko, N.P., Fedoreyev, S.A., Han, J., 2021. Multifaceted clinical effects of echinochrome. *Mar. Drugs* 19, 412.

* Presenting author. E-mail address: skikionis@pharm.uoa.gr (Stefanos Kikionis)

S2.P119 Anti-inflammatory effect of *Glaucium flavum* extracts

Weronika Kozłowska^a, Marcel Białas^{a,†}, Monika Czerwińska^b, Adam Matkowski^c, Sylwia Zielińska^{a,*}

^a Division of Pharmaceutical Biotechnology, Department of Pharmaceutical Biology and Biotechnology, Wrocław Medical University, Borowska 211, 50-556 Wrocław, Poland; ^{a†} Student Scientific Club

^b Department of Biochemistry and Pharmacogenomics, Faculty of Pharmacy, Medical University of Warsaw, Banacha 1, 02-097 Warszawa, Poland

^c Division of Pharmaceutical Biology and Botany, Department of Pharmaceutical Biology and Biotechnology, Wrocław Medical University, Borowska 211A, 50-556 Wrocław, Poland

Glaucium flavum Crantz (yellow hornpoppy, Papaveraceae), distinguished by its vibrant yellow blossoms, is commonly cultivated for ornamental purposes. Yet, its value extends far beyond mere landscape aesthetics. This plant is a rich source of specialized metabolites, offering significant potential in medicine and pharmacy. Consequently, extracts from *G. flavum* obtained from the *in vitro* culture-based biotechnological approach have been explored for their anti-inflammatory and cytotoxic properties. Shoots were cultured on the MH3 medium, supplemented with kinetin (KIN) at 2.5, 4.5, and 7 µM, and indoleacetic acid at 0.5 µM. Additionally, extracts from flowers, roots, and stems of plants cultivated in Kraków and Lublin were also used for comparison. The secretion of IL-1, IL-8, and TNF-α as well as cytotoxic effect were examined using ethanol and methanol extracts [1, 5, 10 µg/mL] on a human neutrophils model. The tested extracts did not show cytotoxicity towards neutrophil cells. The secretion-inhibiting effect was concentration dependent but differed between extracted organs and the particular cytokine. For TNF-α, the secretion was most efficiently inhibited by the extract from stems harvested in Lublin, at a concentration of 1 µg/mL. For IL-1 and IL-8, the highest activity was recorded for 1 µg/mL extracts from *in vitro* shoots cultured on 7 µM KIN MH3 medium and non-supplemented 1/2 MH3 medium, respectively. Additional research is required to elucidate the phytochemical composition of these extracts, thereby pinpointing the specific group of compounds responsible for these observed effects.

Keywords: *Glaucium flavum*, *in vitro* cultures, anti-inflammatory effect

* Presenting author. E-mail address: sylwia.zielinska@umw.edu.pl (Sylwia Zielińska)

S2.P120 Alpha-glucosidase inhibitory profile of sesquiterpenes lactones from *Centaurea* species

Carmen Formisano^a, Elena Serino^a, Daniela Rigano^a, Orazio Tagliatela-Scafati^a, Milena Masullo^b, Gianluigi Lauro^b, Sonia Piacente^b, Giuseppe Bifulco^b

^a Department of Pharmacy, School of Medicine and Surgery, University of Naples Federico II, Via Montesano 49, 80131 Naples, Italy

^b Department of Pharmacy, University of Salerno, Via Giovanni Paolo II, 84084, Salerno, Italy

Centaurea is a genus of flowering plants widely distributed in the Mediterranean region, belonging to Asteraceae family. Many of its species have been used for a long time in traditional medicine as herbal remedies for various purposes, it has been reported that the organic extracts of many *Centaurea* species displayed a high inhibitory activity on α -glucosidase (Fattaheian-Dehkordi et al., 2021), but the bioactive compounds responsible for this activity have not been identified yet. Alpha- glucosidase is a carbohydrate-hydrolase located in the brush border cells of the intestine, which acts on the O-glucosidic α (1 $\rightarrow$ 4) linkage. The enzyme breaks down starch and glucose oligosaccharides into single α -glucose monosaccharides. Its inhibition reduces the digestion rate of carbohydrates, which are not broken down into glucose molecules, reducing glucose levels in the blood. Therefore, the alpha-glucosidase inhibitors represent a class of drugs used to control the glycemic level in non- insulin-dependent diabetes mellitus (type 2), particularly postprandial hyperglycemia. Thus, in order to deepen the potential use of *Centaurea* species to treat metabolic syndrome and identify the bioactive compounds, a screening of various sesquiterpenes lactones isolated from the chloroform extract of different *Centaurea* species aerial parts, including *C. drabifolia*, *C. kotschy* and *C. sicula* has been carried out. As first step, the inhibitory activity of 19 sesquiterpene lactones against α - glucosidase (from *Saccharomyces cerevisiae*) was evaluated by a spectrophotometric assay. The results obtained showed how some compounds exhibited IC₅₀ values in a range from 39.8 to 224.8 μ g/mL. The sesquiterpene melitensin resulted the most active compound with an IC₅₀ of 39.8 μ g/mL, comparable to that observed for acarbose (IC₅₀= 27.9 μ g/mL), the antidiabetic compound used as reference compound. Compounds rhizantholide A, kandavanolide, kotschyol A and 11-epi- 11,13-dihydroartemisiifolin showed IC₅₀ values lower than 50 μ g/mL. Successively, molecular docking experiments were carried out to assess the possibility for selected compounds to form complexes with the α -glucosidase active site. To be consistent with the biological assays performed, they were carried out on the investigated compounds using the 3D protein model of α -glucosidase from *S. cerevisiae*, generated through homology modeling experiments. The analysis of the output poses highlighted the establishment of a network of interactions with residues belonging to the binding site of the enzyme, and specifically with those involved in the catalytic activity, i.e., D214, E276, D349.

Keywords: *Centaurea*, metabolic syndrome, sesquiterpene lactones, α -glucosidase, Asteraceae

References

Fattaheian-Dehkordi, S., Hojjatifard, R., Saeedi, M., Khanavi M., 2021. Review on antidiabetic activity of *Centaurea* spp.: a new approach for developing herbal remedies. Hindawi Evid. Based Complement. Alternat. Med. 2021.

* Presenting author. E-mail address: caformis@unina.it (Carmen Formisano)

S2.P121 Phytochemical profiling and *in vitro* biological activities on *Myrtus communis* L.

Zeliha Parlar^a, Muhammet Cagri Ozkurt^b, Banu Cahide Tel^b, Erdal Bedir^c, I. Irem Tatli Cankaya^a

^a Department of Pharmaceutical Botany, Faculty of Pharmacy, Hacettepe University, 06100, Ankara, Türkiye

^b Department of Pharmacology, Faculty of Pharmacy, Hacettepe University, 06100, Ankara, Türkiye

^c Department of Bioengineering, Faculty of Engineering, Izmir Institute of Technology, 35430, Izmir, Türkiye

Myrtus communis L. (myrtle) is a medicinal plant belonging to the Myrtaceae family. It is an evergreen plant present in most of the Mediterranean countries as wild or cultivated specie. In this study *in vitro* antioxidant and enzyme inhibition tests, as well as cell viability experiments were performed on extracts and fractions. Additionally, the neurorestorative effects were investigated *in vitro* to drive the bioactivity-guided fractionation method (BAGF). Phytochemical analyses were performed on this species. The leaves from wild samples were collected during the flowering period. Using the BAGF method, the antioxidant/antiradical activities of the extracts, and fractions were determined by ABTS, CUPRAC, FRAP, DPPH, and NO tests. Moreover, tyrosinase enzyme inhibition, and cytotoxic activity experiments were carried out. The α -synuclein aggregation model developed in the SH-SY5Y cell line was employed. Phytochemical analyses were performed on 80% EtOH macerate. The structures of the isolated compounds were established by NMR and LC- MS. The phytochemical analysis reported the presence of gallomyrtucommulone C, myricitrin, quercitrin, and ursolic acid, corosolic acid, asiatic acid. All fractions showed antioxidant activity. The infusion was more efficient in reduce the cell viability in the MTT assay. As a result of the tyrosinase enzyme inhibition test, the 80% EtOH macerate was found to be more active than infusion. Further studies on the identified compounds are still ongoing to investigate the pharmacological activities since focusing on the development of natural products.

Keywords: *Myrtus communis*, phytochemistry, BAGF, phenolic compounds, α -synuclein aggregation

References

- Diker, N. Yagmur., 2022. Türkiye'deki Karağaç(*Ulmus* L.) türlerinin farmasötik botanik açısından değerlendirilmesi ve osteoporoz etkilerinin *in silico* yöntemler ile araştırılması.[PhD. Thesis]. Ankara: Ankara University.
- Appendino, G., Maxia, L., Bettoni, P., Locatelli, M., Valdivia, C., Ballero, M., Stavri, M., Gibbons, S., Sterner, O., 2006. antibacterial galloylated alkylphloroglucinol glucosides from myrtle (*Myrtus communis*). J. Nat. Prod. 69, 251- 254.
- Hwang I., Chung S., 2018. Isolation and identification of myricitrin, an antioxidant flavonoid, from daebong persimmon peel. Prev. Nutr. Food Sci. 23(4), 341–346.
- Park J.C., Kim B.W., Young H.S., 1990. Further study on the flavonoids from the leaves of *Machilus thunbergii* in Korea. Kor. J. Pharmacog. 21(3), 197-200.
- Seebacher,W., Simic, N., Weis, R., Saf, R., Kunert, O., 2003. Complete assignments of ¹H and ¹³C NMR resonances of oleanolic acid, 18a-oleanolic acid, ursolic acid and their 11-oxo derivatives. Magn. Reson. Chem. 41, 636–638.
- Lee, et al., 2005. Triterpene acids from the leaves of *Planchonella duclitan* (Blanco) bakhuizen. J. Chin. Chem. Soc. 52, 1275-1280.
- Pérez-Castorena, A., 2014. Triterpenes and other Metabolites from *Tibouchina urvilleana*. J. Mex. Chem. Soc. 58(2), 218-222.

* Presenting author. E-mail address: zelihaparlar@hacettepe.edu.tr (Zeliha Parlar)

S2.P122 Dietary molecules from *Glycyrrhiza foetida* and their modulation on metabolic syndrome pathways

Elena Serino^{a,*}, Hekmat B. Al-Hmadi^{b,c}, Fabio Arturo Iannotti^d, Arianna Pastore^a, Rosa Maria Vitale^d, Giuseppina Chianese^a, Saoussen Hammami^c, Mariano Stornaiuolo^a, Orazio Tagliatela-Scafati^a

^a Department of Pharmacy, School of Medicine and Surgery, University of Naples Federico II, Via D. Montesano 49, 80131 Napoli, Italy

^b Department of Chemistry, College of Medicine, AL-Muthanna University, Samawah, Iraq

^c Laboratory of Environmental Chemistry and Clean Processes (LR21ES04), Faculty of Sciences of Monastir, Monastir University, Monastir, 5000, Tunisia

^d Institute of Biomolecular Chemistry, National Research Council (ICB-CNR), Via Campi Flegrei 34, 80078, Pozzuoli (NA), Italy

Glycyrrhiza is a genus of edible plants, commonly known as licorice, with a long use in folk medicine for their adaptogenic, anti-inflammatory, antioxidant effects, estrogenic activity and hepatoprotective properties. Among the less investigated species of this genus, *G. foetida* seems to be a new promising tool to treat metabolic syndrome (MetS). In fact, this plant has been pointed out as a source of amorfrutins, acetate-deriving polyphenols that explicate a potent insulin-sensitizer effect as PPAR γ agonists that are quite rare in this genus (Weidner et al., 2012). The minor members of this class of compounds have not been in-depth characterized, thus, after a preliminary LC-UV- MS² dereplication step, we have carried out a NMR-based phytochemical screening of *G. foetida* aerial parts that led us to the characterization of 29 compounds, mainly polyphenols, including 11 previously undescribed amorfrutins and an unprecedented flavanone. The amorfrutins were tested on PPAR γ and PPAR α and, besides amorfrutin A, the most active compound on PPAR γ , we have identified the new amorfrutin H as a selective PPAR α agonist and the unprecedented amorfrutin E as a dual PPAR γ/α agonist (Serino et al., 2023). Moreover, all the compounds were tested on other two MetS dysfunctional pathways, highlighting amorfrutin M and decarboxyamorfrutin A as mitochondrial stimulators, while amorfrutin 2 as a glucose uptake promoter. Thus, *G. foetida* is rich source of many bioactive ingredients that, through to the concomitant modulation of different pathways, can ameliorate MetS, emerging as a new potential functional food.

Keywords: *Glycyrrhiza foetida*, metabolic syndrome, amorfrutins, mitochondrial activity, GLUT

References

- Weidner, C., de Groot, J. C., Prasad, A., Freiwald, A., Quedenau, C., Kliem, M., ... & Sauer, S., 2012. Amorfrutins are potent antidiabetic dietary natural products. PNAS. 109(19), 7257-7262.
- Serino, E., Iannotti, F.A., Al-Hmadi, H.B., Caprioglio, D., Moriello, C., Masi, F., ... & Tagliatela-Scafati, O., 2023. PPAR α/γ -targeting amorfrutin phytocannabinoids from aerial parts of *Glycyrrhiza foetida*. J. Nat. Prod. 86(11), 2435- 2447.

* Presenting author. E-mail address: elena.serino@unina.it (Elena Serino)

S2.P123 Smoking hot – the story of the “pepper bark tree” *Warburgia salutaris*

Carmen Leonard^{a,*}, Sandy van Vuuren^b, Alvaro Viljoen^{a,c}

^a Department of Pharmaceutical Sciences, Faculty of Science, Tshwane University of Technology, Private Bag x680, Pretoria 0001, South Africa

^b Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, 7 York Road, Parktown 2193, South Africa

^c SAMRC Herbal Drug Research Unit, Tshwane University of Technology, Private Bag x680, Pretoria 0001, South Africa

In southern Africa, the aromatic, peppery-tasting bark of *Warburgia salutaris* is widely known for its antimicrobial activity especially as treatment for upper respiratory tract infections. Historically, smoke inhalation or smoking of the bark or root alone (or with another plant such as *Cannabis sativa*) has been used in traditional medicine for the treatment of upper respiratory tract infections (Watt and Breyer-Brandwijk, 1962; Hutchings et al., 1996; van Wyk et al., 1997). The bioactive compounds are concentrated in the bark and therefore this plant-part is most often used for infusions, decoctions, and to a lesser extent for smoke therapy in traditional medicine. Recently, in South Africa, a study among the Vhavenda people in Limpopo province, reported the use of *W. salutaris* bark as smoke therapy (11%) (Ramarumo et al., 2019). Furthermore in a recent study, the drimane sesquiterpenes in the smoke fraction of *Warburgia salutaris* bark was reported for the first time (Leonard, 2021). In this study, a smoke recovery system was used to capture these compounds. The sesquiterpenes were isolated using preparative high-performance liquid chromatography (HPLC) and using nuclear magnetic resonance (NMR) were identified as cinnamolide-3 β -acetate, ugandensidial, bemadienolide, and cinnamolide. All these sesquiterpenes have been shown to have anti-infective properties and thus this could offer another therapeutic route of administration as well as assisting in conserving the plant as less material is needed for this therapeutic route of administration.

Keywords: Traditional medicine, Sesquiterpenes, Smoke therapy, *Warburgia salutaris*

References

- Hutchings, A., Scott, A., 1996. Zulu Medicinal Plants. Pietermaritzburg: University of Natal Press.
- Leonard, C.M., 2021. Phytochemical variation, antimicrobial activity and quality control of *W. salutaris* (Pepper bark tree). Doctor of Technologiae. Tshwane University of Technology.
- Ramarumo, L.J., Maroyi, A., Tshisikhawe, M.P., 2019. *Warburgia salutaris* (G. Bertol.) Chiov.: An endangered therapeutic plant used by the vhavenda ethnic group in the soutpansberg, vhembe biosphere reserve, limpopo province, South Africa. Res. J. Pharm. Tech. 12(12), 5893-5898.
- Van Wyk, B.-E., Oudtshoorn, B.V. & Gericke, N., 1997. Medicinal Plants of South Africa. Pretoria: Briza.
- Watt, J., Breyer-Brandwijk, M., 1962. The medicinal and poisonous plants of Southern and Eastern Africa. Edinburgh: Livingstone.

* Presenting author. E-mail address: leonardcm@tut.ac.za (Carmen Leonard)

S2.P124 The effect of cannabidiol, cannabidiol and cannabigerol on the motoric activity of the rat proximal colon – *ex vivo* study

Kijan Crowley^{a,*}, Łukasz Kiraga^a, Edyta Mischuk^a, Magdalena Chłopecka^a, Rafał Rzewuski^b, Joanna Banach^c

^a Division of Pharmacology and Toxicology, Department of Preclinical Sciences, Institute of Veterinary Medicine, Warsaw University of Life Sciences, 8 Ciszewskiego St., 02-786 Warsaw, Poland

^b Medicann LLC, 24/26a Piękna St., 00-549, Warsaw, Poland

^c Department for Research and Processing of Seed, Institute of Natural Fibres and Medicinal Plants, National Research Institution, 71b Wojska Polskiego St., 60-630 Poznań, Poland

The study was aimed at evaluating the effect of cannabidiol (CBD), cannabidiol (CBN) and cannabigerol (CBG) on the motoric activity of the gastrointestinal tract. The experiments were carried out on rat proximal colon preparations incubated in modified Krebs-Henseleit solution (37°C, pH 7.35-7.45) under isometric conditions in an organ bath set under a load of 0.01 N. The strips were treated with solutions of cannabinoids (in the range of 0.08 – 800 µg/mL) and subsequently with acetylcholine (ACh, 1 µM), without flushing. The impact of cannabinoids on contractility of the colon preparations was registered. The spontaneous activity and ACh-reactivity of colon muscles in the presence of solvent (0.5% DMSO) was the control.




Fig. 1. Example recording of a rat proximal colon strip's reactivity to acetylcholine (ACh) in the presence of DMSO (0.5%) and cannabidiol solutions (CBD). F – flushing

Incubation of proximal colon preparations in the presence of cannabinoid solutions caused significant myorelaxation. The obtained results indicate that all cannabinoids dose-dependently affected the spontaneous activity and reactivity of the colon strips, and that the effect differed depending on the type of cannabinoid. The reaction to ACh when cannabinoid solutions were introduced preceding ACh application was reduced. It was demonstrated that the reduction of contractions was strongest in the case of CBG solutions and amounts to $22.1 \pm 21.6\%$ and $7.9 \pm 9.1\%$ when the preparations were treated with 80 and 800 µM of cannabinoid, respectively. The incubation of tissue slices in CBN solutions caused the least changes; in the case of reactivity to ACh, the significant reaction was observed when colon preparations was incubated in the maximum concentration and it amounts to $40.9 \pm 31.4\%$ of control reaction to ACh. The obtained results are a valuable contribution to the area of research on the use of cannabinoids in supportive therapy of gastrointestinal diseases.

Keywords: cannabinoids, rat, colon, *ex vivo*

*Presenting author. E-mail address: kijan_crowley@sggw.edu.pl (Kijan Crowley)

S2.P125 Coumarin polypharmacology and elucidation of an underlying molecular mechanism of action

Sandhya Krishnan^a, Felix Huwyler^{a,*}, Lukasz Kurach^{a,b}, Krystyna Skalicka-Wozniak^b, Jürg Gertsch^a

^a Institute of Biochemistry and Molecular Medicine, University of Bern, Switzerland

^b Medical University of Lublin, Poland

Coumarins, widely present in herbal drugs, spices, and food, are recognized for their diverse pharmacological properties, including anti-inflammatory, anticoagulant, anticancer activities, and effects on central nervous system disorders (Skalicka-Woźniak et al.). *In vivo*, they exert unexpected biphasic pharmacological effects (Kowalczyk et al., Skiba et al.). Our investigation utilizing Nuclear Magnetic Resonance (NMR) reveals the thiophilic nature of the unsubstituted 3,4-double bond in coumarins. Through the creation of clickable coumarin probes (CLICOPs), we demonstrate for the first time that various coumarins form covalent bonds with specific proteins in living cells, murine tissue homogenates, and human blood. Employing a range of different CLICOPs enabled competition with natural coumarins, facilitating the identification and visualization of previously unknown covalent coumarin-protein interactions on SDS-PAGE using chemoproteomics. Notably, attempts to label selected proteins identified in chemoproteomics using purified proteins did not yield covalent reactions. However, the addition of macromolecular crowders, such as Ficoll-400, promoted the thiophilic reaction *in vitro*. Our findings indicate that the presence of artificial or cellular macromolecular crowders is essential for coumarins to react. Further insights revealed that coumarins can elicit no, allosteric, or orthosteric pharmacological effects on specific proteins via this covalent mechanism, underscoring their hitherto underestimated polypharmacological potential and raising questions about their toxicity. Additionally, our study challenges conventional *in vitro* approaches by demonstrating that natural coumarins may not react in certain conditions but covalently interact in a complex cellular environment. This complexity underscores the polypharmacology of coumarins and carries significant toxicopharmacological implications, providing a novel molecular mechanism underlying their diverse effects.

Keywords: Coumarins, mechanism of action, macromolecular crowding, polypharmacology

References

- Skalicka-Woźniak, K., Orhan, I.E., Cordell, G.A., Nabavi, S.M., Budzyńska, B., 2016. Implication of coumarins towards central nervous system disorders. *Pharmacol. Res.* 103, 188-203.
- Kowalczyk, J., Budzyńska, B., Kurach, Ł., Pellegata, D., El Sayed, N.S., Gertsch, J., Skalicka-Woźniak, K., 2022. Neuropsychopharmacological profiling of scoparone in mice. *Sci. Rep.* 12(1), 822.
- Skiba, A., Pellegata, D., Morozova, V., Koziol, E., Budzyńska, B., Lee, S.M., Gertsch, J., Skalicka-Woźniak, K., 2023. Pharmacometabolic effects of pteryxin and valproate on pentylenetetrazole-induced seizures in zebrafish reveal vagus nerve stimulation. *Cells.* 12(11), 1540.

* Presenting author. E-mail address: felix.huwyler@unibe.ch (Felix Huwyler)

S2.P126 Deciphering the targets of trypanocidal chalcones in *Trypanosoma cruzi*

Kaushavi Anil Cholke^{a,*}, Olivier Kirchhoffer^b, Emerson Ferreira Queiroz^b, Jürg Gertsch^a

^a Institute of Biochemistry and Molecular Medicine, University of Bern, Switzerland

^b School of Pharmaceutical Sciences, University of Geneva, Switzerland

The protozoan parasite *Trypanosoma cruzi* (*T. cruzi*) is the causative agent of Chagas disease (CD) which affects millions of primarily indigenous people in Latin America (Bern et al., 2011; Salm and Gertsch, 2019). Infection with *T. cruzi* is usually life-long, and up to 30% of individuals develop chronic Chagas disease, with symptoms that include cardiomyopathy and/or digestive mega syndromes. Treatment of *T. cruzi* infection with the nitroheterocyclic drugs benznidazole and nifurtimox, introduced in the 1970s, is suboptimal. This highlights the urgent need for improved drugs. Based on an extensive screening of 775 extracts of botanical drugs used in Bolivia in the context of CD and botanical drugs from unrelated indications from the Mediterranean De Materia Medica (Salm et al. 2021), we investigate the natural product class chalcones (α , β -unsaturated ketones) (Fig. 1B) which are often present in plants together with flavonoids and showed low micromolar antichagasic effects.




Fig. 1 Chemical structure of the chalcone scaffold

It was recently shown that chalcones mediate covalent interactions with specific enzymes in *Leishmania* spp. (Escrivani et al., 2021). To identify the molecular mechanism of action of trypanocidal chalcones in *T. cruzi*, we employed a chemoproteomics strategy by generating clickable chalcone probes. Through chemical modification, we successfully improved the selective toxicity of chalcones towards *T. cruzi* and established a strategy to identify their targets. Intriguingly, the peroxidase enzymes targeted by chalcones in *T. cruzi* also appeared to have a role in benznidazole and nifurtimox resistance. By employing targeted and untargeted LC-MS/MS based metabolomics we aim to decipher the role of these enzymes in *T. cruzi* biology. Overall, chalcones are an interesting new trypanocidal class of natural products and clickable probes may help to study the biology of peroxidases in this parasite.

Keywords: *T. cruzi*, chalcone, chemoproteomics, antichagasic effects

References

- Bern, C., Kjos, S., Yabsley, M.J. and Montgomery, S.P., 2011. T *Trypanosoma cruzi* and Chagas' disease in the United States. Clin. Microbiol. Rev. 24(4), 655-681.
- Salm, A. and Gertsch, J., 2019. Cultural perception of triatomine bugs and Chagas disease in Bolivia: a cross-sectional field study. Parasit. Vectors. 12(1), 1-19.
- Salm, A., Krishnan, S.R., Collu, M., Danton, O., Hamburger, M., Leonti, M., Almanza, G. and Gertsch, J., 2021. Phylobioactive hotspots in plant resources used to treat Chagas disease. Iscience. 24(4).
- Escrivani, D.O., Charlton, R.L., Caruso, M.B., Burle-Caldas, G.A., Borsodi, M.P.G., Zingali, R.B., Arruda-Costa, N., Palmeira-Mello, M.V., de Jesus, J.B., Souza, A.M. and Abraham-Vieira, B., 2021. Chalcones identify cTXNPx as a potential antileishmanial drug target. PLoS Negl. Trop. Dis. 15(11), p. e0009951.

* Presenting author. E-mail address: kaushavi.cholke@unibe.ch (Kaushavi Anil Cholke)

S2.P127 Innovative methods for analysing flavonolignan complex from *Silybum marianum* L.

Jovana Gjorgjievska^a, Ivana Cvetkovikj^a, Olga Gigopulu^a, Marja Karapandzova^a, Nikola Geshkovski^a, Petre Makreski^b, Gjoshе Stefkov^{a,*}

^a Faculty of Pharmacy, Ss. Cyril and Methodius University in Skopje, Majka Tereza 47, 1000 Skopje, North Macedonia

^b Institute of Chemistry, Faculty of Natural Sciences and Mathematics-Skopje, Ss. Cyril and Methodius University in Skopje, Arhimedova 5, 1000 Skopje, North Macedonia

Silymarin is a mixture of flavonolignans that is mostly present in the fruits of the milk thistle plant (*Silybum marianum* (L.) Gaertn., Asteraceae). Research on *S. marianum* has shown that it has antiatherosclerotic, antidiabetic, hepatoprotective, antihypertensive properties. At least 1.5% w/w of silymarin in dried seeds is required for the European Pharmacopoeia test, which entails a drawn-out and complicated sample preparation process and comprehensive HPLC analysis. As a result, the goal of this study was to ascertain the amount of silymarin contained in milk thistle fruits and extracts from both wild and cultivated plants, in accordance with the Ph. Eur. 11.0 monograph, as well as by utilizing quick sample preparation and quantification techniques to accelerate the analysis. The silymarin concentration of milk thistle fruit samples, which were collected over several years, ranged within a range of 0.3% to 1.8% w/w, according to the acquired HPLC data. The Ph. Eur. sample preparation was substituted with a very effective one-hour subsequent ultrasonic extraction. The fruits and extracts were also examined using spectroscopic methods. A partial least-squares (PLS) model for quantification was constructed utilizing the obtained spectra, with HPLC as a reference technique. The chemometric analysis based on the data obtained from the FTIR and Raman spectra points out satisfactory statistical indicators for employment of these methods for quantitative determination of silymarin.

Keywords: Milk thistle, extraction, spectroscopic methods, chemometric analysis

* Presenting author. E-mail address: gstefkov@yahoo.com (Gjoshе Stefkov)

S2.P128 Phytochemical analysis of *Hyssopus officinalis* L. cultivated in western Turkey

Gulhan Zeybek, Gulen Irem Kaya*

Ege University Faculty of Pharmacy, Department of Pharmacognosy Izmir, Turkey

H. officinalis L. belonging to the family Lamiaceae, possesses antioxidant, antimicrobial and several other pharmacological activities. The essential oil of hyssop is widely used in food, pharmaceutical and cosmetic industries throughout the world (Tahir et al., 2018). In the present work we performed phytochemical analysis on the methanolic extract (yield 9.55 %) and essential oil (yield 0.53%) of *H. officinalis* which was cultivated in western Turkey. A reversed-phase HPLC- DAD method has been used and validated for the determination of rutin and apigenin in the methanolic extract from *H. officinalis*. The content of rutin and apigenin were determined as 3552.913 ± 249.324 µg/g and 12.383 ± 1.102 µg/g extract, respectively. In previous studies, apigenin was detected as the major component (Fathiazad et al., 2011), while rutin was not detected or was detected in very low amounts in *Hyssopus* extracts (Tahir, 2018; Proestos, 2005). The chemical constituents of essential oil obtained by steam distillation from the aerial parts of *H. officinalis* were analyzed by GC-MS. It was determined that hyssop essential oil contained isopinocampone (25.0%), pinocampone (15.6%) and β-pinene (8.5%) as major components. According to the previous studies, so far three types of *Hyssopus* oils have been encountered, namely; monoterpene ketone-type, 1,8-cineole type and methyeugenol-type. In the present study, the essential oil found to be rich in monoterpene ketones and among these, the content of isopinocampone was higher than the content of the same compound found in the oils previously reported from Turkey (Kürkçüoğlu et al., 2016), except in the oil of *Hyssopus* cultivated in southeastern Turkey, with 57.27% (Kızıl, 2010).

Funding: This research was funded by Ege University Research Project YLT-2023-26315 HPLC and GC analysis was carried by EGE MATAL

Keywords: *Hyssopus officinalis*, essential oil, rutin, apigenin

References

- Fathiazad F., Mazandarani M. Hamedeyazdan S., 2011. Phytochemical analysis and antioxidant activity of *Hyssopus officinalis* L. from Iran, Adv. Pharm. Bull. 1(2), 63-67.
- Kızıl, S., Haşimi, N., Tolan, V., Kilinc, E., Karataş, H. 2010. Chemical composition, antimicrobial and antioxidant activities of hyssop (*Hyssopus officinalis* L.) essential oil. Not. Bot. Horti. Agrobo. 38(3), 99-103.
- Kürkçüoğlu M. Arıkan Eser Ş., Başer K.H.C., 2016. Composition of the Essent. Oils essential oil of the *Hyssopus officinalis* L. subsp. *angustifolius* (Bieb.) Arcangeli, Nat. Volatiles & Essent. Oils 3(2). 15-19.
- Proestos, C., Chorianopoulos, N., Nychas, G.J.E., Komaitis, M. 2005. RP-HPLC analysis of the phenolic compounds of plant extracts. Investigation of their antioxidant capacity and antimicrobial activity. J. Agr. Food Chem. 53(4), 1190– 1195.
- Tahir, M., Khushtar, M., Fahad, M., & Rahman, M. A. 2018. Phytochemistry and pharmacological profile of traditionally used medicinal plant Hyssop (*Hyssopus officinalis* L.). J. Appl. Pharm.Sci. 8(7), 132-140.

* Presenting author. E-mail address: gulen.irem.kaya@ege.edu.tr (Gulen Irem Kaya)

S2.P129 Enhancing wound healing: sericin and *Chelidonium majus* L. as potential dressings

Ana Borges^{a,b}, María Luisa Martín Calvo^c, Sandrina A. Heleno^{a,b}, Sónia P. Miguel^d, Maximiano P. Ribeiro^d, Paula Coutinho^d, André F. Moreira^d, Josiana A. Vaz^{a,b}, Lillian Barros^{a,b}, Ricardo C. Calhelha^{a,b}

^a Centro de Investigação da Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

^b Laboratório Associado para a Sustentabilidade e Tecnologia em Regiões de Montanha (SusTEC), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

^c Grupo de Investigación en Fisiología y Farmacología, Facultad de Farmacia, Universidad de Salamanca, Campus Miguel de Unamuno s/n, 37007 Salamanca, Espanha

^d CPIRN-IPG—Center of Potential and Innovation of Natural Resources, Polytechnic Institute of Guarda, 6300-559 Guarda, Portugal

Wound healing, a complex physiological process orchestrating intricate cellular and molecular events, seeks to restore tissue integrity. The burgeoning interest in leveraging the therapeutic potential of natural substances for advanced wound dressings is a recent phenomenon. Notably, Sericin, a silk-derived protein, and *Chelidonium majus* L. (*C. majus*), a botanical agent, have emerged as compelling candidates, providing a unique combination of natural elements that may revolutionize conventional wound care approaches. Sericin, renowned for its diverse properties, displays unique properties that accelerate the wound healing process. Simultaneously, *C. majus*, with its diverse pharmacological compounds, shows promise in reducing inflammation and promoting tissue regeneration. As the demand for innovative wound care solutions increases, understanding the therapeutic potential of natural products becomes imperative. This review synthesizes current knowledge on Sericin and *C. majus*, envisioning their future roles in advancing wound management strategies. The exploration of these natural substances as constituents of wound dressings provides a promising avenue for the development of sustainable, effective, and biocompatible materials that could significantly impact the field of wound healing.

Acknowledgements: FCT, Portugal for financial support to CIMO (UIDB/00690/2020 and UIDP/00690/2020) and SusTEC (LA/P/0007/2021); national funding by FCT, P.I., through the institutional scientific employment program- contract for S.A. Heleno and RC Calhelha, and through the project Wastesilk (<http://doi.org/10.54499/PTDC/BTA-BTA/0696/2020>).

Keywords: tissue repair, biomaterial, medical device, silk, greater celandine

* Presenting author. E-mail address: calhelha@ipb.pt (Ricardo C. Calhelha)

S2.P130 Evaluating the efficacy of topical application of *Genista tridentata* (L.) (*Fabaceae*) leaf extract on wound healing in experimental diabetes

Inês Martins Laranjeira^{a,b,c,d,*}, Beatriz Gonçalves^d, Diana Rodrigues-Fonseca^{a,b}, Alberto C. P. Dias^d, Filipa Pinto-Ribeiro^{a,b}

^a Life and Health Sciences Research Institute (ICVS), School of Medicine, Campus of Gualtar, University of Minho, 4710-057 Braga, Portugal

^b ICVS/3B's - PT Government Associate Laboratory, Braga/Guimarães, Portugal

^c CITAB - Centre for the Research and Technology of Agro-Environmental and Biological Sciences, University of Trás-os-montes e Alto Douro, Portugal

^d Centre of Molecular and Environmental Biology (CBMA), University of Minho, Campus de Gualtar, Braga, 4710-057, Portugal

Diabetic wounds are a serious complication of diabetes along with peripheral neuropathy, macrovascular and microvascular disease, impaired angiogenesis, and chronic inflammation (Pawar et al., 2021). In this work methanolic leaf extracts from *Genista tridentata* (L.), known for their antioxidant and anti-inflammatory properties (Laranjeira et al., 2023), were investigated *in vitro* and *in vivo* for their antioxidant activity, fibroblasts cytotoxicity, and their potential in promoting wound healing in experimental diabetes. The *G. tridentata* extract exhibited strong antioxidant activity both in *in vitro* and in the TBHP test while the MTT test (Oliveira et al., 2018) in fibroblasts showed no toxicity. Using a STZ-NA-induced diabetes model (Diab) (Masiello et al., 1998) in association with an excision wound model (Nasir et al., 2016), 21 rats were divided into three groups (n=7 each; daily treatment for 10 days): (i) Diab+Gt, topically treatment with *G. tridentata* incorporated base-cream; (ii) Diab+Veh, topically treatment with a base-cream; and (iii) Diab, no topical treatment. Skin histopathological analysis demonstrated improvements of several wound healing parameters, namely in wound contraction, decreased granulocytic tissue area, and increased neo-epidermis thickness after treatment with the base-cream with *Genista tridentata* extract. Our data shows the topical application of a methanolic extract from *Genista tridentata* leaves improves wound healing in diabetic rats, highlighting its potential use in the pharmaceutical industry.

Keywords: *Genista tridentata*, *Diabetes mellitus*, medicinal Plants, wound-healing

References

- Laranjeira, I.M., Dias, A.C.P., Pinto-Ribeiro, F.L., 2023. *Genista tridentata* phytochemical characterization and biological activities: a systematic review. *Biology*. 12(11), 1387.
- Masiello, P., Broca, C., Gross, R., Roye, M., Manteghetti, M., Hillaire-Buys, D., Novelli, M. Ribes, G., 1998. Experimental NIDDM: development of a new model in adult rats administered streptozotocin and nicotinamide. *Diabetes*. 47(2), 224-229.
- Nasir, M.A., Mahammed, N.L., Roshan, S., Ahmed, M.W., 2016. Wound healing activity of poly herbal formulation in albino rats using excision wound model, incision wound model, dead space wound model and burn wound model. *Int. J. Res. Dev. Pharm. L. Sci.* 5(2), 2080-2087.
- Oliveira, A.I., Pinho, C., Fonte, P., Sarmiento, B., Dias, A.C., 2018. Development, characterization, antioxidant and hepatoprotective properties of poly (ε-caprolactone) nanoparticles loaded with a neuroprotective fraction of *Hypericum perforatum*. *Int J. Biol. Macromol.* 110, 185-196.
- Pawar, K.B., Desai, S., Bhonde, R.R., Bhole, R.P., Deshmukh, A.A., 2021. Wound with diabetes: present scenario and future. *Curr. Diabetes Rev.* 17(2), 136-142.

* Presenting author. E-mail address: inesmartinslaranjeira@gmail.com (Inês Martins Laranjeira)

S2.P131 *Lepidium meyenii* – isolation of imidazole alkaloids by centrifugal partition chromatography

Dominik Tarabasz*, Wirginia Kukula-Koch

Department of Pharmacognosy with Medicinal Plants Garden, Medical University of Lublin, 1 Chodzki Str., 20-093 Lublin, Poland

Lepidium meyenii (maca) is one of the plants used by Peruvians for over two thousand years as an edible plant (Da Silva Leitão Peres et al., 2020). Most studies indicate its positive effect on fertility, but it has also been shown to have neuroprotective, dermatological, antimicrobial and anticancer effects (Ulloa del Carpio et al., 2024). Lepidilines are a unique and characteristic group of compounds from the 'Peruvian ginseng'. These four imidazole alkaloids that have been isolated by the authors are currently available only as synthesized compounds, which are very expensive. Thus the aim of this study was to use centrifugal partition chromatography to purify all four imidazole alkaloids (Fig. 1) from the total water-methanol extract of *Lepidium meyenii* tubers and, if possible, to separate them from one other. This liquid-liquid separation technique offers low solvent consumption, high selectivity, and repeatability, but also up-scaling potential, which makes the elaborated purification protocols valuable in terms of their potential industrial applications. Due to a common parent structure shared by all lepidilines and only slight differences in the R₁ and R₂ substituents, the selection of a biphasic liquid system (BLS) was a challenge. To achieve the best results in the preparation of promising partition coefficients, we lowered the pH of the BLS, and after establishing appropriate separation conditions, i.e., elution mode, rotation speed, and flow of the mobile phase, we managed to separate lepidilines from the total extract.




Fig. 1. Parent structure of imidazole alkaloids (R₁: –H; –CH₃, R₂: –H; –OCH₃)

Keywords: *Lepidium meyenii*, brassicaceae, lepidilines, isolation, centrifugal partition chromatography

References

- Da Silva Leitão Peres, N., Cabrera Parra Bortoluzzi, L., Medeiros Marques, L.L., Formigoni, M., Fuchs, R.H.B., Droval, A.A., Reitz Cardoso, F.A., 2020. Medicinal effects of Peruvian maca (*Lepidium meyenii*): a review. *Food Funct.* 11(1), 83–92.
- Ulloa del Carpio, N., Alvarado-Corella, D., Quiñones-Laveriano, D.M., Araya-Sibaja, A., Vega-Baudrit, J., Monagas- Juan, M., Navarro-Hoyos, M., Villar-López, M., 2024. Exploring the chemical and pharmacological variability of *Lepidium meyenii*: a comprehensive review of the effects of maca. *Front. Pharmacol.* 15.

* Presenting author. E-mail address: dominikanes@o2.pl (Dominik Tarabasz)

S2.P132 Connecting targets of the standardized herbal combination STW 5-II to gastrointestinal disorders

Gudrun S. Ulrich-Merzenich^{a,*}, Anastasiia Shcherbakova^a, Sabine Rabini^b, Olaf Kelber^b, Ramy M. Ammar^b

^a *University Hospital Bonn (UKB), Medical Clinic III, AG Synergy Research, Bonn Germany*

^b *Bayer Consumer Health, R&D Phytomedicine Development Centre, Darmstadt, Germany*

STW5-II, a standardized combination of six plants, is indicated in Germany for the treatment of Functional Dyspepsia (FD) and the Irritable Bowel Syndrome (IBS). Both are complex and heterogeneous disorders with multiple pathomechanisms. We predicted molecular targets of STW5-II in FD and IBS by constructing compound-target networks between compounds supposed to contribute to the biological activity of STW5-II and relevant targets for FD and IBS. These targets were identified by searching the GEO database for comparative gene expression (GE)-profiles from colon biopsies of IBS or FD patients and healthy controls. The predicted targets were validated by wet lab experiments with NCM460 colon cells treated with STW5-II and its components. Deep sequencing and RT-PCR of these samples confirmed the predicted regulations of AKT1, NOS3, HSP90AA, TRPV and SMAD2 by STW 5-II (Ulrich-Merzenich et al. 2021, Shcherbakova et al. 2022). In a second step gene ontology and reactome pathways were analyzed and indicated significant upregulations of cell motility (GO:2000145), epithelium development (GO:0060429) and lipid metabolism (R-HSA-556833) by STW 5-II, whereas ATP metabolic processes (GO:004634) were downregulated. We extended the comparative GE-analysis of NCM460 colon cells treated with STW5-II to colon biopsies of IBS patients during a 7-day red meat diet. 37 genes were differentially regulated. E.g. Tropomyosin 1, which is considered a novel marker for cancer of the gastrointestinal tract, was decreased. We can provide further evidence that STW5-II may not only addresses disease-specific but also food-related pathomechanisms.

Keywords: plant combination, synergy, compound-target network, IBS, FD

References

- Ulrich-Merzenich, G., Shcherbakova, A., Rabini, S., Thumann, T.A., Pferschy-Wenzig, E.M., Bauer, R. Ammar, R.M., 2021. Elucidation of multi-target mechanisms of STW5-II on functional dyspepsia. *Planta Med.* 87(15), 1281.
- Shcherbakova, A., Kelber, O., Ammar, R.M., Ulrich-Merzenich, G., 2022. STW5-II addresses both- colon targets of the irritable bowel syndrome with predominant constipation and with predominant diarrhea *in vitro*. *Planta Med.* 88(15), 1548.

* Presenting author. E-mail address: gudrun.ulrich-merzenich@ukbonn.de (Gudrun S. Ulrich-Merzenich)

S2.P133 The anti-inflammatory and skin barrier function recovery effects of *Schisandra chinensis* in mice with atopic dermatitis

Yoorae Son^a, Wonjin Yang^a, Sangjun Park^a, Jinkyu Yang^a, Soyeon Kim^a, Ji-Hyo Lyu^b, Hyungwoo Kim^{a,*}

^a Division of Pharmacology, School of Korean Medicine, Pusan National University, Yangsan, 50612, Gyeongnam, Republic of Korea

^b Herbal Medicine Resources Research Center, Korea Institute of Oriental Medicine, Naju, 58245, Jeonnam, Republic of Korea

The fruit of *Schisandra chinensis* (Turcz.) Baill. is widely used medicinally to treat coughs, asthma, exhaustion, eczema, and pruritus in Northeast Asian countries, including Korea, China, and Japan (Kim et al., 1997). This study was designed to investigate the effects of *S. chinensis* on dermatitis and skin barrier dysfunction in mice with calcipotriol (MC903) induced atopic dermatitis (AD) (Moosbrugger-Martinz et al., 2017). In this experiment, the inhibitory effects of an ethanolic extract of *S. chinensis* (EESC) on skin lesions, water content, water holding capacity (WHC), histopathological abnormalities, and inflammatory cytokine and chemokine levels were evaluated in mice with AD induced by MC903. In our results, topical EESC ameliorated skin lesions, reduced skin water content and MC903-induced WHC increases. EESC also prevented MC903-induced histopathological abnormalities such as epidermal disruption, hyperkeratosis, spongiotic changes, and immune cell infiltration in inflamed tissues. Also, topical EESC reduced MC903-induced levels of pro-inflammatory cytokines and chemokines, such as tumour necrosis factor (TNF)- α , interleukin (IL)-1 β , IL-4, IL-6; IL-8, monocyte chemotactic protein (MCP)-1, and thymic stromal lymphopoietin (TSLP). Furthermore, unlike dexamethasone, EESC did not reduce spleen/body weight ratios. In conclusion, these results suggest that *S. chinensis* can be used as an alternative to external corticosteroids and that its anti-inflammatory and skin barrier dysfunction-restoring effects are related to downregulations of pro-inflammatory cytokines and chemokines, such as TNF- α , IL-4, IL-6, IL-8, and TSLP.

Keywords: *Schisandra chinensis*, herbal medicine, inflammation, atopic dermatitis, eczema

References

- Kim, C.M., Shin, M.K., An, D.G., Lee, K.S. 1997. The encyclopedia of oriental herbal medicine. Seoul. Jeongdam. 3058-3066.
- Moosbrugger-Martinz, V., Schmuth, M., Dubrac, S. 2017. A Mouse Model for Atopic Dermatitis Using Topical Application of Vitamin D3 or of Its Analog MC903. Methods Protoc. 1559, 91–106.

* Presenting author. E-mail address: kronos7@pusan.ac.kr (Hyungwoo Kim)

S2.P134 Comparison of amount of tannins in Ericaceae plants using different water extraction methods

Kristina Ložienė*, Evelina Petraitytė, Jurgita Paliunytė

Pharmacy and Pharmacology Center, Institute of Biomedical Sciences, Vilnius University, Geležinio Vilko Str. 29a, Vilnius, Lithuania

Tannins constitute a rich group of phenolic compounds in plants and are soluble in water. However, there is no single tannins extraction with water method suitable for all tannins accumulating plants. Therefore, to extract higher content of tannins from different raw materials, it is necessary to choose a suitable extraction with water method for each species separately (Chuo et. Al., 2022).

The aim was to compare content of tannins and phenolic compounds in *Vaccinium vitis-idaea*, *Vaccinium myrtillus*, *Arctostaphylos uva-ursi*, *Calluna vulgaris*, *Empetrum nigrum*, *Andromeda polifolia*, and *Rhododendron tomentosum* leaves, using hot water extraction, ultrasonic extraction as well as maceration with water. Folin-Ciocalteu method was used to determine tannins and phenolics expressed as tannic acid equivalents (TAE). Tannin content was calculated by subtracting of phenolics remaining after binding of tannins to polyvinyl polypyrrolidone from the total phenolics content.

Hot water was the most effective for extraction of phenolics. For example, hot water extracted four times more phenolic compounds from *A. uva-ursi* than maceration (35.30 ± 0.96 and 8.07 ± 0.46 % (TAE), respectively). Meanwhile, there was no single tannins extraction method suitable for all studied species: extraction with hot water was the most efficiently for *R. tomentosum*, *E. nigrum*, *A. polifolia* and *V. myrtillus* (10.80 ± 1.97 , 3.94 ± 0.04 , 6.81 ± 0.58 , and 5.63 ± 0.54 % (TAE), respectively), ultrasonic extraction – for *A. uva-ursi* and *V. vitis-idaea* (6.54 ± 1.72 and 4.83 ± 0.72 % (TAE), respectively), and maceration – for *C. vulgaris* (3.02 ± 0.36 % (TAE)).

Keywords: Ericaceae, tannins, water, extraction methods

References

Chuo, S.C., Nasir, H.M., Mohd-Setapar, S.H., Mohamed, S.F., Ahmad, A., Wani, W.A., Muddassir, M., Alarifi, A., 2022. A glimpse into the extraction methods of active compounds from plants. Crit. Rev. Anal. Chem. 52, 667-696.

* Presenting author. E-mail address: kristina.loziene@mf.vu.lt, kristina.loziene@gmail.com (Kristina Ložienė)

S2.P135 Biosynthetic relationship and self-assembly nanoparticles of royleanone isolated from *Plectranthus* species

Epole Ntungwe^{a,*}, Chioma M. Ekejiuba^b

^a University of Coimbra, CQC-IMS, Department of Chemistry, 3004-535 Coimbra, Portugal

^b CBIOS – Universidade Lusófona's Research Center for Biosciences & Health Technologies, Universidade Lusófona de Humanidades e Tecnologias, Campo Grande 376, 1749-024 Lisbon, Portugal

The *Plectranthus* genus belongs to the *Lamiaceae* family and consists of around 300 species distributed in Africa, Asia, and Australia. *Plectranthus* spp. has been reported to be rich in diterpenoids, such as different types of royleanones **1** (Fig. 1). These royleanones are known for their different biological activities including antimicrobial, antioxidant, general toxicity, and cytotoxicity activities (Ntungwe et al., 2013). Therefore, this work aimed to isolate large amounts of cytotoxic royleanones from acetone extracts of *plectranthus* species through an ultrasound-assisted extraction method. These royleanones were tested for their cytotoxicity and employed as a starting material for the synthesis of self-assembled squalene and oleic acid nanoparticles. The obtained compound was characterized by spectroscopic methods, mainly NMR techniques. Additionally, the characterization of the nanoparticles (size, zeta potential, and polydispersity using DLS and SEM), the ability to release the drug unit, and the preliminary evaluation of their cytotoxicity in different cell lines were also tested. Due to the similarity in the chemical structure of these royleanones were set to study their biosynthetic relationship using Fukui indexes. This work established that there is a biosynthetic relationship between the isolated abietane diterpenoids and that the synthesized nanoparticles could act as a strategy to vehiculate the drug.




Fig. 1. Royleanone **1** structure

Acknowledgments: This work has run within the research project PTDC/QUI-QFI/1880/2020. The CQC-IMS is supported by the Fundação para a Ciência e a Tecnologia (FCT), through projects UI0313B/QUI/2020 (DOI: 10.54499/UIDB/00313/2020), UI0313P/QUI/2020 (DOI: 10.54499/UIDP/00313/2020) and LA/P/0056/2020.

Keywords: royleanones, self-assembly nanoparticles, biosynthetic relationship

References

Ntungwe, E., Domínguez-Martín, E.M., Bangay, G., Garcia, C., Guerreiro, I., Colombo, E., Saraiva, L., Díaz-Lanza, A.M., Rosatella, A., Alves, M.M., Reis, C.P., Passarella, D., Rijo P, 2021. Self-assembly nanoparticles of natural bioactive abietane diterpenes. *Int. J. Mol. Sci.* 22(19), 10210.

* Presenting author. E-mail address: epole.ntungwe@uc.pt (Epole Ntungwe)

S2.P136 Yamogenin – a steroidal plant metabolite triggers extrinsic and intrinsic pathway of apoptosis in gastric cancer cells

Justyna Stefanowicz-Hajduk*, Anna Hering, Magdalena Gucwa, Piotr Graczyk

Department of Biology and Pharmaceutical Botany, Medical University of Gdansk, Al. Hallera 107, 80-416 Gdansk, Poland

Steroidal saponins are high-molecular weight compounds widely distributed in plants, among which *Agavaceae*, *Liliaceae*, and *Dioscoreaceae* are the main source of these steroids. The compounds have many pharmacological and biological properties, e.g. antimicrobial, anti-inflammatory, antitumor, fungicidal, and insecticidal activity (Hostettmann and Marston, 1995; Qin et al., 2012; Stefanowicz-Hajduk et al., 2022). Yamogenin is one of the spirostane saponins, a stereoisomer of diosgenin and occurs in *Trigonella foenum-graecum*, *Asparagus officinalis*, and *Dioscorea collettii*. In this study, we evaluated cytotoxic activity of yamogenin *in vitro* on gastric cancer AGS cells and determined the role of selected cellular factors in cell death. The viability of the cells was estimated with MTT assay. The cell cycle arrest, mitochondrial membrane depolarization, the level of oxidative stress, and the activity of initiator and executioner caspase-8 and -9 was estimated with flow cytometry and luminometry, respectively. Genes expression analysis at the mRNA level was conducted with Real-Time PCR. The obtained results indicate that yamogenin has cytotoxic activity in AGS cells with an IC_{50} value of 18.50 ± 1.24 μ g/mL and strongly inhibits the cell cycle in the sub-G1 phase. The compound also significantly decreases mitochondrial potential, increases reactive oxygen species (ROS) production, and strongly activates caspase-8 and -9, what indicates that the receptor and mitochondrial way of apoptosis are involved in cell death.

Keywords: cytotoxicity, saponins, receptor and mitochondrial pathway of apoptosis, flow cytometry

References

- Hostettmann, K., Marston, A., 1995. Saponins. Cambridge University Press, New York, NY, USA.
- Qin, X.J., Sun, D.J., Ni, W., Chen, C.X., Hua, Y., He, L., Liu, H.Y., 2012. Steroidal saponins with antimicrobial activity from stems and leaves of *Paris polyphylla* var. *yunnanensis*. *Steroids*. 77, 1242–1248.
- Stefanowicz-Hajduk, J., Hering, A., Gucwa, M., Czerwińska, M., Ochocka, J.R., 2022. Yamogenin-induced cell cycle arrest, oxidative stress, and apoptosis in human ovarian cancer cell line. *Molecules*. 27, 8181.

* Presenting author. E-mail address: justynastef@gumed.edu.pl (Justyna Stefanowicz-Hajduk)

S2.P137 Phytochemical analysis and neuroprotective activity of extracts from *Paliurus spina-christi* (Rhamnaceae)

Melis Kasap^a, Dervis Birim^b, Guliz Armagan^b, Nehir Unver Somer^{a,*}

^a Ege University, Faculty of Pharmacy, Department of Pharmacognosy, 35040 Bornova Izmir

^b Ege University, Faculty of Pharmacy, Department of Biochemistry, 35040 Bornova Izmir

Paliurus spina-christi Mill., the only species of the genus *Paliurus* Mill. growing in Turkey (Davis, 1967), is used traditionally for various medical purposes such as a diuretic, and against kidney stones and diabetes (Sargin et al., 2013). The plant has recently attracted attention due to its phenolic content and promising biological activities (Takım, 2021). In the present study, the total phenolic, total flavonoid and rutin content, a common major compound found in the plant, were determined in the leaf and fruit extracts. In addition, cell viability was evaluated using an MTS/PMS assay in a cellular model of oxidative stress following the exposure to extracts at various concentrations (1-100 µg/mL). Methanolic extracts were prepared using an ultrasonic bath (3 times) and evaporated to dryness. In the leaf extract (yield 25.55%), total phenols and total flavonoids were determined as 102.974 ± 4.511 mg GAE/g and 68.677 ± 3.016 mg QE/g, whereas in the fruit extract (yield 10.57%), total phenolic and flavonoid contents were found to be 62.051 ± 3.307 mg GAE/g and 46.872 ± 1.124 mg QE/g, respectively. A reversed phase HPLC-DAD method was employed and validated for the quantitative determination of rutin. The amount of rutin was detected as 2961.167 ± 88.355 µg/g extract in the leaves (extract yield 21.94%) and 3593.833 ± 247.718 µg/g extract in the fruits (extract yield 9.26%). Neuroprotection by the extracts in the presence of hydrogen peroxide was highlighted by a significant reduction of cellular loss at concentration less than 10 µg/mL ($p < 0.05$).

Keywords: *Paliurus spina-christi*, rutin, HPLC-DAD, neuroprotective activity

Funding: This work was funded by Ege University Research Project YLT-2022-24122 and HPLC analysis was performed by EGE- MATA.

References

- Davis P.H., 1967. *Paliurus* (Rhamnaceae). In: Davis P.H. (Ed.). Flora of Turkey and the East Aegean Islands, Volume 2. Edinburgh University Press, Edinburgh, 523-4.
- Sargin, S.A., Akcicek, E., Selvi, S., 2013. An ethnobotanical study of medicinal plants used by the local people of Alaşehir (Manisa) in Turkey. J. Ethnopharmacol. 139, 626-41.
- Takım, K., 2021. Bioactive component analysis and investigation of antidiabetic effect of Jerusalem thorn (*Paliurus spina-christi*) fruits in diabetic rats induced by streptozotocin. J. Ethnopharmacol. 264, 113263.

* Presenting author. E-mail address: nehir.somer@ege.edu.tr; nehirunver@hotmail.com (Nehir Unver Somer)

S2.P138 Bioactive molecules separation from *Acmella oleracea* (Jambu) from Amazon Florest

Elza Caroline Alves Müller^{a,*}, Clarissa Silva Lima^a, Katia Castanho Scortecci^b, Fabio Boylan^c

^a Department of Biological Sciences and Health of the Federal University of Amapa, 68903-197, Macapa, AP, Brazil

^b Cell Biology and Genetics Dept, Federal University of Rio Grande do Norte, 59078-970, Natal, RN, Brazil

^c School of Pharmacy, Trinity College, University of Dublin. Ireland

Acmella oleracea (L). R. K. Jansen (Asteraceae), popularly known as Jambu, is consumed in aqueous preparations as food in Latin America, more precisely in the north of Brazil. This research focused on the chemical compounds released into the water that can justify its anaesthetic or anti-proliferative medicinal action on tumour cell lines such as spilanthol **1** (Fig. 1), its main constituent (Pereira, 2023; Williams, 2013).




Fig. 1. Structure of spilanthol 1

The initial purpose of this study was to devise the best solvent system to perform the separation of the ethyl acetate and butanol fractions extracted from an infusion prepared with Jambu aerial parts for subsequent use in HSCCC and HPLC. Aerial parts of *Acmella oleracea* were collected at Fazendinha district (S 0°02'30.40"/W 51°06'37.5"), Macapá, Brazil. An infusion was prepared and extracted with ethyl acetate and butanol. After testing four different mobile phases with a combination of ethyl-acetate, acetic acid, formic acid, butanol and water at different proportions, the (butanol 3: acetic acid 1: water 4) was chosen to analyse the compounds in both extracts. Similarly, we could choose the best solvent system separating the extracts on HSCCC. For both cases, after testing four possibilities, the best results were achieved with ethyl acetate: butanol: water 4,5:0,5:5. One run using the reverse phase mode allowed for separating different compounds, and spilanthol was obtained in the final test tubes. It was then purified using Sephadex LH-20. We are currently trying to purify the minor compounds present in the sample, which are believed to be flavonoids.

Keywords: *Acmella oleracea*, HSCCC, spilanthol, flavonoids

Acknowledgements: FAPEAP, UNIFAP, Trinity College Dublin.

References

Williams C. Medicinal Plants in Australia Volume 4: An Antipodean Apothecary. Rosenberg Publishing, 2013. Pereira, F. <https://www.cozinhatradicional.com/tacaca-para/>. Sep, 15. 2023.

* Presenting author. E-mail address: carolinemuller8@hotmail.com (Elza Caroline Alves Müller)

S2.P139 The protective impact of *Aronia melanocarpa* L. berries extract against cadmium-induced changes in the concentrations of calciotropic hormones in the serum

Małgorzata M. Brzóska^a, Joanna Rogalska^a, Małgorzata Gałążyn-Sidorczuk^a, Michał Tomczyk^{b,*}

^a Department of Toxicology, Faculty of Pharmacy with the Division of Laboratory Medicine, Medical University of Białystok, ul. Mickiewicza 2C, 15-222 Białystok, Poland

^b Department of Pharmacognosy, Faculty of Pharmacy with the Division of Laboratory Medicine, Medical University of Białystok, ul. Mickiewicza 2A, 15-230 Białystok, Poland

The results of our studies, conducted in an experimental rat model of current environmental human exposure to cadmium, allowed us to hypothesize that *Aronia melanocarpa* L. (Michx.) Elliott berries (chokeberry) products may be an effective strategy in protection against health outcomes caused by this toxic metal, including skeleton damage (Brzóska et al., 2015; Mężyńska et al., 2019; Ruczaj et al., 2024; Smereczński et al., 2023). However, the protective use of chokeberry products in people exposed to cadmium requires further research. The present study aimed to evaluate, in our experimental model (1 and 5 mg Cd/kg of diet, 3–24 months), the impact of low-level and moderate intoxication with cadmium, on the concentrations of calcium and calciotropic hormones (1,25- dihydroxyvitamin D – 1,25(OH)₂D, parathormone – PTH, and calcitonin – CT) in the serum and whether the supplementation with a 0.1% extract from *A. melanocarpa* berries can influence these parameters. Calcium was determined by atomic absorption spectrometry, whereas calciotropic hormones were assayed with the use of commercial kits. The exposure to cadmium, dose and duration dependently, decreased the serum concentrations of 1,25(OH)₂D and CT and increased PTH concentration. Calcium concentration was unchanged except for its decrease after the 3-month exposure to 5 mg Cd/kg of diet. The co-administration of chokeberry extract prevented all cadmium-induced changes in the concentrations of calciotropic hormones. In conclusion, the supplementation with *A. melanocarpa* berries extract during exposure to cadmium allows for the maintenance of proper levels of calciotropic hormones in the serum but it does not influence the concentration of calcium.

Keywords: *Aronia melanocarpa* berries, cadmium, calcium, calciotropic hormones, protection

References

- Brzóska, M.M., Rogalska, J., Gałążyn-Sidorczuk, M., Jurczuk, M., Roszczenko, A., Tomczyk, M. 2015. Protective effect of *Aronia melanocarpa* polyphenols against cadmium-induced disorders in bone metabolism: a study in a rat model of lifetime human exposure to this heavy metal. *Chem. Biol. Interact.* 229, 132-146.
- Mężyńska, M., Brzóska, M.M., Rogalska, J., Piłat-Marcinkiewicz, B. 2019. Extract from *Aronia melanocarpa* L. berries prevents cadmium-induced oxidative stress in the liver: a study in a rat model of low-level and moderate lifetime human exposure to this toxic metal. *Nutrients* 11, 21.
- Ruczaj, A., Brzóska, M.M., Rogalska, J. 2024. The protective impact of *Aronia melanocarpa* L. berries extract against prooxidative cadmium action in the brain – a study in an in vivo model of current environmental human exposure to this harmful element. *Nutrients* 16, 502.
- Smereczński, N.M., Brzóska, M.M., Rogalska, J., Hutsch T. 2023. The protective potential of *Aronia melanocarpa* L. berry extract against cadmium-induced kidney damage: a study in an animal model of human environmental exposure to this toxic element. *Int. J. Mol. Sci.* 24, 11647.

* Presenting author. E-mail address: michal.tomczyk@umb.edu.pl (Michał Tomczyk)

S2.P140 LC-MS/MS analyses of triterpenes and polyphenols from less-polar fractions of *Rosa rugosa* leaf extract and their biological activity

Marta Olech*, Agnieszka Łubek-Nguyen, Wioleta Dołomisiewicz, Lena Łyko

Department of Pharmaceutical Botany, Medical University of Lublin, 1 Chodźki Str., 20-093 Lublin, Poland

Plants from the rose family (Rosaceae) have high edible and medicinal value and a long history of therapeutical application. Our previous research showed that rose leaves contain considerable amounts of free and bound phenolic acids and flavonoid aglycones (Olech et al., 2020a). In this research, we investigated less-polar rose components. Therefore, air-dried crushed leaves of rugosa rose (*Rosa rugosa* Thunb.) were exhaustively extracted with portions of 80% and 95% ethanol to obtain ethanolic extract (RREt). RREt was then fractionated using liquid-liquid extraction, which yielded several fractions with different polarity levels. Among them, water insoluble (non-polar) and chloroform fractions were obtained with a relatively high yield (13.3 and 1.7%, respectively).

The content of pentacyclic triterpenes was determined with the newly developed method using liquid chromatography-atmospheric pressure chemical ionization-tandem mass spectrometry (LC-APCI-MS/MS). It revealed the presence of pomolic, fupenzic, maslinic, corosolic, betulinic, ursolic, and oleanolic acids as well as uvaol, lupeol, β -amyrin, and α -amyrin. The highest contents of these compounds were found in the water insoluble fraction. The identification and quantification of phenolic acids and flavonoids was carried out using an improved LC-ESI-MS/MS-MRM method (Olech et al., 2020b). The fractions were found to contain twelve phenolic acids (including three caffeoylquinic acids), eleven flavonoid glycosides, and seven flavonoid aglycones. Moreover, enzyme inhibitory and antioxidant activities were determined in the samples in *in vitro* assays. The oxygen radical absorbance capacity varied from 445.12 to 644.63 μ g Trolox/mg. The ability to inhibit α -glucosidase, α -amylase, and pro-inflammatory enzymes (lipoxygenase, hyaluronidase) increased with the sample polarity.

Keywords: rosaceae, enzyme inhibitory activity, anti-inflammatory activity, tandem mass spectrometry

References

- Olech, M., Pietrzak, W., Nowak, R., 2020a. Characterization of free and bound phenolic acids and flavonoid aglycones in *Rosa rugosa* Thunb. leaves and achenes using LC-ESI-MS/MS-MRM methods. *Molecules*. 25, 1804.
- Olech, M., Nowak, R., Ivanova, D., Tashev, A., Boyadzhieva, S., Kalotova, G., Angelov, G., Gawlik-Dziki, U., 2020b. LC-ESI-MS/MS-MRM profiling of polyphenols and antioxidant activity evaluation of junipers of different origin. *Appl. Sci.* 10, 8921.

* Presenting author. E-mail address: marta.olech@umlub.pl (Marta Olech)

S2.P141 Volatiles, triterpenes, phytosterols and anti-inflammatory activity of *Rhododendron luteum* Sweet (Ericaceae) supercritical CO₂ extracts

Lena Łyko^{a,*}, Agnieszka Krajewska^b, Marta Olech^a, Urszula Gawlik^c, Danuta Kalembe^b, Anna Och^a, Renata Nowak^a

^a Department of Pharmaceutical Botany, Medical University of Lublin, 1 Chdźki Street, 20-093 Lublin, Poland

^b Department of Biochemistry and Food Chemistry, University of Life Sciences, Skromna Street 8, 20-704 Lublin, Poland;

^c Institute of General Food Chemistry, Lodz University of Technology, Lodz, Poland

Terpenes are a diverse group of plant secondary metabolites, including monoterpenes, diterpenes, sesquiterpenes, triterpenes, and saponins. This group is also diverse in terms of biological activity. They have been proven to have anti-inflammatory, anti-diabetic, anti-proliferative, and antiviral activities (Castellano et al. 2013; Vázquez et al. 2011; Xiao et al. 2018). The present study aimed to determine the terpene profile of supercritical CO₂ extracts of *Rhododendron luteum* leaf (SCE-RLL) and analyze their anti-inflammatory and antioxidant activity. Extracts were subjected to LC-APCI- MS/MS analysis, revealing the occurrence of some pentacyclic terpenic acids and phytosterols. Using the HS-SPME-GC-FID-MS method allowed the detection of more than 100 volatile compounds in one of the extracts, mainly limonene, eugenol, β-phenylethanol and β-caryophyllene. Next, all samples were subjected to biological studies using ELISA spectroscopic methods. The performed analyses have shown extracts ability to inhibit enzymes involved in inflammation: lipoxygenase, hyaluronidase, and xanthine oxidase. Moreover, they presented high antioxidant potential in the ORAC test. This is the first study of SCE-RLL composition and biological activity. Based on the results, *R. luteum* Sweet can be a valuable source of pentacyclic terpenes, phytosterols, and other phytochemicals with anti-inflammatory and antioxidant activity.

Keywords: *Rhododendron luteum* Sweet, LC-MS, GC-MS, enzyme inhibition

References

- Castellano, J.M., Guinda, A., Delgado, T., Rada, M., Cayuela, J.A., 2013. Biochemical basis of the antidiabetic activity of oleanolic acid and related pentacyclic triterpenes. *Diabetes*. 62(6), 1791–1799.
- Vázquez, L., Palazon, J. Navarro A., 2011. A review of sources and biological activities. *Intechopen*. 426, 487–502.
- Xiao, S., Tian, Z., Wang, Y., Si, L., Zhang, L., Zhou, D., 2018. Recent progress in the antiviral activity and mechanism study of pentacyclic triterpenoids and their derivatives. *Med. Res. Rev.* 38(3), 951–76.

* Presenting author: E-mail address: lenalyko18@gmail.com (Lena Łyko)

S2.P142 Antioxidant and anti-inflammatory activity of the *Berberis vulgaris* L. bark extract

Anna Och, Marta Olech, Lena Łyko*, Renata Nowak

Department of Pharmaceutical Botany, Medical University of Lublin, 1 Chodźki Street, 20-093 Lublin, Poland

Berberis vulgaris L. (Berberidaceae) is a medicinal plant with over 300 years of tradition in folk medicine. *Barberry vulgaris* L. is valuable in Chinese folk medicine, where it is called “Sankezhen”, in Ayurvedic medicine, and has also been used in traditional medicine in Europe in recent centuries (Och et al., 2021). Despite its well-established traditional use, scientific research reveals little about the biological activity of barberry. Modern research has confirmed the anti-inflammatory effect of barberry, but mainly in the field of fruit extracts (Li et al. 2016; Majeed et al. 2016). *Berberis vulgaris* L. is able to accumulate alkaloids in the bark and its pharmacological effect is probably related to its alkaloid composition. However, the broader chemical composition of its bark is still unexplored. The aim of this study was to evaluate the chemical profile of methanol extract from barberry bark and to analyze its antioxidant and anti-inflammatory effects. The extracts obtained with the ASE (accelerated solvent extraction) method were subjected to LC-MS/MS analysis, which showed the presence, among other compounds, of high-level Eleutheroside E – well-known for its anti-inflammatory activity (Och et al. 2023). The *in vitro* biological activity assays demonstrated the ability of the extract to inhibit the enzyme lipoxygenase involved in the inflammatory process and showed high antioxidant potential in the DPPH, ABTS, and ORAC tests. The results suggest that the bark of *Berberis vulgaris* L. can be a valuable source of phytochemicals with anti-inflammatory and antioxidant activity.

Keywords: *Berberis vulgaris* L, antioxidant activity, anti-inflammatory activity

References

- Li, Y., Wang, P., Chai, MJ., 2016. Effects of berberine on serum inflammatory factors and carotid atherosclerotic plaques in patients with acute cerebral ischemic stroke. *Zhongguo Zhongyao Zazhi*. 41, 4066–4071.
- Majeed, W., Aslam, B., Javed, I., Khaliq, T., Muhammad, F., Ali, A., Raza, A. 2015. Histopathological evaluation of gastro protective effect of *Berberis vulgaris* (Zereshk) seeds against aspirin induced ulcer in albino mice. *Pak. J. Pharm. Sci.* 28(6),1953-8.
- Och, A., Nowak, R., 2021. “Barberry (*Berberis vulgaris*)—Traditional and Contemporary Use.” Springer: Cham, Switzerland pp. 797–825.
- Och, A., Olech, M., Bąk, K., Kanak, S., Cwener, A., Cieśla, M., Nowak, R. 2023. Evaluation of the antioxidant and anti-lipoxygenase activity of *Berberis vulgaris* L. leaves, fruits, and stem and their LC-MS/MS polyphenolic profile. *Antioxidants*. 12, 1467.

* Presenting author. E-mail address: lenalyko18@gmail.com (Lena Łyko)

S2.P143 The use of herbal medicines for gynecological ailments: patient-reported outcomes

Alexandra Drebka^{a,*}, Annika J. Scholl^a, Teresa Ochs^a, Olaf Kelber^{b,c}, Ralph Mösges^{d,e}, Esther Raskopf^e, Kija Shah-Hosseini^d, Beatrice E. Bachmeier^a

^a Institute of Pharmaceutical Biology, Goethe University Frankfurt, Max-von-Laue-Str. 9, 60438 Frankfurt am Main, Germany

^b R&D, Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt, Germany

^c Kooperation Phytopharmaka GbR, Bonn, Germany

^d Institute of Medical Statistics and Computational Biology, University of Cologne, Cologne, Germany

^e ClinCompetence Cologne GmbH, Cologne, Germany

Gynecological ailments have a negative impact on quality of life and productivity. Standard treatment is associated with poor tolerability and other issues related to public health. Herbal Medicines (HMs) are used traditionally for the treatment of gynecological complaints for centuries and constitute an excellent addition to current treatment options. HMs are well tolerated and therapeutically efficacious as evidenced by various clinical studies. Aim of this study was to expand this knowledge with real world evidence from patient reported outcomes. We analyzed a data sample taken from the pharmaco-epidemiological database PhytoVIS regarding therapeutic benefit and side effects of HMs used for the treatment of menstrual and menopausal ailments as well as uncomplicated urinary tract infections. We found that more than 80% of the patients in the sample assessed the overall therapeutic benefit of HMs for the treatment of gynecological complaints as very positive. Over 90% of the patients using HMs perceived no or no significant side effects. Treatment habits differed depending on the type of complaint. In this context, the majority of women with menstrual or menopausal ailments preferred to treat for time period of 1 month or longer, while those affected by uncomplicated urinary tract infections reduced the application of HMs to the length of their symptoms. Interestingly women with even strong symptoms relied on the therapeutic benefit of HMs. Real-world outcome data are an important supplement to clinical data. Our results reveal the excellent benefit-risk ratio of HMs and help to implement them into novel therapeutic strategies.

Keywords: gynecological ailments, herbal medicines, real-world data, pharmaco-epidemiological database, therapeutic efficacy

*Presenting author. E-mail address: drebka@em.uni-frankfurt.de (Alexandra Drebka)

S2.P144 An analysis of real-world data regarding therapeutic efficacy and adverse effects of herbal medicines in mental health issues

Annika J. Scholl^{a,*}, Alexandra Drebka^a, Teresa Ochs^a, Olaf Kelber^{b,c}, Ralph Mösges^{d,e}, Esther Raskopf^e, Kija Shah-Hosseini^d, Beatrice E. Bachmeier^a

^a Institute of Pharmaceutical Biology, Goethe University Frankfurt am Main, Max-von-Laue-Str. 9, 60438 Frankfurt am Main, Germany

^b R&D, Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt, Germany

^c Kooperation Phytopharmaka GbR, Bonn, Germany

^d Institute of Medical Statistics and Computational Biology, University of Cologne, Cologne, Germany

^e ClinCompetence Cologne GmbH, Cologne, Germany

Mental health issues (MHI) are commonly treated with psychotropic drugs which can cause central nervous, anticholinergic, and cardiovascular adverse effects (AE) besides compliance issues (Falkai et al., 2021). Herbal medicines (HM) could offer an alternative due to their therapeutic benefits and favorable tolerability (Salm et al., 2023). Nonetheless, they may also provoke AEs including drug interactions. In a data set of patients with MHIs taken from the pharmaco-epidemiological database PhytoVIS, we investigated the therapeutic benefits of HMs (mainly derived from *Valeriana officinalis* L., *Lavandula angustifolia* Mill., *Ginkgo biloba* L. and *Hypericum perforatum* L.) as well as the occurrence of AEs. The data set consisted of Patient-Reported Outcomes reflecting the real world of everyday medical care. Around 53% of the patients applied only HMs, while 45% took at least one additional medication. Over 90% perceived a therapeutic benefit from HMs while only few had unchanged or worsened symptoms. Around 87% did not perceive any AEs and only less than 3% reported significant AEs. Regarding AEs there was no statistically significant difference between patients who took HMs and those who took concomitant medication (Mann-Whitney U- Test: $p=0.084$). Likewise, there was no significant difference between those two groups regarding the severity of AEs (Mann-Whitney U-Test: $p=0.088$). Our results indicate that HMs have a favorable benefit-risk-ratio and a low potential for AEs. The integrated analysis of Real-World outcome data in context of clinical data will be a milestone to implement HMs into novel therapy strategies - not only for MHIs.

Keywords: herbal medicine, adverse effects, interactions, phytovis, real-world data, patient-reported outcomes

References

- Falkai, P., Laux, G., Möller, H., 2021. Duale Reihe Psychiatrie, Psychosomatik und Psychotherapie. (Vol. Hrsg. 7.). Thieme. <https://doi.org/10.1055/b000000071>.
- Salm, S., Rutz, J., van den Akker, M., Blaheta, R.A., Bachmeier, B.E., 2023. Current state of research on the clinical benefits of herbal medicines for non-life-threatening ailments. *Front Pharmacol.* 14, 1234701.

* Presenting author. E-mail address: an.scholl@em.uni-frankfurt.de (Annika J. Scholl)

S2.P145 Antiproliferative activity of homolupane triterpenoids

Lucie Rárová^{a,b,*}, Zbigniew Pakulski^c, Miroslav Strnad^d, Marie Kvasnicová^a, Tereza Štenclová^a, Piotr Cmoch^c

^a Department of Experimental Biology, Faculty of Science, Palacký University, Šlechtitelů 27, CZ- 78371 Olomouc, Czech Republic

^b Institute of Molecular and Translational Medicine, Faculty of Medicine and Dentistry, Palacký University, Hněvotínská 5, CZ-77900 Olomouc, Czech Republic

^c Institute of Organic Chemistry, Polish Academy of Sciences, Kasprzaka 44/52, 01-224 Warsaw, Poland

^d Laboratory of Growth Regulators, Institute of Experimental Botany of the Czech Academy of Sciences, and Faculty of Science, Palacký University, Šlechtitelů 27, CZ-78371 Olomouc, Czech Republic

From our previous studies, we searched for new bioactive triterpenoids and their derivatives based on the lupane scaffold, homolupanes. Source of these lupane-type triterpenes comes from outer part of birch bark (*Betula* species) (Hayek et al., 1989). Antiproliferative activity *in vitro* of these compounds was investigated in four human cancer cell lines and compared to healthy human BJ fibroblasts. In cervical carcinoma HeLa cells, three derivatives were the most promising with lower micromolar IC₅₀s and no toxicity to fibroblasts, thus showing a high therapeutic index. In addition, induction of apoptosis was detected in HeLa cells after 24 h treatment. This new series is more interesting than the previous lupane and homolupane triterpenes and saponins, due to their nontoxicity towards healthy human cells and stronger cytotoxicity to various cancer cell lines. This approach increases their potential as natural products-based anticancer agents. In addition, using other bioassays we proved also more biological properties, such as antiangiogenic or anti-inflammatory activities.

Keywords: lupanes, antiproliferative activity, apoptosis, cancer cells, fibroblasts

The work was supported by the European Union - Next Generation EU (The project National Institute for Cancer Research, Programme EXCELES, Project No. LX22NPO5102).

References

Hayek, E.W.H., Jordis, U., Moche, W., Sauter, F., 1989. A bicentennial of betulin. *Phytochemistry*. 28, 2229-2242.

* Presenting author. E-mail address: lucie.rarova@upol.cz (Lucie Rárová)

S2.P146 Anticancer and antioxidant properties of extracts from *Rubus caesius* L. (Rosaceae)

Anna Hering*, Justyna Stefanowicz-Hajduk, Piotr Graczyk, J. Renata Ochocka

Department of Biology and Pharmaceutical Botany, Medical University of Gdansk, M. Skłodowskiej-Curie 3a Str., 80-210 Gdansk, Poland

The European dewberry (*Rubus caesius* L., Rosaceae) is a medicinal shrub known and used for its pro-health properties in many parts of Europe and Asia. Extracts obtained from the leaves and stems of the European dewberry are a rich source of biologically active compounds with antioxidant, antiaging and antibacterial properties (Hering et al., 2022; Grochowski et al., 2016; Velickovic et al., 2015). Strong or long-lasting oxidative stress is often associated with the development of cancers. The European dewberry is a source of polyphenolic compounds with high ability to capture free radicals. In this poster authors present the results of anticancer and antioxidant properties of extracts obtained from young leaves and stems of the European dewberry. The anticancer properties of the European dewberry extracts (water and 50% ethanol extracts from stems and leaves) were tested *in vitro* on two cell lines - gastric adenocarcinoma (AGS) and human normal fibroblasts. The antioxidant activity of the extracts was estimated with DPPH, ABTS, FRAP and molibden reduction assays. The results indicated that the most active against gastric adenocarcinoma cells was ethanol extract obtained from leaves with $IC_{50} = 11.72 \mu\text{g/mL}$. Fibroblasts used as a control cells were not sensitive to the European dewberry extracts with the exception of water extract obtained from leaves, which reduced cell viability to about 60% at concentration of 400 $\mu\text{g/mL}$. The antioxidant test indicated, high ability of obtained extracts to reduce oxidative stress in similar way to standard- ascorbic acid.

Keywords: European dewberry, *Rubus caesius*, Rosaceae, fibroblasts, gastric cancer, antioxidant

References

- Hering, A., Stefanowicz-Hajduk, J., Hałasa, R., Olech, M., Nowak, R., Kosiński, P., Ochocka, J.R., 2022. Polyphenolic characterization, antioxidant, antihyaluronidase and antimicrobial activity of young leaves and stem extracts from *Rubus caesius* L. *Molecules*. 27, 6181.
- Grochowski, D.M., Paduch, R., Wiater, A., Dudek, A., Pleszczyńska, M., Tomczykowa, M., Granica, S., Polak, P., Tomczyk, M., 2016. *In vitro* antiproliferative and antioxidant effects of extracts from *Rubus caesius* leaves and their quality evaluation. *Evid. Based Complement. Altern. Med.* 2016, 5698685.
- Velickovic, I., Grujic, S., Dzamic, A., Krivosej, Z., Marin, P., 2015. *In vitro* antioxidant activity of dewberry (*Rubus caesius* L. var. *aquaticus* Weihe. & Nees.) leaf extracts. *Arch. Biol. Sci.* 2015, 67, 109.

* Presenting author. E-mail address: anna.hering@gumed.edu.pl (Anna Hering)

S2.P147 Broad antiplatelet activity of the flavones zapotin, chrysin, and tropoflavin in human platelet - rich plasma

Jakub Władysław Strawa^{a,*}, Agata Gołaszewska^b, Tomasz Misztal^c, Michał Tomczyk^a

^a Department of Pharmacognosy, Faculty of Pharmacy with the Division of Laboratory Medicine, Medical University of Białystok, ul. Mickiewicza 2A, 15-230 Białystok, Poland

^b Department of General and Experimental Pathology, Faculty of Medicine with the Division of Dentistry and Division of Medical Education in English Medical University of Białystok, ul. Mickiewicza 2C, 15-230 Białystok, Poland

^c Department of Physical Chemistry, Faculty of Pharmacy with the Division of Laboratory Medicine, Medical University of Białystok, ul. Mickiewicza 2B, 15-222, Białystok, Poland

Flavonoids identified in plants, have been shown to exert cardioprotective properties that can be attributed, at least partially, to their antiplatelet (anti-aggregation) activity (Guerrero et al., 2005). However, whether their potential is (platelet) response- and agonist-specific is still only partially established. Three flavones: zapotin (5,6,2',6'-tetramethoxyflavone), chrysin (5,7- dihydroxyflavone), and tropoflavin (7,8-dihydroxyflavone), were selected for evaluation of their antiplatelet properties in the presence of different platelet agonists, i.e., collagen, adenosine diphosphate (ADP), and the thromboxane A₂ receptor agonist U46619 in platelet-rich plasma (Strawa et al., 2022). Flow cytometry measurements (Gołaszewska et al., 2024) towards platelet secretion (quantified as P-selectin exposure), aggregation (measured as fibrinogen binding) and pro-coagulant response (evaluated as phosphatidylserine (PS) exposure) were performed in the presence of a 200 µM concentration of each flavone. It was established that zapotin reduced collagen-evoked platelet secretion and PS exposure by 35% and 52%, chrysin by 47% and 42%, and tropoflavin by 38% and 60%, respectively. ADP-induced secretion and aggregation was diminished by zapotin by 23% and 46%, by chrysin by 5% and 36%, and by tropoflavin by 37% and 52%, respectively. U46619-triggered secretion and aggregation was inhibited by zapotin by 58% and 60%, by chrysin by 20% and 27%, and by tropoflavin by 27% and 31%, respectively. Our preliminary results point to flavones as broad-range inhibitors of platelet responses triggered by different principal platelet agonists in a close-to-native platelet environment (platelet-rich plasma). Further prospective study comprising structure-activity relationship examination and revealing the specific mechanism of the observed antiplatelet effects is expected.

Keywords: flavones, blood platelets, flow cytometry, pro-coagulant response, P-selectin

References

- Guerrero, J.A., Lozano, M.L., Castillo, J., Benavente-Garcia, O., Vicente, V., Rivera, J., 2005. Flavonoids inhibit platelet function through binding to the thromboxane A₂ receptor. *J. Thromb. Haemost.* 3, 369-376.
- Gołaszewska, A., Misztal, T., Kazberuk, A., Rusak, T., 2024. Study on the mechanism of the adrenaline-evoked procoagulant response in human platelets. *Inter. J. Mol. Sci.* 25, 2997.
- Strawa, J.W., Jakimiuk, K., Kita, Z., Tomczyk, M., 2022. *In vitro* screening for anti-acetylcholinesterase and antioxidant activities of *Hottonia palustris* L. extracts and their unusual flavonoids. *Molecules.* 27, 8034.

* Presenting author. E-mail address: jakub.strawa@umb.edu.pl (Jakub W. Strawa)

S2.P148 Anti-aging, anti-neurodegenerative and healthspan-promoting compounds from *Dendrobium officinale* Kamura et Migo (Orchidaceae)

Okoro Nkwachukwu Oziamara*, Osadebe Patience Ogoamaka, Omeje Edwin Ogechukwu

Department of Pharmaceutical and Medicinal Chemistry, University of Nigeria, Nsukka, Nigeria

Aging is a degenerative process associated with a gradual decline in cellular components and functions, ultimately leading to death. The quest for longevity has led to the exploration of various medicinal substances for their potential anti-aging properties. This research delves into the anti-aging potential of polysaccharides extracted from *Dendrobium officinale* (DOP). The study investigated its effect on lifespan extension, stress resistance, healthspan improvement, among other parameters, in the *Caenorhabditis elegans* model. *D. officinale*, a widely used medicinal herb in Asia, has been reported to possess various medicinal properties, including anti-aging property. However, due to high demand and scarcity, alternative cultivation methods are being explored. In this study, DOP grown in three different environments (tree, greenhouse, and rock) were utilized. The results showed that at 1000 µg/mL, DOP from the greenhouse extended the mean lifespan by 14% and the maximum lifespan by 25%. DOP from all three sources enhanced resistance to H₂O₂-induced stress at 2000 µg/mL, with only rock-grown DOP exhibiting resistance to thermal stress. All DOP sources enhanced Endoplasmic Reticulum stress response, and decreased α-synuclein aggregation. However, only greenhouse-grown DOP delayed β-amyloid-induced paralysis. In conclusion, DOP exhibited promising anti-aging activity, and these findings provide valuable insights into its health benefits, adds to the growing list of potential anti-aging interventions, and suggests best practices for cultivating *D. officinale* for maximum medicinal applications.

Keywords: anti-aging, anti-neurodegenerative, *Dendrobium officinale* polysaccharides

References

López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., Kroemer, G., 2013. The hallmarks of aging. *Cell*, 153(6), 1194.

* Presenting author. E-mail address: nkwachukwu.okoro.pg01008@unn.edu.ng (Okoro, Nkwachukwu Oziamara)

S2.P149 Effects of STW 3-VI (St. John's wort) on the paracrine signaling between neurons and microglia under pro-inflammatory conditions *in vitro*

Laura Much^a, Heba Aziz-Kalbhenn^b, Marlon Palm^{b,*}, Olaf Kelber^b, Gabriel A. Bonaterra^a, Ralf Kinscherf^{1a}

^a Philipps-University of Marburg, Dept. of Medical Cell Biology, Marburg, Germany

^b Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt, Germany

Neurological insults and chronic stress are essential factors for inducing depression, affecting microglia function, so leading to enhanced release of pro-inflammatory cytokines, which causes a reduction of synaptic plasticity in mental disorders. Extracts of *Hypericum perforatum*, like STW 3- VI (SJW), showed protective, neurotrophic, and anti-inflammatory properties and are used in mild to moderate depressive episodes (Jungke et al., 2011). Aim was to determine the effects of SJW on the paracrine signalling under non- and pro-inflammatory conditions between neurons and microglia co-cultured *in vitro*. Co-cultured mouse SIM-A9 microglia and hippocampal HT-22 neurons were treated with different concentrations of SJW, hyperoside (HS), escitalopram (EC), nerve growth factor (NGF) as a positive control or ibuprofen (Ibu) as an anti-inflammatory control, with or without lipopolysaccharides (LPS). BDNF and TNF- α release were quantified by ELISA and neurite outgrowth by a staining kit. Treatment with SJW (250ng/ml) or HS (9 ng/mL) significantly stimulated the neurite formation by 3.8-fold and 2.3-fold, respectively, compared to negative control, similar to EC, Ibu or NGF. After LPS (250ng/ml) activation, neither SJW, HS, EC Ibu nor NGF induced neurite formation compared to LPS control. However, SJW, HS, EC, Ibu or NGF reduced the release of TNF- α for 20-30%. Microglia regulates neurite formation after treatment with SJW or HS, but not under pro-inflammatory conditions. SJW and HS have anti- inflammatory properties and positive effects on neuroplasticity, which might contribute to the clinically proven anti-depressant activity of STW 3-VI.

Keywords: *Hypericum perforatum* L., hypericaceae, anti-depressant, neuroplasticity, microglia, paracrine signalling

References

Jungke P., Ostrow, G., Li, J.L., Norton, S., Nieber, K., Kelber, O., Butterweck, V., 2011. Profiling of hypothalamic and hippocampal gene expression in chronically stressed rats treated with St. John's wort extract (STW 3-VI) and fluoxetine. *Psychopharmacology (Berl)*. 213, 757-772.

* Presenting author. E-mail address: marlon.palm@bayer.com (Marlon Palm)

S2.P150 Gut microbiota - a new aspect for understanding the antidepressant effects of *Hypericum perforatum*?

Maria-Eleni Grafakou^{a,*}, Eva-Maria Pferschy-Wenzig^a, Heba Aziz-Kalbhenn^b, Olaf Kelber^b, Christine Moissl-Eichinger^c, Rudolf Bauer^a

^a Institute of Pharmaceutical Sciences, Department of Pharmacognosy, University of Graz, Austria

^b Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt, Germany

^c Diagnostic and Research Institute of Hygiene, Microbiology and Environmental Medicine, Medical University of Graz, Austria

Hypericum perforatum L. is widely used for mild to moderate depressive disorders. Despite numerous clinical studies and meta-analyses, the mechanism of action and the responsible constituents are not fully elucidated. With the overall aim to investigate the role of gut microbiota for the activity of herbal preparations in mental health, a hydroethanolic extract of *H. perforatum* (STW 3-VI) was subjected to *in-vitro* upper gastrointestinal tract digestion (Brodkorb et al., 2019), and subsequently incubated *ex-vivo* with fecal samples from ten healthy donors (24h, anoxic, 37°C) (Pérez-Burillo et al., 2021). DNA extraction, followed by 16S rRNA gene sequencing, was used to evaluate the shift in microbial community composition during the incubation. Differential abundance analysis was performed using the combination of CLR abundances as input and Wilcoxon for dependent samples/Mann-Whitney U test for independent samples. The analysis showed that 2 phyla, 7 families, 9 genera and 21 OTUs were significantly modulated by treatment with *H. perforatum*. Many of the significantly changing taxa have been associated with depression. For example, *H. perforatum* treatment enhanced the Firmicutes/Bacteroidetes ratio, and decreased the levels of *Odoribacter* (p= 0.0007) and Erysipelotrichaceae_UCG-003 (p= 0.034) (opposite effects have been observed for individuals with depression (Barandouzi et al., 2020)). Provided that they also occur *in-vivo*, these changes in the microbial community composition suggest a potential additional non-CNS mechanism of action for the antidepressant effects of *H. perforatum*.

Conflicts of interest: The investigations and MEG have been funded by, HAK and OK are fully employed by Steigerwald Arzneimittelwerk, Bayer Consumer Health.

Keywords: *Hypericum perforatum*, gut microbiota, differential analysis, depression

References

- Barandouzi, Z.A., Starkweather, A.R., Henderson, W.A., Gyamfi, A., Cong, X.S., 2020. Altered composition of gut microbiota in depression: a systematic review. *Front. Psychiatry*. 11, 541.
- Brodkorb, A., Egger, L., Alming, M., Alvito, P., Assunção, R., Balance, S., Bohn, T., Bourlieu-Lacanal, C., Boutrou, R., Carrière, F., et al., 2019. INFOGEST static *in vitro* simulation of gastrointestinal food digestion. *Nat. Protoc.* 14, 991-1014.
- Pérez-Burillo, S., Molino, S., Navajas-Porras, B., et al., 2021. An *in vitro* batch fermentation protocol for studying the contribution of food to gut microbiota composition and functionality. *Nat. Protoc.* 16, 3186-3209.

* Presenting author. E-mail address: maria.grafakou@uni-graz.at (Maria-Eleni Grafakou)

S2.P151 The cytotoxic effect of chrysosplenetin, a natural product from *Artemisia annua*, investigated with different human cancer cell lines, as well as *in vitro* CYP₄₅₀ enzymatic activity in human liver microsomes

Gael Noel Neh Neba Ambe*, Avninder Bhambra, Randolph Arroo

Natural Product Research Laboratory, Faculty of health and life Sciences, De Montfort University, Leicester, United kingdom.

The polymethoxyflavone Chrysosplenetin [5-hydroxy-2-(4-hydroxy-3-methoxyphenyl)-3',6,7-trimethoxychromen-4-one] (Fig 1) is one of the many chemical compounds found in the aerial parts of the *Artemisia annua* plant (Ferreira et al., 2010). *A. annua* is best known as the source plant for artemisinin, the sesquiterpene precursor for a range of first-line antimalarial drugs (Tu, 2011; Woerdenbag et al., 1990). Our interest is in the polymethoxyflavones which are abundantly present in this medicinal crop plant, and which may find application as drugs in their own right. The cytotoxic effect of chrysosplenetin was tested against the triple negative breast cell line (MDA-MB- 468), a tumorigenic cell line which expresses CYP1, and the normal breast cell line (MCF10A) which is devoid of CYP1 expression.




Fig. 1. Structure of the polymethoxyflavone Chrysosplenetin

The antiproliferative effect of chrysosplenetin was investigated by MTT assay, and it showed greater toxicity in the CYP1-expressing breast cell line (MDA-MB-468) with a significantly lower IC₅₀ of 0.4 μ M as opposed to 20 μ M in the normal MCF10A cells. Additionally, the biotransformation of chrysosplenetin by CYP1 microsomes was investigated using High performance Liquid Chromatography (HPLC-DAD) and mass spectrometry (LC-MS). Two metabolite peaks were observed with retention times at 2.8 and 3.3 minutes, both presenting with an average mass of 361.3g in positive ionisation mode. We therefore propose that the polymethoxyflavone Chrysosplenetin (Fig. 1) may act as a prodrug that is selectively activated in hormone-dependent cancer cells.

Keywords: *Artemisia annua*, chrysosplenetin, cytotoxicity, CYP1 expression, biotransformation

References

- Tu, "The Discovery of Artemisinin (Qinghaosu) and Gifts from Chinese Medicine."
- Woerdenbag, H.J., Lugt, C.B., Pras, N., 1990. *Artemisia annua* L.: a source of novel antimalarial drugs. Pharm. Weekbl. Sci. Ed. 12, 169–181.
- Ferreira, J.F.S., Luthria, D.L., Sasaki, T., Heyerick, A., 2010. Flavonoids from *Artemisia annua* L. as antioxidants and their potential synergism with artemisinin against malaria and cancer. Molecules. 15, 3135.

* Presenting author. E-mail address: gael.nebaambe@dmu.ac.uk (Gael Noel Neh Neba Ambe)

S2.P152 Bioprospecting for plant extracts with protective effects on endothelial function

Alexandra Bujor^a, Anca Miron^{a,*}, Corine Girard^b, Céline Demougeot^b, Perle Totoson^b, Kerstin Jurk^c

^a Grigore T. Popa University of Medicine and Pharmacy, Iasi, Romania

^b PEPITE EA 4267, FHU INCREASE, University Bourgogne Franche-Comté, Besançon, France

^c Center for Thrombosis and Hemostasis, Johannes Gutenberg University of Mainz, Germany

Vascular endothelium is a metabolically active organ which releases substances involved in vasodilation, vasoconstriction, coagulation, fibrinolysis, inflammatory and immunological processes, thus regulating vascular homeostasis. Any alteration in vascular homeostasis leads to endothelial dysfunction which is involved in atherosclerosis, coronary artery disease and hypertension. Plant extracts and pure phytochemicals may protect endothelial function and prevent against various cardiovascular diseases (Bartáková et al., 2021). Extracts from *Crataegus pentagyna* Waldst. et Kit. ex Willd., *Cornus mas* L., *Sorbus aucuparia* L. and *Viburnum opulus* L. were investigated regarding potential vasorelaxant and antiplatelet activities. *V. opulus* fruit and *C. pentagyna* leaf extracts induced strong endothelium-dependent vasorelaxation effects ($EC_{50} = 6.31$ and $1.91 \mu\text{g/mL}$, respectively). The mechanism of action was found to be related to stimulation of endothelial nitric oxide synthase and suppression of arginase. Moreover, the extracts showed a decrease in vasorelaxation following endothelium removal. When investigating the antiplatelet effects, the same extracts proved to be the most active. *C. pentagyna* leaf extract significantly inhibited platelet aggregation ($IC_{50} = 0.109 \text{ mg/mL}$) induced by ristocetin (1 mg/mL), an inductor of platelet agglutination in the presence of von Willebrand factor. *V. opulus* fruit extract showed an inhibitory effect ($IC_{50} = 0.368 \text{ mg/mL}$) on platelet aggregation induced by convulxin (7 ng/mL), a potent activator of collagen receptor GP VI. *V. opulus* and *C. pentagyna* extracts are promising for the development of herbal preparations with potential benefits in cardiovascular diseases related to endothelial dysfunction.

Keywords: extracts, vascular endothelium, vasorelaxation, platelet aggregation

References

Bartáková, A., Nováková, M., 2021. Secondary metabolites of plants as modulators of endothelium functions. *Int. J. Mol. Sci.* 22, 2533.

* Presenting author. E-mail address: anca.miron@umfiasi.ro (Anca Miron)

S2.P153 A purified fraction from 70% hydroethanol extract of whole herb of *Platostoma africanum* P. Beauv. (Lamiaceae) inhibited formation of β -haematin formation

Abiodun Adeyem^{a,*}, Kamila Gromek^b, Rita Serrano^b, Jones Moody^a, Olga Silva^b

^a Department of Pharmacognosy, Faculty of Pharmacognosy, Faculty of Pharmacy, University of Ibadan, Ibadan, Nigeria

^b Research Institute for Medicines and Pharmaceutical Sciences (iMed.UL), Pharmacological Sciences Group, Faculty of Pharmacy, University of Lisbon, Av. Prof. Gama Pinto, 1649-019 Lisbon, Portugal

Compounds inhibiting formation of β -haematin are potential antiplasmodial substances especially when they are natural products from medicinal plants (Li et al., 2019). The inhibitors of β -haematin formation form an important category of partner drugs of ACTs, the WHO-recommended antimalarial regimens. New sources of inhibitors, however are needed because of widespread malaria parasite resistance (WHO, 2023). The whole herb of *Platostoma africanum* P. Beauv. is used in ethnomedicine in South-Western Nigeria to treat malaria. A bioactivity-guided fractionation of the 70% ethanol extract of the whole herb of the plant for inhibition of β -haematin formation has been presented. The 70% ethanol extract of the whole herb prepared by maceration was initially fractionated to give *n*-hexane (NH), diethyl ether (DE), ethyl acetate (EA), *n*-butanol (NB) and aqueous fractions (AQ). The crude extract (25 mg/mL) and fractions (20 mg/mL) were tested for inhibition of β -haematin formation (Vargas et al., 2011), expressed as I_{analysis} using quinine hemi-succinate as reference drug. Values of I_{analysis} considered were positive net absorbance computations and maximal inhibition corresponded to initial absorbance of haematin present. The *n*-butanol and aqueous fractions were fractionated (reversed-phase column chromatography, Sephadex) and analysed (TLC, LC/DAD). The order of inhibition of β -haematin formation was DE (0.046) < NH (0.056) < EA (0.062) < AQ (0.068) < NB (0.083). A purified isolate (10 mg/mL) from AQ showed I_{analysis} of 0.097 mg/mL lower than quinine hemi-succinate (I_{analysis} , 0.971). *Platostoma africanum* showed inhibition of β -haematin formation and an isolate from the aqueous fraction is potential antiplasmodial drug lead.

Keywords: *Platostoma africanum*, inhibition of β -haematin formation, antimalarial

References

- Li, F., Wang, Y., Li, D., Chen, Y. and Dou, Q.P., 2019. Are we seeing a resurgence in the use of natural products for new drug discovery? Expert Opin. Drug Discov. 14(5), 417–420.
- Vargas, S., Ioset, K.N., Hay, A.E., Ioset, J.R., Wittlin, S., Hostettmann, K., 2011. Screening medicinal plants for the detection of novel antimalarial products applying the inhibition of β -hematin formation. J. Pharm. Biomed. Anal. 56(5), 880-886.
- World Health Organisation 2023. World malaria report. Geneva, WHO Press, pp. viii.

* Presenting author. E-mail address: gokeadeyemi11@gmail.com (Abiodun Adeyemi)

S2.P154 Studies of St. Johns's Wort (*Hypericum perforatum* L.) dry extracts: no hints on mutagenicity

Jens Flade^a, Björn Feistel^{b,c}, Olaf Kelber^{c,d,*}, Heather Walker^e, Karen Nieber^f

^a University Leipzig, Postgradual Study Course Toxicology and Environmental Protection, Rudolf Boehm Institute for Pharmacology and Toxicology, Härtelstraße 16-18, Leipzig, Germany

^b Finzelberg GmbH & Co. KG, Koblenzer Str. 48-56, 56626 Andernach, Germany

^c Kooperation Phytopharmaka GbR, Plittersdorfer Str. 218, 53173 Bonn, Germany

^d R&D, Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Havelstraße 5, 64295 Darmstadt, Germany

^e Bayer Health Care, Whippany, New Jersey, USA

^f Institute for Pharmacy, Leipzig University, Brüderstraße 34, 04103 Leipzig, Germany

The genotoxic safety of preparations of *Hypericum perforatum* L., herba have already been assessed and reviewed over the last years (EMA, 2016), given that these preparations are used in the treatment of e.g. mild to moderate depressive episodes (EMA, 2016). One traditional form is the comminuted or powdered herbal substance alone. The way to test their genotoxic potential is the so called “bracketing and matrixing concept”, where extracts covering the whole range of polarities are tested, to obtain a representative sample of all components in the drug in these three extracts as a whole. Additional data on the genotoxic potential are therefore desirable to assess the therapeutic safety of these HMPs, using the Ames test according to the genotoxicity guideline of the Herbal Medicinal Product Committee HMPC of the European regulatory agency EMA. Three dry extracts of *Hypericum perforatum*, representing the whole spectrum of polarities of the extraction solvents (water – 50% ethanol (v/v) – n heptane) were tested in the Ames test, according to the OECD [5] and HMPC guidances. The extracts showed no mutagenic effect, even not in the highest concentrations according to the OECD guidance. The results of the tested dry extracts can be extrapolated by a “bracketing and matrixing concept” for other dry extracts of *Hypericum perforatum* in the tested polarity range of the extraction solvents. The data support the therapeutic safety of the extracts and the drug powder and add the assessment in the HMPC monograph of *Hypericum perforatum* (EMA, 2016).

Keywords: mutagenicity, ames test, genotoxic safety, St. John's wort, depression

References

EMA/HMPC/244315/2016

* Presenting author. E-mail address: olaf.kelber@bayer.com (Olaf Kelber)

S2.P155 Shortcutting iridals to irones transformation in orris root: unleashing the potential of microwave drying for rapid irone production

Mahdi Ayyari^{a,*}, Ghader Ghasemi^a, Mohammad-Hossein Azimi^b, Mohammad-Taghi Ebadi^a

^a Department of Horticultural Sciences, Tarbiat Modares University, Tehran, Iran

^b Ornamental Plants Research Center (OPRC), Horticultural Sciences Research Institute, Agricultural Research, Education and Extension Organization, Mahallat, Iran

Orris root is the *Iris germanica* L. rhizomes known as German iris, is famous for producing valuable irone compounds which are famous in perfume and food industry. Irones compounds normally derived from iridals, traditionally requiring several years of storage for slow oxidation transformation (Roger, et al., 2010). This study explores a shortcut to expedite irone compound production, significantly reducing the time required in German iris rhizomes. In fact, six drying methods (sun, shade, infrared, microwave, oven, and oven vacuum) were employed and compared for both whole and sliced rhizomes. Essential oil extraction was performed using a Clevenger type apparatus on the dried samples, followed by gas chromatography (GC) and gas chromatography- mass spectrometry analysis. The analysis of compounds showed that 35 compounds were identified in different drying methods, of which 8 compounds were found to be the main compounds (above 5%) (Ghasemi, et al., 2023). The results demonstrate that drying methods play a crucial role in irone compound production. cis- α -Irone and γ -Irone, flourished under microwave drying, constituting 71.86% and 18.24% in whole rhizomes (totally 90.1% irones), and 40.91% and 41.44% in sliced rhizomes (totally 82.35% irones), respectively. In contrast, the control fresh sample exhibited a minimal iron content of 0.20%. Other drying techniques could not reach to the microwave drying results. In fact, Microwave drying emerged as a game-changer, serving as a shortcut to rapidly synthesize irone compounds in orris root, by bypassing the traditional years-long oxidation process of iridals. In conclusion, these findings underscore the transformative potential of microwave drying, revolutionizing the traditional timeline for Irone synthesis in German iris rhizomes. By providing a time-saving alternative to the lengthy oxidation process of iridals, microwave drying streamlines fragrance development processes, enabling rapid and enhanced production of captivating scents and fragrances.

Keywords: *Iris germanica*, microwave drying, irone compounds, essential oil, fragrance development

References

- Roger, B., Fernandez, X., Jeannot, V. and Chahboun, J., 2010. An alternative method for irones quantification in iris rhizomes using headspace solid-phase microextraction. *Phytochem. Anal.* 21, 483-488.
- Ghasemi, G., Ayyari, M., Azimi, M-H., Ebadi, M-T., 2023. Orris root diversity and quality assessment: Multivariate analysis of phytochemicals and antioxidant properties. *Ind. Crops Prod.* 202, 116935.

* Presenting author. E-mail address: m.ayyari@nodares.ac.ir; mahdiayyari@gmail.com (Mahdi Ayyari)

S2.P156 Didesmethylocaglamide cytotoxic activity in high grade serous ovarian cancer in preclinical *in vitro* and *in vivo* models

Elizabeth N. Kaweesa^{a,*}, Tyler A. Wilson^b, Jaqueline M. Bazioli^a, Li Pan^b, A. Douglas Kinghorn^b, James R. Fuchs^b, Joanna E. Burdette^a

^a Department of Pharmaceutical Sciences, College of Pharmacy, University of Illinois at Chicago, Chicago, IL 60607, USA

^b Division of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, The Ohio State University, Columbus, OH 43210, USA

Didesmethylocaglamide (DDR) (**1**) is a naturally occurring derivative of rocaglamide with potent antitumor activity isolated from *Aglaia* plant species (Fig. 1) (Chang et al., 2020). DDR displays nanomolar cytotoxic activity in high grade serous ovarian cancer (HGSOC) cell lines including OVCAR8. HGSOC is the most lethal gynecological cause of death in women and requires new treatments to help tackle chemoresistance (Feng et al., 2021). Rocaglamides are known to function as translation inhibitors and trigger apoptosis in other types of solid tumors. We found that racemic **1** induced cytotoxicity in ovarian cancer cell lines as early as 24 hours and activated caspase-3 indicating pro-apoptotic activity. In addition, **1** was cytotoxic for the PEO4 and MCF7ADR cell lines that are resistant to cisplatin and paclitaxel, respectively. In addition, we evaluated **1** in OVCAR8 xenografts and observed a reduction in tumor burden. We hypothesized that combinatorial treatments of **1** with autophagy inhibitors like phyllanthusmin-34 would enhance cell death by blocking translation and the ability to recycle amino acids for protein production through autophagy (Salvi et al., 2022). The data obtained indicates that blocking autophagy or DNA repair in combination with **1** increased cell death in the sensitive and resistant models. Its effective cytotoxicity in drug sensitive and resistant cell models suggests **1** has potential as a new therapeutic strategy against HGSOC.




Fig. 1. Structure of (±)-didesmethylocaglamide (**1**)

Keywords: didesmethylrocaglamide, ovarian cancer, HGSOC, cytotoxicity, chemoresistance

References

- Chang, L.S., Oblinger, J.L., Burns, S.S., Huang, J., Anderson, L.W., Hollingshead, M.G., Shen, R., Pan, L., Agarwal, G., Ren, Y., Roberts, R.D., O'Keefe, B.R., Kinghorn, A.D., Collins, J.M., 2020. Targeting protein translation by rocaglamide and didesmethylrocaglamide to treat MPNST and other sarcomas. *Mol. Cancer Ther.* 19(3),731-741.
- Feng, L.Y., Yan, B.B., Huang, Y.Z., 2021. Abnormal methylation characteristics predict chemoresistance and poor prognosis in advanced high-grade serous ovarian cancer. *Clin. Epigenet.* 13, 141.
- Salvi A., Young A.N., Huntsman, A.C., Pergande, M.R., Korkmaz, M.A., Rathnayake, R.A., Mize, B.K., Kinghorn, A.D., Zhang, X., Ratia, K., Schirle, M., Thomas, J.R., Brittain, S.M., Shelton, C., Aldrich L.N., Cologna, S.M., Fuchs, J.R., 2022. Burdette JE.PHY34 inhibits autophagy through V-ATPase V0A2 subunit inhibition and CAS/CSE1L nuclear cargo trafficking in high grade serous ovarian cancer. *Cell Death Dis.* 13(1), 45.

*Presenting author. E-mail address: ekaweesa@uic.edu (Elizabeth Kaweesa)

S2.P157 The herbal preparation, STW 5, protects against changes in the microbiome in experimental functional dyspepsia

Mohamed T. Khayyal^{a,*}, Olaf Kelber^b, Heba Aziz-Kalbhenn^b

^a Department of Pharmacology, Faculty of Pharmacy, Cairo University, Cairo 11562, Egypt

^b Bayer Consumer Health, Darmstadt, Germany

The herbal preparation, STW 5, consists of nine hydro-alcoholic (31 % alcohol) extracts of *Iberis amara*, *Matricaria recutita*, *Carum carvi*, *Melissa officinalis*, *Mentha piperita*, *Glycyrrhiza glabra*, *Silybum marianum*, *Angelica archangelica* and *Chelidonium majus*. It has been shown to be clinically effective in irritable bowel syndrome (IBS) and functional dyspepsia (FD), but the involvement of the intestinal microbiota has not been adequately studied. The aim of the present study was to examine whether changes in gut microbiota induced in an experimental model of functional dyspepsia could be prevented by STW 5. FD was induced in rats by subjecting them to neonatal maternal separation followed by subjecting them to restraint stress for 90 minutes/day for one week. During that week, one group (8 animals) was treated with the liquid form of STW 5 (5 ml/Kg) and one with its vehicle (31% alcohol) at the same dose level. Animals were sacrificed 24 h after the last drug administration. Fecal samples were taken from the cecum. Genomic DNA was isolated and changes in selected bacterial phyla and genera were assessed using quantitative Real Time-PCR (qPCR). The main phyla studied under the influence of stress in the model were the Bacteroidetes, Firmicutes, Fusibacterium and Actinobacteria. FD led to a 50% drop in the *Lactobacilli* population but to a two-fold rise in *Proteobacteria* abundance but did not affect significantly the changes in *Bacteroidetes*. Furthermore, the *Methanobrevibacter* population was raised more than two-fold while the genomic DNA concentration in *Bifidobacterium* dropped from ca. 3 ng/g feces to nearly nil. In general, STW 5 tended to ameliorate and protect against the changes induced by the stress model. The results lend further support to the use of STW5 in FD and IBS by providing evidence that it acts at least in part through influencing beneficially the changes in intestinal microbiota induced by stress. Furthermore, the findings shed more light on mechanisms involved in potential usefulness of STW 5 in gastro-intestinal disorders.

* Presenting author. E-mail address: mtkhayyal@gmail.com (Mohamed T. Khayyal)

S2.P158 Modulatory effects of *Glycyrrhiza glabra* L. on endocrinological shift in ovariectomy-induced menopausal mice: Network pharmacology guided *in-vivo* study

Poonam Rawat^{*,a,b}, Shekhar Singh^b, Priyanka Kumari Keshri^b, Hagera Dilnashin^b, Bhanu Kumar^a, Ankita Misra^a, Surya Pratap Singh^b, Sharad Srivastava^a

^a Pharmacognosy division, CSIR-National Botanical Research Institute, Lucknow, UP 226001, India

^b Department of Biochemistry, Institute of Science, Banaras Hindu University, Varanasi, UP 221005, India

Menopause is a major endocrinological shift leading to hot flushes, anxiety, mood swings, and increased vulnerability for cognitive impairment, hypertension, and obesity in women. In this condition, an abrupt decline in estrogen and serotonin is directly associated with increased expression of MAO-A. Hormonal replacement therapy in post-menopausal conditions has limitations, including increased risk for estrogen-sensitive cancers and cardiovascular diseases. A more promising therapeutic approach lies in the use of medicinal plants with estrogenic as well as serotonergic activity. The present study investigated the neuromodulatory effects of LC-MS characterized *Glycyrrhiza glabra* extract (GGE) against ovariectomy-induced (OVX) menopausal mice through *in-vivo* study. Four experimental groups of mice were formed: sham (0.9% saline, oral), OVX only (0.9% saline, oral), OVX+E2 (30 µg/ kg b.w., s.c.) and OVX+GGE (150 mg/kg b.w.). After 6 weeks of dosing, mice underwent behavioural analysis for memory, anxiety, depression, hot flushes and body weight. The expression analysis of targeted proteins was done through immunoblotting and immunohistochemistry. Increase in rectal temperature, body weight, anxiety and loss of spatial memory was observed in OVX only group, suggesting severity of menopause. Further, in the hippocampus and hypothalamus region of OVX only group, elevated MAO-A and decreased TPH-2 as well as ERβ expression was observed. However, in OVX+E2 and OVX+GGE groups we found a significant reduction in MAO-A expression while increased expression of TPH-2 and ERβ. GGE also enhanced the blood serum levels of estrogen and serotonin. Altogether, GGE is effective against the endocrinological shift generated through surgical menopause *via* modulating vasomotor symptoms.

Keywords: post-menopausal symptoms, ovariectomy, *Glycyrrhiza glabra*, estrogen, anxiety

* Presenting author. E-mail address: rwtpoonam203@gmail.com (Poonam Rawat)

S2.P159 *Ziziphus nummularia* crude and fractionated extracts attenuate the malignant phenotype of human triple negative breast cancer cells

Abdullah A. Shaito^{a,*}, Rola Abdallah^b, Mansour Sobeh^c, Nihad Sahri^c, Joelle E. Mesmar^b, Elias Baydoun^b

^a Biomedical Research Center, Qatar University, Doha, Qatar

^b Department of Biology, American University of Beirut, Beirut, Lebanon

^c Agrobiosciences Program, College for Agriculture and Environmental Science, Mohammed VI Polytechnic University, Ben Guerir, Morocco

Triple negative breast cancer (TNBC) is the most aggressive subtype of breast cancer and has limited therapeutic options. Effects of *Ziziphus nummularia* against TNBC have not been investigated yet. An ethanolic extract of *Ziziphus nummularia* (Burm.f.) Wight & Arn. (ZNE) was prepared and chromatographically fractionated. Phytochemical composition of ZNE and its chromatographically isolated fraction (F6) was identified both qualitatively by spectrophotometric assays and analytically by HPLC-PDA-MS/MS. Effects of ZNE and F6 on the viability of several cancerous cell lines were tested by MTT assay. The anti-cancerous potential of ZNE and F6 was tested *in vitro* in MDA-MB-231 cells, a TNBC cell line. ZNE and F6 radical scavenging capacity was tested using DPPH assay, and their effects on reactive oxygen species (ROS) generation *in vitro* in cells by DCFDA staining. Propidium iodide-based FACS analysis was used for cell cycle analysis. Scratch wound healing and trans-well migration chamber assays were used to assess MDA-MB-231 cell migration and invasion. Western blotting analysis was used to analyse changes in the levels of cell cycle, apoptosis and autophagy proteins. Findings showed that ZNE and F6 reduced the viability of several cancerous cell lines including MDA-MB-231 cells. F6 decreased MDAMB-231 viability more than crude ZNE or F6. ZNE and F6 are rich in phytochemicals and HPLC-PDA-MS/MS analysis identified several metabolites that were previously reported to have anti-cancerous effects. Both ZNE and F6 showed potent antioxidant capacity in the DPPH assay, but promoted reactive oxygen species (ROS) production in MDA-MB-231 cells; an effect which was blunted by the antioxidant N-acetyl cysteine (NAC). NAC also blunted ZNE- and F6-induced reduction in TNBC cell viability. We also demonstrated that ZNE and F6 induced an arrest of the cell cycle, and triggered apoptosis- and autophagy-mediated cell death. ZNE and F6 inhibited metastasis-related cellular processes by modifying cell migration, invasion, and adhesion. Collectively, our findings reveal that *Z. nummularia* is rich in metabolites that can attenuate the malignant phenotype of TNBC and may provide novel approaches for the discovery of new drug leads for treatment of TNBC and other cancers.

* Presenting author. E-mail address: abdshaito@qu.edu.qa (Abdullah A. Shaito)

S2.P160 Anti-inflammatory properties of jaboticaba peel bioactive compounds during *in vitro* gastrointestinal and colonic digestion

Livia Reguengo^{a,*}, Ana Sofia Saliba^b, Mateus Salgaço^c, Katia Sivieri^c, Severino de Alencar^{b,d}, Mário Roberto Maróstica^a

^a Faculty of Food Engineering, University of Campinas (UNICAMP), Campinas, SP, Brazil

^b Centro de Energia Nuclear na Agricultura, Universidade de São Paulo (USP), Piracicaba, SP, Brazil

^c Postgraduate Program of Food and Nutrition, School of Pharmaceutical Sciences, State University of São Paulo (UNESP), Araraquara, SP, Brazil

^d Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo (USP), Piracicaba, SP, Brazil

Food rich in polyphenols are correlated to health benefits in literature, especially for their anti-inflammatory potential. The bioactivities of these compounds are directly correlated to their bioavailability during digestion. Jaboticaba, aka. "Brazilian berry", is a small fruit with dark purple peel, normally discarded, which is rich in anthocyanins, tannins and phenolic acids. Despite the extensive research on jaboticaba peel (JP) health benefits, the bioavailability of its bioactive compounds is not always considered. Therefore, the objective of this work was to evaluate the influence of the digestion of JP in a Simulator of Human Intestinal Microbial Ecosystem on its phenolic composition, antioxidant capacity and anti-inflammatory effects in RAW 264.7 macrophages. Antioxidant capacity analysis presented a slight divergence, except for JP extract superiority to stomach and duodenum values. However, FRAP presented lower antioxidant results for colonic samples than JP extract, while colonic samples results for ORAC overcame JP extract. As expected, anthocyanins are degraded in stomach, and in duodenum in a greater extent, with a small recovery on colonic samples, followed by a significant decay after 7 hours of colonic fermentation. JP extract (500 µg/mL) effectively reduced NF-κB, TNF-α and MIP-2 levels by 94,2%, 78,6% and 94,2%, respectively. Most of the samples have shown valuable TNF- α modulation, with the lowest inhibition at 10%. However, NF-κB and MIP-2 could only be modulated to a certain extent after JP digestion, being only stomach and duodenum samples effective regarding NF-κB, whereas reductive effects could be observed in MIP-2 until 8 hours of digestion.

Keywords: *Plinia cauliflora*, *in vitro* digestion, macrophages, NF-Kb pathway, antioxidant capacity

* Presenting author. E-mail address: liviareguengo@gmail.com (Lívia Mateus Reguengo)

S2.P161 Improving sleep quality and daytime function with a synergistic combination of tryptophan, magnesium, *Lactuca*, and *Melissa*: A prospective pilot study in adults with sleep disturbances

Franziska Eckert^a, Ramon Weishaupt^{b,*}, Dominic Vogt^a, Nadin Meyer^a, Juliane Hellhammer^a

^a Contract Research Daacro, Trier, Germany

^b A.Vogel AG. Medical Department, Grünaustrasse 4, 9325 Roggwil, Switzerland

Insufficient sleep is a widespread issue, with existing treatments often leading to side effects (1). This monocentric, single-arm, open-label study assessed a novel granulate formulation containing fresh *Lactuca sativa*, *Melissa officinalis*, Tryptophan, and Magnesium in adults with sleep disturbances. Conducted in Germany in 2023, 50 participants consumed the formula nightly for 14 days, with outcomes measured through diaries, questionnaires, cognitive tests, wearables, saliva samples, and in 10 cases, polysomnography (PSG). Descriptive statistics were used to analyze outcomes. The results showed significant improvements: nightly awakenings reduced by 31% ($p < 0.001$) and early morning awakenings by 16% ($p \leq 0.003$). PSG confirmed a 28% increase in overall deep sleep ($p > 0.05$) and a 70% increase in deep sleep phase N4 ($p = 0.042$), while REM sleep decreased by 18% ($p > 0.05$). The Apnea/Hypopnea Index scores also dropped by 26% ($p > 0.05$), suggesting a lower apnea/hypopnea risk. The primary outcome, a 14% improvement in the SF-B/R Sleep Quality Index ($p = 0.003$), highlighted enhanced sleep quality overall, especially in individuals with high anxiety, who saw a 37% improvement ($p \leq 0.001$). Further exploratory analysis using wearables, PSG, and saliva bioanalysis yielded inconsistent results. Furthermore, the study reported a 22-28% increase in restedness ($p \leq 0.001$), up to a 29% reduction in psychological tension ($p \leq 0.01$), and improved daytime performance, including a 13% decrease in daytime sleepiness and a 23% improvement in mood ($p \leq 0.014$). Cognitive function improved by 13% ($p \leq 0.001$), tested by objective computerized test batteries (COMPASS). Adherence to the intervention was high, with no serious adverse events reported. In conclusion, the tested food supplement formulation safely improves sleep quality, deep sleep, and psychological well-being, enhancing daytime mental function and mood.

Keywords: *Lactuca sativa*, clinical trial, food supplement, deep sleep improvement (PSG), cognitive improvements (COMPASS)

References

Morin, C.M., Benca, R., 2012. Chronic insomnia. *Lancet* 379(9821), 1129-1141.

* Presenting author. E-mail address: r.weishaupt@avogel.ag (Ramon Weishaupt)

S2.P162 Harnessing gut microbiota through immunomodulation by LM-1 attenuates HDM-induced asthma

Chuan-Teng Liu^a, Ying-Chyi Song^b, Michael Heinrich^{c,d,e}, Hung-Rong Yen^{a,e,f,*}

^a Research Center for Traditional Chinese Medicine, China Medical University Hospital, Taichung, Taiwan

^b Graduate Institute of Integrated Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^c UCL School of Pharmacy, University College London, London, UK

^d Department of Chinese Pharmaceutical Sciences and Chinese Medicine Resources, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^e Chinese Medicine Research Center, China Medical University, Taichung, Taiwan

School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

The chronic recurrent exacerbated nature of asthma and the side effects of long-term use of conventional medicine for asthma lead patients to seek out novel therapies. In addition to Th2- dominated immune response, the gut microbiome may represent a key target for the management of allergic asthma. Liu-Jun-Zi-Tang (LJZT; decoction containing *Panax ginseng*, *Atractylodes macrocephala*, *Poria cocos*, *Pinellia ternate*, *Glycyrrhiza uralensis*, *Citri reticulatae*, *Zingiber officinale*, and *Zizyphus jujuba*) and Ma-Xing-Gan-Shi-Tang (MXGST; decoction containing *Ephedra sinica*, *Prunus armeniaca*, *Glycyrrhiza uralensis*, and *Gypsum Fibrosum*) are two frequently prescribed Chinese herbal granules for the treatment of pulmonary diseases in Taiwan. Here we investigate whether LJZT combined with MXGST (abbreviated to LM-1) ameliorates asthma through the immunomodulation of the gut-lung axis. LM-1 (0.9 g/kg/day of MXGST + 1.8 g/kg/day of LJZT, oral for 5 days) attenuated pulmonary allergic airway inflammation by inhibiting Th2 cell, eosinophil and mucosubstances accumulation in asthma mice. LM-1 in combination with dexamethasone improved pulmonary function. LM-1 treatment altered gut microbiota, and faecal microbiota transplantation from LM-1-treated mice ameliorated asthma by regulating intestinal IFN- γ /IL-4 ratio (Fig. 1). Furthermore, LM-1 reduced CD4⁺IL-4⁺ cells in the PBMCs of patients with asthma. We conclude that LM-1 attenuates asthma by enhancing IFN- γ /IL-4 ratio through gut immunoregulation. LM-1 is a potential candidate for further clinical investigation.




Fig. 1. Schematic of LM-1's effects on the immunomodulation of the gut-lung axis against asthma

Keywords: traditional Chinese medicine, microbiota, asthma, immunomodulation, fecal microbiota transplantation

* Presenting author. E-mail address: s78981030@gmail.com (Chuan-Teng Liu)

S2.P163 Ethnopharmacological study of the bark traditionally used for diabetes of *Croton guatemalensis* Lotsy: from the field to the action mechanisms

Adolfo Andrade-Cetto*, Sonia Escandón-Rivera, Daniela Moreno-Vargas

Laboratory of Ethnopharmacology, Department of Cellular Biology, School of Sciences, National Autonomous University of Mexico, 04510 Mexico City, Mexico

The aim of this presentation is to address the ethnopharmacological study of the bark of *Croton guatemalensis* Lotsy (Euphorbiaceae). Commonly known as “copalchi”, it was identified as one of the most prominent species traditionally used for the treatment of diabetes in a field study conducted among the Cakchiquels of Chimaltenango in Guatemala (Cruz and Andrade-Cetto, 2015). Posteriorly, the hypoglycemic and antihyperglycemic effects of the ethanol-water extract were confirmed, where its ability to counteract fasting hyperglycemia was highlighted due to its “glibenclamide-like” effect and where the inhibition of α -glucosidase activity was ruled out as a potential effect to manage postprandial hyperglycemia (Andrade-Cetto et al., 2019). Subsequent phytochemical studies revealed the identification of eight compounds: five *ent*-clerodane diterpenoids and three flavonoids (Escandón-Rivera et al., 2022). Among them, junceic acid (**1**) stands out for being the most abundant (Fig. 1)




Fig. 1. Structure of junceic acid (**1**).

Later, it was confirmed the hypoglycemic effect of **1**, which was similar to that exerted by the extract. One of the major mechanisms presumably involved in the control of fasting hyperglycemia was the inhibition of hepatic glucose production. Notably, **1** inhibited the activity of the rate-limiting enzyme glucose-6-phosphatase more potently than the control. Moreover, both extract and **1** showed a similar capacity to reduce postprandial hyperglycemia. In the long term, the extract reduced fasting hyperinsulinemia and improved oral glucose intolerance by restoring the glucose-stimulated insulin secretion response and enhancing insulin signalling in the liver. In addition, it showed to decrease hepatic triglycerides and cholesterol.

Keywords: *Croton guatemalensis*, junceic acid, ethnopharmacology, diabetes, action mechanisms

References

- Andrade-Cetto, A., Cruz, E.C., Cabello-Hernández, C.A., Cárdenas-Vázquez, R., 2019. Hypoglycemic activity of medicinal plants used among the cakchiquels in Guatemala for the treatment of type 2 diabetes. *e-CAM*, 2019, 2168603.
- Cruz, E.C., Andrade-Cetto, A., 2015. Ethnopharmacological field study of the plants used to treat type 2 diabetes among the Cakchiquels in Guatemala. *J Ethnopharmacology*. 159, 238–244.
- Escandón-Rivera, S.M., Andrade-Cetto, A., Rosas-Ramírez, D.G., Arreguín-Espinosa, R., 2022. Phytochemical Screening and Isolation of New *Ent*-Clerodane Diterpenoids from *Croton guatemalensis* Lotsy. *Plants* 11(22), 3159.

* Presenting author. E-mail address: aac@ciencias.unam.mx (Adolfo Andrade-Cetto)

S2.P164 *Eryngium cymosum* F. Delaroche, a Mexican species traditionally used in the treatment of diabetes - ethnopharmacological insights

Adolfo Andrade-Cetto*, Fernanda Espinoza-Hernández

Laboratory of Ethnopharmacology, Department of Cellular Biology, School of Sciences, National Autonomous University of Mexico, 04510 Mexico City, Mexico

Eryngium cymosum F. Delaroche (Apiaceae), commonly known as “piñuela”, was identified as a recurring remedy against diabetes through a field study carried out in the Huejutla market located in Hidalgo, Mexico (Espinoza-Hernández et al., 2021). The aqueous extract elaborated from the aerial parts demonstrated to manage fasting and postprandial hyperglycemia, in which the inhibition of hepatic glucose production and the enhancement of insulin function were implicated. Specifically, it showed to inhibit the activity of two rate-limiting enzymes of liver glucose output (glucose-6-phosphatase and fructose-1,6-bisphosphatase) and protein tyrosine phosphatase 1B. In addition, the extract was considered safe in an acute toxicological test. Phytochemical analysis revealed the presence of phenolic acids, such as chlorogenic and rosmarinic acids, and flavonoids in the aqueous extract (Romo-Pérez et al., 2022). Interestingly, kaempferol-3-O-(2,6-di-O-trans- ρ -coumaryl)- β -D-glucopyranoside (1) (Fig. 1) showed to potently inhibit both glucose-6-phosphatase and fructose-1,6-bisphosphatase, suggesting an important role in the inhibitory effect on hepatic glucose production.




Fig. 1. Structure of kaempferol-3-O-(2,6-di-O-trans- ρ -coumaryl)- β -D-glucopyranoside (1).

When administrated chronically, the extract could reduce glycated hemoglobin, indicating a potential long-term glycemic control (Espinoza-Hernández and Andrade-Cetto, 2022). Moreover, it also enhanced insulin sensitivity by increasing the glucose disappearance rate and improved oral glucose intolerance by restoring the glucose-stimulated insulin secretion response. In addition, the extract decreased Akt hyperphosphorylation and promoted a better insulin signalling response in muscle.

Keywords: *Eryngium cymosum*, ethnopharmacology, diabetes, glycated hemoglobin, insulin sensitivity

References

- Espinoza-Hernández, F., Andrade-Cetto, A., Escandón-Rivera, S., Mata-Torres, G., Mata, R., 2021. Contribution of fasting and postprandial glucose-lowering mechanisms to the acute hypoglycemic effect of traditionally used *Eryngium cymosum* F. Delaroche. J. Ethnopharmacology. 279, 114339.
- Espinoza-Hernández, F.A., Andrade-Cetto, A., 2022. Chronic Antihyperglycemic effect exerted by traditional extracts of three Mexican medicinal plants. eCAM, 2022, 5970358.
- Romo-Pérez, A., Escandón-Rivera, S.M., Miranda, L.D., Andrade-Cetto, A., 2022. Phytochemical study of *Eryngium cymosum* F. Delaroche and the inhibitory capacity of its main compounds on two glucose-producing pathway enzymes. Plants 11(7), 992.

* Presenting author. E-mail address: aac@ciencias.unam.mx (Adolfo Andrade-Cetto)

S2.P165 Onychocolone A produced by the fungus *Onychocola* sp. targets cancer stem cells and stops pancreatic cancer progression by inhibiting MEK2-dependent cell signaling

Maria C Ramos^a, Gloria Crespo-Sueiro^a, Nuria de Pedro^a, Carmen Griñán-Lisón^b, Caridad Díaz^a, Víctor González-Menéndez^a, Francisco Castillo^a, Jose R Tormo^a, Olga Genilloud^a, Fernando Reyes^a, Rosario Fernández-Godino^{a,*}

^a Fundación MEDINA, Centro de Excelencia en Investigación de Medicamentos Innovadores en Andalucía. Parque Tecnológico Ciencias de la Salud, Avda. del Conocimiento 34, 18016, Armilla, Granada, Spain

^b Instituto de Investigación Biosanitaria ibs.GRANADA, University Hospitals of Granada- University of Granada, 18100, Granada, Spain

Pancreatic cancer (PC) remains one of the deadliest cancers and it lacks targeted therapy. PC tumors are enriched with subpopulations of cancer stem cells (CSCs) that are resistant to antiproliferative drugs (Hermann et al., 2007), which raises a need to identify new chemotherapeutic agents. Onychocolones A-F (#1-6) are new anti-tumoral benzophenones compounds identified in a high throughput screening on MEDINA's microbial Natural Product extract library. The compounds were purified by bioassay-guided isolation from an extract of the fungal strain *Onychocola* sp. CF- 107644 and their structures were established by spectroscopic methods. Compounds #1-4 inhibited the growth of the pancreatic tumor cell lines MIA PaCa-2, BxPC-3, and PANC-1 with low- micromolar Median Effective Doses (ED₅₀). Compound #1 (onychocolone A) showed pro-apoptotic effect on pancreatic CSCs and 3D spheroids, which are representative of tumor complexity (Langhans, 2021). The mechanism of action was linked to the inhibition of MEK onco-signaling pathway by protein expression assays. The efficacy of onychocolone A *in vivo* was confirmed in a heterotopic pancreatic xenograft mouse model generated by CSCs (Hernández-Camarero et al., 2019). ADME studies validated the drug-like properties of onychocolone A, including low probability of pharmacologic interaction, prediction of good absorption for oral administration, high protein binding, slow release, and a short half-life compatible with low toxicity. Altogether, the data suggest the potential of onychocolone A as a new small molecule for hit-to-lead development of a new treatment for PC.

Keywords: onychocolone, pancreatic cancer, cancer stem cells, 3D spheroids, MEK

References:

- Hernández-Camarero, P., López-Ruiz, E., Griñán-Lisón, C., García, M.Á., Chocarro-Wrona, C., Marchal, J.A., Kenyon, J., Perán, M., 2019. Pancreatic (pro)enzymes treatment suppresses BxPC-3 pancreatic cancer stem cell subpopulation and impairs tumour engrafting. *Sci. Rep.* 9, 11359.
- Hermann, P.C., Huber, S.L., Herrler, T., Aicher, A., Ellwart, J.W., Guba, M., Bruns, C.J., Heeschen, C., 2007. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem. Cell*, 1, 313–323.
- Langhans, S.A., 2021. Using 3D *in Vitro* cell culture models in anti-cancer drug discovery. *Expert Opin. Drug Discov.* 16, 841–850.

* Presenting author. E-mail address: rosario.fernandez@medinaandalucia.es (Rosario Fernández-Godino)

S2.P166 Human prolyl endopeptidase inhibitory activity of the peptide enriched extracts of *Momordica charantia* (Cucurbitaceae) fruits

Augustine Mbanu^{a,*}, Jasmin Gattringer^b, Roland Hellinger^b, Christian W. Gruber^b

^a Department of Pharmacognosy, University of Ibadan, Oduduwa Road, 200005 Ibadan, Nigeria

^b Centre for Physiology and Pharmacology, Medical University of Vienna, Schwarzschanerstrasse 17A, 1090 Vienna, Austria

Prolyl endopeptidase (PREP) is a proline-specific serine protease that is expressed in several body tissues. In humans, PREP was shown to be involved in blood pressure regulation and central nervous system disorders. Plant peptides are known inhibitors of PREP, for instance the prototypic cyclic cystine-rich peptide psysol 2 from *Psychotria solitudinum* has inhibitory activity in the low micromolar range. *Momordica charantia* has been reported to contain cysteine-rich peptides that can inhibit trypsin, elastase and other serine proteases in nanomolar concentrations. This study will focus on the extraction of *M. charantia* cysteine-rich peptides for protease inhibition assays against human PREP. Four extraction methods for *M. charantia* fruits were applied in a comparative study, where the boiled water extraction proved superior. The crude extract was further purified using C₁₈ solid phase extraction (SPE) to obtain peptide enriched samples by eluting peptides with stepwise increasing concentrations (5 – 45%) of acetonitrile. The presence of peptides was confirmed using MALDI-TOF/TOF-MS and HPLC-UV techniques. PREP inhibition of the peptide enriched fractions was studied using a fluorescent substrate-based *in vitro* assay. Fractions eluted between 15 – 30% showed considerable inhibitory potential (IC₅₀ from 32 – 56 µg/mL) while the 20% fraction exhibited the most significant inhibition (IC₅₀ = 32.4 ± 5.0 µg/mL). The peptide enriched extracts of *M. charantia* fruits demonstrated inhibitory activity against PREP. Next steps, we will isolate peptides from *M. charantia* by bioassay-guided isolation and characterize their potential for PREP inhibition, thus providing new leads for drug development in disease states where PREP is implicated.

Keywords: *Momordica charantia*, natural peptides, Cucurbitaceae, human prolyl endopeptidase, solid phase extraction

* Presenting author. E-mail address: augustinembanu20@gmail.com (Augustine Mbanu)

S2.P167 Impact of phytocannabinoids THC, CBD and their acidic counterparts THCA and CBDA on TGF- β 1-induced epithelial-mesenchymal transition in human conjunctival epithelial cells

Anil Baskan*, Ezzat M. Awad, Ava Elahi, Javeria Pervaiz, Talin Barisani-Asenbauer

Institute of Specific Prophylaxis and Tropical Medicine, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, Vienna, Austria

Epithelial-mesenchymal transition (EMT) is a process in which epithelial cells lose their intercellular connections and acquire a fibroblast-like phenotype with an increased migratory capacity and extracellular matrix secretion (Thiery et al., 2009). EMT was suggested to be involved in the development of fibrosis in various tissues including the eye and TGF- β 1 is known as its potent promoter (Rajić et al., 2020). Endogenous cannabinoids and cannabinoid receptors are found throughout the eye, indicating their potential significance in ocular physiology (Cairns et al., 2016). We assessed the effects of Δ 9-tetrahydrocannabinol (THC), Cannabidiol (CBD), Δ 9-tetrahydrocannabinolic acid (THCA), and Cannabidiolic acid (CBDA) on TGF- β 1-induced EMT in cultured immortalized human conjunctival epithelial (HCjE) cells. Cells were exposed to TGF- β 1 (10 ng/mL) for 72 hours with or without cannabinoids (2 to 10 μ M). TGF- β 1 induced an elongated spindle-shaped appearance in cells. Cells stimulated with TGF- β 1 in presence of THC or CBD remarkably maintained cuboidal epithelial morphology, whereas THCA partially restored the epithelial phenotype of the cells and CBDA showed no effect. E-cadherin, a marker of epithelial phenotype, was assessed by immunocytochemistry. THC and CBD, but not THCA or CBDA, increased E-cadherin expression, which was downregulated by TGF- β 1. A cell migration assay was conducted by introducing a cell free area in confluent cells and subsequently adding TGF- β 1 $\pm$ cannabinoids. At 24 hours, TGF- β 1 induced a complete wound closure. All cannabinoids decreased wound closure in a dose dependent manner, whereas CBD showed the strongest anti-migratory effect. In conclusion, THC and CBD effectively counteracted EMT-related alterations induced by TGF- β 1.

Keywords: cannabinoids, epithelial-mesenchymal transition (EMT), TGF- β 1, conjunctival fibrosis, human conjunctival epithelial cells (HCjE)

References

- Cairns, E.A., Baldrige, W.H., Kelly, M.E.M., 2016. The endocannabinoid system as a therapeutic target in glaucoma. *Neural Plast.* 2016, 9364091.
- Rajić, J., Dinić, S., Uskoković, A., Jovanović, J.A., Tolić, A., Đorđević, M., Đorđević, M., Poznanović, G., Mihailović, M., Inic-Kanada, A., Barisani-Asenbauer, T., Grdović, N., Vidaković, M., 2020. DNA methylation of miR-200 clusters promotes epithelial to mesenchymal transition in human conjunctival epithelial cells. *Exp. Eye Res.* 197, 108047.
- Thiery, J.P., Acloque, H., Huang, R.Y.J., Nieto, M.A., 2009. Epithelial-mesenchymal transitions in development and disease. *Cell* 139(5), 871–890.

* Presenting author. E-mail address: baskan.anil@gmail.com (Anil Baskan)

S2.P168 Development of the immunopharmacognostic interventions on potentiating tumor recognition against prostate cancer

Tzu-Ting Su^a, Jun-Wei Lin^a, Shen-Chieh Wang^a, Tai-Yang Chen^a, Ju-I Chana^a, Pei-Wen Hsiaa^{a,b,c,d}, Hui-Ming Chen^{a,b,c,d,*}

^a Agricultural Biotechnology Research Center, Academia Sinica, Taipei, Taiwan

^b Institute of Biotechnology, National Taiwan University, Taipei, Taiwan

^c Institute of Life Sciences, National Defense Medical Center, Taipei, Taiwan

^d The agricultural college of Tunghai University, Taichung, Taiwan

The most challenge for treating prostate cancer (PCa) is the high incident rate (65~75%) of recurrence and/or bone metastasis. Although immunotherapies become a groundbreaking therapy for metastatic PCa, including antigen-specific dendritic cells (DC) vaccines[i.e. sipuleucel-T (Provenge)], the feature of PCa as the “cold” immune landscap still causes less efficacy or even unresponsiveness of immunotherapies. Here, we’ve identified a semi-purified fractionated extract (Phyto-X) with two chemically identified phytocompounds (Phyto-X-A and Phyto-X-B) from the medicinal plant (*Asteraceae*) confer a strong immunostimulatory activity among other phytoextracts in Taiwan endemic plants-based phytolibrary. We observed that Phyto-X-resultant tumor cell lysate (TCL) retarded the mOVA-RM1 tumor growth in conjunction with DC vaccine and T cell adoptive transfer through increasing DC maturation and CTL activity against tumor cells. Interestingly, we found that Phyto-X and its phytocompounds (Phyto-X-A and Phyto-X-B) have differential effects on tumor cells for induction of ATP release and HMGB1 (hallmarkers of immunogenic cell death (ICD)), autophagosome formation and autophagosomes-mediated DC activation. Besides, we found that Phyto-X can directly limit the expansion of tumor-associated bone marrow myeloid cells and M2 macrophage polarization, which could be through inhibition of PI3K-AKT-mTOR pathway. Most importantly, we found Phyto-X can synergize the effect of chemodrug-Docetaxol on induction of ICD, tumor-specific CTL activity and modulate key immune cells in the intratumoral microenvironment. To sum up, this study is to address the potential development of a full-defined phytoextracts (Phyto-X) as an immunopharmacognostic interventions through eliciting tumor recognition, modulating tumor microenvironment, and thus synergizing immunotherapies or/and chemotherapy.

Keywords: immunopharmacognostic interventions, immunogenic cell death, autophagosome, tumor recognition, dendritic cell

* Presenting author. E-mail address: hmchen@gate.sinica.edu.tw (Hui-Ming Chen)

S2.P169 Investigating the anti-inflammatory effects of a TCM formulation used for the prevention of rheumatoid arthritis

Luke Robertson^{a,*}, Janika Welzel^b, Mingshu Zhang^a, Yang Zhihua^c, Gao Kaixin^c, Huang Runyue^d, Wen Zehuai^e, Marina Korotkova^b, Per-Johan Jakobsson^b, Ulf Göransson^a

^a Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden

^b Division of Rheumatology, Department of Medicine Solna & Karolinska University Hospital, Karolinska Institutet, Stockholm, Sweden

^c The Second Clinical Medical College, Guangzhou University of Chinese Medicine, Guangzhou, China

^d Section of Rheumatology and Immunology Research, The Second Affiliated Hospital of Guangzhou University of Chinese Medicine (Guangdong Provincial Hospital of Chinese Medicine), Guangzhou, China

^e Key Unit of Methodology in Clinical Research, Guangzhou University of Chinese Medicine (Guangdong Provincial Hospital of Chinese Medicine), Guangzhou, China

Chinese medicinal formulations have a long tradition of use against rheumatoid arthritis (RA). RA is an autoimmune, inflammatory, and chronic disease that primarily affects the joints of 0.5%–1% of the population (Zeng et al. 2008). These formulations are based on combinations of typically 8–10 plants, which are usually boiled and administered as a decoction or tea (Yeong-Deug et al. 2004). We have developed a natural products isolation methodology that can be employed to isolate potential anti-rheumatic natural products from a Chinese medicinal formulation using an example formulation called “Decoction A”. These strategies involve a suite of biological assays against relevant anti-inflammatory pathways (e.g., Nf- κ B and JAK/STAT pathways (Schwartz et al. 2017) and bioassay-guided purification. The natural products workflow is based on extraction in MeOH:CH₂Cl₂ (1:1), followed by separation on diol-bonded silica into seven fractions of increasing polarity. After activity testing of these fractions in cell reporter assays, actives are selected for subfractionation with a micro-scale HPLC workflow. Dozens of compounds with activity in various cell reporter assays have been identified. In conclusion, we have developed an efficient isolation workflow for the assessment of the anti-inflammatory activities of a TCM formulation. The functionality of the pipeline was validated through the identification of several compounds with previously reported anti-inflammatory activities.

Keywords: Traditional Chinese medicine, screening, anti-inflammatory, arthritis

References

Zeng, Q.Y., Chen, R., Darmawan, J., Xiao, Z.Y., Chen, S.B., Wigley, R., Le Chen, S., Zhang, N.Z., 2008. *Arthritis Res. Ther.* 10, R17.

Yeong-Deug, Y., Il-Moo, C., 2004. *Evid Based Complement. Alternat. Med.* 1, 125–132.

Schwartz, D.M. Kanno, Y., Villarino, A., Ward, M. Gadina, M., O'Shea, J. J., 2017. *Nat. Rev. Drug Discov.* 16, 843–862.

Grkovic, T., Akee, R.K., Thornburg, C.C., Trinh, S.K., Britt, J.R., Harris, M.J., Evans, J.R., Kang, U., Ensel, S., Henrich, C.J., Gustafson, K.R., Schneider, J.P., O'Keefe, B.R., 2020. *ACS Chem. Biol.* 15, 1104–1114.

* Presenting author. E-mail address: luke.robertson@uu.se (Luke Robertson)

S2.P170 Biological potential of endophytes from liverwort *Marchantia polymorpha* L.

Agnieszka Ludwiczuk^{a,*}, Filip Nowaczyński^a, Łukasz Świątek^b, Beata Zimowska^c

^a Department of Pharmacognosy with the Medicinal Plant Garden, Medical University of Lublin, 20-093 Lublin, Poland

^b Department of Virology with Viral Diagnostics Laboratory, Medical University of Lublin, 20-093 Lublin, Poland

^c Department of Plant Protection, University of Life Sciences, 20-069 Lublin, Poland

The mutualistic coexistence between the host and endophyte is diverse and complex, including host growth regulation, and protection from microbial or herbivore attack. The latter is commonly associated with the production by endophytes of bioactive natural products. The endophytes of higher plants have been extensively studied, but there is a dearth of information on the biodiversity of endophytic microorganisms associated with liverworts and, more importantly, their bioactive metabolites (Stelmasiewicz et al., 2023a,b). This presentation will cover isolation of bacterial and fungal endophytes from liverwort *Marchantia polymorpha* L., analysis of the metabolites present in obtained extracts and their fractions, and evaluation of anticancer potential. The extracts were analysed using gas chromatography or liquid chromatography coupled to mass spectrometry (GC/MS and LC/ESI-QTOF-MS). The cytotoxicity and anticancer potential were assessed against noncancerous VERO cells and cancer cells—namely the HeLa, RKO, and FaDu cell lines. The most characteristic metabolites identified in the ethyl acetate extract and fractions were diketopiperazine derivatives. Among volatiles the most abundant were cyclo(L-phenylalanyl-L- prolyl), cyclo(L-leucyl-L-prolyl), and their stereoisomers (Fig. 1).




Fig. 1. Diketopiperazine derivatives identified in *M. polymorpha* endophytes

The endophyte extracts and isolated fractions showed a potential selective anticancer influence on all tested cancer cell lines. These results reinforce the potential of *Marchantia polymorpha* endophytes as a source of biologically active secondary metabolites.

Keywords: Liverwort endophytes, diketopiperazines, anticancer and antiviral activity

References

- Stelmasiewicz, M., Świątek, Ł., Gibbons, S., Ludwiczuk, A., 2023a. Bioactive compounds produced by endophytic microorganisms associated with bryophytes—the “Bryendophytes”. *Molecules* 28, 3246.
- Stelmasiewicz, M., Świątek Ł., Ludwiczuk, A., 2023b. Chemical and biological studies of endophytes isolated from *Marchantia polymorpha*. *Molecules* 28(5), 2202.

* Presenting author. E-mail address: aludwiczuk@pharmacognosy.org (Agnieszka Ludwiczuk)

Natural product research and the challenges of climate change – ecological and environmental solutions

S2.P171 Small molecule metallophores from bacteria, adaptation and isolation strategies

Paul D. Boudreau^{a,*}, Mohammed M. A. Ahmed^{a,b}, Braxton H. Crumpler^a

^a Department of BioMolecular Sciences, University of Mississippi School of Pharmacy, 405 Faser Hall, University, Mississippi, 38677-1848, USA

^b Department of Pharmacognosy, Al-Azhar University, Nasr City, Cairo, 11651, Egypt

Heavy metal pollution is a persistent environmental threat, but many of these elements are also essential to industry (Binnemans, 2013; Mathivanan, 2021; Reck, 2012). Numerous bioremediation- based strategies are being developed to mitigate heavy metal pollution (Mathivanan, 2021; Yaashikaa, 2022). In this presentation, we will present our laboratory's work to isolate metallophores from bacteria and evaluate their interaction with different heavy metals. As part of this work we have isolated cupriachelin siderophores from *Cupriavidus necator* B-4383 (Ahmed, 2003), and delftibactin siderophores from *Delftia lacustris* DSM 21246 (Ahmed and Boudreau, *in review*). To explore the novel chemical space of small molecule metallophores that bind to metals of interest we have also developed an enrichment workflow to isolate novel metallophore-producing strains from metal polluted environments. In these efforts, we relied on dual metal tolerance and Chrome Azurol S (CAS) siderophore production assays. Here we are hypothesizing that siderophores produced by these strains may have dual role in both iron acquisition and heavy metal tolerance, as has been observed before (Johnston, 2013; Johnstone and Nolan 2015). Initial hits from strains isolated from a former mining site in Montana include a *C. necator* strain which produces a putative taiwachelin homolog (Kreutzer and Nett, 2012), which supports that our approach can identify metallophore producing strains. Future aims for this project include identifying the chemistry that underpins the interaction of these metabolites with heavy metals beyond iron and efforts to understand the biological regulation of the biosynthetic pathways for these compounds under metal stress.

Keywords: siderophores, metallophores, metabolomics, bacteria, *Delftia*

References

- Ahmed, M.M.A.; Tripathi, S.K.; Boudreau, P.D. 2023. Comparative metabolomic profiling of *Cupriavidus necator* B- 4383 revealed production of cupriachelin siderophores, one with activity against *Cryptococcus neoformans*. *J. Nat. Prod.* 11, 1256962.
- Ahmed, M.M.A.; Tripathi, S.K.; Boudreau, P.D. 2024. LCMS-Metabolomic Profiling and Genome Mining of *Delftia lacustris* DSM 21246 Revealed Four New Delftibactin Metallophores. *Manuscript under review*.
- Binnemans, K.; Jones, P. T.; Blanpain, B.; Van Gerven, T.; Yang, Y.; Walton, A.; Buchert, M. 2013. Recycling of Rare Earths: A Critical Review. *J. Clean. Prod.* 51, 1–22.
- Johnston, C.W.; Wyatt, M.A.; Li, X.; Ibrahim, A.; Shuster, J.; Southam, G.; Magarvey, N.A. Gold biomineralization by a metallophore from a gold-associated microbe. *Nat. Chem. Biol.* 2013, 9, 241–243.
- Johnstone, T.C.; Nolan, E.M. Beyond iron: non-classical biological functions of bacterial siderophores. *Dalton Trans.* 2015, 44, 6320.
- Kreutzer, M.; Nett, M. Genomics-driven discovery of taiwachelin, a lipopeptide siderophore from *Cupriavidus taiwanensis*. *Org. Biomol. Chem.* 2012, 10, 9338.
- Mathivanan, K.; Chandirika, A.; Vinothkanna, A.; Yin, H.; Liu, X.; Meng, D. Bacterial adaptive strategies to cope with metal toxicity in the contaminated environment – A review. *Ectotoxicological and Environmental Safety*. 2021, 226, 112863.
- Reck, B. K.; Graedel, T. E. Challenges in Metal Recycling. *Science*. 2012, 337, 690–695.
- Yaashikaa, P.R.; Senthil Kumar, P.; Jeevanantham, S.; Saravanan R. A review on bioremediation approach for heavy metal detoxification and accumulation in plants. *Envrion. Poll.* 2022, 301, 119035.

* Presenting author. E-mail address: boudreau@olemiss.edu (Paul D. Boudreau)

S2.P172 Exploring Tuscany onion (*Allium cepa* L.) heritage: unveiling ancient varieties through molecular network analysis and bioinformatic tools

Emily Cioni^{a,*}, Florent Olivon^{b,c}, Fabiano Camangi^d, Maria Paola Germanò^e, Valeria D'Angelo^e, Marc Litaudon^b, Marinella De Leo^a, Alessandra Braca^a, Cécile Apel^b

^a Dipartimento di Farmacia, Università di Pisa, via Bonanno 33, 56126 Pisa, Italy

^b Institut de Chimie des Substances Naturelles, CNRS, UPR2301, University of Paris-Saclay, Avenue de la Terrasse 1, 91198 Gif-sur-Yvette, France

^c Waters SAS, Rue Jacques Monod 5, 78280 Guyancourt, France

^d Scuola Superiore Sant'Anna di Studi Universitari e di Perfezionamento, Piazza Martiri della Libertà 33, 56127 Pisa, Italy

^e Dipartimento di Scienze Chimiche, Biologiche, Farmaceutiche e Ambientali, Università degli Studi di Messina, Viale Ferdinando Stagno d'Alcontres 31, 98166 Messina, Italy

Preserving plant biodiversity and making sustainable use of genetic resources in agriculture is a topical issue with ecological, economic, and social repercussions (Ebert et al., 2020). *Allium cepa* L. (Amaryllidaceae) is worldwide studied as an important source of dietary phytochemicals (Tedesco et al., 2015). In Italy, some onion varieties, only cultivated by local farmers, are neglected and worth to be valorized. The aim of this work was to investigate for the first time the chemical composition of 10 indigenous Tuscany onion varieties, by comparing with the marketed “Tropea”, “Fiorentina”, and “Di Certaldo” onions. The chemical fingerprint was obtained by processing LC-qTOF-MS/MS data with MZmine3 (Schmid et al., 2023), generating molecular networks with MetGem (Olivon et al., 2018) and using compound annotation tools. Flavonoids such as kaempferol, (dihydro)quercetin and isorhamnetin derivatives, were detected in all the samples, thought in different relative ratio. “Di Certaldo” variety distinguished from the others for a molecular cluster of sulphur-containing compounds. “Acquaviva” and “Treschietto” displayed the presence of fatty acids and amino acids derivatives not detected in other samples. Combining the acquisition of large, high-resolution experimental datasets with powerful informatics tools, it was possible to identify the most known bioactive compounds distribution but also highlight the presence of molecular clusters specific to a few onion varieties, potentially correlated with the biological findings. As future work, the extracts will be investigated for their antiangiogenic activity by two *in vivo* models, chick chorioallantoic membrane/zebrafish embryos.

Keywords: *Allium cepa*, biodiversity, bioinformatics, chemical fingerprint, antiangiogenesis

References

- Ebert, A.W., Engels, J.M.M., 2020. Plant biodiversity and genetic resources matter! *Plants* 9, 1706.
- Olivon, F., Elie, N., Grelier, G., Roussi, F., Litaudon, M., Touboul, D., 2018. MetGem software for the generation of molecular networks based on the t-SNE algorithm. *Anal. Chem.* 90, 13900-13908.
- Schmid, R., Heuckeroth, S., Korf, A., Smirnov, A., Myers, O. et al., 2023. Integrative analysis of multimodal mass spectrometry data in MZmine 3. *Nat. Biotechnol.* 41, 447-449.
- Tedesco, I., Carbone, V., Spagnuolo, C., Minasi, P., Russo, G.L., 2015. Identification and quantification of flavonoids from two southern Italian cultivars of *Allium cepa* L., Tropea (Red Onion) and Montoro (Copper Onion), and their capacity to protect human erythrocytes from oxidative stress. *J. Agric. Food Chem.* 63, 5229-5238.

* Presenting author. E-mail address: emily.cioni@phd.unipi.it (Emily Cioni)

S2.P173 Extraction of *Cannabis sativa*: subcritical water vs. supercritical CO₂ and organic solvents

Katja Schoss*, Nina Kočevar Glavač

University of Ljubljana, Faculty of Pharmacy, Department of Pharmaceutical Biology, Aškerčeva 7, 1000 Ljubljana, Slovenia

The use of hemp (*Cannabis sativa*) has attracted great interest due to its versatile applications in the pharmaceutical, food and cosmetic industries. The plant is also increasingly investigated for use in sustainable extraction. By applying the 2³ full factorial DoE approach, we investigated subcritical water extraction (Chakraborty et al. 2021) for obtaining hemp extracts (var. 'Futura 75') as an alternative to supercritical CO₂ extraction (Jokić et al. 2022), ultrasonic extraction (organic solvents, water), Soxhlet and high-pressure ethanol extraction. The extracts differed significantly in terms of composition and antioxidant activity. In the extraction with subcritical water, we confirmed statistically significant models for the yield, the antioxidant activity obtained in the DPPH and ABTS tests, and the content of polyphenolic compounds for the studied factors (temperature in all models, drug to solvent ratio and product temperature* drug to solvent ratio in the yield). The highest yield was obtained with subcritical water extraction (35.91%), and temperature and drug to solvent ratio were the main parameters affecting yield. Soxhlet extraction (36.06% CBG) and high-pressure ethanol extraction (17.39% CBGA) proved to be the most suitable for the extraction of cannabinoids. Subcritical water extraction yielded extracts with a comparably high antioxidant activity and polyphenol content as the extracts obtained with other methods. Temperature was a key parameter and showed a strong positive correlation with antioxidant activity and polyphenol content.

We conclude that extraction with subcritical water is a good choice for obtaining novel extracts from hemp, as the extracts show relatively high antioxidant activity and good yield.

Keywords: antioxidant activity, cannabinoids, *Cannabis sativa*, subcritical water extraction

References

- Chakraborty, Snehasis, Lubna Shaik, and Jyoti S. Gokhale. 2021. Subcritical Water: An Innovative Processing Technology. Pp. 552–66 in *Innovative Food Processing Technologies*. Elsevier.
- Jokić, Stela, Igor Jerković, Valentina Pavić, Krunoslav Aladić, Maja Molnar, Martina Jakovljević Kovač, and Sanda Vladimir-Knežević. 2022. Terpenes and Cannabinoids in Supercritical CO₂ Extracts of Industrial Hemp Inflorescences: Optimization of Extraction, Antiradical and Antibacterial Activity. *Pharmaceuticals* 15(9):1117.

* Presenting author. E-mail address: katja.schoss@ffa.uni-lj.si (Katja Schoss)

S2.P174 Response surface methodology guided RP-HPLC-PDA method optimization for the quantification of multi-analytes (six) in Indian kudzu and its greenness assessment

Mridul Kant Chaudhary*, Ankita Misra, Sharad Srivastava

Pharmacognosy Division, CSIR- National Botanical Research Institute, Lucknow-226001 (Uttar Pradesh), India

In present study, response surface methodology (RSM) guided RP-HPLC-PDA method was established to maximize the bioactive metabolites viz., gallic acid, p-Coumaric acid, daidzein, biochanin A, genistein and puerarin content in Indian kudzu (*Pueraria tuberosa* DC.). Three independent variables i.e. extraction time (A), extraction temperature (B) and liquid to solid ratio (C) and central composite design (CCD) was used. The experimental design yields 20 runs and the optimum conditions for maximum yield of targeted metabolites were 25.16 minute of (A), 32.47 °C of (B) and 20 mL g⁻¹ of (C). The yield of targeted metabolites viz, gallic acid, p-Coumaric acid, daidzein, biochanin A, genistein and puerarin were 0.166 ± 0.025, 0.0234 ± 0.008, 0.0249 ± 0.004, 0.015 ± 0.002, 0.0375 ± 0.006 and 0.476 ± 0.009 % (dry wt.) respectively and were insignificantly different from predicted (RSM model) values i.e., 0.168, 0.024, 0.024, 0.019, 0.036 and 0.465% respectively. The results demonstrate the RSM aided optimization of RP-HPLC conditions for the highest yield of bioactive metabolites in *P. tuberosa*. The developed RP-HPLC method was also validated according to ICH guidelines (2022). The analytical method was linearly calibrated at 48- 240 ng run⁻¹, LOD and LOQ ranged from 7.88-14.89 and 23.88-45.13 ng among the six metabolites. The repeatability and reproducibility of method was, well within the specified limit of <2% and recovery study varies from 97.12-99.00 %. The greenness of analytical method was estimated using analytical eco scale (AES), NEMI, GAPI, AGREE and AGREEprep, and it was found environment friendly and energy efficient (Gałuszka et al., 2012; Keith et al., 2007; Pena-Pereira et al., 2020; Płotka- Wasyłka, 2018).

Keywords: central composite design (CCD), daidzein, *Pueraria tuberosa*, puerarin, response surface methodology (RSM)

References

- Guideline IH. Validation of analytical procedures Q2 (R2). ICH: Geneva, Switzerland. 2022.
- Gałuszka A, Migaszewski ZM, Konieczka P, Namieśnik J. Analytical Eco-Scale for assessing the greenness of analytical procedures. *TrAC Trends in Analytical Chemistry*. 2012 Jul 1;37:61-72.
- Keith LH, Gron LU, Young JL. Green analytical methodologies. *Chemical reviews*. 2007 Jun 13;107(6):2695-708. Pena-Pereira F, Wojnowski W, Tobiszewski M. AGREE—Analytical GREENness metric approach and software. *Analytical chemistry*. 2020 Jun 15;92(14):10076-82.
- Płotka-Wasyłka J. A new tool for the evaluation of the analytical procedure: Green Analytical Procedure Index. *Talanta*. 2018 May 1; 181:204-9.

* Presenting author. E-mail address: mridulkant1995@gmail.com (Mridul Kant Chaudhary)

S2.P175 Development and validation of RP-HPLC method for estimation of anti-oomycete diterpenoids in *Larix decidua* extracts

Linda Langat^{a,*}, Daniel Driscoll^a, Kevin Stephens^b, Moses Langat^c, Barbara Thuerig^d, Hans-Jakob Schärer^d, Lucius Tamm^d, Dulcie Mulholland^a

^a Natural Products Research Group, Department of Chemistry, University of Surrey, Guildford, GU2 7XH, UK

^b BioExtraction (Wales) Limited, Tafarnaubach, Tredegar, NP22 3AA, Wales

^c Science Department, Royal Botanic Gardens Kew, Kew Green, Richmond, TW9 3AE, UK

^d Research Institute of Organic Agriculture (FiBL), Frick, Switzerland

The study involved the development of reverse phase HPLC methods for quantifying two anti-oomycete diterpenoids, larixyl acetate (**1**), larixol (**2**) and the co-occurring inactive, epimanool (**3**) (Fig. 1) from the Larixyne[®] extract obtained from European Larch bark and Larch oleoresin. Larixyne[®] is effective in controlling grapevine downy mildew, *Plasmopara viticola*, that can cause 100% crop loss if left untreated (Mulholland et al., 2017 and Thuerig et al., 2017). Larixyne[®] offers potential to replace ecotoxicologically unfavourable copper fungicides that are reliably used to control downy mildew in vineyards. Currently, the viticulture industry is the fastest growing sector in UK agriculture and is producing award winning wines. This study, Green Laryxine is an Innovate UK funded project aiming to commercialise the use of Larixyne[®]. The reverse phase HPLC, involved chromatographic separation using Ascentis[®] Express F5 column (10 cm × 2.1 mm, 2.7 µm) with acetonitrile and water: 48:52% v/v as a mobile phase at a constant flow rate of about 0.3 mL/min. The development and validation were carried out at detection wavelength of 203 nm. We developed a robust RP-HPLC method, validated for linearity, precision, accuracy, specificity, and system suitability. The method demonstrated excellent linearity with correlation coefficient value r^2 of > 0.99 with linearity range 10-30 µg/mL for larixyl acetate (**1**), larixol (**2**) and epimanool (**3**). The LOD and LOQ were found to be lower, therefore the method is sensitive. The developed RP-HPLC method is an effective method for the quantification of larixyl acetate, larixol and minor constituents in Larch extracts.




Fig. 1. Structures of larixyl acetate (**1**), larixol (**2**) and epimanool (**3**)

Keywords: *Plasmopara viticola*, diterpenoids, Larixyne, RP-HPLC

References

- Mulholland, D.A., Thuerig, B., Langat, M.K., Tamm, L., Nawrot, D.A., James, E.E., Qayyum, M., Shen, D., Ennis, K., Jones, A., Hokkanen, H., Demidova, N., Izotov, D., Schärer, H., 2017. Efficacy of extracts from eight economically important forestry species against grapevine downy mildew (*Plasmopara viticola*) and identification of active constituents. *Crop Protection*. 102,104-109.
- Thuerig, B., James, E.E., Schärer, H., Langat, M.K., Mulholland, D.A., Treutwein, J., Kleeberg, I., Ludwig, M., Jayarajah, P., Giovannini, O., Markellou, E., Tamm, L., 2017. Reducing copper use in the environment: the use of larixol and larixyl acetate to treat downy mildew caused by *Plasmopara viticola* in viticulture. *Pest Management Science*. 74, 477- 488.

* Presenting author. E-mail address: linda.langat@surrey.ac.uk (Linda Langat)

S2.P176 Eutectic solvents as a sustainable solution for the recovery of high value free fatty acids from microalgal co-products

Laura Wils^a, Solène Odou^a, Louise Van Gheluwe^a, Alia Chevalley^b, Leslie Boudesocque-Delaye^{a,*}

^a SIMBA EA 7502 Université de Tours, Faculté de Pharmacie, 31 avenue Monge 37200 Tours, France

^b AquaEcoCulture, Lamballe, France

Arthrospira platensis and *Porphyridium cruentum* are under intense study due to their high content of phycobiliproteins. The increasing demand for phycobiliproteins has led to an urgent need to develop a biorefinery approach to valorize non-polar metabolites of interest, like Free fatty acids (FFA) (Wils et al., 2021). Natural deep eutectic solvents (NaDES) are an environmentally friendly alternative to conventional solvents. Our laboratory has demonstrated the effectiveness of these solvents in extracting FFA from spirulina freeze-dried biomass (Wils et al, 2021). Based on these promising results, we expanded our research to include by-products of the aqueous extraction of phycobiliproteins (Fig. 1).




Fig. 1. General biorefinery approach for greener valorisation of microalgae cake

Wils et al. (2022) compared the extraction performances of a panel of non-polar NaDES. NaDES based on fatty acids were found to be the most promising for recovering FFAs from microalgae, with high selectivity towards saturated FFAs. The presence of polyols within the NaDES modulates the selectivity in favor of the polyunsaturated FFA. Microalgae cake has been shown to be very effective for FFA recovery. These findings provide a basis for designing a multi-stage microalgae biorefinery scheme using NaDES or sustainable solvents for the cosmetic or food industry.

Keywords: eutectic solvent, microalgae, lipids, green process

References

- Wils L., Leman-Loubière C., Bellin N., Clément-Larosièrè B., Pinault M., Chevalier S., Enguehard-Gueiffier C., Bodet C., Boudesocque-Delaye L. 2021. Natural Deep Eutectic Solvent formulations for spirulina: preparation, intensification, and skin impact. *Algal Research Special Issue on Advanced Technologies* 56, 102317
- Wils, L.; Yagmur, M.; Phelippe, M.; Montigny, B.; Clément-Larosièrè, B.; Jacquemin, J.; Boudesocque-Delaye, L. 2022. Alternative Solvents for the Biorefinery of Spirulina: Impact of Pretreatment on Free Fatty Acids with High Added Value. *Mar. Drugs* 20, 600

* Presenting author. E-mail address: leslie.boudesocque@univ-tours.fr (Leslie Boudesocque-Delaye)

S2.P177 EKO-YAM - investigating *Dioscorea* for a sustainable future

David Krüger*, Alexander Weng

Institute of Pharmacy, Department of Pharmaceutical Biology, Freie Universität Berlin, Königin- Luise-Straße 2+4, 14195 Berlin, Germany

The recently launched EKO-YAM project is presented. It focuses on the Chinese yam *Dioscorea polystachya*. Its tuber is used in Traditional Chinese Medicine given the potential against diabetes and cardiovascular diseases. The tuber is also an important foodstuff due to its nutritional value. That is why cultivation of yam is of interest. However, cultivation becomes increasingly labor-intensive, costly, and thus challenging. Therefore, this project aims at innovative cultivation methods (i.e., mobile raised beds, trench cultivation, mound beds, gutter cultivation). They should allow economical and sustainable production of Chinese yam and maintain or even improve the nutritional quality. These methods are intended to align with the organic regulations in Europe. The different cultivars are profiled regarding the pattern of proteins (using MALDI-TOF MS) as well as primary and secondary metabolites (HPLC-MS). The latter are mainly adduced as molecular markers to monitor breeding experiments and to secure quality assurance. In addition, conventional crossing trials are performed to obtain stronger varieties, e.g., with increased health-promoting effects. Generally, foods with such properties may have contributions to nourishment or health care of society. The EKO-YAM project purposes to promote the establishment of *Dioscorea polystachya* as a foodstuff and possibly as a phyto-pharmaceutical raw material, thereby contributing to the National Bioeconomy Strategy for a sustainable future.

Keywords: *Dioscorea polystachya*, Dioscoreaceae, cultivation, method development, secondary metabolites

References

- Amat, N. et al., 2014. Aqueous extract of *dioscorea opposita thunb.* normalizes the hypertension in 2K1C hypertensive rats. *BMC Complement Altern Med.* 14, 36.
- Epping, J. et al., 2020. An underutilized orphan tuber crop - Chinese yam: A Review. *Planta.* 252, 58.

* Presenting author. E-mail address: d.krueger@fu-berlin.de (David Krüger)

S2.P178 Wound healing and antimicrobial activity of European tree bark extracts

Sabrina Vorderegger^{a,*}, Johanna Rehrl^b, Claudia Hölzl^a, David Sesink^a, Gertie Janneke Oostingh^a, Bettina Stelzhammer^a, Anja Schuster^a

^a Salzburg University of Applied Sciences, Health Sciences Department, Urstein Sued 1, 5412 Puch/Salzburg, Austria

^b Department of Biosciences and Medical Biology, Paris-Lodron University of Salzburg, Hellbrunner Straße 34, 5020 Salzburg, Austria

Various types of wood bark have been identified to possess antimicrobial, antiviral, anti-inflammatory, and wound-healing properties (Emrich et al., 2022). This poster will present the research on wood bark extracts from *Fagus sylvatica*, *Prunus padus*, *Quercus robur*, *Pinus sylvestris*, *Alnus glutinosa* and *Betula pendula* in view of their wound healing and antimicrobial properties. The extracts were retrieved using the Soxhlet extraction method. The minimum inhibitory concentration (MIC) assay was used to determine the effectiveness of antimicrobial agents on *Cutibacterium acnes*, *Staphylococcus epidermidis*, MRSA-*Staphylococcus aureus* and *Escherichia coli*, *K. pneumoniae*, *P. aeruginosa*. To determine the wound healing capacity a scratch assay was performed on a human keratinocyte cell line. Wound closure was measured up to 72 hours in the presence or absence of bark extracts. All bark extracts, except *Prunus padus* show varying degree of bacterial growth inhibition (MIC range from 67 µg/mL to 1073 µg/mL) against gram positive bacteria whereas solely *Alnus glutinosa* demonstrated a partial inhibition against gram negative bacteria at a concentration of 1073 µg/mL. In terms of the wound healing capacity, all bark extracts lead to a significantly accelerated wound closure (timepoint and concentration dependent) compared to untreated cells (e.g. 100% wound closure at 38 hours with *Prunus padus* compared to 72 hours without extract). Their combined positive effect on wound-healing and antimicrobial activity has great potential for the treatment of various skin diseases (Häsler Gunnarsdottir et al., 2023). In addition, the sustainability element is tackled by utilizing bark, considered a byproduct in the forest industry.

Keywords: bark extracts, antimicrobial activity, wound healing capacity, sustainability

References

Emrich, S., Schuster, A., Schnabel, T., Oostingh, G.J., 2022. Antimicrobial Activity and Wound-Healing Capacity of Birch, Beech and Larch Bark Extracts. *Molecules* 27, 2817.

Häsler Gunnarsdottir, S., Sommerauer, L., Schnabel, T., Oostingh, G.J., Schuster, A., 2023. Antioxidative and Antimicrobial Evaluation of Bark Extracts from Common European Trees in Light of Dermal Applications. *Antibiotics* 12,130.

* Presenting author. E-mail address: sabrina.vorderegger@fh-salzburg.ac.at (Sabrina Vorderegger)

S2.P179 An ethnobotanical survey investigating traditional remedies to treat infectious diseases in French Polynesia

Doriane Tittou^{a,b,*}, Jean-François Butaud^c, Raimana Ho^b, Phila Raharivelomanana^b, François Chassagne^a

^a UMR 152 PharmaDev, Université Paul Sabatier, Institut de Recherche pour le Développement (IRD), Toulouse, France

^b UMR 214 SECOPOL, Université de Polynésie Française, IFREMER, ILM, IRD, Faaa, Tahiti, Polynésie française

^c Correspondant du Muséum National d'Histoire Naturelle (PatriNat), Paris & Consultant en foresterie et botanique polynésienne, Tahiti, Polynésie française

In French Polynesia, traditional medicine is still used and considered as a part of the Polynesian culture (Chassagne *et al.*, 2022, 2023; Girardi *et al.*, 2015). Anti-infectious remedies have been reported to treat several disorders, for all types of patients and in many islands, making them an important part of traditional Polynesian medicine. In this context, an ethnobotanical survey was implemented to identify remedies used for infectious diseases, inventory the plants, better understand the local representation of these therapies and provide efficacy and safety data on these uses. A field study was carried out from February to April 2024, on five islands from the Society archipelago (Tahiti, Moorea, Raiatea, Tahaa and Huahine) in French Polynesia. A declaration of research regarding access to biological resources and associated traditional knowledge was recorded according to the local regulations. A total of 83 persons was interviewed including experts in herbalism. This fieldwork led to the identification of 92 plants species for the treatment of 11 infectious diseases (e.g., sinusitis, abscess, urogenital infections). The majority of plants used are native or Polynesian introduced plants. Among them, *Cordia subcordata*, *Rorippa sarmentosa*, *Heliotropium arboreum*, *Coleus barbatus* and *Spondias dulcis* were the most cited and will be subjected to further biological and phytochemical analysis. This study is part of a larger project named “Sustainable Traditional Medicine in French Polynesia”, funded by the French National Research Agency (ANR), and aiming to preserve, support, and valorise the traditional medical knowledge from French Polynesia.

Keywords: Pacific, traditional medicine, ethnobotany, medicinal plants, infectious diseases

References

- Chassagne, F., Butaud, J. F., Ho, R., Conte, E., Hnawia, É., Raharivelomanana, P., 2023. Traditional medical practices for children in five islands from the Society archipelago (French Polynesia). *Journal of Ethnobiology and Ethnomedicine*. 19(1), 44.
- Chassagne, F., Butaud, J. F., Torrente, F., Conte, E., Ho, R., Raharivelomanana, P., 2022. Polynesian medicine used to treat diarrhea and ciguatera: An ethnobotanical survey in six islands from French Polynesia. *Journal of Ethnopharmacology*. 292, 115186.
- Girardi, C., Butaud, J. F., Ollier, C., Ingert, N., Weniger, B., Raharivelomanana, P., Moretti, C., 2015. Herbal medicine in the Marquesas Islands. *Journal of Ethnopharmacology*. 161, 200–213.
- Quenon, C., Hennebelle, T., Butaud, J. F., Ho, R., Raharivelomanana, P., Roumy, V., 2022. Antimicrobial Properties of Compounds Isolated from *Syzygium malaccense* (L.) Merr. and L.M. Perry and Medicinal Plants Used in French Polynesia. *Life* (Basel, Switzerland). 12(5), 733.

* Presenting author. E-mail address: doriane.tittou@ird.fr (Doriane Tittou)

S2.P180 Water as a green solvent to extract polar and non-polar plant pigments

Marine Chambaud^{a,b,*}, Rémi Guyomarch^b, Emilie Destandau^a

^a Institut de Chimie Organique et Analytique, University of Orléans, Orléans, France

^b Terre de Couleur, 6 rue de Châtenay, Rochecorbon, France

Water is a green solvent, harmless to human health and the environment, and commonly used in food and cosmetic products (Chemat et al., 2019). Thus, water is a great solvent choice to obtain plant extracts that could be directly used as ingredients. Plant pigments are a heteroclite group of secondary metabolites reunited by their capacity to absorb light in the visible domain. They often display interesting activities, such as antioxidant or photoprotective (Brudzyńska et al., 2021). However, some have low polarity, making them challenging to extract using water. This work focused on designing efficient methods to extract pigments and active compounds with different polarities using water. For anthraquinones, medium-polarity compounds from madder (*Rubia tinctorum* L.), a pressurised water-based microwave-assisted extraction (PMAE) method was optimised. Microwaves generate much heat in less than a minute via ionic conduction and dipole rotation in the plant matrix and the solvent. Combined with the pressure, these conditions lower the dielectric constant of water and its polarity, enabling the eco-extraction of anthraquinones. For the carotenoids from achiote (*Bixa orellana* L.), which have low polarity, a bio-based co-solvent was added to develop a biphasic oil/water ultrasound-assisted extraction. Combining both solvents allowed the simultaneous extraction of polar and non-polar compounds. The antioxidant activity and pigment content of extracts from both plants were assessed using colorimetric assays and the extract characterisation was completed by UHPLC-HRMS/MS. Water-based extraction methods showed at least similar efficiency as conventional methods with organic solvents. The recovered extracts could be directly formulated into cosmetic products.

Keywords: Water, green extraction, cosmetics, plant pigments, antioxidants

References

- Brudzyńska, P., Sionkowska A., Grisel M., 2021. Plant-Derived Colorants for Food, Cosmetic and Textile Industries: A Review. *Materials*, 14, 3484-3502
- Chemat, F., Abert Vian, M. A., Ravi, H. K., Khadhraoui, B., Hilali, S., Perino, S., Fabiano Tixier, S., 2019. Review of Alternative Solvents for Green Extraction of Food and Natural Products: Panorama, Principles, Applications and Prospects. *Molecules*, 24, 3007-3034.

* Presenting author. E-mail address: marine.chambaud@univ-orleans.fr (Marine Chambaud)

S2.P181 Long term experiments on applying a pulsed electric field (PEF) to living plant roots in order to extract flavonoids in *S. baicalensis* Georgi model

Kajetan Grzelka^{a,*}, Joanna Jaśpińska^b, Adam Matkowski^{a, b}, Sylwester Ślusarczyk^a

^a Department of Pharmaceutical Biology and Biotechnology, Division of Pharmaceutical Biology and Botany, Wrocław Medical University, wyb. Ludwika Pasteura 1, 50-367 Wrocław, Poland

^b Botanical Garden of Medicinal Plants, Wrocław Medical University, al. Jana Kochanowskiego 10, 51-610 Wrocław, Poland

Scutellaria baicalensis Georgi (Lamiaceae) has been used for centuries in Traditional Chinese Medicine and is nowadays listed in Chinese, British and European Pharmacopoeias (Wang *et al.*, 2018). It exhibits antioxidant, cytotoxic, anxiolytic and antimicrobial properties. Among many flavonoids produced in its roots, baicalin (1), wogonoside (2), baicalein (3) and wogonin (4) (Fig. 1) are the most abundant (Zhao *et al.*, 2019).




Fig. 2. Main flavonoids found in *S. baicalensis* root: baicalin (1), wogonoside (2), baicalein (3) and wogonin (4).

This research covers using Pulsed Electric Fields (PEF) to facilitate extraction of flavonoid compounds from living roots of aeroponically cultivated *S. baicalensis* plant model. Most plants survive this treatment, sometimes reaching size and content of root phytochemicals larger than control (Grzelka *et al.*, 2023). Presented work is a result of a two-year-long process of empirically adjusting PEF parameters and composition of Natural Eutectic Solvent (NES) mixture in order to optimize extraction yield. We found that the phytochemical content of extracts acquired in this refined process is quality-wise synonymous with that of control (80% MeOH extract from dried root material). The best extract was obtained using following PEF parameters: 33 fifty-microsecond-long impulses, 1 Hz frequency, 0.5 kV/cm field strength, delivered twice (5 min interval) and a NES mixture of choline chloride : xylose (1:2 + 30% water). It yielded the same amount of wogonin (4) as control (2.15 µg/mL), but with 60% survival rate of plants after 3 weeks of observation. This allows to extract from the same population again, which is much more time- and cost-efficient than cultivating a new one and can be applied to other plant models.

Funding statement: This research is funded by WMU grant to Kajetan Grzelka no. SUBK.D030.24.011, supported with the funds of the National Science Center, Sonata grant to Sylwester Ślusarczyk - UMO-2020/39/D/NZ9/01402.

Keywords: electroporation, extraction, *Scutellaria baicalensis*, NES, flavonoids

References

- Wang, Z. L., Wang, S., Kuang, Y., Hu, Z. M., Qiao, X., and Ye, M., 2018. A comprehensive review on phytochemistry, pharmacology, and flavonoid biosynthesis of *Scutellaria baicalensis*. *Pharm. Biol.* 56, 465–484.
- Zhao, T., Tang, H., Xie, L., Zheng, Y., Ma, Z., Sun, Q., et al., 2019. *Scutellaria baicalensis* georgi. (Lamiaceae): a review of its traditional uses, botany, phytochemistry, pharmacology and toxicology. *J. Pharm. Pharmacol.* 71, 1353–1369.
- Grzelka K., Matkowski A. and Ślusarczyk S., 2023. Electrostimulation improves plant growth and modulates the flavonoid profile in aeroponic culture of *Scutellaria baicalensis* Georgi. *Front. Plant Sci.* 14:1142624.

* Presenting author. E-mail address: kajetan.grzelka@student.umw.edu.pl (Kajetan Grzelka)

S2.P182 Enhancing fire ant bait selectivity: Synthesis and evaluation of feeding deterrent activity of fire ant venom alkaloids and their analogs against invasive fire ants and native ants

H. M. T. Bandara Herath^{a*}, Jian Chen^b, N. P. Dhammika Nanayakkara^a, Ikhlas A. Khan^a

^a National Center for Natural Product Research, School of Pharmacy, University of Mississippi, University, MS 38677, USA

^b USDA-ARS, Biological Control of Pests Research Unit, Stoneville, MS 38776, USA

The red imported fire ant, *Solenopsis invicta*, has significantly impacted public health, agriculture, and ecosystems in the southern United States. Their painful stings pose a threat to humans, pets, livestock, and wildlife, causing allergic reactions and even fatalities in severe cases. These invasive ants aggressively compete with native species for food and territory, often displacing them and disrupting delicate ecological balances. Using bait is one of the most effective methods for controlling fire ants. Baits consist of insecticides mixed with attractive food substances, which ants take back to their colonies, effectively killing the entire colony. The low selectivity of fire ant bait has been the major drawback of this method. Prompted by the observation that fire ants feed their nestmates with their venom (Chen and Du, 2022), we evaluated the feeding deterrent activity of solenopsin A, a major constituent in the fire ant venom, against fire ants and two common native ants, little black ant (*Monomorium minimum*) and thief ant (*Solenopsis molesta*). Solenopsin A had a potent feeding deterrent activity against little black and thief ants but had no effect on fire ants. The addition of fire ant venom alkaloids and their analogs to bait represents a promising strategy for enhancing the selectivity of bait targeting this invasive pest

We prepared the fire ant venom alkaloids (1-6) and their analogs (7-12) and evaluated them for the feeding deterrent activity against little black, thief, and fire ants. Their activity and the structure- activity relationships will be presented

Keywords: red fire ant, *Solenopsis invicta*, venom alkaloids, feeding deterrent activity, bait

References

Chen, J., Du, Y., 2022. Fire ants feed their nestmates with their own venom. *J. Insect Physiol.* 142, 104437,

*Presenting author, E-mail address: bherath@olemiss.edu (H. M. T. Bandara Herath)

S2.P183 Sucrosephenylpropanoid esters and isoflavonoids from *Belamcanda chinensis* (Iridaceae) roots and their potential anti-osteoclastogenic activity

Manh Tuan Ha^a, Jeong Ah Kim^b, Byung Sun Min^{a,*}

^a College of Pharmacy, Drug Research and Development Center, Daegu Catholic University, Gyeongbuk 38430, Republic of Korea

^b College of Pharmacy, Research Institute of Pharmaceutical Sciences, Kyungpook National University, Daegu 41566, Republic of Korea

This presentation will cover the isolation, structure elucidation and biological evaluation of six new sucrosephenylpropanoid esters (**1–6**) and twenty-one known compounds (**7–27**) from the methanol extract of *Belamcanda chinensis* (L.) DC root (Iridaceae). *B. chinensis* was incorporated into the genus *Iris* as *Iris domestica* (L.) Goldblatt & Mabberley by APG (Missouri Botanical Garden) III (Wozniak et al., 2015). In some countries such as Korea, China, Japan, and Vietnam, this herb is used in traditional medicines to treat pharyngitis, coughing, bronchitis, and chronic tracheitis; its mode of action is by reducing pharyngeal swelling, heat-clearing, and detoxifying (Li et al., 2019). Repeated chromatography of the CH₂Cl₂ and EtOAc soluble fractions from the methanol extract of *Belamcanda chinensis* root yielded six new sucrosephenylpropanoid esters (**1–6**) and twenty-one known compounds (**7–27**). The structures of **1–6** were elucidated using diverse nuclear magnetic resonance (NMR) techniques and high-resolution mass spectrometry (HRMS) data analysis, together with chemical methods (Fig. 1).




Fig. 1. Structure of sucrosephenylpropanoid esters **1–6**

All the twenty-seven isolated compounds were evaluated for their anti-osteoclastogenic activities. Preliminary screening results revealed that compounds **1** and **19** exhibited strong effects against RANKL-induced osteoclast formation in RAW264.7 cells. In addition, the treatment of mouse bone marrow macrophages (BMMs) with compounds **1** and **19** significantly decreased RANKL-induced TRAP-positive multinucleated osteoclast formation in a concentration-dependent manner without affecting cell viability. Further bioassay investigation showed that compounds **1** and **19** inhibited the expression of some osteoclast-specific marker genes and the transcription factor nuclear factor of activated T cells cytoplasmic 1 (NFATc1) in response to RANKL. To the best of our knowledge, this is the first investigation of anti-osteoclastogenic activity for compounds isolated from *B. chinensis*.

Keywords: *Belamcanda chinensis*, Iridaceae, sucrosephenylpropanoid esters, osteoclastogenesis, RANKL, NFATc1

References

- Wozniak, D. Matkowski, A., 2015. *Belamcandae chinensis* rhizoma—a review of phytochemistry and bioactivity, Fitoterapia 107, 1–14.
- Li, J. Ni, G. Liu, Y. Mai, Z. Wang, R. Yu, D., 2019. Iridal-Type Triterpenoids with a Cyclopentane Unit from the Rhizomes of *Belamcanda chinensis*, J. Nat. Prod. 82, 1759–1767.

* Presenting author. E-mail address: bsmin@cu.ac.kr (Byung Sun Min)

S2.P184 PTP1B inhibitory activity of serratene-type triterpenoids from *Lycopodium serratum* (Lycopodiaceae)

Manh Tuan Ha^a, Thi Thanh Le^a, Jeong Ah Kim^b, Byung Sun Min^{a,*}

^a Drug Research and Development Center, College of Pharmacy, Daegu Catholic University, Gyeongbuk 38430, Republic of Korea

^b College of Pharmacy, Kyungpook National University, Daegu 41566, Republic of Korea

This presentation will cover the isolation, structure elucidation and biological evaluation of two new serratene-type triterpenoids (**1** and **2**) and nine analogs [seven serratenes (**3–5**, **7–10**) and two serratanes (**6** and **11**)], together with one *Lycopodium* alkaloid (huperzine A, **12**) were isolated and characterized from *Lycopodium serratum* (Lycopodiaceae) (Fig. 1) (Ha et al., 2023). *Lycopodium* plants were reported to show interesting biological activities, such as cytotoxicity, cancer chemo-preventive activity, and the activities involving the treatment of AD such as acetylcholinesterase and butyrylcholinesterase inhibitory activities (Nguyen et al., 2017).




Fig. 1. Structure of compounds 1-12

The structures of isolates from *L. serratum* were elucidated by the HRESI-MS, 1D and 2D NMR analysis, and comparison with NMR data in the literature. Among the isolates, triterpenoids (**4–6**) showed significant inhibitory effects on PTP1B (protein tyrosine phosphatase 1B) with IC₅₀ values of 13.50, 17.65, and 16.64 μM, in comparison with that of the positive control, ursolic acid (IC₅₀ = 5.98 μM) (Ha et al., 2023). While compounds **1**, **7**, and **8** exhibited moderate PTP1B inhibitory effects with IC₅₀ values ranging from 25.30 to 51.56 μM. Enzyme kinetic studies revealed that all these most active compounds (**4–6**) exhibited the same competitive PTP1B inhibition type with K_i values of 7.9, 16.9, and 8.9 μM, respectively. In addition, the probable binding conformation of **4–6** within the PTP1B active site was studied through *in silico* molecular docking. The docking results demonstrated that these active triterpenoids could be docked stably into the catalytic site of the PTP1B enzyme with negative binding energies.

Keywords: *Lycopodium serratum*, Lycopodiaceae, serratene triterpenoids, PTP1B, molecular docking

References

- Ha, M.T., Le, T.T., Nguyen, V.T., Kim, J.A., Choi, J.S., Min, B.S., 2023. Isolation and characterization of PTP1B inhibitory serratene-type triterpenoids from *Lycopodium serratum* Thunb.. *Phytochemistry Letters* 57, 231-238.
- Nguyen, V.T., Zhao, B.T., Seong, S.H., Kim, J.A., Woo, M.H., Choi, J.S., Min, B.S., 2017. Inhibitory effects of serratene-type triterpenoids from *Lycopodium complanatum* on cholinesterases and β-secretase 1. *Chem. Biol. Interact.* 274, 150- 157.

* Presenting author. E-mail address: bsmin@cu.ac.kr (Byung Sun Min)

S2.P185 Introducing NaDES-based biphasic solvent families inspired by the Arizona family: a new approach for the sustainable use of centrifugal partition chromatography

Ramla Sahli^{a,*}, Luke Froment^a, Tsvetelina Mandova^b, Elea Budzy^a, Leslie Boudesocque-Delaye^a

^a EA 7502 Synthèse et Isolement de Molécules BioActives (SIMBA), Université de Tours, 31 Avenue Monge, 37200 Tours, France

^b Gilson Purification SAS, 22 rue Bourseul, 56890 Saint Avé, France

Centrifugal Partition Chromatography (CPC) has a great potential for developing new eco-friendly separation strategies. Being a support-free technique, solvent selection is crucial for CPC. Typically, natural compounds are separated using pre-established families of solvent systems that offer a wide range of polarity. Unfortunately, these families are still dominated by harmful organic solvents. The Arizona family, which is commonly used in separating natural compounds, contain toxic alkanes and alcohols. There are alternatives to these solvents, including Natural Deep Eutectic Solvents (NaDES). However, tailoring NaDES-based solvent systems can be a challenge. In this work, we introduce a new approach to build solvent systems by incorporating hydrophobic NaDES into the ARIZONA family. NaDES with low viscosity, such as terpene-based NaDES, are the most suitable for CPC (Fan et al., 2021). However, their number is limited. Hence, we conducted an extensive screening of terpene-based NaDES to compile a pool of potential candidates, that could be introduced into biphasic solvent systems. Two NaDES were selected as starting points for the creation of two novel families, NADrizona 1 & 2, consisting of 23 biphasic solvent systems following the Arizona family model. The performance of these families was evaluated using various parameters, including volume ratios, settling times, stationary phase retention ratio (Sf), and polarity range predicted by the GUESS method (Friesen & Pauli, 2005). The GUESSmix was also used to evaluate separation capacities within CPC. Plant extract separation assays were used to validate the separation capacities of the NADrizona 1 & 2 families.

Keywords: CPC, green solvents, NaDES, arizona solvent systems

References

- Friesen J. B., Pauli G. F., 2005. G.U.E.S.S.—A Generally Useful Estimate of Solvent Systems for CCC. *J. Liq. Chromatogr. Relat.* 28(17), 2777-2806.
- Fan C., Sebbah T., Liu Y., Cao X., 2021. Terpenoid-capric acid based natural deep eutectic solvent: Insight into the nature of low viscosity. *Clean. Eng. Technol.* 3, 100116.

* Presenting author. E-mail address: ramla.sahli@univ-tours.fr (Ramla Sahli)

S2.P186 Beyond the tuber: crafting high-value molecules from potato residues

Nina Stadler^{a,*}, Marília Bueno da Silva^{b,c}, Anika Wiese-Klinkenberg^b, Sylvia Schleker^c, Florian M. W. Grundler^c, Björn Usadel^{c,d}, Franziska Genzel^b, Thomas Classen^e, Jörg Pietruszka^{a,e}

^a Institute of Bioorganic Chemistry (IBOC), Heinrich-Heine-University Düsseldorf, Germany

^b Institute of Bio- and Geosciences (IBG-4: Bioinformatics), Forschungszentrum Jülich, Germany

^c Institute of Crop Science and Resource Conservation (INRES), Molecular Phytomedicine (MPM), The Rhenish Friedrich Wilhelm University of Bonn, Germany

^d Institute of Biological Data Science, Faculty of Mathematics and Natural Sciences, Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich-Heine-University Düsseldorf, Germany

^e Institute of Bio- and Geosciences (IBG-1: Bioorganic Chemistry), Forschungszentrum Jülich, Germany

Steroidal glykoalkaloids (SGA) are prominent secondary metabolites in potatoes (*Solanum tuberosum* L.) and potato side-streams, like leaves, berries, and flowers (Roosen-Runge *et al.*, 1977). As potato tubers are the fourth most important food globally, the accumulation of these side products is significant (Zaheer *et al.*, 2016). The two major SGAs are α -solanine and α -chaconine, which have biological activity against different types of herbivores, act as natural pesticides and can therefore be used as a green alternative to traditional plant protection (Allen *et al.*, 1971). The aim of this project is to extract of the side-streams from potato production to obtain these compounds and generate new plant-based pesticides by a semi-synthetic (chemical and enzymatic) approach. The focus will be on the aglycon solanidine **1**, which can be obtained by cleaving the sugars (Fig. 1). A simple test- substrate strategy with cholesterol will be performed as a first proof of principle, since both compounds share a similar scaffold. Successful synthetic routes will then be adapted to synthesize derivatives of solanidine. Additionally, the total synthetic approach for solanidine will be performed (Zhang *et al.*, 2016) to compare both the synthetic and isolation-based route by means of economic and sustainability analyses.




Fig. 1: Structural motif of the aglycon solanidine.

Keywords: steroidal glykoalkaloids, plant-based pesticides, circular bioeconomy, *Solanum tuberosum* L., semi-synthesis

References

- Allen, E.H., Feldmesser, J., 1971. Nematicidal Activity of α -Chaconine: Effect of Hydrogen-ion Concentration. *J. Nematol.*, 3, 58-61.
- Roosen-Runge, C., Schneider, E., 1977. Zur Bestimmung der Solanum-Alkaloide Solanin und Chaconin, *Z. Lebensm. Unters.-Forsch.* 164, 96-97.
- Zaheer, K., Akhtar, M.H., 2016, Potato Production, Usage, and Nutrition – A Review. *Crit. Rev. Food Sci. Nutr.* 56, 711- 721.
- Zhang, Z.D., Shi, Y, Wu, J.J., Lin, J.R., Tian, W.S., 2016. Synthesis of Demissidine and Solanidine. *Org. Lett.*, 18, 3038- 3040.

* Presenting author: E-mail address: n.stadler@fz-juelich.de (Nina Stadler)

S2.P187 Mulberry cell cultures: a promising source for health care

Vanessa Dalla Costa*, Anna Piovan, Raffaella Filippini

Department of Pharmaceutical and Pharmacological Sciences, University of Padova, via Marzolo, 5, 35131 Padova, Italy

Plants have long been exploited as a sustainable source of food and chemicals; however, with the increasing global world population and the climate change ending in losses of biodiversity and land, conventional agriculture unlikely can cope with the huge challenges head. These factors could be countered using new and ethically more justifiable technologies such as plant cellular agriculture, a biotechnological tool for the production of nutritious and healthy ingredients, as well as single bioactive compounds (Rischer et al., 2020; Gubser et al., 2021; Bapat et al., 2023). Given the importance of *Morus alba* L. in many traditional medicines due to its health benefits (Chen et al., 2021), the aim of this work was to establish *M. alba in vitro* cell cultures to study their phytochemistry and potential use for nutritional and healthy supplements. Different organs from *in vitro* grown plantlets were used as explant source, and different basal media were tested. The best callogenesis was obtained from stem explants cultured on MS basal medium supplemented with sucrose and 2,4-D. During the following subcultures, the calli resulted in a brownish, juicy, grainy biomass (Fig. 1).



Fig. 1. Calli of *Morus alba*

Considering that *in vitro* cell metabolism can change during the growth cycle, the phytochemical profile (HPLC-DAD), total phenol and flavonoid content and antioxidant activity (colorimetric assays) were analyzed on the juices obtained by squeezing the biomasses at: initial, mid and stationary phase of the growth cycle. The high content of phenols and flavonoids, as well as the antioxidant activity, showed an active metabolism of the culture.

Keywords: *Morus alba*, cell cultures, phenols, flavonoids, antioxidant

References

- Bapat, V.A, Kavi Kishor, P.B., Jalaja, N., Jain, S.M., Penna, S., 2023. Plant cell cultures: biofactories for the production of bioactive compounds. *Agronomy* 13, 858-877.
- Gubser, G., Vollenweider, S., Eibl, D., Eibl, R., 2021. Food ingredients and food made with plant cell and tissue cultures: state-of-the art and future trends. *Eng. Life Sci.* 21, 87-98.
- Rischer, H., Szilvay, G.R., Oksman-Caldentey, K-M., 2020. Cellular agriculture – industrial biotechnology for food and materials. *Curr. Opin. Biotechnol.* 61, 128-134.

* Presenting author. E-mail address: vanessa.dallacosta@phd.unipd.it (Vanessa Dalla Costa)

S2.P188 Spatio temporal dynamics of favourable habitats for *Terminalia leiocarpa*, a traditional plant used in ethnoveterinary medicine in Benin (West Africa)

Fabrice Jean Eudes Crédo Gahounga^{a,*}, Esaïe Tchétan^b, Sèdami Igor Armand Yevide^a, Thierry Dèhouégnon Houéhanou^c, Pascal Abiodoun Olounlade^b, Joëlle Quetin-Leclercq^d, Adandé Berlamain Fandohan^a

^a Unité de Recherche en Foresterie et Conservation des Bioressources, Laboratoire de Sciences Végétales, Horticoles et Forestières, Ecole de Foresterie Tropicale, Université Nationale d'Agriculture, BP 43, Kétou, Bénin

^b Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomic Sciences, University of Abomey-Calavi, Cotonou 01 BP 526, Benin

^c Laboratory of Ecology, Botany and Plant Biology, Faculty of Agronomy, University of Parakou, Parakou, Benin

^d Pharmacognosy Research Group, Louvain Drug Research Institute, Université catholique de Louvain (UCLouvain), Avenue E. Mounier, B1.72.03, B-1200 Brussels, Belgium

This presentation will focus on *Terminalia leiocarpa* (DC.) Baill (Combretaceae), commonly known as *Anogeissus leiocarpa*. It is a plant widely used in West Africa by livestock farming in ethnoveterinary medicine (Tchétan et al., 2022). The harvesting of its various organs, combined with the effects of climate change are leading to a decline in its natural populations (Houéhanou et al., 2013). In order to contribute to the sustainable use of the plant, we analyzed the dynamics of *Terminalia leiocarpa*'s favourable habitats in Benin. Occurrence data were collected in the three climatic regions of Benin and combined with climatic and soil data and subjected to the Maxent ecological niche modelling algorithm to assess the effect of future environmental conditions (RCP 4.5 and RCP 8.5 scenarios to 2055) on the potential distribution of the species in Benin. The results show that the areas currently favourable and very favourable for the presence of the species extend between the three climatic regions, but are much more localised in the Sudano-Guinean region. According to the climate projections, the current favourable areas will increase by 13.16% for RCP 4.5 and decrease by 9% for RCP 8.5. The currently very favourable areas will see a decrease of 9.23% for RCP 4.5 and an increase of 52.88% for RCP 8.5. Taken together, the currently favourable and very favourable areas will increase for both scenarios in all three climate regions. These results offer hope of saving this important species, whose natural populations have been reported to be declining.

Keywords: climate change, ecological niche modelling, sustainable management, *Terminalia leiocarpa*, Benin

References

- Houéhanou, T. D., Glèlè Kakai R. L., Assogbadjo A. E., Kindomihou V., Houinato M., Wittig R., Sinsin B. A., 2013. Change in the woody floristic composition, diversity and structure from protected to unprotected savannahs in Pendjari Biosphere Reserve (Benin, West Africa). *African Journal of Ecology*, 51: 358-365.
- Tchétan, E., Olounlade, P. A., Azando, E. V. B., Khaliq, H. A., Ortiz, S., Houngbeme, A., Quetin-Leclercq, J., 2022. Anthelmintic Activity, Cytotoxicity, and Phytochemical Screening of Plants Used to Treat Digestive Parasitosis of Small Ruminants in Benin (West Africa). *Animals*, 12(19), 2718.

* Presenting author. E-mail address: gahoungafabrice@yahoo.fr (Fabrice Jean Eudes Crédo Gahounga)

S2.P189 From the production of targeted metabolites to the complete valorization of *Coleus forskohlii* Briq. using aeroponic cultivation

Audrey Le Cabec^{a,*}, Pierre-Eric Campos^a, Cyril Colas^{a,b}, Olivier Yzebe^c, Emilie Destandau^a

^a Institut de Chimie Organique et Analytique (ICOA), UMR 7311, Université d'Orléans, Orléans, France

^b Centre de Biophysique Moléculaire (CBM), CNRS, UPR 4301, Orléans, France

^c Comité de Développement Horticole Région Centre (CDHRC), St-Cyr-en-Val, France

Accessing plant resources to extract compounds of interest can sometimes be difficult. To facilitate this access and limit environmental impact of transport, cultivation strategies can be developed. *Coleus forskohlii* Briq. a member of the Lamiaceae family (Ali Khan, 2012), used in Indian and Chinese traditional medicine and rich in diterpenoids, is widely cultivated for forskolin production. Indeed, this diterpenoid which presents anti-inflammatory activity (Suresh *et al.*, 2018) and activate adenylate cyclase (Seamon & Daly, 1981) is the major compound of interest, and it is found in the roots. In this regard, aeroponic cultivation of *C. forskohlii* had been developed in Region Centre Val de Loire (France) to facilitate access to the resources and more particularly to the roots. However, the phytochemical composition and the biological properties of a cultivated plant, that grows off-ground under another climate with less abiotic stress, when compared with the wild one, may vary. The first part of this study focused on optimizing the cultivation conditions to maximize forskolin production. Aeroponic cultivation has been realised for 18 weeks under various stresses. The quantification of forskolin using UHPLC-ELSD indicates that the use of LED lighting compared to the control 3-fold improve the total quantity of forskolin produced. Then a methodology employing non-targeted metabolomics approaches through UHPLC-HRMS/MS have been implemented to study the impact of this stress on other root metabolites. This methodology has been extended to the aerial part to explore potential co-valorization routes.

Keywords: *Coleus forskohlii* Briq., cultivation, aeroponic, forskolin, metabolomics

References

- Ali Khan, B. 2012. Botanical Description of *Coleus forskohlii*: A Review. Journal of Medicinal Plants Research, 6, 4832- 4835
- Seamon, K., Daly, J. W. 1981. Activation of adenylate cyclase by the diterpene forskolin does not require the guanine nucleotide regulatory protein. Journal of Biological Chemistry, 256(19), 9799–9801.
- Suresh, V., Deepika, G., Bantal, V., Beedu, S. R., Rupula, K. 2018. Evaluation of Anti-inflammatory and Anti-nociceptive Potentials of Andrographolide and Forskolin: *In vivo* Studies. Journal of Biologically Active Products from Nature, 8(5), 326–334.

* Presenting author. E-mail address: audrey.le-cabec@univ-orleans.fr (Audrey Le Cabec)

S2.P190 *ViscY* NMR experiments combined with chiral discrimination for complex mixture analysis

Anthony Budai*, Jean-Marc Nuzillard, Pedro Lameiras

Molecular Chemistry Institute of Reims, University of Reims Champagne-Ardenne, CNRS ICMR UMR 7312, 51097 Reims, France

So far, there are only a few approaches for analyzing complex mixtures using liquid-state NMR spectroscopy. One of them is to take profit from spin diffusion under viscous conditions. Using viscous solvents slows the tumbling rate of small molecules so that the longitudinal cross-relaxation becomes very efficient and thus promotes spin diffusion over entire molecular spin networks (Lameiras, P. *et al.*, 2021). Consequently, resonances of every mixed molecule can be grouped according to their ability to share magnetization by spin diffusion. As a result, we can individualize each compound's NMR spectrum using a *ViscY* (Viscosity Enhancement Spectroscopy) experiment (NOESY experiment under viscous conditions, Pedinielli *et al.*, 2020). Enantiomers, characterized by identical chemical shifts owing to their isochronous nuclei, present a challenge in distinguishing the individual component spectrum. The most common technique in NMR is to consider an additional chiral solvating agent (CSA) to induce the formation of diastereomer complexes. Therefore, they present different chemical shifts because of the differences in interaction between the CSA and both enantiomers (Wenzel, T. J., 2018). Our objective was to chirally discriminate and identify individual components of a racemic amino acid mixture made of histidine, tryptophan, threonine, methionine and asparagine by considering the combination of one CSA (Fig. 1) within two viscous solvent blends: DMSO-*d*₆/H₂O at 248 K and DMSO-*d*₆/H₃PO₄ at 288 K, at 500 MHz (¹H). Through a single 2D ¹H *ViscY* NOESY experiment, we successfully chirally discriminated between all amino acids and took advantage of spin diffusion to individualize all proton spectra.




Fig. 1. Chemical structure of the CSA : (-)-(18-crown-6)-2,3,11,12-tetracarboxylic acid.

Keywords: NMR, viscous solvent, spin diffusion, chiral discrimination, chiral solvating agent

References

- Lameiras, P. and Nuzillard, J.-M., 2021. Tailoring the nuclear Overhauser effect for the study of small and medium-sized molecules by solvent viscosity manipulation. *Progress in Nuclear Magnetic Resonance Spectroscopy*. 123: 1-50.
- Pedinielli, F. *et al.*, 2020. Mixture Analysis in Viscous Solvents by NMR Spin Diffusion Spectroscopy: *ViscY*. Application to High- and Low-Polarity Organic Compounds Dissolved in Sulfolane/Water and Sulfolane/DMSO-*d*₆ Blends. *Analytical Chemistry*. 92 (7), 5191-5199.
- Wenzel, T.J., 2018. Differentiation of chiral compounds using NMR spectroscopy. Ed: Wiley.

* Presenting author, E-mail address: anthony.budai@univ-reims.fr (Anthony Budai)

S2.P191 A green approach for preparing alkaloid-rich extracts and fractions

Lucas Apolinário Chibli^a, Bruna Ribeiro de Lima^a, Ariadne Magalhães Carneiro^a, Harley Cardoso Figueiró^a, Serhat Sezai Cicek^b, Verena Spiegler^c, Cristiano Soleo Funari^{a,*}

^a Green Biotech Network, School of Agricultural Sciences, São Paulo State University, Av. Universitária 3780, 18610-034 Botucatu, Brazil

^b Department of Biotechnology, Hamburg University of Applied Sciences, Ulmenliet 20, 21033 Hamburg, Germany

^c Institute of Pharmaceutical Biology and Phytochemistry, University of Münster, Corrensstraße 48, 48149 Münster, Germany

This work aimed to evaluate the performances of HCl and CH₂Cl₂ compared to citric acid (CA) and greener organic solvents (Prat et al., 2016) for preparing alkaloid-rich fractions. Plants were selected to constitute a diverse set of alkaloids: harmine (1), a β -carboline indole alkaloid from *Banisteriopsis caapi* (Spruce ex Griseb.) Morton, boldine (2), an aporphine alkaloid from leaves of *Peumus boldus* Molina, and mescaline (3), a phenylethylamine derivative from *Trichocereus macrogonus* var. *pachanoi* Britton & Rose (Fig. 1). Extractions were performed by dynamic maceration employing a classical acid-base extraction of alkaloids followed by liquid-liquid extractions (LLE). Each procedure was monitored by ultra-high-performance liquid chromatography coupled to a photodiode array detector and a mass spectrometer.




Fig. 1. Structure of alkaloids investigated in this work

The harmful HCl could be replaced with the greener CA (Wang et al., 2020) while keeping similar or better extraction efficiencies. EtOAc could substitute CH₂Cl₂ in at least one of the two LLE steps without reducing efficiencies. *tert*-amyl methyl ether (TAME), *n*-butyl acetate (BuOAc) or EtOAc led to gains in LLE performances when compared to CH₂Cl₂. The cases explored here refute the hypothesis that the hazardous CH₂Cl₂ should be the first option in this type of application. Based on prices currently charged by the main suppliers of chemicals used in this work (i.e., Sigma-Aldrich and Carl Roth), replacing CH₂Cl₂ with TAME, EtOAc, iPrOAc, BuOAc, or anisole would have little or no financial impact on the laboratory carrying out the investigation.

Keywords: green natural products chemistry, ecopharmacognosy, green chemistry

References

- Prat, D., Wells, A., Hayler, J., Sneddon, H., McElroy, C. R., Abou-Shehadeh, S., Dunn, P. J., 2016. CHEM21 selection guide of classical- and less classical-solvents Green Chem. 18, 288-296.
- Wang, J., Cui, Z., Li, Y., Cao, L., Lu, Z., 2020. Techno-economic analysis and environmental impact assessment of citric acid production through different recovery methods. J. Clean. Prod. 249, 119315.

This work was supported by The Sao Paulo Research Foundation (FAPESP, 2017/06216-6, 2018/01786-1)

* Presenting author. E-mail address: cristiano.funari@unesp.br (Cristiano Soleo Funari)

S2.P192 Optimization of a green method for quantifying harmine in *Banisteriopsis caapi* stems used in the preparation of ayahuasca

Geslane Custódio, Bruna Ribeiro de Lima, Lucas Apolinário Chibli, Ariadne Magalães Carneiro, Maria Gabriela de Albuquerque Santiago, Cristiano Soleo Funari*

Green Biotech Network, School of Agricultural Sciences, São Paulo State University, Av. Universitária 3780, 18610-034 Botucatu, Brazil




Fig. 1. Structure of harmine (1)

This work covers the optimization and validation of a green method for quantification of harmine (1) (Fig. 1) in stems of *Banisteriopsis caapi* (Malpighiaceae). Such stems are used in the preparation of a psychotropic drink called “ayahuasca”, being 1 their major biomarker (Katchborian-Neto et al., 2020). A reference extract (Katchborian Neto et al., 2020) was initially used for the optimization of a UHPLC-PAD-ESI-MS condition. Five factors were inserted in a fractional factorial design (FFD). The initial % of EtOH, time of isocratic elution, gradient time, and injection volume were statistically significant. As the separation at the central point of this PFF led to the resolution of harmine ($R_s > 2$), this condition was adopted for subsequent experiments (i.e., 150 x 2.1 mm, 1.8 μ m, ACQUITY Premier HSS T3 column; 40° C; Mobile phase of H₂O and EtOH both with 0.3% HCOOH; Flow rate of 0.27 mL/min; and injection volume of 1 μ L). Then, the extraction process itself was subject to optimization. A new FFD evidenced three significant factors that were enclosed in a central composite design (CCD). The following optimal point was found and confirmed: plant/solvent ratio of 30 mg/mL, solvent at pH 1, for 75 min. The full procedure was subjected to validation until intermediate precision. All relative standard deviations for retention time and peak area of 1 were $\leq 10\%$, attesting to the quality of the new procedure for the intended purpose. A calibration curve was built, leading to 4.8 mg/g of harmine in stems of *B. caapi*.

Acknowledgments: This work was supported by The Sao Paulo Research Foundation (FAPESP, 2017/06216-6, 2018/01786-1).

Keywords: green analytical chemistry, ecopharmacognosy, Ayuasca

References

- Katchborian-Neto, A., Santos, V. T., Nicário, K. J., Corrêa, J. O. A., Murgu, M., Martins, T. M. M., Gomes, D. A., Goes, A. M., Soares, M. G., Dias, D. F., Chagas-Paula, D. A., Paula, A. C. C. 2020. Neuroprotective potential of Ayahuasca and untargeted metabolomics analyses: applicability to Parkinson's disease. *J. Ethnopharmacol.* 255, 112743.

* Presenting author. E-mail address: cristiano.funari@unesp.br (Cristiano Soleo Funari)

S2.P193 Nutrient density and mineral content of radishes grown in black soldier fly frass-enriched compost

Sarah Morgan, Ceri Davies, Darrell Watts, Laura Bystrom*

School of Science, Bath Spa University, Newton Park, Newton St. Loe, Bath, BA2 9BN, UK

More sustainable food production methods are needed for the future, especially considering the challenges of climate change. *Hermetica illucens* (L.), also known as black soldier fly (BSF), shows promise as a tool in the circular economy, as it can convert organic waste into protein-rich feedstocks (Lui et al., 2022). Moreover, a byproduct of BSF is frass, which is a mix of excrement and exoskeletons (rich in minerals, fats, and proteins) that could be an effective fertilizer for food crops (Lopes et al., 2022). Limited information is available about the impact frass has on nutrient density of foods grown in frass-enriched compost. This study evaluates the growth (weight and height), mineral content (Mg, Mn, Zn, Fe, Cu) and nutrient density (% BRIX) of radishes (*Raphanus sativus* cv. Diana) grown in a range of frass concentrations (0-25% frass). Significant growth was found with all frass concentrations ($P < 0.001$), with optimal growth at 10% frass. Leaf BRIX, and whole plant Fe decreased ($P < 0.001$) at all frass concentrations compared to the control. No significant mineral increases were observed except for Mn ($P < 0.001$) in the leaves and roots with all frass concentrations, and Zn ($P < 0.001$) in the roots with 25% frass. Although added frass increased the growth of the radishes, this was offset by decreases in most minerals and consistent with BRIX values. These results indicate more research is needed to evaluate how frass-enriched compost impacts the growth and nutrient density of food crops.

Keywords: frass, nutrients, radish, *Hermetica illucens*, fertilizer

References

- Liu, T., Klammersteiner, T., Dregulo, A.M., Kumar, V., Zhou, Y., Zhang, Z. and Awasthi, M.K., 2022. Black soldier fly larvae for organic manure recycling and its potential for a circular bioeconomy: A review. *Sci. Total Environ.* 833, 155122.
- Lopes, I.G., Yong, J.W. and Lalander, C., 2022. Frass derived from black soldier fly larvae treatment of biodegradable wastes. A critical review and future perspectives. *Waste Manage.* 142, 65-76.

* Presenting author. E-mail address: l.bystrom@bathspa.ac.uk (Laura Bystrom)

S2.P194 Myristoleic acid ameliorates the inhibition of rat dermal papilla cell growth by fine particulate matter via the activation of autophagy and Wnt/ β -catenin pathways

Youn Kyung Choi^{a,b}, Jung-Il Kang^{a,b}, Da Hee Yang^{a,b}, Jin Won Hyun^{a,b}, Eun-Sook Yoo^{a,b}, Hee-Kyoung Kang^{a,b,*}

^a Department of Medicine, College of Medicine, Jeju National University, Republic of Korea

^b Jeju Research Center for Natural Medicine, Jeju National University, Republic of Korea

Air pollution, including fine particulate matter (PM), is recognized as a serious problem worldwide. PM has been reported to affect various organs of the human body, including cardiovascular, respiratory, and skin (Basith et al., 2022). However, its effect on alopecia remain unclear. In this study, we investigated the effects of PM on rat dermal papilla (rDP) cells and the protective effect of myristoleic acid against PM. PM suppressed the growth of rDP cells. To investigate the mechanism by which PM reduces rDP cells growth, we examined the effect of PM on signaling pathways. PM inhibited autophagy by increasing p-mTOR and p62 levels while decreasing the levels of beclin 1, Atg-3, -5, -7, and LC3. Furthermore, PM impairs the Wnt/ β -catenin signaling pathway by reducing p-GSK3 β and p- β -catenin. Myristoleic acid is an omega-5 monounsaturated fatty acid derived from the seeds of plants from the Myristicaceae family, which is already known to promote the growth of rDP cells. Myristoleic acid ameliorated the proliferation of rDP cells inhibited by PM through the activation of Wnt/ β -catenin and autophagy. Our results provide insights into the mechanisms underlying the adverse effects of PM on hair follicles and suggest that myristoleic acid could restore of PM-induced hair follicle damage.

Funding: This research was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(RS-2023-00246959 and RS-2023-00270936)]

Keywords: myristoleic acid, fine particulate matter, dermal papilla cells, autophagy, Wnt/ β -catenin

References

Basith, S., Manavalan, B., Shin, T.H., Park, C.B., Lee, W.S., Kim, J., Lee, G., 2022. The Impact of Fine Particulate Matter 2.5 on the Cardiovascular System: A Review of the Invisible Killer. *Nanomaterials* (Basel). 12, 2656.

* Presenting author. E-mail address: pharmkhk@jejunu.ac.kr (Hee-Kyoung Kang)

S2.P195 Identification of phenolic compounds in three *Veronica* species: *Veronica anagallis-aquatica* L., *Veronica persica* L. and *Veronica polita* L.

Ivana Vrca^{a,*}, Stjepan Orhanović^a, Ivana Pezelj^a, Karolina Sušić^a, Dario Kremer^b, Valerija Dunkić^a, Marija Nazlić^a

^a Department of Biology, Faculty of Science, University of Split, Ruđera Boškovića 33, 21000 Split, Croatia

^b Department of Ornamental Plants, Landscape Architecture and Garden Art, Faculty of Agriculture, University of Zagreb, Svetošimunska cesta 25, 10000 Zagreb, Croatia

Plants are known for the various biological activities which show, and for their specific chemical identity, which allows them to synthesize metabolites such as phenolic compounds, flavonoids, saponins, terpenes, coumarins, and alkaloids that possess protective properties against abiotic and biotic stresses (Cruz et al., 2023). Species of the genus *Veronica* are recognized for their traditional uses and the various biological activities they exhibit (Barreira et al., 2014). The genus *Veronica* (family Plantaginaceae) contains about 450 species, and so far only a few studies have confirmed that certain species possess biological activity such as antibacterial, antioxidant, anti-inflammatory and cytotoxic activities (Ignjatović et al., 2015). Therefore, the aim of this study was the extraction of phenolic compounds using different solvents (methanol, 80% ethanol, water) present in three *Veronica* species (*Veronica anagallis-aquatica* L., *V. persica* L. and *V. polita* L., and their identification and quantification using LC-MS/MS technique. The main phenolic compounds present in all three investigated *Veronica* species were: *p*-hydroxybenzoic acid, apigenin, vanillic acid and caffeic acid. Ethanol as a solvent proved to be better for the extraction of apigenin, while a higher amount of phenolic acids was found in the water extracts. It was extremely important to examine which solvent is the most optimal for the extraction of phenolic compounds and to see the differences in the chemical composition of the three species of the genus *Veronica* due to their influence on biological activity which will be tested.

Keywords: *Veronica* species, extraction, polyphenols, LC-MS/MS technique

References

- Barreira, J.C.M., Dias, M.I., Živković, J., Stojković, D., Soković, M., Santos-Buelga, C., Ferreira, I.C.F.R., 2014. Phenolic profiling of *Veronica* spp. grown in mountain, urban and sandy soil environments. *Food Chem.* 163, 275-283.
- Cruz, A., Sánchez-Hernández, E., Teixeira, A., Martín-Ramos, P., Cunha, A., Oliveira, R., 2024. Antifungal and antioomycete activities of a *Curcuma longa* L. hydroethanolic extract rich in bisabolene sesquiterpenoids. *Horticulturae*. 2024, 10 (124), 1-16.
- Ignjatović, Đ., Živković, J., Tovilović, G., Šavikin, K., Tomić, M., Maksimović, Z., Janković, T. 2015. Evaluation of angiogenic and neuroprotective potential of different extracts from three *Veronica* species. *Front. Life Sci.* 8:2, 107-116.

* Presenting author. E-mail address: ivrca@pmfst.hr (Ivana Vrca)

S2.P196 Turning the Tide: Harnessing the Invasive algae *Rugulopterix okamurae* for Commercial Applications

Amandine D'Unienville^a, Lucas Lasnel^a, Wadi Macquigneau^a, Riccardo Trentin^b, Maria J. Rodrigues^b, Luísa Custódio^{b,*}

^a Centre of Marine Sciences (CCMAR), University of Algarve, Campus de Gambelas, 8005 139 Faro, Portugal

^b Department of Biology, University of Padova, Via U. Bassi 58/B, 35131 Padova, Italy

The brown alga *Rugulopterix okamurae*, originating from the Pacific near Japan and Korea, has spread invasively to the Strait of Gibraltar, impacting coastal areas in Spanish Mediterranean and southern Portugal. Commercial use of invasive algae for their bioactive compounds offers a cost-effective strategy for controlling their spread and creating beneficial products. This study explored the chemical properties of biomass from *R. okamurae* collected in the southern Portugal, focusing on its fatty acids methyl esters profile, proximate and total phenolics composition and minerals. Furthermore, polar (water) and apolar (*n*-hexane) extracts were obtained by overnight maceration and appraised for *in vitro* antioxidant activity by different radical and metal-based assays, and for inhibition of enzymes related to neurological disorders (acetylcholinesterase: AChE, butyrylcholinesterase: BuChE), skin hyperpigmentation (tyrosinase), obesity (lipase), diabetes (α -glucosidase and α -amylase) and skin ageing (elastase). Biomass had higher levels of protein, lipids, and total carbohydrates than other algae, such as *Sargassum horneri*, and moderate total phenolics. It had a prevalence of saturated fatty acids (SFA), especially methyl palmitate (C16:0M). The polar extract had higher *in vitro* antioxidant properties, especially ferric reducing properties. The extracts had reduced enzymatic inhibitory properties.

Funding: Work funded by Foundation for Science and Technology (FCT) through project UIDB/04326/2020, UIDP/04326/2020 and LA/P/0101/2020, and the operational programs CRESC Algarve 2020 and COMPETE 2020 through project EMBRC.PTALG-01-0145-FEDER-022121. FCT supported L.C. by the FCT Scientific Employment Stimulus (CEECIND/00425/2017) and M.J.R. by the FCT program contract (UIDP/04326/2020).

Keywords: marine invasive species, algal invasions, alien species, antioxidant properties

* Presenting author. E-mail address: lcustodio@ualg.pt (Luísa Custódio)

S2.P197 Scale-up production of g-mangostin using Centrifugal Partition Chromatography

Anne-Marie Le Ray^{a,*}, Nadège Blon^a, Estelle Platon^a, Aurélie Guerin^a, Thomas Michel^b, Gregoire Audo^b, Guillaume Viault^a, Pascal Richomme^a

^a SONAS | Substances d'Origine Naturelle et Analogues Structuraux Univ Angers, SONAS, SFR QUASAV, F-49000 Angers, France

^b GILSON, Gilson Purification SAS ZA du Poteau, 22 rue Bourseul, 56890 Saint-Avé, France

g-mangostin, a polyprenylated xanthone isolated from the fruit pericarp of *Garcinia mangostana* (Clusiaceae), has demonstrated its value in agronomy as biocontrol agent acting on the UPR (Unfolded Protein Response) pathway of phytopathogenic fungi (Bataillé-Simoneau et al., 2023 ; Charpentier et al., 2023). To exploit this secondary metabolite industrially, we need to develop a large- scale purification method as green as possible. Since g-mangostin is most often isolated in the laboratory by Flash chromatography, a comparative study using Centrifugal Partition Chromatography (CPC) is currently being carried out on a 250 mL column (« lab scale ») with encouraging preliminary results (Fig.1). This development should enable us to carry out a scale-up on a 1 L column (« pilot scale ») before producing g-mangostin, at multi-kg scale, on a 5 L column (« Process scale »). The solvents used in all these steps are fully recycled. Finally, it should be noted that the initial extract of *G. mangostana* can be significantly enriched in g-mangostin by demethylation of the a-mangostin, the major compound.




Fig.1: γ -mangostin purification by CPC (VERITY® CPC Lab System 250, Gilson)

Keywords: *Garcinia mangostana*, g-mangostin, flash chromatography, centrifugal Partition Chromatography

References

- Bataillé-Simoneau, N., Charpentier, T., Guillemette T., Le Ray, A.-M., Richomme-Peniguel, P., Simoneau, P., Viault G., 2023. Compositions and methods for controlling phytopathogenic infections, WO 2023/099609 A1.
- Charpentier, T., Viault, G., Le Ray, A.-M., Bataillé-Simoneau, N., Helesbeux, J.-J., Blon, N., Bastide, F., Marchi, M., Aligon, S., Bruguère, A., Dinh, C.-P., Benbelkacem, Z., Dallery, J.-F., Simoneau, P., Richomme, P., Guillemette, T., 2023. Natural Products Targeting the Fungal Unfolded Protein Response as an Alternative Crop Protection Strategy. *J. Agric. Food Chem.* 71, 13706–13716.

* Presenting author. E-mail address: anne-marie.leray@univ-angers.fr (Anne-Marie Le Ray)

S2.P198 *In vitro* callus culture of *Mespilus germanica* to improve the production of bioactive compounds from a forgotten fruit

Valentina Parisi^{a,*}, Emanuele Rosa^a, Nunziatina De Tommasi^a, Giuliana Donadio^a, Daniele Fraternale^b

^a Dipartimento di Farmacia, Università degli Studi di Salerno, Via Giovanni Paolo II 132, Fisciano (SA), Italia

^b Dipartimento di Scienze Biomolecolari, DISB, Università di Urbino “Carlo Bo”, via Bramante 28, 61029 Urbino (PU), Italia

The common medlar is the fruit of *Mespilus germanica* L., (Rosaceae). Its consumption is today underestimated but anciently it was very successful; however, it seems that medlar is regaining an interest for its functional properties (Żońnierczyk et al., 2021). The *in vitro* culture of plant cells and tissues can be considered a valid and renewable source of material to obtain bioactive molecules without interferences of external phenomena (pollution, draught, seasonality, etc.) that could interfere in their production (da Silva Santos et al., 2022). Among the various *in vitro* cultures, callus one is interesting because it can be used as a starter for cell cultures on an industrial scale (Lystvan et al., 2018). This study aimed to develop a protocol to obtain production of callus biomass from medlar and subsequently, analyze and identify its composition. In detail, the ripe medlar pulp fruit was used as a source of explants for *in vitro* callus induction for the first time. Once the best conditions for biomass growth have been set, the metabolomic profile of callus was defined and then compared with the pulp and peel of fruits. The obtained results by combining MS and NMR data show that the callus is rich in ursane and olean triterpenoids, unlike pulp and peel. Furthermore, flavonoids, amino acids, sugars, and phenolic acids were also identified. The conspicuous presence of specialized metabolites, especially pentacyclic triterpenes, suggests the potential bioactivity of callus, thus the evaluation of antimicrobial activity on different Gram positive and negative bacterial strains is in progress.

Keywords: *Mespilus germanica*, callus biomass, LC-HRMS/MS, NMR, metabolomics

References

- da Silva Santos, É., Savam, A., Cabral, M. R. P., Castro, J. C., de Oliveira Collet, S. A., Mandim, F., ... & Gonçalves, R. A. C. 2022. Low-cost alternative for the bioproduction of bioactive phenolic compounds of callus cultures from *Cereus hildmannianus* (K.) Schum. J. Biotech. 356, 8-18.
- Lystvan, K., Kumorkiewicz, A., Szneler, E., Wybraniec, S., 2018. Study on betalains in *Celosia cristata* Linn. callus culture and identification of new malonylated amaranthins. J. Agric. Food Chem. 66, 3870–3879.
- Żońnierczyk, A. K., Ciałek, S., Styczyńska, M., Oziembłowski, M. 2021. Functional properties of fruits of common medlar (*Mespilus germanica* L.) extract. Appl. Sci. 11(16), 7528.

* Presenting author. E-mail address: vparisi@unisa.it (Valentina Parisi)

S2.P199 The phytochemical diversity of wild hops from northern France for potential brewing application

Augustin Bildstein^{a,*}, Mathilde Vandenberghe^b, Anne-Sophie Paguet^a, Gabriel Lefèvre^a, Louis Raymond^b, Gaëtan Kientz^a, Jennifer Samaillie^a, Franck Michels^a, Séverine Derbré^c, Cécile Palaric^a, David Mathiron^d, Marie-Laure Fauconnier^a, Jean-Xavier Fontaine^a, Roland Molinié^a, Sylvie Chollet^b, Céline Rivière^a

^a UMRt 1158 BioEcoAgro, Univ. Lille, JUNIA, UPJV, Univ. Liège, INRAE, Univ. Artois, Univ. Littoral Côte D'Opale, ICV—Institut Charles Viollette, F59650 Villeneuve D'Ascq, France

^b ICL, Junia, Université Catholique de Lille, LITL, F-59000 Lille, France

^c Université d'Angers, SONAS, SFR QUASAV, Faculty of Health Sciences, Dpt Pharmacy, 28 rue Roger Amsler, 49045 Angers, France

^d Université de Picardie Jules Verne, Plateforme analytique, Rue Dallery – Passage du sourire d'Avril, 80039 Amiens cedex, France

Hops, *Humulus lupulus* L., is one of the ingredients used in brewing, particularly for its antiseptic, bittering and aromatic properties. Breeding programs are looking for new varieties with original aromatic properties adapted to their terroir in a context of climate change. In our local scale, a study has highlighted the genetic and chemical diversity of wild hops harvested in northern France, suggesting an interest for brewing purposes (Paguet et al. 2024). The present study investigated the chemical diversity of 29 wild accessions collected in northern France and replanted *ex-situ* in an experimental hop field, in comparison with 19 commercial varieties as well as two heirloom varieties. Hop phytochemical analysis included quantification of major prenylated phenolic compounds by UHPLC-MS, untargeted metabolomic and molecular networking studies by UHPLC-HRMS/MS, and analysis of volatile compounds by HS-SPME GC-MS. In addition, experimental beers were brewed and hopped with 13 wild hops from our collection, 14 commercial varieties and 2 heirloom varieties. In order to evaluate the brewing potential of wild hops, experimental beers were analyzed by sensory characterization, performed by a trained panel of 19 people, associated to physico-chemical characterization including analysis of volatile compounds by SBSE GC-MS and bitterness quantification by UV spectroscopy. Wild hops from northern France presented lower cohumulone contents within total alpha acids and a greater quantity and diversity of non-oxygenated sesquiterpenes. The brewing potential of wild hops has been confirmed by sensory analysis and several accessions were comparable to commercial hops on specific descriptors such as yellow fruits or floral.

Acknowledgments: The authors thank the Haut-de-France Region, the French Fondation of Brewery and Maltery, Junia- ISA, the french inter-branch organization of hop “Interhoublon”, the french brewery union “Syndicat des Brasseurs des Hauts-de-France” as well as the CPER project “BiHauts Eco de France” for their financial contribution.

Keywords: Hops, untargeted metabolomics, volatile compounds, sensory analysis, brewing

Reference

Paguet A-S., Siah A., Lefèvre G., Vandenberghe M., Lutun D., Degardin N., Samaillie J., Mathiron D., Dermont C., Michels F., Fauconnier M-L., Chollet S., Molinié., Fontaine J-X., Sahpaz S., Rivière C., 2024. Phytochemical characterisation and aromatic potential for brewing of wild hops (*Humulus lupulus* L.) from Northern France: Towards a lead for local hop varieties. Food Chemistry. 433, 137302.

* Presenting author. E-mail address: augustin.bildstein.etu@univ-lille.fr (Augustin Bildstein)

S2.P200 Effect of the aquaponic system on productivity and quality of *Acmella oleracea* R.K. Jansen

Marta Ferrati^{a*}, Eleonora Spinozzi^a, Cecilia Baldassarri^a, Rita Giovannetti^b, Marco Zannotti^b, Valentina Bolletta^c, Marco Cespi^a, Riccardo Petrelli^a, Filippo Maggi^a

^a Chemistry Interdisciplinary Project (ChIP), School of Pharmacy, University of Camerino, Via Maddona Delle Carceri, 62032 Camerino, Italy

^b Chemistry Interdisciplinary Project (ChIP), School of Science and Technology, University of Camerino, Via Maddona Delle Carceri, 62032 Camerino, Italy

^c Mj Energy srl Società Agricola, Treia, Italy

Acmella oleracea R.K. Jansen (Asteraceae) has recently aroused great interest in several fields (Rondanelli et al., 2020; Uthpala et al., 2021). Thus, great quantities of biomass are required. The *A. oleracea* biological properties are ascribed to *N*-alkylamides, being spilanthol the main representative (Spinozzi et al., 2022). To date, *A. oleracea* production mainly involves in field and hydroponic cultivations, which require large amounts of water, fertilizers, and pesticides. Aquaponics, characterized by a symbiotic ecosystem in which fish, plants and nitrifying bacteria take part, represents an environmentally friendly technique to produce crops. This system potentially overcome environmental issues minimizing nutrient and fertilizer requirements, and water wastage (Yep and Zheng, 2019). On this basis, the quality of two cultivars of *A. oleracea*, i.e., cv. yellow and cv. purple, grown in aquaponic systems was evaluated every two weeks, from 4th July to 29th September 2023. Specifically, capitula production, chlorophyll content by spectrophotometric assay and *N*-alkylamides content by HPLC-DAD-MS analyses were assessed. Monitoring of the main water nutrients through ICP-MS analyses was also carried out. The obtained data were compared to those obtained for plants grown in hydroponics. Statistical analysis revealed that cv purple exhibited notably higher levels of spilanthol (14.85-48.0 mg/g of dried flowers) than cv yellow (10.93-40.77 mg/g). Moreover, a significant increase in flowerhead numbers and *N*-alkylamides concentration was observed during full flowering period (18th August -15th September 2023). Overall, the results suggested that aquaponic systems offer similar efficiency to traditional hydroponic methods, making them a viable alternative for large-scale *A. oleracea* production.

Keywords: *Acmella oleracea*, aquaponics, cultivar, eco-friendliness, *N*-alkylamides

References

- Rondanelli, M., Fossari, F., Vecchio, V., Braschi, V., Riva, A., Allegrini, P., ... & Perna, S., 2020. *Acmella oleracea* for pain management. *Fitoterapia* 140, 104419.
- Spinozzi, E., Ferrati, M., Baldassarri, C., Cappellacci, L., Marmugi, M., Caselli, A., ... & Petrelli, R., 2022. A Review of the Chemistry and Biological Activities of *Acmella oleracea* ("jambù", Asteraceae), with a View to the Development of Bioinsecticides and Acaricides. *Plants* 11(20), 2721.
- Uthpala, T.G.G., Navaratne, S.B., 2021. *Acmella oleracea* plant; identification, applications and use as an emerging food source–review. *Food Reviews International* 37(4), 399-414. <https://doi.org/10.1080/87559129.2019.1709201>
- Yep, B., and Zheng, Y., 2019. Aquaponic trends and challenges–A review. *J. Clean. Prod.* 228, 1586-1599.

* Presenting author. E-mail address: marta.ferrati@unicam.it (Marta Ferrati)

S2.P201 Comparison of free volatile compounds of wild and cultivated species *Veronica catenata* Pennell

Valerija Dunkić^{a,*}, Marija Nazlić^a, Ivana Vrca^a, Karla Akrap^a, Niko Dunkić^b, Dario Kremer^c

^a Department of Biology, Faculty of Science, University of Split, Ruđera Boškovića 33, 21000 Split, Croatia

^b Emergency Medicine, General Hospital Šibenik, Stjepana Radića 83, 22000 Šibenik, Croatia

^c Department of Ornamental Plants, Landscape Architecture and Garden Art, Faculty of Agriculture, University of Zagreb, Svetošiminska cesta 25, 10000 Zagreb, Croatia

Plants of the genus *Veronica* and their extracts are used worldwide in traditional medicine and nutrition (Salehi et al., 2019; Xue et al., 2019). Extracts from the edible species *Veronica catenata* Pennell (Plantaginaceae), the pink water speedwell, are the subject of the present study. The compositions of the extracts comprising the free volatile compounds are compared between plants collected in their natural habitat and those that have been cultivated. The cultivated speedwell was grown from seeds collected the previous year. The isolation of free volatile compounds was performed by microwave-assisted extraction, and each extract obtained consists of two phases: a lipophilic (essential oil) and an aqueous phase (hydrosol). The chemical compositions of the isolates were determined by gas chromatography-mass spectrometry. The essential oil of *V. catenata* collected in the natural habitat was dominated by phytol 42.26% and hexahydrofarnesyl acetone 17.22% (Dunkić et al., 2021). Hexahydrofarnesyl acetone is present in a similar percentage (17.86%) in the essential oil obtained from cultivated plants. With content of 6.47%, phytol is significantly less present in cultivated plants. In the hydrosol phase of *V. catenata* from wild species, the dominant compounds are α -muurolol (35.12%) and hexadecanoic acid (12.85%) (Nazlić et al., 2022). These two compounds also dominated the composition of hydrosol in the cultivated plant with a similar percentage of identification. Water speedwell is an edible species and because of this it is desirable for cultivation. Volatile compounds are a part of specialized metabolites and also contribute to the plant overall biological activity. This study showed that, regardless of whether it is a wild-collected or a cultivated *V. catenata* species, the same basic volatile compounds have been identified, making the cultivated species also suitable for further study.

Keywords: *Veronica catenata*, Plantaginaceae, essential oil, hydrosol

References

- Salehi, B., Shivaprasad Shetty, M., V. Anil Kumar, N., Živković, J., Calina, D., Oana Docea, A., Emamzadeh-Yazdi, S., Sibel Kılıç, C., Goloshvili, T., Nicola, S., et al., 2019. *Veronica* Plants—Drifting from Farm to Traditional Healing, Food Application, and Phytopharmacology. *Molecules*. 24, 2454.
- Xue, H., Chen, K.X., Zhang, L.Q., Li, Y.M., 2019. Review of the Ethnopharmacology, Phytochemistry, and Pharmacology of the Genus *Veronica*. *Am. J. Chin. Med.* 47, 1193–1221.
- Dunkić, V., Nazlić, M., Ruščić, M., Vuko, E., Akrap, K., Topić, S., Milović, M., Vuletić, N., Puizina, J., Jurišić Grubešić, R., Srećec, S., Kremer, D., 2022. Hydrodistillation and Microwave Extraction of Volatile Compounds: Comparing Data for Twenty-One *Veronica* Species from Different Habitats. *Plants*. 11, 902.
- Nazlić, M., Akrap, K., Kremer, D., Dunkić, V., 2022. Hydrosols of *Veronica* Species—Natural Source of Free Volatile Compounds with Potential Pharmacological Interest. *Pharmaceuticals*. 15, 1378.

* Presenting author. E-mail address: dunkic@pmfst.hr (Valerija Dunkić)

S2.P202 Urban parks biowaste as a sustainable source of new antidiabetics

Marijan Marijan, Lejsa Jakupović, Lovorka Vujić, Marijana Zovko Končić*

University of Zagreb Faculty of Pharmacy and Biochemistry, A. Kovačića 1, 10000 Zagreb, Croatia

Biowaste produced in urban parks is composed of large masses of organic matter that is only occasionally used economically. In this work, we aimed to investigate if urban parks biowaste contains phenolic compounds that can be used for production of high-value antidiabetic products. Seven plants, widely distributed in urban parks in Central Europe (*Achillea millefolium*, *Cichorium intybus*, *Centaurea jacea*, *Malva sylvestris*, *Medicago sativa*, *Plantago lanceolata*, and *Trifolium pratense*) were extracted using 10% and 50% ethanol. Chemical composition of the prepared extracts was determined using HPLC and UV-VIS methods, while their antioxidant, anti-inflammatory and antidiabetic activity were determined using chemical and *in vitro* enzymatic methods. The extracts were rich in total phenols, flavonoids and phenolic acids, including caffeic acid, quercetin, kaempferol, luteolin and apigenin derivatives. They demonstrated moderate antioxidant activity as evidenced by their oxygen radical absorbance capacity (ORAC), reducing power and DPPH antiradical activity. The extracts were able to inhibit the activity of lipxygenase, as well as heat- induced ovalbumin coagulation, albeit weaker than the positive controls nordihydroguaiaretic acid and diclofenac. On the other hand, several extracts demonstrated excellent antidiabetic activity, assessed through inhibition α -glucosidase and amylase. For example, anti- α -glucosidase activity of 10% and 50% ethanol extract of *M. sativa* extracts ($9.35 \pm 0.07 \mu\text{g/mL}$ and $3.44 \pm 0.05 \mu\text{g/mL}$, respectively) was higher than the activity of the positive control, acarbose ($156.59 \pm 0.80 \mu\text{g/mL}$). The results indicate that the biowaste obtained from urban parks represents an ecologically acceptable alternative to conventional cultivation for the preparation of ecologically acceptable, high-value antidiabetic products.

Keywords: α -glucosidase, amylase, antidiabetic, phenolics, urban parks

* Presenting author. E-mail address: mzovko@pharma.hr (Marijana Zovko Končić)

S2.P203 Unravelling the secrets of fungicides through actinobacteria-phytopathogen interactions

Kah Yean Lum*, Ling Ding

Department of Biotechnology and Biomedicine, Technical University of Denmark, Søtofts Plads 221, DK-2800 Kgs. Lyngby, Denmark

Phytopathogenic fungi and fungi-like oomycetes contribute to substantial global yield losses in agricultural production annually, with several fungi known to produce toxins that have severe impacts on animal and human health (Ristaino *et al.*, 2021). While synthetic pesticides have been widely used to chemically control microbial infestation in agricultural fields, their extensive use has raised serious concerns such as residual effects and toxicity to human health, biodiversity decline in agricultural areas, and environmental pollution (Kim *et al.*, 2017). Therefore, phasing out synthetic pesticides with proper substitutes is essential to create more resilient and sustainable food systems to feed a growing world population. Biological controls, involving the introduction of naturally occurring agents such as beneficial microorganisms or crude extracts from natural products directly into a natural ecosystem, have emerged as one of the most promising approaches to mitigate crop diseases. As *actinobacteria* are the most important antibiotic producer and co-exist with fungi in nature, we aim to investigate the biotechnological potential of actinobacteria and their secondary metabolites for use in the biocontrol of phytopathogens through advanced chemical ecology and natural product chemistry. This presentation will cover the isolation of the actinobacteria from Danish soil and the antifungal evaluations of the isolated strains against *Alternaria solani* and *Phytophthora infestans*, the causative agents of early blight and late blight diseases, respectively. Additionally, LCMS-based metabolomics analysis of the prioritized strains will be described, and the on-going workflow involves large-scale fermentation and bioassay-guided isolation for the identification of antifungal compounds.

Keywords: actinobacteria, antifungal, *Alternaria solani*, *Phytophthora infestans*, metabolomics

References

- Kim, K.-H.; Kabir, E.; Jahan, S. A. 2017. Exposure to pesticides and the associated human health effects. *Sci. Total Environ.* 575, 525-535.
- Ristaino, J. B.; Anderson, P. K.; Bebber, D. P.; Brauman, K. A.; Cunniffe, N. J.; Fedoroff, N. V.; Finegold, C.; Garrett, K. A.; Gilligan, C. A.; Jones, C. M.; *et al.* 2021. The persistent threat of emerging plant disease pandemics to global food security. *Proc. Natl. Acad. Sci. USA* 118 (23), e2022239118.

* Presenting author. E-mail address: kyelu@dtu.dk (Kah Yean Lum)

S2.P204 The involvement of cyclotides in plant response to heavy metals in *Viola* spp.

Błażej Ślęzak^{a,b,*}, Klaudia Sychta^c, Rebecca Miszczak^{c,d}, Ulf Göransson^b, Anna Nilsson^e, Per E. Andrén^e, Aneta Słomka^c

^a W. Szafer Institute of Botany, Polish Academy of Science, 46 Lubicz St., 31-512, Kraków, Poland

^b Pharmacognosy, Department of Pharmaceutical Biosciences, Uppsala University, P.O. Box 591, 751 24 Uppsala, Sweden

^c Department of Plant Cytology and Embryology, Institute of Botany, Faculty of Biology, Jagiellonian University in Kraków, 9 Gronostajowa St., 30-387 Cracow, Poland

^d Doctoral School of Exact and Natural Sciences, Jagiellonian University in Kraków, 11 prof. S. Łojasiewicza St., 30-348 Cracow, Poland

^e Spatial Mass Spectrometry, Science for Life Laboratory, Department of Pharmaceutical Biosciences, Uppsala University, P.O. Box 591, 751 24 Uppsala, Sweden

Heavy metals are a significant pollutant resulting from the extraction and combustion of fossil fuels (Munawer, 2018). The genus *Viola* is rich in metallophyte species, that can be used in phytoremediation of heavy metals from polluted soils (Sychta et al., 2020). Cyclotides are plant cyclic cysteine knot peptides involved in the host plant's response to various types of stress. Some studies have shown that some cyclotides may have the ability to bind metals (Skjeldal et al., 2002). All the *Viola* species screened for cyclotides showed the production of a wide variety of these peptides (Burman et al., 2015). The present study reviews the results of several studies we have carried out in recent years to test the hypothesis that cyclotides may be involved in mechanisms of tolerance to heavy metals in *Viola* spp. The studies involved several species and populations of violets, both metallophytes and non-metallophytes: *V. uliginosa*, *V. westwalika*, *V. tricolor*, and *V. arvensis*. The effects of heavy metals on cyclotide production were tested *in vitro* and *in vivo*. Quantitative and qualitative analysis of cyclotides was based on mass spectrometry (LC-MS and MALDI-MS). All the studies clearly showed that heavy metal treatment prompts plant response in terms of the change in the production of cyclotides. This indicates that cyclotides play a role in the mechanisms of tolerance. However, the complexity of cyclotide involvement in the tolerance mechanisms requires mechanistically focused studies on the interaction of these peptides with heavy metal ions. Funding: NCN Opus 2021/41/B/NZ8/01737

Keywords: plant peptides, tolerance to heavy metals, abiotic stress

References

- Burman, R., Yeshak, M.Y., Larsson, S., Craik, D.J., Rosengren, K.J., Göransson, U., 2015. Distribution of circular proteins in plants: large-scale mapping of cyclotides in the Violaceae. *Front. Plant Sci.* 6.
- Munawer, M.E., 2018. Human health and environmental impacts of coal combustion and post-combustion wastes. *J. Sustain. Min.* 17, 87–96.
- Skjeldal, L., Gran, L., Sletten, K., Volkman, B.F., 2002. Refined structure and metal binding site of the kalata B1 peptide. *Arch. Biochem. Biophys.* 399, 142–148.
- Sychta, K., Słomka, A., Kuta, E., 2020. Garden pansy (*Viola x wittrockiana* Gams.) - a good candidate for the revitalisation of polluted areas. *Plant, Soil Environ.* 66, 272–280.

* Presenting author. E-mail address: b.slazak@botany.pl (Błażej Ślęzak)

S2.P205 Applying the global nature products social molecular networking (GNPS) platform to identify natural products involved in tolerance to heavy metals in *Viola* spp.

Karpierz K.^{a,*}, Miszczak R.^{b,c}, Sychta K.^b, Słomka A.^b, Göransson U.^d, Slazak B.^{a,d}

^a W. Szafer Institute of Botany, Polish Academy of Science, 46 Lubicz St., 31-512, Kraków, Poland

^b Department of Plant Cytology and Embryology, Institute of Botany, Faculty of Biology, Jagiellonian University in Kraków, 9 Gronostajowa St., 30-387 Cracow, Poland

^c Doctoral School of Exact and Natural Sciences, Jagiellonian University in Kraków, 11 prof. S. Łojasiewicza St., 30-348 Cracow, Poland

^d Pharmacognosy, Department of Pharmaceutical Biosciences, Uppsala University, P.O. Box 591, 751 24 Uppsala, Sweden

The extraction and combustion of fossil fuels are one of major sources of heavy metal pollution. Various potential strategies for coping with heavy metal pollution involve the usage of plants. One of the methods – phytoremediation – involves the use of heavy metal-resistant plants to extract the pollutants from the soil. The genus *Viola*, Violaceae is rich in metallophyte species such as *V. tricolor* L. displaying high resistance to heavy metals, that live and thrive on polluted soils. Multiple classes of different molecules are involved in the mechanisms of tolerance to heavy metals in plants, e.g. phytochelatins or flavonoids. GNPS is a platform that identifies molecules or their classes based on mass spectrometry – LC-MS and LC-MSMS. Our recent study on the metabolomic basis of heavy metal tolerance in *V. tricolor* yielded hundreds of LC-MS peaks corresponding to unknown compounds potentially involved in the response. The current study aimed to identify these molecules using GNPS. Samples (roots and leaves) from two populations (metallicolous, nonmetallicolous) were extracted through maceration in methanol and analyzed using data-dependent analysis (DDA) LC-HR-MSMS. The collected data was subjected to GNPS workflow. The constructed networks allowed for the theoretical identification of several compound classes in the dataset: e.g. flavonoids, lipids, and sterols.

Keywords: tolerance to heavy metals, abiotic stress, metabolomics, bioinformatics

* Presenting author. E-mail address: k.karpierz@botany.pl (Katarzyna Karpierz)

S2.P206 Screening plant extract for biological activity. A work spanning 60° Latitude South

Ivana Barbosa Suffredini^{a,b,*}, Amanda Leal da Silva Teodoro^a, Marcelo Henrique Soller Ramada^c, Sergio Alexandre Frana^{a,b}, Mateus Luís Barradas Paciencia^a, Jefferson de Souza Silva^d, Paulo Eduardo Aguiar Saraiva Câmara^e, Antonio Drauzio Varella^a

^a Center for Research in Biodiversity, Paulista University, 902 Dr. Bacelar Str., 05026-002 São Paulo, São Paulo, Brazil

^b Graduate Program in Environmental and Experimental Pathology, Paulista University, 1212 Dr. Bacelar Str., 05026-002 São Paulo, Brazil

^c BRIOTECH and Graduate Program in Genomic Sciences and Biotechnology, Brasília Catholic University, QS 07, Lote 01, 71966-700 Brasília, Federal District, Brazil

^d Amazônia University, 1544 Ville Roy Ave., 69307-725 Boa Vista, Roraima, Brazil

^e Botany Department, Brasília University, Caixa Postal 04457, 70919-970 Brasília, Federal District, Brazil

This presentation will cover the achievements of accessing the potential of Antarctic mosses' biological activities. The development of the work was based on the methodological algorithm developed during our 30-year experience in prospecting plants in the Amazon rain forest made under rigid protocols involving extract preparation by maceration, disk-diffusion and microdilution-broth assays, thin-layer chromatography, bioautography tests, antioxidant assays, *Artemia salina* toxicity assay, statistical analyses (Suffredini *et al.*, 2023), and phenolic compound quantification by the Folin-Ciocalteu method. So, 24 extracts obtained from the Antarctic moss *Sanionia uncinata* were evaluated for their biological-active potential and their chemical profiles, obtained by gas- chromatography coupled to mass spectrometry (GC-MS). Only two out of 24 extracts showed antibacterial activity against *Staphylococcus aureus*. The extracts showed strong antioxidant activity and high levels of phenolic compounds. Diterpenes, triterpenes, steroids, and fatty acid metabolites have contributed to the biological/chemical achievements. The experience previously acquired with the Brazilian plant extracts was primordial to studying the Antarctic plants due to the small amount of biomass that was possible to collect during fieldwork, and one species was used as a model to be adapted to the Antarctic plants. This species showed antibacterial and antioxidant activities. Carrying out prospecting projects is based on persistence, systematization, and resilience as a way of understanding the biodiversity potential of tropical forests and more distant places in the era of climate change.

Acknowledgments: CNPq, CAPES, FAPESP.

Keywords: Antarctic, Brazilian biodiversity, *in vitro* assays, large-scale screening, statistical analyses

References

Suffredini, I.B., *et al.*, 2023. Screening Amazon rainforest plant extracts for antimicrobial activity: a 15-year commitment to the Brazilian biodiversity. *Front. Antib.* 2, 1122400. DOI:10.3389/frabi.2023.1122400

* Presenting author. E-mail address: ivana.suffredini@docente.unip.br (Ivana Barbosa Suffredini)

S2.P207 Polyphenol and anthocyanin content in early producing maqui hybrids (*Aristotelia chilensis*)

Benita González, Esteban Osores, Ursula Doll, Hermine Vogel*

CENATIV, Facultad de Ciencias Agrarias, Universidad de Talca, Av. Lircay S/N, Casilla 747, Talca, Chile

Maqui berries are known for their high content of polyphenols and anthocyanins (Schreckinger et al., 2010). *Aristotelia chilensis* (Mol.) Stuntz, Elaeocarpaceae, is a bush or tree native to Chile and the western Argentine Patagonia. Its outstanding antioxidant properties (Araya et al., 2006) triggered the overexploitation of natural populations through wild collection to satisfy the growing demand for different maqui berry-based products. To ensure sustainable production, commercial varieties have been developed (Vogel et al., 2016). 69 hybrids have been selected from individuals obtained by crossing plants with outstanding fruit yield and quality characteristics. These genotypes have been evaluated for the proportion of pulp in fruits, and their concentration of polyphenols and anthocyanins. At the same time, the ripening period of each hybrid was recorded to detect early producing genotypes for agricultural production, especially in regions with less water availability during the summer months. The hybrids showed a range of polyphenol concentrations from 0.4 to 6.0 EAG/100 g dry matter (DM), and anthocyanin concentrations from 0.7 to 2.2 g cy-3-glu/100 g DM. The proportion of pulp in the berries ranged from 14 to 82% with an average of 51%. For future agricultural production in regions with water scarcity, early producing genotypes were identified, with a short ripening period, high proportions of pulp, and high concentrations of polyphenols and anthocyanins.

Keywords: maqui berry, proportion of pulp, ripening period, adaptation to water scarcity

References

- Araya L., H., Clavijo R., C., Herrera, C., 2006. Capacidad antioxidante de frutas y verduras cultivados en Chile. Archivos Latinoamericanos de Nutrición 56(4), 361-365.
- Schreckinger, M.E., Lotton, J., Lila, M.A., Gonzalez de Mejia, E., 2010. Berries from South America: A comprehensive review on chemistry, health potential, and commercialization. J Med Food 13(2), 233–246.
- Vogel, H., González, B., Catenacci, G., Doll U., 2016. Domestication and sustainable production of wild crafted plants with special reference to the Chilean Maqui Berry (*Aristotelia chilensis*). Julius-Kühn-Archiv 453, 50-52.

* Presenting author. E-mail address: hvogel@utalca.cl (Hermine Vogel)

S2.P208 Effect of different ethylene levels on other phytohormones and *Arabidopsis* resilience

Mersa Darbandsari*, Michal Karady, Miroslav Strnad

Laboratory of Growth Regulators Palacký University & Institute of Experimental Botany of the Czech Academy of Sciences, Šlechtitelů 27, 78371 Olomouc, Czech Republic

Ethylene, a distinctive gaseous phytohormone, plays a pivotal role in various developmental processes and frequently interacts with other phytohormones. It governs essential plant processes, including fruit ripening and senescence, highlighting its significance in plant physiology. Understanding the nuanced effects of ethylene is crucial for optimizing agricultural practices and effective crop management (Lin et al., 2009). Plants, being immobile, encounter challenges posed by abiotic factors such as drought, salinity, and extreme temperatures. These challenges significantly limit their adaptability, alter growth and development, and consequently reduce crop yield (Zhang et al., 2022). We aim to explore the impact of reduced or elevated ethylene production in *Arabidopsis* mutants on the content of other phytohormones and also possibly explore their relationship under stress conditions. This research introduces a sensitive methodology for assessing levels of ethylene or its direct precursor 1-aminocyclopropane-1-carboxylic acid (ACC) by means of a new liquid chromatography-tandem mass spectrometry method, employing a fully validated method with a specific derivatization technique for accurate ACC and phytohormone profiling. Our results show a clear influence between elevated ethylene and ACC levels and other phytohormones and their selected metabolites content.

Keywords: *Arabidopsis thaliana*, ethylene, phytohormones, abiotic stress

References

- Lin, Zhefeng, Silin Zhong, and Don Grierson. 2009. Recent advances in ethylene research. *Journal of Experimental Botany* 60, 3311-3336.
- Zhang, Huiming, et al. 2022. Abiotic stress responses in plants. *Nature Reviews Genetics* 23, 104-119.

* Presenting author. E-mail address: mersa.darbandsari@gmail.com (Mersa Darbandsari)

S2.P209 Copper-induced siderophore production and copper complex formation in vitamin-producing bacteria at a Vermont copper mine

Lesley-Ann Giddings^{a,b,*}, Robert Samples^b, Pauline Baranger-Hartwig^b

^a Department of Chemistry, Smith College, 100 Green St, Northampton, MA, 01063 USA

^b Biochemistry Program, Smith College, 100 Green St, Northampton, MA, 01063 USA

Ely Copper Mine in Vermont, USA, a metal-polluted Superfund site, has been accumulating acidic water (pH > 2) or acid rock drainage (ARD) for over 50 years, contributing to contamination of nearby water bodies (Fig. 1). Such contamination is exacerbated by rising flooding events associated with climate change. We previously characterized its Proteobacteria-dominated ARD microbiome and isolated amoebae and associated bacteria (*Dyella*) from this copper-contaminated (2 mg/L water; 2 mg/kg sediment) site (Giddings, 2020; Giddings 2022).




Fig. 1. ARD at Ely Copper Mine in Vershire, Vermont and bacterial metabolites detected.

To develop site-specific strategies for remediation, we explored the ARD microbial community for producers of metal-chelators or siderophores. Siderophore-producing *Dyella* were identified using a colorimetric chrome azurol S plate assay and isolates were grown in complex media with increasing copper concentrations (0–100 μ M). Metabolomics analyses of organic extracts of *Dyella* showed an abundance of pantothenic acid (vitamin B5), riboflavin (vitamin B2), and pyridoxine (vitamin B6), which are likely why these bacteria associate with amoebae within this extreme environment (Fig. 1). Notably, siderophores, such as desferrioxamine E, were detected and observed to increase with increasing copper concentrations. Copper also formed stable complexes with several flavins. Our data provide insight into the symbioses and adaptations amoebae employ to survive in nutrient-deficient, metal-polluted environments and offer the potential to find new bioremediation methods.

Keywords: metal pollution, siderophores, *Dyella*, copper-induced metabolites

References

- Giddings, L.A., Chlipala, G., Kunstman, K., Green, S., Morillo, K., Bhawe, K., Peterson, H., Driscoll, H., Maienschein- Cline, M., 2020. Characterization of an acid rock drainage microbiome and transcriptome at the Ely Copper Mine Superfund site. *PLoS One* 15, p.e0237599.
- Giddings, L.A., Kunstman, K., Moumen, B., Asiama, L., Green, S., Delafont, V., Brockley, M. and Samba-Louaka, A., 2022. Isolation and genome analysis of an amoeba-associated bacterium *Dyella terrae* strain Ely Copper Mine from acid rock drainage in Vermont, United States. *Front. Microbiol.* 13, p.856908.

* Presenting author. E-mail address: lgiddings@smith.edu (Lesley-Ann Giddings)

S2.P210 Disentangling metabolite dynamics in extreme environments: life detection strategies for extraterrestrial worlds

Jacob M.C. Shaffer^a, Lesley-Ann Giddings^b, Robert Samples^b, Eli C. Sklute^{c,d}, Abigail Jarratt^a, Katherine Mateos^e, M. Darby Dyar^{c,f}, Peter A. Lee^g, Jill A. Mikucki^{a,*}

^a Department of Microbiology, University of Tennessee - Knoxville, 1311 Cumberland Ave, Knoxville TN 37996, USA

^b Biochemistry Program, Smith College, 100 Green Street, Northampton, MA 01063, USA

^c Planetary Science Institute, 1700 E Fort Lowell Rd STE 106, Tucson, AZ 85719, USA

^d Los Alamos National Laboratory, Los Alamos, NM 87545, USA

^e Ocean Sciences Department, University of California - Santa Cruz, 1156 High Street Santa Cruz, CA 95064, USA

^f Department of Astronomy, Mount Holyoke College, South Hadley, MA 01075, USA

^g Hollings Marine Laboratory, 331 Fort Johnson Rd., Charleston, SC 29412, USA

Microbes from terrestrial extreme environments offer an opportunity to test biosignature production in conditions relevant to astrobiological targets. In Antarctica, iron-rich brine (~3X seawater salinity) leaks through a glacier forming a bright red 'waterfall' known as Blood Falls (Fig. 1). This feature provides a portal into ecosystems hidden beneath the Antarctic ice sheet, one of the least explored biomes on Earth. This site is also analog for ecosystems beyond Earth such as on Enceladus, a moon of Saturn, that harbors a liquid ocean beneath its ice shell, or possible Martian groundwater systems (Sklute et al. 2022).



Figure 1. Blood Falls, McMurdo Dry Valleys, Antarctica (credit: P. Rejeck).

To identify biosignatures produced during microbial iron transformations, we grew *Shewanella* sp. (strain BF02_schw), originally isolated from Blood Falls (Boles et al. 2021), in minimal marine media with amorphous iron hydroxides (100 mM) and analysed the volatilome, metabolome, transcriptome, and mineralogy for indicators of microbe-mineral interactions. Significant tryptophan degradation was observed in iron treatments, suggesting potential Fenton reaction mediated nutrient loss, however this loss may be moderated as genes regulating tryptophan biosynthesis were concomitantly overexpressed. Altering membrane fluidity is a known strategy for growth at low temperatures in psychrotolerant *Shewanella* species (Bowman et al. 1997). In iron treatments, we observed overexpression of eicosapentaenoic acid (EPA) biosynthesis genes, suggesting EPA may also help manage iron toxicity. Collectively our results show the poly- extremophile strain BF02_schw produces diverse molecular biosignatures indicative of growth in a metal-rich environment. Metabolites produced during growth in astrobiological analogs, such as those identified here, may provide potential agnostic biomarkers for life detection.

Keywords: Antarctica, *Shewanella*, iron-mediated metabolism, subglacial environments

References

- Sklute, E.C., Mikucki, J.A., Dyar, M.D., Lee, P.A., Livi, K.J. and Mitchell, S., 2022. A Multi-Technique Analysis of Surface Materials From Blood Falls, Antarctica. *Frontiers in Astronomy and Space Sciences*, 9, p.843174.
- Boles, B.W., Murdoch, R.W., Ohlsson, I. and Mikucki, J.A., 2021. Draft Genome Sequence of *Shewanella* sp. Strain BF02_Schw, Isolated from Blood Falls, a Feature Where Subglacial Brine Discharges to the Surface of Taylor Glacier, Antarctica. *Microbiology Resource Announcements*, 10(22), pp.10-1128.
- Bowman, J.P., McCammon, S.A., Nichols, D.S., Skerratt, J.H., Rea, S.M., Nichols, P.D. and McMeekin, T.A., 1997. *Shewanella gelidimarina* sp. nov. and *Shewanella frigidimarina* sp. nov., novel Antarctic species with the ability to produce eicosapentaenoic acid (20: 5ω3) and grow anaerobically by dissimilatory Fe (III) reduction. *International Journal of Systematic and Evolutionary Microbiology*, 47(4), pp.1040-1047.

* Presenting author. E-mail address: jmikucki@utk.edu (Jill A. Mikucki)

S2.P211 Effect of green-based extraction approaches on antioxidant activity and phenolic compounds of Oregano (*Origanum vulgare* ssp. *hirtum* (Link) Ietswaart)

Christina Panagiotidou, Maria Irakli, Elisavet Bouloumpasi, Paschalina Chatzopoulou*

Hellenic Agricultural Organization – Dimitra – Institute of Plant Breeding and Genetic Resources, Thessaloniki, Greece

Greek Oregano (*Origanum vulgare* ssp. *hirtum* (Link) Ietswaart) is a Mediterranean culinary herb, traditionally used in folk medicine, as far as in food, animal feed and in pharmaceutical industry. The species contain high quality essential oil and is rich source of phenolic compounds, known for their bioactivity, attributed to health-promoting properties; antioxidant, antibacterial, antifungal, etc. Nowadays, there is a growing interest for natural antioxidants that could be used as alternative to the synthetic ones, using at the same time green-based extraction approaches. The impact of microwave- assisted extraction (MAE), ultrasound-assisted extraction (UAE), and accelerated solvent extraction (ASE) on the total phenolics, total flavonoids and total antioxidant activity of oregano extracts was investigated. Dried and ground oregano extracts were subjected to extraction with 60% aqueous methanol. The antioxidant activity of oregano extracts was evaluated using ABTS, DPPH and FRAP assays. MAE proved the most effective extraction technique in exploring the bioactive compounds from oregano. Specifically, MAE extracts showed more total phenolics, total flavonoids and antioxidant activity on ABTS, DPPH and FRAP assays in comparison to ASE and UAE extracts. The HPLC/PDA/MS analyses revealed that rosmarinic acid, salvianolic acid B and carvacrol were the major phenolic compounds in all the obtained extracts, followed by flavonoid glycosides, with their content being highest in MAE extracts, obtained at 70°C, over 10 min. However, higher carvacrol content was favored by UAE and less by ASE. Summarizing, MAE could be employed as green and efficient method for producing oregano extracts rich in bioactive compounds, exhibiting strong antioxidant properties for developing nutraceuticals and pharmaceuticals.

Acknowledgments: This research has been co-financed by the Action “Innovation Investment Plans” of Central Macedonia, Operational Programme 2021-2027, (project code MIS 5179230)

Keywords: Oregano, green extraction, microwave, phenolics, LC-MS analysis

*Presenting author. E-mail address: pchatzopoulou@elgo.gr (Paschalina Chatzopoulou)

S2.P212 Enhancing polyphenol production in *Cistus albidus* L. through strategic mycorrhization and irrigation management: An off-season approach

Daniel Raus de Baviera^{a,*}, Enrique Barrajón-Catalán^{b,c}, Antonio Ruiz-Canales^a

^a Department of Engineering, Area of Agroforestry, Miguel Hernández University, 03312 Orihuela, Spain

^b Institute for Research, Development and Innovation in Health Biotechnology, Miguel Hernández University, 03202 Elche, Spain

^c Department of Pharmacy, Elche University Hospital-FISABIO, 03203 Elche, Spain

Polyphenols, valuable for their health benefits, face challenges in synthetic production, including low yields and toxicity issues (Ofosu et al., 2020). Mediterranean semideciduous shrubs, adapted to abiotic stress, are significant sources of these metabolites. *Cistus albidus* L., a species renowned for its medicinal attributes, thrives under the harsh conditions of the Mediterranean climate, producing a wide array of natural metabolites with minimal resource inputs (Raus de Baviera et al., 2023). Our study focused on enhancing *C. albidus*'s polyphenol production through *Glomus* sp. inoculation and controlled irrigation. We discovered that maintaining water availability at field capacity modulates the plant-mycorrhiza symbiosis so that *Glomus* sp. behaves different as expected, significantly increasing polyphenol and antioxidant levels compared to not inoculated plants. Especially during autumn, season with limited polyphenol production in *C. albidus*, we observed values over four times higher than the control, indicating that strategic water management and *Glomus* sp. inoculation can substantially boost polyphenol and antioxidant content in *C. albidus*. After the plants were exposed to drought stress, typical symptoms of mycorrhizal symbiosis returned. As observed by (Ortuño et al., 2018) mycorrhiza may improve the water status of *C. albidus*, allowing to devote more energy to secondary metabolism and the synthesis of phenolic compounds under well-watered conditions, rather than just focusing on survival. This approach, particularly effective during seasons with limited polyphenol production, highlights the potential of combining mycorrhization with irrigation strategies to enhance the productivity, presenting a viable method for optimizing the production of pharmacological compounds.

Keywords: polyphenols, *Cistus albidus* L., *Glomus* sp. inoculation, water availability

References

- Ofosu, F.K., Daliri, E.B.-M., Elahi, F., Chelliah, R., Lee, B.-H., & Oh, D.-H., 2020. New Insights on the Use of Polyphenols as Natural Preservatives and Their Emerging Safety Concerns. *Front. Sustain. Food Syst.* 4, 525810, 1-14. Raus de Baviera, D., Ruiz-Canales, A., Barrajón-Catalán, E., 2023. *Cistus albidus* L.-Review of a Traditional Mediterranean Medicinal Plant with Pharmacological Potential. *Plants*. 12, 2988, 1-43.
- Ortuño, M. F., Lorente, B., Hernández, J. A., Sánchez-Blanco, M. J. 2018. Mycorrhizal inoculation on compost substrate affects nutritional balance, water uptake and photosynthetic efficiency in *Cistus albidus* plants submitted to water stress. *Braz. J. Bot.* 41, 2, 299-310.

* Presenting author. E-mail address: daniel.raus@goumh.umh.es (Daniel Raus de Baviera)

S2.P213 The potential of chitosan from *Hermetia illucens* as a sustainable biopesticide

Joanita Nangobi^{a,*}, Nicholas Yiga^b

^a Department of Pharmacognosy, School of Pharmacy, 88Makerere University, Mitchell Hall side Rd Kampala, Uganda

^b ISP Uganda, St. Padre Pio Skills Training Centre, Sentema Rd Kampala, Uganda

The long-standing use of chemical pesticides by farmers has enhanced pest resistance while continuing to pose serious environmental risks. As a result, there has been an upsurge in research and development of biopesticide formulations. Farmers are searching for safer, more efficient ways to maintain their harvests while preserving the environment. The use of frass from the *Hermetia illucens* (Black soldier fly) as a potential biofertilizer and biopesticide for crops has previously been crudely reported. Frass is a waste product from the Black soldier fly farm and it is made up of various organic wastes including molten skins and pupal shells of the Black soldier flies from which chitin can be extracted. Chitosan, which is deacetylated chitin is an important commercial molecule with desirable properties and holds immense promise as a sustainable biopesticide. This study aimed at exploring the extraction and application of chitosan in pest management, addressing environmental concerns associated with conventional pesticides. The chitosan extraction process involved pretreatment of the skins and shells from black soldier flies, demineralization, deproteinization followed by the deacetylation of chitin to obtain chitosan while following rigorous quality control parameters, ensuring optimal yield and purity. The resulting chitosan exhibited notable antifungal, antibacterial and antipesticial properties specifically noted for *Colletotrichum sublineolum* (which causes anthracnose in sorghum), *Escherichia coli* (a common pathogenic bacteria and contaminant) and *Aleurodicus dispersus* (whiteflies that transmits begomo viruses which cause Tomato yellow leaf curl) respectively establishing its potential as a multifunctional biopesticide. Our investigations reveal the efficacy of chitosan against a range of agricultural pests. Furthermore, the use of *Hermetia illucens* frass for chitosan production aligns with sustainable practices. The larvae of *Hermetia illucens* thrive on organic waste, providing a dual benefit of waste management and biopesticide production. In addition, chitosan is as well a biodegradable molecule. This integrated approach contributes to a circular economy and reduces the environmental impact associated with conventional pesticide production making it a compelling choice for sustainable agriculture.

Keywords: *Hermetia illucens*, chitosan, biopesticide, sustainable

References

- Huang, W. C., Zhao, D., Xue, C., & Mao, X. (2022). An efficient method for chitin production from crab shells by a natural deep eutectic solvent. *Marine Life Science and Technology*, 4(3), 384–388.
- Kubayi, C. (2022). *Frass from black soldier flies as a valuable fertilizer and biopesticide for crops*.

* Presenting author. E-mail address: joanitanangobi@gmail.com (Nangobi Joanita)

S2.P214 Screening of Nettle Clones with the Ph Eur HPTLC method for identification testing

Evelyn Wolfram^{a,*}, Valentina Kremenovica^a, Samuel Peter^a, Marianne Leupin^b, Corinna Urmann^{cd}

^a Natural Products and Phytopharmacy Research Group, Department of Life Sciences, Zürich University of Applied Sciences, Switzerland, CH-8820 Wädenswil

^b Institute for Environment and Natural Resources, Department of Life Sciences, Zürich University of Applied Sciences, Switzerland, CH-8820 Wädenswil

^c Organic-analytical Chemistry, Weihenstephan-Triesdorf University of Applied Sciences, D-94315 Straubing, Germany

^d TUM Campus Straubing for Biotechnology and Sustainability, Technical University of Munich, D- 94315 Straubing, Germany

The study compares the health-related compounds in 20 clones of stinging nettle leaves (*Urtica dioica* L.) at different harvest times. The cultivars were dedicated for textile fibre production, the leaves are assessed for valorization. High performance thin layer chromatography (HPTLC) fingerprint comparison was used and commercial nettle leaves from a local pharmacy and collected in the wild from the Swiss mountains included. During the growth phase, the morphology of the nettles changed in size as well as in the colour of the leaves and stems. After drying at 40 °C, the nettle leaves were extracted with 70 % methanol (70:30 V/V) in an ultrasonic bath for 15 min. The HPTLC plates derivatized with anisaldehyde sulfuric acid (AS) and ferric chloride (FC) were compared to the PhEur Identity testing prescribed Natural product (NP/PEG) reagent derivatization in order to assess different natural product classes. The ubiquitous reference compounds rutin and chlorogenic acid were identified in every nettle examined, while hyperoside, quercetin, and the Ph Eur relevant scopoletin were not or only faintly detected in any nettle of the summer harvest. Assessment for Scopoletin, as a identity marker content was performed using LC-MS analysis. Complementary to the metabolic comparison, the HPTLC-DPPH radical scavenging assay showed that activity was higher with increasing growth of the plants. Overall a late harvest of the leaves is recommended to achieve the necessary secondary metabolite content for health related benefits.

Keywords: nettle leaves, fibre side stream valorization, HPTLC, DPPH, scopoletin

References:

EDQM, European Pharmacopeia, Edition 11.3, Strassbourg, January 2024

* Presenting author. E-mail address: wola@zhaw.ch (Evelyn Wolfram)

S2.P215 Creation of a medicinal plants collection and research consortium to catalyze drug discovery in San Diego and beyond: An update from 2022-2024

C. Benjamin Naman*

Department of Science and Conservation, San Diego Botanic Garden, Encinitas, California, USA

As many as 40% of all drugs found in modern pharmacies today are directly or indirectly derived from plants (USDA, 2024). Yet, the rate of plant-based drug discovery has declined in recent decades, in part due to lack of readily available medicinal plant collection resources. At the same time, the threat and reality of biodiversity loss has been overwhelmingly increasing. It is now nearly ten years since the Nagoya Protocol went into effect. Many drug discovery programs have yet to effectively implement the aims and scope of the Nagoya Protocol for enhancing the conservation and sustainable use of biodiversity or to recognize the value of traditional ecological knowledge of plants and their uses. The Department of Science and Conservation at San Diego Botanic Garden has undertaken a project initiative “Creation of a National Medicinal Plants Collection and Research Consortium to Catalyze Drug Discovery in San Diego and Beyond” that aims to demonstrate a new way forward for plant natural product drug discovery. As we have embarked on a journey to build a medicinal plants collection in the USA for the purposes of both plant conservation and scientific research, we attempted to proactively create equitable partnerships with meaningful societal contributions including plant conservation efforts, ecosystem restoration projects, and joint development of phytochemical knowledge that aims to bridge traditional and modern medical practices. This project accordingly includes partnerships with members or governments of indigenous communities (i.e. Native American Indians), which brings great opportunities along with operational challenges, and a paramount ethical responsibility. As a non-profit organization, we have created a consortium of research partners that includes other non-profits, academia, industry, and tribal governments to find ways to work together to improve outcomes and remove barriers from more traditional past research programs. We have developed early educational programming to engage community members from diverse backgrounds both in-person and online. Finally, we have undertaken multi-institutional natural product drug discovery and development projects studying two native medicinal plants, California yerba santa (*Eriodictyon californicum*) and California sagebrush (*Artemisia californica*). These plants were grown under varied sets of controlled conditions in our medicinal plant greenhouse and nursery areas to then measure and evaluate the modulation of genomic, transcriptomic, and metabolomic expression. In vitro pharmacological testing has been conducted by partner institutions. Examples from all of these activities will be presented in the conference.

Keywords: Native American ethnobotany, consortium, phytochemistry, *Eriodictyon*, *Artemisia*

References

USDA, U.S. Forest Service, 2024. Medicinal Botany. Accessed 5 April, 2024. <https://www.fs.usda.gov/wildflowers/ethnobotany/medicinal/index.shtml>.

* Presenting author. E-mail address: bnaman@sdbgarden.org (C. Benjamin Naman)

Regulation for the use of natural products in food and drugs

S2.P216 The cannabis/marijuana (*Cannabis sativa* L.) landscape in Africa: an overview of its cultivation and legal aspects

Godwin Anywar*, Esezah Kakudidi, Patience Tugume, Savina Asiimwe

Department of Plant Sciences, Microbiology & Biotechnology, Makerere University, Kampala, Uganda

Cannabis is the most widely used illicit substance globally. However, it remains an important crop in Africa and was originally cultivated by poor farmers who could not benefit from legal agriculture (Merz, 2018). Currently, African farmers produce enough cannabis to meet continental demand while exporting small quantities. We reviewed the literature on cannabis cultivation, legalization, farming, export, production, history, trade, medical and recreational effects of cannabis in Africa between 1975 to 2021. We searched databases such as Web of Science, Google Scholar, PubMed, Scopus for research papers, reports, and books. Also, we searched local newspapers and court rulings for relevant cannabis content. Although many countries in Europe and the Americas have liberalized controls of different aspects along the cannabis value chain. However, most African nations upheld pre- independence laws against cannabis in compliance with the 1925 International Opium Convention, although the legal terrain has and continues to evolve (Carrier and Klantschnig, 2018). Consequently, some African governments, including South Africa, the Democratic Republic of Congo (DRC), Uganda, Lesotho, Malawi, eSwatini, Zimbabwe, Rwanda, Kenya, and Egypt, have authorized limited cannabis farming. We recommend a more inclusive and regulated cannabis farming system so that local farmers benefit from its export while governments from a broader tax base. African governments need to invest more in cannabis research and development. This will pave the way for their citizens to benefit from the therapeutic effects of medical cannabis and value addition which can fetch higher returns than the raw cannabis materials.

Keywords: medicinal cannabis, marijuana, Africa, legalization, trade, farming

References

Merz, F., 2018. United Nations Office on Drugs and Crime: World Drug Report 2017. SIRIUS Z. Strat Anal. 2(1):85–86. Carrier, N., Klantschnig G., 2018. Quasilegality: khat, cannabis and Africa's drug laws. Third World Q. 39(2):350–365.

* Presenting author. E-mail address: godwinanywar@gmail.com (Godwin Anywar)

S2.P217 A comparative toxicity study depending on type of clinical use of the roots of *Asarum heterotropoides* var. *mandshuricum* (Maxim.) Kitag

Kang-Hyun Han^{a,b,*}, Seung Eui Min^{a,b}, Eun-Young Gu^a, Yong Bum Kim^a, Byung Sun Min^c

^a Department of Advanced Toxicology Research, Korea Institute of Toxicology, 141 Gajeong-ro, Yuseong-gu, Daejeon, 34114, Republic of Korea

^b Human and Environmental Toxicology, Korea National University of Science and Technology, 217 Gajeong-ro, Yuseong-gu, Daejeon, 34113, Republic of Korea

^c College of Pharmacy, Daegu Catholic University, Gyeongsan, Gyeongbuk 38430, Republic of Korea

Natural products are used for various purposes such as foods, health supplements and medicines. In the case of the safety assessment of natural products, it is determined based on known toxic substances, but may change depending on the manufacturing method, so there is a limit to simply evaluating them based on toxicity results of the natural product raw materials themselves. The root of *Asarum heterotropoides* var. *mandshuricum* (Maxim.) Kitag (AR) is widely used as a traditional herbal medicine mainly in East Asia. In this study, we presented toxicity comparison results for the main forms of use of AR root, powder and water extraction. Genotoxicity, single- and repeated-dose toxicity studies were conducted in accordance with the OECD testing guidelines. As a result of the study, toxicity changes including positive results in genotoxicity and adenoma in the liver were observed in the powder form of AR. However, no specific toxicity changes were observed in water extraction form of AR. As a result of HPLC analysis for the main components, methyl eugenol and safrole, known as toxic substances, were detected only in powder form. This means that the corresponding component is removed during the water extraction process of AR, and it was considered to be decreased a toxicity effect.



Fig. 1. Hepatocyte adenoma in the liver after administering 2000 mg/kg/day AR powder

Keywords: *Asarum heterotropoides* var. *mandshuricum*, acute toxicity, subchronic toxicity, genotoxicity, toxic components

Acknowledgments

This research was funded by grants from the Ministry of Food and Drug Safety in 2019 (1918MFDS386) and Korea Institute of Toxicology, Republic of Korea (1711195885).

* Presenting author. E-mail address: hankh@kitox.re.kr (Kang-Hyun Hana)

S2.P218 Determination of triterpenoids of the native *Centella asiatica* L. from the Jeju Island, South Korea

Jaehee Hyun*, Soyoung Shin, Wookyu Lee, Jaeuk Seo, Ockjin Paek, Jongwhan Kim, Seungtae Chung

National Center for Medicinal Resources Management, Herbal Medicine Research Division, National Institute of Food and Drug Safety Evaluation, Ministry of Food and Drug Safety, South Korea

Centella asiatica L. (CA) is a (sub)tropical herbal medicine belonging to the Apiaceae (Antoine N Nicolas et al., 2009). It is widely distributed throughout tropical and subtropical regions of Africa, Asia, Australia, and islands in the western Pacific Ocean (Nora E Gray et al., 2018). However, it has been found to have different total triterpenoids depending on the growing ecological environment (Boju Sun et al., 2020). In recent years, its extracts and bioactive components have been reported to have anti-inflammatory, anti-tumor, anti-oxidant, wound healing, cardioprotective and improving- memory effect (Changhe Wang et al. 2022).

In the present study, the aerial part of CA was collected native to Jeju Island, Republic of Korea with Voucher No. MFDS-C-8705, and its dried extract was quantified by high-performance thin-layer chromatography (HPTLC) and high-performance liquid chromatography (HPLC) focusing on four triterpenoids, asiaticoside, madecassoside, asiatic acid, and madecassic acid.

Analysis by HPTLC of the samples was performed on HPTLC aluminum precoated plate (60 F254) by using mobile phase dichloromethane, methanol, water (14:6:1 v/v/v) with 10% sulfuric acid. and we developed a HPLC simultaneous analysis of four triterpenoids in *C. asiatica*.

All the results obtained from in this study help to determine the characteristics and analysis of standardization of CA native to Jeju Island.

Keywords: *Centella asiatica* L., triterpenoids, HPTLC, HPLC, herbal medicine

References

- Antoine N Nicolas, Gregory M Plunkett. 2009. The demise of subfamily Hydrocotyloideae (Apiaceae) and the re- alignment of its genera across the entire order Apiales. Mol. Phylogenet. Evol. Volume 53, Issue 1, Pages 134-151.
- Nora E Gray, Armando Alcazar Magana, Parnian Lak , Kirsten M Wright, Joseph Quinn, Jan F Stevens, Claudia S Maier , Amala Soumyanath . 2018. *Centella asiatica* – Phytochemistry and mechanisms of neuroprotection and cognitive enhancement. Phytochem Rev. 17(1): 161–194.
- Boju Sun, Lili Wu, You Wu, Chengfei Zhang, Lingling Qin, Misa Hayashi, Maya Kudo, Ming Gao and Tonghua Liu. 2020. Therapeutic Potential of *Centella asiatica* and Its Triterpenes: A Review. Front Pharmacol. 11: 568032.
- Changhe Wang, Yanan Zhao, Ruomeng Yang, Haijing Liu. 2020. Simultaneous analysis of five triterpenes in *Centella asiatica* by high performance liquid chromatography with cyclodextrins as the mobile phase additives. *Scientific Reports* volume 10, Article number: 18577.
- Renju Kunjumon, Anil John Johnson, Sabulal Baby. 2022. *Centella asiatica*: Secondary metabolites, biological activities and biomass sources. Phytomedicine Plus. Volume 2, Issue 1, 100176.

* Presenting author. E-mail address: jaeheeh0508@korea.kr (Jaehee Hyun)

S2.P219 Streoselective inhibitory effects of ginsenosides on six uridine 5'-diphosphoglucuronosyl transferases in human liver microsomes

Bae-Gon Kang^{a,*}, Do-Hyeon Gwon^a, Hyunyoung Lee^a, Hyun-Ji Kim^a, So-Young Park^{a,b}, Taeho Lee^a, Kyung-Sik Song^a, Kwang-Hyeon Liu^{a,b}

¹ BK21 FOUR Community-Based Intelligent Novel Drug Discovery Education Unit, College of Pharmacy and Research Institute of Pharmaceutical Sciences, Kyungpook National University, Daegu 41566, Republic of Korea

² Mass Spectrometry Based Convergence Research Institute, Kyungpook National University, Daegu 41566, Republic of Korea

This study investigated the inhibitory potential of both protopanaxadiol (PPD)- and protopanaxatriol (PPT)-type ginsenosides (Fig. 1.) on the activities of six uridine 5'-diphospho- glucuronosyltransferase (UGT) isoforms (1A3, 1A4, 1A4, 1A6, 1A9, and 2B7) in pooled human liver microsomes (HLMs). Among the various active ingredients of ginseng, ginsenosides are the main chemicals responsible for the pharmacological effect of ginseng such as anticancer, antioxidant, and cholesterol-lowering activity. Ginsenosides inhibited the UGT isoform activities in a structure- dependent manner. For example, PPT-type ginsenosides showed minimal inhibition against the activity of the 6 UGT isoforms evaluated ($IC_{50} > 37.1 \mu M$). The activities of UGT2B7 and UGT1A9 were inhibited by the PPD-type ginsenoside having two sugar moieties at the carbon-3 position of the ginsenoside structure (e.g., ginsenoside Rg3) and PPD-type ginsenosides having two sugar moieties at the carbon-3 position of ginsenoside regardless of the presence or absence of the sugar group at carbon-20 position of ginsenosides (e.g., ginsenoside Rb1, Rb2, Rd, and Rg3), respectively. These results suggest that ginsenosides have a structure-specific inhibitory effect on UGTs in HLMs.




Fig. 1. Structure of (a) protopanaxadiol and (b) protopanaxatriol

Keywords: ginsenosides, inhibition, microsomes, uridine 5'-diphospho glucuronosyltransferase

References

- Kim D, Zheng YF, Min JS, Park JB, Bae SH, Yoon KD, Chin YW, Oh AE, Bae SK. 2016. In vitro stereoselective inhibition of ginsenosides toward UDP-glucuronosyltransferase (UGT) isoforms. *Toxicol Lett* 259:1-10.
- Fang ZZ, Cao YF, Hu CM, Hong M, Sun XY, Ge GB, Liu Y, Zhang YY, Yang L, Sun HZ. 2013. Structure-inhibition relationship of ginsenosides towards UDP-glucuronosyltransferases (UGTs). *Toxicol Appl Pharmacol* 267:149-154.
- Lee H, Heo JK, Lee GH, Park SY, Jang SN, Kim HJ, Kwon MJ, Song IS, Liu KH. 2019. Ginsenoside Rc is a new selective UGT1A9 inhibitor in human liver microsomes and recombinant human UGT isoforms. *Drug Metab Dispos* 47:1372- 1379.

* Presenting author. E-mail address: baegon0709@naver.com (Bae-Gon Kang)

S2.P220 Discrimination of herbal medicines originated from soy beans (*Glycine max* Merrill.) using high-performance thin-layer chromatography

Yena Kim^{a,b,*}, Chae Kyung Kim^{b,c}, Geonha Park^b, Min Ju Kim^{a,b}, Young Pyo Jang^{a,c}

^a Department of Integrated Drug Development and Natural Products, Graduate School, Kyung Hee University. Seoul, 02447, Republic of Korea

^b Division of Pharmacognosy, College of Pharmacy, Kyung Hee University. Seoul, 02447, Republic of Korea

^c Department of Oriental Pharmaceutical Sciences, College of Pharmacy, Kyung Hee University, Seoul 02447, Republic of Korea

Glycine Semen Nigra (GN), Glycine Semen Preparata (GP) and Glycine Semen Germinatum (GG) are herbal medicines derived from *Glycine max* Merrill, exhibiting distinct clinical properties due to varied processing methods. GN is recognized for its diuretic and detoxifying effects, while GP alleviates symptoms such as headaches, chest tightness, and insomnia resulting from colds. GG is employed for summer colds, addressing abdominal pain and dysuria by reducing fever and inducing diuresis. This study focuses on the development of a High-Performance Thin-Layer Chromatography (HPTLC) method for the discrimination of GN, GP, and GG. The optimal mobile phase comprised a solvent mixture of dichloromethane, methanol, water, and glacial acetic acid in a ratio of 10:4:1:0.5 (v/v/v/v). HPTLC results revealed a consistent blue spot at an R_f value of 0.6, identified as genistin. (Kim, J. A. et al. 2007). Additionally, spots at R_f values 0.35, 0.55, and 0.65 distinguished the three groups. The identification of compounds within these spots, through TLC-Mass Spectrometry- interface, aimed to correlate them with the specific activities of each herbal medicine. In conclusion, our developed HPTLC method facilitates the simultaneous differentiation of three herbal medicines from the same species, *Glycine Max* Merrill. This approach is expected to contribute to the standardization of herbal medicines through a rapid and cost-effective method.

Keywords: high-performance thin-layer chromatography (HPTLC), *Glycine max* L., glycine semen nigra, glycine semen preparata, glycine semen germinatum

References

Kim, J. A., & Chung, I. M., 2007. Change in isoflavone concentration of soybean (*Glycine max* L.) seeds at different growth stages. Journal of the Science of Food and Agriculture, 87(3), 496-503.

* Presenting author. E-mail address: kimyena2317@naver.com (Yena Kim)

S2.P221 Analytical Quality by Design(AQbD) methodology approach for simultaneous quantification of Hesperidin and Magnolol in Daekumeum-ja using HPLC-PDA

Kyung Ho Lee^{a,b,*}, Sejin Ku^{b,c}, Geonha Park^{b,d}, Minsik Choi^{b,c}, Young Pyo Jang^{a-d}

^a Department of Integrated Drug Development and Natural Products, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

^b Department of Pharmacognosy, College of Pharmacy, Kyung Hee University, Seoul 02447, Republic of Korea

^c Department of Biomedical and Pharmaceutical Sciences, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

^d Department of Life and Nanopharmaceutical Sciences, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

Daekumeum-ja (DKJ) is known for its efficacy in preventing alcohol-induced inflammatory mucosal damage and cognitive impairments (J.J.Kim et al., 2010). To standardize the quality of DKJ, an HPLC-PDA analysis method was developed for the simultaneous quantification of two major components, hesperidin and magnolol. The robustness of this method was improved with the Analytical-Quality-by-Design strategy (AQbD), enhancing efficiency in both time and cost. Initially, a risk assessment was conducted to identify the Critical-Analytical-Procedure-Parameters (CAPPs), such as the ratio of the organic mobile phase and the column oven temperature. To evaluate the impact of CAPPs on Critical-Analytical-Procedure-Attributes (CAPA), peak tailing and resolution, a Design- of-Experiments (DOE) approach using Central-Composite-Design (CCD) was implemented. The optimal chromatographic analysis conditions were established for optimal column oven temperature, detection wavelength of 290 nm, flow rate, and mobile phase composition and gradient condition. In alignment with the International Conference on Harmonization (ICH) guideline, the developed method underwent validation. This study conclusively affirmed the capability for robust and reliable simultaneous quantitative method of DKJ components, thereby proving the successful development of the method.

Keywords: daekumeum-ja, HPLC-PDA, simultaneous quantification, analytical quality by design (AQbD), central composite design (CCD)

References

Kim, Jeong-Ja, et al., 2010. Preventive effects of Daekumeumja on fatty degeneration of liver and immunosuppression induced by alcohol. Korean journal of oriental medicine 16(2), 167-179

* Presenting author. E-mail address: ehrrh500@naver.com (Kyung Ho Lee)

S2.P222 Discrimination between similar herbal medicines of Campanulaceae by high-performance thin layer chromatography (HPTLC)

Won Sok Ha^{a,b,*}, Hyung Yeon Hwang^{b,c}, Roun Lee^{b,c}, Minsik Choi^{c,d}, Geonha Park^b, Young Pyo Jang^{a-d}

^a Department of Integrated Drug Development and Natural Products, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

^b Division of Pharmacognosy, College of Pharmacy, Kyung Hee University, Seoul 02447, Republic of Korea

^c Department of Oriental Pharmaceutical Sciences, College of Pharmacy, Kyung Hee University, Seoul 02447, Republic of Korea

^d Department of Biomedical and Pharmaceutical Sciences, Graduate School, Kyung Hee University, Seoul, 02447, Republic of Korea

Adenophorae Radix (AR), Adenophorae Remotiflori Radix (ARR), and *Codonopsis lanceolata* root (CL), members of the Campanulaceae family, face misidentification due to their morphological similarities, leading to confusion regarding their appropriate utilization in the herbal medicine markets in Korea (Park, E. S., et al., 2017). In response, this study aimed to establish a rapid discrimination method utilizing High-Performance Thin Layer Chromatography (HPTLC). 16 as AR, 7 as ARR, 10 as CL, and 7 samples distributed under unclear names were purchased from the herbal markets. Analysis of components identified in previous research and samples was conducted under the same HPTLC conditions. Results revealed two key components for classification: β -sitosterol with a retardation factor (Rf) value of 0.41 and α -spinasterol with an Rf value of 0.37. Distinct tendencies were confirmed, such as the exclusive presence of the β -sitosterol band in AR, exclusive α -spinasterol in CL, and both components in ARR. Furthermore, AR was discerned by the absence of a blue band with an Rf value of 0.61. Based on this classification, the samples were categorized into three groups: 17 as AR, 5 as ARR, and 18 as CL, leading to the correction of the labeling errors in the origin of 40 samples. Consequently, this study has successfully devised a straightforward and efficient analytical method to differentiate between AR, ARR, and CL, addressing the prevalent issue of misidentification. The implementation of this method is anticipated to play a pivotal role in enhancing the standardization of herbal medicine quality.

Keywords: *Adenophorae Radix*, *Adenophorae Remotiflori Radix*, *Codonopsis lanceolata* root, Campanulaceae, HPTLC

References

Park, E. S., Lee, S. R., Joeng, J. M., Song, M. G., Yoon, J. H., & Ju, Y. S., 2017. Discriminative criteria of *Adenophorae Radix*, *Codonopsis Lanceolatae Radix* and *Adenophorae Remotiflori Radix*. *The Korea Journal of Herbology*, 32(2), 33- 40.

* Presenting author. E-mail address: rmdwjd0220@khu.ac.kr (Won Sok Ha)

S2.P223 Quantification of saponin-rich plants mixture in complete feed with UHPLC-MS/MS using a “one-point” calibration

Maxime Le Bot^{a,b,*}, Mohammed el Amine Benarbia^{a,b}, David Guilet^{b,c}

^a NOR-FEED SAS, 3 rue Amédéo Avogadro, 49070 Beaucouzé, France

^b Joint Lab ANR FeedInTech (FIT: SONAS/Nor-Feed)

^c University of Angers, EA 921 SONAS/SFR 4207 QUASAV, 42 rue Georges Morel, 4970 Beaucouzé, France

Quantification of additives in complete feeds is necessary to estimate the rate of incorporation and homogeneity of the feed, which guarantees the effectiveness of the additive. However, for botanical compounds this remains a challenge due to the complexity of the matrix and the level of incorporation, requiring sensitive and specific methods.

A specific and sensitive method for the quantification of saponin-rich mixtures in complete feed, based on UHPLC-MS/MS, was developed. The additive studied was a natural commercial feed additive (Norponin® XO2-300 Nor-Feed, France) used in animal nutrition for the management of coccidiosis. A mixture consisting of a standardized *Yucca schidigera* and *Trigonella foenum-graecum*, in which protodioscin has been identified and used as a phytomarker for the quantification of the additive at a level of 300 ppm in complete feeds.

Quantification of the sample is carried out with an optimized sample preparation, an internal standard and with a “one-point” calibration. Using the pure additive for calibration permits the direct determination of the additive content in the sample by integrating the area of the phytomarker. The use of an internal standard with a structure close to that of the phytomarker allowed us to overcome the problems of recovery and matrix effects encountered in sample preparation and UHPLC-MS/MS. The analysis method, which is rapid and suitable for routine use, has been validated in-house (Thomson et al., 2002). This method meets all the prerequisites for use by European authorities as part of the registration of a food additive (FEEDAP, 2012).

Keywords: saponins, additive, complete feed, coccidiosis, UHPLC-MS/MS

References

Thompson M, Ellison SLR, Wood R. Harmonized guidelines for single-laboratory validation of methods of analysis (IUPAC Technical Report). Pure Appl Chem 2002; 74: 835–855

FEEDAP. Guidance for the preparation of dossiers for zootechnical additives: Guidance on zootechnical additives. EFSA J 2012; 10: 2536

* Presenting author. E-mail address: maxime.lebot@norfeed.net (Maxime Le Bot)

S2.P224 Interactions of ivy (*Hedera helix*) with other herbs or medicines: a systematic review

Bader A. Al-Suliman*, Faisal A. Al-Madhi, Osama H. Alshehri, Saeed Al-Awadh

Department of herbal evaluation, Saudi food and drug authority, kingdom of Saudi Arabia

Cough is a predominant medical condition and general practitioners are regularly consulted by patients to treat it. Ivy (*Hedera helix* L., Araliaceae) is commonly used for respiratory disorders. It includes saponins which exhibit bronchodilator, mucolytic, spasmolytic, and antibacterial properties. Although it is widely used, the potential interactions with other herbs or medicines are not fully investigated. Through a systematic review of the literature, this study examines the interactions of ivy with other herbs and medicines.

This review utilized multiple databases. Initially, a systematic literature search was performed in PubMed, Embase, and Cochrane Library. Another search was in the databases of several regulatory agencies with excellent records regarding herbal products interactions. These agencies are mainly the European Medicines Agency (EMA) and Health Canada. The search terms used were “ivy”, “*Hedera helix*”, “drug interactions”, “herb-drug interactions”, and “phytochemical interactions”. The search covers the period between January 1990 and December 2021.

In this review, 23 studies were included. These studies (randomized controlled trials, observational studies, case reports and *in vivo* and *in vitro* studies) investigated the interactions of ivy with various herbs and medicines, including anticoagulants, bronchodilators, and corticosteroids. The majority reported no significant interactions between ivy and the other herbs or medicines studied. However, some studies reported potential interactions, such as increased bleeding risk when ivy was used with anticoagulants or antiplatelet agents. Overall, the available evidence suggests that ivy can be used safely with other herbs or medicines, although caution should be used when using ivy with anticoagulants or antiplatelets.

Keywords: ivy, *Hedera helix*, interactions

References

- Holzinger F, Chenot JF. Systematic review of clinical trials assessing the effectiveness of ivy leaf (*Hedera helix*) for acute upper respiratory tract infections. *eCAM*. 2011;2011:382789.
- Irwin RS, Madison JM. The diagnosis and treatment of cough. *N Engl J Med*. 2000;343(23):1715-1721.
- Sieben A, Prenner L, Sorkalla T, et al. Alpha-hederin, but not hederacoside C and hederagenin from *Hedera helix*, affects the binding behavior, dynamics, and regulation of beta 2-adrenergic receptors. *Biochemistry*. 2009;48(15):3477-3482.

* Presenting author. E-mail address: Basuliman@sFDA.gov.sa (Bader A. Alsuliman)

S2.P225 Brazilian regulation for food supplements containing bioactive substances from plant extracts

Andressa Blainski^{a,*}, Eliana Martino Bufaino^b

^a Finzelberg GmbH & Co. KG, Koblenzer Straße 48-56, 56626 Andernach, Germany

^b Finzelberg Ativos Naturais Ltda., Rua do Amor Perfeito 101, 06713-290 Cotia-SP, Brazil

The interest in food supplements is increasing in Brazil. Although the regulation for food supplements has been updated by Brazilian Health Regulatory Agency (ANVISA) in recent years, the regulated sector suffers from challenging requirements for providing compliant products.

Below, we present the latest regulatory changes related to the use of bioactive plant extracts in food supplements and highlight the challenges in approving new ingredients in Brazil.

The Regulation for approving novel food ingredients has existed since 1999 (RES16/1999), however, the food supplement category was not created before 2018 (RDC243/2018). It provided a clear application procedure for such ingredients. Moreover, RDC839/2023 will replace RES16/1999 from March 2024 on, while other older regulations will remain valid, which results in a complex scenario. RDC839/2023 determines the safety requirements for e.g. bioavailability and toxicokinetic in food ingredients as well as in their products of degradation. The chemical characterisation should be as complete as possible. Moreover, it is not clear whether the use of extracts from some species in food supplements will be accepted if the same species have also been used in herbal medicinal products. Insufficient or missing information must be justified, and may or may not be accepted by ANVISA. Lastly, only plant extracts that have been approved by ANVISA can be used in supplements. The approved ingredients are listed in IN28/2018 and its updates. ANVISA's decision normally takes around 2 years after application. Publishing the summarised manufacturing process and specifications will be a disadvantage for own developed ingredients.

Keywords: Brazilian regulation, food supplement, plant extract, bioactive substances

References

RDC 16/1999 – Regulamento referente a procedimentos para registro de alimentos e ou novos ingredientes. Brazil, 30 April 1999.

RDC 243/2018 – Dispõe sobre os requisitos sanitários dos suplementos alimentares. Brazil, 26 July 2018.

RDC 839/2023 – Dispõe sobre a comprovação de segurança e a autorização de uso de novos alimentos e novos ingredientes. Brazil 14 December 2023.

IN 28/2018 – Estabelece as listas de constituintes, de limites de uso, de alegações e de rotulagem complementar dos suplementos alimentares. Brazil, 26 July 2018.

* Presenting author. E-mail address: andressa.blainski@finzelberg.com (Andressa Blainski, PhD)

S2.P226 Chemical profile of natural musk (*Moschus moschiferus*) by GC-MS/MS

Seong Mi Lee, Jisu Ha^a, Eun Hye Ko, Ji-Young Park, Seh Wook Oh, Jinhee Hwang*

Herbal Medicine Research Division, National Institute of Food and Drug Safety Evaluation, Ministry of Food and Drug Safety, Chungcheongbuk-do 28159, Korea

Since 'natural musk' is one of the most expensive herbal medicine, it's essential to develop an adequate quality control method in order to prevent forgery and maintain its quality in the view of regulatory authorities. Analytical methods using indicator compounds have been reported, but with limitations that make it difficult to reflect the overall chemical characteristics of musk. In this study, we presented a standardized chemical profile, which is expected to be utilized as a basis for the development of quality evaluation methods based on chemical similarity. Musk samples were collected without removing the abdominal connective tissue from *Moschus moschiferus* of different areas and ages. Twelve standard samples were obtained by confirming that no traces of artificial manipulation were found through genetic analysis and organoleptic tests. Samples were analyzed using GC-MS/MS (scan mode), which can efficiently identify the constituents of musk. Seven peaks that were detected at 1% area of the total peak area or more, were selected as the standard chemical profile. The sum of the areas of the seven peaks was more than 75% of the total peak area and the relative standard deviation (% RSD) of the relative retention time [reference: IS (methandienone)] of each peak was consistent at less than 5%. Three peaks were identified as etiocholanolone, dihydroandrosterone, and epicholesterol by comparing retention times and mass spectra with commercial standards. Among the seven peaks in the standardized chemical profile, four peaks were not individually identified, but they are commonly detected peaks that accounts for 45% of the total area and are judged to be very important in reflecting the overall chemical properties of musk. Also, the relative standard deviation (% RSD) of the relative peak area of each peak [reference: IS (methandienone)] was 25% or less, indicating that there is a difference in the composition of each individual musk.




Fig. 1. Chromatogram of a standardized chemical profile for 'natural musk'. (1) unknown 1, (2) unknown 2, (3) etiocholanol one, (4) unknown 3, (5) dihydroandrosterone, (6, I.S.) methandienone, (7) unknown 4, (8) epicholesterol,

Keywords: Natural musk(*Moschus moschiferus*), Chemical fingerprint analysis, Herbal medicine

References

The Korean Herbal Pharmacopoeia 6th edition, 2021

Jin X. -f., Lu Y.-h., Wei D.-z., Wang Z.-t., 2008. Chemical fingerprint and quantitative analysis of *Salvia plebeia* R. Br. by high-performance liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis*. 48, 100-104

Chiriac E. R., Chitescu C. L., Geană E. I., Grid C. E., Socoteanu R. P., Boscencu R., 2021. Advanced analytical approaches for the analysis of polyphenols in plants matrices-A review. *Separations*. 8, 65

Wang Q., Yu X., Sun L., Tian R., He H., Wang S., Ma S., 2021. Fingerprint analysis of phenolic acid extract of *Salvia miltiorrhiza* by digital reference standard analyzer with one or two reference standards. *Chinese Medicine*, 16:8

Acknowledgements

This research was supported by a grant (22201MFDS144) from Ministry of Food and Drug Safety in 2024.

* Presenting author. E-mail address: rain01@korea.kr (Jinhee Hwang)

S2.P227 Advances in the Brazilian Regulation for the use of plant extracts in medicines

Eliana Martino Bufaino^{a,*}, Andressa Blainski^b, Cynthia Suarez^b

^a *Finzelberg Ativos Naturais Ltda., Rua do Amor Perfeito 101, 06713-290 Cotia-SP, Brazil*

^b *Finzelberg GmbH & Co. KG, Koblenzer Straße 48-56, 56626 Andernach, Germany*

The use of herbal medicines dates back to ancient times and is now increasing globally. Street markets in Brazil offer plants for any treatment, but the regulated sector must meet requirements to register herbal medicines.

We aim to present the regulatory advances related to herbal medicines, compare them with European Regulations, and highlight the main difficulties in approving new products in Brazil.

Herbal medicinal products are separately regulated and different from other medicinal products. As in Europe, they are distinct categories related to traditional use or scientific evidence. ANVISA (Brazilian Health Regulatory Agency) is updating the legislation regarding herbal medicines to harmonise with European concepts (DOU95/2021; Port1.409/2023).

The current RDC26/2014 and IN02/2014 focus on the chemical markers and do not take into account the classification into standardised, quantified and other extracts. On the other hand, HMPC- monographs are accepted for a simplified process, and many registered herbal active ingredients come from the European tradition. The harmonisation of concepts will make many aspects of quality and pharmacological interpretations very similar to those in Europe.

However, there are still challenges in terms of distinguishing the concepts and characteristics that differentiate herbal medicines from those that are synthetic. This has led to a discussion that has now lasted for more than 6 years. This period of uncertainty, coupled with the lengthy registration process, discourages the development of new herbal medicines and slows down the market potential.

Regulatory harmonisation bodes well for a promising development of herbal medicines in Brazil in the near future.

Keywords: herbal medicines, plant extracts, ANVISA

References

RDC 26/2014 – Determines about registration of herbal medicines and the registration and notification of traditional herbal products. Brazil, May 13, 2014.

IN 02/2014 – Publishes the “List of herbal Medicines with simplified registration” and the “List of Traditional Herbal Products with simplified registration”, Brazil May, 13 2014.

DOU 95/2021 – Approves ANVISA's Regulatory Agenda for the 2021-2023 three-year period. Item 8.25 – Review of regulatory aspects related to the production and quality control of herbal medicines (IN 4/2024 and specific parts of RDC 26/2014). Brazil, 21 May 2021.

Port 1.409/2023 – Approves ANVISA's Regulatory Agenda for 2024-2025. Item 8.2 – Review of regulatory aspects related to the production and quality control of herbal medicines (IN 4/2024 and specific parts of RDC 26/2014). Brazil, 15 December 2023.

* Presenting author. E-mail address: eliana.bufaino@finzelberg.com.br (Eliana Martino Bufainoa)

S2.P228 Comparing the volatilome of Spanish-style and natural cv. Chalkidiki green table olives at industrial scale

Panagiota Alvanoudi^a, Alexandros Nakas^{b,c,*}, Andreana N. Assimopoulou^{b,c}, Fani Th. Mantzouridou^{a,c}

^a *Laboratory of Food Chemistry and Technology, School of Chemistry, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece*

^b *Laboratory of Organic Chemistry, School of Chemical Engineering, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece*

^c *Natural Products Research Center of Excellence (NatPro-AUTH), Center for Interdisciplinary Research and Innovation (CIRI-AUTH), 57001 Thessaloniki, Greece*

Table olives are one of the most popular fermented foods, with great nutritional and sensory characteristics. For the preparation of cv. Chalkidiki green table olives, Spanish-style processing is almost exclusively used despite diminishing the content in health-promoting compounds and increasing the content in salt of the edible part (Mastralexi et al., 2019). On the contrary, the natural processing of olives involves minimal chemical input and retains better the nutritional value of olive drupes (Conte et al., 2020). These different practices also shape the fermenting microbial communities that are implemented in the quality attributes of the products (Ruiz-Barba et al., 2023). Volatilomics approach has been used to identify specific molecules, originating from the microbial metabolism taking place on table olive fermentation and affecting table olives' aroma and flavor (Alvanoudi et al., 2023).

The present study aims to shed light on the compositional characteristics of volatile organic compounds (VOCs) of Spanish-style and natural cv. Chalkidiki green table olives at industrial scale. VOCs were determined using headspace SPME-GC-MS. Microbiological analyses and sensory evaluation of the final products were also conducted via appropriate methods. More than 50 VOCs were detected (acids, alcohols, aldehydes, esters, phenols, etc.) and the final products were differentiated depending on the process applied. Furthermore, results from the microbiological and sensory quality assessment of the products showed that all of them were safe and organoleptically acceptable. Consumers' preference for the odor attribute found to be significantly different between the samples. The Greek table olive industry may benefit from the current findings.

Keywords: Spanish-style green olives, natural olives, volatile organic compounds, headspace SPME- GC-MS, Chalkidiki cultivar

References

- Alvanoudi, P., Ordoudi, S.A., Nakas, A., Assimopoulou, A.N., Mantzouridou, F.T., 2023. Brine Volatilome Changes Along the Spontaneous Fermentation of Spanish-Style cv. Chalkidiki Green Olives Under High and Low NaCl Conditions. *Food Bioprocess Technol.* 1-17.
- Conte, P., Fadda, C., Del Caro, A., Urgeghe, P.P., Piga, A., 2020. Table Olives: An Overview on Effects of Processing on Nutritional and Sensory Quality. *Foods*. 9, 514.
- Mastralexi, A., Mantzouridou, F.T., Tsimidou, M.Z., 2019. Evolution of Safety and Other Quality Parameters of the Greek PDO Table Olives "Prasines Elies Chalkidikis" During Industrial Scale Processing and Storage. *Eur. J. Lipid Sci. Technol.* 121, 1–13.
- Ruiz-Barba, J.L., Sánchez, A.H., López-López, A., Cortés-Delgado, A., Montañó, A., 2023. Microbial Community and Volatilome Changes in Brines along the Spontaneous Fermentation of Spanish-Style and Natural-Style Green Table Olives (Manzanilla Cultivar). *Food Microbiol.* 113.

* Presenting author. E-mail address: anakas@cheng.auth.gr (Alexandros Nakas)

S2.P229 Addressing regulatory challenges and ensuring safety in the e-commerce of herbal remedies in Mexico

José-Luis Echevarría-Becerra^{a,*}, Guadalupe Yáñez-Ibarra^b, Ávila-Villarreal Gabriela^{b,c}

^a Instituto Tecnológico de Monterrey, Av. Gral Ramón Corona No 2514, Colonia Nuevo México, 45201 Zapopan, Jalisco

^b Centro Nayarita de Innovación y Transferencia de Tecnología, A.C. Unidad Especializada en I+D+i en Calidad de Alimentos y productos Naturales. Universidad Autónoma de Nayarit. 63000. Tepic, Nayarit, México

^c Unidad Académica de Ciencias Químico Biológico y Farmacéuticas-Universidad Autónoma de Nayarit. Ciudad de la Cultura s/n. 63000. Tepic, Nayarit, México

In Mexico, plant-based products are classified as herbal medicines, herbal remedies (HR), and food supplements. Regulatory scrutiny focuses solely on herbal medicines, complicating oversight of remedies (Rojas et al., 2022). Ambiguous standards contribute to quality issues, such as mislabeling and adulteration, posing risks to safety and public health (Biesterbos et al., 2019). Online purchases through e-commerce platforms worsen the situation by lacking essential information, such as plant identity and origin (Alwhaibi et al., 2021). Producers frequently exploit legal loopholes to prioritize economic gains over rigorous studies.

This study aimed to evaluate HR obtained from physical health food stores and online platforms, adhering to Mexican regulations. Three HR were purchased nationwide: dry plants (MV-A, MV-B, MV-S), and infusion drops (ID-A, ID-B, ID-S) online. Compliance with NOM-072-SSA1-2012 for labeling and General Methods of Analysis for foreign matter, botanical identification, and thin-layer chromatography (TLC) following the guidelines of the Mexican Herbal Pharmacopoeia (FHEUM, for its acronym in Spanish).

Results revealed labeling discrepancies, with MVs lacking labels and IDs having deficient ones. MV-S samples exceeded the 2 % foreign matter limit, while the other samples complied. Only two MV-B samples did not match the specified species, affecting their macroscopic characters and chromatographic profiles. TLC analysis of IDs showed lower concentrations, indicating dilution but consistent species identification.

In conclusion, the analyzed samples do not fully adhere to national regulations, posing a significant issue as they are distributed nationwide in both physical and online stores.

Keywords: e-commerce, general methods of analysis, Herbal Mexican Pharmacopoeia, herbal remedies, safety

References

- Alwhaibi, M., Asser, W. M., Al Aloola, N., Alsalem, N., Almomen, A., & Alhawassi, T. M. 2021. Evaluating the frequency, consumers' motivation and perception of online medicinal, herbal, and health products purchase safety in Saudi Arabia. *Saudi Pharm J.* 29(2), 166–172.
- Biesterbos, J. W. H., Sijm, D. T. H. M., van Dam, R., & Mol, H. G. J. 2019. A health risk for consumers: the presence of adulterated food supplements in the Netherlands. *Food Addit Contam. - Part A*, 36(9), 1273–1288.
- Ministry of Health, 2012. Mexican Official Standard NOM-072-SSA1-2012, labeling of drug products and herbal remedies. Official Journal of the Federation, Mexico.
- Ministry of Health, 2021. Herbal Pharmacopoeia of the United Mexican States, 3rd ed.
- Rojas, P., Jung-Cook, H., Ruiz-Sánchez, E., Rojas-Tomé, I. S., Rojas, C., López-Ramírez, A. M., & Reséndiz-Albor, A.
- A. 2022. Historical Aspects of Herbal Use and Comparison of Current Regulations of Herbal Products between Mexico, Canada and the United States of America. *Int J Environ Res Public Health*, 19(23), 15690.

* Presenting author. E-mail address: joseechevabec@gmail.com (José-Luis Echevarría-Becerra)

S2.P230 Questions for consumer reporting on risks and benefits from herbal medicine usage: A preliminary international Delphi study

Peerapong Meesomperm^a, Chuenjid Kongkaew^{b,*}

^a Medical Division of General Support Office, Armed Forces Development Command, Bangkok, Thailand

^b Centre for Safety and Quality in Health, Department of Pharmacy Practice, Faculty of Pharmaceutical Sciences, Naresuan University, Thailand

Information about rational uses of allopathic drugs exists in abundance, yet global primary healthcare continues to make use of herbal medicines (HMs) for which data are sparse. We began to address this issue by validating a questionnaire on consumer experiences.

Thirty-nine questions were constructed from existing pharmacovigilance (PV) reporting forms that could ask HM consumers about 5-domains: reporter information, consumer's health status, the HMs being consumed, perceived HM benefits, and adverse experiences. Experts in PV and HMs identified from the International Society of Pharmacovigilance (ISoP) and online resources were recruited into a 2-round Delphi study to score importance of questions on herbal medicine usage.

The 16 experts were recruited with ~20 yr experience in PV or HMs. Both rounds had 100% response rates. Twenty-one consensus questions from 39 questions were highly important. Most other questions were scored as moderately important. Experts made more than 400 comments about reasons for scores, use of the reported data, and recommendations for effectively reporting.

In conclusion, the international Delphi indicated important questions for consumers to report HM experiences. These needed formatting to help reporters to easily provide structured and reliable data and could be implemented in countries lacking HM reporting. The study demonstrated the value of small cohorts of experienced panelists who freely comment.

Keywords: herbal medicines, consumers, spontaneous reporting system, pharmacovigilance, Delphi technique

References

- Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in pharmacology*. 2014;4:177 DOI: 10.3389/fphar.2013.00177.3.
- Shaw D, Graeme L, Pierre D, Elizabeth W, Kelvin C. Pharmacovigilance of herbal medicine. *Journal of ethnopharmacology*. 2012;140(3):513-8 DOI: 10.1016/j.jep.2012.01.051.
- WHO. Consumer reporting of adverse drug reactions: general policy issues. 2000.

* Presenting author. E-mail address: chuenjiddk@nu.ac.th (Chuenjid Kongkaew)

S2.P231 Analysis of the chromatographic profile and *in vitro* antiurolytic activity of species of the genus *Phyllanthus* (Phyllanthaceae), endemic and non-endemic of the flora of Santa Catarina

Adrielli Tenfen Voltolini^{a,*}, Lucas Natã Pereira^a, Gabrieli Zolett^b, Rita de Cássia Melo Vilhena de Andrade Fonseca da Silva^a, Priscila de Souza^a, Valdir Cechinel Filho^a, Luiz Adriano Funez^c, Luiz Carlos Klein Junior^{a,b}

^a Postgraduate Program in Pharmaceutical Sciences - Universidade do Vale do Itajaí (PPGCF - UNIVALI), Itajaí, SC, Brazil

^b Undergraduate course in Pharmacy at the School of Health Sciences (ECS) at UNIVALI, Itajaí, SC, Brazil

^c Herbário Barbosa Rodrigues – UNIVALI, Itajaí, SC, Brazil

Phyllanthus spp. popularly known as “stone breaker,” is used in folk medicine for kidney stones. However, the interchangeability between different species of the genus is described in reports of popular use, also stimulated by the difficulty of botanical differentiation, especially by laypeople. Therefore, the objective of this work aims to evaluate the chemical profile of *Phyllanthus* species and the antiurolytic activity in an *in vitro* model. For this, the aerial parts of *Phyllanthus* species were studied, including *P. niruri*, *P. tenellus*, *P. urinaria*, *P. hyssopifolioides*, and two recently described species endemic to the State: *P. eremitus*, and *P. timboensis*. The 70% ethanolic extracts were obtained by maceration from dried plant material. For subsequent analyses, the extracts were freeze-dried. The chromatographic profiles were obtained by High Performance Liquid Chromatography coupled to Diode Array (HPLC-DAD). The antiurolytic activity was carried out using the model described by Zanovello et al., 2021. The evaluation of the chromatographic profiles obtained showed different chemical compositions between the species. By comparing the profiles with standards (gallic acid, corilagin, luteolin and quercetin), it was possible to identify quercetin and luteolin in *P. tenellus*. Gallic acid was found in *P. niruri* and *P. tenellus*, corroborating with Brazilian Pharmacopeia (BF) Regarding the prediction of antiurolytic activity measured *in vitro*, the species *P. niruri*, *P. tenellus*, *P. urinaria*, *P. hyssopifolioides* significantly decreased ($p < 0.001$) the number of calcium oxalate crystals at concentrations of 0.1 and 1.0 mg/mL, respectively, relative to the negative control.

Keywords: Stonebreaker, antiurolytic activity, chromatographic profile

References

ANVISA. Agência Nacional de Vigilância Sanitária. Farmacopeia Brasileira. 6. ed. Brasília, 2019. v.2.

Brasil. Ministério da Saúde, Secretaria de Ciência, Tecnologia e Insumos Estratégicos. Departamento de Assistência Farmacêutica. RENISUS – Relação Nacional de Plantas Medicinais de Interesse ao SUS. DAF/SCTI/MS – RENISUS – fev/2009. Disponível em <https://www.gov.br/saude/pt-r/composicao/sectics/daf/pnppmf/ppnppmf/arquivos/2017/renisus1.pdf>

Zanovello, M.; Mariano, L.N.B.; Cechinel-Zanchett, C.C.; Boeing, T.; Tazinaffo, G.C.; Silva, L.M.; Silva, D.B.; Gasparotto Junior, A; Souza, P. *Tagetes erecta* L. flowers, a medicinal plant traditionally used to promote diuresis, induced diuretic and natriuretic effects in normotensive and hypertensive rats. *Journal of Ethnopharmacology*, v. 279, 114393, 2021.

* Presenting author. E-mail address: adrielli.tenfen@gmail.com (Adrielli Tenfen Voltolini Univali)

S2.P232 Hot chips - is it a real 'challenge'?

Zuzana Bínová^{a,*}, František Beneš^a, Jana Hajšlová^a

Department of Food Analysis and Nutrition, University of Chemistry and Technology, Technická 3, 160 00 Prague, Czech Republic

Capsaicinoids are biologically active secondary metabolites found in the placental tissue of chili peppers. These fruits are utilized globally in cuisine to impart a range of flavors and levels of “heat,” with the character and intensity varying across different varieties. The most common capsaicinoids include capsaicin, dihydrocapsaicin and nordihydrocapsaicin. The Scoville Heat Scale (SHU) serves as a conventional method to evaluate the pungency of chili-pepper-based products, spanning from 0 to 16,000,000 SHU for pure capsaicin (Angelov T., et al., 2023). The increasing interest in spicy foods among the youth has motivated producers to incorporate highly pungent chili powders into their products. An example of this trend is the “Hot Chips Challenge”, which led to severe health issues, including gastrointestinal pain and discomfort, after consuming just one chip. This incident led to alerts through the RASFF and the subsequent removal of this product from the EU market.

Employing a validated U-HPLC-HRMS/MS method, we analyzed the capsaicinoid profile in a selection of individual chips. The total capsaicinoid content was found to exceed 2000 mg/kg. In the absence of regulatory caps for capsaicinoid content in chili-based products, guidance documents from the BfR and EFSA are available for reference. Given the BfR's proposed NOAEL (No Observed Adverse Effect Level) for capsaicin at 0.083 mg/kg and LOAEL (Lowest Observed Adverse Effect Level) at 0.25 mg/kg of body weight, our findings indicate that the NOEL was surpassed by 20% for adults and 100% for young people.

Keywords: chilli peppers, capsaicinoids, health risk

References

- Angelov T, Gavrilova A, Panayotov N, Dyakova G, Pashev A, Gavrilov G, Yotova M (2023) Capsaicinoids content in some Bulgarian varieties of *Capsicum annuum* L. obtained by RP-HPLC. *Pharmacia* 70(3): 771-777. <https://doi.org/10.3897/pharmacia.70.e109767>.
- Bundesinstitut für Risikobewertung (BfR). (2011). Zu scharf ist nicht gesund - Lebensmittel mit sehr hohen Capsaicingehalten können der Gesundheit schaden [Too spicy is not healthy - Foods with very high levels of capsaicin can be harmful to health]. Stellungnahme Nr. 053/2011. Retrieved from <https://www.bfr.bund.de/cm/343/zu-scharf-ist-nicht-gesund-lebensmittel-mit-sehr-hohen-capsaicingehalten-koennen-der-gesundheit-schaden.pdf>.
- Hensel, A. (2023). Hot chilli peppers: Health risks of capsaicin(oids). Presentation at the 90th Advisory Forum Meeting. German Federal Institute for Risk Assessment (BfR). URL <https://www.efsa.europa.eu/en/events/90th-advisory-forum-meeting>.
- European Commission, Health & Consumer Protection Directorate-General. (2002). Opinion of the Scientific Committee on Food on Capsaicin (SCF/CS/FLAV/FLAVOUR/8 ADD1 Final). Brussels, Belgium. Retrieved from https://ec.europa.eu/food/fs/sc/scf/out120_en.pdf.

* Presenting author. E-mail address: bobkovaz@vscht.cz (Zuzana Bínová)

S2.P233 Botanical adventure in pharmacy education: implementing an escape game for quality control training

Ledoux Allison*, Bonnet Olivier, Leroy Charline, Frédérick Michel, Hamann Carla, Francotte Pierre

University of Liège Faculty of Medicine, CIRM, Building B36 Quartier Hôpital 4000, Liège, Belgium

Pharmacy education continuously seeks inventive methods to engage and educate students in response to the dynamic pharmaceutical landscape. Active and experiential learning approaches have emerged as effective strategies in this pursuit. This study introduces a novel application of escape game principles in pharmacy education, tailored specifically for third-year bachelor's students.

Our escape game intricately simulates quality control procedures in phytomedicines, with a focus on analyzing mint leaves and essential oils in accordance with European Pharmacopoeia guidelines. Quality control plays a pivotal role in pharmaceuticals, particularly those derived from botanical sources, necessitating a thorough understanding from students. Traditional teaching methods often lack immersive experiences in this specialized field.

Comparative analysis utilizing Gaussian curves demonstrates discernible differences in student performance between the years 2022 and 2023. The introduction of the escape game resulted in improved grades and reduced variability, indicating its positive impact on student learning outcomes. Importantly, consistent outcomes for a standardized task across both years, including unchanged practical assignments, underscore the game's influence on grades, rather than solely students' inherent skill enhancement. Our study underscores the potential of innovative pedagogical approaches in pharmacy education, suggesting further investigation into their long-term efficacy across diverse educational settings.

Keywords: escape game, pharmacognosy education, *Mentha piperita*

* Presenting author. E-mail address: allison.ledoux@uliege.be (Allison Ledoux)

S2.P234 Call for a contextual reform of the regulations on traditional herbal medicine in Burkina Faso

Kampadilemba Ouoba^{a,b,c}, Jean-Yves Pabst^c, Rasmané Semdé^a

^a Laboratory for Drug Development (LADME), Centre for Training, Research and Expertise in Drug Sciences (CEA-CFOREM), Joseph Ki-Zerbo University, 03 BP 7021 Ouagadougou, Burkina Faso ^bNational Agency for Pharmaceutical Regulation (NRA), Ministry of Health, 03 BP 7009 Ouagadougou, Burkina Faso

^c EA 7307, Centre for International and European Studies (CEIE), Faculty of Pharmacy, University of Strasbourg, 67400 Illkirch, France

Medicinal plants and traditional phytotherapy were legally recognised in Burkina Faso in 1994, under the Public Health Code, as being able to contribute to access to healthcare. Subsequently, a regulatory framework on the conditions for practising traditional medicine, traditional plant-based medicines and traditional health establishments was adopted with a view to promoting their integration into the health system. However, adherence to these regulations by traditional medicine practitioners has not been documented. The aim of this study was to assess the regulatory compliance of the practices of these traditional practitioners in relation to traditional herbal medicine and to formulate prospects for improvement. A mixed cross-sectional study using individual semi-directive interviews was conducted with traditional practitioners in four health regions of the country.

Sixty-seven (67) traditional healers, with an average age of 56 years, the majority of whom were men (71.6%), took part in the study. Herbalists (43.3%) and naturopaths (37.3%) were the main professional categories. The majority (82%) did not have legal authorisation to practise traditional medicine, and most herbalist establishments (92%) did not have a licence to open. The main reason for these regulatory non-compliances was that the regulation of traditional herbal medicine is very bureaucratic and orthodox, and is not adequate to the socio-cultural realities of traditional health practitioners. The results of this study therefore call for national reform of the regulations on traditional herbal medicine in Burkina Faso, with a view to facilitating its integration into the conventional health system.

Keywords: Traditional herbal medicine, Access to healthcare, Regulation, Burkina Faso

* Presenting author. E-mail address: ouobakampa@gmail.com (Kampadilemba Ouoba)

S2.P235 The in vitro modulatory effects of *Artemisia sieberi* on phase-I cytochrome P-450 isoenzyme gene expression using human hepatoma cells (HepG2)

Bader Hazazi^a, Amani Alsharidah^b, Aljowhara Almogren^a, Marwah Alruzaihi^b, Abir Alamro^b, Zeyad Alehaideb^{a,c,d,*}, Mohamed Boudjelal^{a,c,d}

^a King Abdullah International Medical Research Centre, Riyadh, Saudi Arabia

^b King Saud University, Riyadh, Saudi Arabia

^c King Saud bin Abdulaziz for Health Sciences, Riyadh, Saudi Arabia

^d Ministry of National Guard Health Affairs, Riyadh, Saudi Arabia

The modulation of cytochrome P-450 (CYP) gene expression by natural products can lead to an unwanted clinical situation and drug adverse effect(s). We investigated if the consumption of *Artemisia sieberi* (*A. sieberi*) could modulate five CYP enzymes using Hep G2 cells. The plant was extracted twice as boiled water and water by sonication to mimic the traditional consumption methods. The endotoxin levels were determined in the aqueous extracts using the LAL assay. The modulation of CYP gene expression was determined using the RT q-PCR assay. The aqueous extracts were screened for Pregnane X Receptor (PXR) ligands using the competitive binding assay. The endotoxin levels in the plant extracts were determined to be below the maximum allowable endotoxin concentration and deemed suitable for Hep G2 cell culture. The IC₅₀ values for the boiled and sonicated extracts of *A. sieberi* extracts were 110.6 and 191.9 µg/mL, respectively. Thus, an extract concentration of 16.5 µg/mL, and lower, were chosen for the subsequent gene expression experiments. The aqueous extracts were found to modulate the gene expression of two CYP isoenzymes significantly. The aqueous extract of boiled *A. sieberi* were able to induce the gene expression of CYP 1A2 and 3A4 for more than two-fold in comparison to untreated control. The receptor binding assay show that constituent(s) of *A. sieberi* boiled extract bind strongly to PXR. The study warns consumers from the concomitant consumption of boiled *A. sieberi* aqueous extract and narrow-therapeutic indexed pharmaceutical drugs that are mainly metabolized by CYP 1A2 and 3A4 enzymes.

Keywords: *Artemisia sieberi*, cytochrome P-450 enzyme induction, PXR, herb-drug Interaction

* Presenting author. E-mail address: alehaidebz@kaimrc.edu.sa (Zeyad Alehaideb)

S2.P236 Comparison of qNMR and HPLC-UV techniques for quantitation of secoiridoid phenolics in olive oil

Sofia Lympieropoulou, Panagiotis Diamantakos, Nikoleta Anna Madelou*, Eleni Melliou, Prokopios Magiatis

Laboratory of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimioupolis 15771, Athens, Greece

Extra virgin olive oil (EVOO), a principal component of the Mediterranean diet, is distinguished by its rich concentration in secoiridoid phenolic compounds with antioxidant properties. Despite its recognized association with cardiovascular disease prevention (Health claim EU 432/2012), the lack of a standardized analytical method in EU legislation for the measurement of the secoiridoid phenolics, creates a notable gap. The International Olive Council requires the application of an HPLC-UV method for phenolic compounds quantitation in *Olea europaea* L.

The aim of this study was the comparison of HPLC-UV (1) and qNMR (2) techniques for the quantitation of secoiridoid phenolics in EVOO. qNMR directly measures signals proportional to the number of atoms and molecules, while HPLC-UV indirectly correlates signals with the number of molecules using tyrosol equivalents or other response factors (1). Notably, discrepancies in quantitation results between the two techniques were observed. For instance, equimolar quantities of certain phenolic compounds, verified by NMR, yielded significantly different peak areas when measured by HPLC-UV due to different light absorption capacities for each compound (Fig. 1). For the first time we have clarified the quantitative correlation between NMR and HPLC signals for the EVOO phenolics related with the EU health claim regulation. Additionally, some EVOO secoiridoid phenolic compounds may undergo transformation or isomerization reactions during chromatography which may result in inaccurate assessment of their true content in olive oil when measured by HPLC.




Fig. 1. Comparison of NMR and HPLC-UV techniques for quantifying secoiridoids in equimolar quantities in olive oil.

Keywords: secoiridoids, olive oil, liquid chromatography, nuclear magnetic resonance, UV-Vis spectrometry

References

1. https://www.internationaloliveoil.org/wp-content/uploads/2022/06/Doc.-No-29-REV-2_ENK.pdf
2. Karkoula E., Skantzari A., Melliou E., & Magiatis P., 2012. Direct Measurement of Oleocanthal and Oleacein Levels in Olive Oil by Quantitative ^1H NMR. Establishment of a New Index for the Characterization of Extra Virgin Olive Oils. *Journal of Agricultural and Food Chemistry*, 60(47), 11696–11703.

* Presenting author. E-mail address: nikoletamadelou@gmail.com (Nikoleta Anna Madelou)

S2.P237 Anticancer efficacy of Korean medicine and sorafenib (concomitant drug) for hepatic tumors

Mi Rae Shin, Jin A. Lee, Min Ju Kim, Min Hyuk Oh, Seong Soo Roh*

College of Korean Medicine, Daegu Haany University, Daegu, South Korea

This study evaluated the therapeutic effects of a *Dioscoreae rhizome* and *Polygalae radix* mixture in combination with sorafenib (MIX) on a transplantation tumor model induced using hepatocellular carcinoma HepG2 cells.

The HepG2 cells transplantation tumor model has been established through HepG2 subcutaneously injection in Balb/c nude mice and the effect of MIX on the growth of the transplantation tumor was observed. Natural killer (NK) cell activity was measured using Granzyme B. Telomerase reverse transcriptase (TERT) and alpha-fetoprotein (AFP), known as prognostic markers of hepatocellular carcinoma, were analyzed by real-time PCR. Also, apoptosis and inflammation-related protein expressions were assayed by Western blotting.

Among the MIX groups, MIX25 (sorafenib 10 mg/kg + *Dioscoreae rhizome* and *Polygalae radix* mixture 25 mg/kg) showed surprisingly the best effect. MIX25 inhibited the tumor volume and weight. MIX25 enhanced apoptosis-promoting proteins such as Bax, cytochrome c, and caspase-3 and decreased apoptosis-inhibiting proteins such as survivin. In addition, MIX25 inhibited MAPK inflammatory response and inflammatory markers including TNF- α and COX-2. NK cell activity in the MIX25 group showed a significant difference compared with other drug-treated groups. Moreover, cancer marker genes including AFP and TERT in the MIX25 groups showed a better suppressive effect compared with MIX-treated groups.

MIX25 inhibited HepG2 cell transplantation tumor proliferation in Balb/c nude mice and could be suggested as a potential candidate for the treatment of patients with hepatocellular carcinoma.

Keywords: *Dioscoreae rhizome*, *Polygalae radix*, sorafenib, HepG2 cell, anticancer

References

Davis, J., Caldwell, S., 2020. Healing gone wrong: convergence of hemostatic pathways and liver fibrosis? Clin. Sci. (Lond) 34, 2189-2201.

Acknowledgements

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No.2018R1A5A2025272).

* Presenting author. E-mail address: ddede@dhu.ac.kr (Seong Soo Roh)

S2.P238 *Erythrina* spp. as natural anti-inflammatory ingredients for cosmeceuticals: An *in-silico* study

Natalia Blicharska^{a,*}, Dino Rotondo^a, Simon Jackson^b, Paolo Goi^b, Veronique Seidel^a

^a Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, 161 Cathedral Street, Glasgow UK

^b Davines Group - Botanical Research Department, Via Don Angelo Calzolari 55/A, 43126 Parma, Italy

Many inflammatory skin conditions depend upon activation of the nuclear factor NF- κ B pathway which triggers the transcription of genes coding for pro-inflammatory mediators (Bell *et al.*, 2003). Considering the rising demand for natural cosmeceuticals, developing products containing anti-inflammatory botanical ingredients capable of modulating the NF- κ B pathway is worth investigating. *Erythrina* spp. are used traditionally to treat skin ailments and alleviate inflammation. Their extracts and phytochemicals have demonstrated anti-inflammatory activity (Jiménez-Cabrera *et al.*, 2020; Khumalo *et al.*, 2021; Susilawati *et al.*, 2023). These anti-inflammatory phytochemicals include flavonoids, a class of compounds already known to target the NF- κ B pathway (Kadioglu *et al.*, 2015). In this study, an *in-silico* molecular docking approach using AutoDock Vina was employed to predict the binding affinities and ligand efficiencies of 19 flavonoid derivatives, from *Erythrina* spp. with anti-inflammatory activity, against three key proteins of the NF- κ B pathway. These included the NF- κ B p50/p65 heterodimer, the I κ B kinase subunit b (IKKb) and the NEMO/IKK association domain protein.

The isoflavonoids alpinumisoflavone (1), alpumisoflavone (2) and wighteone (3) showed the best ligand efficiencies towards the NF- κ B p50/p65 complex, while the pterocarpan erybraedin A (4) and phaseollidin (5) showed the strongest ligand efficiencies towards IKK β and NEMO/IKK, respectively. To the best of our knowledge, this is the first report on the interactions of *Erythrina* flavonoids with proteins of the NF- κ B pathway.

Further *in-vitro* studies are warranted to validate the effects of these compounds on the NF- κ B pathway and confirm the potential of *Erythrina* spp. as natural anti-inflammatory ingredients for cosmeceuticals.

Keywords: *Erythrina*, anti-inflammatory, NF- κ B, *in-silico*, molecular docking

References

- Bell, S., Degitz, K., Quirling, M., Jilg, N., Page, S., Brand, K., 2003. Involvement of NF-kappaB signalling in skin physiology and disease. *Cell Signal.* 15, 1-7
- Kadioglu, O., Nass, J., Saeed, M.E., Schuler, B., Efferth, T., 2015. Kaempferol is an anti-inflammatory compound with activity towards NF- κ B pathway proteins. *Anticancer Res.* 35, 2645-2650
- Jiménez-Cabrera, T., Bautista, M., Velázquez-González, C., *et al.*, 2020. Promising antioxidant activity of *Erythrina* genus: An alternative treatment for inflammatory pain? *Int. J. Mol Sci.* 22, 1-19
- Khumalo, G.P., Sadgrove, N.J., Van Vuuren, S.F., Van Wyk, B.E., 2021. South Africa's best bark medicines prescribed at the Johannesburg Muthi markets for skin, gut and lung infections: MIC's and brine shrimp lethality. *Antibiotics.* 10, 1-27
- Susilawati, E., Levita, J., Susilawati, Y., Sumiwi, S.A., 2023. Pharmacology activity, toxicity and clinical trials of *Erythrina* genus plants (Fabaceae): an evidence-based review. *Front. Pharmacol.* 14, 1-9

* Presenting author. E-mail address: natalia.blicharska@strath.ac.uk (Natalia Blicharska)

Medicinal Plants and Natural Products in Animal Healthcare and Veterinary Medicine

S2.P239 Chemical composition of resin extract from *Corymbia* spp. an indigenous Australian native plant

A. F. Ruiz^{a,*}, Matthew Perry^a, Gerry Turpin^c, Kate Miller^b, Phurpa Wangchuk^a

^a Centre for Molecular Therapeutics, Australian Institute of Tropical Health and Medicine, James Cook University, Building E4, McGregor Rd, Smithfield, Cairns, QLD 4878, Australia

^b College of Medicine and Dentistry James Cook University Smithfield, Queensland 4878, Australia

^c Tropical Herbarium of Australia, James Cook University, Building E1, McGregor Rd, Smithfield, Cairns, QLD 4878, Australia

Corymbia spp. an indigenous Australian plant found across regions from northern Queensland to New South Wales, has attracted considerable attention due to its medicinal properties, particularly its essential oils (Schuster et al., 2018). However, there remains a dearth of research concerning the compounds derived from its resin. To bridge this knowledge gap, this study employed various techniques such as column chromatography, size exclusion, and multiple rounds of preparative high-performance liquid chromatography (HPLC) to isolate compounds from the resin. The isolated compounds were determined using mass spectroscopy and Nuclear Magnetic Resonance (Bruker 600 MHz AVANCE III NMR spectrometer). 1D NMR (¹H and ¹³C NMR) and 2D NMR (COSY, HMBC, HSQC, NOESY) experiments were used in this identification (Ritmejeriyte et al., 2022). The analysis revealed the presence of (E)-3-(4-hydroxyphenyl) prop-2-enoic acid (**C1**), 3,5-dihydroxy-2-(4-hydroxyphenyl)-7-methoxy-chroman-4-one (**C2**), 5-hydroxy-2-(4-hydroxyphenyl)-7-methoxy-chroman-4-one (**C3**), 3,5-dihydroxy-2-(4-hydroxyphenyl)-6-[1-(4-hydroxyphenyl)ethyl]-7-methoxy-chroman-4-one (**C4**). The physicochemical properties of these compounds agreed with Lipinski rule of 5 for small molecule therapeutics, making them promising candidates for the development of new drugs (Daina et al., 2017).

Keywords: *Corymbia* spp., HPLC, NMR, physicochemical properties

References

- Daina, A., Michielin, O., & Zoete, V. (2017). SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* 7(1), 42717.
- Ritmejeriyte, E., Ryan, R. Y. M., Byatt, B. J., Peck, Y., Yeshi, K., Daly, N. L., Zhao, G., Crayn, D., Loukas, A., Pyne, S. G., Ruscher, R., & Wangchuk, P. (2022). Anti-inflammatory properties of novel galloyl glucosides isolated from the Australian tropical plant *Uromyrtus metrosideros*. *Chem. Biol. Interact.*, 368, 110124-110124.
- Schuster, T. M., Setaro, S. D., Tibbits, J. F. G., Batty, E. L., Fowler, R. M., McLay, T. G. B., Wilcox, S., Ades, P. K., & Bayly, M. J. (2018). Chloroplast variation is incongruent with classification of the Australian bloodwood eucalypts (genus *Corymbia*, family Myrtaceae). *PloS One*, 13(4), e0195034-e0195034.

* Presenting author. E-mail address: andres.ruizalvarez@my.jcu.edu.au (Andres Ruiz)

S2.P240 *Melissa officinalis* (Lemon balm) extract to promote calming effect before stressful events

Anaïs Rongeon^a, Sekhou Cisse^{b,c}, Fatima Aberkane^{b,c,*}, Nathalie Gauthier^d, Mohammed El Amine Benarbia^{b,c}, Yassine Mallem^a

^a ONIRIS, 101, Route de Gachet, 44307 Nantes Cedex 3, France

^b Nor-Feed SAS, 3 rue Amédéo Avogadro, 49070 Beaucouze, France

^c Joint Lab ANR FeedInTech (FIT: SONAS/Nor-Feed), France

^d Clinique vétérinaire des écoles de Lanester, 6 Av. François Billoux, 56600 Lanester, France

Behavioral disorders, due to stress, are common during veterinary consultation for dogs. Even if plant-based solutions are commonly used to calm dogs, data supporting their efficacy are scarce. *Melissa officinalis* is not an exception. This study aims to highlight its calming effect on dogs.

Fifteen dogs with behavioral disorders linked to stress were divided into 3 groups:

- Control group (CTL) fed with standard diet and supplemented with placebo (dextrose, 5 mg/kg/day)
- Lemon Balm (LB) group fed with standard diet and supplemented with *Melissa officinalis* extract (5 mg/kg/day)
- Positive control group (PC) fed with standard diet and treated with α cazosepin (15 mg/kg/day).

In collaboration with a local veterinary clinic, 15 dogs were recruited and randomly allocated to the 3 treatment groups. The treatment duration was 3 days. Physiological parameters (heart rate, respiratory rate, and body temperature) and behavioral parameters (body signs and special behavior) were monitored before and after the treatment.

After 3 days of treatment, LB dogs significantly reduced the intensity of the expression of their behavioral signs of stress between the two consultations compared to CTL dogs. The observed effect was at least similar to PC dogs. However, there is no significant differences between the 3 groups in terms of physiological parameters.

This study demonstrates the calming effect of *Melissa officinalis* before stressful events. However, the sample size was quite small. It may be worthwhile to extend this study by replicating it on a larger number of dogs to confirm the initial results obtained.

Keywords: *Melissa officinalis*, stress, behavior, pet

* Presenting author. E-mail address: sekhoul.cisse@norfeed.net (Sekhou Cisse)

S2.P241 Equine Stromal Cells: effects of non-ribosomal peptides on proliferation and viability

Taylor Blair^a, Laurie Goodrich^b, Kalindi Morgan^{c,*}

^a University of Northern British Columbia, 3333 University Way, Prince George, BC, Canada

^b Colorado State University, College of Veterinary Medicine, Fort Collins, Colorado, USA

^c University of Northern British Columbia, 3333 University Way, Prince George, BC, Canada

Equine stem cells (ESC) are used extensively in treatment of osteoarthritis and tendon and ligament injuries in horses. Two of the most common cell types used in treatments are bone marrow derived mesenchymal cells (BMSC) and articular chondroprogenitor cells (ACP) (McCarthy et al., 2012). Non-ribosomal peptides (NRP) have also been used throughout the medical field in drug development and antibiotic use (Felnagle et al., 2008). Virginiamycin is a pure NRP that was specifically fed to cattle to increase growth and milk production (McEwan and Fedorka-Cray 2002). The objective for this project is to compare doses of both non-pure and pure NRP's on the proliferation and viability of ESC's when growing in culture. The hypothesis is that proliferation and viability of ESC's will be increased with the addition of NRP's.

A non-pure crude NRP and Virginiamycin were added to the ESC's at doses 5ug, 10ug, 20ug, and 40ug for either 3 or 6 days. ESC's were then harvested and counted using a hemocytometer. A live/dead stain measured by flow cytometry was performed for viability. The flow cytometry revealed all cells maintained good viability. Proliferation results showed significant increases for doses of 10ug and 20ug Virginiamycin on day 3 ACP's. In the BMSC's, all dose amounts of Virginiamycin increased proliferation at day 3 but only the 5ug dose continued to increase proliferation by day 6. Based on these results, Virginiamycin can be added to culture to increase proliferation of ACP's and BMSC's. This can be beneficial when high cell counts are needed.

Keywords: Non-ribosomal peptides, stem cells, equine, *Virginiamycin*

References

- Felnagle EA, Jackson EE, Chan YA, Podevels AM, Berti AD, McMahon MD, Thomas MG. 2008. Nonribosomal Peptide Synthetases Involved in the Production of Medically Relevant Natural Products. *Mol Pharm.* 5(2):191–211.
- McCarthy, H.E., Bara, J.J., Brakspear, K., Singhrao, S.K., Archer, C.W. 2012. The Comparison of Equine Articular Cartilage Progenitor Cells and Bone Marrow-Derived Stromal Cells as Potential Cell Sources for Cartilage Repair in the Horse. *Veterinary Journal*, 192 (3), 345–351.
- McEwen, S.A., Fedorka-Cray, P.J. 2002. Antimicrobial Use and Resistance in Animals. *Clinical Infectious Diseases*, 34 (Supplement_3), S93–S106.

*Presenting author. E-mail address: kalindi.morgan@unbc.ca (Kalindi Morgan)

S2.P242 Extracts from the roots of *Scutellaria baicalensis* reduce honeybee death caused by nosemosis

Jakub Gębalski^{a,*}, Aneta A. Ptaszyńska^b, Daniel Załuski^{a,*}

^a Department of Pharmaceutical Botany and Pharmacognosy, Nicolaus Copernicus University, Marie Skłodowska Curie 9, 85-094 Bydgoszcz, Poland

^b Department of Immunobiology, Institute of Biological Sciences, Faculty of Biology and Biotechnology, Maria Curie-Skłodowska University, Akademicka 19 Str., 20-033 Lublin, Poland

Honeybees are very important pollinators and their role in the ecosystems can not be overestimated. *Apis mellifera* L. workers pollinate approximately 80% of flowering plants and only in the United States their effort is estimated up to \$15 billion worth of crops annually. Moreover, they produce honey, wax, pollen, royal jelly, and propolis, products eagerly used by many people. Unfortunately, honeybees suffer from pesticides, immunodeficiencies, beekeeping practice in which antibiotics are used, malnutrition, and disease. These factors can drastically reduce honeybee lifespan (Ptaszyńska and Załuski, 2020).

The aim of the study was to check if *Scutellariae baicalensis radix* extract can prolong honeybee lifespan especially after infected with pathogenic fungi (*Nosema* spp.). The 75% ethanolic extracts were prepared from the freshly collected roots after their lyophilization using classical maceration (CM) and ultrasound-assisted extraction (UAE).

After 20 days of administration of extracts (0,01; 0,1; 1,0; 10%) to non-infected honeybees, no toxic changes were observed, it did not shorten the honeybee lifespan.

After treatment of *Nosema*-infected honeybees with the extracts, the number of *Nosema* spores decreased to $2,05 \times 10^6$ and $4,20 \times 10^6$ for CM and UAE comparing to a control group $15,20 \times 10^6$. The 10% extracts appeared to reduce spores the strongest.

To sum up, the extracts decreased the level of honeybee nosemosis in infected insects, as well as when used as prophylaxis. All these state-of-the-art research findings, from *in vitro* to nature, should be taken into account when drafting protocols for prospective honeybee medicament tests.

Keywords: *Scutellaria baicalensis*, honeybees, *Nosema*

References

Ptaszyńska, A.A., Załuski D., 2020. Extracts from *Eleutherococcus senticosus* (Rupr. et Maxim.) Maxim. roots: a new hope against honeybee death caused by nosemosis. *Molecules*, 25, 1-25.

* Presenting author. E-mail address: daniel.zaluski@cm.umk.pl (Daniel Załuski)

S2.P243 Larch (*Larix decidua* L.) fibre waste: characterization of a functional feed ingredient

Doriana Eurosia Angela Tedesco^{a,*}, Alessandro Guerrini^a, Alessio Massironi^a, Stefania Marzorati^a

Department of Environmental Science and Policy, Via Celoria 2, 20133 Milan, University of Milan, Italy

In the larch woodworking industry, wood shavings or sawdust is a fibre waste product. In previous studies, larch fibre waste (*Larix decidua* L.) was evaluated for health-beneficial compounds, flavonols (taxifolin, TXF; dihydrokaempferol, DHK), and terpenoides (larixol, LX; larixyl acetate, LXA) and the antioxidant and anti-inflammatory potential effects (Pferschy-Wenzig et al., 2008; Stockhammer et al., 2009) also enhancing its use in animal nutrition (Tedesco et al., 2015; Tzika et al., 2017).

This study aimed to evaluate and compare the bioactive compounds and antioxidant potential of larch fiber waste (native to the mountains of central Europe) obtained from five woodworking companies, to verify possible differences in their composition. The hexane, methanol and water extracts were fractionated by UPLC-PDA instrument, to quantify the presence of TXF, DHK, LX, and LXA. ABTS assay was used to determine the antioxidant activity (EC_{50} mg/mL).

In the hexane extract, LX and LXA were quantified in the range 0.80-15.7 % w/w, and 8.70-14.50 % w/w respectively, while in the methanol extract, TXF and DHK were present in the range 16.20-23.20 % w/w and 6.40-24.00 % w/w respectively. The mean antioxidant activity of the methanol extracts was $EC_{50} = 0.002$ mg/mL, and $EC_{50} = 0.012$ mg/mL of the water extract.

Although there were differences in bioactive compound content, it can be concluded that larch fibre waste from woodworking companies requires further attention as a functional feed ingredient in supporting metabolism and animal health.

Keywords: wood waste, biomass valorization, sustainable feed

References

- Pferschy-Wenzig, E.M., Kunert, O., Presser, A., Bauer, R., 2008. In vitro anti-inflammatory activity of larch (*Larix decidua* L.) sawdust. J Agric Food Chem 24;56(24):11688-93.
- Stockhammer, S., Stolze, K., Rohr-Udilova, N., Chizzola, R., Zitterl-Eglseer, K., Franz, C., 2009. Antioxidant activity of phytochemicals from industrial waste and derived extracts for the production of feed and food additive. Int. J. Food Sci. Technol. 44, 702-710.
- Tedesco, D., Garavaglia, L., Spagnuolo, M.S., Pferschy-Wenzig, E.M., Bauer, R., Franz, C., 2015. In vivo assessment of an industrial waste product as a feed additive in dairy cows: effects of larch (*Larix decidua* L.) sawdust on blood parameters and milk composition. Vet J. 206, 322-326.
- Tzika, E.D., Tassis, P.D., Papatsiros, V.G., Pferschy-Wenzig, E.M., Siochu, A., Bauer, R., Alexopoulos, C., Kyriakis, S.C., Franz, C., 2017. Evaluation of in-feed larch sawdust anti-inflammatory effect in sows. Pol J Vet Sci. 1;20(2):321-327.

Acknowledgements

This study was carried out within the Agritech National Research Center and received funding from the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)–MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4–D.D. 1032 17/06/2022, CN00000022). This manuscript reflects only the authors' views and opinions, neither the European Union nor the European Commission can be considered responsible for them.

* Presenting author. E-mail address: doriana.tedesco@unimi.it (Doriana Eurosia Angela Tedesco)

S2.P244 Identification of novel caffeoyl- and feruloyl-derivatives in *Solanum glaucophyllum*

Sabrina Autzen^{a,*}, Thomas Heymann^b, Marcus A. Glomb^b

^a Herbonis Animal Health GmbH, Rheinstraße 30, 4302 Augst BL, Switzerland

^b Institute of Chemistry, Department of Food Chemistry, Martin-Luther-University Halle-Wittenberg, Kurt-Mothes-Str. 2, 06120 Halle (Saale), Germany

Solanum glaucophyllum Desf. (SG) is one of the few plants that produce the source of the bioactive form of vitamin D as 1,25(OH)₂D₃. However, plants are always a multi-component system with synergistic effects, and it is difficult to correlate efficacy data with a unique component. The aim of this study was to expand the knowledge of phenolic compounds in SG, which have mainly been reported with a focus on quercetin derivatives (Rappaportt et al., 1977). For the first time, several chlorogenic acids were detected in leaf extracts of SG. Specifically, 3-O-caffeoylquinic acid (chlorogenic acid) and 5-O-caffeoylquinic acid (neochlorogenic acid) were unequivocally verified via reference material, collision-induced dissociation (CID) experiments and high-resolution mass spectrometry (HR-MS).

The workup included extraction with acetone-water, size exclusion and reversed phased chromatography leading to further caffeoyl- and feruloyl-derivatives. LC-MS/MS experiments indicated conjugations to glucose and glucaric acid. Interestingly, fragmentation of the molecular ions led to the very same patterns at different retention times suggesting several isomers, e.g. those known for tomatoes (Larbat et al., 2014). Independent organic syntheses performed for various isomers of caffeoyl glucoside, feruloyl glucoside, caffeoyl glucaric acid and feruloyl glucaric acid esters allowed verification of these substances in SG for the first time. All substances were fully verified by NMR, CID and HR-MS experiments. Compared to current literature the phenolic spectrum of SG was significantly increased by the work reported here.

Keywords: *Solanum glaucophyllum*, polyphenols, caffeoyl derivatives, feruloyl derivatives

References

- Rappaportt, I., Giacomello, G., Seldes, A.M., Blanco, M.C., Deulofeu, V., 1977. Phenolic glycosides from *Solanum glaucophyllum*: a new quercetin triglycoside containing D-Apiose. *Phytochem.* 16, 1115–1116.
- Larbat, R., Paris, C., Le Bot, J., Adamowicz, S., 2014. Phenolic characterization and variability in leaves, stems and roots of Micro-Tom and patio tomatoes, in response to nitrogen limitation. *Plant Sci.* 224, 62–73.

* Presenting author. E-mail address: s.autzen@herbonis.com (Sabrina Autzen)

S2.P245 Phytochemical screening and study of the hypotensive effect of the aqueous extract of *Rubus ulmifolius* Schott leaves

Afaf Mehio^{1,2*}, Chaimae Alla^{1,2}, Zachee L. E. Akissi², Anca Lucau-Danila², Abdelkhaleq Legssyer¹, Jean-Louis Hilbert², Abderrahim Ziyat¹, Sevser Sahpaz²

¹ Laboratory of Bioresources, Biotechnologies, Ethnopharmacology and Health, Department of Biology, Faculty of Sciences, University Mohammed First, Oujda-Morocco

² BioEcoAgro Joint Cross-Border Research Unit 1158, Univ. Lille, INRAE, Univ. Liège, UPJV, JUNIA, Univ. 11 Artois, Univ. Littoral Côte d'Opale, 59650 Villeneuve d'Ascq, France;

Hypertension (hypertension) is a cardiovascular pathology characterized by systolic blood pressure (SBP) ≥ 140 mmHg or diastolic blood pressure (DBP) ≥ 90 mmHg (Yousefi et al., 2021). Several antihypertensive agents have many side effects, hence herbal medicine as an alternative (Landazuri et al., 2017). *Rubus ulmifolius* Schott (*Rosaceae*) is a thorny shrub, used in traditional medicine for the treatment of several pathologies such as hypertension (Ali et al., 2017; Fakchich and Elachouri, 2021).

The objective of this study is to investigate its hypotensive effect and analyze its chemical composition. The study of the hypotensive effect of the aqueous extract of *R. ulmifolius* leaves (AERu), administered intravenously, was conducted on anesthetized Wistar rats maintained at a temperature of 37°C. Compound separation from AERu was conducted using ultra-high performance liquid chromatography coupled with mass spectrometry (UHPLC-MS). The results showed that the intravenous administration of the aqueous extract in anesthetized normotensive rats induced a dose-dependent hypotensive effect, with a reduction in systolic blood pressure (SBP), diastolic blood pressure (DBP), and mean blood pressure (MBP) by 47.77%, 51.98%, and 48.73%, respectively, at a dose of 50 mg/kg of body weight. Furthermore, the extract showed no effect on heart rate. Phytochemical analysis revealed the presence of chlorogenic acid, quercetin-3-O- β -D-glucuronoside, and kaempferol-3-O- β -D-glucuronoside.

In conclusion, our results demonstrated that the aqueous extract of *R. ulmifolius* leaves has a significant hypotensive effect. The richness of this plant in bioactive phytochemical compounds may be responsible for this effect.

Keywords: *Rubus ulmifolius*, hypertension, hypotensive, polyphenols, tannins

References

- Ali, N., Shaoib, M., Shah, S. W. A., Shah, I., Shuaib, M. 2017. Pharmacological profile of the aerial parts of *Rubus ulmifolius* Schott. BMC complementary and alternative medicine, 17, 1-7. DOI 10.1186/s12906-017-1564-z.
- Fakchich, J., ElachourI, M. 2021. An overview on ethnobotanico-pharmacological studies carried out in Morocco, from 1991 to 2015: Systematic review (part 1). J Ethnopharmacol, 267, 113200. doi.org/10.1016/j.jep.2020.113200.
- Landazuri, P., Chamorro, N., Cortes, B. 2017. Medicinal plants used in the management hypertension. J Anal Pharm Res, 5, 00134. DOI: 10.15406/japlr.2017.05.00134.
- Yousefi, M., Shadnoush, M., Khorshidian, N., Mortazavian, A. M. 2021. Insights to potential antihypertensive activity of berry fruits. Phytotherapy Research, 35, 846-863. doi.org/10.1002/ptr.6877

* Presenting author. E-mail address: afafmehiou@outlook.fr; afaf.mehiou.etu@univ-lille.fr (Afaf Mehio)

S2.P246 Evaluating the antibacterial properties of macroalgae against swine pathogens

Peter O'Hara*, James Blee, Thomas J. Smyth, Owen Kenny

Atlantic Technological University Sligo. Department of Health and Nutritional Science, Ireland

The EU swine industry is the second largest in the world, while also being the largest exporter of pork and pork products (€13.8bn in 2022). However, the recent banning of zinc oxide as an antibacterial agent in pig production may have a knock-on effect in the sustainment and future growth of this industry. Although the supplementation of animal feed with seaweed is common for increasing key performance indicators (i.e. Feed intake, weight gain and prebiotics), less is known about their efficacy as antibacterial agents. In this study, the antibacterial properties of extracts, derived from thirteen macroalgae species, were investigated against five priority swine pathogenic bacteria, using a broth microdilution assay. Of the 65 extracts tested, 38 were shown to have widespread activity at minimum inhibitory concentrations (MICs) ≤ 1 mg/ml. Overall, extracts from Rhodophyta and Phaeophyceae genera macroalgae demonstrated greater activity (1000 – 31.25 $\mu\text{g/mL}$) compared to the Chlorophyta genus macroalgae, respectively. In particular, non-polar extracts of *Alaria esculenta* (Phaeophyceae), *Asparagopsis armata* & *Asparagopsis taxiformis* (Rhodophyta) displayed the strongest antibacterial activity against the Gram-positive *S. aureus* (62.5 – 31.25 $\mu\text{g/mL}$). While the dichloromethane (DCM) extracts of *A. esculenta* & *A. taxiformis* displayed the strongest antibacterial activity against the Gram-negative *S. enterica* subsp & *E. coli* (31.25 $\mu\text{g/mL}$). The results of this study highlight the antibacterial potential of macroalgae, which may further support its future application as an antibacterial agent in supplemented animal feed.

Keywords: Macroalgae, swine, antibacterial, supplemented animal feed

* Presenting author. E-mail address. Peter.OHara@research.atu.ie (Peter O'Hara)

S2.P247 A survey-based study on knowledge, attitude, and perception of pet owners towards the use of plant extracts from bio-refinery in pet food formulations

Talal Hassan^{a,*}, Joana Nery^a, Mohammad Zaker Rehan^b, Muhammad Irfan Malik^a, Silvia Mioletti^a, Luigi Bertolotti^a, Alba Maria Ramos^c, Natascia Bruni^b, José M. Pinilla-Rosas^c, Maria Teresa Capucchio^a

^a Department of Veterinary Sciences, University of Turin, 10095 Grugliasco (Torino), Italy

^b Acelpharma, 10092 Beinasco (Torino), Italy

^c NATAC BIOTECH S.L., Alcorcón, Madrid (Spain)

Plant biomass extracts contain bioactive compounds that can be used in pet food, and therefore promote production cycle sustainability of biorefineries. Therefore, the use of plant extracts in pet food formulation can improve pet's gut-health while contributing to environmental sustainability (Buff et al., 2014). Understanding pet owners' thoughts on pet food quality is critical to knowing how much interest they demonstrate towards more sustainable solutions (Goswami et al., 2022). This study used a web based cross-sectional survey to evaluate Italian pet owners' knowledge, attitudes, and perceptions of innovative pet food containing plant based functional feed ingredients. Preliminary findings based on 169 responses revealed a predominantly female respondent base (63.3%), aged-51-64 years (38%), mainly from the northwest region of Italy (80%), and primarily employed (47%). Most respondents owned one pet (48.2%), predominantly dogs (50.6%), and gastrointestinal issues were the most reported (39.2%) health disorders. The majority sought advice from veterinarians (80.1%) and purchased pet food from specialized shops (65.1%). Notably, a significant proportion expressed unfamiliarity with natural extracts in pet food (61.4%) but showed interest in extracts, particularly from pomegranate (33.7%) and olive leaves (22.3%), to enhance gastrointestinal health (72.9%). A noteworthy percentage (42.2%) expressed willingness to pay a premium of 5-10% for pet food containing plant extracts.

This study highlights the growing interest among Italian pet owners in the addition of natural plant extracts to pet foods. Despite limited knowledge amongst consumers, there is a positive trend toward using plant extracts, particularly pomegranate and olive leaves, to improve gut-health.

Keywords: natural by-products, pet owners, questionnaire, animal-health

References

- Buff, P. R., Carter, R. A., Bauer, J. E., & Kersey, J. H. (2014). Natural pet food: A review of natural diets and their impact on canine and feline physiology. *J. Anim. Sci*, 92, 3781–3791.
- Goswami, L., Kayalvizhi, R., Dikshit, P. K., Sherpa, K. C., Roy, S., Kushwaha, A., Kim, B. S., Banerjee, R., Jacob, S., & Rajak, R. C. (2022). A critical review on prospects of bio-refinery products from second and third generation biomasses. *Chem. Eng. J.*, 448, 137677.

Funding

Funded by the European Union. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or CBE JU. Neither the European Union nor the CBE JU can be held responsible for them (Grant agreement ID: 101112434).

* Presenting author. E-mail address: talal.hassan@unito.it (Talal Hassan)

S2.P248 Anti-inflammatory effects of Cinnamon essential oil assessed on diverse cell models

D. Gardan-Salmon^{a,*}, M. Arturo-Schaan^a, A. Bruyère^b

^a Deltavit, CCPA Group, 35150 Janzé, France

^b Irset, Inserm UMR1085, Université Rennes, France

The One Health approach advocates cutting antibiotic use in livestock feed to boost human and animal health. Phytochemicals, with anti-inflammatory properties, offer valuable health benefits as natural alternatives. Cinnamaldehyde is a compound naturally found in Cinnamon (*Cinnamomum cassia*) with various therapeutic properties (Da Silveira et al., 2014). In the context of designing functional feed in antibiotic-free diets for animals, it is mandatory to evaluate its effects on livestock models, thereby reducing *in vivo* studies. This study aimed to assess the efficacy of a Cinnamon essential oil (Cinna; 5 to 100 µg/mL) in modulating inflammation *in vitro* across three cell models. Chicken macrophage-like HD11 and porcine enterocyte IPEC-J2 cells were used to study cytokine gene expression in livestock cell models. Mechanisms underlying such effects in enterocytes were further investigated with a human reference cell line (Caco-2), by cell imagery. Cinna alleviated inflammation by significantly reducing IL-1β and IL-8 gene expressions in HD11 cells (50 and 100 µg/mL, $p < 0.01$), and IL-6, IL-8, TNF-α, CCL20 gene expressions in IPEC-J2 cells (-2 to -30-fold depending on cytokines and doses; $p < 0.05$), compared to pro-inflammatory group (Fig. 1). In the Caco-2 cells, Cinna (50 and 100 µg/mL) alleviated inflammation by significantly reducing the % of NF-κB positive nuclei (-70%; $p < 0.01$). Using cellular models, these results demonstrate Cinna's anti-inflammatory effects by regulating pro-inflammatory cytokines in livestock models. This regulation involves the NF-κB pathway, as confirmed in Caco-2 cells. Cinna shows promise as a candidate for developing functional feed for livestock. The authors declare no conflicts of interest.




Fig. 1. Effect of Cinnamon essential oil on induced inflammation in 3 different cell models. HD-11 cells (A) or IPEC-J2 cells (B) were pre-treated with 5 to 100 µg/mL of Cinnamon essential oil (Cinna) for 2h (HD11) or 1h (IPEC-J2) and then co-incubated with LPS (10 ng/mL) for 2 or 1h respectively (n=3). IL-8 gene expression (A and B) was measured by real-time qPCR. Caco2 cells (C) were pre-treated with 5 to 100 µg/mL of Cinna for 1h and then co-incubated with IL-1β for 1h. Inflammation was assessed using fluorescent NF-κB immunocytochemical labeling (C).

Keywords: Cinnamon essential oil, HD11, IPEC-J2, cytokines, NF-κB

References

R. D. C. Da Silveira E Sá, L. N. Andrade, R. D. R. B. De Oliveira, and D. P. De Sousa, 2014. A review on anti-inflammatory activity of phenylpropanoids found in essential oils. *Molecules*. Vol. 19, no. 2, 1459–1480.

* Presenting author. E-mail address: dgardan-salmon@ccpa.com (Delphine Gardan-Salmon)

S2.P249 Induction of apoptosis in pancreatic cancer cells by *Albizia julibrissin* extract through inhibition of an oncogenic signaling pathway

Ji Su Han^{a,b,*}, Sang-Chul Han^a, Seon-A Yoon^c, Hyejin Hyeon^c, Young-Min Ham^c, Yong-Hwan Jung^c, Hye-Jin Boo^{a,b}

^a Department of Histology, College of Medicine, Jeju National University, Jeju, Republic of Korea

^b Jeju Research Center for Natural Medicine, Jeju National University, Jeju, Republic of Korea

^c Biodiversity Research Institute, Jeju Technopark, Jeju, Republic of Korea

Pancreatic cancer is one of the most fatal solid tumors due to poor prognosis. *Albizia julibrissin* exhibits anti-inflammatory and antioxidant against a variety of diseases. However, little is known about its anti-cancer effects in human pancreatic cancer cells. In the present study, we aimed to investigate the anti-cancer potential of *Albizia julibrissin* extract against human pancreatic cancer cells and define its molecular mechanism of action. Anti-cancer effects of *Albizia julibrissin* extract were determined by WST-1 assay, colony formation assay, western blot analysis, Annexin V/PI staining, and animal experiments using xenograft mouse model. *Albizia julibrissin* extract significantly suppressed the growth and colony formation of pancreatic cancer cells. *Albizia julibrissin* extract also induced apoptosis in pancreatic cancer cells by enhancing the expression levels of cleaved PARP and cleaved caspase-3 as well as inhibiting oncogenic signaling such as MAPK and PI3K/AKT pathway. In addition, *Albizia julibrissin* extract suppressed tumor growth and induced apoptosis in xenograft mouse model used human pancreatic cancer cells. Our findings provide crucial evidence that *Albizia julibrissin* extract may have potential as a therapeutic drug for pancreatic cancer.

Keywords: natural product, *Albizia julibrissin*, npoptosis, pancreatic cancer

References

- Halbrook CJ, Lyssiotis CA, Magliano MP, Maitra A., 2023. Pancreatic cancer: Advances and challenges. *Cell*. 13;186(8):1729-1754.
- Hu ZI, O'Reilly EM. 2024. Therapeutic developments in pancreatic cancer. *Nat Rev Gastroenterol. Hepatol*. 21(1):7-24.
- Naeem A, Hu P, Yang M, Zhang J, Liu Y, Zhu W, Zheng Q. 2022. Natural Products as Anticancer Agents: Current Status and Future Perspectives. *Molecules*. 30;27(23):8367.

Funding

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2022R1I1A1A01073350, RS-2023-00270936) and was supported by “Regional Innovation Strategy (RIS)” through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (MOE) (2023RIS-009).

* Presenting author. E-mail address: gkswlndjaak@jejunu.ac.kr (Ji Su Han)

S2.P250 Antibacterial, antibiofilm and quorum sensing inhibitory activities of fractions of the ethanol extract of *Searsia lancea* against pathogens causing bovine mastitis

Ayodele O. Akinboye^a, Fikile N. Makhubu^a, Ibukun M. Famuyide^a, Joanne Karzis^b, Inge-Marie Petzer^b, Lyndy J. McGaw^{a,*}

^a Phytomedicine Programme, Department of Paraclinical Sciences, Faculty of Veterinary Science, Private Bag X04, Onderstepoort 0110, South Africa

^b Department of Production Animal Studies, Faculty of Veterinary Science, University of Pretoria, Pretoria, South Africa

Bovine mastitis is a significant problem in the dairy industry exacerbated by antibiotic-resistant pathogens able to form protective biofilms via quorum sensing. This study aimed to determine antibacterial, antibiofilm and anti-quorum sensing activities, and *in vitro* safety, of fractions of the ethanol extract of *Searsia lancea* against isolates of major multidrug resistant bacteria of mastitis origin. Antibacterial, antibiofilm and anti-quorum sensing activities of the fractions were determined against pathogenic bacteria isolated from clinical cases of mastitis. Cytotoxicity was determined against bovine dermis cells. A Waters UPLC-QTOF-MS (XEVO-G2) instrument was used to tentatively identify compounds in the most active fractions.

DCM and ethyl acetate fractions had MIC values of 0.001 and 0.01 mg/mL respectively, better than that of the antibiotic ciprofloxacin. All fractions were relatively non-cytotoxic. The DCM fraction had excellent selectivity index values up to 90, but the ethyl acetate fraction was least toxic. The ethyl acetate fraction had the best antibiofilm and preformed biofilm disrupting activities. Notably, the DCM and water fractions demonstrated the lowest minimum quorum sensing inhibitory concentration (MQSIC), with DCM MQSIC₅₀ = 0.01 mg/mL, suggesting that fractions have potential to modulate virulence factors. Compounds including myricetin 3-O-glucoside were identified in active fractions.

These findings underscore the potential of fractions of *S. lancea* extract as valuable resources in combating antimicrobial resistant mastitis pathogens. Further investigation of their mechanisms, identification of active compounds, and thorough *in vivo* assessments are essential steps toward their development as effective preventative and treatment options for microbial infections such as bovine mastitis.

Keywords: bovine mastitis, antibiotic resistance, quorum sensing, antibiofilm, *Searsia lancea*

* Presenting author. E-mail address: lyndy.mcgaw@up.ac.za (Lyndy McGaw)

S2.P251 Potential use of *Paulownia* leaves as feed additive to enhance animal health condition

Marta Mendel^{a,*}, Wiesław Ptach^b, Joanna Polak^a, Urszula Latek^a

^a Department of Preclinical Sciences Institute of Veterinary Medicine, Warsaw University of Life Sciences, 8, Ciszewskiego Str., 02-876 Warsaw, Poland

^b Department of Remote Sensing and Environmental Assessment. Institute of Environmental Engineering. Warsaw University of Life Sciences, 166 Nowoursynowska Str. 02-787 Warsaw, Poland

Paulownia is used as a medicinal plant in traditional Chinese medicine against numerous diseases (Dzugań et al. 2021). Previous studies have reported that the use of *Paulownia* leaves (PL) is also suitable for feeding domestic animals and has similar nutritional values to alfalfa (Al-Sagheer et al 2019). PL are reported to be slightly bitter but palatable to sheep and cattle, can be used in feed for pigs without adverse effects, and are processed into pellets for fish and chicken feed containing up to 20% PL (Al-Sagheer et al 2019). The use of up to 15% PL meal in the diet of rabbits had no negative effects on performance, nutrient digestibility, and blood components, in addition, a remarkable reduction in pathogenic bacteria was observed in both the caecum and the feed itself (Al-Sagheer et al 2019). In feeding trial in sheep, the number of leukocytes and erythrocytes was significantly reduced and the blood glucose level decreased significantly (Vaslyakov et al. 2013). Studies with broilers showed that *Paulownia* polysaccharides can improve the immune system by enhancing cellular and humoral immunity (Wang et al. 2019). Enriched broiler feeds with 0.5 g PL/kg diet is the most acceptable dosage for supplementation in broilers to optimize performance, improve oxidative remarks activity, and enhance immunity without compromising productivity (Sakr, et al. 2022). However, further studies are needed to provide insights into the role of PL in alleviating oxidative stress, modulating intestinal microbial biodiversity, and mediating metabolic activities associated with gut function activity in animals.

Keywords: *Paulownia* leaves, animal feeding, health condition

References

- Al-Sagheer, A.A., Abd El-Hack, M.E., Alagawany, M., Naiel, M.A., Mahgoub, S.A., Badr, M.M., Hussein, E.O.S., Alowaimier, A.N.; Swelum, A.A. 2-19. *Paulownia* leaves as a new feed resource: chemical composition and effects on growth, carcasses, digestibility, blood biochemistry, and intestinal bacterial populations of growing rabbits. *Animals*, 9, 95.
- Dzugań, M., Miłek, M., Grabek-Lejko, D., Hęclik, J., Jacek, B., Litwińczuk, W., 2021. Antioxidant activity, polyphenolic profiles and antibacterial properties of leaf extract of various *Paulownia* spp. clones. *Agronomy* 11, 2001.
- Sakr, S.A., EL-Emam, H.A., Naiel, M.A.E., Wahed, N.M., Zaher, H.A., Abougabal, M.S., Alghamdi, Y.S., Albogami, S., Soliman, M.M., Shukry, M., Elghareeb, M.M., 2022. The impact of *Paulownia* leaves extract on performance, blood biochemical, antioxidant, immunological indices, and related gene expression of broilers. *Front Vet Sci.* 9: 882390.
- Varlyakov, I., Radev, T., Slavov, T., Gancho, S., Ganchev G., 2013. Blood parameters in yearling sheep fed *Paulownia* (*Paulownia* spp.) leaves. *Agricult. Sci. Technol.*, 5, 405-409.
- Wang, Q., Meng, X., Zhu, L., Xu, Y., Cui, W., He, X., Wei, K., Zhu, R., 2019. A polysaccharide found in *Paulownia fortunei* flowers can enhance cellular and humoral immunity in chickens. *Int. J. Biol. Macromol.* 1, 213-219.

* Presenting author. E-mail address: marta_mendel@sggw.edu.pl (Marta Mendel)

S2.P252 Exploring ethnoveterinary medicine in dogs in Ireland, Norway and Poland

Lorraine McKenna, Joanna Polak, Urszula Latek, Marta Mendel*

Department of Preclinical Sciences, Institute of Veterinary Medicine, Warsaw University of Life Sciences, 8, Ciszewskiego Str, 02-786 Warsaw, Poland

The study aimed to investigate the use of ethnoveterinary medicine by dogs' owners. It has been studied in various regions worldwide (Mayer et al., 2017), but ethnoveterinary medicine's application in Northern and Central-Eastern Europe remains relatively understudied (Disler et al., 2014). A survey was conducted in late 2023 and early 2024, in three countries: Ireland, Poland and Norway. A questionnaire about the use of medicinal plants (MP) and their preparations in pets, was distributed to dog owners who visited veterinary clinics. The results were not surprising; out of 102 responders less than 30% used medicinal plants as therapy in their dogs but in those that had used these therapies, there was a report of more than 85% in effectiveness. According to the survey, the owners learn about the use of medicinal plants mostly from internet and in case of Poland also from the veterinarians who take care of their pets. In total the owners indicated the use of 17, 5 and 8 different herbs in Poland, Ireland and Norway, respectively. Linseed, valerian and chamomile were the most frequently mentioned plants, whereas digestive issues, skin problems and anxiety were the most common circumstances of MP application. In general, the owners indicated interest but lack of knowledge about how to use herbal medicines. The survey indicates high need of promotion of veterinary herbal medicine among animal owners and urgent need of education of present and future veterinarians in phytotherapy.

Keywords: ethnoveterinary medicine, dogs, medicinal plants

References

- Mayer, M., Zbinden, M., Vogl, C. R., Ivemeyer, S., Meier, B., Amorena, M., Walkenhorst, M., 2017. Swiss ethnoveterinary knowledge on medicinal plants – a within-country comparison of Italian speaking regions with North-Western German speaking regions. *J. Ethnobiol. Ethnomed.* 13, 1.
- Disler, M., Ivemeyer, S., Hamburger, M., Vogl, C. R., Tesic, A., Klarer, F., Walkenhorst, M., 2014. Ethnoveterinary herbal remedies used by farmers in four North-Eastern Swiss cantons (St. gallen, Thurgau, Appenzell Innerrhoden and Appenzell Ausserrhoden). *J. Ethnobiol. Ethnomed.* 10, 1.

* Presenting author. E-mail address: marta_mendel@sggw.edu.pl (Marta Mendel)

S2.P253 Polish perspective on phytotherapeutic approaches to pet anxiety: an analysis of popular preparations

Joanna Polak*, Urszula Latek, Marta Mendel

Division of Pharmacology and Toxicology, Warsaw University of Life Sciences, Nowoursynowska 166, 02-787 Warsaw, Poland

Herbal medicines and supplements are gaining growing popularity among veterinarians and animal owners in Poland. They are thought to be effective, safe and environmentally friendly, with causes their high social acceptance. The common belief, especially among animal owners, is that remedies of natural origin have mild calming effect without causing serious sedation. This leads to many cases of self-introduced therapy of pets, as herbal supplements are also easily and widely accessible. There are several dozen preparations of this type available on the market in Poland.

This work aims to compare popular calming preparations for dogs and cats available on the market in Poland and their composition in terms of the substances of plant origin used and its concentration (as stated by the producer). They are available in different formulations: oral preparations, diffusers, sprays, calming collars, or rectal suppositories. We decided to narrow our list only to various oral preparations: tablets, twist off capsules, oils and suspensions. We analyzed 50 products, some of which were medicinal products, others complementary food or dietary supplements. All of them are available in pet stores and online without a prescription.

The compared products differed considerably in terms of composition and available information on the content of individual additives/substances. Most of them were multi-component. The most commonly used extracts were: *Valeriana officinalis*, *Melissa officinalis*, *Passiflora incarnata*, *Camellia sinensis* and *Chamomilla recutita*. The popularity of CBD oils is also noteworthy: among the 50 products analyzed, there were as many as 16 oils from various manufacturers.

Keywords: calming preparations, pet anxiety, herbal extracts, CBD oil

* Presenting author. E-mail address: joanna_polak@sggw.edu.pl (Joanna Polak)

S2.P254 Anti-tumor effects of *Auricularia polytricha* extracts on human breast cancer cell lines

Sunita Nilkhet^a, Chanin Sillapachaiyaporn^a, Kuljira Mongkolpobsina^{a,c}, Tewin Tencomnao^b, Siriporn Chuchawankul^{c,d,*}

^a Graduate program in Clinical Biochemistry and Molecular Medicine, Department of Clinical Chemistry, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^b Department of Clinical Chemistry, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^c Department of Transfusion Medicine and Clinical Microbiology, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^d Immunomodulation of Natural Products Research Unit, Chulalongkorn University, Bangkok 10330, Thailand

Auricularia polytricha (AP), a cloud ear mushroom, has been reported for several medicinal properties such as anti-inflammation (Xiang et al., 2021), anti-oxidation (Chiu et al., 2014), and immunomodulation (Yu et al., 2009) as well as anti-human lung cancer activity (Yu et al., 2014). In this study, anti-tumor activity of AP was demonstrated *in vitro* against human breast cancer cell lines. APH, APE and APW were extracted by sequential maceration with hexane, ethanol, and water, respectively. All extracts were investigated for their toxicity against human breast cancer cells (MDA-MB-231 and MCF-7). The cytotoxicity of APH on MDA-MB-231 and MCF-7 after 24 h of treatment showed IC₅₀ values at 24.31 ± 0.80 and 13.13 ± 1.60 mg/mL, respectively. Moreover, the biological results showed that the APH inhibited migration, invasion, and colony formation and induced apoptosis. In addition, the pro-apoptotic genes (*CASP3*, *CASP8* and *CASP9*) were upregulated by APH treatment on breast cancer cells. In conclusions, APH displayed the anti-tumor effect on breast cancer cells by apoptosis induction. These results will lead to the future development of drug discovery of AP.

Keywords: *Auricularia polytricha*, anti-tumor activity, apoptosis, colony formation

References

- Chiu, W.-C., Yang, H.-H., Chiang, S.-C., Chou, Y.-X., Yang, H.-T. *Auricularia polytricha* aqueous extract supplementation decreases hepatic lipid accumulation and improves antioxidative status in animal model of nonalcoholic fatty liver. *BioMedicine* 4 (2014).
- Yu, M. et al. Isolation of an anti-tumor polysaccharide from *Auricularia polytricha* (jew's ear) and its effects on macrophage activation. *European Food Research and Technology* 228, 477-485 (2009).
- Yu, J., Sun, R., Zhao, Z. & Wang, Y. *Auricularia polytricha* polysaccharides induce cell cycle arrest and apoptosis in human lung cancer A549 cells. *International journal of biological macromolecules* 68, 67-71 (2014).
- Xiang, H., Sun-Waterhouse, D. & Cui, C. Hypoglycemic polysaccharides from *Auricularia auricula* and *Auricularia polytricha* inhibit oxidative stress, NF-κB signaling and proinflammatory cytokine production in streptozotocin-induced diabetic mice. *Food Science and Human Wellness* 10, 87-93 (2021).

* Presenting author. E-mail address: siriporn.ch@chula.ac.th (Siriporn Chuchawankul)

S2.P255 Analysis of veterinary historical German-language textbooks regarding medicinal plants used for the treatment of gynecological diseases and fertility disorders

Luzia Steiner^a, Urs Jenny^b, Gaby Hirsbrunner^c, Michael Walkenhorst^{d,*}

^a Tierarztpraxis Luzia Steiner, Bahnmatt 27, 6340 Baar, Switzerland

^b Rietwiesstrasse 135, 8810 Horgen, Switzerland

^c Wiederkäuerklinik, Vetsuisse-Fakultät, Universität Bern, Länggassstrasse 120, 3012 Bern, Switzerland

^d Forschungsinstitut für biologischen Landbau FiBL, Ackerstrasse 113, 5070 Frick, Switzerland

Diseases of the female genital tract cause antibiotic and hormonal treatments as well as high culling rates in cattle. We analyzed five books published in Germany and Switzerland between 1878 and 1921, and handwritten notes of the veterinarian Ammann-Honegger (1879–1960) (Steiner et al., 2022). We systematically examined the sources regarding use reports (UR) based on medicinal plants to treat gynecological diseases. Herbal ingredients, target animal species, type of administration and indication were detailed documented for each UR. The sources contained 103 UR (79 administered orally (o), 13 locally (l), and 11 both o and l). Fifty-five different plant species were recorded. The most frequently mentioned medicinal plants were *Juniperus communis* L. (19 UR, l and o), *Linum usitatissimum* L. (18 UR, o), *Juniperus sabina* L. (13 UR, l and o), *Matricaria chamomilla* L. (13 UR, l and o) and *Gentiana lutea* L. (12 UR, o). *Retentio secundinarum* was most frequently mentioned as indication (44 UR), followed by parturition preparation (17 UR) and endometritis (15 UR). Based on their ingredients, the recorded plant species could be divided into (a) high-quality fodder, (b) appetizer and digestives, (c) specific gynecologicals and (d), predominantly toxic plants. Active defense against bacterial infections as well as a stable barrier of the endometrium contribute both to uterine health (Sheldon et al., 2019). The plants listed under a)-c) could at least indirectly support these functions.

References

- Sheldon, I.M., Cronin, J.G., Bromfield, J.J. (2019). Tolerance and Innate Immunity Shape the Development of Postpartum Uterine Disease and the Impact of Endometritis in Dairy Cattle. *Annu Rev Anim Biosci* 7, 361–384
- Steiner, L., Jenny, U., Hirsbrunner, G., Walkenhorst, M. (2022). Phytotherapeutische Behandlungen gynäkologischer Erkrankungen und Fruchtbarkeitsstörungen von Rindern – eine veterinärhistorische Analyse. *Schweizer Archiv für Tierheilkunde* 164: 645-659. doi.org/10.17236/sat00367

* Presenting author. E-mail address: michael.walkenhorst@fibl.org (Michael Walkenhorst)

S2.P256 Phyto-synergy: a holistic approach to tackle *paratuberculosis*

Varsha Srivastava^{a,*}, Manthena Navabharath^b, Saurabh Gupta^b, Shoor Vir Singh^b, Sayeed Ahmad^a

^a Centre of excellence in Unani Medicine (Pharmacognosy and Pharmacology), Bioactive Natural Product Laboratory, Department of Pharmacognosy and Phytochemistry, School of Pharmaceutical Education and Research, Jamia Hamdard, New Delhi 110062, India

^b Department of Biotechnology, Institute of Applied Sciences & Humanities, GLA University, Mathura 281406, Uttar Pradesh, India

The presented work will provide a comprehensive overview of our research methodology, key findings associated with the development of a potential herbal therapy against *Mycobacterium avium* subspecies *paratuberculosis* (MAP). MAP is considered to be the causative agent of Johne's disease, an intestinal chronic infection in ruminants (Feller et al., 2007). These bacilli upon transmission through milk and milk products lead to development of Crohn's disease like autoimmune disorders in human beings (Naser et al., 2004). Till date, no reliable treatments are available for this pathogenic infection.

The conducted research focuses on the development of a novel herb-based synergistic combination as a potential anti-MAP therapy. Our study involved screening a diverse range of medicinal herbs for their anti-MAP potential using *in vitro* Resazurin microtitre assay (REMA). The obtained best active extracts (BAEs) i.e. hydroalcoholic extracts of *O. sanctum* and *S. xanthocarpum* were further assessed for synergy using fractional inhibitory concentration index (FICI). Subsequently, a synergistic combination was formulated to enhance the overall efficacy through the potentiation of individual herb properties. The results demonstrate significant antimycobacterial activity of the herb-based synergistic combination against MAP, with no significant cytotoxic effects on host cells. Moreover, the combination exhibited promising immunomodulatory properties, suggesting a dual mechanism of action for enhanced efficacy.

Keywords: *Mycobacterium avium* subspecies *paratuberculosis*, Johne's disease, Crohn's disease, REMA assay

References

- Feller, M., Huwiler, K., Stephan, R., Altpeter, E., Shang, A., Furrer, H., et al. (2007). *Mycobacterium avium* subspecies *paratuberculosis* and Crohn's disease: a systematic review and meta-analysis. *Lancet Infect. Dis.* doi: 10.1016/S1473-3099(07)70211-6.
- Naser, S. A., Ghobrial, G., Romero, C., and Valentine, J. F. (2004). Culture of *Mycobacterium avium* subspecies *paratuberculosis* from the blood of patients with Crohn's disease. *Lancet.* doi: 10.1016/S0140-6736(04)17058-X.

* Presenting author. E-mail address: varshasrivastav2796@gmail.com (Varsha Srivastava)

S2.P257 Preliminary study of the neuroprotective capacity of *Corema album* (L.) D.Don ex Steud (Ericaceae) juice and its butanolic fraction

Antonio Canoyra^a, Nuria Acero^{a,*}, Dolores Muñoz-Mingarro^b, Carmen Martín-Cordero^c, Antonio-José León-González^c

^a Departamento de CC Experimentales y de la Salud, Facultad de Farmacia, Universidad San Pablo-CEU, CEU Universities, Urb. Montepríncipe, 28668 Boadilla del Monte, Madrid, Spain

^b Departamento de Química y Bioquímica, Facultad de Farmacia, Universidad San Pablo-CEU, CEU Universities, Urb. Montepríncipe, 28668 Boadilla del Monte, Madrid, Spain

^c Departamento de Farmacología. Faculty of Pharmacy, Seville University, C/ P. García González 2, 41012 Seville, Spain

Corema album (L.) D.Don ex Steud (Ericaceae) berries are an edible fruit which have traditionally been consumed in those geographical areas where the plant is endemic (Iberian Peninsula). In the present study, the ability of this fruits juice and its butanolic extract to decrease ROS levels in SHSY-5Y cell line culture was analysed. Both its direct effect on this cell line, and its preventive effect against H₂O₂-induced oxidative stress were assessed (León-González *et al.*, 2012). On the other hand, SH-SY5Y cells were differentiated into dopaminergic (Retinoic acid) and cholinergic (BDNF), to replicate an *in vitro* model of both Parkinson's and Alzheimer's disease respectively (de Medeiros *et al.*, 2019). Dopaminergic differentiated SHSY-5Y were stressed with 6-hydroxydopamine which induce a massive oxidative stress in this cell line (Lopes *et al.*, 2017) in the presence or absence of *C. album*. On the other hand, for mimicking tau and amyloid- β pathology associated with Alzheimer's disease, BDNF-differentiated cells were treated with okadaic acid, with or without the juice and the extract of the fruits. The plant capacity to increase cell viability under these toxic conditions was studied. Finally, the inhibitory effect of juice and butanolic juice extract on different enzymes related to neuroprotection was also investigated. The results show an interesting neuroprotective capacity of *C. album* by being able to reduce induced oxidative stress, to inhibit MAO-B *in vitro*, and to increase the survival of differentiated cholinergic or dopaminergic cells under toxicity conditions. UHPLC-MS/MS analysis of the juice and the extract was performed.

Keywords: *Corema album*, SHSY-5Y, neuroprotection, MAO, antioxidant

References

- León-González, A.J. Mateos, R. Ramos, S. Martín, M.T. Sarriá, B. Martín-Cordero, C. López-Lázaro, M. Bravo, L. Goya, L. 2012. Chemo-protective activity and characterization of phenolic extracts from *Corema album*. Food Res Int., 49, 728-738.
- de Medeiros, L. M., De Bastiani, M. A., Rico, E. P., Schonhofen, P., Pfaffenseller, B., Wollenhaupt-Aguiar, B., Grun, L., Barbé-Tuana, F., Zimmer, E. R., Castro, M. A. A., Parsons, R. B., Klamt, F. 2019. Cholinergic differentiation of human neuroblastoma SH-SY5Y cell line and its potential use as an *in vitro* model for Alzheimer's disease studies. Mol. Neurobiol, 56(11), 7355-7367.
- Lopes, F. M., da Motta, L. L., De Bastiani, M. A., Pfaffenseller, B., Aguiar, B. W., de Souza, L. F., Zanatta, G., Vargas, D. M., Schönhofen, P., Londero, G. F., de Medeiros, L. M., Freire, V. N., Dafre, A. L., Castro, M. A., Parsons, R. B., Klamt, F. 2017. RA Differentiation enhances dopaminergic features, changes redox parameters, and increases dopamine transporter dependency in 6-hydroxydopamine-induced neurotoxicity in SH-SY5Y cells. Neurotox Res, 31(4), 545-559.

* Presenting author. E-mail address: nacemes@ceu.es (Nuria Acero)

S2.P258 Anti-inflammatory and antioxidant ability of *Vitis vinifera* L. leaves

Dolores Muñoz-Mingarro^a, Teresa Ortega^b, Nuria Acero^{c,*}, Olga Palomino^b, Francisco Bosch-Morell^c, Isabel Martínez Solís^d

^a Departamento de Química y Bioquímica, Universidad San Pablo-CEU, CEU Universities, Boadilla del Monte, Madrid, Spain

^b Departamento de Farmacología, Farmacognosia y Botánica, Facultad de Farmacia, Universidad Complutense de Madrid, Spain

^c Departamento de CC Farmacéuticas y de la Salud, Universidad San Pablo-CEU, CEU Universities, Boadilla del Monte, Madrid, Spain

^d Departamento de CC Biomédicas, Universidad Cardenal Herrera-CEU, CEU Universities, Alfara del Patriarca, Valencia, Spain

^e Departamento de Farmacia, Universidad Cardenal Herrera-CEU, CEU Universities, Alfara del Patriarca, Valencia, Spain

Vitis vinifera L. is a species belonging to the Vitaceae family, which fruit has been extensively studied due to its use since ancient times. Grapes and their seeds are considered source of pharmacological and cosmetic products. However, other parts of the plant such as the leaves, have been less studied, despite having been used for different therapeutic purposes (Mansour *et al.* 2013). Currently, the European Medicines Agency accepts the well-established use of a dried aqueous grapevine leaf extract for the treatment of chronic venous insufficiencies symptoms. It is interesting to deepen the study of this part of the plant to take advantage of the large quantities of them that are generated as agricultural waste (Maia *et al.*, 2018). In the present study, the anti-inflammatory ability of a methanolic extract of the *V. vinifera* leaves has been analysed. Radical scavenging ability *in vitro* and in cell-based assays was determined. The direct effect and the protective effect of the extract after induced oxidative stress were assessed in HepG2 cell line. On the other hand, its anti-inflammatory potential was evaluated through the impact upon NO levels produced by RAW 264.7 stimulated with *E. coli* lipopolysaccharides. In addition, different concentrations of the extract were approached on its effect upon lipoxygenase activity *in vitro*. The phytochemical profile of the extract was analyzed using UHLC-QTOF-MS. Results showed an interesting antioxidant and anti-inflammatory ability of the leaves, pointing out this part of the grapevine plant as a valuable resource for phytochemical metabolites of pharmacological and nutritional value.

Keywords: *Vitis vinifera*, antioxidant capacity, anti-inflammatory, phenolic content

References

- Mansour, R., Haouas, N., Ben Kahla-Nakbi, A., Hammami, S., Mighri, Z., Mhenni, F., & Babba, H. 2013. The effect of *Vitis vinifera* L. leaves extract on *Leishmania infantum*. IJPR, 12(3), 349–355.
- Maia, M., Ferreira, A. E. N., Laureano, G., Marques, A. P., Torres, V. M., Silva, A. B., Matos, A. R., Cordeiro, C., Figueiredo, A., Sousa Silva, M. (2019). *Vitis vinifera* 'Pinot noir' leaves as a source of bioactive nutraceutical compounds. Food Funct., 10(7), 3822–3827.

* Presenting author. E-mail address: nacemes@ceu.es (Nuria Acero)

S2.P259 Pre- and post- treatment effect of water-soluble phenolics from date palm (*Phoenix dactylifera* L.) fruits in cisplatin-induced hepatotoxicity in rats

Omowumi Adewale^{a,*}, Roseline Oyelola^a, Adeola Adenmosun^a, Oluwaseun Adebisi^{a,b}, Isaac Babatunde^a, Adedayo Adekomi^c, Johnson Oladele^d

^a Department of Biochemistry, Osun State University, P.M.B. 4494, Osogbo, Osun State, Nigeria

^b Department of Biochemistry and Molecular Biology, University of Nebraska Medical Center, Omaha, NE, USA

^c Department of Anatomy, Faculty of Basic Medical Sciences, Osun State University, Osogbo, Nigeria

^d Veterinary Physiology and Pharmacology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, Tx, USA

This presentation will cover the pre- and post-treatment for the effects of water-soluble phenolics from date (*Phoenix dactylifera* L) fruits (DFP) on cisplatin-induced hepatotoxicity in rats. Natural products have been proposed to be better agents in the suppression of chemotherapy-induced toxicities due to their inherent bioactive values and lack of toxicity. (Asma et al., 2022; Dziadek et al., 2019; Zhang et al., 2018).

As part of continuing search for novel natural products with potent bioactive principles capable of suppressing toxicities associated with cancer therapy, we employed the pre- and post-treatment strategies with DFP on cisplatin-induced rats. Treatment of rats with DFP before or after exposure to a single therapeutic dose of cisplatin (5mg/kg) led to enhanced antioxidant status, in addition to significant ($p < 0.05$) suppression of lipid peroxidation in cisplatin-induced rats. Furthermore, a concomitant reduction in activities and levels of liver function markers was observed after treatment with DFP. Photomicrographs from hematoxylin/eosin stained liver tissues revealed the reversal of cisplatin-induced pathology at both time points. Chromatographic characterization of DFP revealed peaks of phenolics that have been widely reported for medicinal values (Fig. 1). DFP at both time points suppressed cisplatin-induced hepatotoxicity in this study, though, pre-treatment strategy showed greater significance than post-treatment strategy. Hence, pre-treatment with DFP may provide protection from the hepatic damage induced as a result of cisplatin-based chemotherapy.




Fig. 1: GC-MS chromatogram of Date Fruit phenolics (DFP)

Keywords: hepatotoxicity, cisplatin, DFP, liver markers, antioxidant status

References

- Asma, S.T., Acaroz, U., Imre, K., Morar, A., Shah, S.R.A., Hussain, S.Z., Arslan-Acaroz, D., Demirbas, H., Hajrulai-Musliu, Z., Istanbulgil, F.R., Soleimanzadeh, A., 2022. Natural products/bioactive compounds as a source of anticancer drugs. *Cancers*, 14(24), 6203.
- Dziadek, K., Kopeć, A., Tabaszewska, M., 2019. Potential of sweet cherry (*Prunus avium* L.) by-products: bioactive compounds and antioxidant activity of leaves and petioles. *European Food Research and Technology*, 245(3), 763-772.
- Zhang, Q. Y., Wang, F. X., Jia, K. K., & Kong, L. D. (2018). Natural product interventions for chemotherapy and radiotherapy-induced side effects. *Front Pharmacol* 9: 1253.

* Presenting author. E-mail address: omowumi.adewale@uniosun.edu.ng (Omowumi Adewale)

S2.P260 Determination of anticancer properties of *Moringa oleifera* seed bioactive fraction and the interaction effect of its combination with the classical cytotoxic drugs

Patrycja Wińska^a, Joanna Cieśla^a, Jan Zając^a, Siriya Faith^a, Eryk Grzechnik^b, Omowumi Adewale^{a,c,*}

^a Chair of Drug and Cosmetics Biotechnology, Faculty of Chemistry, Warsaw University of Technology, 00-664 Warsaw, Poland

^b Chair of Polymer Chemistry and Technology, Faculty of Chemistry, Warsaw University of Technology, Nowakowskiego 3, 00-664 Warsaw, Poland

^c Cancer Research and Molecular Toxicology Laboratory, Department of Biochemistry, Osun State University, Osogbo, Nigeria

This presentation will cover the determination of anticancer properties of *Moringa oleifera* seed bioactive fraction (MSBF) and the interaction effect of its combination with 5-Fluoro-uracil, Cisplatin or CX-4945 (CK2-inhibitor) in breast and colon cancer cell lines. *Moringa oleifera* is known for its diverse range of biological activities including anticancer. Recently, drug combination strategy has been recommended to defeat the associated challenges with chemotherapy and combination of natural products with conventional anticancer drugs has been supported by several studies (Naeem et al., 2022; Gilad et al., 2021; Talib et al., 2021)

As part of continuing search for novel natural products as drug combination candidate in cancer therapy, we investigated the interaction effects of MSBF with 5-fluorouracil, Cisplatin or CX-4945 (a CK2-inhibitor) in breast cancer (MDA-MB-231 (triple-negative breast cancer (TNBC)) and MCF-7 (estrogen receptor-positive (ER+)) and colon cancer (HCT-116 and HT-29) cell lines. MSBF was obtained through defatting, depigmentation and ethanol precipitation. The results reveal the synergistic interaction between MSBF with 5-FU, or Cisplatin, while antagonism was observed between MSBF with CX-4945 in both cancer cells lines. However, treatment with MSBF/5-FU combination potentiated the highest significant ($P < 0.05$) DRI (103.7 ± 5.63) for 5-FU at F_a of 0.96 with CI, 0.72 ± 0.01 in MDA-MB-231 (Fig.1). Assessment of the effect of MSBF /5-FU treatment on pro-apoptotic properties and cell cycle arrest on MDA-MB-231 cell lines is ongoing to understand the mechanism behind this interaction. Hopefully, treatment with MSBF/5-FU combination may cause dose reduction and consequently, reduction in toxicity of 5-fluorouracil, thereby, improving chemotherapy.




Fig 1. Synergistic interaction between MSBF and (5-FU) in MDA-MB-231 cell lines

Keywords: Chemotherapy, *Moringa oleifera* seed, 5-fluorouracil, combination index, synergism

References

- Gilad, Y., Gellerman, G., Lonard, D. M., O'malley, B. W., 2021. Drug combination in cancer treatment—From cocktails to conjugated combinations. *Cancers*, 13(4), 669.
- Talib, W. H., Alsayed, A. R., Barakat, M., Abu-Taha, M. I., Mahmood, A. I., 2021. Targeting Drug Chemo-Resistance in Cancer Using Natural Products. *Biomedicines*, 9, 1353.
- Naeem, A., Hu, P., Yang, M., Zhang, J., Liu, Y., Zhu, W., Zheng, Q. 2022. Natural products as anticancer agents: Current status and future perspectives. *Molecules*, 27(23), 8367.

* Presenting author. E-mail address: omowumi.adewale@pw.edu.pl (Omowumi Adewale)

S2.P261 Fertility parameters of dairy cows after intrauterine treatment of clinical endometritis with an antibiotic or an herbal veterinary medicinal product

Victoria Scheuber^{a,b}, Valérie Menoud^{a,b,c}, Ariane Maeschli^a, Mirjam Holinger^a, Sandra Graf-Schiller^d, Philipp Mayer^d, Luc Gerber^c, Gaby Hirsbrunner^b, Michael Walkenhorst^{a,*}

^a Research Institute of Organic Agriculture (FiBL), Frick, Switzerland

^b Vetsuisse Faculty, University of Bern, Bern, Switzerland

^c Clinique du Vieux-Château/JuraVet, Delémont, Switzerland

^d SaluVet GmbH, Bad Waldsee, Germany

Reproductive disorders represent challenging problems for dairy cattle. We compared fertility parameters of cows after the intrauterine treatment of clinical endometritis (CE) with cephapirin (Metricure®; cefapirin benzathin 500 mg per dosis; CEPH) or an herbal product (25 ml of EucaComp®PlantaVet containing alcoholic extracts of *Calendula officinalis* L., *Mellissa officinalis* L., *Origanum majorana* L. and the essential oil of *Eucalyptus globulus* Labill. (EUC)).

Totally, 169 cows between 21 and 35 days after calving were included and randomly assigned to one of the both treatment groups (Menoud et al., 2024). For final analysis, the reproductive performance of 140 cows (73 CEPH and 67 EUC) from 27 farms was monitored from 24 days after the first treatment until the next calving or culling date. No significant differences between treatment groups were found regarding first service conception rate, calving to conception interval for cows pregnant at 120 days p.p., pregnancy rate at 120 days p.p., number of inseminations per calving, intercalving period, calving rate, culling reasons and number of gynaecological treatments later than 24 days after the first treatment of the clinical endometritis. There was a trend ($p=0.06$) for a slightly longer first service period in the treatment group for CEPH (80 days postpartum (p.p.)) compared to EUC (74 days p.p.). In conclusion, EUC is with a high probability not inferior to CEPH. This finding could help to reduce the use of antimicrobials in the daily routine veterinary treatment of CE.

Keywords: Veterinary phytotherapy, on farm study, clinical endometritis treatment

References

Menoud, V., Holinger, M., Graf-Schiller, S., Mayer, P., Gerber, L., Walkenhorst, M., Hirsbrunner, G., 2024. Comparison between intrauterine application of an antibiotic and an herbal product to treat clinical endometritis in dairy cattle – A randomized multicentre field study. Res. Vet. Sci. In Press, doi.org/10.1016/j.rvsc.2024.105250

* Presenting author. E-mail address: michael.walkenhorst@fibl.org (Michael Walkenhorst)

S2.P262 Medicinal plants for animal health care: translating tradition into modern veterinary medicine - the COST Action MedPlants4Vet

Theresa Schlittenlacher^a, Marta Mendel^{b,*}, Michael Walkenhorst^a

^a Research Institute of Organic Agriculture (FiBL), Ackerstrasse 113, post box 2019, 5070 Frick, Switzerland

^b Department of Preclinical Sciences, Institute of Veterinary Medicine, Warsaw University of Life Sciences, 8, Ciszewskiego St, 02-786 Warsaw, Poland

The demand for natural products supporting animal health and welfare is increasing in the post-antibiotic era. This is in line with the goals of national and international action plans on antimicrobial resistance, the One Health paradigm and the European green deal. It addresses not only farm animals, but also companion animals living in proximity with their owners. Herbal veterinary medicinal products (HVMP) are instrumental in achieving these goals. In the current EU legislation, HVMP are only mentioned to the extent that there is insufficient information for a simplified authorization (Regulation (Eu) 2019/6).

Therefore, the interdisciplinary and international network MedPlants4Vet was founded to identify and close significant gaps in knowledge about the use of traditional HVMP. The network consists of over 250 people from science, industry, practice and regulatory authorities from around 39 countries. MedPlants4Vet aims to significantly promote interdisciplinary cooperation as well as professional training and further education. The creation of a comprehensive decision tree that could serve as a guide for future regulations and authorization procedures is a key focus. In line with current EU legislation, the creation of a basis for a simplified registration of traditional HVMP at EU level with a corresponding legal framework is a matter of priority. In human medicine, a corresponding legal framework was already established in 2004 (Directive 2004/24/Eg).

To achieve these goals, MedPlants4Vet is organized in six working groups (WG; Fig. 1).



Fig. 1. MedPlants4Vet is organized in six working groups.

Further information is available on the homepages www.medplants4vet.eu and www.cost.eu/actions/CA22109/. MedPlants4Vet (CA22109) is supported by COST (European Cooperation in Science and Technology).

Keywords: herbal veterinary medicinal products, legal framework, ethnoveterinary research - veterinary phytotherapy, phytochemistry

References

Regulation (Eu) 2019/6 Of The European Parliament And Of The Council of 11

December 2018: <https://eur-lex.europa.eu/eli/reg/2019/6/oj>

Directive 2004/24/Eg Of The European Parliament And Of The Council: <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:136:0085:0090>

* Presenting author. E-mail address: marta_mendel@sggw.edu.pl (Marta Mendel)

S2.P263 Effectiveness of replacing antibiotics with botanicals in the treatment of bovine mastitis

Marcela Šperanda^{a,*}, Mislav Đidara^b, Berislav Vulić^c

^a Central Agrobiotechnical Analytical Unit Faculty of Agrobiotechnical Science Osijek, Vladimira Preloga 1, Osijek, Croatia

^b Department of Livestock Production and Biotechnology Faculty of Agrobiotechnical Science Osijek, Osijek, Croatia

^c Dairy farm Salaš, J. J. Strossmayer street 111, Marijanci, Croatia

Due to the growing resistance to antibiotics in the human population, it is necessary to use antibiotics prudently in controlling diseases in humans and animals. Bovine mastitis is considered the most common disease leading to economic loss in dairy industries. The goal of this research was to check whether antibiotic therapy can be successfully replaced with botanicals that inhibit biofilm formation (Singh et al., 2017), to reduce or disrupt quorum sensing (Boyen et al., 2009) in the mammary gland.

Cows with clinical signs of mastitis in one or more quarters were divided into two groups. The first was treated intramammary beta-lactam antibacterial Ampicillin sodium+Cloxacillin sodium (A), second with AHV Extra bolus (AHV International) for per os one-time (B). Some cows from the first treatment got combined therapy intramammary with Amoxicillin 150 mg/ml parenterally, (A+A). On the first day after the appearance of clinical signs of the disease and day 5th following parameters in milk were determined: somatic cell count (SCC), haptoglobin, lactoferrin, total protein, albumin, and aspartate aminotransferase (AST) activity.

Lower ($P<0.05$) SCC and haptoglobin concentration after 5 days of treatment were found in cows from the B and A+A groups. Considering that in the A+A group, the therapy was for 4 days, and in the B group only once, we conclude that botanicals had a prominent effect in treating mastitis following standard parameters like SCC and haptoglobin in milk. The use of antibiotics did not appear to be a factor of choice. The potential of supportive therapy with botanicals should be explored.

Keywords: bovine mastitis, botanicals, antimicrobials

References

- Boyen F., Eeckhaut V., Van Immerseel F., Pasmans F., Ducatelle R., Haesebrouck F. Quorum sensing in veterinary pathogens: mechanisms, clinical importance and future perspectives. *Vet Microbiol.* 2009 Mar 30;135(3-4):187-95.
- Singh A.K., Prakash P., Singh R., Nandy N., Firdaus Z., Bansal M., Singh R.K., Srivastava A., Roy J.K., Mishra B., Singh R.K., 2017. Curcumin Quantum Dots Mediated Degradation of Bacterial Biofilms. *Front Microbiol.* 9;8:1517.

* Presenting author. E-mail address: marcela.speranda@fazos.hr (Marcela Šperanda)

S2.P264 Pumpkin seed cake waste as a sustainable functional feed ingredient

Stefania Marzorati, Alessio Massironi, Alessandro Guerrini, Doriana Eurosia Angela Tedesco*

Department of Environmental Science and Policy, Via Celoria 2, 20133 Milan, University of Milan, Italy

The valorization of biomass residuals as feed ingredients is a key aspect of the circular economy. This study aimed to evaluate the characteristics of pumpkin (*Cucurbita pepo* L.) seed waste, commonly called pumpkin seed cake in the feed industry, obtained after cold-press oil extraction.

Using green chemistry methodologies, the biomass was characterized in its apolar and polar fractions by supercritical CO₂ and water extraction, respectively. The qualitative and quantitative characterization was conducted using GC-MS and UPLC-PDA. Klason lignin (KL), and acid-soluble lignin (ASL) were determined following the ISO standards (ISO, 2020). ABTS assay was used for the antioxidant capacity evaluation.

Among apolar components, fatty acids composition indicates high relative percentage (81%) of unsaturated fatty acids, with a higher percentage of oleic (36%) and linoleic acid (45%). The unsaponifiable fraction (4.4% of the apolar extract), was represented by 97-98% of sterols (spinasterol, $\Delta^7,22,25$ -stigmastatrienol, $\Delta^7,25$ -stigmastadienol, β -sitosterol), and squalene traces (3%). The characterization of the polar fraction detected the presence of cucurbitin at 1% of the extract, 54.51 mg/g_{biomass} of total lignin, 1.98 mg/g_{biomass} of KL, and 52.52 mg/g_{biomass} of ASL. The antioxidant assay showed a higher antioxidant capacity of the polar fraction ($EC_{50} = 0.075$ mg/mL), compared to the apolar fraction ($EC_{50} = 0.24$ mg/mL), confirming that the presence of added-value compounds could be beneficial in the development of sustainable feed ingredients. *In vivo* studies are needed to expand and confirm the effectiveness of pumpkin seed cake waste in the perspective of the use of this waste as a functional feed ingredient.

Acknowledgements

This study was carried out within the Agritech National Research Center and received funding from the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR)–MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4–D.D. 1032 17/06/2022, CN00000022). This manuscript reflects only the authors' views and opinions, neither the European Union nor the European Commission can be considered responsible for them.

Keywords: *Cucurbita pepo* L., biomass valorization, sustainable feed

References

- Massironi, A., Biella, S., de Moura Pereira, P.F., Scibona, F., Feni, L., Sindaco, M., Emide, D., Jim'enez-Quero, A., Bianchi, C.L.M., Verotta, L., S. Marzorati, S., 2023. Valorization of pumpkin seed hulls, cucurbitin extraction strategies and their comparative life cycle assessment. *J. Clean. Prod.* 427 (2023) 139267.
- International Organization for Standardization (ISO)., 2020. Pulps—Determination of Lignin Content—Acid Hydrolysis Method; *ISO 21436*; ISO: Geneva, Switzerland.

* Presenting author. E-mail address: doriana.tedesco@unimi.it (Doriana Eurosia Angela Tedesco)

S2.P265 Herbal bioenhancers: an example of inhibition of ABCB1 using *Brugmansia* tincture

Begum Yurdakok-Dikmen^a, Sevcan Yangın^b, Gizem Erdogan^c, Sinem Aslan-Erdem^d, Semanur Yurduseven^a, Ayhan Filazi^a

^a Department of Pharmacology and Toxicology, Faculty of Veterinary Medicine, Ankara University, 06070, Ankara, Türkiye

^b Graduate School of Natural and Applied Sciences, Ankara University, 06070, Ankara, Türkiye

^c Graduate School of Health Sciences, Ankara University, 06070, Ankara, Türkiye

^d Department of Pharmacognosy, Faculty of Pharmacy, 06560, Ankara, Türkiye

The therapeutic efficacy of numerous pharmacological agents is often compromised by the active efflux of drugs across cellular membranes, mediated by the ATP-binding cassette (ABCB1) transporters (Yurdakok-Dikmen et al., 2018). This study explores the potential of *Brugmansia* (angel's trumpet) tincture, derived from *Brugmansia suaveolens* (Willd.) Sweet known for its tropane alkaloids majorly scopolamine (De Feo, 2004; Oğraş et al., 2022), as a bioenhancer through the inhibition of ABCB1 expression *in vitro*. *Brugmansia suaveolens*, belonging to the Solanaceae family, is widespread throughout the world, mainly in areas with tropical, subtropical, and temperate climates, both as a spontaneous species and as an ornamental (Petricevich et al., 2020). Using the L929 mouse fibroblast cell line as a model system, we investigated the effects of the tincture on the mRNA expression levels of ABCB1 using quantitative PCR (qPCR). We applied ethanolic tincture at 9.68-0.005 mg/ml concentrations and determined percent cytotoxicity regression curve by MTT. At tested concentrations (Inhibitory Concentration₁₅₋₂₅) a dose dependent decrease by 68.57-76.27% of mRNA expression at 302.42- 604.85 ug/ml concentrations were observed; which is also confirmed at protein level by western blot assays. Our preliminary results indicate a dose-dependent decrease in ABCB1 expression upon treatment with *Brugmansia* tincture, suggesting its potential role as a bioenhancer by facilitating greater intracellular drug accumulation. These findings provide a promising foundation for further research into the application of natural compounds in overcoming drug resistance mechanisms, potentially leading to more effective therapeutic strategies.

Keywords: *Brugmansia* tincture, L929 cell line, ABCB1 expression, *Brugmansia suaveolens*

References

- De Feo, V., 2004. The ritual use of *Brugmansia* species in traditional Andean medicine in northern Peru. *Economic Botany*, 58(1), pp.S221-S229.
- Oğraş, T.T., Tahtasakal, E. and ÖZTÜRK, S., 2022. In vitro production of tropane alkaloids from *Brugmansia suaveolens*. *International Journal of Secondary Metabolite*, 9(1), pp.1-13.
- Yurdakok-Dikmen, B., Turgut, Y. and Filazi, A., 2018. Herbal bioenhancers in veterinary phytotherapy. *Frontiers in Veterinary Science*, 5, p.249.
- Petricevich, Vera L., David Osvaldo Salinas-Sánchez, Dante Avilés-Montes, Cesar Sotelo-Leyva, and Rodolfo Abarca-Vargas. 2020. "Chemical Compounds, Pharmacological and Toxicological Activity of *Brugmansia suaveolens*: A Review" *Plants* 9, no. 9: 1161.

* Presenting author. E-mail address: byurdakok@yahoo.com (Begum Yurdakok-Dikmen)

S2.P266 Antibacterial activity of essential oils against streptococci and staphylococci causing bovine mastitis

Alice Caneschi^{a,*}, Donatella Scarafile^b, Anna Zaghini^a, Paola Mattarelli^b, Maurizio Scozzoli^c

^a Department of Veterinary Medical Sciences, Alma Mater Studiorum-University of Bologna, Via Tolara di Sopra 50, 40064 Bologna, Italy

^b Department of Agricultural and Food Sciences, University of Bologna, Viale Fanin 42, 40127 Bologna, Italy. University

^c Italian Society for Research on Essential Oils (Società Italiana per la Ricerca sugli Oli Essenziali-SIROE), Rome, RM, Italy

Bovine mastitis is a major concern for the dairy cattle community worldwide, leading to high economic loss (Gomes et al., 2016). Mastitis, subclinical or clinical, can be caused by contagious or environmental pathogens. Although the use of antibiotics is still the primary approach for treatment, its efficacy is limited. Moreover, the development of antimicrobial resistance (AMR) leads to the need for alternatives to antibiotic therapy (Yang et al., 2019). Due to their antibacterial effects, essential oils (EOs) have gained much attention in this field as candidates to fight against bacterial infections and control further development of AMR (Arbab et al., 2022).

This research aimed at testing the antibacterial activity of eleven EOs and two EO blends against *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Streptococcus uberis*, *Staphylococcus aureus* and *Staphylococcus epidermidis*, by evaluating the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC), using microbroth dilution assay. The MIC values of the EOs tested ranged between <0.001% and >3.125% v/v. The effect was higher against *Streptococcus spp* than *Staphylococcus spp* bacteria in general. The EOs of *Origanum vulgare*, *Cinnamomum zeylanicum*, *Thymus vulgaris*, and Blend BR were the most effective against *S. agalactiae*, *S. dysgalactiae*, and *S. uberis* showing MIC and MBC values between <0.001% and 0.390% v/v, and <0.001% and 0.780% v/v, respectively. Against *S. aureus* and *S. epidermidis*, the lowest MICs occurred with *Cinnamomum zeylanicum*: 0.098% and 0.195% v/v, respectively.

This study highlights the importance of evaluating EOs as effective antibiotic alternatives and complementary resources for treatment of bacterial infections.

Keywords: Essential oils, bovine mastitis, antimicrobial resistance, streptococci, staphylococci

References

- Arbab, Safia, Hanif Ullah, Iqra Bano, Ka Li, Inam Ul Hassan, Weiwei Wang, Abdul Qadeer, e Jiyu Zhang. 2022. «Evaluation of in Vitro Antibacterial Effect of Essential Oil and Some Herbal Plant Extract Used against Mastitis Pathogens». *Veterinary Medicine and Science* 8 (6): 2655–61. <https://doi.org/10.1002/vms3.959>.
- Gomes, Fernanda, e Mariana Henriques. 2016. «Control of Bovine Mastitis: Old and Recent Therapeutic Approaches». *Current Microbiology* 72 (4): 377–82. <https://doi.org/10.1007/s00284-015-0958-8>.
- Yang, Wan-Ting, Chun-Yen Ke, Wen-Tien Wu, Ru-Ping Lee, e Yi-Hsiung Tseng. 2019. «Effective Treatment of Bovine Mastitis with Intramammary Infusion of *Angelica dahurica* and *Rheum officinale* Extracts». *Evidence-based Complementary and Alternative Medicine: eCAM* 2019: 7242705. <https://doi.org/10.1155/2019/7242705>.

* Presenting author. E-mail address: alice@greenvet.com (Alice Caneschi)

S2.P267 Absorption, distribution, accumulation and excretion of thymol after its sustained administration in rabbits

Kristina Bacova^{a,*}, Karin Zitterl Eglseer^b, Gesine Karas Räuber^b, Iveta Placha^a

^a Centre of Biosciences, Slovak Academy of Sciences, Institute of Animal Physiology, Soltesovej 4–6, 040 01 Kosice, Slovakia

^b Centre for Animal Nutrition and Welfare, University of Veterinary Medicine Vienna, Veterinärplatz 1, A-1210 Wien, Austria

The metabolic pathway of thymol, a major constituent of *Thymus vulgaris* L., was studied for the first time in the rabbit organism after its sustained oral administration.

Forty-eight rabbits were allocated to control and experimental groups with thymol addition (250 mg/kg feed) for 21 days and withdrawal for 7 days. Significant correlation of the levels of thymol determined by GC/MS after solid phase microextraction ($r_s = -1.000$, $p < 0.01$) between intestinal wall (IW) and plasma points to the intensive absorption of thymol from the intestine. The significant correlation between plasma and liver ($r_s = 0.786$, $p < 0.05$) indicates intensive biotransformation and excretion processes in liver, and between liver and kidney ($r_s = 0.738$, $p < 0.05$) the intensive metabolism of thymol in the kidney. Even though thymol was determined only in trace amounts during the period without thymol addition, its amount in IW was significantly higher than in plasma. Thymol as lipophilic substance was found only in trace amounts in fat and muscle. In faeces it was detected in both periods above trace amounts showing significantly higher levels than in colon as a consequence of its conversion into hydrophilic metabolites and efficient elimination from organism. Our results showed some metabolic processes also after thymol withdrawal as consequence of caecotrophy, a characteristic feature of rabbit digestion processes responsible for better utilization of nutrients from feed.

Insufficient information about bioactivity of plant compounds in the animal organism represents a challenge for scientists to understand their metabolic processes at molecular level.

Funding: VEGA 2/0009/20, 2/0005/24; CNR-SAS-2022-02, Open-Mob-2022-01, COST CA22109, Action Austria-Slovakia ICM-2019-13685

Keywords: thymol, rabbit, bioavailability

* Presenting author. E-mail address: bacovak@saske.sk (Kristina Bacova)

S2.P268 *In-vitro* investigation of the gastroprotective effects of a natural extract blend: a promising avenue for canine gastritis prevention

Nicolò Lonigro^{a,b}, Talal Hassan^c, Mohammad Zaker Rehan^b, Mauro Bigliati^b, Maria Teresa Capucchio^{c,*}

^a Department of Drug Science and Technology, University of Turin, 10125 Torino, Italy

^b Candioli Pharma, 10092 Beinasco (Torino), Italy

^c Department of Veterinary Sciences, University of Turin, 10095 Grugliasco (Torino), Italy

Gastric hyperacidity is a common ailment in dogs that requires gastroprotection. Natural compounds, such as (psyllium, *aloe vera*) have anti-inflammatory properties (Szweda et al., 2014). The purpose of this study was to evaluate the gastroprotective efficacy of a natural extract (NE) blend containing inactivated yeast, psyllium seed cuticle (*Plantago ovata* L.), lupin protein flour, fenugreek, licorice, and *Aloe vera* (L.) Burm.f. extracts, in gastric adenocarcinoma (AGS) cells under hyperacidic conditions. Three NE concentrations were investigated based on a medium-sized dog's (15-20kg) stomach volume to deliver a maximum 2 g dose (0.3/0.15/0.075 mg/ml). AGS cells (10⁴ cells/well) were treated with NE in 96 well plates and examined with the MTT assay (3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyl tetrazolium bromide) using a Tecan M NANO+ plate reader (Ghazanfari et al., 2013). Cell viability was quantified against a positive control (sodium dodecyl sulfate, 1 mg/ml), followed by an acid pH gastroprotection test to determine NE's effectiveness under established positive control, hyperacidic conditions (pH=2). One-way ANOVA with Dunnett's post-hoc test was used for statistical analysis. NE showed 100% cell viability (0.554 ± 0.009 mg/ml, p<0.05), while the positive control had 40% viability (0.213 ± 0.003 mg/ml) compared to the negative control (0.534 ± 0.013). NE showed considerable gastroprotection (0.837 ± 0.009, p<0.05) at 0.3 mg/ml compared to negative control (0.852 ± 0.057) and positive control (0.686 ± 0.027). In conclusion, NE showed promising gastroprotective properties in AGS cells, although in vivo experiments are required to confirm NE's efficacy for clinical use in the prevention of gastritis in dogs.

Keywords: bioactive-compounds, dogs, gastroprotection

References

- Ghazanfari, T., Yaraee, R., Shams, J., Rahmati, B., Radjabian, T., & Hakimzadeh, H. 2013. Cytotoxic effect of four herbal medicines on gastric cancer (AGS) cell line. Food and Agri. Immu., 24, 1–7.
- Szweda, M., Szarek, J., Dublan, K., Męcik-Kronenberg, T., Kielbowicz, Z., & Bigoszewski, M. 2014. Effect of mucoprotective plant-derived therapies on damage to colonic mucosa caused by carprofen and robenacoxib administered to healthy dogs for 21 days. Vet. Quar., 34, 185–193.

* Presenting author. E-mail address: mariateresa.capucchio@unito.it (Maria Teresa Capucchio)

S2.P269 Acaricidal activity of essential oil of *Satureja montana* L. against *Dermanyssus gallinae* in vitro conditions

Radomir Ratajac^{a,*}, Aleksandra Pavličević^b, Jelena Petrović^a, Igor Stojanov^a, Dejan Orčić^c, Filip Štrbac^d, Nataša Simin^c

^a Scientific Veterinary Institute "Novi Sad" Novi Sad, Serbia

^b Aves mit d.o.o., Bajmok – Subotica, Serbia

^c Department of Chemistry, Biochemistry and Environmental Protection, Faculty of Sciences, University of Novi Sad, Novi Sad, Serbia

^d Institute for Multidisciplinary Research, University of Belgrade, Belgrade, Serbia

Dermanyssus gallinae commonly known as the poultry red mite causes great damage in the poultry industry. Plant-derived acaricides offer a strong alternative to chemical acaricides in the control of ectoparasites such as *D. gallinae* (Radsetoulalova et al., 2020; Ratajac et al. 2024). The aim of this study was to investigate the acaricidal activity of essential oil (EO) *Satureja montana* L. from the Balkan Peninsula and to evaluate their potential and limitations to be included in formulations for rational control of *D. gallinae* in the poultry industry. The chemical profile of EO was analyzed by gas chromatography coupled with mass spectrometry. The acaricidal efficacy of EO (0.1; 0.3; 0.5; 1; 3; 5 and 6% concentration) was tested on adult mites using the Petri-dish method, through direct exposure for 1 minute (contact toxicity) and subsequent exposure for 1 hour (residual toxicity). The EO efficiency ranged, in direct exposure-contact, after 48 hours of observation, from 3 to 100% toxicity, depending on the tested concentration, while the residual effect were negligible. George et al. (2010) also demonstrated efficacy after 24h exposure as 77.80% for *S. montana* L. Thus, the results of the present and other studies suggest a strong effect of EO against *D. gallinae* through direct contact and absence of prolonged effect. Plant based formulations are most likely less susceptible to resistance development and more acceptable in terms of the residues in animal products and the environment in comparison with chemicals (Abbas et al., 2018).

Keywords: Acaricides, *Dermanyssus gallinae*, *Satureja montana* L., Contact effect, Residual effect

References

- Abbas A, Abbas RZ, Masood S, et al., 2018. Acaricidal and insecticidal effects of essential oils against ectoparasites of veterinary importance. Bol Latinoam Caribe Plantas Med Aromat 17:441-52
- George DR, Sparagano OAE, Port G, et al., 2010. Environmental interactions with the toxicity of plant essential oils to the poultry red mite *Dermanyssus gallinae*. Med Vet Entomol 24:1-8.
- Radsetoulalova I, Hubert J, Hampel D, et al., 2020. Active components of essential oils as acaricides against *Dermanyssus gallinae*. Br Poult Sci 61:169-172
- Ratajac R, Pavličević A, Petrović J, Stojanov I, Orčić D, Štrbac F and Simin N, 2024. In vitro evaluation of acaricidal efficacy of selected essential oils against *Dermanyssus gallinae*. Pak Vet J, 44(1): 93-98

* Presenting author. E-mail address: ratajac@niv.ns.ac.rs (Radomir Ratajac)

S2.P270 A novel phytogetic supplement that prevents coccidiosis in chickens

Ming-Chu Cheng^a, Yuan-Bin Cheng^b, Meng-Ting Chang^c, Chi-Ting Chang^c, Yu-Hsin Chen^d, Guan-Hua Lai^a, Hsing-Yu Liao^a, Yi-Yang Lien^{a,*}, Lie-Fen Shyur^{c,*}

^a Department of Veterinary Medicine, National Pingtung University of Science and Technology, Pingtung 912, Taiwan

^b Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804, Taiwan

^c Agricultural Biotechnology Research Center, Academia Sinica, Taipei 115, Taiwan

^d Taichung District Research and Extension Station, Council of Agriculture, Executive Yuan, Changhua 515, Taiwan

Coccidiosis, caused by a protozoan parasite of the genus *Eimeria*, is one of the most severe contagious parasite diseases affecting the poultry industry worldwide. Using phytochemicals to prevent chicken coccidiosis is a novel strategy aimed at combating the increasing issue of drug-resistant strains of *Eimeria* spp. This study demonstrated the anticoccidial activities of a *Leguminosae* medicinal herb TP showing significant activity against *Eimeria* spp. The chemically defined TP extract exhibited significant suppressive activity against *E. maxima* oocyst sporulation and *E. tenella* sporozoite invasion and reproduction in Madin-Darby bovine kidney (MDBK) cells. Furthermore, administration of basal chicken diets containing TP to *Eimeria*-infected chickens significantly reduced the output of oocysts and severity of intestinal lesions. Dietary supplementation with TP significantly improved relative weight gain in *E. tenella*- and *E. acervulina*-infected chickens. The anticoccidial activities of TP on *E. acervulina*, *E. tenella* and *E. maxima* were further supported by anticoccidial index scores, which showed greater efficacy than those of amprolium, a commercial coccidiostat used in poultry. Furthermore, TP supplementation positively impacted the primary metabolism of chickens challenged with *E. tenella* or *E. acervulina*. In conclusion, the TP-derived chemical ingredients are demonstrated to be a novel phytochemical supplement that can be used to control *Eimeria*-induced coccidiosis in chickens.

Keywords: *Eimeria* spp., Coccidiosis, Anticoccidial index, Phytochemical supplement, Primary metabolism

* Presenting author. E-mail address: jaclyn@gate.sinica.edu.tw (Lie-Fen Shyur)

Poster Session 3, Tuesday, July 16, 2024

Cutting edge technologies in natural product drug discovery, formulation and development

S3.P1 *In silico* investigation of *Pterocarpus marsupium* against type-2 diabetes mellitus through network pharmacology approach

Hitender Sharma, Munish Garg*

Natural Products Research Laboratory, Department of pharmaceutical sciences, Maharshi Dayanand University, Rohtak-124001, Haryana, India

Pterocarpus marsupium heartwood (PMH) is a widely used plant in the management of diabetes in India. This presentation will highlight the potential of PMH-derived phytochemicals to control T2DM through network pharmacology approach. In this study, the bioactive compounds were retrieved from authentic scientific domains related to heartwood. Thereafter, the scientific tools were used to identify phytochemicals that meet Lipinski's Rule of Five parameters and oral absorption standards. Target identification for these phytochemicals was conducted using databases such as Binding DB, SwissTargetPrediction, STITCH, and TargetNet. Targets associated with T2DM were obtained from databases such as GeneCard, TherapeuticTargetDisease, NCBI Gene, and DrugBank. The STRING database was used for the PPI study. KEGG and GO enrichment studies were performed using the DAVID database to identify the pathways, biological pathways, cellular components, and molecular functions. The common targets between the PMH-phytochemicals and T2DM were used for network formation, including phytochemical targets, protein-protein interactions, and target-pathway networks. Finally, molecular docking between anti-T2DM targets and PMH phytochemicals was performed. The results of this study revealed that PMH may regulate blood glucose levels by directly interacting with target proteins such as SLC5A2, DPP4, DRD2, PPARG, and AKR1B1, which was verified using the molecular docking studies. Moreover, several networks like PT, TT, and TP were constructed and introduced modifications to each network. Subsequently, these networks were integrated to form a phytochemical-target-pathway network, which we hypothesize to offer enhanced precision in information retrieval. Several PMH targets showed protein-protein interactions which are involved in the signaling pathways such as insulin resistance and the T2DM signaling pathway. These findings generate further interest in the drug development process against T2DM.

Keywords: Network pharmacology, *Pterocarpus marsupium*, T2DM, Insulin resistance, molecular docking, biotargets, phytochemicals

References

- Abdul-Ghani, M.A., 2013. Type 2 diabetes and the evolving paradigm in glucose regulation. *Am. J. Manag. Care* 19, S43-50.
- Chellappan, D.K., Yap, W.S., Bt Ahmad Suhaimi, N.A., Gupta, G., Dua, K., 2018. Current therapies and targets for type 2 diabetes mellitus. *Panminerva Med.* 60, 117-131.
- Chen, Y., Ma, H., Zhu, D., Zhao, G., Wang, L., Fu, X., Chen, W., 2017. Discovery of novel insulin sensitizers: promising approaches and targets. *PPAR Res.* 2017, 8360919.
- Zhang, W., Huai, Y., Miao, Z., Qian, A., Wang, Y., 2019. Systems pharmacology for investigation of the mechanisms of action of traditional chinese medicine in drug discovery. *Front. Pharmacol.* 10, 743.

*Presenting author (email address): munishgarg@mdurohtak.ac.in (Munish Garg)

S3.P2 Integration of High-Throughput Screening, AI-ML Strategies, and ECD/DP4+ for the Discovery and Elucidation of Novel Natural Products

Yihao Zhuang^a, Arya Menon^b, Bahar Behsaz^c, Pamela J. Schultz^a, Hosein Mohimani^c, Amanda L. Garner^a, Ashootosh Tripathi^{a,*}

^a Department of Medicinal Chemistry, College of Pharmacy, University of Michigan, 1600 Huron Parkway, NCRC B520, Ann Arbor, Michigan 48109, USA; Natural Product Discovery Core, Mary Sue Coleman Hall, University of Michigan, 210 Washtenaw Avenue, Ann Arbor, Michigan 48109, USA

^b Department of Medicinal Chemistry, College of Pharmacy, University of Michigan, 1600 Huron Parkway, NCRC B520, Ann Arbor, Michigan 48109, USA


^c Computational Biology Department, School of Computer Science, Carnegie Mellon University, PA 15213, USA

Identifying and characterizing novel natural products holds immense potential for discovering bioactive compounds with diverse therapeutic applications. This presentation focuses on the integration of high-throughput screening (HTS), artificial intelligence-machine learning (AI-ML), and electronic circular dichroism (ECD)/DP4+ for the efficient discovery and structural elucidation of novel natural products. HTS enables the rapid evaluation of sizeable natural product libraries against biological targets, facilitating the identification of potentially bioactive compounds. Furthermore, combining HTS with advanced separation techniques allows for the isolation and purification of active compounds.

AI-ML strategies enhance the discovery process by predicting de novo structures and bioactivities and prioritizing compounds for further investigation. These approaches aid in exploring chemical space and identifying novel structural motifs. AI-ML algorithms trained on large datasets of known natural product structures and activities enable efficient compound screening.

Once potential hits are identified, ECD coupled with DP4+ becomes crucial for structural elucidation. ECD spectroscopy provides information on absolute configuration, aiding in stereochemistry determination. DP4+ is a probabilistic method for relative configuration determination, enabling accurate structural elucidation, even in complex mixtures.

In this pipeline, we incorporated ECD and GIAO NMR calculations coupled with a DP4+ probability measure, enabling the structure revision of phenazinolin D (4), izumiphenazine A (5), and baraphenazine G (7) and the structure characterization of two new diphenazines, baraphenazine H (3) and izumiphenazine E (6). The advancements discussed have the potential to revolutionize the field, facilitating the identification of compounds with significant therapeutic potential.



Keywords: ECD, NMR/DP4+, cytotoxicity, eIF4e, high-throughput screening, natural product

References

An ECD and NMR/DP4+ Computational Pipeline for Structure Revision and Elucidation of Diphenazine-Based Natural Products. Yihao Zhuang, Fei Yang, Arya Menon, James M. Song, Rosa V. Espinoza, Pamela J. Schultz, Amanda L. Garner, and Ashootosh Tripathi. *J. Nat. Prod.* 2023, 86, 7, 1801–1814.

*Presenting author. E-mail address: ashtri@umich.edu (Ashootosh Tripathi)

S3.P3 On-tissue derivatization and MALDI–MS Imaging reveal hemlock alkaloid distribution in *Conium maculatum*

Diana A. Barrera A.^{a,b,*}, Sabine Schuster^a, Timo H. J. Niedermeyer^b

^a Institute of Pharmacy, Martin-Luther-University Halle-Wittenberg, Hoher Weg 8, 06120 Halle (Saale), Germany

^b Institute of Pharmacy, Freie Universität Berlin, Königin-Luise-Straße 2+4, 14195 Berlin, Germany

The plant *Conium maculatum* is well-known for containing piperidine alkaloids, mainly coniine (**1**) and γ -coniceine (**2**). Its ingestion has strong effects on the central nervous system of mammals, causing ataxia and convulsions. (López et al., 1999; Talaty et al., 2005). However, the distribution of these alkaloids in plant tissues has not been studied in detail, yet. To close this knowledge gap, we analyzed the distribution of the hemlock alkaloids using MALDI mass spectrometry imaging (MSI), a technique that allows the determination of the localization of specific compounds in samples. However, it has limited applications for low molecular weight compounds, as the matrices can have background interference as well as overlapping signals with the analytes (Chen et al., 2012). Also, the hemlock alkaloids are rather volatile, which makes them difficult to analyze using MALDI-MSI. To overcome these limitations by modifying the physicochemical characteristics of these compounds, three derivatizing agents were studied. Derivatization with coniferyl aldehyde (CA), showed the best results, and the alkaloids were successfully visualized in the plant tissues, detecting **1** mainly in the midrib and margin of the leaf and in the endocarp of the fruit, while **2** was mainly detected in the dorsal ribs of the fruit (Fig. 1).




Fig. 1. MS Images of hemlock alkaloids in leaf and fruit tissues of *C. maculatum*. Midrib (mi), margin (m), dorsal secondary rib (dsr) and endocarp (en)

Keywords: *Conium maculatum*, spatial distribution, piperidine alkaloids, derivatization

References

- Chen, S., Chen, L., Wang, J., Hou, J., He, Q., Liu, J., Wang, J., Xiong, S., Yang, G., Nie, Z., 2012. 2,3,4,5-Tetrakis (3',4'- dihydroxylphenyl)thiophene: a new matrix for the selective analysis of low molecular weight amines and direct determination of creatinine in urine by MALDI-TOF MS. *Anal. Chem.* 84, 10291–10297.
- López, T.A., Cid, M.S., Bianchini, M.L., 1999. Biochemistry of hemlock (*Conium maculatum* L.) alkaloids and their acute and chronic toxicity in livestock. A review. *Toxicon* 37, 841–865.
- Talaty, N., Takáts, Z., Cooks, R.G., 2005. Rapid in situ detection of alkaloids in plant tissue under ambient conditions using desorption electrospray ionization. *Analyst* 130, 1624–1633.

* Presenting author. E-mail address: diana.barrera-adame@fu-berlin.de (Diana Astrid Barrera Adame)

S3.P4 New world and old world *Salvia* species (sages): searching of diterpenoids with cytotoxic activity

Diana Pérez-Juanchi^{a,*}, Baldomero Esquivel-Rodríguez^a, Celia Bustos-Brito^a, Leovigildo Quijano^a, Teresa Ramírez-Apan^a, Brenda Y. Bedolla-García^b, Sergio Zamudio^c

^a Departamento de Productos Naturales, Instituto de Química, Universidad Nacional Autónoma de México, Circuito Exterior, Ciudad Universitaria, Ciudad de México, 04510, México

^b Instituto de Ecología, A.C., Centro Regional del Bajío, Apartado Postal 386, 61600 Pátzcuaro, Michoacán, México

^c Apartado Postal 392, 61600 Pátzcuaro, Michoacán, México

This presentation will cover the isolation, structure elucidation and cytotoxic activity of diterpenoids of three *Salvia* species, two from the New World (NW) *S. involucrata* (SI) and *S. fulgens* (SF) (subgenus *Calosphace*), and one from the Old World (OW) *S. rosmarinus* (SR, subgenus *Rosmarinus*). *Salvia* is the most abundant genus of the Lamiaceae family, and it is divided into 11 subgenera (Kriebel et al., 2019). The main secondary metabolites isolated from these plants are diterpenoids, mainly *neo*-clerodanes from the NW species, and abietanes from the OW species (Wu et al., 2012).

We made collections of the three species, SI collected at Xilitla, San Luis Potosí (May 2017), SF at Los Azufres, Michoacán (November 2018), and SR at Milpa Alta, Mexico City (August 2022). The dried and powdered leaves of the collected species were extracted by percolation with CH₂Cl₂. The isolation was carried out by column chromatography, preparative TLC and semipreparative HPLC to produce eight diterpenoids (1-8, Fig. 1). The chemical structures were elucidated by extensive analysis of NMR spectroscopy and HR-DART-MS. The absolute configuration was established by experimental and theoretical ECD data or single-crystal X-ray diffraction.




Fig. 1. Isolated compounds from SI (1-6), SF (7) and SR (8).

The isolated diterpenoids were screened *in vitro* against six human cancer cell lines: MCF-7, K562, U251, SKLU-1, HCT-15 and PC-3. The highest growth inhibition was observed against K562, U251 and SKLU-1 (with values of 33.0-81.4 %), and the most active diterpenoid was 8 with 81.4 % against K562 (IC₅₀ = 22.04±1.0 μM).

Keywords: *Salvia*, diterpenoids, abietane, *neo*-clerodane, cytotoxicity

References

- Kriebel, R., Drew, B.T., Drummond, C.P., González-Gallegos, J.G., Celep, F., Mahdjoub, M.M., Rose, J.P., Xiang, C.L., Hu, G.X., Walker, J.B., Lemmon, E.M., Lemmon, A.R., Sytsma, J.S. 2019. Tracking temporal shifts in area, biomes, and pollinators in the radiation of *Salvia* (sages) across continents: leveraging anchored hybrid enrichment and targeted sequence data. *Am. J. Bot.* 106(4), 573-597.
- Wu, Y.B., Ni, Z.Y., Shi, Q.W., Dong, M., Kiyota, H., Gu, Y.C., Cong, B. 2012. Constituents from *Salvia* species and their biological activities. *Chem. Rev.* 112(11), 5967-6026.

* Presenting author. E-mail address: djuanchi_1404@live.com.mx (Diana Pérez Juanchi)

S3.P5 Use of *Galleria* moth larvae model to assess the pathogenesis of AIEC and the therapeutic potential of Geraniol as an antibacterial agent

Ryath Al-Hemedawi*

Teaching Laboratories Department - Medical City Hospitals Campus/Baghdad, Iraq

Crohn's disease (CD) is an inflammatory bowel disease (IBD) caused by abnormal immune responses to microbial infections in genetically susceptible individuals. Antibiotic-resistant adherent-invasive *Escherichia coli* is related to Crohn's disease (AIEC). Antibiotic resistance is one of the most important problems in modern healthcare, especially when it comes to illnesses of the gastrointestinal tract. There is an urgent need for new medications because the effects of these bacteria cannot be treated with current antibiotics. To address the issues of bacterial resistance and the decline of beneficial gut bacteria, the study concentrated on substances known as terpenoids (geraniol), which can reduce the action of bacteria rather than exterminate them. Gene deletions were put together utilizing the phage red gamma recombination technique in the adherent invasive *E. coli* background (Datsenko & Wanner 2000).

The moth larvae were bought from Live Food UK. Each larva weighed somewhere between 250 and 300 mg. To reach the doses of 10^5 , 10^6 , and 10^7 CFU/larvae, bacteria obtained from the mid-log phase were diluted with 1xPBS to 10^5 , 10^6 , 10^7 , and 10^8 CFU/ml, which is nearly equivalent to OD= inapplicable, 0.01, 0.02, and 0.13 respectively based on the free online Lab tools (<https://www.labtools.us/bacterial-cell-number-od600/>). Following injection, the larvae were kept at 37°C and kept in the dark. Over the course of five days, the coloration and mortality of the larvae were monitored twice daily. If the larvae did not react to touch, it was considered that the larvae had died. A minimum of three different iterations of each experiment were conducted. The Kaplan-Meier estimator was used to examine survival data.



Fig. 1 AB. (A) In the larvae geraniol safety test, 10 μ L of the mixture (HM605-107 + geraniol) was injected at the prescribed doses to the right of each larva's final proleg. Each larva has been observed for five days. Death of larvae is predicted by melanisation and loss of mobility. There were three replications of each experiment (n=3), and the graphs were constructed using pooled data. Using a significance level of P 0.0001, the Gehan-Breslow-Wilcoxon test was used to compare untreated and treated groups infected with AIEC (HM605).

(B) Image of *Galleria mellonella* larvae. *Galleria mellonella* larvae were injected with AIEC 10^7 CFU — HM605(WT) and indicative doses of geraniol or geraniol alone in the last proleg.

As a result, it was shown that the wild type of AIEC was more effective at killing larvae at the recommended dose than the *dsbA*-mutated strains. DsbA is required for the HtrA/DegP protein's protease activity, which is essential for AIEC survival in phagosomes. Geraniol can target the *dsbA* gene and prevent bacterial growth by competitive inhibition. This implies that the substance geraniol was discovered to possess antibacterial activity against the strains employed in this experiment.

Keywords: Adherent invasive *E. coli* (AIEC), Crohn's disease, geraniol, antibiotic resistance, Larvae, *Galleria mellonella*

References

1. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A*. 2000 Jun 6;97(12):6640-5. doi: 10.1073/pnas.120163297. PMID: 10829079; PMCID: PMC18686.
2. <https://www.labtools.us/bacterial-cell-number-od600/> E. coli Cell Culture Concentration from OD 600 CalculatorFree online Labtools.

* Presenting author. E-mail address: dr_ryathalhemedawi@yahoo.com (Ryath Al-Hemedawi)

S3.P6 Inhibition of transforming growth factor β signaling by prodigiosin through disruption of receptor recycling and subcellular translocation in epithelial cells

Chun-Lin Chen^{a,*}, Jing-Ru Weng^b, Jui-Hsin Su^c, Ping-Jyun Sung^c

^a Department of Biological Science, National Sun Yat-sen University, Kaohsiung 804, Taiwan, ROC

^b Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804, Taiwan, ROC

^c National Museum of Marine Biology and Aquarium, Pingtung 944, Taiwan, ROC

Prodigiosin (PG), a naturally occurring polypyrrole red pigment produced by various microorganisms, including certain *Serratia* and *Streptomyces* strains, has shown promising anticancer activity (Perez-Tomas et al., 2003). However, the molecular mechanisms underlying its action on malignant cells remain unclear. Transforming growth factor- β (TGF- β) is a multifunctional cytokine that governs various cellular processes in development and tissue homeostasis, with dysregulation of TGF- β signaling associated with numerous human cancers. Recent evidence highlights the importance of internalized TGF- β receptors and their intracellular trafficking in initiating signaling cascades (Heldin and Moustakas, 2016).

In this study, we identified PG as a potent inhibitor of the TGF- β pathway. PG hinders TGF- β signaling by targeting multiple sites in this pathway. It facilitates the sequestration of TGF- β receptors in the cytoplasm by impeding the recycling of type II TGF- β receptors to the cell surface. Furthermore, PG induces a reduction in receptor abundance on the cell surface by disrupting receptor glycosylation. In lung cancer A549 and HepG2 cells, nanomolar concentrations of PG significantly reduce TGF- β -triggered phosphorylation of Smad2 protein. This reduction is also reflected in the suppression of downstream target gene expression, including genes encoding fibronectin, plasminogen activator inhibitor-1 (PAI-1), and N-cadherin.

Keywords: Prodigiosin, cancer, TGF- β , intracellular trafficking

References

Perez-Tomas R, Montaner B, Llagostera E and Soto-Cerrato V (2003) The prodigiosins, proapoptotic drugs with anticancer properties. *Biochem Pharmacol* 66(8): 1447-1452.

Heldin CH and Moustakas A (2016) *Signaling Receptors for TGF-beta Family Members*. Cold Spring Harb Perspect Biol 8(8).

* Presenting author. E-mail address: chunlinchen@mail.nsysu.edu.tw (Chun-Lin Chen)

S3.P7 Assessing intra- and inter-variability of Macedonian wild-types and commercial *Cannabis* strains using short tandem repeat markers

Veronika Stoilkovska Gjorgievska^{a,*}, Nadica Matevska-Geskovska^b, Elizabeta Krstevska Bozinovikj^b, Ivana Cvetkovikj Karanfilova^a, Ana Trajkovska^a, Marija Karapandzova^a, Svetlana Kulevanova^a, Aleksandar Dimovski^b, Gjoshе Stefkov^a

^a Institute of Pharmacognosy, Faculty of Pharmacy, Ss. Cyril and Methodius University, Majka Tereza 47, 1000 Skopje, Republic of North Macedonia

^b Center for biomolecular pharmaceutical analyses, Faculty of Pharmacy, Ss. Cyril and Methodius University, Majka Tereza 47, 1000 Skopje, Republic of North Macedonia

Developing consistent Cannabis varieties capable of reliably producing a wide range of potentially beneficial cannabinoids is crucial for the research and medical communities (Nutt et al., 2013). Macedonian wild-growing cannabis possess cannabidiol potential and can be prospectively registered as a variety, but extensive UPOV (International Union for the Protection of New Varieties of Plants) descriptor-based testing is required. Genetic individuality and stability can be assessed utilizing STR (short tandem repeat) marker analysis as an alternative method. The aim of this study was to assess intra and inter-variability of outdoor cultivated Macedonian wild-types and commercial Cannabis strains. Automatic DNA extraction using MagCore kit was performed on 52 cannabis samples (flower/leaf tissue material) from five groups of wild-type plants and four groups of commercial strains that were outdoor cultivated from seeds. The wild-type seeds were collected from areas in Eastern Macedonia. Polymerase chain reaction and fragment analysis conditions were according to Houston et al. (2017) using 12 STR markers (ANUCS501, 9269, 4910, 5159, ANUCS305, B05- CANN1, 1528, 3735, 9043, D02-CANN, C11-CANN1, H06-CANN2). All of the samples were same homozygotes for marker 9269. STR markers ANUCS501, 4910, ANUCS305, 1528, 3735 and C11-CANN1 exhibited high inter- and intra- group variability, whereas only wild plants from Dolni and Gorni Podlog, were distinctive from all other groups with different alleles of ANUCS501, 1528 and D02-CANN. The STR analysis of cultivated seed plant material reveals significant variability within groups, underscoring the necessity for thorough genetic analysis correlated with phenotypic and phytochemical traits to identify the optimal genotype for registering wild plant groups as varieties in the European Union database.

Keywords: *Cannabis sativa*, hemp, DNA, molecular markers, genetic variability

References

- Nutt, D.J., King, L.A., Nicols, D.E., 2013. Effects of Schedule I drug laws on neuroscience research and treatment innovation. *Nat Rev Neurosci.* Vol. 14, 557-585.
- Houston, R., Birck, M., Hughes-Stamm, S., Gangitano, D., 2017. Developmental and internal validation of a novel 13 loci STR multiplex method for Cannabis sativa DNA profiling. *Leg Med.* Vol. 26, 33-40.

* Presenting author. E-mail address: vsgjorgievska@ff.ukim.edu.mk (Veronika Stoilkovska Gjorgievska)

S3.P8 Preclinical development of natural products into agents for pediatric brain cancers

Jehad Almaliti^a, Evgenia Glukhov^a, Momen Al-Hindy^a, Pavla Fajitova^b, Anthony J. O'Donoghue^b, Lena Gerwick^a, William H. Gerwick^{a,b}

^a Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California San Diego, 9500 Gilman Drive, La Jolla, California, 92093, USA

^b Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California San Diego, 9500 Gilman Drive, La Jolla, California, 92093, USA

Pediatric CNS tumors are the most common malignancy in childhood and adolescence, and occur with highest incidence in infants and children 5 years of age or younger. Therefore, anticancer therapeutics for the treatment of pediatric brain tumors are urgently needed. Selective human constitutive proteasome inhibitors have shown early stage promise for treating CNS tumors. We present here a project that was initiated with cytotoxic natural products, such as the carmaphycins and salinosporamide, as leads for the synthetic medicinal chemistry development of novel brain blood barrier penetrant proteasome inhibitors. These synthetic small molecules possess improved profiles of selectivity and metabolic stability in several different cell lines and microsome experiments. Extensive SAR studies, including incorporation of fluorinated amino acid residues, have achieved the production of highly potent small molecules at 5-6 μ M fully inhibiting constitutive proteasome at 0.5-1 nM range with favorable pharmaceutical properties for the development of a CNS pharmaceutical agent.

Keywords: proteasome inhibitors, cytotoxicity, cell lines

References

Jehad Almaliti, Pavla Fajtová, Jaeson Calla, Gregory M LaMonte *etc.*, 2023. Development of Potent and Highly Selective Epoxyketone-Based Plasmodium Proteasome Inhibitors. *Chemistry – A European Journal*. 29 (20).

* Presenting author. E-mail address: eglukhov@ucsd.edu (Evgenia Glukhov)

S3.P9 Exploring the anti-mycobacterial potential of French Guiana's flora through a metabolomic study

Célia Breaud^{a,*}, Clémentine Saunier^b, Stéphane Canaan^b, Béatrice Baghdikian^a, Sok-Siya Bunnelopet^a and Elnur Garayev^a

^a Aix Marseille Univ, CNRS 7263, IRD 237, Avignon Université, IMBE, 27 Blvd Jean Moulin, Service of Pharmacognosy, Faculty of Pharmacy, 13385 Marseille, France

^b CNRS, Aix-Marseille Univ, LISM UMR 7255, IMM FR3479, 13402 Marseille, France

Among the unique flora of the Amazonian Forest, numerous species have captured researcher's attention for their therapeutic potential, demonstrating promising activities against infectious diseases. Notably, mycobacterial infections such as tuberculosis stand as a significant health concern in French Guiana. With a high incidence rate and the emergence of drug-resistant strains, there is an urgent need to explore therapeutic alternatives (Chaptal et al., 2022; Succo, 2020).

This study delves into French Guiana's biodiversity, to explore the chemical composition of 11 plant species, and uncover novel metabolites showcasing anti-mycobacterial properties. Drawing from the literature, preliminary results indicate interesting activities in the selected plant species (Pavan et al., 2009).

The use of a methodology combining recent computational metabolomics tools and a MS-based Molecular Network approach, facilitates structural dereplication and allows to work on large datasets, ensuring rapid and accurate characterization of metabolites of therapeutic interest. Using multiple analytical techniques, such as UHPLC-HRMS/MS and NMR, we conducted a comprehensive metabolomic analysis to profile the chemical composition of the plant extracts. Simultaneously, bioassays against multiple strains of mycobacteria were performed. The integration of bioassays results on a Molecular Network facilitated the identification of bioactive molecules, which were prioritized for further investigation and bioassay-guided purification.

This multidisciplinary approach provides insights into the diverse chemical landscape of the studied plant extracts, and proposes a systematic approach to uncover novel anti-bacterial agents. This presentation will highlight the methodologies employed, initial findings, and the potential contribution of these plants in discovering therapeutic alternatives against mycobacterial infections.

Keywords: French Guiana, medicinal plants, mycobacteria, metabolomics, molecular network

References

- Carrion, L., Ramos, D., Martins, D., Osorio, M., Cursino, L., de Oliveira, W., Nunez, C., Silva, P. (2013). Antimycobacterial activity of Brazilian Amazon plants extracts. *International Journal of Phytomedicine*, 5, 479.
- Chaptal, M., Andrejak, C., Bonifay, T., Beillard, E., Guillot, G., Guyomard-Rabenirina, S., Demar, M., Trombert- Paolantoni, S., Jacomo, V., Mosnier, E., Veziris, N., Djossou, F., Epelboin, L., French Guiana PNTM working group. (2022). Epidemiology of infection by pulmonary non-tuberculous mycobacteria in French Guiana 2008-2018. *PLoS Neglected Tropical Diseases*, 16(9)
- Pavan, F. R., Sato, D. N., Higuchi, C. T., Santos, A. C. B., Vilegas, W., Leite, C. Q. E. (2009). In vitro anti-Mycobacterium tuberculosis activity of some Brazilian « Cerrado » plants. *Revista Brasileira de Farmacognosia*, 19, 204-206.
- Succo, T. (2020). Tuberculosis control in French Guyana: a public health priority. monitoring data, 2005-2017,7.

* Presenting author. E-mail address: celia.breaud@imbe.fr (Célia Breaud)

S3.P10 Arnica of Provence: an alternative to *Arnica montana* L.? Investigations about polyphenolic composition and antioxidant potential

Clémentine Achard-Baccati*, Charifat Saïd Hassane, Elnur Garayev, Béatrice Baghdikian

Aix Marseille Univ, CNRS 7263, IRD 237, Avignon Université, IMBE, 27 Blvd Jean Moulin, Service of Pharmacognosy and Ethnopharmacology, Faculty of Pharmacy, 13385 Marseille, France

In the Provence region (France), an Asteraceae species named *Pentanema montanum* (L.) D.Gut.Larr., Santos-Vicente, Anderb., E.Rico & M.M.Mart.Ort. (syn. *Inula montana* L.) is traditionally used with the same indications as *Arnica montana* L., against bruises, wounds, rheumatisms and muscular pain, and is even locally called “Arnica de Provence” (Garayev et al., 2018; Gyawali et al., 2022). Both species contain mainly sesquiterpene lactones and phenolic compounds, which are usually implicated in the defence against oxidative stress, involved in acute or chronic diseases (Vuolo et al., 2019).

Hydro-ethanolic extracts of these two species underwent evaluation of antioxidant potential and phytochemical studies. The radical scavenging activity was assessed using both 96 well-plate DPPH and DPPH-On-Line-HPLC assays. Well-plate DPPH assay showed higher activity for *P. montanum* extracts, with an IC₅₀ twice lower than *A. montana* extracts, respectively 23.66 ± 0.75 µg/mL and 45.71 ± 3.05 µg/mL. DPPH-On-Line-HPLC assay allowed to identify 1,5-dicaffeoylquinic acid as the main compound responsible for the anti-oxidant activity in both extracts. This compound is more abundant in *P. montanum* extracts, supporting the well-plate DPPH assay results.

Metabolomic data were obtained through UHPLC-HRMS/MS analysis and processed using computational metabolomics and MS-based Molecular Networking. The two species only share 16% of their chemical composition, most metabolites being species-specific.

This presentation will emphasize the comparative analysis by molecular networking of *A. montana* and *P. montanum* chemical profiles, with a focus on exclusive metabolites of each species and those contributing to radical scavenging activity.

Keywords: metabolomics, molecular networking, antioxidant, DPPH, traditional medicine

References

- Breud, C., Lallemand, L., Mares, G., Mabrouki, F., Bertolotti, M., Simmler, C., Greff, S., Mauduit, M., Herbette, G., Garayev, E., Laverne, C., Cesari, M., Bun-Llopet, S.-S., Baghdikian, B., & Garayev, E. (2022). LC-MS Based Phytochemical Profiling towards the Identification of Antioxidant Markers in Some Endemic *Aloe* Species from Mascarene Islands. *Antioxidants (Basel, Switzerland)*, 12(1), 50.
- Garayev, E., Di Giorgio, C., Herbette, G., Mabrouki, F., Chiffolleau, P., Roux, D., Sallanon, H., Ollivier, E., Elias, R., & Baghdikian, B. (2018). Bioassay-guided isolation and UHPLC-DAD-ESI-MS/MS quantification of potential anti-inflammatory phenolic compounds from flowers of *Inula montana* L. *Journal of ethnopharmacology*, 226, 176-184.
- Gyawali, N., Rayamajhi, A., Karki, D., Pokhrel, T., & Adhikari, A. (2022). *Arnica montana* L.: Traditional Uses, Bioactive Chemical Constituents, and Pharmacological Activities. In H. P. Devkota & T. Aftab (Éds.), *Medicinal Plants of the Asteraceae Family: Traditional Uses, Phytochemistry and Pharmacological Activities* (p. 61-75). Springer Nature.
- Vuolo, M. M., Lima, V. S., & Maróstica Junior, M. R. (2019). Chapter 2 - Phenolic Compounds: Structure, Classification, and Antioxidant Power. In M. R. S. Campos (Éd.), *Bioactive Compounds* (p. 33-50). Woodhead Publishing.

* Presenting author. E-mail address: clementine.baccati@imbe.fr (Clémentine Achard-Baccati)

S3.P11 Biosynthesis of anti-inflammatory compounds from *Calendula officinalis*

Daria Golubova^{a,*}, Melissa Salmon^a, Connor Tansley^{a,b}, Honghao Su^a, Maria O'Connell^c, Nicola Patron^{a,b}

^a Engineering Biology, Earlham Institute, Norwich Research Park, Norwich, NR4 7UZ, UK

^b Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK

^c School of Pharmacy, University of East Anglia, Norwich, Norwich Research Park, Norwich, NR4 7TJ, UK

Plants have been used in traditional medicine for thousands of years. In some cases, therapeutic properties have been attributed to specific metabolites, enabling their use as drugs in modern medicine. However, the exact molecule(s) responsible for the bioactivity of many plant extracts remains unknown. Further, access to natural products is often complicated by low abundance, occurrence in complex mixtures, or by structural complexity, which can complicate chemical synthesis. In this project, we are investigating the previously reported anti-inflammatory bioactivity of *Calendula officinalis* (pot marigold) floral extracts. Studies have suggested bioactivity may be associated with the presence of a class of molecules known as triterpene fatty acid esters (TFAEs) (Colombo et al., 2015; Nicolaus et al., 2017). To investigate this, and to identify specific molecules that contribute to bioactivity, we combined comparative metabolic profiling of Asteraceae species with bioactivity assays in model human cell lines. Our data suggests that TFAEs with C:16 hydroxylated scaffolds may exert an anti-inflammatory effect via tumour necrosis factor (TNF) and the interleukin 6 (IL-6) pathways. Additionally, we have combined metabolomics, genomics, and transcriptomics with rapid, transient expression in *Nicotiana benthamiana* to identify and characterize the biosynthetic pathways of *C. officinalis* TFAEs (Fig. 1). Our work provides a route to the identification and production of novel plant bioactives.



Fig. 1. *Calendula officinalis* flowers are rich in triterpene fatty acid esters (TFAEs) which have been proposed to confer anti-inflammatory bioactivity. Heterologous expression of candidate genes in *Nicotiana benthamiana* enabled us to elucidate the biosynthetic pathways of these molecules and investigate their bioactivity.

Keywords: *Calendula officinalis*, anti-inflammatory bioactivity, bioassays, triterpenes, *Nicotiana benthamiana*

References

- Colombo, E., Sangiovanni, E., D'Ambrosio, M., Bosisio, E., Ciocarlan, A., Fumagalli, M., Guerriero, A., Harghel, P. and Dell'Agli, M., 2015. A bio-guided fractionation to assess the inhibitory activity of *Calendula officinalis* L. on the NF- κ B driven transcription in human gastric epithelial cells. *Evidence-based complementary and alternative medicine*, 2015.
- Nicolaus, C., Junghanns, S., Hartmann, A., Murillo, R., Ganzera, M. and Merfort, I., 2017. *In vitro* studies to evaluate the wound healing properties of *Calendula officinalis* extracts. *Journal of Ethnopharmacology*, 196, pp.94-103.

Presenting author. golubova@nbi.ac.uk (Daria Golubova)

S3.P12 Development and validation of an analytical method for the analysis of standardized red propolis extract and its chemopreventive effect against colon carcinogenesis

Sergio Ricardo Ambrosio^{a,*}, Iara Silva Squarisi^a, Victor Pena Ribeiro^a, Arthur Barcelos Ribeiro^a, Leticia Teixeira Marcos De Souza^a, Marcelo de Melo Junqueira^a, Katia Mara de Oliveira^a, Jairo Kenupp Bastos^b, Gaelle Hayot^c, Thomas Dickmeis^c, Denise Crispim Tavares^a

^a Research Group on Natural Products, Center for Research in Sciences and Technology, University of Franca, Franca, SP, Brazil

^b School of Pharmaceutical Sciences of Ribeirão Preto – University of São Paulo, Ribeirão Preto, Brazil

^c Institute of Biological and Chemical Systems - Biological Information Processing - Karlsruhe Institute of Technology, Karlsruhe, Germany

Nowadays, Brazilian red propolis (BRP) has gained prominence in the scientific literature due to its several pharmacological applications. Despite this fact, some toxicological effects have been observed from BRP, in which is credited to the presence of polyphenylated benzophenones, such as gutiferone E, xanthochimol and oblongifolin B (Ccana-Ccapatinta et al. 2020).

In view of this context, our research group has obtained a standardized red propolis extract without the presence of polyphenylated benzophenones using a sequence of *n*-hexanes and then methanol (SRPE). In addition, an analytical method using RP-HPLC-PDA was developed and validated according to the guidelines proposed by Brazilian National Health Surveillance Agency (ANVISA, 2017). For this purpose, the main compounds from SRPE were isolated, identified, and considered as chemical markers (Fig. 1). The developed method showed to be precise, sensitive, and reliable for analyzing SRPE.




Fig.1. Compounds used as chemical markers in the development and validation of the analytical method: 1: isoliquiritigenin; 2: vestitol; 3: neovestitol; 4: medicarpin; 5: 7-O-methylvestitol.

Considering the biological properties of SRPE, different doses of this extract (3, 6, and 12 mg/kg b.w.) were administered in combination with the carcinogen 1,2-dimethylhydrazine (DMH, 40 mg/kg b.w.) using an *in vivo* colon carcinogenesis model (da Silva et al., 2022). The animals treated with 6 mg/kg b.w. showed a significant reduction (41.57%) in pre-neoplastic lesions induced by DMH, thus revealing a preventive effect against colon carcinogenesis of SRPE.

Keywords: Brazilian red propolis, chemopreventive effect, colon carcinogenesis, analytical method

References

- ANVISA, 2017. Resolução da diretoria colegiada - RCD No. 166, Ministério da Saúde.
- Ccana-Ccapatinta, G.V., Mejía, J.A.A., Tanimoto, M.H., Groppo, M., Carvalho, J.C.A.S., Bastos, J.K., 2020. *Dalbergia ecastaphyllum* (L.) Taub. and *Symphonia globulifera* L.f.: the botanical sources of isoflavonoids and benzophenones in Brazilian red propolis. *Molecules*. 25, 2060-2067.
- da Silva, L.H.D., Squarisi, I.S., de Freitas, K.S., Barcelos, A., Ozelin, S.D., Aldana-mejía, de Oliveira, L.T.S., Rodrigues, T.E., Nicolella, H.D., Alves, B.S., Ccana-ccapatinta, G.V., Bastos, J.K., Tavares, D.C., 2022. Toxicological and chemoprevention studies of *Dalbergia ecastaphyllum* (L.) Taub. stem, the botanical source of Brazilian red propolis. *J. Pharm. Pharmacol.* 74, 1-10.

* Presenting author. E-mail address: sergio.ambrosio@unifran.edu.br (Sergio Ricardo Ambrosio)

S3.P13 Extract rich in polyprenylated benzophenones from Brazilian red propolis: obtaining, analytical characterization and *in vitro* evaluation against breast cancer lines

Rodrigo Cassio Sola Veneziani^{a,*}, Raquel Alves dos Santos^a, Loren Monielly Pires^a, Laisla Rodrigues Figueiredo^a, Renata Faleiros^a, Karolinne Beloti Silva^a, Mario Ferreira Conceição Santos^a and Jairo Kenupp Bastos^b

^a Universidade de Franca, Av. Dr. Armando Salles de Oliveira 201, 14404600 Franca, Brazil

^b Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Av. do Café s/n, 14040903 Ribeirão Preto, Brazil

The present work describes the obtaining of an extract rich in polyprenylated benzophenones from Brazilian red propolis (ERP), the development and validation of an HPLC-UV method to characterize it as well as its evaluation against breast cancer cell lines MCF-7 and MDA-MB-231 and a normal counterpart MCF-10A. A mixture of guttiferone E + xanthochymol (1+2), and isolated oblongifolin B (3) (Fig. 1) were used as chemical markers of ERP.



Fig. 1. Structure of the polyprenylated benzophenones guttiferone E 1, xanthochymol 2, and oblongifolin B 3

ERP was obtained by maceration of Brazilian red propolis using *n*-hexanes followed by concentration and suspension in cold methanol to remove waxy material. The RP-HPLC-DAD analytical method was established based on initial experiments using gradient scouting run evaluation (Snyder and Dolan, 1996) and validated according to the guidelines of the Brazilian National Health Surveillance Agency (Santos et al., 2021). The concentrations of 1+2 and 3 were 101,822 µg/mL and 257,918 µg/mL, respectively, corresponding to 16.68% and 42.25% of the total content of the extract and the validation parameters evaluated were satisfactorily met. The cytotoxic effects of ERP were assessed through XTT assay (Gonçalves et al., 2021) and the obtained IC₅₀ were 19.58 µg/mL (MCF-10A), 11.56 µg/mL (MCF-7) and 5.22 µg/mL (MDA-MB-231) thus resulting in selectivity index values of 3.75 (MCF-10A/MDA-MB-231) and 1.69 (MCF-10A/ MCF-7). In conclusion, ERP exhibit promising cytotoxic effects on the tested breast cell lines. However, further investigation to elucidate its potential therapeutic applications and safety profiles should be conducted.

Keywords: Brazilian red propolis, polyprenylated benzophenones, analytical method, breast cancer cell lines

References

- Snyder, L.R., Dolan J.W., 1996. Initial experiments in high-performance liquid chromatographic method development .1. Use of a starting gradient run. J. Chromatogr. A. 721, 3-14.
- Santos M.F.C., Oliveira L.C., Ribeiro V.P., Soares M.G., Moraes G.D.I., Sartori A.G.D., Rosalen P.L., Bastos J.K., de Alencar S.M., Veneziani R.C.S., Ambrosio S.R., 2021. Isolation of diterpenes from *Araucaria* sp Brazilian brown propolis and development of a validated high-performance liquid chromatography method for its analysis. J. Sep. Sci. 44, 3089– 3097.
- Gonçalves, N.S., de Mello, T.M.S.P., Mizuno, C.S., Haider, S., dos Santos, R.A., 2021. *Cis*-trimethoxystilbene, exhibits higher genotoxic and antiproliferative effects than its isomer *trans*-trimethoxystilbene in MCF-7 and MCF-10A cell lines. Genet. Mol. Biol. 44, e20200477.

* Presenting author. E-mail address: rodrigo.veneziani@unifran.edu.br (Rodrigo Cassio Sola Veneziani)

S3.P14 Computational NMR enhances metabolomic analysis and sustainability

Matthias Niemitz^{a,*}, Petrus Korhonen^a, Juuso Lehtivarjo^a, G. Joseph Ray^b, David C. Lankin^b, Charlotte A. Corbett^c, Shao-Nong Chen^b, Guido F. Pauli^{bz}

^a NMR Solutions Ltd., 70100 Kuopio, Finland

^b Pharmacognosy Institute and Department of Pharmaceutical Sciences, University of Illinois Chicago, College of Pharmacy, Chicago, IL 60612, USA

^c Drug Enforcement Administration, Special Testing and Research Laboratory, Dulles, VA 20166, USA

In the absence of crystals for diffraction measurements, rigorous stereospecific structural analysis of natural products (NPs) is the domain of NMR and chiroptical spectroscopies. The stereochemical significance and near ubiquitous occurrence of hydrogen atoms make ¹H NMR a powerful analytical tool, but the wealth of information comes at the price of complexity - even in seemingly simple small molecules.

The solution to this conundrum is applying the NMR theory. Its foundation was laid in the 1920s and proven experimentally in the 1950s. Since then, the lack of computational resources has been the main, if not only, reason why applying NMR theory to spectral analysis has been unfeasible. This has now changed dramatically: adequate computational power and software tools are available that allow automated computational spectral analysis - even of highly complex ¹H NMR spectra - in seconds to minutes. While 2D NMR is invaluable for spectral analysis, much of the valuable coupling information in NPs remains ignored in current practice, as is evident from the abundant use of “multiplet” annotations. However, the ability to extract the entire ¹H NMR information can transform both qualitative and quantitative analysis of NPs for several reasons:

1. Couplings provide essential stereochemical information and allow independent verification of chemical (shift) assignments from 2D correlation experiments.
2. As sets of chemical shifts and couplings must be self-consistent and coherent with the observed spectra, confidence in the correctness of the results increases dramatically.
3. Even overly complex spectral patterns and peak overlap can be fully explained using the additional constraints provided by the NMR theory.
4. Chemical shifts and especially couplings are instrument independent and represent highly reproducible physical constants under well-defined experimental conditions.
5. Collectively, this allows automated and highly specific dereplication of known NPs based on shared raw 1D ¹H spectra - regardless of NMR instrumentation.

Results from computational analyses of newly discovered and challenging NPs with available raw NMR data will be shared to explain the methodology and its capabilities. We will exemplify the growing experimental evidence for capability of the fastest and most affordable NMR measurement to extract new structural and quantitative insights and potentially transform metabolomic and biomedical NP analysis. Applying computational analysis particularly to 1D ¹H NMR spectra serves scientific sustainability for several reasons: (A) Exploitation of the full NMR information content enhances the re-use of essential prior knowledge. (B) Increased throughput and reduced per-sample costs of today's capable NMR instruments. (C) Widened capability of NMR as a non-destructive analytical technique with reduced solvent footprint, especially on benchtop instruments without deuterated solvents.

Keywords: Structural analysis, quantitative analysis, dereplication, NMR

* Presenting author. E-mail address: niemitz@nmrsolutions.io (Matthias Niemitz)

S3.P15 Exploring on the networking of chemical compositions and their cytotoxicity toward KBCC *Nepenthes* plants

Szu-Yin Yu^{a,b,*}, Hao-Chun Hu^{a,c,d}, Attila Hunyadi^e, Fang-Rong Chang^a

^a Graduate Institute of Natural Products, College of Pharmacy, Kaohsiung Medical University, Kaohsiung 80708, Taiwan

^b Institute of Pharmacognosy, University of Szeged, Szeged 6720, Hungary

^c Graduate Institute of Natural Products, Chang Gung University, Wenhua 1st Rd, Kwei-Shan District, Taoyuan 333, Taiwan

^d Institute of Pharmaceutical Chemistry and HUN-REN-Stereochemistry Research Group, University of Szeged, Szeged 6720, Hungary

^e Institute of Pharmacognosy, Interdisciplinary Excellence Centre, University of Szeged, Szeged 6720, Hungary

In collaboration with the Dr. Cecilia Koo Botanic Conservation Centre, we collected 60 *Nepenthes* species for further investigation. Preliminary activity screening showed that the methanol extract of *Nepenthes mirabilis* suppressed elastase release, indicating anti-inflammatory activity.

In addition, activity-guided isolation, molecular networking and mass spectrometry were used to isolate, purify and identify potentially active natural products. Various analytical experiments were performed using instruments such as HPLC-PDA, spectrophotometer, and LCMS/MS. The study used activity-guided isolation, chemical analysis, and supplemented by computer simulation of the Global Natural Product Molecular Network (GNPS) for ancillary analysis.

This study found that the naphthoquinone component, plumbagin, in *Nepenthes* was effective in inhibiting the growth of human liver cancer cells (HepG2), human triple negative breast cancer cells (MDA-MB-231) and human non-small cell lung cancer cells (A549) (Wójciak et al., 2023). The compound also exhibited anti-inflammatory (human neutrophil) properties. The dried methanol and *n*-hexane partition layers of 60 *Nepenthes* species were screened for cytotoxicity and anti-inflammatory activity, and the chemical composition and cytotoxic network were explored with the indicator compound, plumbagin. Besides, we carried out the carbon dots synthesis of plumbagin and found that it became a nano-probe with wide spectral performance.

Keywords: *Nepenthes mirabilis*, *Nepenthes*, carbon dots, cytotoxicity, anti-inflammatory, molecular networking

References

Wójciak, M., Feldo, M., Stolarczyk, P., Płachno, B. J., 2023. Carnivorous Plants from Nepenthaceae and Droseraceae as a Source of Secondary Metabolites. *Molecules*. 28(5), 2155.

* Presenting author. E-mail address: u105531007@gap.kmu.edu.tw (Szu-Yin Yu)

S3.P16 Chemical composition and bioactivity of *Uvaria micrantha* from Vietnam

Fang-Rong Chang^{a,*}, Hao-Chun Hu^{a,b,c}, Szu-Yin Yu^{a,d}, Yi-Hong Tai^{a,e}, Hsin-Yi, Tang^a

^a Graduate Institute of Natural Products, College of Pharmacy, Kaohsiung Medical University, Kaohsiung 80708, Taiwan

^b Graduate Institute of Natural Products, Chang Gung University, Wenhua 1st Rd, Kwei-Shan District, Taoyuan 333, Taiwan

^c Institute of Pharmaceutical Chemistry and HUN-REN-Stereochemistry Research Group, University of Szeged, Szeged 6720, Hungary

^d Institute of Pharmacognosy, University of Szeged, Szeged 6720, Hungary

^e Department of Pharmacy and Master Program, College of Pharmacy and Health Care, Tajen University, Pingtung County 907101, Taiwan

Uvaria (Family Annonaceae) was reported to have anti-inflammatory, anti-cancer, and anti-malaria activities. Among them, *Uvaria micrantha* is distributed in Southeast Asia and Australia. However, only one paper revealed the isolated compounds from the stems of *Uvaria micrantha* and their potential effects on anti-leukemia and cervical cancers (Boonsombat et al., 2021). Traditionally, this plant can be used to treat back pain and digestive problems. In this study, 18 folk Vietnamese herbal plants were collected from Ho Chi Minh City, Vietnam. After extraction and partition, the *n*-hexane, *n*-butanol, water, and methanol extracts of these plants were screened for their anti-Alzheimer's disease, anti-osteoporosis, anti-stroke, and anti-inflammatory activities, in collaboration with National Research Institute of Chinese Medicine, Taiwan. This preliminary screening found that VN018 showed good activities in the multiple assays, especially its methanol layer. After gene sequencing and plant specimen identification, VN018 was identified as *U. micrantha*. Hence, the methanol extract of branches and leaves from *U. micrantha* was chosen for further separation, purification, and exploring their potential bioactivities. In the current study, one new and twelve known compounds were isolated (Fig. 1). The activities of the extracts and pure compounds for anti-stroke activity will be shown.




Fig. 1. Structure of compound 1

Keywords: *Uvaria*, *Uvaria micrantha*, uvarisinyicone, Alzheimer's disease, osteoporosis, stroke, anti-inflammatory

References

Boonsombat J, Thongnest S, Kheawchaum S, Mahidol C, Ruchirawat S, Prawat H, 2021. Uvarmicranones A and B, two new benzoquinones and cytotoxic constituents from the stems of *Uvaria micrantha* (A. DC.) Hook. f. & Thomson. Nat Prod Res. 35(24), 5643–5652.

* Presenting author. E-mail address: aaronfrc@kmu.edu.tw (Fang-Rong Chang)

S3.P17 **Ontology based ethnobotany data to support natural product-based drug discovery – growth of data, large-scale insights and case studies**

Vijay Ingalalli^a, Jui-Hung Yuan^a, Lila Khederlarian^b, Lukas Hermann^a, Kamen Petrov^a, Yoann Gloaguen^a, Niklas Leuchtenmüller^c, Sona Chandra^a, Andreas Bender^a

^a PangeAI, Pangea Botanica GmbH, Hardenbergstraße 32, 10623 Berlin, Germany

^b Pangea Botanica Ltd, 15 Southampton Pl, London WC1A 2AJ, UK

^c Wilde Ventures GmbH, In der Rehwiese 3, 40629 Düsseldorf, Germany

For millennia and across cultures plants have been used to treat human diseases. Despite efforts to organize this knowledge, attempts have so far been either limited to specific types of traditional medicines or unable to practically support drug discovery tasks. In this work we describe our work of creating ontologies and controlled vocabularies with the aim to achieve interoperability of ethnobotany data with chemical, biological, and medical datasets. This dataset comprises >500m data points, >2m species-disease links, >500,000 natural products, and >34m target-disease links, covering areas from botany, historical use, plant metabolites, bioactivity, to target-disease links. We performed analyses on the data to understand and quantify chemotaxonomy principles as well as to develop real-world use cases of relevance for drug discovery. We investigated to what extent plants within the same genus (or family) contain the same metabolite, as well as to estimate the novelty of the chemistry that can be discovered in each plant (both in terms of absolute novelty, and novelty in a given organism). Furthermore, we established a real-world use case for the pain disease area, and quantified in which way we can re-identify chemistry associated with *in vivo* activity from data across different mode-of-action classes. Overall, compiling ethnobotany information on a large scale allowed us to quantify relationships between historical use, plants, chemistry, compounds, and bioactivity, and to use such data in real-world drug discovery projects. Conflict of interest statement: The authors are employees of Pangea Botanica GmbH, Pangea Botanica Ltd and Wilde Ventures GmbH.

Keywords: ethnobotany, chemotaxonomy, drug discovery, artificial intelligence, machine learning

* Presenting author. E-mail address: andreas@pangeabio.com (Vijay Ingalalli)

S3.P18 Optimization formulation of *Rhodiola rosea* L. nanoemulsion containing natural oil using response surface method (RSM): preparation, characterization, stability test and antioxidant activity

Benni Iskandar^{a,b}, Hsiu-Mei Lin^c, Ching-Kuo Lee^{a,d,*}

^a School of Pharmacy, College of Pharmacy, Taipei Medical University, Taipei 11031, Taiwan

^b Department of Pharmaceutical Technology, Riau College of Pharmaceutical Sciences (STIFAR), Pekanbaru, Riau, Indonesia, 28292

^c Department of Bioscience and Biotechnology, National Taiwan Ocean University, Keelung 202, Taiwan

^d Graduate Institute of Pharmacognosy, Taipei Medical University, Taipei 11031, Taiwan

The oil phase plays a vital role in nanoemulsion formulation. *Rhodiola rosea* L. root extract offers a wide range of pharmacological advantages, such as antioxidative, anti-collagenase, anti-aging, neuroprotective, anti-inflammatory, cardioprotective, and anti-cancer activities. Sunflower oil, chosen as the natural oil phase, offers various advantages such as its antioxidant properties and its safety for skin application. This study involved preparing 13 variations of *Rhodiola rosea* L. nanoemulsions with sunflower oil and a combination of surfactant and co-surfactant using the spontaneous emulsification method. The nanoemulsion was analyzed for several parameters, such as Particle Size, Polydispersity Index (PDI), and Zeta Potential. The analytical parameters for the nanoemulsion of *Rhodiola rosea* L. were calculated using Design Expert software. The nanoemulsion was optimized using the response surface method (RSM) to determine the optimal amounts of surfactant and cosurfactant. The optimized formulation consists of 21.5% transcutool, 30% labrasol, and 3% sunflower oil. These proportions yield optimal response values of 177.63 nm for particle size,

0.25 for PDI, and -47.69 for zeta potential. The optimized *Rhodiola rosea* L. nanoemulsion exhibits the characteristics of a nanoemulsion, viscosity, stability studies and demonstrates antioxidant activity. Therefore, this study provided the guidelines for a method and combination of formulation stable nanoemulsions for cosmetic and pharmaceutical fields in the future by a combination of surfactant-cosurfactant.

Keywords: nanoemulsion, *Rhodiola rosea* L., response surface method, natural oil

References

- Zhang, M., et al., *Functional nanoemulsions: Controllable low-energy nanoemulsification and advanced biomedical application*. Chinese Chemical Letters, 2023: p. 108710.
- Lee, T.H., et al., *Anti-MMP-2 Activity and Skin-Penetrating Capability of the Chemical Constituents from Rhodiola rosea*. Planta Med, 2016. 82(8): p. 698-704.
- Wang, C.-H., et al. *Protective Evaluation of Compounds Extracted from Root of Rhodiola rosea L. against Methylglyoxal-Induced Toxicity in a Neuronal Cell Line*. Molecules, 2020. 25, DOI: 10.3390/molecules25122801.
- Ma, R., et al., *Extraction, purification, structure, and antioxidant activity of polysaccharide from Rhodiola rosea*. Journal of Molecular Structure, 2023. 1283: p. 135310.
- Masi, F., et al., *Phytochemical profile and anti-inflammatory activity of a commercially available Rhodiola rosea root extract*. Fitoterapia, 2023. 166: p. 105439.
- Iskandar B, Mei H-C, Liu T-W, Lin H-M, Lee C-K: Evaluating the effects of surfactant types on the properties and stability of oil-in-water *Rhodiola rosea* nanoemulsion. *Colloids and Surfaces B: Biointerfaces* 2024, 234:113692.

* Presenting author. E-mail address: cklee@tmu.edu.tw (Ching-Kuo Lee)

S3.P19 Oxidoreductase FSP1-mediated redox-regenerating natural product E18C6 potently suppresses ferroptotic cell death

Minh Bui Hoang^{a,b,*}, Stefan Schwaiger^b, Claudia Kostner^a, Alexander Weiss^c, Hermann Stuppner^b, Andreas Koeberle^a, Solveigh Koeberle^a

^a Michael Popp Institute and Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, 6020, Innsbruck, Austria

^b Pharmacognosy - Institute of Pharmacy and Center for Molecular Biosciences Innsbruck (CMBI), University of Innsbruck, 6020, Innsbruck, Austria

^c Research Institute for Biomedical Aging Research, University of Innsbruck, 6020, Innsbruck, Austria

Ferroptosis, a form of regulated cell death dependent on iron and driven by lipid peroxidation, contributes to degenerative diseases and organ injury (Stockwell, 2022; Angeli et al., 2017; Koeberle et al., 2023). Agents with anti-ferroptotic properties hold great promise in preventing excessive cell death in these pathologies. Bioactivity-guided isolation from 19 traditional Vietnamese medicinal plant extracts identified the natural product E18C6 as a potent anti-ferroptotic compound ($EC_{50} = 10$ nM) standing out among the most efficient ferroptosis inhibitors described to date. Subsequent screening of 59 structural analogues from our in-house library identified E18C6 as the most potent derivative in this structural compound class. Mechanistically, the oxidoreductase FSP1 converts E18C6 to its reduced form, a potent radical scavenger that efficiently prevents lipid peroxidation and ferroptosis. The FSP1-dependent inhibition of lipid peroxidation was confirmed in the FENIX assay (Shah et al., 2019), which assesses radical scavenging efficiency within phospholipid bilayers *in vitro*. These results strongly suggest that E18C6 acts as a robust inhibitor of ferroptosis through FSP1-dependent reduction to its active radical-scavenging form, with continuous efficient regeneration by FSP1. *In vivo* efficacy of E18C6 was evaluated in a *Caenorhabditis elegans* model employing the GSH-depleting compound diethyl maleate. E18C6 demonstrated a marked reduction in the toxicity associated with diethyl maleate, providing compelling evidence for its efficacy in preventing ferroptosis in an *in vivo* context. In conclusion, our study identified E18C6 as a promising anti-ferroptotic lead that has the potential to prevent extensive pathological cell death in degenerative diseases and inhibit disease progression.

Keywords: ferroptosis, cell death, FSP1, FENIX assay, *C. elegans*, degenerative disease

References

- Angeli, J. P. F., Shah, R., Pratt, D. A., Conrad, M., 2017. Ferroptosis inhibition: mechanisms and opportunities. *Trends Pharmacol. Sci.*, 38(5), 489-498.
- Koeberle, S. C., Kipp, A. P., Stuppner, H., Koeberle, A., 2023. Ferroptosis-modulating small molecules for targeting drug-resistant cancer: Challenges and opportunities in manipulating redox signaling. *Med. Res. Rev.*, 43(3), 614-682.
- Shah, R., Farmer, L. A., Zilka, O., Van Kessel, A. T., Pratt, D. A., 2019. Beyond DPPH: use of fluorescence-enabled inhibited autoxidation to predict oxidative cell death rescue. *Cell Chem. Biol.*, 26(11), 1594-1607.
- Stockwell, B. R., 2022. Ferroptosis turns 10: Emerging mechanisms, physiological functions, and therapeutic applications. *Cell*, 185(14), 2401-2421.

* Presenting author. E-mail address: minh.bui-hoang@student.uibk.ac.at (Minh Bui Hoang)

S3.P20 Chinese traditional medicine for preventing hair loss in androgenic alopecia mice

Ching-Peng Wei^a, Man-Hsiu Chu^a, Ching-Kuo Lee^{a,b,*}

^a School of Pharmacy, College of Pharmacy, Taipei Medical University, Taipei 11031, Taiwan

^b Graduate Institute of Pharmacognosy, Taipei Medical University, Taipei 11031, Taiwan

Hair has a crucial role in protecting the skin from detrimental environmental elements, regulating body temperature, and contributing to an individual's personal identity in contemporary culture. The human hair follicle undergoes sequential stages, namely anagen, catagen, and telogen, throughout its lifespan. *Panax ginseng*, *Angelica sinensis*, and *Ligusticum chuanxiong* are frequently employed in Asia as a conventional botanical treatment for persons afflicted with alopecia. However, the effectiveness of this herbal ingredient has not been conclusively demonstrated. We aimed to ascertain the hair growth-stimulating impact of this botanical compound using an animal model. A cohort of male C57BL/6 mice, aged seven weeks, was randomly assigned to three groups: a blank group, a positive control group, and an experimental group. Each group consisted of five animals, except for the blank group. Each group was given Testosterone 0.05% by subcutaneous injection to induce male alopecia in the first seven days; the positive control group was given Finasteride 1%, the experimental group was given 1% of each extracts. The assessment of hair growth was conducted over a period of 21 days. The herbal mixture demonstrated hair growth-stimulating effects in C57BL/6 mice. The hair regrowth area was 3.04 (± 0.6)% for the blank sample, 4.66 (± 1.9)% negative control, and 5.98 (± 3.8)% positive control, experimental group containing herbal extracts are 5.14 (± 1.2) % for *Ligusticum chuanxiong*, 6.42 (± 1.2) % *Angelica sinensis* and 7.18 (± 2.3) % *Panax ginseng*. *Panax ginseng* has the best effect to improve the hair growth than control group and extracts.

Keywords: Androgenic Alopecia, *Panax ginseng*, *Angelica sinensis*, *Ligusticum chuanxiong*, hair growth

References

- Kishimoto, J. and Y. Nakazawa, Chapter 48 - Hair Physiology (Hair Growth, Alopecia, Scalp Treatment, etc.), in *Cosmetic Science and Technology*, K. Sakamoto, et al., Editors. 2017, Elsevier: Amsterdam. p. 767-780.
- Lévêque, M., B. Sandrine, and N. Castex-Rizzi, 1366 Interest of *Panax Ginseng* and *Albizia Julibrissin* extracts to prevent age-related hair density reduction. *Journal of Investigative Dermatology*, 2018. 138(5, Supplement): p. S232.
- Lourith, N. and M. Kanlayavattanakul, Hair loss and herbs for treatment. *J Cosmet Dermatol*, 2013. 12(3): p. 210-22. Ntshingila, S., et al., *Androgenetic alopecia: An update*. *JAAD International*, 2023. 13: p. 150-158.
- Tan, Y.-Z., et al., Structurally diverse phthalides from fibrous roots of *Ligusticum chuanxiong* Hort. and their biological activities. *Fitoterapia*, 2024. 175: p. 105882.

* Presenting author. E-mail address: cklee@tmu.edu.tw (Ching-Kuo Lee)

S3.P21 Qualitative and semi-quantitative screening of triterpenes in *Vaccinium uliginosum* L. fruits using supercritical fluid chromatography

Zuzana Vaneková*, Judith M. Rollinger

Division of Pharmacognosy, Department of Pharmaceutical Sciences, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

Bog bilberry (*Vaccinium uliginosum* L.) is a small arctic-alpine bush that produces berries similar to commercial blueberries. Hydrophilic constituents of bog bilberries are well researched, but there is sparse information about the lipophilic substances (Vaneková et al. 2024).

Freeze-dried and ground berries originating from Alaska, Norway, Finland, Estonia, Austria, Czechia and Slovakia were defatted with n-hexane and extracted 3x with dichloromethane. These extracts were analyzed using environmentally friendly, ultra-high-performance supercritical fluid chromatography (SFC) device coupled with PDA, ELSD and QDa detectors using supercritical CO₂ and ethanol as mobile phase. A sufficient separation of constituents was achieved on a Waters Torus 1-Aminoanthracene 1.7 µm column in 10 minutes (Goels et al. 2022).

The main constituents of the dichloromethane extract of all berry samples were triterpenes oleanolic and ursolic acid in an approximate ratio of 1:6, respectively. In comparison, a dichloromethane extract of European bilberry (*Vaccinium myrtillus* L.) fruits contains these major triterpenes in a ratio of 1:1. The content of oleanolic and ursolic acid in lyophilized samples ranged from 0.22 ± 0.06 mg/g and

1.04 ± 0.34 mg/g (in Delta Junction, Alaska) to 1.64 ± 0.17 mg/g and 5.67 ± 0.25 mg/g DW (in Seefeld, Austria), respectively. All samples originating from Alaska also contain micromeric acid (20(30) dehydrousolic acid), which is absent in all samples of European origin, aside from a single location in Austrian Waldviertel. All European samples contain a set of plant sterols, likely isomers of stigmastanol and cholestanol, which are absent in the Alaskan samples.

Keywords: *Vaccinium uliginosum*, Ericaceae, triterpenes, SFC

References

- Goels, T., Eichenauer, E., Langeder, J., Aichner, G. F., Mauser, G., Amtmann, L., Grienke, U., & Glasl, S. (2022). Ultra high-performance supercritical fluid chromatography for the quantitation of diterpene resin acids in Norway spruce samples. *Frontiers in Pharmacology*, 13. <https://doi.org/10.3389/fphar.2022.906411>
- Vaneková, Z., Holloway, P., & Rollinger, J. M. (2024). *Vaccinium uliginosum* L. (bog bilberry) and the search for its alleged toxicity: A Review. *Frontiers in Toxicology*, 6.

* Presenting author. E-mail address: zuzana.vanekova@univie.ac.at (Zuzana Vaneková)

S3.P22 Matrix free laser desorption ion mobility mass spectrometry in chemometrics: A new approach to the rapid identification of activity markers in complex mixtures of natural products

Manon Meunier^a, Martina Haack^b, Dania Awad^b, Thomas Brück^b, Khalijah Awang^c, Marc Litaudon^d, Frédéric Saubion^e, Marc Legeay^e, Dimitri Bréard^a, David Guilet^a, Séverine Derbré^a, Andreas Schinkovitz^{a,*}

^a Univ Angers, SONAS, SFR QUASAV, Faculty of Health Sciences, Dpt Pharmacy, 16 Bd Daviers, 49045 Angers CEDEX 01, France

^b Technical University of Munich, TUM School of Natural Sciences, Werner Siemens-Chair of Synthetic Biotechnology (WSSB), Lichtenbergstraße 4, 85748 Garching, Germany

^c University of Malaya, Faculty of Sciences, Department of Chemistry, Malaysia

^d Institut de Chimie des Substances Naturelles, CNRS-ICSN, UPR 2301, Université Paris-Saclay, 91198, Gif-sur-Yvette, France

^e UNIV Angers, LERIA, EA2645, SFR MathSTIC, Faculty of Sciences, 2 boulevard Lavoisier, 49045 Angers cedex 01, France

Matrix free laser desorption ionization mass spectrometry (LDI-MS) is a highly efficient tool for the quick chemical characterization of complex mixtures of Natural Products (Nps) (Islam et al. 2022). Showing very close structural similarities to compounds used as matrices in matrix assisted LDI-MS, many NPs can be easily ionized by simple laser irradiation. Compared to LC-MS the method yields instant results, works with any volatile solvent, and hardly requires sample preparation (Schinkovitz et al. 2018).

With this in mind, the current presentation highlights recent advances in LDI-MS and its integration into chemometric workflows. As a practical example, the correct prediction and consecutive isolation of three bioactive xanthenes (rubraxanthone, isocowanol and parvixanthone) from a crude bark extract of *Garcinia parvolia* (Miq) (Clusiaceae) is presented (Meunier et. al. 2023). All isolated compounds exhibited moderate inhibitory effects on the formation of advanced glycation end products (IC₅₀: 123-170 mM). In addition, the hyphenation of LDI tandem Mass Spectrometry with Ion Mobility Spectrometry (LDI-IMS-MS²) will be discussed. The latter facilitated the distinct differentiation of xanthone isomers without chromatographic separation and permitted the creation of the first molecular network that is entirely based on LDI-MS².




Fig. 1. The concept of high resolution LDI-IMS-MS²

Keywords: Matrix free laser desorption mass spectrometry, chemometrics, xanthenes, ion mobility, molecular networks

References

- Islam S., Alam R., Kim S., 2022. Improved coverage of plant metabolites using powder laser desorption/ ion-ization coupled with Fourier-transform ion cyclotron mass spectrometry. Food Chem. 373, 131541.
- Meunier M., Bréard D., Awang K., Boisard S., Guilet D., Richomme P., Derbré S., Schinkovitz A., 2023. Matrix free laser desorption ionization assisted by ¹³C NMR dereplication: An alternative approach to LC-MS² based chemometrics. Talanta 253, 123988.
- Schinkovitz A., Boisard S., Freuze I., Osuga J., Mehlmer N., Brück T., Pascal Richomme P., 2018. Matrix free laser desorption ionization mass spectrometry as a functional tool for the analysis and differentiation of complex phenolic mixtures in propolis: A new approach to quality control. Anal. Bioanal. Chem. 410, 6187–6195.

* Presenting author. E-mail address: andreas.schinkovitz@univ-angers.fr (Andreas Schinkovitz)

S3.P23 Metabolomic study of tea tree oil and *Melaleuca* spp. using HPTLC, LC-MS and GC-MS

Yi Zhao^{a,b,*}, Naomi Nzisa^c, Xavier Cheseto^d, Anne Osano^e, Jiangnan Peng^f

^a Department of Biological Sciences, Lehman College, The City University of New York, Bronx, 10468 NY, USA

^b Biology Ph.D. Program, The Graduate Center, The City University of New York, New York, 10016 NY, USA

^c Department of Analytical Chemistry and Food Safety, Ministry of Livestock Department, 30028 Nairobi, Kenya

^d International Centre of Insect Physiology and Ecology, 30772 Nairobi, Kenya

^e Department of Natural Sciences, Bowie State University, Bowie, 20715 MD, USA

^f Department of Chemistry, Morgan State University, Baltimore, 21251 MD, USA

Tea tree oil (TTO) is an essential oil obtained from the leaves and terminal branches of *Melaleuca alternifolia*, and is widely used in cosmetics, household products, and dermatology treatments (Yadav et al., 2017). The concentration range of the main components in TTO including 15 terpenes and terpenoids is defined by an international standard (International Organization of Standardization, 2017). This study aims to evaluate the quality of commercial TTOs and explore the commercial potential of oils from other *Melaleuca* species using metabolomic techniques for chemical profiling. TTOs purchased from the United States and Kenya, along with terminal branches of *M. salicina* and *M. citrina* were extracted, diluted, and analyzed using high-performance thin-layer chromatography (HPTLC), liquid chromatography-mass spectrometry (LC-MS), and gas chromatography-mass spectrometry (GC-MS). A total of 120 compounds were tentatively identified, including all 15 markers. The results showed that the commercial oils purchased from markets did not meet the international standard in comparison with the TTO provided by Kenyan Forestry Research Institute. The oils from two closely related species, *M. salicina* and *M. citrina*, are distinct from TTO based on their chemical profile. Three polar compounds, grandinol, multifidol, and 3,3-di-*O*-methylellagic acid--*O*-D-glucopyranoside were detected in *M. citrina* using LC-MS, while only the first two compounds were detected in *M. salicina*. None of the samples showed potent antioxidant activity through a DPPH assay on an HPTLC plate.

Keywords: tea tree oil, *Melaleuca* spp., HPTLC, LC-MS, GC-MS

References

Yadav, E., Kumar, S., Mahant, S., Khatkar, S., Rao, R, 2017. Tea tree oil: a promising essential oil. J. Essent. Oil Res. 29 (3), 201–213.

International Organization for Standardization, ISO 4730:2017, 2017. Essential oil of *Melaleuca*, terpinen-4-ol type (tea tree oil).

* Presenting author. E-mail address: yi.zhao@lehman.cuny.edu (Yi Zhao)

S3.P24 Preclinical evaluation of oral nanoformulation based on natural extract as a potential treatment for type 2 diabetes mellitus

Elkin Escobar^{a,b}, Natalia Arbelaez^c, Adriana Restrepo^c, Diana Muñoz^d, Jahir Orozco^b, Norman Balcazar^{a,d*}

^a GENMOL Group, ^cPECET Group, Faculty of Medicine, ^dDepartment of Physiology and Biochemistry, Faculty of Medicine. Universidad de Antioquia, Calle 70, N° 52-21, A.A. 1226, Medellín, Colombia

^b Max Planck Tandem Group in Nanobioengineering, University of Antioquia. Complejo Ruta N, Calle 67 N° 52-20, Medellín, 050010, Colombia

An oral nanoformulation (ONF) based on natural extracts (OBE100), ursolic acid, oleanolic acid, and ursolic acid lactone obtained by hexane: methanol: water 4:3:1(v/v) extraction from *Eucalyptus tereticornis* Sm. (Myrtaceae) leaves reduce metabolic alterations in a diet-induced obese mouse (DIO) model (1)(2). The work aims to develop preclinical safety and biological efficacy tests for the ONF. Three groups were established: vehicle, empty nanoparticles, and ONF OBE100 encapsulated in nanoparticles; 214 mg/kg OBE100 for repeated dose oral toxicity for 28, 90 days and combine chronic toxicity/carcinogenicity (3). The ophthalmological response to proprioception, threat reflex, and ocular evaluations were performed. At the endpoint, blood samples for hemogram and blood chemistry were taken. All mice were euthanised and necropsied, and samples of the internal organs were taken for histopathological analysis.

Results showed that ONF is biologically effective in decreasing metabolic abnormalities in a DIO model. No clinical abnormalities were recorded during the administration of repeated doses of oral toxicity for 28, 90 days and chronic toxicity. Body weight gain was stable, and all groups' hemograms and blood chemistry analyses were within normal ranges. No signs of neurological or ocular toxicity were recorded during the study. Macroscopically, no lesions were observed in any organ, and the weight of the internal organs was homogeneous for the different treatments.

The findings of acute oral toxicity, oral toxicokinetics, repeated doses of oral toxicity for 28 and 90 days, and chronic toxicity indicate that the compound ONF is safe based on clinical and paraclinical evaluation and preliminary histopathological analysis.

Keywords: *Eucalyptus tereticornis*, nanoformulation, preclinical studies, type 2 diabetes mellitus

References

- (1) Escobar E., Acin S., Muñoz, DL., Fernandez, M., Echeverri, A., Echeverri, F., Orozco, J., Balcazar N. 2023. Polymeric nanoformulation prototype based on a natural extract for the potential treatment of type 2 diabetes mellitus. Journal of Drug Delivery Science and Technology. 81: 104264. <https://doi.org/10.1016/j.jddst.2023.104264>.
- (2) Norman Balcazar, Sergio Acin, Luis Alberto Gonzalez. Non-targeted Profiling Analysis by GC/MS and LC/MS/MS of extracts derived from *Eucalyptus tereticornis* with antiobesogenic and antidiabetic activity. POSTER ICNPR 2024.
- (3) <https://www.oecd.org/env/ehs/testing/section4-health-effects-replaced-and-cancelled-test-guidelines.htm>

*Presenting author. E-mail address: norman.balcazar@udea.edu.co (Norman Balcazar)

S3.P25 Employing Plant-Bacterial Interactions to Access the Metabolic Capacity of Bacteroidetes

Barbara I. Adaikpoh*

Department of Pharmaceutical Sciences, and Center for Biomolecular Sciences, University of Illinois at Chicago, Chicago IL 60607, USA

To improve the discovery of unusual bioactive natural products, it is necessary to explore untapped bacterial taxa. Bacteroidetes are often found interacting with eukaryotic hosts like plants, and several species have been shown to have plant-beneficial attributes (Pan et al., 2023). The metabolic capacity to produce bioactive natural products by this phylum is yet to be explored (Brinkmann et al., 2022). Previous studies prove metabolic impacts from a bacterial perception of plant signals (Adaikpoh et al., 2020). We aim to explore the metabolic and transcriptomic responses of our Bacteroidetes strains to various plant metabolites often found within the rhizosphere. We hypothesize that exposure to plant metabolites will result in the production of natural products not observed under traditional cultivation methods. Phenotypic changes observed in the lab in response to selected plant metabolites could be evidence of significant metabolic changes. Analysis of RNA sequence data in addition to LC-MS/MS datasets and antimicrobial assays of extracts from supplemented cultures will be presented. This project provides data for optimizing culture conditions for the discovery of novel bioactive natural products from Bacteroidetes.

Keywords: plant-bacteria interactions, bacteroidetes, natural products, transcriptomics, metabolomics

References

- Adaikpoh, B.I., Akbar, S., Albataineh, H., Misra, S.K., Sharp, J.S., Stevens, D.C., 2020. Myxobacterial Response to Methyljasmonate Exposure Indicates Contribution to Plant Recruitment of Micropredators. *Front. Microbiol.* 11, 34.
- Brinkmann, S., Spohn, M.S., Schäberle, T.F., 2022. Bioactive natural products from Bacteroidetes. *Nat. Prod. Rep.* 39, 1045–1065.
- Pan, X., Raaijmakers, J.M., Carrión, V.J., 2023. Importance of Bacteroidetes in host–microbe interactions and ecosystem functioning. *Trends in Microbiology* 31, 959–971.

*Presenting author. E-mail address: biada@uic.edu (Barbara I. Adaikpoh)

S3.P26 *Epilobii herba*: Quality assessment using chromatographic techniques

Olha Mykhailenko^{a,b}, Banaz Jalil^a, Michael Heinrich^{a,c,*}

^a Pharmacognosy and Phytotherapy, UCL School of Pharmacy, 29-39 Brunswick Square, WC1N 1AX, London, UK

^b Department of Pharmaceutical Chemistry, National University of Pharmacy, 4-Valentynivska Str., 61168, Kharkiv, Ukraine

^c Chinese Medicine Research Center, College of Chinese Medicine, China Medical University, Taichung City 404, Taiwan

The genus *Epilobium* (willow herb) includes ca. 160 species distributed throughout the world. The most rudely used species are *E. angustifolium* L., *E. parviflorum* Schreb., and *E. hirsutum* L. as herbal teas, which have been reported to have prostate-protective and anti-inflammatory properties (1). However, despite their wide use, these plants are not included in key pharmacopoeias. Only the Herbal Medicines Committee (HMPC) has assessed the botanical drug (2, 3) regarding *E. parviflorum*'s medical use, but not in terms of metabolites or analysis.

Here, we focus on a comparative qualitative analysis for *E. hirsutum* herb and the assessment of the homogeneity of samples during a growing season. From April until October 2023 *E. hirsutum* samples (n=78), including leaves and stems, were collected in the United Kingdom. Polyphenols, e.g., chlorogenic acid, gallic acid, caffeic acid, avicularin, guajaverin, isoquercitrin and hyperoside used as reference standards. The analysis was carried out in HPTLC plates Si 60 F₂₅₄ (Merck) in mobile phase: ethyl acetate: formic acid: water (68:8:8) and 2-aminoethyldiphenylborinate and macrogol 400 for derivatization.




Fig. 1. HPTLC profile under UV 254 nm (A), UV 366 nm (B) prior to derivatization, and under UV 366 nm after derivatization (C), and white light after derivatization (D)

The HPTLC analysis showed all reference substances in *E. hirsutum* extracts but in different concentrations, e.g., yellow fluorescent zones ($R_f=0.4$; $R_f=0.38$; $R_f=0.85$; $R_f=0.52$) were in line with isoquercitrin, hyperoside, avicularin and guajaverin, and these were the dominant compounds in leaves. The light blue, fluorescent zone ($R_f=0.7$) was identified as gallic acid, and it accumulated more in the stems (Fig. 1). The presented method can be used for assessing the *Epilobium* quality, including botanical drugs or finished commercial products.

Keywords: *Epilobium hirsutum*, Onagraceae, quality control, HPTLC

References

1. Vlase AM, Toiu A, Tomuț I, Vlase, L.; Muntean, D.; Casian, T.; Fizesan, I.; Nădăraș, G.C.; Novac, C.S.; Tămaș, M.; et al. *Epilobium* Species: From Optimization of the Extraction Process to Evaluation of Biological Properties. *Antioxidants* 2023, 12, 91.
2. EMA/HMPC/72238/2022. Call for scientific data for the periodic review of the monograph on *Epilobii herba* (PDF/107.1 KB), published: 15/02/2022. European Medicines Agency, 2022. Amsterdam, The Netherlands.
3. EMA/HMPC/615965/2022. Addendum to Assessment report on *Epilobium angustifolium* L. and/or *Epilobium parviflorum* Schreb., published: 17/11/2022. European Medicines Agency, 2022. Amsterdam, The Netherlands.

* Presenting author. E-mail address: o.mykhailenko@ucl.ac.uk (Olha Mykhailenko)

S3.P27 Phytochemical analysis and quality control study of *Dendrobium fimbriatum* Hook. flowers

Anham Abbasi^{a,*}, Yuan-Shiun Chang^b, Wu-Chang Chuang^c, Hung-Rong Yen^{d,e,f,#}, Michael Heinrich^{b,d,g,#}, Chia-Lin Lee^{d,h,*}

^a School of Pharmacy, College of Pharmacy, China Medical University, Taichung, Taiwan

^b Department of Chinese Pharmaceutical Science and Chinese Medicine Resources, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^c Brion Research Institute of Taiwan, New Taipei City, Taiwan

^d Chinese Medicine Research Center, China Medical University, Taichung, Taiwan

^e School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^f Department of Chinese Medicine, China Medical University Hospital, Taichung, Taiwan

^g Pharmacognosy and Phytotherapy Group, UCL School of Pharmacy, University College of London, UK

^h Department of Cosmeceutics, College of Pharmacy, China Medical University, Taichung, Taiwan

Dendrobium fimbriatum Hook (Orchidaceae) is a renowned edible and medicinal plant in Chinese medicine used traditionally for the treatment of yin deficiency, clearing heat from the body, and nourishing the stomach. A literature review suggests that there has been no major study on pharmacognostic and phytochemical aspects of the flowers.

This study aims to develop an analytical framework for the chemical characterization of the *Dendrobium fimbriatum* flowers in combination with defining basic pharmacognostic parameters. We investigated different extracts/fractions including crude methanol and ethanol extracts sourced from a commercial production site in Taiwan to characterize the phytochemical profile using chromatography (HPTLC, LC and GC), spectroscopic (NMR) and spectrometric (MS) analysis. The samples were analyzed using proton-NMR and carbon-NMR to identify the metabolite composition and to assess similarities and differences between crude methanol and ethanol extracts/fractions. The samples were subsequently analyzed using HPTLC to assess if there were any benefits gained by employing a dual analysis strategy.

No significant difference was found in the chemical composition of both methanol and ethanol extracts/fractions by NMR. Nevertheless, HPTLC detected the existence of metabolites such as terpenes and flavonoid markers, including quercetin, rutin, ursolic acid, and lupeol. The combination of chromatographic, spectroscopic, and spectrometric techniques forms the basis for the authentication and quality control of *Dendrobium fimbriatum* flowers. Further research is ongoing to define the best marker compounds.

Using HPTLC technique to analyze the crude methanol and ethanol extracts of *Dendrobium fimbriatum* flowers showed the presence of terpenes when tested in comparison with markers lupeol and ursolic acid.

Fig. 1. HPTLC of crude MeOH and EtOH extracts of *Dendrobium fimbriatum* flowers.

Keywords: *Dendrobium fimbriatum* flowers, Standardization, NMR, Quality control study, HPTLC

References

1. da Costa Mendes, D. S. (2013). *Chromatographic Fingerprinting and Quality Control of Herbal Medicines: Comparison of two official Chinese pharmacopoeia species of Dendrobii based on High-Performance Liquid Chromatography and Chemometric analysis* (master's thesis, The University of Bergen and Central South University, PR China).
2. Sunita Shailajan, S. S., Suman Kumaria, S. K., Deepti Gurjar, D. G., Mayuresh Joshi, M. J., Prasenjit Paul, P. P., & Khongthaw, N. (2015). Variation in the marker content of five different *Dendrobium* species: comparative evaluation using validated HPTLC technique.
3. Sinha, S. N., & Biswas, K. (2020). Phytochemical Screening and Evaluation of Antimicrobial Potential of *Dendrobium fimbriatum* Hook. *Orchid Biology: Recent Trends & Challenges*, 531-538.

*Presenting author. E-mail address: anhamabbasi1@gmail.com (Anham Shahid Abbasi)

S3.P28 Identification of chemical defenses of a tropical tree *Sextonia rubra* using molecular networks from GC-MS, LC-MS/MS and MALDI-MS/MS imaging data

Levasseur Marceau^a, Eparvier Véronique^a, Houël Emeline^b, Touboul David^{c,*}

^a Université Paris-Saclay, CNRS, Institut de Chimie des Substances Naturelles, UPR 2301, 91198, Gif-sur-Yvette, France

^b Sorbonne Université, CNRS, Laboratoire de Biodiversité et Biotechnologies Microbiennes, LBBM, Observatoire Océanologique, 66650 Banyuls-sur-Mer, France

^c LCM, CNRS, Ecole polytechnique, Institut Polytechnique de Paris, Route de Saclay, 91120 Palaiseau, France

Sextonia rubra is a tropical tree endemic to the Guiana Shield and the Brazilian Amazon. It is known for its heartwood natural durability that comprises numerous lactone derivatives (Rodrigues et al., 2010). However, its fruits have not been studied chemically. Here we propose analytical methods to explore the chemical diversity of these fruits using SPME-GC-MS, LC-HRMS/MS and MALDI-FT-ICR MS imaging. For the first time, GC-MS and MALDI-MS/MS (Levasseur et al., 2024) data were analyzed using molecular networks for efficient annotation using MetGem software (Olivon et al., 2018).

The most abundant volatile compounds identified by SPME-GC-MS in the hydrolate were eucalyptol (13.5%), α calamenene (7.4%), β caryophyllene (7.4%), α copaene (7.3%) and δ cadinene (5.8%). The antimicrobial activity of the hydrolate was tested against pathogens with a relative MIC equal to 5%.

In parallel of chemical analysis, wood decay resistance was assessed using long-term soil bed tests. Ethyl acetate extracts of different parts of the tree were tested against 6 Glutathione-S-Transferase (GST) of *Trametes versicolor*, corollary of their anti-fungal activity (Barbier et al., 2020).

Our results suggest that for pith and heartwood, the higher the concentration ratios of lactones towards alkaloids, the higher the reactivity of GSTs, and the more the natural durability against wood-decaying fungi is positively affected. Although the involvement of lactone derivatives in the natural durability of this species was known, the involvement of alkaloids is new and suggests that *S. rubra* specializes its chemical defenses according to the tissue.

Keywords: *Sextonia rubra*, metabolomic, tandem mass spectrometry, MALDI imaging, molecular network

References

- Barbier, M., et al., 2020. Glutathione Transferases: Surrogate Targets for Discovering Biologically Active Compounds. *J. Nat. Prod.* 83, 2960-2966.
- Levasseur, M., Nicol, E., Elie, N., Houël, E., Eparvier, V., Touboul, D., 2024. Spatialized Metabolomic Annotation Combining MALDI Imaging and Molecular Networks. *Anal. Chem.* 96(1), 18-22.
- Olivon, F., Elie, N., Grelier, G., Roussi, F., Litaudon, M., Touboul, D., 2018. MetGem Software for the Generation of Molecular Networks Based on the t-SNE Algorithm. *Anal. Chem.* 90, 13900-13908.
- Rodrigues, A., Theodoro, P., Eparvier, V., Basset, C., Silva, M., Beauchêne, J., Espíndola, L., Stien, D., 2010. Search for antifungal compounds from the wood of durable tropical trees. *J. Nat. Prod.* 73, 1706-1707.

* Presenting author. E-mail address: david.touboul@cnrs.fr (David Touboul)

S3.P29 An ethnobotanical, pharmacological and phytochemical review on flowers of Shi-hu (*Dendrobium* spp.)

Anham Abbasi^{a#}, Michael Heinrich^{b,c,d}, Hung-Rong Yen^{d,e,f,*}

^a School of Pharmacy, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^b Department of Chinese Pharmaceutical Science and Chinese Medicine Resources, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^c UCL School of Pharmacy, University College London, London, UK

^d Chinese Medicine Research Center, China Medical University, Taichung, Taiwan

^e School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^f Department of Chinese Medicine, China Medical University Hospital, Taichung, Taiwan

The genus *Dendrobium* is a medicinal plant with nutraceutical importance. *Dendrobium* species have been used to nourish “Yin” and tonify the five viscera in traditional Chinese medicine. The Taiwan Herbal Pharmacopoeia (4th English Edition, 2022) includes the fresh or dried stem of seven *Dendrobium* species (Shi-hu in Chinese) in the monograph. However, it is not clear how the flower part of these seven species could be used.

We reviewed various aspects of the chemical composition and pharmacological characteristics of *Dendrobium* flowers aiming at defining future research and development priorities. Additionally, traditional uses, conservation status and cultivation techniques relevant to ascertain a sustainable use of the species were summarized.

Web of Science, SciFinder, Scopus, PubMed, Google Scholar and CNKI were used to gather data on the phytochemistry and pharmacological studies of *Dendrobium* flower usage. The keywords for this study are *Dendrobium*, Pharmacology and Chemistry.

Open column chromatography, GC-MS, UPLC, HPLC and GC-FID are common separation and analytical techniques that have been utilized for the isolation, characterization and quantification of polysaccharides, alkaloids, essential oils, and flavonoids. They have prominent pharmacological/biological activities like pro-apoptotic, immune-modulatory, antioxidant and anti-microbial activities. The most investigated species of flowers are *D. officinale* Kimura & Migo. and *D. chrysanthum* Wall. ex Lindl.

This study sheds light on the historical uses of *Dendrobium* flowers, their current clinical significance and future potential. The study systematically assesses the link between chemical composition, pharmacological activity and current or potential uses. According to our findings, in-depth investigations are needed in areas where there is a paucity of knowledge on the potential of flowers, such as quality control, phytochemical characterization, and dosage form development.

Acknowledgment

This research is partially supported by the Yushan Fellow Program by the Ministry of Education (MOE), Taiwan. This work is also financially supported by “Chinese Medicine Research Center, China Medical University” from the Featured Areas Research Center Program within the framework of the Higher Education Sprout Project by the Ministry of Education (MOE), Taiwan.

#Presenting author. E-mail address: anhamabbasi1@gmail.com (Anham Shahid Abbasi)

S3.P30 Patients' use of medicinal plants in the urology service of Ibn-Sina hospital in Rabat, Morocco: integrating ethnobotanical and pharmacovigilance approaches

Khadija Dabra, Souad Skalli

Center of Plant and Microbial Biotechnology, Biodiversity and Environment, Department of Biology, Faculty of Sciences, Mohammed V University in Rabat, Morocco

Phytotherapy is a very old medical discipline, based on the use of medicinal plants or traditional pharmacopoeia products to treat various pathologies. It depends on the socio-economic context, and given the wealth of medicinal plants in Morocco, the use of phytotherapy has only increased (Bouzouita, 2016). While popular belief assumes that herbal remedies are natural and therefore mistakenly considered harmless, scientific research has shown that these plants can contain powerful chemical compounds that can induce intoxication and serious, even fatal, side effects. The aim of this work is to carry out an ethnobotanical study on the use of medicinal and aromatic plants and traditional Moroccan pharmacopoeia products among patients with disorders and diseases of the urinary system, in the urology department of the Ibn Sina hospital in Rabat. The Rabat-Salé-Kenitra region is one of the 12 regions of Morocco, created in 2015 following the new territorial division of the regions. It is limited to the north by the region of Tangier-Tetouan-Al Hoceima, to the southeast by the region of Fez-Meknes and to the south by the two regions of Casablanca-Settat and Beni-Mellal-Khénifra and to the west by The Atlantic Ocean. This region covers a total area of 17,569 km², or 2.5% of that of Morocco (3). Note that the study region was chosen for the traditional know-how of its local population (4) and for its floristic and ecological diversity (2). A semi-structured questionnaire was used, targeting patients in the urology department of the Ibn Sina Rabat hospital. Data was collected from 649 patients, in full respect of patient confidentiality and anonymity, and the questionnaire used consists of 44 questions and four items: patient identity, patient pathology, use of medicinal plants, and adverse effects (AEs) related to the use of plants. Although the questionnaire was written in French, oral interviews with patients were conducted in Moroccan dialect. In addition to completing the questionnaire designed for this purpose, additional information was recorded for each day of the survey, namely:

- the total number of patients hospitalized;
- the total number of consulting patients
- the total number of patients who refused to take part in the survey;
- the total number of patients who took part in the survey but did not use plants to treat their illness;
- the total number of patients who took part in the survey and used herbs to treat their illness, but did not experience any adverse effects as a result;
- the total number of patients who took part in the survey and who used plants to treat their illness and who experienced undesirable side-effects as a result.
- patient encountered in the various health facilities surveyed;
- patients who voluntarily agreed to take part in the survey;
- patients who use plants alone or in mixtures with or without prescribed conventional treatments.
- patients who have experienced undesirable effects as a result of using plants.

For socio-demographic data, the age ranges used in this study were established with reference to the age range classification provided by the WHO in its latest World Diabetes Report (WHO, 2016). Among 656 patients, 29,88 % used medicinal plants. Floristic analysis of the results obtained identified 37 species belonging to 23 plant families. 23 patients from the urology department reported the occurrence of one or more AEs following the use of medicinal plants, with a prevalence of 3.54%. 15 plants linked to these AEs were listed. *Eruca sativa* L. was associated with 3 (13,04 %) cases of AEs, and *Petroselinum crispum* L. with 2 (8,69 %).

References

- Bouzouita, K. (2016). *Phytovigilance: enquête auprès des pharmaciens officinaux d'Oujda* (Doctoral dissertation).
- DGCL. Direction Générale de Collectivités Locales, Maroc. La Région de Rabat-Salé-Kénitra: Monographie générale. 2015. <https://www.regions-maroc.ma/wp-content/uploads/2020/10/MONOGRAPHIE-DE-LA-REGION-DE-RABAT-SALE-KENITRA-FR.pdf>
- HCP. Haut-Commissariat Au Plan du Royaume du Maroc. Présentation de la Région Rabat-Salé-Kénitra. 2015. Main report. <https://www.hcp.ma/region-rabat/>
- Ministry of agriculture, Fisheries, Rural Development, Water and Forests, Morocco. Monographie de la région de Rabat-Salé-Kenitra. 2023. <https://www.agriculture.gov.ma/fr/region/rabat-sale-kenitra>

*Presenting author. E-mail address: khadijadabra1996@gmail.com (Khadija Dabra)

S3.P31 Unveiling the multifaceted efficacy of Gu Sui Bu in osteoporosis treatment: insights from a zebrafish model and advanced analytical approaches

Wen-Ying Lin^a, Cheng-Huan Peng^b, Ming-Der Lin^{a,c,*}

^a Department of Molecular Biology and Human Genetics, Tzu Chi University, Hualien, 97004, Taiwan

^b Department of Orthopedics, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, 97002, Taiwan

^c Institute of Medical Science, Tzu Chi University, Hualien, 97004, Taiwan

Osteoporosis significantly increases the risk of fractures and adversely affects the quality of life, particularly in the elderly. Existing osteoporosis treatments are often expensive and may lead to severe side effects with long-term use (Peng et al., 2021). In this scenario, herbal medicine, celebrated for its diverse bioactive components and multifaceted action, presents a promising alternative with potentially fewer side effects. Specifically, Gu Sui Bu (GSB), the dried rhizome of *Drynaria fortunei* J. Sm., is acclaimed in traditional Chinese medicine for its bone-healing capabilities. Our prior research using a zebrafish model has revealed GSB's dual role in reducing bone resorption and enhancing bone formation (Peng et al., 2022). Zebrafish, as an *in vivo* model, are instrumental in assessing both the direct and indirect effects of the drug, providing comprehensive insights into its mechanisms. To decipher GSB's complex therapeutic mechanisms, we implemented a sophisticated multidimensional analytical approach, including RNA-seq analysis, a network pharmacology approach, and molecular docking for the active ingredients of GSB on the zebrafish osteoporosis model. By comparing the gene expression profiles before and after the treatment with 1 mg/ml GSB in a glucocorticoid-induced osteoporosis zebrafish model, we identified 469 genes whose adverse expression changes due to glucocorticoid treatment were reversed by GSB. According to the results of Gene Ontology enrichment analysis, genes involved in osteoblast differentiation and ossification were identified. Among them, TMEM119 particularly piqued our interest; it could stimulate the expression of *runx2*, *sp7*, *collagen 1a1*, *osteopontin*, and *alkaline phosphatase*, crucial for osteoblast differentiation. The expression of TMEM119 decreased 1.6-fold after dexamethasone treatment but returned to normal levels after GSB treatment. In addition to exploring GSB's effects using RNA-seq analysis, we also employed network pharmacological analysis to investigate the direct targets of GSB. We collected 117 chemical ingredients of GSB from TCM- ID, LTM-TCM, INPUT 2.0, and TCMSP databases, and predicted 1011 possible action targets using the Swiss Target Prediction and Similarity Ensemble Approach web servers. Interestingly, over-representation analysis revealed key signalling pathways, such as Hedgehog and FGF, as potential GSB targets to activate *runx2* expression for osteoblast differentiation. We further performed molecular docking to demonstrate that Naringin, an active ingredient in GSB, could bind to FGF receptors. Considering the multi-active component and multi-target nature of herbal drugs, adopting a multidimensional analytical approach could facilitate the understanding of the action mechanisms of herbal drugs and establish a foundation for the innovative development of cocktail therapy, combining prescription drugs with herbal medicine. This aims to improve treatment outcomes and reduce side effects in clinical practice.

Keywords: *Drynaria fortunei*, herbal medicine, osteoporosis, osteogenesis, zebrafish

References

- Peng, C.H., Lin, W.Y., Li, C.Y., Dharini, K.K., Chang, C.Y., Hong, J.T., Lin, M.D., 2022. Gu Sui Bu (*Drynaria fortunei* J. Sm.) antagonizes glucocorticoid-induced mineralization reduction in zebrafish larvae by modulating the activity of osteoblasts and osteoclasts. *J Ethnopharmacol* 297, 115565.
- Peng, C.H., Lin, W.Y., Yeh, K.T., Chen, I.H., Wu, W.T., Lin, M.D., 2021. The molecular etiology and treatment of glucocorticoid-induced osteoporosis. *Tzu Chi Med J* 33, 212-223.

* Presenting author. E-mail address: mingder@gms.tcu.edu.tw (Ming-Der Lin)

S3.P32 Phytochemical profiling and the wound healing potential of five *Bulbine* species using *in vitro* and *in vivo* assays

Ramoagi Segone^{a,*}, Alvaro Viljoen^{a,b}, Maxleene Sandasi^{a,b}

^a Department of Pharmaceutical Sciences, Tshwane University of Technology, Private Bag X680, Pretoria 0001, South Africa

^b SAMRC Herbal Drugs Research Unit, Faculty of Science, Tshwane University of Technology, Pretoria 0001, South Africa

Traditionally, the leaf exudates of *Bulbine* species are used to treat skin conditions (Bodede and Prinsloo, 2020), which has led to the development of bulbine-containing cosmetic products. However, scientific evidence to support the claimed benefits is still lacking, hence the need for evidence-based methods to investigate the medicinal properties of some indigenous *Bulbine* species. In the current study, the *in vitro* scratch assay and *in vivo* zebrafish caudal fin amputation assay were used to evaluate the wound healing properties of five ethnobotanically important species; *Bulbine abyssinica*, *B. asphodeloides*, *B. frutescens*, *B. latifolia* and *B. narcissifolia*. Phytochemical profiling was achieved using ultra-performance liquid chromatography coupled to mass spectrometry, and knipholone was identified as the marker compound which varied quantitatively across the five species. Pharmacologically, *B. frutescens* and *B. abyssinica* (100 µg/mL) extracts accelerated HaCaT cell migration and subsequent wound closure in the scratch assay (96% and 94% acceleration after 24hrs, respectively). Similarly, the two species increased caudal fin regeneration by 90% and 83% after 72 hrs for *B. frutescens* and *B. abyssinica*, respectively. The results of the study align with the traditional use of *B. frutescens* and *B. abyssinica* in wound healing, which could inform further research for cosmetic and pharmaceutical product development involving these species.

Keywords: *Bulbine* species, wound healing, zebrafish larvae, scratch assay, knipholone

References

Bodede, O., Prinsloo, G., 2020. Ethnobotany, phytochemistry and pharmacological significance of the genus *Bulbine* (Asphodelaceae). J. Ethnopharmacol. 260, 112986.

* Presenting author. E-mail address: segoner@tut.ac.za (Ramoagi Segone)

S3.P33 Translating community-wide mass spectral libraries into actionable chemical knowledge: a proof of concept with monoterpene indole alkaloids

Sarah Szwarc^{a,*}, Yassine M. Mejri^{a,b}, Meltem Öztürk^b, Olivier Cailloux^b, Vincent Courdavault^c, Kyungha Lee^d, Kyo Bin Kang^d, Adriano Rutz^e, Justin J.J. van der Hooft^f, Florian Huber^g, Erwan Poupon^a, Pierre Le Pogam^a, Mehdi A. Beniddir^a

^a Université Paris-Saclay, CNRS, BioCIS, 91400 Orsay, France

^b Université Paris Dauphine, PSL Research University, 75016 Paris, France

^c University of Tours, EA2106, Faculty of Pharmacy, Tours, France

^d College of Pharmacy, Sookmyung Women's University, Seoul 04310, Republic of Korea

^e Institute of Molecular Systems Biology, ETH Zurich, Otto-Stern-Weg 3, Zürich, 8093, Switzerland

^f Bioinformatics Group, Wageningen University & Research, 6708 PB, Wageningen, the Netherlands

^g Centre for Digitalisation and Digitality, Düsseldorf University of Applied Sciences, 40476 Düsseldorf, Germany

Mass spectral libraries play a key role in annotating metabolite features in untargeted metabolomics profiles. Here, we will present the latest update of the MIADB (Fox Ramos et al., 2019), a public spectral library containing fragmentation spectra of more than 400 unique monoterpene indole alkaloids (MIAs) and the tools we developed to extract chemical knowledge from it.

MIAs represent one of the most complex and rich classes of natural products with more than 4000 representatives to date. This chemical class is also renowned for the valuable biological activities associated with some of its representatives such as vinblastine, quinine and camptothecin. One objective of this collaborative project was to chart the MS/MS spectral space of this chemically diverse library using several mass spectral similarity scores. We used molecular networking and heatmaps to map those similarities. MIAs were first classified according to their structural similarity, determined by Tanimoto scoring. As anticipated, compounds with the highest chemical similarities generally demonstrated greater MS/MS similarities, irrespective of the chosen spectral similarity score. For each MIA skeleton, the commonly shared mass fragments and neutral losses were then extracted and used as MassQL (Jarmusch et al., 2022) queries against a dataset of 71 MIAs-producing plants and 4 non-producing plants adding a further layer to the many available dereplication tools. Finally, this database was confronted to state-of-the-art MS/MS annotation tools. Overall, we believe that the MIADB is a great example highlighting a general strategy for extracting spectral and chemical information from a variety of metadata-informed spectral libraries.

Keywords: Metabolomics, spectral library, monoterpene indole alkaloids, chemo-informatics, MS/MS

References

- Fox Ramos, A. E.; Le Pogam, P.; Fox Alcover, C.; Ootogo N'Nang, E.; Cauchie, G.; Hazni, H.; Awang, K.; Bréard, D.; Echavarren, A. M.; Frederich, M., 2019. Collected mass spectrometry data on monoterpene indole alkaloids from natural product chemistry research. *Scientific data*. 6, 1-6.
- Jarmusch, A. K.; Aron, A. T.; Petras, D.; Phelan, V. V.; Bittremieux, W.; Acharya, D. D.; Ahmed, M. M.; Bauermeister, A.; Bertin, M. J.; Boudreau, P. D., 2022. A universal language for finding mass spectrometry data patterns. *bioRxiv*. 2022.08. 06.503000.

* Presenting author. E-mail address: sarah.szwarc@universite-paris-saclay.fr (Sarah Szwarc)

S3.P34 Non-targeted profiling analysis by GC/MS and LC/MS/MS of extracts derived from *Eucalyptus tereticornis* with antiobesogenic and antidiabetic activity

Norman Balcazar^{a,b,*}, Jennifer Quinchia^c, Luis Alberto Gonzalez^d

^a GENMOL Group

^b Department of Physiology and Biochemistry, Faculty of Medicine

^c Max Planck Tandem Group in Nanobioengineering, University of Antioquia. Complejo Ruta N, Calle 67 N° 52–20, Medellín, 050010, Colombia

^d Organic chemistry of natural products group, Faculty of Natural and Exact Sciences, Universidad de Antioquia, Calle 70, N° 52-21, A.A. 1226, Medellin, Colombia

Previous studies have shown that triterpene-enriched natural extracts (OBE100), ursolic acid, oleanolic acid, and ursolic acid lactone obtained from *Eucalyptus tereticornis* Sm (Myrtaceae) leaves, reduces metabolic alterations in a diet-induced obese mouse model (1). The present work aims to chemically characterize the additional molecules in the OBE100 to obtain a standardized extract.

The chromatography conditions for HPLC measurements were using a column for polycyclic aromatic hydrocarbons (PAHs, Agilent) of 5µm pore and readings at 210nm wavelength on diode array detector.

Gas chromatography (GC) and Liquid chromatography (LC) coupled with mass spectrometry (MS) were utilized to identify constituents present in OBE100. The GC–MS analysis involved a derivatization step using bis(trimethylsilyl)trifluoroacetamide (BSTFA) for polar compounds prior to MS analysis. The full Scan method was applied to screen and identify small molecular metabolites (<850 Daltons) by comparing spectral data with the NIST 14 Mass Spectral Library. GC–MS analysis was developed using a mass spectrometer (Agilent 7890/MSD 5975C) equipped with a capillary column (HP-5MS Agilent). The results revealed the presence of over 140 small acids, alcohols, hydroxy acids, sugars, fatty acids, sterols, and terpenoids. The LC–(ESI+)–MS/MS was developed using an HPLC-QTOF (Bruker Impact II). A tentative identification of principal constituents was achieved by analyzing the MS/MS data and exact mass. Compounds identified include germacrene, glycyrrhetaldehyde, loxanic acid, dehydrousolic acid lactone, sirenin, ophiopogonone B, 1,3,6- trihydroxy-8-(3-methylbutyl) anthraquinone, coumafuryl, 2-O-E-p-coumaroyl alphetolic acid, ursolic acid, ursolic acid lactone, and oleanolic acid. These compounds have also been reported previously in other *Eucalyptus* species.

Keywords: *Eucalyptus tereticornis*, antiobesogenic, antidiabetic, chemical profiling analysis

References

- (1) Acin, S., Muñoz, DL., Guillen, A., Soscue, D., Castaño, A., Echeverri, F., Balcazar, N. 2021. Triterpene-enriched fractions from *Eucalyptus tereticornis* ameliorate metabolic alterations in a mouse model of diet-induced obesity. *J. Ethnopharmacol.* 265: 113298.

* Presenting author. E-mail address: norman.balcazar@udea.edu.co (Norman Balcazar)


S3.P35 Design, synthesis and screening of novel prenylated chalcones as optimised anti-cancer agents

Mouna Hind Laiche^{a,*}, Brona Murphy^b, James W. Barlow^a

^a Department of Chemistry, Royal College of Surgeons in Ireland, 123 St Stephen's Green, Dublin, Ireland

^b Department of Physiology and Medical Physics, Royal College of Surgeons in Ireland, 123 St Stephen's Green, Dublin, Ireland

The anti-cancer potential of simple chalcones has been an intense area of research for some years, which has allowed structure-activity relationships to be well defined. Although many chalcones display cytotoxic properties, important predictors of activity are the presence and position of hydroxy and methoxy groups. An important example is the liquorice metabolite, licochalcone A (LA), a prenylated chalcone. LA has been shown to have interesting cytotoxic activity against glioma (U87) cells, at concentrations of $>5 \mu\text{M}$. LA and related prenylated chalcones thus represent potentially valuable lead compounds. To prepare novel chalcones derived from LA, we reacted O-prenylated aldehydes with various acetophenone derivatives, substituted at positions 3' and 4' of the phenyl ring. We also replaced the phenyl ring with heteroaromatic systems. Rearrangement of the novel O-prenylated chalcones via 3,3-sigmatropic reaction resulted in the novel licochalcone A analogues (**Scheme 1**).



Scheme 1. [3,3]-sigmatropic rearrangement reaction.

This poster presents the synthetic methods developed to prepare the aldehyde and ketone precursors of a family of LA and prenylated chalcone analogues, and to describe the reaction conditions necessary for the condensation of these precursors. Requisite prenylated aldehydes were condensed with either monocyclic acetophenone derivatives or various heterocycles (including chromans and chromanones), with a key subsequent reaction being Claisen rearrangement of the substituted alkenyls. The resultant products represent novel heteroaryl systems, some of which display interesting fluorescence properties. We further show the results of cytotoxicity testing of the compounds on brain cancer cell lines, including the U87 cell line, revealing initial structure-activity relationships.

Keywords: prenylated chalcones, licochalcone, condensation, cytotoxicity, glioblastoma

References

- Jeon, Jae-Ho, Mi Ran Kim, and Jong-Gab Jun. "Concise synthesis of licochalcone A through water-accelerated [3, 3]- sigmatropic rearrangement of an aryl prenyl ether." *Synthesis* 2011.03 (2011): 370-376.
- Lu, Wan Jung, et al. "Licochalcone A attenuates glioma cell growth in vitro and in vivo through cell cycle arrest." *Food & function* 9.8 (2018): 4500-4507.
- Huang, Chien-Feng, et al. "Licochalcone A inhibits the invasive potential of human glioma cells by targeting the MEK/ERK and ADAM9 signaling pathways." *Food & function* 9.12 (2018): 6196-6204.

* Presenting author. E-mail address: mounahindlaiche22@rcsi.ie (Mouna Hind Laiche)

S3.P36 Does the combined effect of natural abietane diterpenes with temozolomide have therapeutic potential against glioblastoma?

Mariana Magalhães^{a,b,c}, Eva María Domínguez-Martín^{d,e}, Mário Pedro Marques^{b,c,*}, Ana Cristina Gonçalves^{b,c,f}, Thomas Efferth^g, Patrícia Rijo^{d,h}, Célia Cabral^{b,c,i}

^a University of Coimbra, III/UC - PDBEB & CNC/UC, Coimbra, Portugal

^b University of Coimbra, iCBR/CACC, FMUC, Coimbra, Portugal

^c University of Coimbra, CIBB/UC, Coimbra, Portugal

^d Universidade Lusófona, CBIOS, Lisbon, Portugal

^e Universidad de Alcalá de Henares, Facultad de Farmacia, Madrid, Spain

^f University of Coimbra, iCBR/CIMAGO/FMUC, Coimbra, Portugal

^g Johannes Gutenberg University, IPBS, Mainz, Germany

^h University of Lisbon, iMED.UL/FFUL, Lisbon, Portugal

ⁱ University of Coimbra, CEF, Coimbra, Portugal

Glioblastoma (GB) is the most aggressive and malignant glioma of the Central Nervous System. GB standard of care involves surgical resection followed by radiotherapy and chemotherapy (Temozolomide-TMZ). However, these treatments are often ineffective. Despite intensive efforts, less than 10% of patients survive beyond 5 years, due to late diagnosis, poor prognosis, and lack of effective therapies. Concerning this, there is a growing interest in exploring new molecules isolated from natural sources.

In this work, we investigated the potential of combined therapy using two abietane diterpenes with TMZ in U87 cells. We isolated the abietane diterpenes 7 α -acetoxy-6 β -hydroxyroyleanone (Roy) and Parvifloron D (ParvD) from the acetonic extract of *Plectranthus hadiensis* Schweinf. and *Plectranthus ecklonii* Benth., respectively. After, we assessed the cytotoxic/antiproliferative activity of co-administration of Roy (16-64 μ M) or ParvD (4-14 μ M) with TMZ (100-500 μ M) using Alamar Blue[®] assay. Cell death and cell cycle regulation was analyzed by flow cytometry. This work evidenced that treatment with these natural compounds alone is comparable with those observed in cells treated with the co-administration approach, with strong synergistic effects only emerging at higher concentrations of TMZ. Also, Roy and ParvD induced apoptosis and cell cycle arrest at S/G₂M phase, which was also observed after a co-administration at high concentrations of TMZ. Surprisingly, our results unveil that both natural compounds could be used in monotherapy, showing no advantages at co-administration with TMZ. Roy and ParvD emerge as promising drug leads for future therapeutic strategies against GB, rather than only being adjuvants to the existing chemotherapy.

Keywords: glioblastoma, *Plectranthus* L'Hér., abietane diterpenes, temozolomide, antitumor effect

Acknowledgements

This work was funded by FCT, Portugal, through the following projects: UIDB/04539/2020 - DOI: 10.54499/UIDB/04539/2020, UIDP/04539/2020 - DOI: 10.54499/UIDP/04539/2020, LA/P/0058/2020 - DOI: 10.54499/UIDP/04539/2020, UIDB/04567/2020 and UIDP/04567/2020. MM is supported by a Foundation for Science and Technology (FCT) PhD grant (Reference: SFRH/BD/146441/2019). MPM is supported by a Foundation for Science and Technology (FCT) PhD grant (Reference: PRT/BD/153391/2021). These PhD grants are financed by national budget and co-financed through the European Social Fund (ESF) and the Regional Operational Por_Centro. EMD-M is supported by a predoctoral FPU-UAH 2019 fellowship from University of Alcalá de Henares.

*Presenting author. E-mail address: mpedromarques1995@gmail.com (Mário Pedro Marques)

S3.P37 Assay-directed method development for natural product purification

Hui-Hui Xiao^a, Jack Silver^{b,*}

^a State Key Laboratory of Chinese Medicine and Molecular Pharmacology (Incubation), The Hong Kong Polytechnic University Shenzhen Research Institute, Shenzhen 518057, PR China

^b Teledyne LABS, 4700 Superior Street, Lincoln, NE 68504

Natural products are an important source of lead compounds in drug discovery research, presenting such unique challenges to drug discovery as screening, purification, and identification. Purification can be streamlined using Assay-Directed Method Development, where small extractions and scouting gradients are used to determine an efficient purification strategy.

Extraction and dissolution solvents are used to select initial column media and solvents in liquid chromatography. The retention times of active fractions collected from scouting gradients are used to calculate efficient focused gradients using tools such as the Time-on-Target algorithm. Scouting gradient active-fraction retention times are used for assay-directed method development for liquid chromatography and supercritical fluid chromatography. Assay-directed method development allows rapid purification of active compounds from extracts by providing data that enables decision-making about extraction solvents, column and mobile phase selection, and efficient chromatography gradient methods.

Keywords: chromatography, purification, method development, extraction, *assay*

References

Silver, J. Overview of Analytical-to-Preparative Liquid Chromatography Method Development. *ACS Comb. Sci.* 2019, 21, 9, 609–613.

* Presenting author. E-mail: jack.silver@teledyne.com

S3.P38 Comparative analysis of the anti-inflammatory properties of nanoparticles derived from the medicinal plants *Zingiber officinale* (Ginger) and *Caralluma edulis* (Chunga), in the treatment of Inflammatory Bowel Disease

Abayomi Yusuf^{a,*}, Sarah O'Neill^a, Zeibun Ramtoola^a, James W. Barlow^b

^a School of Pharmacy and Biomolecular Sciences, Royal College of Surgeons in Ireland, Dublin, Ireland

^b Department of Chemistry, Royal College of Surgeons in Ireland, Dublin, Ireland

Inflammatory Bowel Disease (IBD) is a chronic inflammatory condition affecting the gastrointestinal tract of patients (Baumgart and Carding, 2007). Diverse approaches have been proposed to develop novel treatments for IBD, including targeted delivery of therapeutics using nanomedicine and/or nanoparticles. Recently, plant-derived exosome-like nanoparticles (PENs), a group of multifunctional nanoparticles, have been explored for their potential as drug delivery vectors, in addition to having intrinsic biological activities (Kim et al., 2021). Ginger-derived nanoparticles (GDNPs) are one of the more investigated PENs, while Chunga nanoparticles (ChNPs) have recently been isolated by our group. The aim of this study was to evaluate and compare the anti-inflammatory properties of both PENs *in vitro*.

Both PENs were isolated using a top-down approach. An inflamed intestinal epithelial barrier of Caco-2 cells was established using a cocktail containing pro-inflammatory cytokines. Inflamed monolayers were treated with different sizes of GDNPs and ChNPs. Confocal Laser Microscopy, transepithelial electrical resistance (TEER) monitoring and cytokine quantification were used to compare the effects of these nanoparticles on the monolayer.

Nanoparticles of different average particle sizes were isolated from both parent plants. Inflamed monolayers treated with ChNPs had the highest increase in TEER value among the various nanoparticles compared. ChNP treated cells also showed a higher reduction in pro-inflammatory cytokines when compared to GDNP treated cells.

Overall, this study showed that ginger- and Chunga-derived PENs have anti-inflammatory effects on the inflamed intestinal Caco-2 epithelial barrier *in vitro*. The results of this study also suggest that ChNPs may exhibit stronger anti-inflammatory effect than GDNPs.

Keywords: *Zingiber officinale*, *Caralluma edulis*, multifunctional nanoparticles, Inflammatory Bowel Disease, intestinal epithelial barrier model

References

Baumgart, D.C. and S.R. Carding, Inflammatory bowel disease: cause and immunobiology. *The Lancet*, 2007. **369**(9573): p. 1627-1640.

Kim, J., et al., Plant-derived Exosome-like Nanoparticles and their Therapeutic Activities. *Asian Journal of Pharmaceutical Sciences*, 2021.

* Presenting author. E-mail address: abayomiyusuf21@rcsi.com (Abayomi Yusuf)

S3.P39 Isolation, characterization, and bioactivity of compounds from the stem bark of *Prunus africana*

Nambooze Jennifer*, Anke Wilhelm, Susan L. Bonnet

Department of Chemistry, University of the Free State, 9300, Bloemfontein, South Africa

Prunus africana is a forest tree that grows on hillsides and flourish in humid, semitropical, and tropical central areas (Komakech et al., 2017), and a wide range of therapeutic benefits have been reported (Ting et al., 2014). However, limited information on the bioactive compounds responsible for specific therapeutic effects is available. Thus, several antioxidant and cytotoxic compounds were isolated from the DCM and methanol extracts of the stem bark of *Prunus africana* via a bioassay- guided approach.

The crude DCM extract showed the highest cytotoxicity against MCF7, CEM, HeLa, and BJ cell lines, with IC₅₀ values of 3.5 ± 0.4 µg/mL, 17.1 ± 2.0 µg/mL, 12.9 ± 3.8 µg/mL and 23.7 ± 1.4 µg/mL, respectively. The MeOH crude extract with a total phenol content of 381 mg GAE/g dry weight and a total flavonoid content of 67 mg QE/g dry weight demonstrated the best antioxidant activity. The DPPH and FRAP antioxidant tests were used and IC₅₀ values of 26.94 µg/mL and 77.28 µg/mL were obtained, respectively. Compounds **1**, **2** and **3** were isolated from the DCM extract, and compounds **4** and **5** from the MeOH extract (Fig. 1)




Figure 1: Structures of compound **1**, **2**, **3**, **4** and **5**

Against HeLa and MCF7 cells compound **1** had the best cytotoxic activity, showing percentage cell proliferation inhibition of 4% at the test concentration of 16.7 µg/mL for the HeLa cells, and cell proliferation inhibition of 15% against MCF7 at a concentration of 50 µg/mL. Computational docking calculations confirmed these results and suggested that the extract and some isolated compounds of *P. africana* has the potential to be developed as anti-cancer preparations or drugs.

Keywords: *Prunus africana*, antioxidant, cytotoxicity, anticancer activity

References

- Komakech, R., Kang, Y., Lee, J., & Omujal, F. (2017). A Review of the Potential of Phytochemicals from *Prunus africana* (Hook f.) Kalkman Stem Bark for Chemoprevention and Chemotherapy of Prostate Cancer. 2017
- Ting, H C. Agarwal, and R. Agarwal, G. D. (2014). H. Ting, G. Deep, C. Agarwal, and R. Agarwal, "The strategies to control prostate cancer by chemoprevention approaches," *Mutation Research*, 760, 1–15.

* Presenting author. E-mail address: n.jennifer33@gmail.com (Nambooze Jennifer)

S3.P40 Lanostane triterpenoids of *Inonotus obliquus* are modulators of inflammation regulators ROR γ and GPBAR1

Benjamin Kirchweger^{a,b,*}, Patrik F. Schwarz^{a,c}, Ulrike Grienke^a, Alexander F. Perhal^a, Karmen Kapp^{a,d}, Verena M. Dirsch^a, Judith M. Rollinger^a

^a University of Vienna, Department of Pharmaceutical Sciences, Division of Pharmacognosy, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^b Ludwig-Maximilians-University Munich, Department of Pharmacy- Center for Drug Research, Pharmaceutical Biology, Butenandtstraße 5-13, 81377 Munich, Germany

^c University of Vienna, Vienna Doctoral School of Pharmaceutical, Nutritional and Sport Sciences (PhaNuSpo), Josef-Holaubek-Platz 2, 1090 Vienna, Austria

^d University of Helsinki, Faculty of Pharmacy, Division of Pharmaceutical Biosciences, Viikinkaari 5 E, 00014, Helsinki, Finland

The sclerotic conks of the birch-parasite mushroom *Inonotus obliquus* (Fr.) Pilát called chaga, have a rich history in traditional medicine across various cultures, with applications ranging from adaptogenic properties to cancer and inflammation treatment (Duru et al., 2019). In light of its historical use in traditional medicine, we included two *I. obliquus* extracts to screen for new natural modulators of GPBAR1 and ROR γ , promising targets in anti-inflammatory research (Ladurner et al., 2021, Pols, 2014).

The two extracts, differing in lipophilicity, were obtained via supercritical liquid extraction. The lipophilic extract was acquired using CO₂ with 10% EtOH, while the more hydrophilic extract was obtained with CO₂ and 20% EtOH. Our screening revealed distinct activities: the lipophilic extract inhibited ROR γ (0.6-fold at 10 μ g/mL), while the hydrophilic extract activated GPBAR1 (8.8-fold at 30 μ g/mL). To identify the different compounds contributing to these effects, we performed a dereplication with UPSFC-ESI-MS. The tentatively identified constituents were forwarded to molecular docking which indicated lanostane triterpenoids as active principles. Subsequently, we isolated seven lanostane triterpenoids along with four additional compounds and tested them for ROR γ and GPBAR1 modulation. Notably, the lanostane triterpenoids inonotsutriol A and inonotsutriol epoxide demonstrated potent and concentration-dependent inverse agonistic activity with IC₅₀s of 1.3 μ M and 1.8 μ M in a luciferase assay employing ROR γ -Gal4. Additionally, trametenolic acid B was identified as a modest GPBAR1 activator.

These findings contribute to a better mechanistic understanding of the pharmacological profile of *I. obliquus*, shedding light on its potential mechanisms of action against inflammatory diseases.

Keywords: *Inonotus obliquus*, ROR γ , GPBAR1, lanostane, supercritical fluid chromatography

References

- Duru, K.C., Kovaleva, E.G., Danilova, I.G. & van der Bijl, P. 2019. The pharmacological potential and possible molecular mechanisms of action of *Inonotus obliquus* from preclinical studies. *Phytother. Res.*, 33, 1966-1980.
- Ladurner, A., Schwarz, P.F., Dirsch, V.M. 2021. Natural products as modulators of retinoic acid receptor-related orphan receptors (RORs). *Nat. Prod. Rep.*, 38, 757-781.
- Pols, T.W.H. 2014. TGR5 in inflammation and cardiovascular disease. *Biochem. Soc. Trans.*, 42, 244-249.

* Presenting author. E-mail address: benjamin.kirchweger@cup.uni-muenchen.de (Benjamin Kirchweger)

S3.P41 COSMO-RS as cutting-edge technology for design and pre-screening of Natural Deep Eutectic Solvents for selective extraction: Spirulina as case study

Mervé Yagmur^{a,*}, Alia Chevalley^b, Bénédicte Montigny^c, Johan Jacquemin^d, Leslie Boudesocque-Delaye^a

^a EA7502 SIMBA, Faculty of Pharmacy, University of Tours, 31 avenue Monge, 37200 Tours, France

^b AquaEcoCulture, 7 rue d'Armor Maroué, 22400 Lamballe, France

^c EA 6299 PCM2E, University of Tours, Parc de Grandmont, 37200 Tours, France

^d MSN Department, Mohammed VI Polytechnic University, Lot 660, Hay Moulay Rachid, Ben Guerir 43150, Morocco

The aim of this work is to demonstrate the selective extraction of high value compounds of nutraceutical or cosmetic interest from Spirulina using an advanced methodology combining experimental and simulation approaches. The SIMBA laboratory uses NaDES (Natural Deep Eutectic Solvents) to extract these compounds. NaDES are combinations of two or three primary plant metabolites that are considered biocompatible (Dai et al., 2013). NaDES are also known for their versatile selectivity, which can be modulated by the nature or ratio of components that enter into NaDES composition.

NaDES modulation was previously achieved experimentally, resulting in time-consuming and chemical-intensive procedures. However, this work introduces a dual approach that combines solvent restriction regulation with the utilization of simulation tools, such as COSMO-RS (Conductor-like Screening Model for Real Solvents) method, as an innovative pre-screening tool to achieve this modulation. By determining *ab-initio* key descriptors such as 3D shape, charge density, and sigma profiles onto the molecule surface for each pre-selected bio-compatible compound, we were able to formulate NaDES compatible with our application *in-silico*. This approach saves time and minimizes the ecological impact of our experimental trials.

Several NaDES based on octanoic acid were selected for experimental testing to validate this screening approach. Furthermore, experimental works were done to not only validate the simulated eutectic window but also to determine if the eutectic ratio was required to achieve optimum recovery. Additionally, various experimental conditions were tested to guide process selection and define the best spirulina extraction conditions for future scale-up.

Keywords: Simulation, COSMO-RS, Green Chemistry, NaDES, extraction

References

- Wils, L., Leman-Loubière, C., Bellin, N., Clément-Larosière, B., Pinault, M., Chevalier, S., Enguehard-Gueiffier, C., Bodet, C., Boudesocque-Delaye, L., 2021. Natural deep eutectic solvent formulations for spirulina: Preparation, intensification, and skin impact. *Algal Res.*, 56, 102317.
- Dai, Y., van Spronsen, J., Witkamp, G.-J., Verpoorte, R., Choi, Y.H., 2013. Natural deep eutectic solvents as new potential media for green technology. *Analytica Chimica Acta* 766, 61–68.

* Presenting author. E-mail address: merve.yagmur@univ-tours.fr (Mervé Yagmur)

S3.P42 Tetrahydrofuran sesquiterpenes and the bronchodilator effect of *Artemisia judaica*

Ali ElGamal^{a,*} Abdullah AlDosari^a, Najeebb Rehman^b, Omer Basudan^a, Maged AbdelKader^c

^a Dept. of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, Saudi Arabi

^c Dept. of Pharmacology and Toxicology, College of Pharmacy, Sattam Bin Abdulaziz University, Al- kharj, Saudi Arabia

^c Dept. of Pharmacognosy, College of Pharmacy, Sattam Bin Abdulaziz University, Al-kharj, Saudi Arabia

Bronchodilators are employed for both acute and ongoing treatment of respiratory ailments such as asthma and chronic obstructive pulmonary disease (Barnes, 2010). The genus *Artemisia*, belonging to the family Asteraceae, is a large diverse genus with approximately 500 species. *Artemisia judaica*. is an aromatic perennial herbaceous plant. In Saudi Arabia, *A. judaica* is commonly used for treating bronchitis and bronchial asthma. Other species of *Artemisia* have been found to have beneficial effects in treating asthma and epilepsy (Sapkota, 2008). The previously reported secondary metabolites identified in *Artemisia* species are sesquiterpene lactones, flavonoids and essential oil (Abou El Hamd, et al. 2010). The objective of the present study is to provide scientific proof supporting the bronchodilator effects of *A. judaica*. isolating, identifying its active ingredients and exploring the mechanism of the bronchodilator effect. The bronchodilator effect was assessed using isolated guinea-pig tracheal tissues. Bronchoconstriction was induced by carbachol (CCh, 1 M), and the inhibitory effect of the plant extract and fractions was tested against these induced contractions. The findings demonstrated that the petroleum ether fraction had greater potency (0.5 mg/ml) in inhibiting CCh-induced contraction compared to the other fractions. Several tetrahydrofuran sesquiterpenes (**1-6**) (Fig. 1) were obtained by bioassay-guided fractionation employing various normal CC and reversed phase HPLC chromatographic techniques. The identification of the isolated compounds was achieved using spectroscopic techniques, such as HRESIMS, 1D and 2DNMR. Compounds **5-7** are reported here for the first time. An investigation was also conducted to investigate the mechanism of action of the isolated compounds. The bronchodilator effect involved the activation of multiple K⁺ channels.




Fig. 1. Structures of isolated sesquiterpene

Keywords: *Artemisia judaica*, Asteraceae, bronchodilator effect

References

- Barnes, P. J. 2010. New therapies for chronic obstructive pulmonary disease. *Med. Princ. Pract.*, 19, 330-338.
- Sapkota, P. P. 2008. Ethno-ecological Observation of Magar of Bukini, Baglung, Western, Nepal. *Dhau. Jour. of Soci. and Anthro.*, 2, 227-252.
- Abou El Hamd, H. M., El-Sayed, M. A., Hegazy, M. E., et al. 2010. Chemical constituents and biological activities of *Artemisia herba-alba*. *Rec. of Nat. Prod.*, 4, 1-25.

* Presenting author. E-mail address: aelgamel@ksu.edu.sa (Ali ElGamal)

S3.P43 Development of non-targeted metabolomic approaches for the screening of bioactive molecules in *Clematis* species

Audrey Le Cabec^{a,*}, Pierre-Eric Campos^a, Pierre-Marie Allard^b, Edouard Brüelhart^b, Emmanuel Defossez^c, Emilie Destandau^a

^a Institute of Organic and analytical chemistry (ICOA), UMR 7311, University of Orléans, France

^b Department of Biology, University of Fribourg, Fribourg, Switzerland

^c Institute of Biology, University of Neuchâtel, Switzerland

Recent advancements in analytical techniques and data treatment have led to the development of new workflows for discovering natural bioactive compounds. UHPLC-HRMS/MS crude extract profiling allows rapid and sensitive analysis of small amounts of plants, but annotation of complex mixture, even using molecular networking (MN) (Wang et al., 2016) can be time-consuming. Furthermore, increasing annotation confidence is a challenge, requiring the integration of diverse approaches, such as implementing taxonomic information through filters, scoring (Tima-R) and comparison to experimental or predicted databases (Rutz et al., 2019).

Clematis genus, third largest of Ranunculaceae, is widespread throughout the northern hemisphere, used in traditional medicines or ornamental purposes. These different uses can be explained by a wide diversity of compounds leading to different bioactivities (Chawla et al., 2012). To highlight phytochemical differences and identify potential way of valorizing the huge biomass from these crops, 10 species of *Clematis* were analysed by UHPLC-HRMS/MS. Data were represented using MN, and annotation tools as SIRIUS4 (Dührkop et al., 2019) and Tima-R were employed to refine the compound identification.

Shared clusters of O-glycosylated flavonoids, triterpenoids and lignans had been determined between the 10 species. However, specific clusters had been also identifying for different species for example acetylated O-glycoside flavonoids for *C. chinensis* and C-glycoside flavonoids for *C. viticella* which may lead to a modulation of the bioactivity of plant extract (Xiao et al., 2016), while *C. koreana*, is the only species with a chemical profile rich in alkaloids directing to a different way of valorization.

Keywords: *Clematis* genus, metabolomics, mass spectrometry, molecular network, bioactivity

References

- Chawla, R., Kumar, S., Sharma, A., 2012. The genus *Clematis* (Ranunculaceae): Chemical and pharmacological perspectives. J Ethnopharmacol 143, 116–150.
- Dührkop, K., Fleischauer, M., Ludwig, M., et al. 2019. SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure information. Nat Methods 16, 299–302.
- Rutz, A., Dounoue-Kubo, M., Ollivier, S., et al 2019. Taxonomically Informed Scoring Enhances Confidence in Natural Products Annotation. Front Plant Sci 10, 1–15.
- Wang, M., Carver, J.J., Phelan, V. V, et al. 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. Nat Biotechnol 34, 828–837.
- Xiao, J., Capanoglu, E., Jassbi, A.R., Miron, A., 2016. Advance on the Flavonoid C-glycosides and Health Benefits. Crit Rev Food Sci Nutr 56, S29–S45.

* Presenting author. E-mail address: audrey.le-cabec@univ-orleans.fr (Audrey Le Cabec)

S3.P44 Semisynthetic pyrazole ring containing hydrocurcumin derivatives as potential new antitumor agents

Gábor Girst^{a,*}, Benedek Szűcs^a, Sándor Bartha^b, Bizhar A. Tayeb^b, István Zupkó^b, Renáta Minorics^b, Attila Hunyadi^a

^a Institute of Pharmacognosy, Faculty of Pharmacy, University of Szeged, Eötvös utca 6, H-6720 Szeged, Hungary

^b Institute of Pharmacodynamics and Biopharmacy, Faculty of Pharmacy, University of Szeged, Eötvös utca 6, H-6720 Szeged, Hungary

Hydrocurcumins occur in the rhizomes and leaves of *Curcuma longa*, as well as some other species of the Zingiberaceae family. They are reduced metabolites of the deeply studied diarylheptanoid curcumin, the main bioactive compound of the well-known eastern spice and traditional medicine curcuma. Despite the promising effect of curcumin, it has not become a modern medicine due to its very poor pharmacokinetic properties. Hydrocurcumins possess much better bioavailability, and in many cases have similar bioactivities, so they have a higher drug discovery potential (Girst et al., 2021).

Tetrahydrocurcumin (4HC) inhibits tumour metastasis and tumour angiogenesis in mouse models. Synthetic modification of the diketo moiety of 4HC has led to pyrazole containing antiproliferative agents (Mahal et al., 2017).




Fig. 1. General structure of pyrazole-containing hydrocurcumins

In this study, our aim was to synthesise new, more potent nitrogen-containing hydrocurcumin derivatives, as well as the most potent one from Mahal's work, and investigate *in vitro* their cytotoxic activities in different cancer cell lines (HeLa, A2780 and MCF-7).

Synthetic reactions were followed by chromatographic purifications (flash and HPLC) and the structures were confirmed by one- and two-dimensional NMR and HRMS measurements. Starting from curcumin, in this project 9 nitrogen-containing hydrocurcumin derivatives were synthesised, among which 6 are newly described. The A2780 ovarian cell line was found to be the most sensitive to them, with the best compounds having IC₅₀ values of 11.2 and 11.4 μM.

Keywords: curcumin, hydrocurcumins, semisynthesis, pyrazole, ovarian cancer

References

- Girst, G., Ötvös, S.B., Fülöp, F., Balogh, G.T., Hunyadi, A., 2021. Pharmacokinetics-Driven Evaluation of the Antioxidant Activity of Curcuminoids and Their Major Reduced Metabolites—A Medicinal Chemistry Approach. *Molecules*. 26, 3542.
- Mahal A., Wu, O., Jiang Z.-H., Wei, X., 2017. Synthesis and Cytotoxic Activity of Novel Tetrahydrocurcumin Derivatives Bearing Pyrazole Moiety. *Nat. Prod. Bioprospect.* 7, 461-469.

Acknowledgements

Grants NKFI K-134704 and TKP2021-EGA-32

* Presenting author. E-mail address: girst.gabor.02@szte.hu (Gábor Girst)

S3.P46 A decision theory-based aggregation tool for tandem mass spectrometry multi-annotation outputs

Yassine M. Mejri^{a,b,*}, Meltem Öztürk^b, Olivier Cailloux^b, Mehdi A. Beniddir^a

^a Université Paris-Saclay Bâtiment Henri Moissan, CNRS, BioCIS, 17 Av. des Sciences, 91400 Orsay, France

^b Université Paris-Dauphine, PSL Research University, CNRS, LAMSADE, 75016 Paris, France

The accurate annotation of molecular structures in metabolomics is crucial across diverse chemistry fields, marked by explosive growth and ongoing improvements. Predictions of MS/MS spectra and chemical class using in silico approaches have emerged as powerful complementary tools for annotation. Breakthrough strategies include molecular networking (GNPS (Nothias et al., 2020)), fragmentation trees (SIRIUS (Dührkop et al., 2019)), artificial neural networks (ISDB (Wang et al., 2021)), and chemical ontology. However, the increasing number of outputs generated by these tools poses challenges in prioritization for natural product chemists. Thus, there's an urgent need for a tool that processes outputs from common annotation tools (e.g., GNPS, SIRIUS, ISDB) to compute a "knownness" score. This score would aid chemists in targeting new compounds effectively. This communication presents the development of a decision-support tool for the in-depth analysis of multi-annotation of tandem mass spectrometry data in natural product chemistry. We will utilize decision theory modeling and a decision-making approach to design a function that translates what the annotation tools say about the similarity of an MS/MS spectrum to known molecules. With this aggregated knownness score, natural product chemists can evaluate if a molecule warrants isolation and further structure elucidation efforts. Real-case scenarios, inspired by previously published phytochemical reports, will be further assessed from the angles of decision theory and targeted isolation.

Keywords: Annotation tools, multi-informative data, decision theory, aggregation

References

- Nothias, L. F., Petras, D., Schmid, R., Dührkop, K., Rainer, J., Sarvepalli, A., ... & Dorrestein, P. C. 2020. Feature-based molecular networking in the GNPS analysis environment. *Nat. Methods*. 17, 905-908.
- Dührkop, K., Fleischauer, M., Ludwig, M., Aksenov, A.A., Melnik, A.V., Meusel, M., Dorrestein, P.C., Rousu, J. and Böcker, S., 2019. SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure information. *Nat. Methods*. 16, 299-302.
- Wang, F., Liigand, J., Tian, S., Arndt, D., Greiner, R., & Wishart, D. S. 2021. CFM-ID 4.0: more accurate ESI-MS/MS spectral prediction and compound identification. *Anal. Chem*. 93, 11692-11700.

* Presenting author. E-mail address: Mohamed-yassine.mejri@dauphine.eu (Yassine M. Mejri)

S3.P47 *Plectranthus*: an important source of bioactive diterpenoids for therapeutic applications

Vera M. S. Isca^{a,b}, Gabrielle Bangay^{a,b}, Marcia Filipe^{a,b}, Rebeca André^{a,b}, Ahmed A. Hussein^c, Patricia Rijo^{a,b,*}

^a Center for Research in Biosciences & Health Technologies (CBIOS), Universidade Lusófona, Lisboa, Portugal

^b Instituto de Investigação do Medicamento (iMed.Ulisboa), Faculdade de Farmácia, Universidade de Lisboa, Portugal;

^c Chemistry Department, Cape Peninsula University of Technology, Symphony Rd., Bellville 7535, South Africa

Plectranthus spp., commonly used in traditional medicine, offer a rich source of bioactive abietane diterpenoids, with diverse biological activities supported by recent scientific research. (Matias, 2016). Several approaches utilizing bioactive lead molecules from *Plectranthus* spp. for therapeutic applications are reported here. Examples include Parvifloron D (ParvD) from *P. ecklonii* and 7 α -acetoxy-6 β -hydroxyroyleanone (Roy) from *P. grandidentatus*. One notable instance involves the patented diterpenoid dibenzoylroyleanone (RoyBz; Bessa, 2016), derived from natural Roy. RoyBz has been identified as a PKC-selective activator, exhibiting potent anti-proliferative effects in colon cancer by triggering a PKC-dependent mitochondrial apoptotic pathway. Remarkably, the PKC δ -dependent anticancer activity of RoyBz was confirmed in vivo using xenograft mouse models of both control and PKC δ -knockdown human colon cancer cells (Bessa, 2018). Further investigation delves into the molecular mechanism underlying Roy-Bz's reported antitumor activity, particularly its impact on glucose metabolism in colon cancer cells (Bessa, 2023). This study sheds light on the role of PKC in tumor cell metabolism and underscores Roy-Bz's potential to target ATP-producing pathways of glucose metabolism in cancer cells, bolstering its candidacy as an anticancer agent.

Keywords: *Plectranthus*, Lamiaceae, diterpenoids, cancer, PKC

References

- D. Matias, C. Bessa, M.F. Simões, C.P. Reis, L. Saraiva, P. Rijo*, Natural Products as Lead Protein Kinase C Modulators for Cancer Therapy, in: Atta-ur-Rahman (Ed.), Studies in Natural Products Chemistry, 2016, 45–79
- Bessa C., Soares J., Raimundo L., Simões M.F., Gonçalves J., Rijo P., Saraiva L.. International Patent nº 109140: “Roy- Bz: A small molecule selective activator of Protein Kinase Cd”, PCT/IB2017/050633, 2016.
- Bessa C., Soares J., Raimundo L., Loureiro J. B., Gomes C, Reis F, Soares M., Santos D, Dureja C., Chaudhuri S.R., Lopez-Haber C., Kazanietz M.G., Gonçalves J., Simões M.F., Rijo P., Saraiva L. (2018). Cell death & disease. 18;9(2):23. Bessa C., Loureiro J.B., Barros M., Isca V.M.S., Sardão V.A., Oliveira P.J., Bernardino R.L., Herman-de-Sousa C., Costa M.A., Correia-de-Sá P., et al. 2023. “Counteracting Colon Cancer by Inhibiting Mitochondrial Respiration and Glycolysis with a Selective PKC δ Activator” International Journal of Molecular Sciences 24, no. 6: 5710.

* Presenting author. E-mail address: patricia.rijo@ulusofona.pt (Patricia Rijo)

S3.P48 Characterization of volatile and non-volatile compounds in *Larix decidua* (Pinaceae) oleoresin and its nanoemulsion development

João V.C. Batista^{a,b,*}, Anna Bourguignon^c, Bettina Leonhard^b, Jakob Maier^b, Aline Carvalho^d, Rafael Garrett^d, Fabio Boylan^c, Carla Holandino^{b,e}, Stephan Baumgartner^{b,f,g}, Jörg Huwyler^a

^a Department of Pharmaceutical Sciences, Division of Pharmaceutical Technology, University of Basel, Klingelbergstrasse 50, 4056 Basel, Switzerland

^b Society for Cancer Research, Hiscia Institute, Kirschweg 9, 4144 Arlesheim, Switzerland

^c School of Pharmacy and Pharmaceutical Sciences, Panoz Institute, Trinity Biomedical Sciences Institute, Trinity Natural Products Research Centre, Trinity College Dublin, D02 PN40, Dublin, Ireland

^d Metabolomics Laboratory, Chemistry Institute, Universidade Federal do Rio de Janeiro, Av. Horácio Macedo 1281, Ilha do Fundão, 21941-598 Rio de Janeiro, Brazil

^e Multidisciplinary Laboratory of Pharmaceutical Sciences, Faculty of Pharmacy, Universidade Federal do Rio de Janeiro, Av. Carlos Chagas Filho 373, 21941-902 Rio de Janeiro, Brazil

^f Faculty of Health, Institute of Integrative Medicine, University of Witten/Herdecke, Gerhard-Kienle-Weg 4, 58313 Herdecke, Germany

^g Faculty of Medicine, Institute of Complementary and Integrative Medicine, University of Bern, Freiburgstrasse 40, 3010 Bern, Switzerland

Chronic wounds continue to present a significant medical challenge (Falanga et al., 2022). Among natural products, *Larix decidua*'s oleoresin has long been recognized for its wound-healing properties and has traditionally been incorporated into ointment formulations (Batista et al., 2022). However, as modern pharmaceutical formulations increasingly favour emulsions for encapsulating lipophilic compounds, there is a growing interest in utilizing this technology to deliver the benefits of oleoresins. Thus, this study aimed to characterize the phytochemical profile of volatile and non-volatile compounds in the European Larch oleoresin, and to develop a nanoemulsion containing it as an innovative approach to delivering it.

Gas chromatography-mass spectrometry analysis revealed the volatile and non-volatile chemical composition of the oleoresin (Fig. 1). The monoterpenes alpha-pinene (**1**; >70%), beta-pinene (**2**; >8%), and limonene (**3**; >5%) were the major volatile compounds. The labdane-type diterpenes larixyl acetate (**4**; >23%) and larixol (**5**; >5%), and the pimarane diterpene isopimaric acid (**6**; >17%) were identified as the major non-volatile compounds.




Figure 1. Structure of the main monoterpenes and diterpenes in the *Larix decidua* oleoresin.

A nanoemulsion comprising 10% oleoresin was developed. Two sterilization methods, membrane filtration and X-ray exposure, were employed, and the resulting nanoemulsions were evaluated over 14 weeks. The formulation subjected to sterile membrane filtration without X-ray exposure demonstrated optimal characteristics for further investigation. Throughout the evaluation period, the droplet size (~200 nm), PDI (<0.2), pH (~4.2), and the spherical morphology of the droplets were observed. The developed nanoemulsion effectively encapsulates the oleoresin from *L. decidua*, demonstrating potential for treating chronic wounds.

Keywords: *Larix decidua*, nanoemulsion, diterpenes, monoterpenes, GC-MS

References

- Batista, J.V.C., Uecker, A., Holandino, C., Boylan, F., Maier, J., Huwyler, J., et al., 2022. A Scoping Review on the Therapeutic Potential of Resin From the Species *Larix decidua* Mill. [Pinaceae] to Treat Ulcerating Wounds. *Front. Pharmacol.* 13, 895838. doi: 10.3389/fphar.2022.895838.
- Falanga, V., Isseroff, R.R., Soulika, A.M., Romanelli, M., Margolis, D., Kapp, S., et al., 2022. Chronic wounds. *Nat. Rev. Dis. Primers* 8(1), 50. doi: 10.1038/s41572-022-00377-3.

* Presenting author. E-mail address: joao.dacostabatista@unibas.ch (João Vitor da Costa Batista)

S3.P49 *In vitro* antihypertensive effects of *Viscum album* mother tinctures in kidney proximal tubule cells

Rodrigo dos Santos Pinto Duarte^a, Michelle Nonato de Oliveira Melo^a, João V.C. Batista^b, Giovanna Gomes Martins^a, Adriana Passos Oliveira^a, Rosilane Taveira da Silva^c, Marcelo Einicker Lamas^c, Stephan Baumgartner^b, Carla Holandino^{a,b,*}

^a Multidisciplinary Laboratory of Pharmaceutical Sciences, Faculty of Pharmacy, Universidade Federal do Rio de Janeiro, Av. Carlos Chagas Filho 373, 21941-902 Rio de Janeiro, Brazil

^b Society for Cancer Research, Hiscia Institute, Kirschweg 9, 4144 Arlesheim, Switzerland

^c Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

Alcoholic extracts (mother tinctures) from *Viscum album* (Santalaceae), a semiparasitic plant, have a centenary use in folk medicine to treat hypertension and cardiovascular diseases (Melo et al., 2023). However, the biological mechanisms related to this hypotensive property are fewer investigated. In this study, we evaluated the effects of *V. album* mother tincture (VAMT) in the Na⁺/K⁺ ATPase using the porcine proximal tubule kidney cells (LLC-PK1). The VAMTs were prepared using fresh European mistletoe from 3 different subspecies (*album*, *abietis*, *austriacum*), harvested in Summer and Winter, from five host trees (*M. domestica*, *Q. petrea*, *U. carpinifolia*, *P. sylvestris*, *A. alba*) (Holandino et al., 2020). The MTT assay indicated the cytotoxic effect triggered only by Summer VAMT from *Q. petrea* (p<0.05), after 24 h of incubation. Using the same incubation time, Summer VAMT from *A. alba* induced a significant decrease in the Na⁺/K⁺ ATPase of LLC-PK1 cells (p<0.05), highlighting the promising antihypertensive potential of this mother tincture. These interesting results support the traditional use of VAMT and suggesting, for the first time, one of its mechanism of action in hypertension. Besides, the influence of seasons and the host trees in LLC-PK1 points out the importance to investigate the biological influences regarding the mistletoe's origins. The effects in Na⁺ pump expression are under investigation in order to evaluate if changes in Na⁺/K⁺ ATPase activity and expression are responsible for the hypotensive effects of VAMT on kidney cells.

Keywords: *Viscum album* mother tinctures, kidney enzymes, *in vitro* research

References:

- Melo, M.N.d.O., Batista, J.V.d.C., Peñaloza, E.M.C., Oliveira, A.P., Garrett, R., Baumgartner, S., Holandino, C., 2023. A Scoping Review of Genus *Viscum*: Biological and Chemical Aspects of Alcoholic Extracts. *Plants*, 12, 1-35.
- Holandino, C., de Melo, M.N.d.O., Oliveira, A.P., da Batista, J.V.C., Capella, M.A.M., Garrett, R., Grazi, M., Ramm, H., Torre, C.D., Schaller, G., Urech, K., Weissenstein, U., Baumgartner, S., 2020. Phytochemical Analysis and *in Vitro* Anti- Proliferative Activity of *Viscum album* Ethanolic Extracts. *BMC Complement. Med. Ther.* 20, 1-11.

* Presenting author. E-mail address: cholandino@gmail.com (Carla Holandino)

S3.P50 ^1H NMR-based metabolomics on 30 east asian traditional herbal medicines for quality control

Eunah Jeong, Huong T. Pham, Kyo Bin Kang*

College of Pharmacy and Research Institute of Pharmaceutical Sciences, Sookmyung Women's University, 100 Cheongpa-ro 47 gil, Yongsan-gu, Seoul 04310, Republic of Korea

East Asian traditional medicine typically consists of multi-herbal formulations consisting of up to 20 different herbs, each of which contributes to synergistic effects through complex interactions between their bioactive components. However, they possess chemical complexity issues, making it necessary to develop appropriate analytical methods for their identification and standardization (Jianping Zhao et al., 2022). Herein, we performed a comparative nuclear magnetic resonance (NMR)-based metabolomics analysis to consider applicability of the method for quality control of multi-herbal formulations. A total of 86 samples composed of 30 formulation samples were analyzed by ^1H NMR spectroscopy, and chemical fingerprints were compared to each other in batch-to-batch and formulation-to-formulation manners. We could identify specialized metabolites with high content along with polar primary metabolites such as amino acids, sugars and some phenolic compounds. ^1H NMR offered comprehensive insights on the multiple formulations and has the potential to be applied for quality control.

Keywords: ^1H NMR spectroscopy, traditional herbal medicines, quality control, metabolomics

References

Jianping Z, Mei W, Seethapathy G. S, Ikhlas A. K. NMR technique and methodology in botanical health product analysis and quality control. *Journal of Pharmaceutical and Biomedical Analysis*. Volume 207.2022.114376.

* Corresponding author. E-mail address: kbkang@sookmyung.ac.kr (Kyo Bin Kang)

S3.P51 Unveiling terpenoid diversity in *Sideritis* spp. through molecular networks

Ekaterina-Michaela Tomou^{a,b,*}, Helen Skaltsa^a, Corinna Urmann^{b,c,*}

^a Section of Pharmacognosy & Chemistry of Natural Products, Department of Pharmacy, School of Health Sciences, National and Kapodistrian University of Athens, Panepistimiopolis, Zografou, 15771 Athens, Greece

^b Organic-analytical Chemistry, Weihenstephan-Triesdorf University of Applied Sciences, 94315 Straubing, Germany

^c TUM Campus Straubing for Biotechnology and Sustainability, Technical University of Munich, 94315 Straubing, Germany

The genus *Sideritis* L. (Lamiaceae family) is traditionally used for the treatment of various ailments. These plants are phytochemically rich and represent an inexhaustible source of natural products (e.g. flavonoids, phenylethanoid glycosides, and others) (González-Burgos et al., 2011). However, the terpenoids profile of *Sideritis* species has not yet been fully determined, and due to the complexity of these compounds, their characterization remains challenging.

As part of our ongoing research on *Sideritis* species (Tomou et al., 2023; Tomou et al., 2024), the main aim of the present study was to use the molecular network approach to investigate the terpenoid constituents of several *Sideritis* taxa (belonging to the *Empedoclia* section) and to highlight the relevance of molecular networks in exploring the chemotaxonomy of these plants. Therefore, GC/MS and the GNPS platform were used to analyze several different dichloromethane extracts.



Fig. 1. Molecular network cluster showing diterpenoids of the dichloromethane extracts of different *Sideritis* species.

The generated molecular network shows differences among the investigated extracts. The characteristic *ent*-kaurene type diterpenoids (such as siderol) of these plants were identified (Fig.1) and our results revealed clusters common to all samples, as well as taxa-specific clusters. This work demonstrates the ability of *in silico* tools such as molecular networks to profile the terpenoids of *Sideritis* plants and to determine possible characteristic markers relevant to chemotaxonomy.

Keywords: *Sideritis* spp., Lamiaceae, terpenoids, GC-MS, molecular networks, GNPS

References

- González-Burgos, E., Carretero, M.E., Gómez-Serranillos, M.P., 2011. *Sideritis* spp.: Uses, chemical composition and pharmacological activities-A review. J. Ethnopharmacol. 135, 209-225.
- Tomou, E-M., Bieler, L., Spöttl, T., Couillard-Despres, S., Skaltsa, H., Urmann, C., 2023. Metabolic fingerprinting of different *Sideritis* taxa infusions and their neurogenic activity. Planta Med. 89, 1087-1096.
- Tomou, E-M., Goula, K., Skaltsa, H., Urmann, C., 2024. Comparative phytochemical analysis of cultivated and wild *Sideritis raeseri* Boiss. & Heldr. subsp. *raeseri* infusions. Nat. Prod. Res. 1-6.

* Presenting author. E-mail address: corinna.urmann@hswt.de (Dr. Corinna Urmann)

S3.P52 Targeted discovery of sesquiterpene indole alkaloids from Gabonese *Greenwayodendron suaveolens*

Rany Berty Mbeng Obame^{a,*}, Pierre Le Pogam^a, Jean-François Gallard^b, Elvis N’Nang Otogo^c, Véronique Fontaine^d, Guillaume Bernadat^a, Mehdi A Beniddir^a

^a Université Paris-Saclay, CNRS, BioCIS, 91400 Orsay, France

^b CNRS, Université Paris-Saclay UPR 2301, Institut de chimie des substances naturelles, ICSN, 1 avenue de la Terrasse, 91198 Gif-sur-Yvette, France

^c Medical research centre of Lambaréné, CERMEL, Biomedical et social sciences, Lambaréné, Gabon

^d Microbiology, Bioorganic and Macromolecular Chemistry Unit, Université Libre de Bruxelles (ULB), Faculty of pharmacy, Boulevard du Triomphe, 1050 Bruxelles, Belgium

Annonaceous plants are of interest to natural product chemists due to their therapeutic value and to their isoquinoline alkaloids content. A few years ago, a MS/MS database of 322 isoquinolines and other annonaceous metabolites had been implemented to the GNPS repositories (*viz.* IQAMDB) (Agnès et al., 2022). We herein report on the dereplication of known alkaloids from stem barks of *Greenwayodendron suaveolens* (Engl. & Diels) Verdc. leveraging IQAMDB-informed feature-based molecular networking further refined by *in silico* annotation and taxonomic weighting based on Tima- R (Rutz et al., 2019). This strategy annotated over 30 compounds and streamlined the isolation of diverse sesquiterpene indole alkaloids (SIA) (**1-3**) (Fig. 1). Structure elucidation and absolute configuration assignment (by TDDFT-ECD and X-ray), determined these compounds to be greenwaylactam A (**1**), a recently reported Witkop-Winterfeldt oxidized SIA (Kemgni et al., 2021), and a newly reported diastereoisomer, namely greenwaylactam D (**2**). *N,O*-diacetylpolyveoline (**3**), is here first reported as a natural product (Kouam et al., 2014). The antibacterial activity of these isolates was assayed against *Staphylococcus aureus*, *S. epidermidis* and *Pseudomonas aeruginosa*.




Fig. 1. Chemical structure of sesquiterpene indole alkaloids isolated from *G. suaveolens*

Keywords: *Greenwayodendron suaveolens*, Annonaceae, sesquiterpene indole alkaloid, molecular networking

References

- Agnès, S.A., Okpekon, T., Kouadio, Y.A., Jagora, A., Bréard, D., et al. 2022. Implementation of a MS/MS database for isoquinoline alkaloids and other annonaceous metabolites. *Sci. Data.* 9, 270.
- Rutz, A., Dounoue-Kubo, M., Ollivier, S., Bisson, J., Bagheri, M., Saesong, T., Allard, P. M. 2019. Taxonomically informed scoring enhances confidence in natural products annotation. *Front. Plant Sci.* 10, 1329.
- Kemgni, M.F., Chenda, L.B.N., Tchamgoue, J., Kenfack, P.T., Ngandjui, Y.A.T., Wouamba, S.C., Kouam, S.F. 2021. Greenwaylactams A, B and C, the First Group of Sesquiterpene Alkaloids with an Eight-Membered Lactam Ring from *Greenwayodendron oliveri*. *ChemistrySelect.* 6, 1705-1709.
- Kouam, S.F., Ngouonpe, A.W., Lamshöft, M., Talontsi, F.M., Bauer, J.O., Strohmman, C., Spiteller, M. 2014. Indolosesquiterpene alkaloids from the Cameroonian medicinal plant *Polyalthia oliveri* (Annonaceae). *Phytochemistry.* 105, 52-59.

* Presenting author. E-mail address: rany.mbeng-obame@universite-paris-saclay.fr (Rany Berty Mbeng Obame)

S3.P53 The benefit of classification machine learning in virtual drug discovery for beta-lactamase inhibitors identification based on an in-house library

Thanet Pitakbut^{a,*}, Jennifer Munkert^a, Wenhui Xi^b, Yanjie Wei^b, Gregor Fuhrmann^a

^a Friedrich-Alexander-Universität Erlangen-Nürnberg, Department of Biology, Pharmaceutical Biology, Staudtstr. 5, 91058 Erlangen, Germany

^b Shenzhen Key Laboratory of Intelligent Bioinformatics and Center for High Performance Computing, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055, China

Machine learning (ML) is a cutting-edge computational technology that solves complex problems. Many studies utilized ML to improve virtual drug screening (Chan et al., 2023). However, few studies focused on identifying beta-lactamase inhibitors (Anant et al., 2022 and Papastergiou et al., 2022), even though beta-lactamase is the leading cause of antibiotic resistance and causes global medical problems. Therefore, the authors aim to fill this gap by proposing an alternative classification ML (random forest) integrated into a virtual screening to identify beta-lactamase inhibitors, validated by experimental data from an in-house chemical library. The authors performed a virtual screening via molecular docking using AutoDock Vina and DOCK6. A modified protocol from a previous report was used to evaluate beta-lactamase inhibitory activity (Everaert and Coenye, 2016). The results indicated that our proposed virtual screening with an ML-based quantitative structure-activity relationship model with a ROC AUC performance score of 0.67 outperformed a non-ML model, a score of 0.56, and better than some previously reported ML models with the best score of 0.51 in identifying bio-inhibitors against beta-lactamase (Anant et al., 2022). In addition, the anti-beta-lactamase activity of some aromatic bioactive molecules from the library was found for the first time. This hints at the potential of natural products in drug discovery. The authors' proposed alternative ML model integrated into a virtual screening initially proves to benefit the identification of beta-lactamase inhibitors based on the in-house chemical library and experiment. However, modification is needed to improve the ML model's performance.

Keywords: Machine learning, virtual drug screening, beta-lactamase inhibitors

References

- Anant, P.S. and Gupta, P., 2022, May. Application of machine learning in understanding bioactivity of beta-lactamase AmpC. In *Journal of Physics: Conference Series* (Vol. 2273, No. 1, p. 012005). IOP Publishing.
- Chen, W., Liu, X., Zhang, S. and Chen, S., 2023. Artificial intelligence for drug discovery: Resources, methods, and applications. *Molecular Therapy-Nucleic Acids*.
- Everaert, A. and Coenye, T., 2016. Effect of β -Lactamase inhibitors on in vitro activity of β -Lactam antibiotics against *Burkholderia cepacia* complex species. *Antimicrobial Resistance & Infection Control*, 5(1), pp.1-8.
- Papastergiou, T., Azé, J., Bringay, S., Louet, M., Poncelet, P. and Gavara, L., 2022, July. Multiple Instance Learning Based on Mol2vec Molecular Substructure Embeddings for Discovery of NDM-1 Inhibitors. In *International Conference on Practical Applications of Computational Biology & Bioinformatics* (pp. 55-66). Cham: Springer International Publishing.

* Presenting author. E-mail address: thanet.pitakbut@fau.de (Thanet Pitakbut)

S3.P54 A bioactive chlorophyll-derived yellow phyllobilin from senescent leaves of *Carica papaya* (Caricaceae) and its potential role in the ethnopharmacological use of withered leaf extracts

Christian Elvert^{a,*}, Rao Zhigang^b, Oriol Ruiz-Catalan^c, Johanna M. Gostner^c, Andreas Koeberle^b, Cornelia Karg^a, Awodayo O. Adepiti^d, Simone Moser^a

^a Institute of Pharmacy, Pharmacognosy, University of Innsbruck, Center for Chemistry and Biomedicine (CCB), Innrain 80-82, 6020 Innsbruck Austria

^b Michel Popp Institute, University of Innsbruck, Mitterweg 24, 6020 Innsbruck Austria

^c Institute of Medical Biochemistry, Medical University of Innsbruck, Innrain 80, 6020 Innsbruck, Austria

^d Department of Pharmacognosy, Obafemi Awolowo University, Ile-Ife, Osun, Nigeria

Extracts of *Carica papaya* leaves, especially withered ones, have been used traditionally by practitioners in Nigeria for the treatment of various diseases (Owoyele et al., 2008).

Phyllobilins are the degradation products of the green plant pigment chlorophyll and accumulate inside the vacuoles of the leaves during senescence of the plant. So far, over 70 different phyllobilins from more than 30 plant species have been identified.

These breakdown products have been completely overlooked as bioactive compounds. In our group, pharmacologically relevant effects of different phyllobilins from several plants could be demonstrated including anti-oxidative properties, anti-proliferative effects on cancer cells (Wang et al., 2021) as well as anti-inflammatory activities (Karg et al., 2021).

In this study we have isolated a yellow phyllobilin from senescent *Carica papaya* leaves, named Cp- PxB, and elucidated its structure with 1D- and 2D NMR experiments. Further we tested the anti-inflammatory potential of the compound in different assays studying the acute as well as the chronic phase of inflammation. The investigation on the senescence-related occurrence of phyllobilins and their bioactivities could provide a scientific rationale for the traditional use of senescent papaya leaves.

Keywords: *Carica papaya*, Caricaceae, phyllobilins, ethnopharmacology, anti-inflammatory

References

- Karg, C.A., Doppler, C., Schilling, C., Jakobs, F., Dal Colle, M.C.S., Frey, N., Bernhard, D., Vollmar, A.M., Moser, S., 2021, A yellow chlorophyll catabolite in leaves of *Urtica dioica* L.: An overlooked phytochemical that contributes to health benefits of stinging nettle. *Food Chemistry*. 359, 129906.
- Owoyele, B.V., Adebukola, O.M., Funmilayo, A.A., Soladoye, A.O., 2008, Anti-inflammatory activities of ethanolic extract of *Carica papaya* leaves. *Inflammopharmacology*. 16, 168-173
- Wang, P., Karg, A.C., Frey, N., Frädriich, J., Vollmar, A.M., Moser, S., 2021, Phyllobilins as a challenging diverse natural product class: Exploration of pharmacological activities. *Archiv der Pharmazie*. 354, 2100061

* Presenting author. E-mail address: christian.elvert@uibk.ac.at (Christian Elvert)

S3.P55 Evaluating the use of Raman spectroscopy for the characterization of volatile compounds by direct measurement on the oregano leaf glands

Galateia Grigoriou^a, Elli Kampasakali^a, Alexandros Nakas^{b,d}, Dimitrios Mertzaniadis^{c,d}, Stella Kokkini^{c,d}, Andreana Assimopoulou^{b,d,*}, Dimitrios Christofilos^a

^a School of Chemical Engineering & Physics Laboratory, Faculty of Engineering, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece

^b Laboratory of Organic Chemistry, School of Chemical Engineering, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece

^c Laboratory of Systematic Botany and Phytogeography, School of Biology, Aristotle University of Thessaloniki 54124 Thessaloniki, Greece

^d Natural Products Research Centre of Excellence (NatPro-AUTh), Center for Interdisciplinary Research and Innovation, Aristotle University of Thessaloniki, 57001 Thessaloniki, Greece

Essential oils (EOs) distilled from leaves and flowers of aromatic plants include volatile organic compounds such as terpenes and phenols. Owing to these compounds, EOs possess numerous and diverse biological properties, namely anti-inflammatory, antimicrobial, antioxidant and anticancer ones (Dhifi et al., 2016). In combination with their natural origin, they find extensive biomedical, cosmetic, food and domestic applications. Unfortunately, adulteration cases of EOs with synthetic products or cheaper EOs are not rare. Hence, the development of simple and reliable analytical techniques for their authentication is of great importance for the industry and commerce. Amongst them, Raman spectroscopy is a versatile, non-invasive method that requires minimal sample preparation, effectively utilized for identifying and quantifying EO constituents (Kampasakali et al., 2023).

In this work, the feasibility of using Raman spectroscopy in the identification of the main volatile components of oregano directly from the leaf glands has been explored for the cases of two μ -Raman systems, a standard benchtop spectrometer (LabRAM HR, Horiba) and a portable one (HE785, Horiba). Measurements on the glands have been compared to those of the respective EOs distilled from the leaves. For both instruments, the identification of volatile components is straightforward. On the other hand, the main volatile components can be identified from the oregano leaf glands with the standard system, but only the presence of carvacrol or thymol can be deduced using the portable one. Improvements in the spatial and spectral resolution of portable instruments may allow their use for quality control of raw oregano material.

Keywords: essential oils, Raman spectroscopy, *Origanum*, identification, adulteration

References

- Dhifi, W., Bellili, S., Jazi, S., Bahloul, N., Mnif, W., 2016. Essential Oils' Chemical Characterization and Investigation of Some Biological Activities: A Critical Review. *Medicines* 3, 25-42.
- Kampasakali, E., Nakas, A., Mertzaniadis, D., Kokkini, S., Assimopoulou, A. N., Christofilos, D., 2023. μ -Raman Determination of Essential Oils' Constituents from Distillates and Leaf Glands of *Origanum* Plants. *Molecules* 28, 1221- 1237.

* Presenting author. E-mail address: adreana@cheng.auth.gr (Andreana Assimopoulou)

S3.P56 New benzothiazines with pain-relieving activity from marine bacterium *Croceibacter atlanticus*

Erin M. Marshall^{a,*}, Amelia Bunnell^a, Lucas Cathey^a, Chenxi Zhu^a, Haoyi Yao^b, Giuliana Panieri^b, James A. Strother^a, Sandra Loesgen^a

^a The Whitney Laboratory for Marine Bioscience, University of Florida, Gainesville, FL, USA

^b Department of Geosciences, CAGE, UiT The Arctic University of Norway, 9037, Tromsø, Norway

Chronic pain is a debilitating and widespread disease, affecting approximately 20% of adults (Cohen et al., 2021). Current therapeutic approaches are either not able to completely relieve pain or have severe side effects such as dependence and addiction (Obeng et al., 2021). Microbial secondary metabolites possess a wide range of biological activities and are a highly promising source for novel drug leads. Here we explore the bioactivity of two *Croceibacter atlanticus* strains and the discovery of sulfur-containing natural products with pain relieving potential.

We employed a novel zebrafish behavior-based assay developed by the Strother Lab able to detect pain mediating activity even in complex extracts. 85 extracts from our collection of Arctic Ocean bacteria were screened for *in vivo* pain-relieving activity. ‘Behavior-guided’ isolation resulted in the discovery of metabolites with pain-relieving activity of the rare benzothiazine class of compounds. Subsequently, *in vivo* zebrafish whole-brain calcium imaging experiments were used to pinpoint targeted neuron populations to elucidate mode of action. Compounds are tested for receptor specificity *in vitro* using cell lines expressing ion channels and submitted to the NIMH’s Psychoactive Drug Screening Program (PDSP) for receptor binding and functional studies.

Keywords: Pain, marine, Gram-negative, zebrafish

References

Cohen, S. P., Vase, L. & Hooten, W. M. Chronic pain: an update on burden, best practices, and new advances. *The Lancet* **397**, 2082–2097 (2021).

Obeng, S., Hiranita, T., León, F., McMahon, L. R. & McCurdy, C. R. Novel Approaches, Drug Candidates, and Targets in Pain Drug Discovery. *J. Med. Chem.* **64**, 6523–6548 (2021)

* Presenting author. E-mail address: erinmarshall@ufl.edu (Erin Marshall)

S3.P57 3D printed polymeric scaffolds loaded with antibiotic and anti-inflammatory drugs for tissue engineering applications

Konstantinos Theodoridis^a, Athanasios Arampatzis^{a,b}, Georgia Liasi^a, Christiana Pliakou^a, Lazaros Tsalikis^c, Panagiotis Barmapalexis^{d,b}, Dimitrios Christofilos^e, Andreana Assimopoulou^{a,b,*}

^a Aristotle University of Thessaloniki, School of Chemical Engineering, Laboratory of Organic Chemistry, Thessaloniki 54124, Greece

^b Center for Interdisciplinary Research and Innovation of Aristotle University of Thessaloniki, Natural Products Research Center of Excellence (NatPro-AUTH), Thessaloniki 57001, Greece ^c Aristotle University of Thessaloniki, School of Dentistry, Thessaloniki 54124, Greece

^d Aristotle University of Thessaloniki, School of Pharmacy, Laboratory of Pharmaceutical Technology, Thessaloniki 54124, Greece

^e Aristotle University of Thessaloniki, Faculty of Engineering, School of Chemical Engineering & Physics Laboratory, Thessaloniki 54124, Greece

Critical skeletal defects and bone loss are common clinical scenarios in orthopedics and craniofacial surgery due to trauma, infection, and tumor resection. For the treatment of bone defects, engineered bone tissue can provide an alternative to conventional bone grafts. The use of innovative biomaterials, 3D bioprinting and autologous mesenchymal stem cells can provide 3D bioscaffolds with regenerative bone tissue properties (Ashammakhi et al., 2022). These 3D-bioscaffolds can further incorporate antimicrobial and/or anti-inflammatory active pharmaceutical ingredients, as modern drug delivery systems. The incorporation of antimicrobial agents is vital for preventing or treating existing infections upon implantation, which is critical for the success of tissue regeneration (Dos Santos et al., 2022).

In this work, tetracycline hydrochloride (TCH), a well-established and broad-spectrum antibiotic agent (produced by a fermentation process, involving microbes), and diclofenac sodium (DIC), a well-established anti-inflammatory drug, were incorporated into a poly(ϵ -caprolactone) (PCL) biomaterial. Drug-free and drug-loaded scaffolds (1 wt.% drug content based on polymer's weight) were manufactured using a bioprinter (BIO-X, Cellink, Sweden). The scaffolds were evaluated for their morphology by scanning electron microscopy (SEM). Their physicochemical properties were determined by means of differential scanning calorimetry (DSC) and thermogravimetric analysis (TGA). Additionally, entrapment efficiency and *in vitro* drug release characteristics were evaluated, while cell biocompatibility was assessed using MC3T3-E1 pre-osteoblast cells. Both neat and drug-loaded scaffolds were tested for their antibacterial activity against *Staphylococcus aureus*.

Keywords: tissue engineering, bone remodelling, 3D-bioprinted scaffolds, antimicrobial activity

References

- Ashammakhi, N., GhavamiNejad, A., Tutar, R., Fricker, A., Roy, I., Chatzistavrou, X., et al., 2022. Highlights on Advancing Frontiers in Tissue Engineering. *Tissue Eng. Part B Rev.* 28, 633-664.
- Dos Santos, D.M., de Annunzio, S.R., Carmello, J.C., Pavarina, A.C., Fontana, C.R., Correa, D.S., 2022. Combining Coaxial Electrospinning and 3D Printing: Design of Biodegradable Bilayered Membranes with Dual Drug Delivery Capability for Periodontitis Treatment. *ACS Appl. Bio Mater.* 5, 146–159.

* Presenting author. E-mail address: adreana@cheng.auth.gr (Andreana Assimopoulou)

S3.P58 ChromAnnot – a webserver for reproducible LC-HRMS/MS chromatogram annotation

Samuel Bertrand^{a,b,*}, Jean-Marie Mottu^a, Catherine Roullier^b, Yann Guitton^c

^a Laboratoire des Sciences du Numérique de Nantes-UMR6004, Université de Nantes, France

^b Institut des Substances et Organismes de la Mer, ISOMer, Nantes Université, UR 2160, Nantes, France

^c Laboratoire d'Etude des Résidus et Contaminants dans les Aliments (LABERCA), LUNAM Université, Oniris, Nantes 44307, France

One main challenge in the exploration of organisms' chemical diversity remains the annotation of compounds detected within LC-HRMS/MS data, to be able to highlight known and/or unknown metabolites (Wolfender *et al.* 2019). Having information on all observed/detected peaks from a LC- HRMS/MS chromatogram remains challenging.

At ISOMer laboratory, we developed an entire workflow taking into consideration each step of the general annotation strategy that was particularly adapted for dereplication of specialized metabolites in LC-HRMS/MS profiles. This automated approach based on R (Cran), XCMS (Tautenhahn *et al.* 2008), IPO (Libiseller *et al.* 2015), CAMERA (Kuhl *et al.* 2012), SIRIUS (Böcker *et al.* 2009), database search, Taxize (Chamberlain *et al.* 2013) and CFMID (Wang *et al.* 2021). Biological origin and MS/MS *in silico* fragmentation comparison were added into the workflow to improve compound discrimination.

The workflow, which analyses directly raw LC-HRMS/MS data (in open format), was integrated into a web-based interface to ease exploring the annotation result. The ChromAnnot webserver is accessible at <https://chromannot.univ-nantes.fr/>.

This deep annotation of LC-HRMS/MS data strategy is currently applied to characterize and highlight known/unknown compounds produced various *Penicillium* strains.

Keywords: annotation, dereplication, LC-HRMS/MS

References

- Böcker, S., M. C. Letzel, Z. Lipták, A. Pervukhin, 2009. SIRIUS: decomposing isotope patterns for metabolite identification. *Bioinformatics* 25, 218-224.
- Chamberlain, S., E. Szöcs, 2013. Taxize: Taxonomic search and retrieval in R. *F1000Research* 2, 191.
- Kuhl, C., R. Tautenhahn, C. Böttcher, T. R. Larson, S. Neumann, 2012. CAMERA: An Integrated Strategy for Compound Spectra Extraction and Annotation of Liquid Chromatography/Mass Spectrometry Data Sets. *Analytical Chemistry* 84, 283-289.
- Libiseller, G., M. Dvorzak, U. Kleb, E. Gander, T. Eisenberg, *et al.*, 2015. IPO: a tool for automated optimization of XCMS parameters. *BMC Bioinformatics* 16, 118.
- Tautenhahn, R., C. Böttcher, S. Neumann, 2008. Highly sensitive feature detection for high resolution LC/MS. *BMC Bioinformatics* 9, 504.
- Wang, F., J. Liigand, S. Tian, D. Arndt, R. Greiner, *et al.*, 2021. CFM-ID 4.0: More Accurate ESI-MS/MS Spectral Prediction and Compound Identification. *Analytical Chemistry* 93, 11692-11700.
- Wolfender, J.-L., J.-M. Nuzillard, J. J. van der Hooft, J.-H. Renault, S. Bertrand, 2019. Accelerating metabolite identification in natural product research: toward an ideal combination of LC-HRMS/MS and NMR profiling, *in silico* databases and chemometrics. *Analytical Chemistry* 91, 704-742.

Acknowledgements

This project is funded by the ANR-18-CE43-0013-01 - FREE-NPs project.

* Presenting author. E-mail address: samuel.bertrand@univ-nantes.fr (Samuel Bertrand)

S3.P59 Exploring the anticancer potential of *Polygonum barbatum* (L.) against colorectal cancer: *In vitro* and *in vivo* investigations

Pi-Kai Chang^a, I-Chuan Yen^b, Wei-Cheng Tsai^c, Kuen-Tze Lin^{d,e}, Shih-Yu Lee^{c,*}

^a Division of Colon and Rectal Surgery, Department of Surgery, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan

^b School of Pharmacy, National Defense Medical Center, Taipei, Taiwan

^c Graduate Institute of Aerospace and Undersea Medicine, National Defense Medical Center, Taipei, Taiwan

^d Department of Radiation Oncology, Cardinal Tien Hospital, New Taipei City, Taiwan

^e School of Medicine, College of Medicine, Fu-Jen Catholic University, New Taipei City, Taiwan

Polygonum barbatum (L.), a medicinal herb, exhibits promising anticancer properties, yet its mode of action remains elusive. This study aimed to elucidate the anti-colorectal cancer (CRC) effects of *Polygonum barbatum* extract (PBE) both *in vitro*, utilizing HCT-116 (KRAS^{G13D} mutation) and HT-29 (BRAF mutation) human CRC cell lines, and *in vivo*, employing a BALB/c nude mice tumour xenograft model. Various analyses including RNA sequencing, western blotting, and co-immunoprecipitation assays were conducted. Additionally, assessments of cell viability, colony formation, migration, invasion abilities, and apoptosis were performed through Cell Counting Kit-8, colony formation, wound healing, Transwell invasion, and flow cytometry assays, respectively. Results demonstrated a significant reduction in CRC cell growth by PBE without impacting non-cancerous cells, accompanied by the downregulation of genes associated with extracellular matrix (ECM) organization, cell motility, and proliferation. Furthermore, PBE affected ECM interactions and focal adhesion pathways, modulating the expression of migration, invasion, and epithelial-mesenchymal transition markers. Notably, in the mouse xenograft model, PBE exhibited tumour growth inhibition, particularly synergizing with oxaliplatin. This study concludes that PBE impedes CRC cell motility and tumorigenicity via the YAP- β -catenin axis, suggesting its potential in addressing CRC resistant to epidermal growth factor receptor inhibitors.

Keywords: colorectal cancer, *Polygonum barbatum*, xenograft mouse model, cell-matrix adhesion

* Presenting author. E-mail address: leeshihyuno1@mail.ndmctsgh.edu.tw (Shih-Yu Lee)

S3.P60 Physical interactions between *Streptomyces* (Streptomycetaceae) and competing microbes trigger the production of natural products

Heli Tirkkonen*, Keith Yamada, Arina Koroleva, Mikko Metsä-Ketelä

University of Turku, Department of Life Technologies, Tykistökatu 6, FIN-20014, Turku, Finland

Streptomyces (Streptomycetaceae) are known for producing numerous natural products, including antimicrobial agents. However, antimicrobial resistance is rising, and the discovery of new antimicrobial agents has slowed down. While genome sequencing reveals a vast reservoir of secondary metabolites in *Streptomyces*, these often remain unexpressed in laboratory monocultures (Lewis, 2013). In our work, we have discovered predatory behavior by *Streptomyces* when encountering *Saccharomyces cerevisiae* (Saccharomycetaceae). Physical interactions with *S. cerevisiae* trigger *Streptomyces* to produce numerous lytic enzymes and antifungal compounds, such as pentamycin and filipin III, which are not produced in monoculture settings. RNA-seq data uncovered activation of 40 % of biosynthetic gene clusters of *Streptomyces lavendulae*, including unknown secondary metabolite biosynthetic gene clusters (Yamada et al., 2023).

To expand our research, we aimed to utilize cutting-edge technologies to study the interactions between *Streptomyces* and competing microbes. We have selected 50 *Streptomyces* strains and four non-pathogenic microbes that are related to important human pathogens: *Candida tropicalis*, *Aspergillus nidulans*, *Mycobacterium smegmatis*, and *Pseudomonas syringae*. Our preliminary results show that 21 of the selected *Streptomyces* strains react to the presence of the competing microbes by producing different compounds depending on the microbe. We utilized Global Natural Products Social Molecular Networking (Wang et al., 2016) and SIRIUS (Dührkop et al., 2019) to uncover the identity of the produced secondary metabolites, such as pamamycins, puromycins, and lydicamycins. We aim to further uncover the identity of all the unknown compounds by combining transcriptomics, metabolomics, and knock-outs.

Keywords: *Streptomyces*, drug discovery, microbial interactions

References

- Dührkop, K., Fleischauer, M., Ludwig, M., Aksenov, A. A., Melnik, A. V., Meusel, M., Dorrestein, P. C., Rousu, J., Böcker, S., 2019. SIRIUS 4: A Rapid Tool for Turning Tandem Mass Spectra into Metabolite Structure Information. *Nat Methods*. 16 (4), 299–302.
- Lewis, K., 2013. Platforms for antibiotic discovery. *Nat Rev Drug Discov*. 12, 371–387.
- Wang, M., Carver, J. J., Phelan, V. V., Sanchez, L. M., Garg, N., Peng, Y., Bandeira, N., 2016. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat Biotechnol*. 34, 828– 837.
- Yamada, K., Koroleva, A., Tirkkonen, H., Siitonen, V., Laughlin, M., Akhgari, A., Mazurier, G., Niemi, J., Metsä-Ketelä, M., 2023. Physical interactions trigger *Streptomyces* to prey on yeast using natural products and lytic enzymes. *bioRxiv* 2023.06.15.545052

* Presenting author. E-mail address: hmtirk@utu.fi (Heli Tirkkonen)

S3.P61 Energy Saving Efficacy and Optimized condition of pulse energy fields technology on *Centella asiatica* extraction

Kyung-Eon Oh^{a,*}, Na Young Lee^a, Do-Hyeong Kim^b, Hyuk Joon Choi^b, You Ah Kim^a,
Byoungjun Park^{a,*}

^a Skin&Natural products Lab, Kolmar Korea, 61 Heolleung-ro 8 gil, Seocho-gu, Seoul, Korea

^b BK Bio, 2706-38 Iljudong-ro, Gujwa-eup, Jeju-si, Jeju-do, Korea

Eco-friendly technologies are focused worldwide to deal with climate crisis and resource depletion. As pulsed electric fields (PEF) treatment changes permeability of cell membrane is used as energy- saving green technology for sterilization in the food industry (Wei et al., 2023). To extend the application of PEF into the industrial realm, we attempted PEF on *Centella asiatica* (L.) Urban (Apiaceae).

C. asiatica, which is Southeast Asian native medicinal plant, is known to have several biological activities including antioxidation, anti-inflammation etc. (Orhan, 2012). *C. asiatica* has identical compounds, madecassoside, asiaticoside, madecassic acid, asiatic acid, which are reported to have wound healing activity and collagen stimulation in human fibroblast (Maquart et al., 1990).

To compare energy consumption and extracted contents of four well-known actives between PEF treated and non-treated extract of *C. asiatica* leaves, extract was sampled every five minutes. As the result, the maximum levels of active compounds were higher at PEF treated extract and found out the energy consumption to extract madecassic acid same level of the maximum extracted at non-treated extract showed about 20 % lower in PEF treated extract.

Next, optimization of PEF treatment was established with Box-Benhken design. Time, voltage, and frequency were selected as variables to maximize extraction of madecassoside, asiaticoside and yield. The optimized *C. asiatica* extract showed 18.39 ppm of madecassoside, 22.98 ppm of asiaticoside, 1.30 % of yield, similar to expected values. Therefore, PEF can be considered as new environmentally friendly solution for cosmetics with maximizing active compounds.

Keywords: pulsed electric fields, *Centella asiatica*, energy saving, optimization

References

- Wei, S., Chen, T., Hou, H., Xu, Y., 2023. Recent advances in electrochemical sterilization. *Journal of Electroanalytical Chemistry*, 937, 117419.
- Orhan, I. E., 2012. *Centella asiatica* (L.) Urban: From traditional medicine to modern medicine with neuroprotective potential. *Evidence-based complementary and alternative medicine*, 2012, 946259.
- Maquart, F.X., Bellon, G., Gillery, P., Wegrowski, Y., Borel, J.P., 1990. Stimulation of collagen synthesis in fibroblast cultures by a triterpene extract from *Centella asiatica*. *Connect. Tissue Res.* 24, 107-120

* Presenting author. E-mail address: ohkyung91@kolmar.co.kr (Kyung-Eon Oh)

S3.P62 Identification of a leoligin-inspired synthetic lignan as a novel TGR5 agonist

Alexander F. Perhal^{a,*}, Patrik F. Schwarz^a, Thomas Linder^b, Marko D. Mihovilovic^b, Michael Schnürch^b, Verena M. Dirsch^a

^a Department of Pharmaceutical Sciences, University of Vienna, Vienna, Austria

^b Institute of Applied Synthetic Chemistry, TU Wien, Vienna, Austria

The G protein-coupled receptor TGR5 binds bile acids as endogenous ligands and emerged as an important regulator of metabolic and immunologic functions. It was shown to be an effective target in alleviating inflammatory diseases (Pols et al., 2011). The naturally occurring lignan leoligin and synthetic derivatives were shown to be effective anti-inflammatory compounds *in vitro* (Linder et al., 2021). However, the underlying molecular targets for these activities are still not fully understood and require further investigations. In this study we present the identification of a natural product- inspired synthetic derivative of leoligin, LT-188A, as a novel TGR5 agonist. Both, a CRE-Luciferase reporter gene assay and cAMP accumulation assay supported TGR5 agonism of LT-188A (3-30 μ M). Specificity for TGR5 was shown by comparing activities in HEK293A cells expressing TGR5 *versus* non-expressing cells. In line with this agonistic activity, the transactivation activity of the pro- inflammatory transcription factor NF κ B was inhibited by LT-188A concentration- (3-20 μ M) and TGR5-dependently as shown by luciferase reporter assays. Functional consequences of these activities were studied in LPS-activated macrophages: the expression levels of pro-inflammatory cytokines (IL-1 β , IL-6) and the release of nitric oxide were concentration-dependently (3-20 μ M) reduced in response to LT-188A as shown by qPCR and the Griess Assay, respectively. Cytotoxicity was excluded by the Resazurin conversion assay. Taken together, LT-188A shows promising *in vitro* anti-inflammatory bioactivities in relevant cellular assays and provides a starting point for further optimization and development.

Keywords: TGR5, anti-inflammatory, macrophages, leoligin, lignans

References

- Linder, T., Papaplioura, E., Ogurlu, D., Geyrhofer, S., Hummelbrunner, S., Schachner, D., Atanasov, A.G., Mihovilovic, M.D., Dirsch, V.M., Schnürch, M., 2021. Investigation of Leoligin Derivatives as NF- κ B Inhibitory Agents. *Biomedicines*. 10(1), 62.
- Pols, T.W., Noriega, L.G., Nomura, M., Auwerx, J., Schoonjans, K., 2011. The bile acid membrane receptor TGR5 as an emerging target in metabolism and inflammation. *J Hepatol*. 54(6), 1263-1272.

* Presenting author. E-mail address: alexander.perhal@univie.ac.at (Alexander Perhal)

S3.P63 Nano-phytosome delivery system loaded with *Crataegus laciniata* (Rosaceae) flowers extract: whitening properties

Shiva Pouramin Arabi^c, Salvatore Mirabile^{a b}, Valeria D'Angelo^a, Maria Paola Germanò^a, Valentina Parisi^c, Francesco Maria Raimondo^d, Emanuele Rosa^c, Fatemeh Adinehpour^e, Seyed Fakhreddin Hosseini^f, Nunziatina De Tommasi^c

^a Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Viale F. Stagno d'Alcontres 31, 98166 Messina, Italy

^b Foundation Prof. Antonio Imbesi, University of Messina, Piazza Pugliatti 1, 98122 Messina, Italy

^c Department of Pharmacy, University of Salerno, 84084 Fisciano, Italy

^d PLANTA/Centro autonomo di Ricerca, Documentazione e Formazione, Via Serraglio Vecchio 28, 90123 Palermo, Italy

^e Department of Food Materials and Process Design Engineering, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

^f Department of Seafood Processing, Faculty of Marine Science, Tarbiat Modares University. PO Box 46414-356, Noor, Mazandaran, Iran

Hawthorn is a large genus of small trees belonging to Rosaceae family, including approximately 200 species. In recent years, hawthorn has been demonstrated to be an excellent source of natural bioactive molecules, which have promising benefits for human health (Venskutonis, 2018). In the present study, we focused on *Crataegus laciniata* Ucria, a species distributed through the western Mediterranean area, northern Algeria, and Italy (Puglia and Sicilia). Nowadays there is still a great interest for the research of novel non-toxic, effective and stable tyrosinase inhibitory agents, from plant sources (Pillaiyar et al., 2017). Our previous study demonstrated the efficacy of *C. laciniata* flower's (CLF) ethanolic (70%) extract as new sources of antimelanogenic agents (Mirabile, et al., 2023). The chemical profile of extract showed the presence of both C-flavonoid and O-flavonoid derivatives and hyperoside and vitexin. Phytosome system could enhance skin penetration and bioavailability of phytoconstituent like polyphenol-rich extract. The objective of this research was to formulate phytosome containing CLF extract to enhance skin-whitening properties of the extract. In order to improve the physicochemical properties of the extract, the obtained CLF extract was developed into phytosome system by thin-layer method using soy lecithin. FT-IR and XRD diffraction confirmed the successful loading of CLF in nano-phytosomes. The phytosome complex containing CLF extract and lecithin (1:1) has good entrapment efficiency (94.8%) with spherical shape as confirmed by SEM. The CLF-loaded phytosome displayed particle size in the range of 338-534 nm and polydispersity index (PDI) values in the range of 0.39-0.47. This herbal preparation could be developed as tyrosinase inhibitory agent in skin pharmacology and cosmetics.

Keywords: tyrosinase, *Crataegus*, Phytosome, whitening properties, thin-layer method

References

- Venskutonis, P.R. Phytochemical composition and bioactivities of hawthorn (*Crataegus* spp.): Review of recent research advances. 2018. J. Food Bioact. 4, 69-87.
- Pillaiyar, T., Manickam, M., Jung, S.-H. 2017. Recent development of signaling pathways inhibitors of melanogenesis. Cell. Signal. 40, 99-115.
- Mirabile, S., D'Angelo, V., Germanò, M. P., Pouramin Arabi, S., Parisi, V., Raimondo, F. M., & Rosa, E. 2023. Chemical Profile and Health-Promoting Activities of *Crataegus laciniata* (Rosaceae) Flowers. Plants, 13(1), 34.

* Presenting author. E-mail address: spouraminarabi@unisa.it (Shiva Pouramin Arabi)

S3.P64 Leoligin derivatives as novel phosphodiesterase 4 inhibitors

Christine Coffey^{a,*}, Alexander F. Perhal^{a,#}, Lorenza Bertaina^a, Thomas Linder^b, Marko D. Mihovilovic^b, Michael Schnürch^b, Verena M. Dirsch^a

^a Department of Pharmaceutical Sciences, University of Vienna, Vienna, Austria

^b Institute of Applied Synthetic Chemistry, TU Wien, Vienna, Austria

Phosphodiesterase 4 (PDE4) is the primary hydrolysing enzyme of the ubiquitous second messenger, cyclic adenosine monophosphate (cAMP). PDE4 plays a crucial role in physiological and metabolic functions, including pro-inflammatory pathways (Francis et al., 2011). Marketed PDE4 inhibitors, namely *Roflumilast* and *Apremilast*, have been used to treat chronic obstructive pulmonary disease and psoriasis, respectively (Baillie et al., 2019). Identification and characterisation of PDE4 inhibitors provides a promising strategy in targeting a variety of pathological processes.

In this study, leoligin, a natural lignan found in the roots of Edelweiss (*Leontopodium nivale ssp. alpinum*, Asteraceae) and 169 structural derivatives (Mihovilovic et al., 2015) were screened for their PDE inhibitory activity at 30 μ M using a cAMP accumulation assay employing an EPAC-based FRET biosensor (Perhal, 2020). Seven compounds, including leoligin itself, were identified as causing a significant accumulation of cAMP and a structure-activity relationship was deduced. One derivative, designated LT-104A, showed a dose-dependent activity with the highest determined potency ($EC_{50} = 2 \mu$ M) in the EPAC assay. LT-104A was further characterised using a CRE- Luciferase assay, showing comparable activity to known PDE4 inhibitors in inducing the cAMP- PKA-CREB pathway. Lastly, a functional study was carried out in LPS-activated macrophages in which LT-104A reduced nitric oxide release in the Griess assay, indicative of its anti-inflammatory properties.

In conclusion, extensive *in vitro* screening of leoligin derivatives led to the identification and characterisation of a novel PDE4 inhibitor, LT-104A, with promising *in vitro* anti-inflammatory properties.

Keywords: cyclic AMP, phosphodiesterase, PDE4, leoligin, lignans

References:

- Francis, S.H., Blount, M.A., Corbin, J.D., 2011. Mammalian cyclic nucleotide phosphodiesterases: molecular mechanisms and physiological functions. *Physiol. Rev.* 91, 651-690.
- Baillie, G.S., Tejada, G.S., Kelly, M.P., 2019. Therapeutic targeting of 3',5'-cyclic nucleotide phosphodiesterases: inhibition and beyond. *Nat. Rev. Drug Discov.* 18, 770-796.
- Mihovilovic, M.D., Linder, T., Atanasov, A., Dirsch, V.M., Stuppner, H., Schwaiger, S., Bernhard, D., 2015. Novel leoligin derivatives for applications as proliferation inhibitors. Austrian Patent Application AT 2015/050160
- Perhal, A., 2020. Elucidation of Signal Transduction Mechanisms with Pathological Relevance for Cardiovascular Diseases in-vitro and in-vivo. Doctoral Thesis, ETH Zurich. DOI: 10.3929/ethz-b-000440925

* Presenting author. E-mail address: christine.coffey@univie.ac.at (Christine Coffey)

S3.P65 The search for nontraditional synergy: pairing *in vitro* omics and *in silico* machine learning methods to discover novel combinations of antimicrobial agents from natural products

Katherine L. Lev^{a,b,*}, Margaret M. Reuter^{f,g,*}, David H. Sherman^{a,b,c,d,e}, Sriram Chandrasekaran^{b,f,g}

^a Life Sciences Institute, University of Michigan, Ann Arbor, MI, USA

^b Program in Chemical Biology, University of Michigan, Ann Arbor, MI, USA

^c Department of Medicinal Chemistry, University of Michigan, Ann Arbor, MI, USA

^d Department of Microbiology & Immunology, University of Michigan, Ann Arbor, MI, USA


^e Department of Chemistry, University of Michigan, Ann Arbor, MI, USA

^f Department of Biomedical Engineering, University of Michigan, Ann Arbor, MI, USA

^g Michigan Medicine

In response to lagging antibiotic development pipelines and to combat multi-drug resistance, antimicrobial combination therapy is a promising therapeutic option, and novel compounds that act synergistically are of clinical interest. However, combinations are often chosen empirically leading to suboptimal treatment outcomes and spread of resistance. Antibacterial adjuvants, such as natural products (NPs), have been understudied as most NP library campaigns screen individual compounds for bioactivity. The vast sample size of combinations make it impossible to systematically screen combinations *in vitro*. Recent methods to develop synergistic treatments depend on computational models to reduce the vast sample size of combinations available. These methods are limited by (1) requiring inaccessible, costly datasets for novel NP compounds and (2) black-box AI which provides little insight into mechanisms of synergy that govern drug interactions.

We hypothesized that by pairing computational drug – protein and NP – protein interactions with experimentally obtained chemogenomics via CRISPR interference (CRISPRi) screens, we could 1. predict synergy/antagonism of FDA approved antimicrobials with NPs, 2. undertake an unbiased exploration of the underlying biological mechanisms of synergy via profiling perturbed bacterial genetic networks. Our model uses a unique combination of computationally calculated drug – protein interactions, multi-omics studies, and machine learning (ML) to predict effective drug combination therapies.



(Fig. 1). The model was trained and tested on 648 two-way and three-way drug interactions in *E. coli*. To assess drug modes of action and mechanisms of synergy, we cross-referenced all five datasets to the Kegg database of 113 *E. coli* functional pathways and known drug targets.

Keywords: machine learning, antibiotics, synergy, target, CRISPRi-chemogenomics

*Presenting author. Email address: katlev@umich.edu (Katherine Lev)

S3.P66 Electron diffraction-guided natural product discovery: identification and isolation of new scaffold class from *Podospora australis*

David A. Delgadillo^{a,*}, Lin Wu^b, Jessica E. Burch^a, Lygia Silva de Moraes^a, Caroline Wang^d, Benjamin P. Tu^d, Yi Tang^{b,c}, Hosea M. Nelson^a

^a Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA 91125, USA

^b Departments of Chemical and Biomolecular Engineering, ^c Chemistry and Biochemistry, University of California, Los Angeles, Los Angeles, CA 90095, USA

^d Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, TX 75390-9152, USA

This presentation will highlight the utility of ArrayED (Delgadillo et. al., 2024) as an effective screening platform to rapidly discover novel natural products. The presentation will also showcase how microED is employed in conjunction with high-resolution electron spray ionization mass spectrometry (HRESIMS) to elucidate the structures of novel scaffolds, as well as the use of nuclear magnetic resonance (NMR) spectroscopy for further supporting evidence.

As part of our continuing search for novel natural products, we identified several crystalline entities from a MeOH extract of *Podospora australis* utilizing ArrayED. Routine microED analysis provided the structural solution of a scaffold previously undescribed in the literature. HRESIMS of the well of interest revealed the presence of eight unique mass signatures, indicating that microED had produced a structural solution from a highly mixed fraction. Careful analysis of bond lengths and geometry, coupled with the observed m/z 477.1960 [M+H]⁺, enabled the elucidation of the structure of **1** (Fig. 1). Subsequent regrowth of the culture and isolation efforts yielded several analogs of **1**, which were characterized *via* 2D NMR to further support the structural solution provided by microED and HRESIMS.




Fig. 1. Structure of the first novel natural product discovered via ArrayED **1**

Since the *Podospora australis* strain is known to produce anti-fungal compounds (Li et. al., 2016), the newly discovered compounds were submitted for anti-fungal activity screening. Two of the six compounds exhibited modest activity in a general screen against *S. cerevisiae*. These preliminary biological activity results are encouraging, and further biological evaluation is underway.

Keywords: microED, ArrayED, antifungal, *Podospora*, drug discovery

References

- Delgadillo, D.A., Burch, J.E., Kim, L.J., de Moraes, L.S., Niwa, K., Williams, J., Tang, M.J., Lavallo, V.G., Chhetri, B.K., Jones, C.J., Rodriguez, I.H., Signore, J.A., Marquez, L., Bhanushali, R., Woo, S., Kubanek, J., Quave, C., Tang, Y., Nelson, H.M., 2024. High-Throughput Identification of Crystalline Natural Products from Crude Extracts Enabled by Microarray Technology and microED. *ACS Cent. Sci.*, 10(1), 176-183.
- Li, Y., Yue, Q., Krausert, N. M., An, Z., Gloer, J. B., & Bills, G. F. (2016). Emestrins: anti-*Cryptococcus* epipolythiodioxopiperazines from *Podospora australis*. *Journal of Natural Products*, 79(9), 2357-2363.

* Presenting author. E-mail address: ddelgadi@caltech.edu (David A. Delgadillo)

S3.P67 Metaboseek 2.0: Deconvoluting the complexity of MS/MS Big Data through interactive data processing

Alexandros Polyzois*, Maximilian J. Helf, Goncalo Gouveia, Frank C. Schroeder

Boyce Thompson Institute and Department of Chemistry and Chemical Biology, Cornell University, Ithaca, New York 14853, United States

In the current era of technological advancements in mass spectrometry, we are now capable of acquiring thousands of high-quality features in a single analysis. This breakthrough signifies that the solution to a drug discovery or chemical ecology question could already be within our grasp. Significant efforts in the computational mass spectrometry field have been devoted to unlocking the full potential of this vast amount of recorded data (Nothias et al., 2020, Shah et al., 2023). In this context, Metaboseek (Helf et al., 2022) represents a notable step forward by integrating the primary advantages of these computational approaches in a user-friendly interface, and it additionally incorporates an effective data filter. Specifically, it offers an interactive workflow that seamlessly begins with data preprocessing using an integrated XCMS step (Tautenhahn et al., 2012). The processed data undergo comprehensive statistical analysis, culminating in the prediction of the prioritized features' identity through SIRIUS (Dührkop et al., 2019). Metaboseek's updated version pivotal characteristic lies in a sophisticated set of filters, including fold change and machine learning prioritization, designed to streamline the feature table. These filters ensure that only features meeting specific criteria are retained, thereby enhancing the efficiency and relevance of subsequent analyses. Here, we demonstrate the software's effectiveness in feature prioritization, elucidating metabolic pathways, and identifying novel targets.

Keywords: Metaboseek, computational mass spectrometry, metabolomics

References

- Dührkop, K., Fleischauer, M., Ludwig, M., Aksenov, A.A., Melnik, A.V., Meusel, M., Dorrestein, P.C., Rousu, J., Böcker, S., 2019. SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure information. *Nature methods*, 16, 4, 299-302.
- Helf, M.J., Fox, B.W., Artyukhin, A.B., Zhang, Y.K., Schroeder, F.C., 2022. Comparative metabolomics with Metaboseek reveals functions of a conserved fat metabolism pathway in *C. elegans*. *Nature Communication*, 13, 1, 782.
- Nothias, L.F., Petras, D., Schmid, R., Dührkop, K., Rainer, J., Sarvepalli, A., Protsyuk, I., Ernst, M., Tsugawa, H., Fleischauer, M., Aicheler, F., et al., 2020. Feature-based molecular networking in the GNPS analysis environment. *Nature methods*, 17, 9, 905-908.
- Shah, A.K.P., Walter, A., Ottosson, F., Russo, F., Navarro-Díaz, M., Boldt, J., Kalinski, J.C., Kontou, E.E., Elofson, J., Polyzois, A., González-Marín, C., et al. 2023. The Hitchhiker's Guide to Statistical Analysis of Feature-based Molecular Networks from Non-Targeted Metabolomics Data. *ChemRxiv*
- Tautenhahn, R., Patti, G.J., Rinehart, D., Siuzdak, G., 2012. XCMS Online: a web-based platform to process untargeted metabolomic data. *Analytical chemistry*, 84, 11, 5035-5039.

* Presenting author. E-mail address: a.polyzois92@gmail.com (Alexandros Polyzois)

S3.P68 Cytotoxic activities of terpenoids from genera *Euphorbia rowlandii* and *Euphorbia grandicornis* (Euphorbiaceae) against the Triple-negative breast cancer (TNBC)

Jacqueline Tembu^{a,*}, Kemboi Magozwi^a, Mmabatho Monica Dinala^a, Wilma Augustyn^a, Xavier Siwe-Noundou^b, Getrude Nyemba^c, Candance Davidson^c, Jo Anne De LaMare^c

^a Department of Chemistry, Tshwane University of Technology, Private Bag x680, Pretoria, South Africa

^b School of Pharmacy, Department of Pharmaceutical Science, Sefako Makgatho Health Sciences University, PO Box 218, South Africa

^c Department of Biochemistry and Microbiology, Female Cancers Research at Rhodes University (FemCR2U), 6140, Makhanda, South Africa

Triple-negative breast cancer cells (TNBC) are a heterogeneous group of malignancies that affect women. This subtype accounts for approximately 10 to 15% of diagnosed breast cancers worldwide and in Africa, more than 50% of women diagnosed with (TNBC) succumb to the disease. The genus *Euphorbia* contains terpenoids with anticancer properties, (Damoun et al, 2015; Lebert et al., 2018; Chang-Qing et al., 2020; Yan et al., 2020; Anyawu 2008), hence exploring *Euphorbia rowlandii* R.A Dyer and *Euphorbia grandicornis* L, phytochemicals and evaluated their cytotoxic properties. Structures of the isolated terpenoids from the genera *E. grandicornis* and *E. rowlandii* were characterised using IR, NMR, and MS. *E. grandicornis* yielded, hexyl(E)-3-(4-hydroxy-3-methoxyphenyl)-2-propenoate (1), tirucalla-8,25-diene-3 β ,24R-diol (2), 24-methylenetirucalla-8-en-3 β -ol (3), β -glutanol, β -amyryn (4) whereas *E. rowlandii*, afforded phorbol ester, 16-Hydroxy-12-deoxy-phorbol (5), a tentative phorbol ester, 3 β -hydroxy-5-glutinene (6). Extracts and compounds were tested for cytotoxicity using MTT against MCF-7, HCC70, and MCF-12A cells. Dichloromethane extract of *E. rowlandii* showed high toxicity against HCC70 and MCF-7 cells at 4.97 and 1.177 Mm, while *E. grandicornis* extract exhibited IC₅₀ values of 1.03, 0.301 μ g/mL. Compound 1 showed activity against HCC70 cells with an IC₅₀ = 29.45 μ M and MCF-7 with IC₅₀ = 23.41 μ M and compound 5 showed activity against HCC70 and MCF-7 cells with IC₅₀ values of 0.592 and 1.003 μ M respectively. Terpenoids from *Euphorbia* genus demonstrated cytotoxic activities against the breast cancer cells.

Keywords: *Euphorbia*, TNBC, HCC70, MCF-7, terpenoids

References

- Damoun, S., D. Fattaneh, D., Tavassoli., A. 2015. A targetable androgen receptor-positive breast cancer subtype hidden among the triple-negative cancers. Arch. Pathol. Lab. Med., 139, pp. 612-617.
- Lebert, J., Lester, R., Powell, E., Seal, M., McCarthy, J., 2018. Advances in the systemic treatment of triple-negative breast cancer, Curr. Oncol., 25, pp. 142-150, 10.3747/co.25.3954
- Chang-Qing, Y L. Jie, L Shi-Qi, Z., Kun, Z., Zi-Qian, G., Ran, X., Ren-Ben, Z., Gang, Z., Da-Chuan, Y., Chen-Yan, Z., 2020. Recent treatment progress of triple-negative breast cancer. Prog. Biophys. Mol. Biol., 151, pp. 40-53.
- Anyanwu, S., 2008. Temporal trends in breast cancer presentation in the third world. J Exp Clin Cancer Res, Jul 11;27(1):17

* Presenting author. E-mail address: Tembuwj@tut.ac.za (Jacqueline Tembu)

S3.P69 Simultaneous extraction and synthesis of new cannabinoid acid esters and their evaluation on TRPA1, cell proliferation and in MDA-MB-231 xenograft model of cancer

Evangelos Dadiotis^{a,*}, Aikaterini Papakonstantinou^{a,b}, Meng Cui^c, Eleni Melliou^a, Vangelis Mitsis^d, Diomedes Logothetis^c, Haralabia Boleti^b, Prokopios Magiatis^a

^a Department of Pharmacy, National and Kapodistrian University of Athens, Greece

^b Intracellular Parasitism Laboratory, Department of Microbiology, Hellenic Pasteur Institute, Athens, Greece

^c Department of Pharmaceutical Sciences, School of Pharmacy, Bouvé College of Health Sciences, Northeastern University, Boston, USA

^d Ekati Alchemy Lab SL, Barcelona, Spain

Natural phytocannabinoids have a privileged pharmacophore scaffold with antitumor properties (Hinz and Ramer, 2022). In this study we present the simultaneous extraction and synthesis of cannabinoid acid esters. The synthesis method was performed for esters of cannabidiolic acid (CBDA), cannabigerolic acid (CBGA) and cannabichromenic acid (CBCA) with C1-5 alcohols. Cannabinoid acid esters and their neutral forms (CBD, CBG and CBC) were evaluated for their effect on TRPA1 channels, which are implicated in several cancer forms (Takahashi et al., 2018), and for their cytotoxicity. Trichomes of *Cannabis sativa* containing either CBDA or CBGA, were used for the synthesis of each ester. Each time a different alcohol was used as solvent/reagent for esterification in the presence of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide and 4-dimethylaminopyridine. The reaction with ultrasonication led to 90% yield of each ester. CBCA esters were synthesized by oxidizing CBGA esters with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone with 80% yield. Cannabinoid esters showed similar effect on TRPA1 activation, with their respective neutral cannabinoids in a single dose experiment (in 50 μ M, 2-fold higher activation compared to AITC). The *in vitro* study of the cytotoxic activity with the MTT assay showed that CBG and its respective esters were the most potent with $EC_{50} < 40 \mu$ M in MDA-MB-231 cancer cells. CBGA butyl ester ($EC_{50} = 8.7 \mu$ M) was also evaluated in an *in vivo* MDA-MB-231 xenograft mouse model of cancer (oral administration, 50 mg/kg). Additionally, CBGA butyl ester was classified to “Category 5” of GHS (2000 mg/kg) in an oral acute toxicity assessment in mice (Fig.1).

Fig. 1. Structure of cannabigerolic acid butyl ester (CBGA butyl ester)

Keywords: cannabinoid acid esters, cytotoxicity, cannabinoids, antitumor, TRPA1 channels

References

- Hinz, B., Ramer, R., 2022. Cannabinoids as anticancer drugs: current status of preclinical research. *Br J Cancer* 127, 1–13
- Takahashi, N., Chen, H.-Y., Harris, I.S., Stover, D.G., Selfors, L.M., Bronson, R.T., Deraedt, T., Cichowski, K., Welm, A.L., Mori, Y., Mills, G.B., Brugge, J.S., 2018. Cancer Cells Co-opt the Neuronal Redox-Sensing Channel TRPA1 to Promote Oxidative-Stress Tolerance. *Cancer Cell* 33, 985–1003

* Presenting author. E-mail address: edadioti@pharm.uoa.gr (Evangelos Dadiotis)

S3.P70 Exploring potential breast and melanoma cancer drug candidates from *Crocus sativus*: molecular docking insights

Olha Mykhailenko^{a,b,*}, Victoriya Georgyants^b

^a Pharmacognosy and Phytotherapy, UCL School of Pharmacy, 29-39 Brunswick Square, WC1N 1AX, London, UK

^b Department of Pharmaceutical Chemistry, National University of Pharmacy, 4-Valentynivska Str., 61168, Kharkiv, Ukraine

Molecular docking can be particularly useful in identifying potential drug candidates from natural sources, exploring the binding mechanisms of natural products with therapeutic targets, and guiding the optimization of lead compounds derived from natural sources [1]. The current molecular docking aimed the analysis of descriptors with the greatest cytotoxic activity and the study of the possible mechanism of action of selected phenolic compounds of *Crocus sativus*. Previously [2], we identified the following compounds: chlorogenic acid, caffeic acid, mangiferin, isoorientin, ferulic acid, rutin, tectoridin, quercetin, t-cinnamic acid, genistin, apigenin, kaempferol, iristectorigenin B, nigricin and irigenin. The docking simulations were performed with the SCIGRESS software package (Fujitsu, Fukuoka, Japan (license 742F6852C191)). Among estrogen receptors that play an important role in cancer pathogenesis, we selected the human ER α -LBD (PDB ID 3ERT) complex with 4- hydroxytamoxifen, which is an active metabolite of tamoxifen. According to the docking studies, almost all natural compounds showed affinity to the active sites of the selected enzymes that were similar or even better than their native ligands. Such compounds as chlorogenic acid, isoorientin, ferulic acid, tectoridin, quercetin, cinnamic acid, genistin, apigenin, kaempferol, and irigenin showed, better affinity compared with the native ligands of breast cancer proteins 4RJ3, 2IOK, and 4XYF. Moreover, the affinity of many compounds (chlorogenic acid, ferulic acid, cinnamic acid, iristectorigenin B) to the estrogen receptor (3ERT) was comparable to hydroxytamoxifen. Subsequent *in vitro* assays of *Crocus* extracts and individual substances correlated with the results of docking.

Keywords: natural compounds, breast cancer, melanoma, docking studies, drug designing

References

1. Asiamah et al. 2023. Applications of molecular docking in natural products-based drug discovery. Scientific African, 20: e01593.
2. Mykhaolenko et al.2022. Pharmacological Potential and Chemical Composition of *Crocus sativus* Leaf Extracts. Molecules. 27: 10.

* Presenting author. E-mail address: o.mykhailenko@ucl.ac.uk (Olha Mykhailenko)

S3.P71 Metabolic fate of oleocanthal and oleacein after oral consumption: Novel biomarkers, human bioactivity insights and TRPA1 channel activation

Nikoleta Anna Madelou^{a,*}, Evangelos Dadiotis^{a,b}, Maria Nikolantonaki^c, Eleni Melliou^a, Tzortzis Nomikos^d, Prokopios Magiatis^a

^a Laboratory of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimioupolis Zografou 15 771, Athens, Greece

^b Department of Pharmaceutical Sciences, School of Pharmacy and Pharmaceutical Sciences, Bouvé College of Health Sciences, Boston, MA 02115, USA

^c Univ. Bourgogne Franche-Comté, AgroSup Dijon, PAM UMR A 02.102, Institut Universitaire de la Vigne et du Vin – Jules Guyot, F-21000 Dijon, France

^d Department of Nutrition and Dietetics, Harokopio University, Athens, Greece

Oleocanthal and oleacein are secoiridoids found in olive oil, which have gained strong scientific interest due to their pharmacological properties. Although some efforts have been done to investigate their pharmacokinetics in mice or rats (Darakjian et al., 2021; López-Yerena et al., 2021), their metabolic fate in humans after oral consumption remains unknown. In this study, we present the results of a human-based protocol involving the oral administration of a commercially available olive oil extract enriched in oleocanthal and oleacein, in the form of a food supplement capsule (OLEOPROTECT®) in comparison with high-phenolic olive oil. UHPLC MS/MS analysis, for the first time succeeded to identify metabolites of oleocanthal and oleacein in human plasma: oleocysteine, oleotaurine and hydroxyoleocysteine, all making a peak at 30 min and being detectable even 4 h after oral consumption. Oleocanthal and oleacein were not detected in plasma, confirming the previously described spontaneous reactivity with amino acids. Interestingly, the administration of the encapsulated product showed higher rate of formation of the metabolites in comparison with the respective olive oil. Extensive studies investigated the spontaneous reactions with all plasma amino acids. Real-time structural elucidation through 1D and 2D NMR experiments revealed the biochemical transformation of oleocanthal and oleacein into a series of novel products. Furthermore, we investigated the activity of oleocanthal and some of the formed metabolites in TRPA1 channels. Notably, oleocysteine activated TRPA1 similarly with oleocanthal. This discovery provides a new perspective on understanding the well-documented bioactivities of olive oil in humans, offering valuable insights for future research.

Keywords: oleocanthal, oleacein, pharmacokinetics, biological fluids, TRPA1

References

- Darakjian, L.I., Rigakou, A., Brannen, A., Qusa, M.H., Tasiakou, N., Diamantakos, P., Reed, M.N., Panizzi, P., Boersma, M.D., Melliou, E., El Sayed, K.A., Magiatis, P., Kaddoumi, A., 2021. Spontaneous In Vitro and In Vivo Interaction of (–)-Oleocanthal with Glycine in Biological Fluids: Novel Pharmacokinetic Markers. *ACS Pharmacol. Transl. Sci.* 4, 179– 192.
- López-Yerena, A., Vallverdú-Queralt, A., Jáuregui, O., Garcia-Sala, X., Lamuela-Raventós, R.M., Escribano-Ferrer, E., 2021. Tissue Distribution of Oleocanthal and Its Metabolites after Oral Ingestion in Rats. *Antioxidants* 10, 688.

* Presenting author. E-mail address: nikoletamadelou@gmail.com (Nikoleta Madelou)

S3.P72 An innovative methodology for the production of a new food supplement enriched in oleocanthal and oleacein with multi target protective properties

Panagiotis Diamantakos, Nikoleta Anna Madelou, Eleni Melliou, Prokopios Magiatis*

Laboratory of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Athens, Greece

Olive oil aldehydic phenols (OOPs: oleocanthal, oleacein, oleuropein aglycone, ligstroside aglycone) are highly bioactive secoiridoids found in olive oil. Nutritional interventions in humans using olive oil with or without OOPs have provided strong evidence about their unique therapeutic role. The most important clinical trials have been performed in patients against chronic lymphocytic leukemia (Rojas et al., 2022), mild cognitive impairment (Tsolaki et al, 2020) and against platelet aggregation (Agrawal et al., 2017). The commercialization of OOPs as food supplements or active pharmaceutical ingredients has been hampered by the lack of economically viable methods for their large-scale isolation or synthesis. Towards this direction, we developed a new, fast and cheap method for their large-scale selective extraction from olive oil using polyethyleneglycol 400. PG400 is a bio-compatible solvent that is not miscible with olive oil and interestingly can react reversibly with olive oil aldehydic phenolics and transform them selectively to hydrosoluble hemiacetals (Fig.1). This unique extraction/reaction procedure has been applied to multi tons scale affording a commercially available extract of OOPs in PEG400 (OLEOPROTECT®). The extract has been formulated to solid or liquid food supplements with excellent stability. When the extract encounters aqueous biological fluids in the human digestive system it spontaneously reacts with water affording 1,1-diols that can further react with plasma or bile aminoacids and enter in the blood circulation. The extract has shown fascinating antitumor, neuroprotective and cardioprotective properties in animal models and currently several clinical trials are in progress.




Fig. 1. Selective extraction/reaction of oleocanthal with PEG400

Keywords: oleocanthal, oleacein, olive oil, food supplement, Oleoprotect

References

- Agrawal, K., Melliou, E., Li, X., Pedersen, T.L., Wang, S.C., Magiatis, P., Newman, J.W., Holt, R.R., 2017. Oleocanthal- rich extra virgin olive oil demonstrates acute anti-platelet effects in healthy men in a randomized trial. *J. Funct. Foods* 36, 84-93
- Rojas Gil, A.P., Kodonis, I., Ioannidis, A., Nomikos, T., Dimopoulos, I., Kosmidis, G., Melliou, E., Magiatis, P., 2022. The Effect of Dietary Intervention With High-Oleocanthal and Oleacein Olive Oil in Patients With Early-Stage Chronic Lymphocytic Leukemia: A Pilot Randomized Trial. *Front. Oncol.* 5746.
- Tsolaki, M., Lazarou, E., Kozori, M., Petridou, N., Tabakis, E., Lazarou, I., Karakota, M., Saoulidis, I., Melliou, E., Magiatis, P., 2020. A Randomized Clinical Trial of Greek High Phenolic Early Harvest Extra Virgin Olive Oil in Mild Cognitive Impairment: The MICOIL Pilot Study. *J. Alzh. Disease* 78.

* Presenting author. E-mail address: magiatis@pharm.uoa.gr (Prokopios Magiatis)

S3.P73 Online dried spot LC-Orbitrap analysis for the quality assessment of cannabis products

Varvara Papaioannou*, Petros Tzimas, Stavros Beteinakis, Leandros A. Skaltsounis, Maria Halabalaki

Division of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimioupoli, Zografos, 15771, Athens, Greece

Dried Blood Spot (DBS) is commonly employed in blood handling and analysis through microsampling on specialized paper cards form. It offers convenient sample collection, storage combined with increased analyte stability and easy transportation to testing laboratories (Tey et al., 2021). To our knowledge, there are no online applications to this day in natural products substrates, where it could confer significant advantages especially when combined with high-resolution MS platforms. This is particularly relevant to Cannabis sativa L. extracts and products based on non- psychoactive cannabinoids like CBD, which have recently inundated the market. A major concern regarding these products is inadequate or misleading labeling about the content of CBD and other cannabinoids, and absence of the psychoactive THC (EMCDDA, 2020). In view of potential risks for consumers, authorities have been prompted to issue regulations. These vary from requiring novel food authorization for CBD edibles in the EU (EU Regulation, 2021) to even prohibition of THC and CBD in food by the FDA (FD&C Act, 2023); however, in the latter, cannabis-derived ingredients without CBD or THC might be allowed in foods and supplements. Consequently, there is a necessity for high-capacity workflows that ensure accurate quantification of cannabinoids in cannabis extracts and end-products, while relieving limitations related to sample collection, delivery, and processing. This study presents an integrated protocol encompassing fully automated DBS-based extraction and LC-IT-Orbitrap MS analysis of cannabis products. The proposed strategy combines robust determination, speed, and cost-effective sample handling, which is particularly suitable for quality control of cannabis-derived materials.

Keywords: Cannabis sativa, DBS, CBD, THC, LC-IT-Orbitrap

References

- EMCDDA (European Monitoring Centre for Drugs and Drug Addiction), 2020. Low-THC cannabis products in Europe. Publications Office of the European Union, Luxembourg.
- FD&C (Federal Food, Drug, and Cosmetic) Act -Regulation of dietary supplement & conventional food products containing cannabis and cannabis derived compounds
- Regulation (EU) 2015/2283 of the European Parliament and of the Council on novel foods
- Tey Hui Yin, Hong Heng See, A review of recent advances in microsampling techniques of biological fluids for therapeutic drug monitoring, Journal of Chromatography A, January 2021, 1635, 461731,

* Presenting author. E-mail address: vpapaioa@pharm.uoa.gr (Varvara Papaioannou)

S3.P74 Anti-prostatic effects of *V. album* L extracts are influenced by season and the use of hydrogel delivery system

Mariana Souza Rocha¹; Sascha D. Markowitsch²; Olesya Vakhrusheva²; Larissa Albrecht²; João VC Batista³; Adriana Passos Oliveira¹; Stephan Baumgartner³; Carla Holandino^{1*}; Roman A. Blaheta²

¹ Multidisciplinary Laboratory of Pharmaceutical Sciences, Faculty of Pharmacy, Universidade Federal do Rio de Janeiro, Av. Carlos Chagas Filho 373, 21941-902 Rio de Janeiro, Brazil

² Department of Urology and Pediatric Urology, University Medical Center Mainz, 55131 Mainz, Germany

³ Society for Cancer Research, Hiscia Institute, Kirschweg 9, 4144 Arlesheim, Arlesheim, Switzerland

Viscum album L. extracts (VA) are widely used in complementary cancer therapy because of its cytotoxicity, immunostimulating, and anti-inflammatory properties (Oei et al., 2019). The cytotoxic activity of a hydrogel containing VA from *Abies alba* host tree (VA-hydrogel) was previously reported by our group (Batista et al., 2022; Rocha et al., 2022). In the present work, the effects of these VA dry extracts from Summer (VAS) and Winter (VAW), as well as VA-hydrogels, were investigated for the first time, in *in vitro* human prostate cancer (PCa) cells.

The following analyses were performed using DU145, PC3, LNCaP prostate cells: cytotoxicity; clonogenic growth; apoptosis and cell cycle progression. The chemical profile of extracts was performed by UV spectrophotometer and thin-layer chromatography.

VAS and VAS-hydrogel presented highest cytotoxic effects, reducing tumor cell growth and colony formation, in a dose dependent manner. The number of PCa cells in early and late apoptosis increased after VAS and VAS-hydrogel treatment. A cell cycle arrest in the S phase was observed in DU145 cells, but not for the other cell lines. The VAW and VAW-hydrogel did not presented significant biological effects. Both extracts (VAS and VAW) showed phytochemical contents in accordance with the pharmacopoeia specifications.

This study highlights the antitumor effect of VAS and VAS-hydrogel on PCa cells. In addition to the importance of harvesting period traceability, the different effects observed in the tested cell lines should be further investigated to elucidate the role of VA and VA-hydrogel in prostate cancer cells.

Keywords: *Viscum album* L., cytotoxic assays, mistletoe, prostate cancer, thermoresponsive hydrogel

References

Oei, S.L., Thronicke, A., Schad, F. 2019. Mistletoe and Immunomodulation: Insights and Implications for Anticancer Therapies. Evidence-Based Complementary and Alternative Medicine 2019.

Batista, J.V.D.C., Matos, A.P.S., Oliveira, A.P., et al. 2020. Thermoresponsive Hydrogel Containing *Viscum album*

Extract for Topic and Transdermal Use: Development, Stability and Cytotoxicity Activity. *Pharmaceutics*. 14, 37. Rocha, M.S., Batista, J.V.C., Melo, M.N.O., Campos, V.E.B., Toledo, A.L.M.M., Oliveira, A.P., Picciani, P.H.S., Baumgartner, S., Holandino, C., 2022. Pluronic® F127 Thermoresponsive *Viscum album* Hydrogel: Physicochemical Features and Cellular *In Vitro* Evaluation. *Pharmaceutics*. 12;14(12):2775.

* Presenting author. E-mail address: cholandino@gmail.com (Carla Holandino)

S3.P75 A “head-to-tail” approach towards foodomics advancement: LC-HRMS and NMR correlation in the case of olive products

Stavros Beteinakis^{a,*}, Anastasia Papachristodoulou^a, Emmanuel Mikros^b, Maria Halabalaki^a

^a Division of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis, Zografou, 15771 Athens, Greece

^b Division of Pharmaceutical Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis, Zografou, 15771 Athens, Greece

Olive derived products, i.e. olive oil, and table olives, comprise sources of high nutritional and economic value, especially for the Mediterranean basin area. Thus, they are prone to adulteration and mislabeling. Conventional approaches commonly adopted by official organizations for their quality assessment, such as the International Olive Council (IOC), are trait-specific and time-consuming. Therefore, there is a need for more holistic approaches, namely metabolite profiling, using advanced analytical instrumentation like NMR and HRMS. Hence, the aim of the present study was to employ both analytical platforms towards the quality and authentication evaluation of Greek EVOOs and table olives, while exploiting statistical tools in the biomarkers' identification process.

Statistical Total Correlation Spectroscopy (STOCSY) and Statistical Heterospectroscopy (SHY) were applied for the first time in the dereplication process of such samples helping with the lack of commercial standards and public databases, especially in the case of NMR spectroscopy. The former generates a pseudospectrum by correlating peaks with the same fluctuation across NMR spectra of the respective samples (Beteinakis et al., 2020, 2023), while the latter detects correlations between spectroscopic and spectrometric data, in this case NMR and HRMS (Crockford et al., 2006). Herein, more than 130 EVOO samples and 60 table olive samples were analyzed and subjected to MVA. Following discrimination based on botanical and geographical origin, tentative biomarkers were detected in each case. The employment of STOCSY and SHY aided in the identification of several biomarkers. Both statistical tools could find further application in foods' profiling approaches.

Keywords: NMR & LC-HRMS profiling, STOCSY, SHY, olive oil, table olives

References

- Beteinakis, S., Papachristodoulou, A., Gogou, G., Katsikis, S., Mikros, E., Halabalaki, M., 2020. NMR-Based Metabolic Profiling of Edible Olives—Determination of Quality Parameters. *Molecules*. 25, 3339.
- Beteinakis, S.; Papachristodoulou, A.; Kolb, P.; Rösch, P.; Schwarzing, S.; Mikros, E.; Halabalaki, M., 2023. NMR- Based Metabolite Profiling and the Application of STOCSY toward the Quality and Authentication Assessment of European EVOOs. *Molecules*. 28, 1738.
- Crockford, D.J., Holmes, E., Lindon, J.C., Plumb, R.S., Zirah, S., Bruce, S.J., Rainville, P., Stumpf, C.L., Nicholson, J.K., 2006. Statistical heterospectroscopy, an approach to the integrated analysis of NMR and UPLC-MS data sets: Application in metabonomic toxicology studies. *Anal. Chem.* 78, 363–371.

* Presenting author. E-mail address: sbeteinakis@pharm.uoa.gr (Stavros Beteinakis)

S3.P76 Systematic cheminformatics analysis of herbal plants in malawi: uncovering chemical scaffolds with anti-candidal potential

King Bwaila^{a,b,*}, John Mponda^a, Ibrahim Chikowe^a

^a Department of Pharmacy, Kamuzu University of Health Sciences, Mahatma Gandhi Str, P/Bag 360, Blantyre, Malawi

^b Department of Pharmacy, Malawi College of Health Sciences, Chewonga Str, P.O Box 122, Zomba, Malawi

Novel medicines for various diseases, including antimicrobial agents against infections have been discovered from herbal plants. However, antifungal drug discovery has received limited attention compared to its antibacterial counterparts (Perfect, 2017). Ethnobotanical surveys remain a conventional approach to collect traditional knowledge on the medicinal uses of herbal plants by local communities (Vogl et al., 2004).

In this study, a literature review on herbal plants species used in Malawi was conducted. A validated algorithm, based on reported uses' similarity to clinical symptoms of candidiasis, identified 133 plant species likely to have anticandidal activities. PubChem queries using identified plants species' names returned 1909 compounds. The 166 Molecular Access System (MACCS) fingerprints and the Murcko chemical scaffolds of the compounds were calculated using RDKit in knime analytics (Khalid et al., 2022). The three most frequent scaffolds were flavanone (19%), piperidine (7%) and (2S)-flavanone (7%) (Yu et al., 2023). Moreover, the 40 most common chemical scaffolds in the dataset of 1909 compounds were screened for their presence in compounds active against *C.albicans* (Fig. 1). This analysis revealed 355 active compounds, a finding which underscores the chemical scaffolds' anti- candidal potential.

Table 1: The three most frequent chemical scaffolds

Chemical Scaffold	Chemical Structure	# Compounds
flavanone		36 (19%)
Piperidine		28 (7%)
(2S)-Flavanone		28 (7%)

ut: 40 chemical scaffolds



Fig. 1: ChEMBL database searching workflow

Chemical Scaffolds screened on the bioactive compounds against *C.albicans* using the Data Warrior module that connects with Chembl database.

In conclusion, future research should focus on bio-guided fractionation, structural elucidation, synthesis, pharmacophore modelling, and toxicity studies on extracts from these plant species.

Keywords: Herbal medicines, antifungal drug discovery, cheminformatics, chemical scaffolds

References

- Khalid, B., Ghorab, H., Benkhemissa, A., 2022. Automated QSPR modeling and data curation of physicochemical properties using KNIME platform: Prediction of partition coefficients. *J. Indian Chem. Soc.* 99, 0019-4522.
- Perfect, J.R., 2017. The antifungal pipeline: a reality check. *Nat. Rev. Drug Disco.* 16, 603–616.
- Vogl, C.R., Vogl-Lukasser, B., Puri, R.K., 2004. Tools and Methods for Data Collection in Ethnobotanical Studies of Homegardens. *Field Methods* 16, 285–306.
- Yu, T., Nantasenamat, C., Kachenton, S., Anuwongcharoen, N., Piacham, T., 2023. Cheminformatic Analysis and Machine Learning Modeling to Investigate Androgen Receptor Antagonists to Combat Prostate Cancer. *ACS Omega* 8, 6729–6742.

* Presenting author. E-mail address: 202440360001@kuhes.ac.mw (King Bwaila)

S3.P77 From nature's palette to drug candidates: exploring the chemodiversity of *Piper fimbriulatum* and expanding the bioactive potential for the development of novel drug leads

Milana Perković^{a,b,*}, Tito Damiani^a, Téó Hebra^a, Marijo Čičak^{a,b}, Tomáš Pluskal^a

^a Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Prague

^b Faculty of Chemistry and Technology, Prague

Piperaceae plants are rich sources of bioactive alkaloids with anticancer properties. *Piper fimbriulatum* is a member of the Piperaceae family from Central America living in symbiosis with *Pheidole bicornis* ants. By exploring *P. fimbriulatum* alkaloid chemodiversity using high resolution mass spectrometry and molecular networking, we discovered that this plant produces piperlongumine, a potent alkaloid that exhibited anticancer activities in numerous preclinical studies. Piperlongumine inhibits the GSTP1 enzyme, causing accumulation of reactive oxygen species and consequently selectively inducing apoptotic and autophagic cell death in cancer cells. Recently, this molecule also demonstrated selective anti-COVID activity in a mouse model. Through the structural annotation of compounds within the piperlongumine cluster in the molecular network representing the plant's chemical diversity, we discovered a diverse array of piperlongumine analogues and cyclobutane dimers produced by the plant. Among those identified, piperlongumine cyclobutane dimer demonstrated *in vitro* activities in the micromolar range against melanoma and breast carcinoma. Afterwards, we performed the synthesis of naturally occurring piperamides for confirmation of their presence in the plant and to investigate UV-light catalyzed 2+2 cycloaddition of these molecules. Our findings revealed that piperamide compounds undergo UV-light induced dimerization forming cyclobutane dimers, which can be synthesized via a straightforward procedure using light irradiation. Furthermore, drawing inspiration from nature's pharmacological treasures, we aim to synthesize new-to-nature analogues and dimers to systematically evaluate their bioactivities, thereby advancing the exploration of plant natural products as potential drug candidates.

Keywords: *Piper fimbriulatum*, alkaloids, piperlongumine, dimers, drug leads

References

- Fisher, R.C., et al., 2002. Plants feed ants: food bodies of myrmecophytic *Piper* and their significance for the interaction with *Pheidole bicornis* ants. *Oecologia* 133, 186-192.
- Harshbarger, W., et al., 2017. Structural and Biochemical Analyses Reveal the Mechanism of Glutathione S-Transferase Pi 1 Inhibition by the Anti-cancer Compound Piperlongumine. *J. Biol. Chem.* 292, 112-120.
- Tang, C. et al., 2023. A Selective SARS-CoV-2 Host-Directed Antiviral Targeting Stress Response to Reactive Oxygen Species. *ACS Cent. Sci.* 9, 109-121.
- Sommerwerk, S. et al., 2016. Synthesis, characterization and cytotoxicity of new pipartine dimers. *Tetrahedron* 72, 1447- 1454.

*Presenting author. E-mail address: milana.perkovic@uochb.cas.cz (Milana Perković)

S3.P78 Inhibition of autotaxin by quercetin and its derivatives

Nam-Hui Yim^{a*}, Jin Yeul Ma^a, Manh Tuan Ha^b, Byung Sun Min^b

^a Korean Medicine (KM) Application Center, Korea Institute of Oriental Medicine, Daegu, Republic of Korea, 41062

^b College of Pharmacy, Drug Research and Development Center, Daegu Catholic University, Gyeongbuk, Republic of Korea, 38430

Autotaxin (ATX), an extracellular enzyme catalyzes the hydrolysis of lysophosphatidylcholin (LPC) to lysophosphatidic acid (LPA), is associated with several disease conditions including inflammation, fibrosis, and cancer (Magkrioti et al., 2019). Quercetin is one of the most potent anti-oxidants of plant origin as a member of flavonoids (Ganbold et al., 2019). In this study, we investigated the ATX inhibitory activities of seven kinds of flavonoids. We selected six flavonoids as quercetin-derived flavonoids (QDF), such as apigenin, kaempferol, azaleatin, rhamnetin, isorhamnetin, and tamarixetin, measured the blockage of ATX secretion by QDF using ELISA. We also determined cellular ROS generation by QDF using DCFDA assay and ATX protein expression change by QDF in vitro. In addition, we performed a protein-ligand docking simulation and examined the hydrophobic interactions and hydrogen bonds formed at the interfaces between the flavonoids and ATX using AutoDock 4.2 software. Four flavonoids, including apigenin, rhamnetin, tamarixetin, and isorhamnetin, showed the inhibition of ATX secretion by more than 30% at maximum concentration (10 μ M). Among them, rhamnetin demonstrated the highest inhibitory effects. In cells, rhamnetin also strongly inhibited ROS generation and ATX protein expression comparing those of quercetin. QDF were further identified for their interaction efficacy with ATX, which supported our results demonstrating the inhibition of QDF against ATX binding interactions. Collectively, our findings suggest that QDF including rhamnetin inhibit ATX activity at a cellular level, which may prove useful as part of an overall modulating the diseases related to ATX.

Keywords: autotaxin, quercetin, quercetin-derived flavonoids (QDF), molecular docking

References

- Magkrioti, C., Galaris A., Kanellopoulou P., Stylianaki E.A., Kaffe E., Aidinis V., 2019. Autotaxin and chronic inflammatory diseases. *J. Autoimm.* 104, 102327.
- Ganbold M., Shimamoto Y., Ferdousi F., Tominaga K., Isoda H., 2019. Antifibrotic effect of methylated quercetin derivatives on TGF β -induced hepatic stellate cells. *BBRC.* 20, 100678.

* Presenting author. E-mail address: nhyim23@kiom.re.kr (Nam-Hui Yim)

S3.P79 Exploring vibrational spectroscopic approaches for quantifying main phytocannabinoids in medicinal *Cannabis sativa* inflorescence

Olga Gigopulu^{a,*}, Gjoshe Stefkov^a, Nikola Geshkovski^a, Petre Makreski^b

^a Faculty of Pharmacy, Ss. Cyril and Methodius University in Skopje, Majka Tereza 47, 1000 Skopje, North Macedonia

^b Institute of Chemistry, Faculty of Natural Sciences and Mathematics-Skopje, Ss. Cyril and Methodius University in Skopje, Arhimedova 5, 1000 Skopje, North Macedonia

The exponential increase of medical cannabis consumption has resulted in higher demand for its effective large-scale cultivation and manufacturing. Additionally, quality criteria must be continually assessed throughout the manufacturing process since the growing procedures have a significant impact on the variability of phytocannabinoid content in plant material. Rapid technological advancements in recent years have resulted in a notable surge in the application of vibrational spectroscopic techniques, including Raman, mid-infrared (MIR), and near-infrared (NIR) spectroscopy, in conjunction with chemometric analysis, as highly effective for industrial research, process monitoring, and quality control (Deidda et al., 2022).

Thus, the goal of our study was to evaluate and compare these techniques for quantitative measurement of the main phytocannabinoids, tetrahydrocannabinol (THC) and cannabidiol (CBD), in medical cannabis flowers in order to further explore their potential for application as process analytical technology (PAT) in the medical cannabis industry.

MIR, NIR, and Raman spectra were obtained for large sets of dried cannabis flower, accompanied with HPLC determined content of total THC ($\text{THC\%} + 0.877 \cdot \text{THCA\%}$) and total CBD ($\text{CBD\%} + 0.877 \cdot \text{CBDA\%}$). Partial least-squares (PLS) analysis was carried out on the collected spectra in order to create calibration models for the determination of THC and CBD content in cannabis flowers.

Based on the acquired results, vibrational spectroscopic methods should be considered as a promising PAT tool for rapid and non-destructive estimation of THC and CBD in the medical cannabis industry.

Keywords: THC, CBD, spectroscopy, medical cannabis, PAT

References

Deidda, R., Dispas, A., De Bleye, C., Hubert, P., Ziemons, É., 2022. Critical review on recent trends in cannabinoid determination on cannabis herbal samples: From chromatographic to vibrational spectroscopic techniques. *Anal. Chim. Acta* 1209, 339184.

* Presenting author. E-mail address: ogigopulu@gmail.com (Olga Gigopulu)

S3.P80 Exploring the therapeutic synergy: comprehensive assays unveil the potential of sericin and *Chelidonium majus* L. in developing a possible wound dressing

Ana Borges^{1,2,3}, Sandrina A. Heleno^{1,2,*}, Sónia P. Miguel⁴, Maximiano P. Ribeiro⁴, Paula Coutinho⁴, André F. Moreira⁴, María Luisa Martín Calvo⁵, Josiana A. Vaz^{1,2}, Lillian Barros^{1,2}, Ricardo C. Calhella^{1,2}

¹ Centro de Investigação da Montanha (CIMO), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

² Laboratório Associado para a Sustentabilidade e Tecnologia em Regiões de Montanha (SusTEC), Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

³ Grupo de Investigación en Desarrollo y Evaluación de Formas Farmacéuticas y Sistemas de Liberación Controlada, Facultad de Farmacia, Universidad de Salamanca, Campus Miguel de Unamuno s/n, 37007 Salamanca, Espanha

⁴ CPIRN-IPG—Center of Potential and Innovation of Natural Resources, Polytechnic Institute of Guarda, 6300-559 Guarda, Portugal

⁵ Grupo de Investigación en Fisiología y Farmacología, Facultad de Farmacia, Universidad de Salamanca, Campus Miguel de Unamuno s/n, 37007 Salamanca, Espanha

Termed as a “silent epidemic,” wounds exert a significant social and economic toll, detrimentally affecting the quality of life for millions worldwide. Natural substances like Sericin and *Chelidonium majus* L. (*C. majus*) have shown potential in wound care (1–4). Sericin, a silk protein, accelerates wound healing by promoting cell proliferation, collagen synthesis, and angiogenesis (5–7). On the other hand, *C. majus*, a medicinal plant, has anti-inflammatory effects and supports tissue regeneration (8–11). Combining these substances in wound dressings could address multiple aspects of the healing process. This investigation aimed to assess the potential synergy between Sericin and *C. majus* in the context of wound healing. Cytotoxicity, anti-inflammatory, antimicrobial, and wound healing assays were conducted to observe the healing rates under distinct concentrations (1:1, 1:2, 2:2, and 2:1 ratios) of the two extracts combined. These extracts, in all ratios, included various combinations of *C. majus* and Sericin sourced from Bragança, Castelo Branco, and two commercial types. Through these assays, valuable insights were gained into the collaborative influence of Sericin and *C. majus* on the wound healing process. Across all assays, results varied among the different combinations tested. However, the combination that consistently stood out across multiple assays was the 2:2 ratio with Sericin sourced from Castelo Branco. The exploration of these natural substances as constituents of wound dressings presents a hopeful path for crafting sustainable, efficacious, and biocompatible materials, that could significantly impact the field of wound healing.

Keywords: wound healing, biomaterial, dressing, sericin, greater celandine

Acknowledgements

FCT, Portugal for financial support to CIMO (UIDB/00690/2020 and UIDP/00690/2020) and SusTEC (LA/P/0007/2021); national funding by FCT, P.I., through the institutional scientific employment program-contract for S.A. Heleno and RC Calhella, and through the project Wastesilk (<http://doi.org/10.54499/PTDC/BTA-BTA/0696/2020>).

* Presenting author. E-mail address: sheleno@ipb.pt (Sandrina A. Heleno)

S3.P81 Synergistic impact of natural stilbene-linked mono-substituted nitroarenes in hormone- dependent breast cancer: a greener approach

Lairikyengbam Deepti Roy*, Jyotsna Kumar

Department of Chemistry, Faculty of Mathematical and Physical Sciences, M.S. Ramaiah University of Applied Sciences, Bangalore-560058, Karnataka, India

Resveratrol, a natural stilbene, is a promising multi-target drug with diverse biological effects. However, its low bioavailability has led researchers to develop novel derivatives to address pharmacokinetic issues (Turcov et al., 2022). In the present study, a series of nitroarene substituted resveratrol analogues (NSRAs) were synthesized using resveratrol isolated from green grapes to study their anti-breast cancer properties on a panel of hormone-dependent (HR+) and hormone-independent (HR-) breast cancer (BC) cells. Structures of NSRAs were elucidated by ^1H -NMR, ^{13}C -NMR, FT-IR, and HPLC/MS (Krishnamurthy et al., 2023).

The cell viability of both HR+ T-47D and HR- MDA-MB-157 decreased in a dose and time- dependent manner when treated with isolated resveratrol and NSRAs, individually. Interestingly, NSRA-I demonstrated augmented and selective antiproliferative activity against HR+ T-47D cells ($\text{IC}_{50} = 17.92 \pm 0.072 \mu\text{M}$) compared to isolated resveratrol ($\text{IC}_{50} = 27.26 \pm 0.087 \mu\text{M}$). Further investigations unveiled that both the NSRAs induced apoptosis in T-47D cells, leading to cell cycle arrest at the G_0/G_1 phase after 48 hours of incubation.

The study emphasises the curative potential of the synthesised novel resveratrol analogues, particularly NSRA-I, establishing it as a potential scaffold to design and develop novel stilbene analogues with improved therapeutic effectiveness and selectivity towards HR+ breast cancer cells. Additionally, the compound's antiproliferative capabilities, combined with its ability to cause programmed cell death along with cell cycle arrest, nominate it as a promising therapeutic candidate. Advancing research and uncovering potent inhibitors, utilizing resveratrol as a precursor for HR+ breast cancer with enhanced efficacy, remains a viable direction. Ongoing investigations are currently underway to delve deeper into these possibilities.

Keywords: stilbene, resveratrol, nitroarenes, MTT, HR+ breast cancer

References

- Krishnamurthy, G., Roy, L.D., Kumar, J., Gour, P., Arland, S.E., Prabu, M., Rahman, N., Srinivasa, G.R., Shreenivas, M.T., 2023. Design, preparation, and *in silico* study of novel curcumin-biphenyl carbonitrile conjugate as novel anticancer drug molecules. *Int. J. Appl. Pharm.* 15, 143–159.
- Turcov, D., Barna, S., Profire, L., Iacob, A.T., Lişă, G., Puiţel, A.C., Zbranca, A., Suteu, D., 2022. Physico-chemical characterization of the antioxidant mixture resveratrol-ferulic acid for applications in dermato-cosmetic products. *Farmacia* 70, 410-416.

* Presenting author. E-mail address: deepti07roy@gmail.com (Lairikyengbam Deepti Roy)

S3.P82 The natural diversity of acyltransferases reveals versatility and specificity in the synthesis of gene-encoded lipopeptides

Johannes A. Eckert^{a,*}, Nina M. Bösch^a, Florian Hubrich^a, Serina L. Robinson^b, Jörn Piel^a, Anna L. Vagstad^a

^a Eidgenössische Technische Hochschule (ETH) Zürich, Institute of Microbiology, Vladimir-Prelog- Weg 4, 8093 Zürich, Switzerland

^b Swiss Federal Institute of Aquatic Science and Technology (Eawag), Department of Environmental Microbiology, Überlandstrasse 133, 8600 Dübendorf, Switzerland

Lipopeptides comprise biochemical compounds that can be naturally produced or chemically synthesized. They exhibit diverse bioactive functions and are widely used in health care (Hamley, 2015). Despite their versatility, the production and diversification of lipopeptide compounds continue to be a challenge for state-of-the-art biological and chemical synthesis. In contrast, we offer a more streamlined and adaptable route to lipopeptide natural products via ribosomally synthesized and post-translationally modified peptides (RiPPs). Their primary structure is determined by a genetically encoded precursor protein, which is modified by maturases within the same biosynthetic gene cluster (Montalbán-López et al., 2021). The crucial lipidation reaction is catalyzed by a novel class of maturases of the GCN5-related *N*-acetyltransferase (GNAT) family, which introduce medium-chain fatty acids onto the sidechain amino group of lysines or arginine-derived ornithines (Hubrich et al., 2022) (Fig. 1).




Fig. 1. Transfer of fatty acids to the sidechain of ornithines and lysines by RiPP GNATs.

Here we present the natural diversity of this GNAT family and their corresponding gene-encoded lipopeptides by characterizing the substrate scope of selected GNATs with their cognate precursor peptides. We could show that a variety of fatty acids can be introduced including chain lengths from C10 to C18. We are currently exploring this enzyme class to enable custom peptide engineering efforts towards generating bioactive lipopeptide compounds. We envision this platform to be used to facilitate the production of crucial lipopeptide compounds for human health and to quickly screen for antimicrobial lipopeptide compounds and non-ribosomal lipopeptide mimics.

Keywords: GNAT, RiPPs, lipopeptides, synthetic biology

References

- Hamley, I. W., 2015. Lipopeptides: from self-assembly to bioactivity. *Chem. Comm.* 51 (41), 8574-8583.
- Montalbán-López, M., et al., 2021. New developments in RiPP discovery, enzymology and engineering. *Nat. Prod. Rep.* 38, 130-239.
- Hubrich, F. & Bösch, N. M., et al. 2022. Ribosomally derived lipopeptides containing distinct fatty acyl moieties. *PNAS* 119 (3), e2113120119.

* Presenting author. E-mail address: johannes.eckert@biol.ethz.ch (Johannes Eckert)

S3.P83 Exploring new dimensions: analytical method development for enhanced quality control of natural products

Justine Raeber*, Christian Steuer

Institute of Pharmaceutical Sciences, Department of Chemistry and Applied Biosciences, ETH Zurich, 8093 Zurich, Switzerland

Natural products are substances produced by living organisms. They serve as a valuable source for new leads in drug discovery and development and are widely used in medicine and phytotherapy as isolated plant compounds or as crude extracts. (Rodrigues et al., 2016) Natural products contain secondary metabolites that are not essential for basic life but play crucial roles, such as deterring predators and attracting pollinators. Natural products can occur as complex mixtures that vary in composition depending on the species, growing location, genetics and/or harvest time. Terpenes are the largest class of phytochemicals. They consist of isoprene units, can be linear or cyclic and can be oxygenated. The small subtleties in the carbon backbone give rise to the diversity of terpenes. (Raeber and Steuer, 2023; Hüsni et al., 2007) However, these subtleties pose a challenge for analysis and structure determination. Terpenes may exist as enantiomers, exhibit further isomer structures, share identical masses depending on the terpene class, and have similar fragmentation patterns when analysed by mass spectrometry.

Consequently, the exact composition of terpenes can influence the pharmacological efficacy of plants extracts. In this study, we applied three strategies to enhance the current analysis of terpene patterns in plants. These strategies range from the analysis of enantiomeric ratios, to the training and testing of classification algorithms using machine learning, and the examination of adduct patterns using ambient ionisation methods followed by mass spectrometric analysis. (Raeber et al., 2023; Raeber and Steuer, 2023) These strategies are designed to overcome the existing limitations of terpene analysis and enable in-depth characterisation of plant extracts.

Keywords: chemometrics, terpenes, ambient ionization, classification, quality control

References

- Hüsni K, Başer C, Demirci F (2007) Chemistry of essential oils. In: Flavours and fragrances: chemistry, bioprocessing and sustainability. Springer, pp 43-86
- Raeber J, Favrod S, Steuer C (2023) Determination of Major, Minor and Chiral Components as Quality and Authenticity Markers of Rosa damascena Oil by GC-FID. *Plants* 12 (3):506
- Raeber J, Steuer C (2023) Exploring new dimensions: Single and multi-block analysis of essential oils using DBDI-MS and FT-IR for enhanced authenticity control. *Analytica Chimica Acta* 1277:341657
- Rodrigues T, Reker D, Schneider P, Schneider G (2016) Counting on natural products for drug design. *Nature Chemistry* 8 (6):531-541. doi:10.1038/nchem.2479

* Presenting author. E-mail address: justine.raeber@pharma.ethz.ch (Justine Raeber)

S3.P84 Simultaneous quantification of terpenes and cannabinoids in *Cannabis sativa* by reversed phase LC-APCI-MS/MS

Justine Raeber^{a,*}, Michael Poetzsch^b, Anina Schmidli^a, Sina Favrod^a, Ana Curavic^a, Christian Steuer^a

^a Institute of Pharmaceutical Sciences, Department of Chemistry and Applied Biosciences, ETH Zurich, 8093 Zurich, Switzerland

^b Swiss Drug Testing GmbH, Technoparkstrasse 2, 8406 Winterthur, Switzerland

In recent years, *Cannabis sativa* L. has undergone a significant paradigm shift, transitioning from a prohibited drug to being recognized as a medicinal candidate for a wide range of diseases. This change has also led to changes in legal regulations in Europe and other countries (Pattnaik *et al.*, 2022). As a consequence, the plant was included in various pharmacopoeias, such as the Pharmacopeia Europea 11.5. (Ph. Eur.). The Ph. Eur. distinguishes between different chemotypes of *C. sativa* based on their ratio of THC to CBD (Council of Europe, 2023). While widely accepted, this classification method is overly simplistic and does not do justice to the complex chemical composition of *C. sativa*. Other minor cannabinoids are of increasing interest, and interactions between cannabinoids (terpenophenols) and terpenes have been proposed, which may influence pharmacological effects (Russo, 2011).

Current state of the art analytical techniques favour liquid chromatography (LC) for cannabinoid analysis, as the cannabinoid acid forms are preserved. Gas chromatography (GC) is preferred for volatile compounds such as terpenes and can also be used for simultaneous cannabinoid analysis. However, this approach requires prior derivatization of cannabinoid acids, a timely process suffering from incomplete reactions (Zivovinic *et al.*, 2018).

In this study, we present the first method to our knowledge that combines terpene and cannabinoid analysis in a single LC-MS/MS method. This method allows the quantification of both terpenes and cannabinoids without the need for prior derivatization of their acids. This technique offers a fast, reliable and holistic approach that allows a deeper understanding of chemotype classification.

Keywords: chemotype, classification, LC-MS/MS, ambient ionization, *Cannabis sativa*

References

- Council of Europe, 2023. Cannabis flower. European Pharmacopeia (Vol. 11.5.). Strasbourg.
- Pattnaik, F., Nanda, S., Mohanty, S., Dalai, A. K., Kumar, V., Ponnusamy, S. K., & Naik, S., 2022. Cannabis: Chemistry, extraction and therapeutic applications. *Chemosphere*, 289, 133012.
- Russo, E. B., 2011. Taming THC: potential cannabis synergy and phytocannabinoid-terpenoid entourage effects. *British Journal of Pharmacology*, 163(7), 1344-1364.
- Zivovinic, S., Alder, R., Allenspach, M. D., & Steuer, C., 2018. Determination of cannabinoids in *Cannabis sativa* L. samples for recreational, medical, and forensic purposes by reversed-phase liquid chromatography-ultraviolet detection. *Journal of Analytical Science and Technology*, 9(1), 27.

* Presenting author. E-mail address: justine.raeber@pharma.ethz.ch (Justine Raeber)

S3.P85 **LumiNose: a novel instrument to apply the receptomics biosensor technology for identifying and quantifying medicinal bioactivities in plant extracts**

Lorenzo Corsico^{a,*}, Maurice G.L. Henquet^{b,*}, Aria Samimi^a, Margriet Roelse^b, Maarten A. Jongsma^b

^a *InsectSense, Plus Ultra II Building, Bronland 10, 6708 WH Wageningen, The Netherlands*

^b *BU Bioscience, WageningenUR, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands*

G-protein coupled receptors (GPCRs) are important targets for various plant and food components. The members of this large family of membrane proteins are involved in virtually every physiological process. By analysing the interactions between food components and receptors in the body, food scientists can determine the health potential of a particular food product. LumiNose, a powerful novel biosensor platform based on the Receptomics technology developed at Wageningen Research, mimics the exceptional sensing mechanism of insects for detection purposes by using insect olfactory receptors. Previous research by Wageningen Research has already demonstrated the technology to work with various human taste and health-related GPCRs and ion channels. This allows for the analysis of plant extract interactions with receptors in the human body, and the identification and quantification of the specific bioactive compounds. The implementation of LumiNose holds the potential to revolutionize the industry, offering more efficient and effective methods of discovery and product monitoring. In this regard, plant extract bioactivity has been detected for a range of different receptors, including satiety (GLP-1), dopamine, serotonin, and free fatty acid receptors. This information can also be used to optimize product formulations and improve bioavailability.

Keywords: GPCR, receptomics, bioactivity, health, food

References

- Henquet, M.G.L., Roelse, M., de Vos, R.C.H., Schipper, A., Polder, G., Verhoeven, H.A., de Ruijter, N.C.A., Jongsma, M.A. 2016. Metabolomics meets functional assays: coupling LCMS and microfluidic cell-based receptor-ligand analyses. *Metabolomics*. 12: 1-13.
- Roelse, M., Henquet, M.G.L., Jongsma, M.A. 2021. Receptomics: Tongue-on-a-chip with novel opportunities for food screening. *Proc. of the 16th Weurman Flavour Research Symposium*.
- Roelse, M., Henquet, M.G.L., Verhoeven, H.A., de Ruijter, N.C.A., Wehrens, R., van Lenthe, M.S., Witkamp, R.F., Hall, R.D., Jongsma, M.A. 2018. Calcium imaging of GPCR activation using arrays of reverse transfected HEK293 cells in a microfluidic system. *Sensors*. 18, 602.
- Roelse, M., Wehrens, R., Henquet, M.G.L., Witkamp, R.F., Hall, R.D., Jongsma, M.A. 2019. The effect of calcium buffering and calcium sensor type on the sensitivity of an array-based bitter receptor screening assay. *Chemical Senses*. 44, 497-505.
- Wehrens, R., Roelse, M., Henquet, M.G.L., van Lenthe, M., Goedhart, P., Jongsma, M.A. 2019. Statistical models discriminating between samples measured with microfluidic receptor cell arrays. *PLOS One*. 14(4): e0214878.

* Presenting author. E-mail address: lorenzo@insectsense.com (Lorenzo Corsico), maurice.henquet@wur.nl (Maurice Henquet)

S3.P86 pH-Dependent degradation of andrographolide; kinetics, product identification and anti-inflammatory and cytotoxic activity assessment

Wuttichai Jaidee^{a,*}, Narawadee Rujanapun^a, Kulawadee Malee^a, Geoffrey A. Cordell^b, Gudun Ulrich-Merzenich^c, Rawiwan Charoensup^a

^a Medicinal Plant Innovation Center of Mae Fah Luang University, Mae Fah Luang University, Chiang Rai 57100, Thailand

^b Natural Products Inc., Evanston, IL 60201, USA; Department of Pharmaceutics, College of Pharmacy, University of Florida, Gainesville, FL, 32610, USA

^c University Hospital Bonn (UKB), Medical Clinic III, AG Synergy Research and Experimental Medicine, Bonn, Germany

Andrographolide, the abundant labdane-type diterpene lactone in the leaves of *Andrographis paniculata* (Burm. f.) Wall. (Acanthaceae), is increasingly under investigation for its biological properties. Consequently, the kinetics of the chemical reactions of andrographolide under acid and basic condition, the identification of the degradation products, and their preliminary biological assessment, are of interest. The thermal degradation kinetics of andrographolide at pH 2 (70, 77, and 85 °C), 6 (60, 70, and 80 °C), and 8 (50, 60, and 70 °C) were determined. The calculated values of the activation energies (E_a), shelf-life ($t_{90\%}$), and rate constant (k) were determined. The effect of pH on andrographolide was assessed using the Arrhenius equation. First-order reaction kinetics were fitted to the data using linear regression analysis to determine the reaction rate constant at each pH and temperature combination which increased with increasing temperature and pH. The optimum pH for andrographolide stability was between pH 2 and 4.




Fig. 1. Structures of andrographolide and its degradation products formed under acidic (AGA-1-2) and basic (AGB-1-3) conditions

The major degradation products in pH 2 and 6 solutions were isolated and characterized by NMR spectral analysis. Two degradation products, isoandrographolide (AGA-1) (Lui et al., 2019) and 8,9-didehydroandrographolide (AGA-2) were formed under acidic conditions, whereas the three degradation products generated under basic conditions (pH 8) were 15-seco-andrographolide (AGB-1), 14-deoxy-15-methoxy-andrographolide (AGB-2) (Gui-Yang, et al., 2017) and 14-deoxy-11,12-didehydroandrographolide (AGB-3) (Fig. 1) (Phattanawasin et al., 2018). The anti-inflammatory and cytotoxic activities of the degradation products were also evaluated.

Keywords: Andrographolide, pH-dependency, degradation products, structure determination, formation kinetics

References

- Gui-Yang, W., Ting, W., Fei-Fei, L., Hai-Yan, T., Chun-Lin, F., Xiao-Jun, H., Wen-Cai, Y., Ying, W. 2017. Two new diterpenoid lactones isolated from *Andrographis paniculata*. *Chin. J. Nat. Med.*, 2017, 15(6): 458-462.
- Kumar, S., Singh, B., Bajpai, V. 2021. *Andrographis paniculata* (Burm.f.) Nees: Traditional uses, phytochemistry, pharmacological properties and quality control/quality assurance. *J. Ethnopharmacol.* 275, 114054.
- Liu, L., Yan, Y., Zheng, L., Jia, H., Han, G. 2020. Synthesis and structure anti-inflammatory activity relationships studies of andrographolide derivatives. *Nat. Prod. Res.* 34(6):782-789.
- Phattanawasin, P., Sotanaphun, U., Burana-Osot, J., Piyapolrungrroj, N. 2018. Isolation and characterization of the acid and base degradation products of andrographolide. *Pharmazie* 73, 559-562.

* Presenting author. E-mail address: wuttichai.jai@mfu.ac.th (Wuttichai Jaidee)

S3.P87 Immunohistochemical evaluation of the effect of *Kigelia africana* stem-bark extracts on benign prostatic hyperplasia induced by testosterone and estradiol in albino Wistar rats

Mbaka GO¹, Olubi OO², Oyejide JO¹, Osiagwu D³

¹ Dept. of Anatomy, Lagos State University College of Medicine, Ikeja, Lagos State. Postal Code - 234, Nigeria

² Dept. of Ear, Nose and Throat Surgery, Lagos State University College of Medicine, Ikeja, Lagos State. Postal Code - 234, Nigeria

³ Dept. of Anatomic and Molecular Pathology, College of Medicine of the University of Lagos, Idi- Araba. Postal Code - 234, Nigeria

Stem-bark extract of *Kigelia africana* (KA) is traditionally used for treating various health conditions including benign prostatic hyperplasia (BPH). However, its use for treating BPH has not been validated.

Male rats weighing above 200g body weight (bw) had exogenous administration of testosterone and estradiol in staggered doses (three times weekly) for three weeks. Induced animals were divided into five groups of 3 per group. Group 1 was positive control while the induced animals were distributed between groups 2 to 6. Group 2 was untreated (negative control). Group 3 (prophylactic study) received by gavages the extract (200 mg/kg bw) simultaneously with the steroid administration for thirty days. Groups 4 and 5 received extract at 100 and 200 mg/kg respectively by gavages for thirty days while group 6 received finasteride (0.1 mg/kg). The assays conducted include dihydrotestosterone level (DHT) and prostate-specific antigen (PSA). Immunohistochemical (IHC) techniques were performed on the sections using primary antibodies TNF- α , a potent pro- inflammatory cytokine involved in wide spectrum of activity including cell proliferation.

The extract exerted marked decrease in the prostate weight (0.23 g) compared to the untreated with average weight of 0.70 g. Dihydrotestosterone level showed marked decrease in the extract treated (1352 ng/L) compared to the untreated with 1913 ng/L. IHC staining showed high intensity of the marker expression in the negative control but decreased very significantly with dose in the extract treated group.

The extract treated showed decrease in DHT and prostate weight with the IHC staining indicating less intensity.

* Presenting author. E-mail address: mbaakagm@yahoo.com (Godwin Mbaka)

S3.P88 Styryl lactone derivatives and aristolactam alkaloids from *Goniothalamus tapis* Miq. and their α -glucosidase inhibitory activity

Piyaporn Phukhatmuen^a, Passakorn Teerapongpisan^b, Virayu Suthiphasilp^c, Tharakorn Maneerat^{b,d}, Thidarat Duangyod^{a,d}, Rawiwan Charoensup^{a,d}, Sarot Cheenpracha^e, Jin-Lin Zhu^f, Yan Alexander Wang^f, Suwanna Deachathai^b, Raymond J. Andersen^{f,g}, Surat Laphookhieo^{b,d}

^a School of Integrative Medicine, Mae Fah luang university, Chiang rai, Thailand

^b Center of chemical Innovation for Sustainability (CIS) and School of Science, Mae Fah Luang University, Chiang rai, Thailand

^c Department of Industrial Technology and Innovation Management, Faculty of Science and Technology, Pathumwan Institute of Technology, Bangkok, Thailand

^d Medicinal Plants Innovation Center of Mae Fah luang university, Chiang rai, Thailand ^e Division of Chemistry, School of Science, University of Phayao, Phayao, Thailand

^f Department of Chemistry, University of British Columbia, Vancouver, BC, Canada

^g Department of Earth, Ocean & Atmospheric Sciences, University of British Columbia, Vancouver, BC, Canada

Two recently discovered styryl lactone derivatives, namely goniothapic acids A (**1**) and B (**2**), along with 18 previously identified compounds, were extracted from the twig and leaf samples of *Goniothalamus tapis* Miq. The new compounds' structures were determined through spectroscopic methods and high resolution electrospray ionization time-of-flight mass spectrometry (HRESITOFMS), and their absolute configuration was confirmed by comparing experimental and calculated electronic circular dichroism (ECD) spectra. The α -glucosidase inhibitory activity of eleven compounds was assessed, revealing that (–)- goniothalamin (**5**) and oldhamactam (**16**) exhibited the most potent inhibition with IC₅₀ values of 54.8 and 57.9 μ M, respectively (Phukhatmuen et al., 2023).



Fig. 1. Structure of goniothapic acids A (**1**) and B (**2**)

Keywords: *Goniothalamus tapis*, Annonaceae, styryl lactones, aristolactam alkaloids, α -glucosidase inhibitory activity

References

Phukhatmuen, P., Teerapongpisan, P., Suthiphasilp, V., Maneerat, T., Duangyod, T., Charoensup, R., Cheenpracha, S., Zhu, J.L., Wang, Y.A., Deachathai, S., Andersen, R.J., Laphookhieo, S., 2024. Styryl lactone derivatives and aristolactam alkaloids from *Goniothalamus tapis* Miq. and their α -glucosidase inhibitory activity. Nat.Prod.Res.1-10.

* Presenting author. E-mail address: piyaporn.phu@mfu.ac.th (Piyaporn Phukhatmuen)

S3.P89 Exploring synergies between hemp cannabinoids and β -caryophyllene on triple-negative breast cancer cell line

Carla Hamann^{a,*}, Olivia Jansen^a, Kateline Jullien^a, Allison Ledoux^a, Erik Maquoi^b, Michel Frédérick^a

^a Laboratory of Pharmacognosy (CIRM), University of Liege, Avenue Hippocrate 15, 4000 Liege, Belgium

^b Laboratory of Tumour and Development Biology (GIGA-Cancer), University of Liege, Avenue Hippocrate 15, 4000 Liege, Belgium

Hemp, also known as *Cannabis sativa*, contains over 500 compounds that include cannabinoids and terpenes. Some noteworthy studies have highlighted that cannabis inflorescence extracts were more active than the single purified cannabinoids on breast cancer cell lines (Blasco-Benito et al., 2018). Furthermore, the interplay between cannabinoids and terpenes has been emphasised, with β -caryophyllene, the major sesquiterpene, emerging as an agonist of the CB2 receptor, which is part of the endocannabinoid system.

Our study aims to determine the anticancer properties of combinations of hemp components and contributes to the understanding of the synergies in hemp. Neutral/acidic cannabinoids and β -caryophyllene were combined and tested in triplicate against a human triple-negative breast cancer cell line (MDA-MB-231). Cell proliferation was quantified kinetically for 3 days using an Incucyte SX-5 live-cell imaging system (Sartorius) and at a single time point (72 hours) with a resazurin-based metabolic assay. The SynergyFinder Plus web application was used for analysing and visualising multi-drug combination response data (Zheng et al., 2022) (Fig. 1).

The highest synergy score was obtained with cannabidiol and β -caryophyllene, highlighting the potential of a cannabis extract rich in these compounds. Nonetheless, the presence of cannabinoid acids in the extract is undesirable, as our results showed the lowest synergy score, indicative of an antagonistic interaction. These findings not only contribute to the understanding of the entourage effect in hemp but also guide the formulation of extracts that harness the full synergistic power of *Cannabis sativa* for enhanced therapeutic efficacy.




Fig. 1. Experimental flowchart

Keywords: *Cannabis sativa*, synergy, breast cancer, cannabinoids, β -caryophyllene

References

- Blasco-Benito, S., Seijo-Vila, M., Caro-Villalobos, M., Tundidor, I., Andradás, C., García-Taboada, E., Wade, J., Smith, S., Guzmán, M., Pérez-Gómez, E., Gordon, M., Sánchez, C., 2018. Appraising the “entourage effect”: Antitumor action of a pure cannabinoid versus a botanical drug preparation in preclinical models of breast cancer. *Biochem Pharmacol* 157, 285–293.
- Zheng, S., Wang, W., Aldahdooh, J., Malyutina, A., Shadbahr, T., Tanoli, Z., Pessia, A., Tang, J., 2022. SynergyFinder Plus: Toward Better Interpretation and Annotation of Drug Combination Screening Datasets. *Genomics, Proteomics & Bioinformatics* 20, 587–596.

* Presenting author. E-mail address: chamann@uliege.be (Carla Hamann)

S3.P90 Response Surface Methodology as a tool to obtain a potential satiating bitter enriched extract from *Gentiana lutea* L.

Vittorio Carlucci^{a,*}, Maria Ponticelli^a, Ludovica Lela^a, Claudia Mangieri^a, Federica Santopietro^a, Immacolata Faraone^a, Daniela Russo^a, Luigi Milella^a

Department of Science, Università degli Studi della Basilicata, Via dell'Ateneo Lucano 10, 85100 Potenza, Italy

In recent years, there has been a growing interest in bitter compounds due to their ability to regulate energy intake and hunger (Rezaie et al., 2021). Obesity is becoming a real challenge as current treatments are very limited with associated adverse effects. Nature offers several bitter compounds and in particular, *Gentiana lutea* L. roots represent a rich source of bitter secoiridoid glycosides like gentiopicrin or amarogentin, the most bitter natural compound (Ponticelli et al., 2023). Bitter compounds are able to interact with bitter taste receptors (*Tas2rs*), highly related to body weight regulation (Mennella et al., 2016).

Therefore, the aim of this work is to optimize the extraction parameters to recover bitter compounds from gentian roots and identify the molecular mechanism in the secretion of hormones that reduce the feeling of hunger in STC-1 cells. To obtain an extract rich in bitter compounds, three different parameters (time, solvent, and temperature) were considered using the Response Surface Methodology.

The optimal extractive conditions were 85 min, 95% EtOH, and 75°C. The main bitter compounds have been identified and quantified by HPLC-DAD. Treatment of STC-1 cells with optimized gentian extract increased the release of the incretins GLP-1 and CCK. These effects are mediated by the variation in intracellular calcium concentrations following the upregulation of the *Tas2r138* receptor or of other genes involved in the bitter taste transduction mechanism including *Plcβ2*, *Ip3r3* or *Trpm5*. Furthermore, molecular docking studies have allowed us to define the interactions between the main bitter compounds and the *Tas2r138* receptor.

Keywords: *Gentiana lutea* L., bitter compounds, Response Surface Methodology, *Tas2r138*, obesity

References

- Mennella, I., Fogliano, V., Ferracane, R., Arlorio, M., Pattarino, F., Vitaglione, P., 2016. Microencapsulated bitter compounds (from *Gentiana lutea*) reduce daily energy intakes in humans. *Br. J. Nutr.* 116, 1841-1850.
- Ponticelli, M., Lela, L., Moles, M., Mangieri, C., Bisaccia, D., Faraone, I., Falabella, R., Milella, L., 2023. The healing bitterness of *Gentiana lutea* L., phytochemistry and biological activities: A systematic review. *Phytochem.* 206, 113518.
- Rezaie, P., Bitarafan, V., Horowitz, M., Feinle-Bisset, C., 2021. Effects of Bitter Substances on GI Function, Energy Intake and Glycaemia-Do Preclinical Findings Translate to Outcomes in Humans?. *Nutrients.* 13, 1317.

* Presenting author. E-mail address: vittorio.carlucci@unibas.it (Vittorio Carlucci)

S3.P91 The Botanical Safety Consortium: collaborative effort to improve botanical safety methods

Connie Mitchell^a, Michelle Embry^a, Julie Krzykwa^a, Amy Roe^b, Olaf Kelber^c

^a Health and Environmental Sciences Institute, Washington, DC, USA

^b Procter & Gamble Company, Cincinnati, OH, USA; 3. R&D, Phytomedicines Supply and Development Center, Bayer Consumer Health, Steigerwald Arzneimittelwerk GmbH, Darmstadt, Germany

The use of natural health products, including botanical dietary supplements, is increasing, making their safety a crucial issue for public health. The Botanical Safety Consortium (BSC), has been formed through a Memorandum of Understanding as a collaborative effort between the US FDA, the National Institute of Environmental Health Sciences, and the Health and Environmental Sciences Institute. It aims to improve safety evaluation methods for botanicals, operating as an international platform where experts from various sectors, including government, academia, industry, and non-profits, collaborate to advance and apply new approach methodologies (NAMs) for the safety assessment of botanical products (1, 2).

The BSC is 1) collaborating with global stakeholders to advance scientific safety methods; 2) determining the right level of chemical characterization for complex botanical mixtures; 3) finding practical, suitable NAMs for safety evaluation; 4) testing these methods against existing safety data on certain botanicals; 5) incorporating these methods into a systematic framework for assessing botanicals.

Results representing the past year's work include a set of in vitro studies for selected herbs, supporting safety evaluations and exemplifying BSC's structure, ambitions, and methodologies.

Initially focusing on oral intake from dietary supplements, the BSC may broaden its scope in future work phases.

References:

1. Mitchell CA et al. , Embry MR, Gafner S, Johnson H, Marsman DS, Kelber O, Oketch-Rabah HA, Rider CV, Sudberg E, Welch C. Enhancing the Botanical Safety Toolkit: The Strategy of the Botanical Safety Consortium. Poster contribution, 33. GUM Meeting 04.10. – 07. 09. 2022, Würzburg, Germany
2. Patel D, Sorkin BC, Mitchell CA, Embry MR, Rina-Kong S, Adams RE, DeTemple ER, Reddam A, Gafner S, Kelber O, Rider CV, Oketch-Rabah H, Roe AL, Marles RJ, Dever J, Dentali S. Improving the rigor and utility of botanical toxicity studies: Recommended resources. *Regul Toxicol Pharmacol.* 2023 Oct;144:105471. doi: 10.1016/j.yrtph.2023.105471.

* Presenting author. E-mail address: cmitchell@hesiglobal.org (Connie Mitchell)

S3.P92 Natural digestion mimicking curcuminoid formulation enhances oral bioavailability of curcuminoids

Johannes-Paul Fladerer^{a,b,*}, Thomas Klein^b, Albert Kompek^b

^a University of Graz, Institute of Pharmaceutical Sciences, Department of Pharmacognosy, Beethovenstraße 8, 8010 Graz, Austria

^b Apomedica Pharmazeutische Produkte GmbH, Roseggerkai 3, 8010 Graz, Austria

Curcumin, the fat-soluble, active ingredient of the curcuma root, is known for its low absorption and bioavailability and therefore various compositions and galenic technologies like addition of cyclodextrin or pepper extracts are on the market. The galenic formulation tested in this study was intended to increase bioavailability of curcuminoids and to mimic the natural digestion process for fat-soluble substances additionally. Thus, the curcuminoids were provided in an oily matrix mixed with phospholipids without integration of new technology or artificial excipients. This study evaluated the oral bioavailability of curcuminoids in this natural galenic formulation compared to the unformulated native curcuminoid powder. A randomized, cross-over, single oral dose study (200 mg curcuminoids) in 12 healthy subjects was performed under fasting conditions. Pharmacokinetic parameters were analysed from individual concentration-time curves of total curcuminoids as well as curcumin. Data showed significantly higher AUC_{0-4h} levels after the intake of the new phospholipid curcuminoid formulation for total curcuminoids (106.7 vs. 43.17 ng/mL*h, $p=0.0001$) as well as for Curcumin (77.8 vs. 20.66 ng/mL*h, $p<0.0001$) in comparison to the native curcuminoid powder. C_{max} was also significantly higher for both parameters analysed (total curcuminoids: 47.54 vs. 21.16 ng/mL, $p=0.0001$; Curcumin: 25.33 vs. 6.98 ng/mL, $p<0.0001$). In addition, the uptake kinetic of total curcuminoids was significantly faster with the new phospholipid curcuminoid formulation than that of the unformulated native curcuminoid powder (total curcuminoids: 2.21 vs. 5.17 h, $p<0.05$). In conclusion, the study confirms good tolerability of the new phospholipid curcuminoid formulation and its enhancing impact on curcuminoid uptake.

Keywords: Curcumin, bioavailability, phospholipid matrix, uptake kinetik

* Presenting author. E-mail address: johannes.fladerer@uni-graz.at (Johannes-Paul Fladerer)

S3.P93 Reveal the structure of plants by X-ray imaging and biomedical beamline at SSRF

Guohao Du^{a,b,*}, Jianfeng Ji^{a,b}, Han Guo^{a,b}, Yanling Xue^{a,b}, Yanan Fu^{a,b}, Biao Deng^{a,b}, Honglan Xie^{a,b}, Tiqiao Xiao^{a,b}

^a Shanghai Synchrotron Radiation Facility, Shanghai Advanced Research Institute, Chinese Academy of Sciences, Shanghai 201204, People's Republic of China

^b Shanghai Institute of Applied Physics, China Academy of Science, Shanghai 201800, People's Republic of China

X-ray Imaging and Biomedical Applications Beamline (BL13W) at Shanghai Synchrotron Radiation Facility (SSRF) has been opened to users since 2009. It is a powerful tool for non-destructive, high-resolution, and three-dimensional imaging research on samples from biology, materials, archaeology, geophysics, and other fields (Xie et al., 2015). With the construction of the phase II project of SSRF, BL13W underwent relocation to BL13HB in 2021 (Ji et al., 2023). The multilayer monochromator significantly enhanced the photon flux at the experimental station by over an order of magnitude, making it more suitable for dynamic X-ray imaging. The high resolution of 0.8 μm has greatly assisted in the identification of traditional Chinese medicinal materials (Ye et al., 2013) and the study of drug structures in pharmaceuticals. Additionally, dynamic X-ray imaging has provided valuable insights into the transport of water in plant roots and stems (Xu et al., 2023).

Projection images of Saw Palm was collected at X-ray Imaging and biomedical Beamline. The tomographic image (Fig.1) was calculated from the projections. The internal structure of the seed is clearly visible (Jaiswal et al., 2019).




Fig. 1. Tomography which shows the structure of the seed

BL13HB is like a high-resolution X-ray microscope that can help us reveal the internal structure of plants and many other types of samples.

Keywords: *X-ray Imaging*, multilayer monochromator, tomography

References

- Hong-Lan Xie., Biao Deng., Guo-Hao Du., Ya-Nan Fu., et al., 2015. Latest advances of X-ray imaging and biomedical applications beamline at SSRF. *Nuclear. Sci. Tech.*, 26, 020102.
- Jian-Feng Ji., Han Guo., Yan-Ling Xue., Rong-Chang Chen., et al., 2023. The new X-ray imaging and biomedical application beamline BL13HB at SSRF. *Nuclear Science and Techniques*, 34,197.
- L.L. Ye., Y.L. Xue., L.H. Ni., H. Tan., Y.D. Wang., T.Q. Xiao., 2013. Application of X-ray phase contrast micro- tomography to the identification of traditional Chinese medicines. *JINST*, 8, C07006.
- Mingwei Xu., Ke Li., Yanling Xue., Feixiang Wang., et al., 2023. Water refilling along vessels at initial stage of willow cuttage revealed by move contrast CT. *FRONTIERS IN PHYSICS*, 11.
- Yogini Jaiswal., Daniel Weber., Aaron Yerke., Yanling Xue., Danielle Lehman., et al., 2019. A substitute variety for agronomically and medicinally important *Serenoa repens*. *Scientific Reports*, 9,4709.

* Presenting author. E-mail address: dugh@sari.ac.cn (Guohao Du)

S3.P94 A cyclodextrin based tertiary complex of curcuminoids to improve bioavailability -its basic characteristics and immunomodulatory properties

Wuttichai Jaidee^a, Subhadip Banerjee^a, Narawadee Rujanapun^a, Kulwadee Malee^a, Thidarat Duanyong^{a,b}, Tharakorn Maneerat^{a,c}, Pravaree Phuneerub^{a,b}, Gudrun Ulrich-Merzenich^c, Rawiwan Charoensup^{a,b,*}

^a Medicinal Plant Innovation Center of Mae Fah Luang University, Mae Fah Luang University, Chiang Rai 57100, Thailand

^b School of Integrative Medicine, Mae Fah Luang University, Chiang Rai 57100, Thailand

^c School of Sciences, Mae Fah Luang University, Chiang Rai 57100, Thailand

^d University Hospital Bonn (UKB), Medical Clinic III, AG Synergy Research and Experimental Medicine, Bonn, Germany

The work aims to develop a water-soluble formulation from the rhizomes of *Curcuma longa* L. with immunomodulatory properties. Curcuminoids (CUR) were extracted using ethanol, and the total curcumin content was quantified by high-performance liquid chromatography (HPLC) utilizing the Agilent Technologies 1260 Infinity Series (Pawar et al., 2013). CUR (Curcumin equivalent 0.02 mmol) were complexed with beta-cyclodextrin (Poly- β -CD) at a 2:1 ratio. Polysaccharides (PS) isolated by hot water extraction from the rhizome (Yue et al., 2010), along with ascorbic acid (A), were included. Complexation was achieved through freeze-drying technique (Mohan et al., 2012). The CUR-A-Poly- β -CD complexes exhibited an average particle size of 517.9 nanometers, determined by laser light scattering using Horiba, Partica LA-950V2, with a solubility of 13.65 mg/L and stability over 45 days. The complexation process was analyzed using X-ray powder diffraction (XRPD). FTIR spectroscopy (Spectrum GX, Perkin Elmer) was utilized to confirm the presence of CUR, PS, A, and CD by comparing the FTIR spectrum of a sample containing the CUR-A-Poly- β -CD complex with reference spectra of standard compounds. The release kinetics of curcuminoids from the CUR-A-Poly- β -CD complex were assessed *in vitro* (Monmai et al., 2018 and Tang et al., 2019). Different ratios of CUR and PS within the CUR-A-Poly- β -CD complex were evaluated using biological and immunomodulatory activity assays (viability, phagocytosis, CD86/CD69 determination) in THP1, Jurkat T, and fibroblast cells. None of the tested ratios exhibited cytotoxic effects, although they differentially influenced immune cell functions.

Keywords: *Curcuma longa* L., water-soluble formulation, immunomodulatory activity, CUR-A- Poly- β -CD complex

References

- Mohan, P.K.; Sreelakshmi, G.; Muraleedharan, C.; Joseph, R. 2012. Water soluble complexes of curcumin with cyclodextrins: Characterization by FT-Raman spectroscopy. *Vib. Spectrosc.* 62, 77–84.
- Monmai, C., Park, S. H., You, S., & Park, W. J. 2018. Immuno-enhancement effect of polysaccharide extracted from *Stichopus japonicus* on cyclophosphamide-induced immunosuppression mice. *Food science and biotechnology*, 27(2), 565-573.
- Pawar HA, Gavasane AJ, Choudhary PD. 2018. A Novel and Simple Approach for Extraction and Isolation of Curcuminoids from Turmeric Rhizomes. *Nat Prod Chem Res.* 6 (300), 1-4.
- Tang, C., Sun, J., Liu, J., Jin, C., Wu, X., Zhang, X., ... & Zhang, N. 2019. Immune-enhancing effects of polysaccharides from purple sweet potato. *International journal of biological macromolecules*, 123, 923-930.
- Yue, G. G., Chan, B. C., Hon, P. M., Kennelly, E. J., Yeung, S. K., Cassileth, B. R., ... & Lau, C. B. 2010. Immunostimulatory activities of polysaccharide extract isolated from *Curcuma longa*. *International journal of biological macromolecules*, 47(3), 342-347.

* Presenting author E-mail address: rawiwan.cha@mfu.ac.th (Rawiwan Charoensup)

S3.P95 Phytochemical analysis of artichoke (*Cynara scolymus*) herb: Implications for *in vitro* neuroprotective and anti-obesity potential

Ewa Garbiec^a, Kornelia Budnik^a, Przemysław Zalewski^{a,b,*}, Judyta Cielecka-Piontek^{a,b}

^a Department of Pharmacognosy and Biomaterials, Poznan University of Medical Sciences, 3 Rokietnicka Str., 60-806 Poznan, Poland

^b Department of Pharmacology and Phytochemistry, Institute of Natural Fibres and Medicinal Plants—National Research Institute, 2 Kolejowa Str., 62-064 Plewiska, Poland

The artichoke (*Cynara scolymus*) is recognized for its positive impact on health, supported by experimental and clinical research. Its traditional application, renowned for addressing gastrointestinal issues, stems from its choleric and cholagogic effects. Beyond this, the artichoke herb, enriched with active compounds like phenolic acids and flavonoids, holds potential in addressing neurodegenerative diseases and obesity (Nourhan et al., 2021, Rondanelli et al., 2020). Hence, the aim of this study was to assess the inhibitory effects on acetylcholinesterase and lipase, along with examining the antioxidant activity of a dry, spirit extract derived from artichoke herb.

The methodology involved analyzing the tested extract using HPLC to determine the percentage content of chlorogenic acid (Carneiro et al., 2017). The total polyphenol content in the extract was quantified. The antioxidant activity of the extract was measured using DPPH, CUPRAC, and ABTS methods. Furthermore, the extract's ability to inhibit lipase and acetylcholinesterase was determined (Stasiłowicz-Krzemień et al., 2023).

The obtained results show that the tested extract contains 1.11% of the chlorogenic acid, meeting the European Medicines Agency requirement of at least 0.8% of this compound in artichoke herb extracts. The extract demonstrated antioxidant activity, as assessed including the use of DPPH, ABTS, and CUPRAC assays. It also showed inhibition of lipase ($EC_{50} = 53.144 \mu\text{g/mL}$) and acetylcholinesterase ($EC_{50} = 1.501 \text{ mg/mL}$), albeit less potent than standard substances. The average polyphenolic compound content was found to be 10.052 mg GAE/g, suggesting these compounds contribute to antioxidant and enzyme inhibitory activities.

Keywords: *Cynara scolymus*, phytochemical analysis, in vitro biological activity screening

References

- Carneiro, A., Barros, Y., Freitas, M., Simeoni, L.A., Magalhães, P.O., Silveira, D., Fonseca-Bazzo, Y.M., 2017. Identification and quantification of caffeoylquinic acid derivatives in *Cynara scolymus* L. tablets and capsules. *African Journal of Pharmacy and Pharmacology* 11, 94-102.
- Nourhan, M., Olfat M., Mohamed G. Shehata, Sobhy A., 2021. Antioxidant and anti-acetylcholinesterase potential of artichoke phenolic compounds. *Food Bioscience*, 41, 101006.
- Stasiłowicz-Krzemień, A., Sip, S., Szulc, P., Walkowiak, J., Cielecka-Piontek, J., 2023. The Antioxidant and Neuroprotective Potential of Leaves and Inflorescences Extracts of Selected Hemp Varieties Obtained with scCO₂. *Antioxidants* 12, 1827.
- Rondanelli, M., Riva, A., Petrangolini, G., Allegrini, P., Bernardinelli, L., Fazio, T., Peroni, G., Gasparri, C., Nichetti, M., Faliva, M.A., 2020. The Metabolic Effects of *Cynara* Supplementation in Overweight and Obese Class I Subjects with Newly Detected Impaired Fasting Glycemia: A Double-Blind, Placebo-Controlled, Randomized Clinical Trial. *Nutrients* 12, 3298.

* Presenting author. pzalewski@ump.edu.pl: (Przemysław Zalewski)

S3.P96 A preliminary study investigating the wound healing potential of *Cedrus brevifolia* extracts in mice

Georgia Lyggitsou^{a,b}, Christina Barda^a, Maria Anagnostou^a, Dimitra Statha^b, Christina Karampasi^b, Anastasia–Ioanna Papantonaki^b, Ioannis Svoliantopoulos^b, Andreas Vitsos^b, Michail Ch. Rallis^b, Helen Skaltsa^{a,*}

^a Department of Pharmacognosy and Chemistry of Natural Products, Faculty of Pharmacy, National & Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771, Athens, Greece;

^b Department of Pharmaceutical Technology, Faculty of Pharmacy, School of Health Sciences, National & Kapodistrian University of Athens, 15771 Athens, Greece

Wound healing stands as a paramount therapeutic pursuit, imposing significant challenges on healthcare, particularly for vulnerable populations such as diabetic patients and the elderly. Notably, wounds in diabetic individuals may escalate to the point of limb excision, while bedsores (pressure ulcers) among the elderly can culminate tragically and may lead to death. In this context, our study aims to investigate the potential of resin, leaves and bark extracts from *Cedrus brevifolia* (Hook.f.) Elwes & A. Henry in promoting wound healing in mice model. *C. brevifolia* is an endemic species of Cyprus, grown in Tripylos region, commonly known as Cedar Valley, within the Paphos forest. Despite its endemism, this species exhibits negligible genetic divergence from its Mediterranean related species. Previous studies revealed the chemical profile of extracts of different plant parts, i.e. needles, sapwood, heartwood and resin, focusing on potential active constituents, namely phenolic acids and flavonoids from polar bark extracts, while resin was rich in lignans and phenolic acids. These investigations were followed by assessments of the antioxidant and anti-inflammatory potential of all extracts and derived isolates (Douros et al., 2018; id., 2019). The findings underscore the need for further exploration in an *in vivo* setting, as indicated by Charalambous et al. (2022). Our experimental design employed 40 male SKH-hr2 black and brown mice aged 2–4 months. Wounds measuring 1 cm² were meticulously induced in anesthetized mice and the potential healing effect of the extracts was evaluated. The healing potential of *C. brevifolia* resin and extracts was then rigorously assessed through daily application of gel formulations containing resin concentrations of 5% and 10% w/w, alongside bark methanol-water extracts at concentrations of 0.5% and 1% w/w. These formulations, meticulously prepared with 10% propylene glycol and 3% carbomer 940, were subjected to comparative analysis. Evaluation of the treatments encompassed a multifaceted approach, incorporating clinical observations, skin biophysical parameter assessments utilizing the Antera 3D camera, and FT-IR spectroscopy, in addition to histopathological examination. Through this comprehensive evaluation, we aimed to elucidate the effects of *C. brevifolia* extracts on wound healing dynamics. Our study bridging the gap between *in vitro* observations and *in vivo* outcomes attempts to shed light on the potential therapeutic benefits of *C. brevifolia* in wound care.

Keywords: *Cedrus brevifolia*, Pinaceae, resin, bark methanol-water extract, wound healing

References

- Charalambous, D., Eliades, N.G.H., Christoforou, M., Kakouri, E., Kanakis, C., Tarantilis, P.A., Pantelidou, M., 2022. Chemical characterization, antioxidant and antimicrobial properties of different types of tissue of *Cedrus brevifolia* Henry extracts. *Molecules*, 27(9), 2717.
- Douros, A., Hadjipavlou-Litina, D., Nikolaou, K., Skaltsa, H., 2018. The Occurrence of Flavonoids and Related Compounds in *Cedrus brevifolia* A. Henry ex Elwes & A. Henry Needles. Inhibitory Potencies on Lipoxigenase, Linoleic Acid Lipid Peroxidation and Antioxidant Activity. *Plants*, 7(1), 1.
- Douros, A., Christopoulou, A., Kikionis, S., Nikolaou, K., & Skaltsa, H., 2019. Volatile components of heartwood, sapwood, and resin from a dated *Cedrus brevifolia*. *Nat. Prod. Comm.*, 14(6).

Presenting author. E-mail address: skaltsa@pharm.uoa.gr (Skaltsa Helen)

S3.P97 Preformulation studies of lycopene-rich tomato extract

Anna Kulawik^{a,b}, Kamil Wdowiak^a, Natalia Rosiak^a, Przemysław Zalewski^{a,c,*}, Judyta Cielecka-Piontek^{a,c}

^a Department of Pharmacognosy and Biomaterials, Poznan University of Medical Sciences, 3 Rokietnicka Str., 60-806 Poznan, Poland

^b Phytopharm Kłęka S.A., Kłęka 1, 63-040 Nowe Miasto nad Wartą, Poland

^c Department of Pharmacology and Phytochemistry, Institute of Natural Fibres and Medicinal Plants—National Research Institute, 2 Kolejowa Str., 62-064 Plewiska, Poland

The most common sources of lycopene in diets are tomatoes and products containing tomatoes. More than 85% of this ingredient in our diet comes from these sources. Tomatoes are also the cheapest source for lycopene production. The tomato-based products are a better source of this compound than raw tomatoes. Different varieties of tomatoes, as well as other fruits and vegetables, contain different amounts of this ingredient (Kulawik et al., 2023).

The methodology involved analyzing the tested tomato extract using HPLC to determine the content of lycopene (1.5%) (Olives Barba et al., 2006). Hot-melt extrusion technique was used to obtain solid dispersions of tomato extract with polyvinylpyrrolidone and Soluplus with following extract concentrations 10%, 20%, 30% and 40%. The prepared extrudates were further studied by XRPD to confirm their amorphous state. The intermolecular interactions were investigated by FT-IR/ATR. To determine the impact on biopharmaceutical properties, solubility as well as dissolution studies of lycopene were performed. The antioxidant activity of the extract was measured using the DPPH method.

The obtained results show the improvement of pharmaceutically important parameters. Formulations showed great potential for further development.

Keywords: Tomato extract, lycopene, hot-melt extrusion, amorphous state, solubility

References

- Kulawik, A., Cielecka-Piontek, J., Zalewski, P., 2023. The Importance of Antioxidant Activity for the Health-Promoting Effect of Lycopene. *Nutrients* 15, 3821.
- Olives Barba, A.I., Cámara Hurtado, M., Sánchez Mata, M.C., Fernández Ruiz, V., López Sáenz de Tejada, M., 2006 Application of a UV-vis detection-HPLC method for a rapid determination of lycopene and β -carotene in vegetables. *Food Chemistry* 95, 328–336

Acknowledgments

This research was funded by the Ministry of Science and Higher Education, grant number DWD/6/0002/2022.

* Presenting author. E-mail address: pzalewski@ump.edu.pl: (Przemysław Zalewski)

S3.P98 Investigating the mechanism of action of the anti-aging ability of geraniin nano-phytosome

Asly Poh-Tze Goh^a, Uma Devi Palanisamy^a, Usha Sundralingam^{b,*}

^a Jeffrey Cheah School of Medicine and Health Sciences, Monash University Malaysia, Jalan Lagoon Selatan, Bandar Sunway 47500, Selangor, Malaysia

^b School of Pharmacy, Monash University Malaysia, Jalan Lagoon Selatan, Bandar Sunway 47500, Selangor, Malaysia

This study explores geraniin, an ellagitannin from fruit waste, as a sustainable anti-aging solution in cosmetics. To overcome absorption challenges, a topical approach leveraging geraniin's potent antioxidant properties is pioneered. By unraveling its anti-aging mechanisms and introducing an innovative geraniin nano-phytosome, this study aligns with sustainability goals and addresses bioavailability issues, aiming to redefine skincare standards with eco-conscious formulations.

Geraniin from Rambutan rind was purified and assessed for its anti-aging effects on key skin enzymes. Nano-phytosome formulations containing geraniin, soy phosphatidylcholine, squalene, and edge activators (Tween 80, Span 80, sodium cholate hydrate) were developed. The optimized formulation was characterized for stability, entrapment efficiency, morphology, particle size, and permeability through ex vivo permeation studies.

Geraniin displayed significant anti-collagenase, anti-elastase, anti-hyaluronidase, and anti-tyrosinase effects compared to the positive control. The geraniin nano-phytosome with Tween 80 showed stability, elliptical morphology, entrapment efficiency of 67.61%, average particle size of 95.60 nm, and polydispersity index of 0.167. Although geraniin did not penetrate the skin, it accumulated within the membrane (53.87%), suggesting suitability for topical application.

This study underscores geraniin's promising anti-aging properties, demonstrating significant effects on key skin enzymes. Utilizing a nano-phytosome formulation with Tween 80 enhances its characteristics for topical application, indicating its potential in eco-conscious skincare formulations despite limited skin penetration.

Keywords: nanophytosome, geraniin, skin anti-aging, mechanism of action, permeability

* Presenting author. E-mail address: usha.sundralingam@monash.edu (Usha Sundralingam)

S3.P99 QbD approach for the development of CBD-loaded polymeric micelles for oral delivery

Lucia Grifoni*, Serena Orlandini, Essadia Rebbani, Giulia Vanti, Maria Camilla Bergonzi, Anna Rita Bilia

Department of Chemistry, University of Florence, 6 Ugo Schiff Str., 50019, Sesto Fiorentino, Florence, Italy

As part of our studies dedicated to the development of new drug delivery systems (Vanti, 2021) of cannabidiol, (CBD) a main constituent of *Cannabis sativa*, we report on QbD approach for preparation and optimisation of polymeric micelles for oral delivery. To date, only one prescription medicine has been approved. Oral administration of cannabidiol has been reported to have two major drawbacks: very low water solubility (12 mg/L) and extensive first-pass metabolism (Grifoni, 2022). To overcome these limitations, the present study developed micelles made of Poloxamer 407 (P407) and D- α -tocopherol polyethylene glycol succinate (TPGS). P407 is a GRAS excipient, while TPGS has been approved by FDA as a safe adjuvant and widely used in drug delivery systems. It has multiple advantages including high biocompatibility, enhancement of drug solubility, and improvement of drug permeation and inhibition of the activity of ATP dependent P-glycoprotein. The experimental design was carried out using D-optimal design; mixture process-variable (MPV) approach and mixture design were used for screening and optimization, respectively. Micelles loaded with CBD showed a size of 21.14 ± 0.30 nm, polydispersity index (PDI) of 0.040 ± 0.004 , recovery % of 98.24 ± 2.52 % and encapsulation efficiency % of 83.00 ± 2.52 %. Micelles loaded with CBD were evaluated for the stability in simulated gastrointestinal fluids in the presence of enzymes; a slight increase in PDI (0.280) was registered after intestinal dilution, while the size remained unchanged. A loss of up to 10% of CBD was observed in the simulated intestinal environment.

Keywords: design of experiments (DoE), cannabidiol, polymeric micelles, simulated gastrointestinal fluids

References

- Vanti, G., Grifoni, L., Bergonzi, M.C., Antiga, E., Montefusco, F., Caproni, M., Bilia, A.R. (2021). Development and optimisation of biopharmaceutical properties of a new microemulgel of cannabidiol for locally-acting dermatological delivery. *Int J Pharm.*, 607:121036.
- Grifoni, L., Vanti, G., Donato, R., Sacco, C., Bilia, A.R. (2022). Promising Nanocarriers to Enhance Solubility and Bioavailability of Cannabidiol for a Plethora of Therapeutic Opportunities. *Molecules*. 27(18):6070.

* Presenting author. E-mail address: lucia.grifoni@unifi.it (Lucia Grifoni)

S3.P100 Native metabolomics for the discovery of new small molecule gai protein modulators

Amira Naimi^{a,*}, Tim Berger^a, Daniel Petras^b, Daniel Hilger^c, Raphael Reher^a

^a Institute for Pharmaceutical Biology and Biotechnology, Department of Pharmacy, Philipps- University Marburg, Robert-Koch-Straße 4, 35037 Marburg, Germany

^b Department of Biochemistry, University of California Riverside, Riverside

^c Department of Pharmaceutical Chemistry, Philipps-University Marburg, Marbacher Weg 6, 35037 Marburg, Germany

The cost- and time-consuming process of finding a single bioactive molecule for a target protein from multiple complex crude extracts hampers natural product discovery. To accelerate and facilitate this iterative process of fractionating and bioactivity testing, we present a scalable native metabolomics approach, which combines non-targeted liquid chromatography-tandem mass spectrometry with the detection of protein binding by native mass spectrometry (Reher et al., 2022).

Heterotrimeric G proteins represent an interesting, but so far “undruggable” pharmacological target despite being involved in the regulation of a wide range of physiological and pathophysiological processes in living organisms. Currently, selective small molecule modulators specifically target G_{αq} and more recently G_{αs} (Dai et al., 2022), leaving the remaining G_α subfamilies without such modulators (Li et al., 2020). To date, only two highly selective G_{αq} inhibitors of natural origin have been identified: FR900359 (FR) and YM-254890 (YM). FR is derived from the endophytic *Candidatus* Burkholderia crenata, which is produced in the leaf nodes of the Traditional Chinese Medicine plant *Ardisia crenata* and underscores the remarkable diversity of interactions between organisms (Reher et al., 2018).

We use the native metabolomics approach to screen Gai against a diverse library of complex extracts derived from marine fungi. This led to the discovery, isolation and structure elucidation of the first selective small molecule inhibitor of Gai protein that paves the way for fundamental studies as well as new therapeutic strategies towards Gai-mediated diseases.

Keywords: native metabolomics, G protein, marine fungi

References

- Dai, S. A.; Hu, Q.; Gao, R.; Blythe, E. E.; Touhara, K. K.; Peacock, H.; Zhang, Z.; Von Zastrow, M.; Suga, H.; Shokat, K. M. State-Selective Modulation of Heterotrimeric G_{αs} Signaling with Macrocyclic Peptides. *Cell* 2022, 185 (21), 3950- 3965.e25.
- Li, J.; Ge, Y.; Huang, J.-X.; Strømgaard, K.; Zhang, X.; Xiong, X.-F. Heterotrimeric G Proteins as Therapeutic Targets in Drug Discovery. *J. Med. Chem.* 2020, 63 (10), 5013–5030.
- Reher, R.; Aron, A. T.; Fajtová, P.; Stincone, P.; Wagner, B.; Pérez-Lorente, A. I.; Liu, C.; Shalom, I. Y. B.; Bittremieux, W.; Wang, M.; Jeong, K.; Matos-Hernandez, M. L.; Alexander, K. L.; Caro-Díaz, E. J.; Naman, C. B.; Scanlan, J. H. W.; Hochban, P. M. M.; Diederich, W. E.; Molina-Santiago, C.; Romero, D.; Selim, K. A.; Sass, P.; Brötz-Oesterhelt, H.; Hughes, C. C.; Dorrestein, P. C.; O'Donoghue, A. J.; Gerwick, W. H.; Petras, D. Native Metabolomics Identifies the Rivulariapeptolide Family of Protease Inhibitors. *Nat Commun* 2022, 13 (1), 4619.
- Reher, R.; Kühl, T.; Annala, S.; Benkel, T.; Kaufmann, D.; Nubbemeyer, B.; Odhiambo, J. P.; Heimer, P.; Bäuml, C. A.; Kehraus, S.; Crüsemann, M.; Kostenis, E.; Tietze, D.; König, G. M.; Imhof, D. Deciphering Specificity Determinants for FR900359-Derived G_{αq} α Inhibitors Based on Computational and Structure-Activity Studies. *ChemMedChem* 2018, 13 (16), 1634–1643.

* Presenting author. E-mail address: amira.naimi@pharmazie.uni-marburg.de (Amira Naimi)

S3.P101 NMR-HetCA uncovers bioactive secondary metabolites from propolis extracts prior to isolation

Maria-Ioanna Stavropoulou, Antigoni Cheilari, Konstantia Graikou, Ioanna Chinou, Nektarios Aligiannis*

Department of Pharmacognosy and Natural Products Chemistry, Faculty of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou, 15771, Greece

Propolis is known for its antioxidant, antimicrobial and anti-inflammatory properties and it has been used traditionally to treat cuts, burns, wounds, and other skin conditions such as eczema and psoriasis. In our previous study (Stavropoulou et al., 2021), several propolis samples originating from Greece showed strong tyrosinase and high collagenase inhibition, respectively. In the current study, the methanolic extract of a sample from Mainland was fractionated by FCPC and the obtained fractions were evaluated *in-vitro* for their ability to inhibit DPPH free radical and collagenase enzyme. In parallel, the ¹H-NMR spectra of the fractions were recorded and the biological activity was statistically correlated with spectral data through the HeteroCovariance Approach (NMR-HetCA). NMR-HetCA is a MATLAB based toolbox that correlates NMR resonances and activity values through generated NMR pseudospectra, the HetCA plots. Each point (NMR peak) that is highly correlated is positive and color coded according to the respective correlation values (ranging from blue for those that show low correlation to deep red for those that show high correlation). HetCA plots revealed prior to isolation, high correlation of phenolic substances for DPPH and collagenase inhibition. In particular, the analysis suggested that polyphenols contribute to the DPPH free radical scavenging activity, while caffeic acid derivatives (e.g. CAPEs) seemed to be responsible for the anti- collagenase activity. The direct identification of the bioactive constituents was achieved in the complex propolis extract by the NMR-HetCA approach, a promising tool that can be used to accelerate the discovery of bioactive natural compounds prior to isolation.

Keywords: HeteroCovariance approach, propolis, antioxidant, anti-collagenase, caffeic acid derivatives

References

Stavropoulou, M.I., Stathopoulou, K., Cheilari, A., Benaki, D., Gardikis, K., Chinou, I., Aligiannis, N., 2021. NMR metabolic profiling of Greek propolis samples: Comparative evaluation of their phytochemical compositions and investigation of their anti-ageing and antioxidant properties. *Journal of Pharmaceutical and Biomedical Analysis* 194, 113814.

* Presenting author. E-mail address: aligiannis@pharm.uoa.gr (Nektarios Aligiannis)

S3.P102 Discovery of new bioactive compounds: cloning, activation and engineering of specialized metabolites biosynthetic gene clusters

Iryna Ostash*, Olena Kurylenko, Anna Kachor, Maxym Deneka, Sofiia Melnyk, Yana Sydorko, Markiyan Samborskyy, Yuriy Rebets

Explogen LLC, Zelena Street 20, Lviv, 79005, Ukraine

Genomes of bacteria are considered an inexhaustible source of new structurally diverse specialized metabolites. However, most of this chemical diversity is hidden within bacterial genomes, since corresponding genes are rarely expressed efficiently under laboratory conditions. Genomes of soil-dwelling *Actinomycetes* (in first turn *Streptomyces*), each harboring approximately 30 SM-BGCs, are the richest and validated source of secondary metabolites that are still awaiting to be discovered (Lee et al., 2019).

Heterologous expression is one of the most efficient approaches for the discovery of new specialized metabolites. However, it is limited by the access to the biosynthetic gene clusters (BGCs) to be expressed. We developed two different but complementary approaches for cloning the bacterial natural products BGCs. The first approach is based on optimized TAR (transformation-associated recombination) cloning. The procedure was modified by introducing the direct selection of clones carrying chromosomal DNA inserts. As a showcase, chelocardin (35 kbp), kanamycin (42 kbp), and daptomycin (67 kbp) BGCs were cloned and expressed in *Streptomyces albus* and *Streptomyces lividans* hosts. Another approach is based on massive BGCs cloning by cosmid library construction, sequencing, and mapping. This way multiple BGCs were recovered including eight aminoglycoside BGCs expressed in the aforementioned hosts.

In conclusion, the approaches presented here significantly increase the chances of discovering novel specialized metabolites and could be efficient for cloning large bacterial genome fragments. The details of both approaches will be presented.

Keywords: specialized metabolites, TAR cloning, heterologous expression

References

Lee N, Hwang S, Lee Y, Cho S, Palsson B, Cho BK. Synthetic Biology Tools for Novel Secondary Metabolite Discovery in *Streptomyces*. J Microbiol Biotechnol. 2019 May 28;29(5):667-686. doi: 10.4014/jmb.1904.04015. PMID: 31091862.

* Presenting author. E-mail address: iryna.kozarevska@gmail.com (Iryna Ostash)

S3.P103 *Origanum dictamnus* infusion characterization by HPLC-PDA-MS and nanoformulation development

Charikleia Paloukopoulou^{a,*}, Giulia Vanti^{b,*}, Anastasia Karioti^a, Anna Rita Bilia^b

^a Laboratory of Pharmacognosy, School of Pharmacy, Aristotle University of Thessaloniki, University Campus, 54124, Thessaloniki, Greece

^b Department of Chemistry "Ugo Schiff", University of Florence, Via Ugo Schiff 6, 50019 Sesto Fiorentino, Florence, Italy

Origanum dictamnus L. (Lamiaceae) is a Greek species endemic to the island of Crete, commonly known as dittany. In 2014, dittany was authorized by EMA (EMA, 2014) under the category of Traditional Use Herbal Medicinal Products in the form of herbal teas for the treatment of skin inflammations and bruises, as a wound healing agent, for mild gastrointestinal disorders and relief of cough associated with cold.

In the present work, dittany infusion was prepared according to EMA and characterized by HPLC- PDA-MS. Identification of the constituents was assisted by chromatographic isolations (Paloukopoulou et al., 2023). Rosmarinic acid, salvianolic acid P, and other depsides were the main constituents, whereas apigenin and luteolin glycosides, as well as neolignans, were also present. The infusion was quantified based on its rosmarinic acid content. In continuation of our investigation on this important medicinal plant and given the EMA indications, preliminary studies were carried out with dittany infusion, which showed a relaxing effect on the rabbit intestine. Since infusions need to be consumed instantly to maintain their nutritional features, the inclusion of aqueous herbal extracts in advanced technological forms was important to improve their chemical stability (Bilia et al., 2019). Specifically, this study focused on developing a nanoformulation for oral infusion delivery. The infusion was quantified in terms of rosmarinic and salvianolic acids by HPLC-DAD. The physical characterization of the nanoformulation was performed by dynamic light scattering and transmission electron microscopy. Both techniques showed nanoparticles of about 130 nm and optimal homogeneity in dimension distribution. The infusion encapsulation efficiency was evaluated by the dialysis method and showed a percentage of 93.041 for rosmarinic acid and 92.12 for salvianolic acid. Chemical and physical stability studies on the storage and incubation of the nanoformulation in simulated biological fluids (saliva, gastric, intestinal) are ongoing. Preliminary findings showed a good stability of the nanoformulation during the first two weeks of storage.

Keywords: *Origanum dictamnus*, Lamiaceae, infusion, HPLC-PDA-MS, nanoformulation

References

- EMA, 2014. EMA/HMPC/200429/2012 Corr. Committee on Herbal Medicinal Products (HMPC) Community herbal monograph on *Origanum dictamnus* L., herba.
- Paloukopoulou, C., Tsadila, C., Govari, S., Soulioti, A., Mossialos, D., Karioti, A., 2023. Extensive analysis of the cultivated medicinal herbal drug *Origanum dictamnus* L. and antimicrobial activity of its constituents. *Phytochemistry*, 208, 113591. Advance online publication.
- Bilia, A. R., Piazzini, V., Risaliti, L., Vanti, G., ... & Bergonzi, M. C., 2019. Nanocarriers: A successful tool to increase solubility, stability and optimise bioefficacy of natural constituents. *Curr Med Chem*, 26(24), 4631-4656.

Acknowledgments

This research was funded by the project EthnoHERBS (H2020-MSCA-RISE-2018, Grant Agreement No. 823973). The financial support provided by the MUR - Dipartimenti di Eccellenza 2023-2027 (DICUS 2.0) to the Department of Chemistry "Ugo Schiff" of the University of Florence is acknowledged.

* Presenting author. E-mail address: giulia.vanti@unifi.it (Giulia Vanti)

S3.P104 Extract from the seedlings of *Brassica oleracea var. italica* (Brassicaceae) reduces tissue damage and alters inflammatory markers over time: development of new medications

Katherine M Warpeha^{a,*}, Cimran Nakum^a, Yahya Najjar^a, Aleksandra A. Gurgul^b, Chun-Tao Che^b, Thomas J. Park^a

^a Department of Biological Sciences, University of Illinois at Chicago, 900 S Ashland Ave, Chicago, IL USA

^b Department of Pharmaceutical Sciences, College of Pharmacy, University of Illinois Chicago, Chicago, IL, USA

This presentation outlines the utility of a combinatorial of natural broccoli chemicals processed by liver microsomal S9 fraction to reduce direct tissue injury and produce a more favorable cytokine and inflammation profile. Pain is a global issue needing more effective, affordable, and non-addictive and non-opiate therapies. Many chemicals derived from phenylalanine have antioxidant and anti-inflammatory properties, yet many structures are as yet unstudied (Perkowski & Warpeha, 2019). Using the edible plant broccoli, a growth protocol was developed to induce a concentrated combinatorial of potential phenylpropanoid anti-inflammatories extracted from seedlings, then treated with liver microsomal S9 fraction (Gurgul et al. 2023). A mouse model of pain behavior was used for an in vivo test of early and late pain responses in a paw. The vehicle control mice demonstrated tissue damage, and cytokine and NOS2 levels indicating negative aspects of inflammation. Pre-treatment of mice with S9 broccoli extract (ip injection 30 min prior to pain stimulus) indicated improved tissue integrity in early and late responses examined at the tissue level in the paw. In addition, cytokine levels showed changes right after pain stimulus, but also positive affects 35 min later. Preliminary data suggest that this non-opiate inflammation and pain-reducing treatment could have multiple systemic affects to reduce pain.

Keywords: phenylpropanoid, broccoli, pain, anti-inflammatory, opiate

References

Gurgul, A.A., Najjar, Y., Chee, A. An H., Che C-T, Park T.J., Warpeha K.M. 2023. Phenylpropanoid-enriched broccoli seedling extract can reduce inflammatory markers and pain behavior. J. Transl. Med. 21, 922.

Perkowski, M.C., Warpeha, K.M. 2019. Phenylalanine roles in the seed-to-seedling stage: Not just an amino acid. Plant science: an international journal of experimental plant biology. 289, 110223.

* Presenting author. E-mail address: kwarpaha@uic.edu (Katherine M Warpeha)

S3.P105 Identification of naturally-derived lipids as GPR84 agonists

Listiana Oktavia^{a,*}, Pinqi Wang^{a,b}, Vincent B. Luscombe^c, David R. Greaves^c, Angela J. Russell^{a,b}

^a Department of Chemistry, University of Oxford, Mansfield Road Oxford OX1 3TA, UK

^b Department of Pharmacology, University of Oxford, Mansfield Road, Oxford OX1 3QT, UK

^c Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford OX1 3RE, UK

G Protein-Coupled Receptor 84 (GPR84) has emerged as a promising target for therapeutic interventions in metabolic dysfunction, inflammation-related diseases, cancer, and bacterial infections (Luscombe et al. 2023, Wang et al., 2024) due to its predominant expression on immune cells (Lucy et al., 2019). However, understanding the pathophysiological role of GPR84 is hindered by the absence of a verified endogenous ligand. The suggested endogenous ligands of GPR84 (C₁₀- C₁₄ Medium Chain Fatty Acids) show low potency *in vitro* cAMP assay (Marsango et al., 2022). This study aimed to develop suitable methods to identify potent endogenous or exogenous GPR84 ligands from natural resources.

As a test system, we focused on Goat Milk (GM) to discover novel GPR84-activating lipids through bioactivity-guided isolation and a biochemometric approach. Polar lipid fractions (S30_a and S30_b) from GM exhibited over 10-fold higher potency in cAMP assays compared to the most active reported medium chain fatty acid (capric acid C₁₀), suggesting the presence of previously unreported GPR84 agonist(s). Component analysis and bioactivity-NMR correlations tentatively identified sphingolipid- like ligands.

Subsequently, five sphingolipid analogues with various alkyl chain lengths (R:C₆ to C₁₀) were synthesized and tested. The findings revealed that the Sphinganine analogue (R:C₉) showed better activity (EC₅₀:215 nM) than capric acid EC₅₀: 4.5 µM). Moreover, this discovery unveils the previously unreported potency of sphinganine analogues on GPR84 activation. Further efforts will involve additional separation and analysis of bioactive compound in the GM fraction, and development of omic-based platforms to robustly identify bioactive compound(s) in the natural product mixture.

Keywords: goat milk, GPR84, lipid, sphingolipids

References

- Lucy, D., Purvis G.S.D, Zeboudj L., Chatzopoulou, M., Recio, C., Bataille, C.J.R., Wynne, G.M., Russell, A.J., 2019. A Biased Agonist at Immunometabolic Receptor GPR84 Causes Distinct Functional Effects in Macrophages. *ACS. Chem. Biol.* 14(9), 2055–64. Luscombe, V.B., Wang, P., Russell, A.J., Greaves, D.R., 2024. Biased agonists of GPR84 and insights into biological control. *Br. J. Pharmacol.* 181, 1509–1523.
- Wang, P., Raja A., Luscombe V.B., Bataille C.J.R., Lucy, D., Rogga, V.V., Russell, A.J., 2023. Development of Highly Potent, G-Protein Pathway Biased, Selective, and Orally Bioavailable GPR84 Agonists. *J. Med. Chem.* 67, 1, 110–137. Marsango, S., Barki, N., Jenkins, L., Tobin, A.B., Milligan, G., 2022. Therapeutic validation of an orphan G protein- coupled receptor: The case of GPR84. *Br. J. Pharmacol.* 179(14), 3529–41.

* Presenting author. E-mail address: listiana.oktavia@chem.ox.ac.uk (Listiana Oktavia)

S3.P106 Quantitative analysis of endogenous steroids by UPLC-MS/MS in various medicinal plant species

Petra Barnášová, Lucie Rárová, Petra Amakorová, Jana Oklestková*

Laboratory of Growth Regulators, The Czech Academy of Sciences, Institute of Experimental Botany & Palacký University, Faculty of Science, Šlechtitelů 27, 78371 Olomouc, Czech Republic

Traditionally, mammalian steroids were thought to be a diverse group of bioactive compounds found only in animal tissues. However, recent research has confirmed that mammalian steroids are also present in several plant species. These steroids, including pregnenolone, progesterone, testosterone and estrogens, have been identified in different parts of plants such as leaves, stems, roots and reproductive organs. Their localisation in specific tissues suggests a potential role in plant development and reproductive processes (Tarkowská 2019).

Plant steroids are known to modulate a wide range of physiological responses in plants, leading to improvements in both the quality and yield of food crops. Treatment of plants with steroids or their precursors affects plant development: cell division, root and shoot growth, embryo growth and flowering. These compounds also have many interesting medicinal and pharmaceutical effects, such as anti-cancer, anti-angiogenic, antiviral and antibacterial bioactivities (Janeczko 2021).

The aim of this work was to determine the profile of mammalian sex hormones (progestagens, androgens and estrogens) in medicinal plants (in their flowers, leaves, buds). A combination of SPE and ultra-high performance liquid chromatography with tandem mass spectrometry (UHPLC- MS/MS) was used for the determination of steroid compounds. The antiproliferative activity of selected plant extracts was also tested on cancer cell lines.

Keywords: steroids, UHPLC-MS/MS, anti-cancer activity, medicinal plants

References

Tarkowská, D. Plants are Capable of Synthesizing Animal Steroid Hormones. *Molecules* 2019, 24, 2585. Janeczko, A. Estrogens and Androgens in Plants: The Last 20 Years of Studies. *Plants* 2021, 10, 2783.

* Presenting author. E-mail address: jana.oklestkova@upol.cz (Jana Oklestková)

S3.P107 Enhancing drug discovery efficiency in plant extracts through polyphenol removal

Christina Davidson*, Katherine Schoenhardt, Pelle Simpson, Venkat Macherla

Enveda Biosciences Inc., 5700 Flatiron Pkwy Boulder, CO 80301, USA

Polyphenols, such as flavonoids, tannins, and phenolic acids, are common compounds found in plant extracts. In the drug discovery process polyphenols in plant extracts are often promiscuous hitting compounds due to assay interference, non-specific binding, and redox cycling resulting in false positives. Moreover, their low *in vivo* oral bioavailability makes most polyphenols unsuitable as drug leads. In addition, the high abundance of these nuisance compounds can make the discovery of more attractive drug leads difficult. This study presents a method using polyvinylpolypyrrolidone (PVPP) resin in solid-phase extraction to reduce polyphenol content, thereby improving the detection of minor bioactive compounds. Evaluation of the process via HPLC-QTOF and MS/MS using Compound Discovery in Agilent MassHunter confirmed that approximately 80% polyphenols have been removed, and many previously obscured compounds have been identified. This method enhances confidence and quality in drug discovery screening outcomes and provide true hits that were previously suppressed due to excess polyphenols.

Keywords: *polyphenols*, polyvinylpolypyrrolidone, drug discovery, HPLC-QTOF, solid-phase extraction

* Presenting author. E-mail address: christina.davidson@envedabio.com (Christina Davidson)

S3.P108 Bioactive *Lycopodium* alkaloids from clubmoss cultivars and transformation to enhance anti- acetylcholinesterase activity

Nopporn Thasana^{a,b,c,*}, Wanlaya Thamnarak^a, Chuleeporn Ngernnak^a, Wachirasak Thaisaeng^a, Shisanupong Anukanon^b, Somsak Ruchiwat^{a,b,c}

^a Laboratory of Medicinal Chemistry, Chulabhorn Research Institute, Laksi, Bangkok 10210, Thailand

^b Chemical Sciences Program, Chulabhorn Graduate Institute, Laksi, Bangkok 10210, Thailand

^c Center of Excellence on Environmental Health and Toxicology (EHT), OPS, Ministry of Education, Science, Research and Innovation, Bangkok, 10400, Thailand

The *Phlegmariurus* genus (Lycopodiaceae) is cultivated as ornamental plants in Thailand, with extracts serving as a rich source of *Lycopodium* alkaloids (Ngernnak et al., 2021). This presentation encompasses the isolation, structure elucidation, transformation, and biological evaluation of isolated *Lycopodium* alkaloids from Thai clubmoss cultivars. Eleven unprecedented *Lycopodium* alkaloids, namely phlegcarines A-C and nummulines A-H, were isolated from *Phlegmariurus carinatus* (Desv. ex Poir.) Ching. (Thamnarak et al., 2023) and *P. nummulariifolius* (Blume) Ching, respectively. Additionally, an acetylcholinesterase (AChE) inhibitor, huperzine A, was identified. The structural assignments were established through comprehensive spectroscopic techniques, including HR-ESI- MS, NMR techniques, chemical correlations, and further confirmed by X-ray analysis specifically for nummuline D. While the isolated *Lycopodium* alkaloids were evaluated for anti- acetylcholinesterase activity, none demonstrated efficacy surpassing that of huperzine A (IC_{50} eeAChE = 0.03 μ M). Huperzine A derivatives were further synthesized to enhance potency (Anukanon et al., 2021).

Fig. 1. Structure of new *Lycopodium* alkaloids

Keywords: Lycopodiaceae, *Lycopodium* alkaloid, *Phlegmariurus*, huperzine A, acetylcholinesterase

References

- Anukanon, S., Pongpamorn, P., Tiyaabhorn, W., Chatwichien, J., Niwetmarin, W., Sessions, R.B., Ruchirawat, S., Thasana, N., 2021. In silico-guided Rational Drug Design and Semi-synthesis of C(2) Functionalized Huperzine A Derivatives as Acetylcholinesterase Inhibitor. ACS Omega 6, 19924-19939.
- Ngernnak, C., Thamnarak, W., Thaisaeng, W., Ruchirawat, S., Thasana, N., 2021. *Lycopodium* Alkaloids from Thai Club Mosses. Prayogik Rasayan 5, 7-16.
- Thamnarak, W., Thaisaeng, W., Batsomboon, P., Eurtivong, C., Chainok, K., Wannarit, N., Ruchirawat, S., Thasana, N., 2023. Phlegcarines A-C, Three *Lycopodium* Alkaloids from *Phlegmariurus carinatus* (Desv. ex Poir.) Ching. Phytochemistry 206, 113553.

* Presenting author. E-mail address: nopporn@cri.or.th. (Nopporn Thasana)

S3.P109 Depicting the chemical diversity of bioactive meroterpenoids produced by the largest organism on earth

Marc Stadler*

Department Microbial Drugs, Helmholtz Centre for Infection Research (HZI), Inhoffenstrasse 7, 38124 Braunschweig, Germany

We have explored the diversity of melleolide-type meroterpenoids produced by *Armillaria ostoyae*, one of the largest and oldest organisms on Earth, using extracts from liquid and solid fermentation media. The study unveiled three unprecedented dimeric bismelleolides (1-3) and three novel fatty acid-substituted congeners (4-6), along with 11 new (7-17) and 21 known (18-38) derivatives. Structure elucidation was done by 1D- and 2D-NMR spectroscopy, HRESI-MS data, and ROESY spectral analysis for relative configurations. Absolute configurations were determined through crystal structures and ECD spectra comparison. A compound library of melleolide-type meroterpenoids facilitated metabolomics-wide associations, revealing production patterns under different culture conditions. The library enabled assessments of antimicrobial and cytotoxic activities, unveiling that the $\Delta^{2,4}$ double bond is not crucial for antifungal activity. Cytotoxicity was linked to the presence of an aldehyde at C-1, but lost with a hydroxylation at C-13. Chemoinformatic analyses demonstrated the intricate interplay of chemical modifications on biological properties. This study marks the first systematic exploration of *Armillaria* spp. meroterpenoid diversity via MS-based untargeted metabolomics, offering insights into structure-activity relationships through innovative chemoinformatics.




Fig. 1. Some chemical structures of newly identified metabolites from cultures of *A. ostoyae*.

Keywords: Basidiomycota, metabolic diversity, metabolomics, cheminformatics

Reference

Pfütze, S., Charria-Girón, E., Schulzke, E. et al.. 2024. Depicting the chemical diversity of bioactive meroterpenoids produced by the largest organism on Earth. *Angew Chem Int Ed*, in press. DOI:10.1002/ange.202318505

* Presenting author. E-mail address: marc.stadler@helmholtz-hzi.de (Marc Stadler)

S3.P110 Comparative *In Vitro* enzyme inhibitory activities of different plant parts of five *Albizia* species (Fabaceae)

Abdullahi Adeyemi Adegoke*, Queeneth Abiola Ogunniyi, Abraham O. Nkumah, Omonike O. Ogbole

Department of Pharmacognosy, University of Ibadan, Ibadan, Nigeria

The number of diabetic patients in Africa is expected to rise to 41 million in 2045. Management of diabetes without any side effects is still a challenge to the medical community. The study aims to evaluate the comparative *in vitro* enzymatic inhibitory activities (α -amylase and α -glucosidase) of different plant parts of five *Albizia* species (*Albizia lebbbeck*, *Albizia ferruginea*, *Albizia saman*, *Albizia coriaria*, *Albizia zygia*).

The plants were selected based on previous studies, and were further authenticated at Forest Herbarium Ibadan, where voucher specimens were deposited. The plant parts were macerated for 72 hours at room temperature using methanol. The extract was filtered and concentrated to dryness and stored in a refrigerator at 4 °C. The phytochemical screening of the crude extract was done following a standard procedure (Trease and Evans, 2002). Ethyl acetate fractions from partitioned extracts showing the best activity in α –amylase and α –glucosidase inhibitory assays were subjected to vacuum liquid chromatography (VLC). Data were analyzed using Graph Pad Prism 5.0.

The phytochemical screening of the crude extract revealed the presence of secondary metabolites in some species, and some are absent or present in little quantities. *Albizia lebbbeck* bark had the lowest α - glucosidase and α - amylase inhibitory values (IC_{50} <31.20 μ g/mL and 42.33 μ g/mL, respectively) when compared with the standard acarbose (IC_{50} 121.50 μ g/mL) and was subjected to solvent-solvent partitioning. Ethyl acetate fraction which showed the best activity in both α - glucosidase and α - amylase inhibitory assay was subjected to VLC.

Albizia lebbbeck bark extracts could be regarded as a source of phytochemicals with potential use in the management of postprandial hyperglycemia.

Keywords: *Albizia lebbbeck* (L.) Benth, Diabetes, Maceration, Vacuum liquid chromatography, α -glucosidase inhibitory assay

References

Trease, GE; and Evans, WC (2002). Pharmacognosy (15th Edition). Saunders. pp 214-393.

*Presenting author: aadegoke0278@stu.ui.edu.ng (Abdullahi Adeyemi Adegokea)

S3.P111 Study and nutritional screening of some notanicals used as herbal remedies in managing eye and ear ailments in Ado-Odo Ota Local Government Areas, Ogun State

Abdullahi Adeyemi Adegoke^{a,c,d,*}, Anthony Wale Ojewumi^b, Adenike Mariam Saliu^c, Queeneth Abiola Ogunniyi^a, Mary Ololade Akinlade^c, Maryam Titilope Agbaje^e, John Oluwafemi Oyeyode^f, Olubunmi Olaitan Mada^a, Abraham Oluwalana Nkumah^a

^a Department of Pharmacognosy, University of Ibadan, Ibadan, Nigeria

^b Department of Botany, Lagos State University, Nigeria

^c Department of Biological Sciences, Federal University of Agriculture, Abeokuta, Nigeria

^d Department of Biological Sciences, Westland University Iwo, Osun State, Nigeria

^e Department of Food Technology, University of Ibadan, Ibadan, Nigeria

^f Department of Chemistry, Federal University of Agriculture, Abeokuta, Nigeria

Traditional medicine is one of the most well-established and widely used types of healthcare. It has long been known that phytochemicals can be used to treat infectious disorders. Several ailments, including those concerning vision and auditory senses, are treated with a variety of natural products. This research aims to source the natural products used in the management of ear and eye ailments.

An ethnobotanical survey was conducted in Ado-Odo, Ota Local Government. A total of hundred structured and validated questionnaires were administered to a hundred respondents by purposeful sampling method. Also, nutritional, phytochemical, and antinutrient constituents of the most mentioned plants were investigated. Ethnobotanical data were analyzed by ANOVA and DMRT at $p < 0.05$.

Results revealed that 37 plant species belonging to 26 families being used to treat eye and ear ailments were collated. The most frequently mentioned plants were *Cucumeropsis mannii*, *Allium cepa*, and *Allium sativum*. Leaves (37.0%) seeds (30.0%), roots (10.0%), flowers value (7.0%) and fruits (7.0%), stems (5.0%) and barks (4.0%) were the most exploited plant part used for the diseases. The preparations of these plants were administered by dropping the preparations of the plants into the affected eye and ear, bathing with the leaf preparations, and oral application. Anti-nutrient, proximate values, and Vitamins were documented in *Cucumeropsis mannii*, *Allium cepa* and *Allium sativum*.

The effectiveness of plants like *Cucumeropsis mannii*, *Allium cepa*, and *Allium sativum* for the treatment of ear and eye diseases could be due to the presence of alkaloids, flavonoids, saponin, and Vitamin C.

Keywords: nutritional, conjunctivitis, phytochemicals, glaucoma, deafness

* Presenting author. E-mail address: adegokeabdullahi24@gmail.com (Abdullahi Adeyemi Adegoke)

S3.P112 New data about chemical composition of *Cynanchum acutum* L.

Vazha Nebieridze*

The University of Georgia, Department of Pharmacy, 77a, M. Kostava str., 0171, Tbilisi, Georgia

This presentation will cover the isolation and structure elucidation of a new data about chemical composition from *Cynanchum acutum* (L.). This genus comprises of about 200 species in *Asclepiadaceae* family and is distributed worldwide. *C. acutum* is quite poisonous with few medical applications.

The roots of *C. acutum* were powdered and extracted three times with MeOH at room temperature. The collected extracts were dried under reduced pressure, and the concentrate was partitioned between CHCl₃ and H₂O. A part of the aqueous extract was subjected to column chromatography.

Column chromatography was carried out on Diaion HP-20, Sephadex LH-20 and silica gel yielding 7 known compounds: 3 cardenolide glycosides - lanatoside A, lanatoside C (Yochu Fujii et al., 1990) and corchoroside C (Nathan P Mirtallo Ezzone et al., 2022), 2 flavonoids - kaempferol 3-O-β-D- rutinoside and quercetin 7-O-β-D-glucopyranoside (Mona A. Mohamed et al., 2008) and 2 pregnans – atratosides A and atratoside D (Zhuang-Xin Zhang et al., 1988) The full chemical structures were determined by using one- and two-dimensional nuclear magnetic resonance spectroscopy (1H, C13, HSQC, HMBC, COSY) and mass spectrometry.

It is significant that cardenolide glycosides in plants of the genus *Cynanchum* were described for the first time by us.

Keywords: *Cynanchum acutum*, cardenolide glycosides, flavonoids, pregnans

References

- Mona A. Mohameda, Wafaa S. Ahameda, Mortada M. El-Saida and Heiko Hayen. 2008. New Acylated Flavonol Diglycosides of *Cynanchum acutum*. *Natural Product Communications* Vol. 3 No. 2, 193 – 198.
- Nathan P Mirtallo Ezzone, Gerardo D Anaya-Eugenio, Ermias Mekuria Addo, Yulin Ren, A Douglas Kinghorn, Esperanza J Carcache de Blanco. 2022. Effects of Corchoroside C on NF-κB and PARP-1 Molecular Targets and Toxicity Profile in Zebrafish. *Molecular Sciences.*, 23(23), 145-146.
- Yochu Fujii, Yukari Ikeda, Ichiro Okamoto and Mitsuru Yamazaki. 1990. Separation and determination of lanatosides in *Digitalis lanata* leaves by high-performance liquid chromatography. *Journal of Chromatography*, , 241-246.
- Zhuang-Xin Zhang, Jun Zhou, Koji Hayashi and Koh Kaeko. 1988. Atratosides A, B, C and D, steroid glycosides Of *Cynanchum atratum*. *Phytochemistry*, Vol. 27, No. 9, pp. 2935-2941.

* Presenting author. E-mail address: v.nebieridze@ug.edu.ge (Vazha Nebieridze)

S3.P113 A comparative study between CPC and flash chromatography for natural product purification

Michel Thomas*, Justine Cadoret, Grégoire Audo

Gilson Purification SAS, 22 rue Bourseul, 56890 Saint Avé, France

Centrifugal Partition Chromatography (CPC) and flash chromatography are both chromatography- based techniques that are widely used in the pharmaceutical and cosmetic industries for drug discovery or enrichment of active ingredients. Further, CPC and flash chromatography are easily applicable to scale up to meet industrial applications (Sutherland, 2007) (Fernando et al., 2022). They differ from each other in the nature of their stationary phase. The CPC is liquid-liquid chromatography using two immiscible liquid phases, collectively referred to as the solvent system. Whilst flash chromatography requires the use of a cartridge filled with a solid stationary phase (e.g. silica, C₁₈). Due to its liquid nature, CPC is often seen as a solvent-consuming technique compared to other preparative strategies.

Based on these findings, the main objective of this study was to compare CPC and flash chromatography for the purification of piperine (**1**, Fig. 1), an alkaloid that confers the pungent taste to black pepper (*Piper nigrum*). Piperine can be easily purified in one step from crude extract of black pepper using CPC or flash chromatography. Both purification techniques were compared at laboratory and pilot scales in terms of purity, yield, solvent consumption, and injected quantity.




Fig. 1. Structure of Piperine 1

Results highlight CPC at laboratory and pilot scale for piperine purification. A larger quantity of sample can be injected in CPC, with lower solvent consumption and no solid waste to treat (i.e. silica cartridge). Furthermore, the purities obtained by CPC were always higher than those obtained by flash chromatography. Therefore, this work addresses the issue of solvent consumption and recommends CPC for purification of natural products.

Keywords: piperine, purification, CPC, flash chromatography

References

- Sutherland I,A, 2007. Recent progress on the industrial scale-up of counter-current chromatography. J. Chrom. A. 1151, 6-13.
- Fernando, G.S.F. et al., 2022. Novel approach for purification of major betalains using flash chromatography and comparison of radical scavenging and antioxidant activities. Food Chem. 385, 132632.

* Presenting author. E-mail address: tmichel@gilson.com (Thomas Michel)

S3.P114 Molecular docking and dynamics simulations study of Cathecin from *Mentha cordifolia* as potential HIV-1 protease inhibitor and Tat1-Cyclin competitor

Jeffrey Clement Bungar^{a,b,*}, Rod Vincent Borromeo^a, Ranelle Janine Asi^a, Sheriah Laine de Paz-Silava^c, Ahmad Reza Mazaherya, Ricky Nellas^b

^a Institute of Biology, University of the Philippines Diliman, Diliman, Quezon City, Philippines

^b Institute of Chemistry, University of the Philippines Diliman, Diliman, Quezon City, Philippines

^c College of Public Health, University of the Philippines Manila, Pedro Gil, Manila, Philippines

There has been a 411% increase in daily incidence of Human Immunodeficiency Virus (HIV) cases in the Philippines from 2012 to 2023 (Gangciangco and Eustaquio, 2023). There are antiretroviral drugs that are available which increases a patient's survivability and prolonging their lifespan. However, the emergence of drug-resistant HIV strains and discovery of detrimental side effects of these drugs threatens the effectiveness of it.

A study showed that *Mentha cordifolia*, a plant included in the Philippines' Department of Health's Ten Medicinal Plants, can reduce the p24 production in HIV-1 latently infected cells (OM10.1) *in vitro* (dePaz-Silava et al., 2022). The same group was able to identify that cathecin, one of the known compounds found in the plant extract, can strongly bind to HIV-1 protease and Tat-Cyclin, and potentially inhibit their functions, through *in silico* analysis.

Researchers performed 100 ns of Molecular Dynamics simulation for both HIV-1 Protease and Tat- Cyclin to obtain and capture the movement of the proteins in the said timeframe. The conformations obtained will then undergo k-means clustering using CPPTRAJ-Amber. This is done to cluster the conformations into 99 cluster and obtain one representative conformation from each cluster. The conformations were then used for Molecular Docking Analysis of the proteins and cathecin. In the following weeks, the top five (5) cathecin-protein interactions that had the highest binding energy will be identified and further analyzed. This will be done to determine the important interactions present between the proteins and cathecin. The results that will be obtained from this will give insights on how cathecin can potentially inhibit both HIV-1 Protease and Tat-Cyclin. This will give insights on how *Mentha cordifolia* plant extract can potentially prevent the viral replication in the virus' life cycle at a molecular level.

Keywords: HIV, *Mentha cordifolia*, molecular docking, molecular dynamics simulations

References

- Gangciangco, L.M.A, and Eustaquio, P.C. 2023. The State of the HIV Epidemic in the Philippines: Progress and Challenges in 2023. Tropical Medicine of Infectious Diseases, 8(5), 258.
- de Paz-Silava, S. L. M., Victoriano-Belvis, A. F. B., Gloriani, N. G., Hibi, Y., Asamitsu, K., & Okamoto, T. 2022. In Vitro Antiviral Activity of *Mentha cordifolia* Plant Extract in HIV-1 Latently Infected Cells Using an Established Human Cell Line. AIDS research and human retroviruses, 38(1), 64–72.

*Presenting author. E-mail address: jcbungar1@up.edu.ph (Jeffrey Clement C. Bungar)

S3.P115 Phytochemical profiling of *Acalypha australis* and screening for its most potent anti-inflammatory compounds with bio-affinity ultrafiltration LC-MS

Mingquan Guo^{a,b,*}, Rakotondrabe Tojofaniry Fabien^b, Minxia Fan^b

^a Cixi Institute of Biomedical Engineering, Ningbo Institute of Materials Technology and Engineering, Chinese Academy of Sciences, Ningbo 315300, China

^b CAS Key Laboratory of Plant Germplasm Enhancement and Specialty Agriculture, Wuhan Botanical Garden, Chinese Academy of Sciences, Wuhan 430074, China

Acalypha australis, commonly known as Copperleaf or Tiexian Cai, has served as an expectorant and for managing bacterial diseases and skin disorders. (Seebaluck et al., 2015). However, its phytochemical profiles along with its anti-inflammatory compounds have not yet been explored (Kim et al., 2020; Ma et al., 2017). In order to identify its potential anti-inflammatory compounds, an in vitro activity-guided fractionation followed by bio-affinity ultrafiltration LC-MS(UF/LC-MS) screening with the COX-2 enzyme, was successfully developed. By UPLC-QTOF/MS, 95 chemical compounds were characterized from *A. australis*, out of which 49 were reported for the first time in this species. Bioactivity-guided fractionation revealed that AAEA and AANB exerted similar COX- 2 inhibition with IC₅₀ values of 0.08 ± 0.01 µg/mL and 0.20 ± 0.02 µg/mL respectively, which showed better activity than the reference drug indomethacin with an IC₅₀ value of 0.47 ± 0.07 µg/mL. To further screening out its most potent compounds, the AAEA-3 fraction was subjected to UF/LC-MS targeting COX-2, and aurantiamide was then screened out and identified as the most promising COX- 2 ligand. Thereafter, the most potent ligand candidates were isolated and subjected to docking simulation and activity substantiation towards COX-2 to comprehend their probable mode of action. The docking interaction pf aurantiamide within the COX-2 active pocket exhibited a binding energy (BE) value of -9.56 Kcal/mol. The substantiation of aurantiamide activity towards COX-2 indicated a moderate inhibitory effect, with an IC₅₀ calculated to be 512.11 ± 21.80 µM.

Keywords: *Acalypha australis*, anti-inflammatory compounds, UF/LC-MS, COX-2

References

- Kim, H. J., Joe, H.I., Zhang, Z., et al., 2020. Anti-inflammatory effect of *Acalypha australis* L via suppression of NF-κB signaling in LPS-stimulated Raw 264.7 macrophages and LPS-induced septic mice. *Mol. Immunol.* 119, 123–131.
- Ma, Q., Wei, R., Liu, W., et al., 2017. Cytotoxic anthraquinones from the aerial parts of *Acalypha australis*. *Chem. Nat. Compd* 53, 949–952.
- Seebaluck, R., Gurib-Fakim, A., Mahomoodally, F., 2015. Medicinal plants from the genus *Acalypha* (Euphorbiaceae) a review of their ethnopharmacology and phytochemistry. *J. Ethnopharmacol.* 159, 137–157.

* Presenting author. E-mail address: guomingquan@nimte.ac.cn (Mingquan Guo)

S3.P116 The natural products magnetic resonance database (NP-MRD): comprehensive resource for natural products NMR data enabling discovery and understanding

John R. Cort^{a,b,*}, Jessica L. Bade^a, Roger G. Linington^c, Matthew Pin^c, Ella F. Poynton^c, Lloyd W. Sumner^d, James T. Koller^d, Zachary M. Tretter^d, Bharat Goel^d, David S. Wishart^e, Vasuk Gautam^e, Scott McKay^e, Harrison Peters^e, Tanvir Sajed^e, Brian L. Lee^e, Mark Berjanski^e

^a Earth and Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland, Washington, USA

^b Institute of Biological Chemistry, Washington State University, Pullman, Washington, USA

^c Department of Chemistry, Simon Fraser University, 8888 University Drive, Burnaby, British Columbia, Canada

^d Department of Biochemistry, University of Missouri, Columbia, Missouri, USA

^e Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

The Natural Products Magnetic Resonance Database (NP-MRD, np-mrd.org) is a database and repository for natural products and specialized metabolites NMR data. NP-MRD contains raw (time domain) data, derived data (chemical shift assignments, coupling constants) curated from the scientific literature or raw data, predicted data from density functional theory (DFT) calculations or machine learning, and simulated spectra. The database also provides structures, synonyms, search and other tools, links to other databases, and deposition interfaces. NP-MRD is an open database complying with FAIR (findable, accessible, interoperable, reusable) database principles. The NP-MRD mission is to create an enduring research resource to facilitate discovery and understanding in natural products and secondary metabolites research.

NP-MRD accepts raw data collected in support of novel structure elucidations and characterization of mixtures, through an intuitive deposition interface; deposition is now required by some journals and funding agencies. NP-MRD also accepts legacy data still residing on computers in individual laboratories. Curation of derived data (e.g. chemical shifts) published in the literature is problematic due to ambiguous or erroneous assignments, incorrect referencing, or typesetting mistakes. Curation itself, particularly from older literature in pdf format, is slow and may introduce additional errors. These limitations are less problematic as chemical shift predictions *via* both AI/ML and DFT calculations are increasingly accurate and can be used to validate published chemical shifts and locate and correct outliers. NP-MRD has predicted chemical shifts and simulated spectra for most natural products.

Insofar as this abstract describes a scientific database resource, rather than a research investigation, and in the interest of brevity, we do not have a conclusion. I have added an acknowledgment of the major funding source. References can be found on the database web site which appears at the beginning of the abstract.

NP-MRD is supported by the National Institutes of Health (USA), award 5U24AT010811-04.

* Presenting author. E-mail address: john.cort@pnnl.gov (John R. Cort)

S3.P117 Exploring the impact of moderate water stress on anthocyanin, flavonol glycosides and terpene dynamics across fruit development and ripening in *Pistacia lentiscus* L.

Lucrezia Muti^{a,*}, Diana Vanacore^b, Francesca Alderotti^a, Cassandra Detti^a, Giulia Goracci^a, Cecilia Brunetti^{b,a}, Francesco Ferrini^{a,b}, Anna Rita Bilia^c, Antonella Gori^{a,b}

^a Department of Agriculture, Food, Environment and Forestry (DAGRI), University of Florence, Sesto Fiorentino, Florence 50019, Italy

^b National Research Council of Italy, Institute for Sustainable Plant Protection (IPSP), Sesto Fiorentino, Florence and Turin 50019 and 10135, Italy

^c Department of Chemistry, University of Florence, Sesto Fiorentino, Firenze 50019, Italy

Pistacia lentiscus L. (Anacardiaceae) is a Mediterranean shrub whose fruits can be a source of new nutraceuticals (Milia et al., 2021). Our experiment evaluated the effects of water stress on the biochemical composition of *P. lentiscus* fruits at two ripening stages (November, T1, and December, T2). The experiment involved twelve potted plants, six of which (WW, well-watered) received adequate watering to pot capacity. The remaining plants underwent moderate water-stress treatment (WS, water-stressed), providing 70% of the Fraction of Transpirable Soil Water. At T1 and T2, Total Terpenes Content (TTC) was analyzed by GC-MS in the oil extracted from fruits. In the remaining oil cake, Anthocyanins and Flavonol Contents (TAC and TFC, respectively) were quantified through HPLC-DAD. Plants physiological status was monitored through measurements of stomatal conductance, chlorophyll fluorescence, leaf Relative Water Content, and leaf water potential, alongside leaf flavonol and chlorophyll indexes.

Results of oil cake biochemical analyses revealed that the best ripening time was T1, with TAC and TFC being 5.6 mg/g DW and 1.2 mg/g DW, respectively. In contrast, T2 was the best ripening stage for TTC (0.25 mg/g DW). Overall, TAC, TFC, and TTC were negatively influenced by the deficit-irrigation treatment at both ripening stages. However, a significant increase in β -myrcene in the oil collected from WS plants at T1 was observed. This emphasizes the potential use of a moderate water deficit treatment to enhance specific metabolite pathways. These findings offer valuable insights into strategic agricultural practices aimed at optimizing the nutraceutical potential of *P. lentiscus* oil and oil cake extracts.

Keywords: *Pistacia lentiscus*, fruits, ripening stages, oil, polyphenols

References

Milia, E., Bullitta, S. M., Mastandrea, G., Szotáková, B., Schoubben, A., Langhansová, L., Eick, S. 2021. Leaves and fruits preparations of *Pistacia lentiscus* L.: a review on the ethnopharmacological uses and implications in inflammation and infection. *Antibiotics*, 10(4), 425.

* Presenting author. E-mail address: lucrezia.muti@unifi.it (Lucrezia Muti)

S3.P118 Isolation and Characterization of novel compound from *Anaphalis triplinervis*, its HPTLC quantification and antioxidant potential

Ankush Sundriyal*, Yogita Dobhal

School of Pharmaceutical Sciences and Research, Sardar Bhagwan Singh University 248001, Balawala Dehradun, India

The genus *Anaphalis* (Asteraceae), comprised of herbaceous, annual and perennial flowering plants comprise of 110 species *Anaphalis* genus belongs to family Asteraceae, which is well reported for its medicinal importance. Traditionally many species of genus *Anaphalis* used as antiasthmatic, anticoughing, expectorant, sedative, antiepileptic, as anti-inflammatory agents and to cure many disorders (Murugan et al., 2021). The aim of the present study was Isolation and Characterization of novel compound from *Anaphalis triplinervis*, its HPTLC quantification and antioxidant potential. *Anaphalis triplinervis* (aerial part) was taken for extraction (successive solvent extraction scheme) and further ethyl acetate extract was taken for isolation by column chromatography. For isolation, the column was packed with silica gel (60-120) and eluted in toluene and ethyl acetate in gradient manner. Each fraction of was collected, concentrated pooled together according RF values and left for crystallization. Purification of the isolated compound was carried out by repeated recrystallization technique. The isolated compound was characterized by using various spectroscopic techniques, including UV, IR, NMR and MASS. Further qualitative and quantitative estimation (Murai et al., 2021) of isolated compound was done by HPTLC and antioxidant potential was done by DPPH assay (Akter et al., 2019). In summary, the isolated compound shows significant antioxidant efficacy, supported by comprehensive characterization efforts. This finding underscores its potential as a therapeutic agent against oxidative stress-related ailments.

Keywords: anaphalis triplinervis, characterization, isolation, antioxidant, chromatography

References

- Murugan, P., Murugan, C., Karthigeyan, K., 2021. Typification of twelve names in the genus *Anaphalis* DC. (Asteraceae). *Adansonia*. 43(17), 197-204.
- Murai, Y., Yangzom, R., Wangmo, C., Dorji, R., 2021. Phenolic Compounds from Three Alpine *Anaphalis* Species in Japan and Bhutan. *Bulletin of the National Museum of Nature and Science. Series B, Botany*. 47(2), 109-14.
- Akter, J., Hossain, MA., Takara, K., Islam, MZ., Hou, DX., 2019. Antioxidant activity of different species and varieties of turmeric (*Curcuma* spp): Isolation of active compounds. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*. 215, 9-17.

*Presenting author. E-mail address: ankushsundriyal@gmail.com (Ankush Sundriyal)

S3.P119 Trapping and detecting electrophilic metabolites of botanical natural products as glutathione conjugates

Richard B. van Breemen*, Jialin Liu, Ruth N. Muchiri

Department of Pharmaceutical Sciences. College of Pharmacy, Linus Pauling Institute, Oregon State University, 373 Linus Pauling Science Center, Corvallis, Oregon 97331 USA

Hepatic metabolism of drugs and natural products is usually a detoxification process that inactivates and increases polarity of xenobiotic compounds for excretion. However, oxidation by the cytochromes P450 sometimes forms electrophilic products that can react with cellular nucleophiles and cause toxicity. In the liver, the endogenous nucleophile glutathione (GSH) serves as a natural scavenger that deactivates reactive metabolites such as epoxides, arene oxides, alkyl halides, and quinones. Trapping of these short-lived metabolites with GSH can form stable products that may be analyzed using ultrahigh-pressure liquid chromatography-tandem mass spectrometry (UHPLC- MS/MS). Originally developed to detect reactive drug metabolites (Nikolic et al., 1999), we pioneered the application of this assay to test for the formation of electrophilic metabolites in botanical extracts. UHPLC-MS/MS (Huang, et al., 2015) was used to screen ethanolic extracts of 16 botanicals for hepatic cytochrome 450 formation of electrophiles that could be trapped as GSH conjugates. Botanical authentication and standardization of the test materials were carried out by the Botanical Safety Consortium, which is an international group of experts working to identify assays for the evaluation of the safety of botanicals used as dietary supplements. Botanicals included those with a safe history of human use as well as some with known toxic effects. Botanicals such as *Panax ginseng* and milk thistle were negative in this assay, whereas botanicals known to form toxic metabolites such as *Aristolochia* and kava produced GSH conjugates. Some toxic plants such as blue cohosh did not form GSH conjugates, indicating they have different mechanisms of toxicity.

Keywords: Metabolism, toxicity, safety, botanicals, glutathione

References

- Nikolic, D., Fan, P.W., Bolton, J.L., van Breemen, R.B., 1999. Screening for xenobiotic electrophilic metabolites using pulsed ultrafiltration-mass spectrometry. *Comb. Chem. High Throughput Screen.* 2(3):165-175.
- Huang, K., Huang, L., van Breemen, R.B., 2015. Detection of reactive metabolites using isotope-labeled glutathione trapping and simultaneous neutral loss and precursor ion scanning with ultra-high-pressure liquid chromatography, triple quadrupole mass spectrometry. *Anal. Chem.* 87, 3646-3654.

* Presenting author. E-mail address: richardvanbreemen@oregonstate.edu (Richard B. van Breemen)

S3.P120 Validation of artificial intelligence tools for expedited innovation and globalization of medicinal plants in the dietary supplement industry

The aim of this study was to identify a valid and reliable AI tool for fostering Innovation and globalization of medicinal plants in the dietary supplement industry with artificial intelligence.

We have surveyed several available tools. These include Google DeepMind, MIT Media Research Lab, Aradeepopsis, PlantCV, AyurAI and NutrifyGenie. Two of these tools were selected for this study - AyurAI and NutrifyGenie. These tools are not open access and require a paid subscription for their use. The AyurAI is available as mobile version while NutrifyGenie offers both desktop and mobile versions. The study is divided in three key steps. The first step involved the establishment of the theoretical background, outlining the essential standards for quantitative and qualitative effectiveness in AI tools for medicinal plants in the dietary supplement industry. The second step included a quantitative study to assess the performance of the identified tools. Specifically, the tools were evaluated against established standards for data management, accuracy, presentation, consistency, validation, and adaptability. The third step involved a qualitative study, comprehensively exploring the capabilities and features of each tool, with a particular focus on evidence verification, data exploration, regulatory compliance, feature analysis, automated label claims, integration of the supply chain, and more.

The experiment on NutrifyGenie yielded significant outcomes, identifying 320 potential ingredients for prevalent health conditions like diabetes, polycystic ovary syndrome, and osteoporosis backed by sound published clinical, mechanistic, safety, and efficacy data. The data included PubMed and peer-reviewed human clinical trials with validated findings. We found an average of three supporting pieces of evidence per ingredient. The NutrifyGenie tool also produced some systematic reviews and intellectual property data for evaluation. The NutrifyGenie contains regulatory compliance in eleven countries, offering overview of ingredient compliance. There are some other notable features available, including personalized label creation and integrated supply chain information for manufacturing a commercial product. These results underscore Nutrify Genie's transformative potential in facilitating evidence-based herbal ingredient formulation for self-care and eventually a fast-track option for product development. The AyurAI on the other hand is based more on the principals of Ayurveda with focus on 5 areas - personalized, preventive, predictive, participatory, and promotive aspects of the Ayurvedic therapies. This tool uses digital biomarkers, blood biochemistry, and genomics markers to reach a recommended product.

AyurAI and NutrifyGenie emerge as transformative AI-powered solutions for medicinal plants in the dietary supplement industry's global expansion. NutrifyGenie, operating as a hybrid model, caters to both B2B and partial B2C segments and is a regulatory complaint. AyurAI offers personalized wellness experiences directly to consumers through the B2C segment. These tools address challenges, enhance insights, and streamline industry processes, supported by accurate data sourcing and user-friendly interfaces.

* Presenting author. E-mail address: anand@cepham.com (Anand Swaroop)

S3.P121 Modified proanthocyanidin DESIGNER materials with dentin biomodification activity

Shu-Xi Jing^{a,b}, Walleska Liberato^c, Guilherme Neves^c, James B. McAlpine^{a,b}, Shao-Nong Chen^{a,b}
Ana K. Bedran-Russo^c, Guido F. Pauli^{a,b,*}

^a Pharmacognosy Institute and

^b Department of Pharmaceutical Science, College of Pharmacy;

^c Department of Restorative Dentistry, College of Dentistry; University of Illinois Chicago, Chicago, IL, 60612, USA

Proanthocyanidins (PACs, *syn.* condensed tannins) are oligomers to polymers of flavan-3-ols, a group of plant constituents that impact humans via their health, nutritional, and agronomic value. To date, quantitative analysis of PAC-rich materials depend on colorimetric assays or phloroglucinolysis/thiolysis combined with UV-HPLC analysis - all methods with limited accuracy, extended sample preparation, degradation, and/or need for identical reference standards. In the past decade, high-field and low-temperature NMR as well as via precise assignments using computer-aided ¹H full spin analysis (HifSA) have jointly advanced the analysis of intact, genuine PACs. Towards our broader goal of developing new plant-sourced biomaterials that modulate the mechanical properties of dental tissue for clinical interventions, we have characterized 12 new PAC DESIGNER (Depletion and Enrichment of Select Ingredients Generating Normalized Extract Resources) materials. The DESIGNER approach involves centrifugal partition chromatography (CPC) or size-exclusion chromatography (SEC) for the selective enrichment of trimeric and tetrameric PACs. Moreover, the rare but biologically interesting all-A-type PAC DESIGNER materials were successfully produced via conversion of their natural AB-type precursors via an improved method using a free radical oxidative reagent. Quantification of the total PAC content by quantitative NMR (qNMR) utilized an internal calibrant and diol-HPLC to ensure quality and stability of the PAC DESIGNER materials. Diol-HPLC deduced the polymerization profiles, and qNMR determined the total PAC content to range from 61% to 95%. This highlights the complementarity of diol-HPLC and qNMR to accurately assess the amount of PACs across a range of concentrations as well as PAC stability in the DESIGNER materials. Presented dental bioactivity data will demonstrate the unique ability of these materials to enhance dental biomechanical properties comparable to isolated PACs. The newly available quantitative methodology paves the way to standardized DESIGNER materials via rigorous quality control of PAC-based dental (pre-)clinical interventions.

* Presenting author. E-mail address: gfp@uic.edu (Guido F. Pauli)

S3.P122 Antimicrobial activity of 2'-hydroxychalcones with a chlorine atom and their glycosides

Agnieszka Krawczyk-Łebek^{a,*}, Barbara Żarowska^b, Monika Dymarska^a, Tomasz Janeczko^a, Edyta Kostrzewa-Susłow^a

^a Department of Food Chemistry and Biocatalysis, Faculty of Biotechnology and Food Science, Wrocław University of Environmental and Life Sciences, 50-375 Wrocław, Poland

^b Department of Biotechnology and Food Microbiology, Faculty of Biotechnology and Food Science, Wrocław University of Environmental and Life Sciences, 50-375 Wrocław, Poland

Chalcones are intermediate products in the biosynthesis of flavonoids, which possess a wide range of biological properties, including antimicrobial activity. The introduction of a chlorine atom and the glucosyl moiety into their structure may increase their bioavailability and bioactivity (Lin et al. 2002, Guvenalp et al. 2015). To receive such flavonoids we combined chemical and biotechnological methods (Krawczyk-Łebek et al. 2021). Firstly, we obtained four flavonoid aglycones in the Claisen- Schmidt condensation reaction, i.e., 2-chloro-2'-hydroxychalcone, 3-chloro-2'-hydroxychalcone, 4-chloro-2'-hydroxychalcone, and 5'-chloro-2'-hydroxychalcone. Secondly, we biotransformed them in the cultures of two entomopathogenic filamentous fungi strains, i.e., *Isaria fumosorosea* KCH J2 and *Beauveria bassiana* KCH J1.5 in order to obtain their glycosides. The structures of the resulting compounds were determined by NMR spectroscopy and confirmed by MS spectroscopy. The received compounds along with 2'-hydroxychalcone were used for the tests of their antimicrobial activity against three bacteria strains *Escherichia coli* 10536 (Gram-), *Pseudomonas aeruginosa* DSM 939 (Gram-), *Staphylococcus aureus* DSM 799 (Gram+), one strain of yeast *Candida albicans* DSM 1386, and three strains of lactic acid bacteria *Lactococcus acidophilus* KBiMZ 01 (Gram+), *Lactococcus rhamnosus* GG (Gram+), *Streptococcus thermophilus* KBM – 1 (Gram+). In general, chlorinated chalcones were more effective in the inhibition of the tested microbial strains than their unchlorinated counterparts and aglycones were similarly or a little more effective than their glycosides. The highest antibacterial potential among the tested compounds was demonstrated by 5'-chlorodihydrochalcone 2'-O-β-D-(4"-O-methyl)-glucopyranoside.

Keywords: chalcones with a chlorine atom, flavonoid glycosylation, *Isaria fumosorosea*, *Beauveria bassiana*, antimicrobial activity

References

- Guvenalp, Z.; Ozbek, H.; Karadayi, M.; Gulluce, M.; Kuruuzum-Uz, A.; Salih, B.; Demirezer, O., 2015. Two Antigenotoxic Chalcone Glycosides from *Mentha Longifolia* Subsp. *Longifolia*. *Pharm. Biol.* 53(6), 888-896
- Lin, Y. M.; Zhou, Y.; Flavin, M. T.; Zhou, L. M.; Nie, W.; Chen, F. C., 2002. Chalcones and Flavonoids as Anti- Tuberculosis Agents. *Bioorg. Med. Chem.* 10(8), 2795-2802.
- Krawczyk-Łebek, A.; Dymarska, M.; Janeczko, T.; Kostrzewa-Susłow, E., 2021. New Glycosylated Dihydrochalcones Obtained by Biotransformation of 2'-Hydroxy-2-Methylchalcone in Cultures of Entomopathogenic Filamentous Fungi. *Int. J. Mol. Sci.* 22(17), 9619.

Acknowledgements

The work was created as a result of research project No 2021/41/N/NZ9/01195 financed by the National Science Center.

* Presenting author. E-mail address: agnieszka.krawczyk-lebek@upwr.edu.pl (Agnieszka Krawczyk-Łebek)

S3.P123 Novel ardisiphenol D derivatives 1279P1 and 1279P2: promising anti-colorectal cancer agents targeting the PI3K/Akt pathway and lipogenesis enzymes

Bai-Sen Chen*, Wai-I Chik, Hong-Jie Zhang

School of Chinese Medicine, Hong Kong Baptist University, 7 Baptist University Road, Kowloon Tong, Hong Kong SAR, People's Republic of China

This study investigates the potential efficacy of two novel derivatives of ardisiphenol D (**1**) as anti-colorectal cancer agents with improved therapeutic outcomes and decreased toxicity profiles in comparison to the parent compound. The efficacy of the novel ardisiphenol D derivatives, 1279P1 (**2**) and 1279P2 (**3**) (Fig.1), has been validated in anti-colorectal cancer treatment through comprehensive *in vitro* and *in vivo* experiments. Both derivatives have been shown to effectively reduce phosphorylated PI3K and AKT protein expression, inhibiting the critical PI3K/AKT signaling pathway associated with cell transformation, tumor progression, and drug resistance (Bahrami et al., 2018).

Furthermore, both compounds exhibit significant inhibitory effects on key lipogenic enzymes ACCA and FASN, essential for *de novo* lipogenesis (Röhrig et al., 2016), which could disrupt cell energy supply, particularly in cancer cells (Krauß et al., 2022). This dual mechanism of action highlights their promising role as potent inhibitors of cancer-promoting pathways. Notably, the cytotoxic effects of **2** and **3** on colorectal cancer cells underscore their therapeutic potential. The synergistic blockade of the PI3K/Akt pathway and suppression of lipogenesis enzymes present a novel mechanism for their anti-cancer activity.




Fig. 1. Structures of ardisiphenol D (**1**), 1279P1 (**2**), and 1279P2 (**3**)

These findings underscore the synergy between PI3K/Akt pathway blockade and lipogenesis enzyme suppression delineates a novel mechanism for the anti-cancer potential of ardisiphenol D derivatives. These results warrant further exploration and clinical translation of **2** and **3** as potential agents for innovative colorectal cancer therapy. The work was financially supported by Hong Kong Baptist University, Research Committee, Initiation Grant – Faculty Niche Research Areas (IG-FNRA) 2021/22, Hong Kong (RC-IG-FNRA/21-22/SCM/01).

Keywords: Ardisiphenol D derivatives, PI3K/AKT pathway, *de novo* lipogenesis, anti-colorectal cancer

References

- Bahrami, A., Khazaei, M., Hasanzadeh, M., ShahidSales, S., Mashhad, M.J., Farazestanian, M., Sadeghnia, H.R., Majid Rezayi, Maftouh, M., Hassanian, S.M., Avan, A., 2018. Therapeutic potential of targeting PI3K/AKT pathway in treatment of colorectal cancer: rational and progress. *J. Cell. Biochem.* 119, 2460-2469.
- Krauß, D., Fari, O., Sibilia, M., 2022. Lipid metabolism interplay in CRC—An update. *Metabolites* 12, 213.
- Röhrig, F.; Schulze, A., 2016. The multifaceted roles of fatty acid synthesis in cancer. *Nat. Rev. Cancer* 16, 732–749.

* Presenting author. E-mail address: 22483136@life.hkbu.edu.hk (Bai-Sen Chen)

S3.P124 Antifibrotic activity study of natural stilbenoids from *Dendrobium nobile* Lindl

Chu Shing Lam*, Hong-Jie Zhang

School of Chinese Medicine, Hong Kong Baptist University, 7 Baptist University Road, Kowloon Tong, Hong Kong SAR, People's Republic of China

This presentation will cover the biological evaluation and mechanism of action study of dihydro-resveratrol (**1**), a natural stilbenoid from *Dendrobium nobile* Lindl. (Orchidaceae). The compound can be found in many different plant sources, while it is also regarded as a microbial metabolite of resveratrol (**2**) by gut microbiota. Dihydro-resveratrol possesses higher solubility and bioavailability than resveratrol (Li et al., 2022), suggesting that **2** possesses better potential in developing as an antifibrotic agent than **1**.

We compared the cytotoxicity of **1** and **2** in different pancreatic cell lines, suggesting that **1** possesses lower cytotoxicity than **2** in primary murine acini cells (Tsang et al., 2022), both human and rat pancreatic stellate cells (HPSC, LTC-14) and murine pancreatic beta cells (NIT-1) (Fig.1).



		Cytotoxicity: IC ₅₀ (μM)			
Compounds / Cell lines		LTC-14	HPSC	NIT-1	Primary acini cells ¹
	Dihydro-resveratrol (1)	384.6	>200	~125	>250
	Resveratrol (2)	63.4	>200	~50	<250

Fig. 1. Structures and cytotoxicity of dihydro-resveratrol (**1**) and resveratrol (**2**)

Our *in vitro* studies revealed that **1** is a potent antifibrotic agent against pancreatic fibrosis and chronic pancreatitis by inhibiting the activation of pancreatic stellate cells. Our *in vivo* findings indicated that **1** could significantly ameliorate pancreatic fibrosis and macrophage infiltration and protect the islets of the pancreas in a murine chronic pancreatitis model. Moreover, **1** is shown to have an anti-inflammatory effect on macrophage cell lines and a protective effect on pancreatic beta-cell and pancreatic acini cell lines against apoptosis and ferroptosis, respectively. Preliminary mechanistic studies and HuProt human proteome array suggested that **1** modulates PI3K/AKT, NFκB, and MAPK cell signaling pathways. The work described in this study was financially supported by the Food and Health Bureau, the Government of the Hong Kong Special Administrative Region (grant number COVID190214).

Keywords: *Dendrobium nobile*, dihydro-resveratrol, resveratrol, pancreatic fibrosis, chronic pancreatitis

References

- Li, F., Han, Y., Wu, X., Cao, X., Gao, Z., Sun, Y., Wang, M., Xiao, H., 2022. Gut microbiota-derived resveratrol metabolites, dihydroresveratrol and lunularin, significantly contribute to the biological activities of resveratrol. *Front. Nutr.* 9, 912591.
- Tsang, S.W., Guan, Y.F., Wang, J., Bian, Z.X., Zhang, H.J., 2016. Inhibition of pancreatic oxidative damage by stilbene derivative dihydro-resveratrol: implication for treatment of acute pancreatitis. *Sci. Rep.* 6, 22859.

* Presenting author. E-mail address: 18482619@life.hkbu.edu.hk (Chu Shing LAM)

S3.P125 Planar chromatography method development for the herbal metabolomic study on *Achillea millefolium* (Asteraceae)

Simon J. Jackson^{a,*}, Paolo Goi^a, Dario Fornara^b, Tim Yeomans^c, Katie Shiels^c, Emma Murphy^c

^a Davines Group - Botanical Research Department, Via Don Angelo Calzolari 55/A 43126 Parma, Italy

^b Davines Group - EROC (European Regenerative Organic Centre), Via Don Angelo Calzolari 55/A 43126 Parma, Italy

^c Shannon ABC, TUS (Technical University Shannon), Limerick, Ireland

This study covers both quantitative and qualitative analysis of extracts of *Achillea millefolium*, using planar chromatography. Comparing and contrasting extracts of biomass grown regeneratively in the EROC^b, with organic, and wild harvested commercial samples. The isolation and structure elucidation will be undertaken separately, but this study will focus on the known reference standards, Chamazulene (**1**), Salicylic acid (**2**), Quercetin (**3**) and using these marker compounds for qualitative analysis.

Achillea as a source of phytochemicals for the cosmetic Industry has gained increased interest (Becker et al, 2016). This study focuses on flowering tops of the *Achillea* plants and comparative samples were obtained commercially and grown regeneratively at EROC. Comparing and contrasting monographs published in the British Herbal Pharmacopeia (BHP 1996) and monographs from Herbal drugs and phytopharmaceuticals (Bisset et al 1994), and the European Pharmacopeia. We determined a simple method to identify qualitative parameters for commercial batch sampling of extracts. Working with Shannon ABC^c using CAMAG HPTLC apparatus, we developed a solvent system of Toluene: Ethyl Acetate: Formic Acid (70:30:1) using a stationary phase of Merck HPTLC Plates, silica gel 60F 254 and using Methanol as a sample solvent and pre dosage volume of 0.2 µl visualising with VisionCats software, white light, and UV 254nm and 366 nm.

As part of our ongoing phytochemical research project developing simple monographs for commercially interesting herbal extracts the aim is to use these results in producing open access best practice for laboratory guidance documentation.




Fig. 1. Structure of Chamazulene **1** Fig. 2. Structure of Salicylic acid **2**. Fig. 3. Structure of Quercetin **3**

Keywords: *Achillea millefolium*, Asteraceae, HPTLC, planar chromatography, Chamazulene, Salicylic acid, Quercetin

References

- Becker, L.C et al 2016. Safety Assessment of *Achillea millefolium* as used in cosmetics. International Journal of Toxicology, Vol. 35(Supplement 3) 5S-15S
- British Herbal Pharmacopoeia (BHP). 1996. Exeter, U.K.: British Herbal Medicine Association.
- Wichtl, M. and N.G. Bisset (eds.). 1994. Herbal Drugs and Phytopharmaceuticals. Stuttgart: Medpharm Scientific Publishers.

* Presenting author. E-mail address: s.jackson@davines.it (Simon James Jackson)

S3.P126 Metabolomics and microbiomics analysis reveals the dynamic changes during *Citrus depressa* Hayata fermentation

Ta-Wei Liu*, Ching-Kuo Lee

School of Pharmacy, College of Pharmacy, Taipei Medical University, Taipei, 11042, Taiwan

In recent years, biotransformation has emerged as a new source for small molecule natural drugs, expanding chemical diversity. Many traditional Chinese medicines are produced through biotransformation to generate active ingredients. Representative examples include Chen-Pi, which undergoes microbial transformation through aging to produce more flavonoids and volatile components (Wang et al., 2015). *Citrus depressa* Hayata is renowned for its high content of polymethoxyflavones (PMFs) and used as raw materials of Chen-Pi in early Taiwan (Wu et al., 1983). Therefore, this study aims to explore the interaction between metabolites produced during the fermentation process of *C. depressa* and microorganisms, and to develop a more stable fermentation method. The changes in microbiota within *C. depressa* endophyte fermentation products at each time point were analyzed using 16S rRNA, and analyzed for differential metabolites using non-targeted LC-MS/MS. Additionally, *Bacillus amyloliquefaciens* was isolated from endophyte fermentation samples, and the same method was employed for metabolomic analysis as with endophyte fermentation. In metabolomic analysis, carbohydrates, amino acids, and PMFs were the three most abundant categories among differential metabolites, and these compounds were subjected to quantitative analysis and metabolic pathway analysis. In the end, we found that *B. amyloliquefaciens* can produce a large amount of PMFs more rapidly and steadily, including nobiletin, 5-demethylnobiletin, and tangeretin, reaching concentrations of 8, 0.8, and 3 µg/mg, respectively, in 12 weeks.

Keywords: *Citrus depressa* Hayata, metabolomics, microbiomics, LC-MS/MS

References

- Wang, F., Zhang, X., Lu, J.Y., Mei, G.R., Chen, H.P., Chen, L., Liu, Y.P., 2015. Study of reason of “the older, the better” of Pericarpium Citri Reticulatae (PCR) which contributed increase of flavonoids. China Journal of Chinese Materia Medica, 01 Dec 2015, 40(24):4890-4896
- Wu, T.S., Hiroshi, F., Kuoh, C.S., 1983. The constituents of *Citrus depressa*. Isolation and structure elucidation of new acridone alkaloids from *Citrus* genus. Chem. Pharm. Bull. 31 (3), 895–900.

* Presenting author. E-mail address: d301110010@tmu.edu.tw (Ta-Wei, Liu)

S3.P127 LC-MS analysis-based molecular networking for identifying anthelmintic compounds in *Terminalia leiocarpa*, a traditional plant used in livestock farming in West Africa

Esaïe Tchetan^{a,b,*}, Sergio Ortiz^b, Pascal Abiodoun Olounladé^a, Kristelle Hughes^b, Patrick Laurent^c, Erick Virgile Bertrand Azando^a, Sylvie Mawule Hounzangbe-Adoté^a, Fernand Ahokanou Gbaguidi^d, Joëlle Quetin-Leclercq^b

^a Laboratoire d'Ethnopharmacologie et de Santé Animale, Faculté des Sciences Agronomiques, Université d'Abomey-Calavi, Cotonou 01 BP 526, Benin

^b Pharmacognosy Research Group, Louvain Drug Research Institute, Université catholique de Louvain (UCLouvain), Avenue E. Mounier, B1.72.03, B-1200 Brussels, Belgium

^c Laboratory of Neurophysiology, ULB Neuroscience Institute (UNI), Université Libre de Bruxelles (ULB), 808 route de Lennik, 1070, CP 601, Brussels, Belgium

^d Laboratoire de Chimie Organique et Chimie Pharmaceutique, UFR Pharmacie, Faculté des Sciences de la Santé, Université d'Abomey-Calavi, Cotonou 01 BP 188, Benin

Terminalia leiocarpa (Combretaceae) is a medicinal plant widely used in ethnoveterinary medicine to treat digestive parasitosis (Tchetan et al., 2021). Its extracts were shown to be active against gastrointestinal nematodes of domestic ruminants (Ndjonka et al., 2012; Tchetan et al., 2022). The objective of our study was to identify compounds responsible for this activity. Open column fractionation was performed, and the activity of the fractions was assessed *in vitro* on *Haemonchus contortus* and *Caenorhabditis elegans* as well as their cytotoxicity on WI38 fibroblasts. Two fractions were the most active on both nematode models and less cytotoxic. LC-MS/MS analysis and manual dereplication coupled to molecular networking allowed identification of the main compounds: ellagic acid and derivatives, gallic acid, astragalin, rutin, quinic acid, and fructose. Other potentially identified compounds such as shikimic acid, 2,3-(*S*)-hexahydroxydiphenyl-D-glucose or an isomer, quercetin-3-O-(6-O-galloyl)- β -D-galactopyranoside or an isomer, and rosamultin or an isomer are reported in this plant for the first time. Evaluation of the anthelmintic activity of the major compounds available showed that ellagic and gallic acids were the most effective in inhibiting the viability of *C. elegans*. Ellagic acid and its derivatives may have additive or synergistic effects when combined, but other unidentified compounds could also be implicated in the observed activity.

Keywords: anthelmintic activity, molecular networking, *Terminalia leiocarpa*, *Haemonchus contortus*, *Caenorhabditis elegans*

References

- Ndjonka, D., Ajonina-Ekoti, I., Djafsia, B., Lüersen, K., Abladam, E., Liebau, E. 2012. *Anogeissus leiocarpus* extract on the parasite nematode *Onchocerca ochengi* and on drug resistant mutant strains of the free-living nematode *Caenorhabditis elegans*. Vet. Parasitol., 190, 136–142.
- Tchetan, E., Olounlade, A. P., Houehanou, T. D., Azando, E. V. B., Kaneho, J. A., Houinato, M. R. B. 2021. Ethnoveterinary knowledge of sheep and goat farmers in Benin (West Africa): effect of socioeconomic and environmental factors. Heliyon, 7(7), e07656.
- Tchetan, E., Olounladé, P. A., Azando, E. V. B., Khaliq, H. A., Ortiz, S., Hounbeme, A., & Quetin-Leclercq, J. 2022. Anthelmintic Activity, Cytotoxicity, and Phytochemical Screening of Plants Used to Treat Digestive Parasitosis of Small Ruminants in Benin (West Africa). Animals, 12(19), 2718.

* Presenting author. E-mail address: esaïetchetan@gmail.com <mailto:kskalicka@pharmacognosy.org>(Esaïe Tchetan)

S3.P128 Integration of Wnt-inhibitory activity and metabolomics structural novelty results to discovery novel bioactive natural products: new bicyclo[3.3.1]non-3-ene-2,9-diones from the leaves of *Hymenocardia punctata*

Luis-Manuel Quiros-Guerrero^{a,b,*}, Laurence Marcourt^{a,b}, Nathareen Chaiwangrach^c, Alexey Koval^d, Emerson Ferreira Queiroz^{a,b}, Bruno David^e, Antonio Grondin^e, Vladimir L. Katanaev^{d,f}, Jean-Luc Wolfender^{a,b}

^a Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, CMU, 1211 Geneva, Switzerland

^b School of Pharmaceutical Sciences, University of Geneva, CMU, 1211 Geneva, Switzerland

^c Centre of Excellence in Cannabis Research, Department of Pharmaceutical Chemistry and Pharmacognosy, Faculty of Pharmaceutical Sciences, Naresuan University, Phitsanulok 65000, Thailand

^d Department of Cell Physiology and Metabolism, Translational Research Centre in Oncohaematology, Faculty of Medicine, 1211 Geneva, Switzerland

^e Green Mission Department, Herbal Products Laboratory, Pierre Fabre Research Institute, Toulouse, France

^f School of Medicine and Life Sciences, Far Eastern Federal University, 690091 Vladivostok, Russia

In natural products (NP) research, efficiently prioritizing samples in natural plant extract (NEs) libraries is crucial to accelerate the discovery of original bioactive NPs. In this study a biodiverse collection of 1600 NEs, previously analyzed by UHPLC-HRMS² metabolite profiling (Allard *et al.* 2023) was screened for Wnt pathway regulation (Fig. 1). The results of the biological screening initially drove the selection of NEs. Only non-toxic NEs with an inhibitory IC₅₀ ≤ 5 µg/mL were considered, resulting in a subset of 30 NEs. To increase the chance of finding structurally novel bioactive NPs, *Inventa*, a computational tool for automated scoring of NEs based on structural novelty was used to mine the HRMS² analysis and dereplication results (Quiros-Guerrero *et al.* 2022). This procedure, which is aimed at targeting NE with novel structures, resulted in a 2nd step selection of 4 NEs out of the 1st step post-screening selection of 30 bioactive NEs. The most promising candidate was *Hymenocardia punctata* (Phyllanthaceae). Further phytochemical investigations of this species resulted in the targeted isolation of three known prenylated flavones and ten novel bicyclo[3.3.1]non-3-ene-2,9- diones, named *Hymenotamayonins*. Assessment of the Wnt inhibitory activity of these compounds revealed that two prenylated flavones and three novel bicyclic compounds showed interesting activity without apparent cytotoxicity. This study highlights the potential of combining *Inventa*'s structural novelty scores with biological screening results to effectively discover novel bioactive NPs in large NE collections (Quiros-Guerrero *et al.* 2024).



Fig. 1. Overview of the general strategy for the selection of promising NEs for the discovery of structurally novel bioactive NPs in collections of NEs.

Keywords: Natural products, Wnt-pathway modulators, structural novelty discovery, 'Inventa' scoring, *Hymenocardia punctata*

References

- Allard, P-M., Gaudry, A., Quirós-Guerrero, L-M., Rutz, A., Dounoue-Kubo, M., Walker, T., Defossez, E. et al. 2023. Open and Reusable Annotated Mass Spectrometry Dataset of a Chemodiverse Collection of 1,600 Plant Extracts. *GigaScience* 12. <https://doi.org/10.1093/gigascience/giac124>.
- Quiros-Guerrero, L-M., Nothias, L-F., Gaudry, A., Marcourt, L., Allard, P-M., Rutz, A., David, B., Ferreira-Queiroz, E., Wolfender, J-L. 2022. Inventa: A Computational Tool to Discover Structural Novelty in Natural Extracts Libraries. *Frontiers in Molecular Biosciences* 9: 1028334. <https://doi.org/10.3389/fmolb.2022.1028334>.
- Quiros-Guerrero, L-M., Marcourt, L., Chaiwangrach, N., Koval, A., Ferreira-Queiroz, E., David, B., Grondin, A., Katanaev, V., Wolfender, J-L. 2024. Integration of Wnt-Inhibitory Activity and Structural Novelty Scoring Results to Uncover Novel Bioactive Natural Products: New Bicyclo[3.3.1]non-3-Ene-2,9-Diones from the Leaves of *Hymenocardia Punctata*. *ChemRxiv*. <https://doi.org/10.26434/chemrxiv-2024-96nz6>.

Acknowledgements

The authors are thankful to the Swiss National Science Foundation (SNSF) for the financial support of this project (no. CRSII5_189921/1).

* Presenting author. E-mail address: luis.guerrero@unige.ch (Luis-Manuel Quiros-Guerrero)

S3.P129 Enhancing prioritization strategies of natural extracts: integrating heterogeneous data from metabolomics datasets and biological screenings into knowledge graphs

Luis-Manuel Quiros-Guerrero^{a,b,*}, Frederic Burdet^c, Olivier Kirchhoffer^{a,b}, Paola Haemmerli^{a,b}, Louis-Felix Nothias^{a,b,d}, Pierre-Marie Allard^{a,b,e}, Arnaud Gaudry^{a,b}, Jahn Nitschke^f, Nabil Hanna^f, Sébastien Moretti^c, Robin Engler^c, Florence Mehl^c, Antonio Grondin^g, Bruno David^g, Chunyan Wu^h, Erick Carreiraⁱ, Thierry Soldati^f, Christian Wolfrum^h, Marco Pagni^c, Jean-Luc Wolfender^{a,b}

^a Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, 1211 Geneva 4, Switzerland

^b School of Pharmaceutical Sciences, University of Geneva, 1211 Geneva 4, Switzerland

^c Vital-IT, SIB Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland

^d Université Côte d'Azur, Institut de Chimie de Nice, Campus Valrose, Nice, France

^e Department of Biology, University of Fribourg, 1700 Fribourg, Switzerland

^f Department of Biochemistry, Faculty of Science, University of Geneva, 1211 Geneva, Switzerland ^g Green Mission Department, Herbal Products Laboratory, Pierre Fabre Research Institute, Toulouse, France

^h Department of Health Sciences and Technology, ETHZ, 8092 Zürich, Switzerland

ⁱ Department of Chemistry and Applied Biosciences, ETHZ, 8092 Zürich, Switzerland

Integrating heterogeneous data in natural products (NPs) research, like metabolite profiling (UHPLC-HRMS²), annotations and biological screening results is challenging. Typically, data is processed in an *aligned* manner to create feature tables across samples for explorative techniques like Molecular Networking. This facilitates precise comparisons within homogeneous sets. However, integration of new samples and comparisons across batches are hindered by experimental variations. To address these limitations, novel *sample-centric* approaches have been developed to explore diverse datasets of Natural Extracts (NEs) over time (Gaudry *et al.* 2022). Additionally, the need for a more comprehensive *knowledge-driven* framework integrating all type of data led to the implementation of Knowledge Graphs (KG) in metabolomic projects (Gaudry *et al.* 2023). KG provides structured representations of complex datasets through RDF semantic web data standardization as subject-predicate-object triples (RDF-SemanticWebStandards 2014). This facilitates exploration and interconnects information (Caufield *et al.* 2023). For example, it enables to link sample's taxonomy, spectral annotations, bioactivity with prior information. In this context a multidimensional KG containing over 200 million triples was generated in the frame of a collaborative project incorporating UHPLC-HRMS² data of over 3,000 NEs, fractions and pure compounds; taxonomical information, chemo-informatics results, and bioassay outcomes for tuberculosis, obesity, anticancer, and antiviral models. We will describe key element for the constitution of the KG based on NEs experimental data. Additionally, several examples will showcase the potential of using SPARQL queries (SPARQLQueryLanguageforRDF 2018) to explore this KG, demonstrating their efficacy in guiding sample selection and accelerating NP discovery.

Keywords: knowledge graph, RDF, sample-centric, SPARQL query, drug discovery

References

- Caufield, J. Harry, Tim Putman, Kevin Schaper, Deepak R. Unni, Harshad Hegde, Tiffany J. Callahan, Luca Cappelletti, *et al.* 2023. KG-Hub-Building and Exchanging Biological Knowledge Graphs. *Bioinformatics* 39 (7). <https://doi.org/10.1093/bioinformatics/btad418>.
- Gaudry, Arnaud, Florian Huber, Louis-Félix Nothias, Sylvain Cretton, Marcel Kaiser, Jean-Luc Wolfender, and Pierre-Marie Allard. 2022. MEMO: Mass Spectrometry-Based Sample Vectorization to Explore Chemodiverse Datasets. *Frontiers in Bioinformatics* 2: 842964.
- Gaudry, Arnaud, Marco Pagni, Florence Mehl, Sébastien Moretti, Luis-Manuel Quiros-Guerrero, Adriano Rutz, Marcel Kaiser, *et al.* 2023. A Sample-Centric and Knowledge-Driven Computational Framework for Natural Products Drug Discovery. *ChemRxiv*, July. <https://doi.org/10.26434/chemrxiv-2023-sljbt>.
- "RDF-SemanticWebStandards" February 25, 2014. <https://www.w3.org/RDF/>.
- "SPARQLQueryLanguageforRDF" January 15, 2018. <https://www.w3.org/TR/rdf-sparql-query/>

Acknowledgements: The authors are thankful to the Swiss National Science Foundation (SNSF) for the financial support of this project (no. CRSII5_189921/1).

* Presenting author. E-mail address: luis.guerrero@unige.ch (Luis-Manuel Quiros-Guerrero)

S3.P130 The impact of various technological parameters and a place of origin of *Nigella sativa*. L seeds on the thymoquinone content in black cumin oil

Benita Hryć^{a,*}, Michał Musiał^b, Elwira Sieniawska^a

^a Department of Chemistry of Natural Products, Medical University of Lublin, 1 Chodzki Str., 20-093 Lublin, Poland

^b Michał Musiał Agricultural Farm, Wielkie Rychnowo 67, 87-410 Kowalewo Pomorskie, Poland

A black cumin oil has a great therapeutical potential and is commonly used as a food supplement. Very important feature of black cumin oil is content of thymoquinone (**1**) (Fig. 1), which can vary from 0,45% to 4,57% depending on a genotype of *Nigella sativa* (Telci et al., 2023). Thymoquinone exerts very broad spectrum of effects among others: antioxidant, anti-inflammatory, antimicrobial, immunomodulatory, antineoplastic, antidiabetic, hepatoprotective and neuroprotective activity (Darakhshan et al., 2015).




Fig. 1. Structure of thymoquinone **1**

The aim of this study was to evaluate the impact of various technological features of oil making procedure and a place of seeds origin on the content of thymoquinone. For this purpose, the validated HPLC/PDA method was used. The validation procedure included such elements as repeatability, intermediate precision, linearity, accuracy, LOD and LOQ assessment according to ICH guidelines. The biggest impact on the **1** content has the seeds humidity. The greatest difference was observed between 10 and 15/16% of water content in the seeds. There was almost no difference between sedimentation and centrifuging procedure. The type of press machine had smaller influence than seeds humidity. Increase in pressing temperature from 40 to 48°C had small impact on the **1** amount in oil. The place of origin was the most important factor. In conclusion the biggest impact on the **1** content in the oil had the black cumin origin. Nevertheless, the technological parameters especially the seeds humidity can have significant influence on the quality of the *Nigella sativa* oil represented by thymoquinone level.

Keywords: *Nigella sativa*, black cumin oil, thymoquinone, validation, HPLC

References

- Darakhshan, S., Bidmeshki Pour, A., Hosseinzadeh Colagar, A., & Sisakhtnezhad, S., 2015. Thymoquinone and its therapeutic potentials. *Pharmacol Res*, 95–96, 138–158.
- Telci, İ., Özek, T., Demirtaş, İ., Özek, G., Yur, S., Ersoy, S., Yasak, S., Gül, F., & Karakurt, Y., 2023. Studies on black cumin genotypes of Türkiye: Agronomy, seed and thymoquinone yields. *J Appl Res Med Aromat Plants*, 35(May), 100494.

* Presenting author. E-mail address: benita.hryc@umlub.pl (Benita Hryć)

S3.P131 Unravelling enzymatic hydrolysis inhibitors in lignocellulosic biomass samples: a combined LC-HRMS/MS and NMR multiblock modelling approach.

Ian Ramtanon^{a,*}, Marion Lacoue-Nègre^{a,b}, Alexandra Berlioz-Barbier^a, Simon Remy^c, Agnès Le Masle^a, Jean-Hugues Renault^c

^a IFP Energies nouvelles, Rond-point de l'échangeur de Solaize, BP 3, 69360 Solaize, France

^b ChemHouse Research Group, Montpellier, France

^c Université de Reims Champagne-Ardenne, CNRS, ICMR 7312, 51097, Reims, France

Considering the current environmental challenges, transformation of lignocellulosic biomass into bioethanol offers a sustainable alternative to petroleum fuels (Wyman, 1999). The pretreatment step, applied in some industrial processes, generates aqueous phases called hydrolysates. These complex mixtures contain a variety of oxygenated compounds which may interfere with the enzymatic hydrolysis (EH) step and hinder the conversion of cellulosic feedstock into glucose (Kim, 2018). The confident identification of these EH “inhibitors” is therefore crucial for the optimization of the bioconversion dynamics.

LC-HRMS and NMR spectroscopy are complementary techniques for the molecular characterization of complex samples (Bhinderwala *et al.*, 2018). Indeed, HRMS sensitivity allows detection of a large number of compounds while NMR provides crucial information about species' abundance. Moreover, trend determination and biomarker identification from these large analytical datasets have been markedly improved by the use multivariate analyses (Deng *et al.*, 2016). In this work, we propose the simultaneous exploration of LC-ESI(±)-HRMS, ¹³C & ¹H NMR and enzymatic reactivity datasets, through a multiblock modelling strategy (Beniddir *et al.*, 2022), for the determination of candidate inhibitory species in biomass hydrolysates.

This study focuses on the comparison of data from multi-informative analytical techniques to pinpoint descriptors of EH inhibition in biomass samples. The first part of the presentation will focus on sample preparation and applied analytical data preprocessing approach. The core “Data fusion and Multiblock modelling” strategy will then be thoroughly covered. Ultimately, the potential of this methodology to assist biomarker discovery in complex NP matrices will be highlighted.

Keywords: Biomass conversion, inhibitors, mass spectrometry, nuclear magnetic resonance, Multiblock modelling

References

- Beniddir, M.A., Le Moyec, L., Triba, M.N., Longeon, A., Deville, A., Blond, A., van Pham, C., Voogd, N.J. de, Bourguet- Kondracki, M.-L., 2022. Metabolomics with multi-block modelling of mass spectrometry and nuclear magnetic resonance in order to discriminate Haplosclerida marine sponges. *Anal Bioanal Chem* 414, 5929–5942.
- Bhinderwala, F., Wase, N., DiRusso, C., Powers, R., 2018. Combining Mass Spectrometry and NMR Improves Metabolite Detection and Annotation. *Journal of Proteome Research* 17, 4017–4022.
- Deng, L., Gu, H., Zhu, J., Nagana Gowda, G.A., Djukovic, D., Chiorean, E.G., Raftery, D., 2016. Combining NMR and LC/MS Using Backward Variable Elimination: Metabolomics Analysis of Colorectal Cancer, Polyps, and Healthy Controls. *Analytical Chemistry* 88, 7975–7983.
- Kim, D., 2018. Physico-Chemical Conversion of Lignocellulose: Inhibitor Effects and Detoxification Strategies: A Mini Review. *Molecules* (Basel, Switzerland) 23, 309.
- Wyman, C.E., 1999. Biomass Ethanol : Technical Progress, Opportunities, and Commercial Challenges. *Annu. Rev. Energy. Environ.* 24, 189–226.

* Presenting author. E-mail address: ian-emmanuel.ramtan@ifpen.fr (Ian Ramtanon)

S3.P132 Obtaining new [6]-gingerol derivatives with potential antiparasitary activity

Daniel Stoffle Trancoso^a, William Cezar de Lima Silva^a, Thalita Bachelli Riul^b, Warley de Souza Borges^{a,*}

^a Department of Chemistry, Universidade Federal do Espírito Santo, 29075-910, Vitória, Brazil

^b Department of Pharmaceutical Sciences, Food and Nutrition, Universidade Federal do Mato Grosso do Sul, 79070-900, Campo Grande, Brazil

Some parasitic infections are recognized as affecting predominantly the most vulnerable population's parcel due to the state of need and difficulty of access to health services (da Silva et al., 2013). Leishmaniosis is a prevalent pathology in Brazil, and the current treatment options are faced with many obstacles, such as long therapy duration and high toxicity (Trancoso et al., 2022). To mitigate these problems, it becomes necessary to develop new pharmaceutical inputs that are accessible and effective, such as natural product derivatives. [6]-Gingerol is a strong contender to meet such a need, due to the vast range of positive biological effects already visualized, including against parasitic diseases. A form of achieving a better biological activity is through chemical reactions aiming to result in modified products. In this way of thinking, its chemical structure lets different compounds be made by replacing two molecules of nucleophiles at its phenolic hydroxyl. Given that, the goal of this study was to submit [6]-gingerol to modifications using structurally diverse reagents, resulting in seven different and new products. Once synthesized and characterized, these new compounds were tested against the parasitic species *Leishmania amazonensis* to verify the possibility of the enhancement of the [6]-gingerol's activity, something that was achieved even in a moderate form. This work seeks to find new, promising inputs to be used in leishmaniosis treatment.

Keywords: *Leishmania amazonensis*, biological activity, semisynthesis

References

- da Silva, M.A., Fokoue, H.H., Fialho, S.N., Dos Santos, A.P.A., Rossi, N.R.D.L.P., Gouveia, A.J., Ferreira, A.S., Passarini, G.M., Garay, A.F.G., Alfonso, J.J., Soares, A.M., Zanchi, F.B., Kato, M.J., Teles, C.B.G., Kuehn, C.C., 2021. Antileishmanial activity evaluation of a natural amide and its synthetic analogs against *Leishmania (V.) braziliensis*: an integrated approach *in vitro* and *in silico*. *Parasitol Res.* 120, 2199-2218.
- Trancoso, M. E., Germanó, M. J., Arrieta, V.J., Bustos, M.F.G., Cifuentes, D., Cargnelutti, D.E., Lozano, E. S., 2023. Antiparasitic activity of two natural terpenes from *Salvia cuspidate* against *Leishmania amazonenses*. *J. Nat. Prod.* 86, 797-803.

* Presenting author. E-mail address: warley.borges@ufes.br (Warley de Souza Borges)

S3.P133 Identification of the C2-sulfonamidyl chromone scaffold for developing anti-inflammatory agents targeting hyperactivation of human neutrophils

Yi-Han Chang^{a,b}, Po-Jen Chen^{c,d}, Hsin-Yi Hung^{a,*}

^a School of Pharmacy, College of Medicine, National Cheng Kung University, Tainan, 710, Taiwan

^b Biomedical Translation Research Center (BioTReC), Academia Sinica, Taipei, 11571, Taiwan

^c Department of Medical Research, E-Da Hospital, I-Shou University, Kaohsiung 824410, Taiwan

^d Graduate Institute of Medicine, I-Shou University, Kaohsiung 824410, Taiwan

Neutrophil hyperactivation is implicated in a spectrum of inflammatory pathologies, which implies the potential for targeting neutrophils as a pharmacological management to treat inflammatory disorders (van der Linden and Meyaard, 2016). In continue of the previous study that C2-ether bridge of capillarisin was replaced by a more active C2-thioether linkage (Chang et al., 2021), herein, a structure-activity relationship study focusing on the different C-2 bridging moiety was leveraged with the bioisostere-replacing and scaffold- hopping approaches. Among various chemotypes, the C2-sulfonamidyl chromone derivatives emerged as the potential inhibitors against superoxide anion generation and elastase release from fMLF-activated human neutrophils. The most active compound exhibited IC₅₀ value of single digit micromolar range against both neutrophilic superoxide anion production and elastase release. Pharmacophore model predicted the most active compound to be a p38a inhibitor, which is validated by direct p38a inhibition assay, p38a cascade assay as well as western blot analyses. This study provides a successful example of natural product-derived new p38a inhibitor ameliorating neutrophilic inflammation.

Keywords: capillarisin, C2-sulfonamidyl chromone, p38a, neutrophilic inflammation

Reference

van der Linden, M., Meyaard, L., 2016. Fine-tuning neutrophil activation: Strategies and consequences, *Immunol. Lett.* 178 (2016) 3-9.

Chang, Y.H., Fang, S.Y., Lai, H.Y., Hwang, T.L., Hung, H.Y., 2021. The study on structure-activity relationship between chromone derivatives and inhibition of superoxide anion generating from human neutrophils. *Bioorg. Med. Chem. Lett.* 36, 127822.

* Presenting author. E-mail address: z10308005@email.ncku.edu.tw (Hsin-Yi Hung)

S3.P134 New xanthone analogues with potent intracellular reactive oxygen species inhibition effects

Siau Hui Mah^{a,*}, Natalie Vivien Gunter^a, Soek Sin Teh^b, Ibrahim Jantan^c, Kung Pui Law^a

^a School of Biosciences, Taylor's University, 47500 Subang Jaya, Malaysia

^b Engineering and Processing Division, Energy and Environment Unit, Malaysian Palm Oil Board, 43000 Kajang, Malaysia

^c Institute of Systems Biology, University Kebangsaan Malaysia, 43600 Bangi, Malaysia

An elevated level of reactive oxygen species (ROS) often results in oxidative stress, subsequently leading to the initiation and progression of inflammation. As a consequence, compounds that could suppress the ROS levels by upregulating intrinsic antioxidant Nrf2/ARE signaling pathway are needed. The polyketides xanthenes are potential candidate due to their reported antioxidant, anti-inflammatory and anticancer activities (Minami et al. 1994; Gunter et al. 2023). Their biological activities rely on the types and position of substituent group present on the molecular scaffold of xanthone (Loh et al. 2021). Thus, this study aimed to synthesize new xanthone analogues starting from 3-hydroxyxanthone to provide further insight on their structure-activity relationship on intracellular ROS inhibition level. The analogues obtained were purified by chromatography techniques and structurally characterized via NMR, MS and FTIR spectroscopy. Moreover, the xanthone analogues were evaluated for their inhibitory effects towards intracellular levels of ROAs as a consequence through the DCFH-DA assay in H₂O₂-induced oxidative damage model using SNU-1 cells. Furthermore, the modulation of the gene and protein expression of Nrf2 and downstream proteins were assessed. Remarkably, the xanthone analogues inhibited the ROS by restoring the levels to that of negative control. Their activities were found to be comparable to an established Nrf2 activator, sulforaphane. Molecular investigations showed concentration-dependent increases in the mRNA expression of Nrf2 and downstream antioxidant enzymes HO-1, NQO1, SOD, and CAT. Similar results were obtained for the protein expression of these enzymes even though no marked increase was observed for Nrf2. In summary, xanthone analogues are potential lead compounds for anti-inflammatory agents and thus further *in vivo* studies are highly recommended.

Keywords: antioxidant, Nrf2/ARE signaling pathway, structure-activity relationship

References

- Minami, H. et al., 1994. Antioxidant xanthenes from *Garcinia subelliptica*. *Phytochem.* 36, 501-506.
- Gunter, N.V. et al., 2023. The mechanisms of action of prenylated xanthenes against breast, colon, and lung cancers, and their potential application against drug resistance. *Phytochem. Rev.* 22, 467-503.
- Loh, Z.H. et al., 2021. New 3-O-substituted xanthone derivatives as promising acetylcholinesterase inhibitors. *J. Enz. Inhib. Med. Chem.* 36, 627-639.


* Presenting author. E-mail address: SiauHui.Mah@taylors.edu.my (Siau Hui Mah)

S3.P135 A new complementary process toward functionalized dihydropyridinones: application in total synthesis of phenanthroindolizidine alkaloids

Sanghee Kim*

College of Pharmacy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

Due to their interesting biological properties, functionalized dihydropyridinones have attracted considerable interest from drug research fields. Although several synthetic strategies have been developed for preparing these structural motifs, only a limited number of these procedures can be applied for the direct and selective synthesis of polysubstituted dihydropyridinones. Thus, the development of new synthetic strategies that enable an efficient and selective synthesis of highly substituted dihydropyridinones are highly desirable. Herein, we report our research on a new process toward functionalized dihydropyridinones through the intramolecular cyclization of amides with a tethered alkyne moiety and its application. An unexplored reaction between amide and alkyne was realized through an O-silyl N,O-ketene acetal to give functionalized dihydropyridinones in high yields. Applications of this reaction for the synthesis of various functional dihydropyridinones and investigation of its extension to the total synthesis of phenanthroindolizidine and phenanthroquinolizidine alkaloids were achieved. Our strategy could be extended to realize the related heterocyclic substrate series.



Keywords: dihydropyridinones, total synthesis, phenanthroindolizidine, phenanthroquinolizidine, alkaloids

References

Lee, S., Shin, J.E., Yoon, R., Yoo, H. and Kim, S., 2023. Annulation of O-silyl N,O-ketene acetals with alkynes for the synthesis of dihydropyridinones and its application in concise total synthesis of phenanthroindolizidine alkaloids. *Front. Chem.* 11, 1267422.

* Presenting author. E-mail address: pennkim@snu.ac.kr (Sanghee Kim)

S3.P136 Divergent synthesis of prenylated xanthenes as novel fungal IRE1 inhibitors

Thomas Charpentier^a, Elise Oger^a, Anne-Marie Le Ray Richomme^a, Nelly Bataillé-Simoneau^b, Philippe Simoneau^b, Thomas Guillemette^b, Pascal Richomme^a, Guillaume Viault^{a,*}

^a Univ Angers, SONAS, SFR QUASAV, 49000 Angers, France

^b Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, 49070 Beaucouzé, France

The development of fungal infections is a major threat to people's health and agriculture. Over the past decades, with profound environmental changes, fungal pathogens infecting plants and humans have evolved resistance to several antifungal drugs (Fisher et al., 2022). The unfolded protein response (UPR) is essential in controlling the folding and maturation of endoplasmic reticulum (ER) proteins in eukaryotes. UPR has been shown to be involved in plant pathogenic fungal virulence (Joubert et al., 2011), whereas IRE1 protein appears as its only effector in fungi. Therefore, we considered the IRE1 protein as a promising target for the development of new antifungal drugs. The evaluation of a small but high-diversity library of 76 natural products (terpenoids, alkaloids, and polyphenols) through a cell-based screening assay allowed us to identify seven potential inhibitors of IRE1, including polyhydroxylated and prenylated xanthenes. Antifungal activity of the latter was finally assessed in planta on cabbage leaves infected by *Alternaria brassicicola* (Charpentier et al., 2023).




Fig. 1. Structure of synthesized xanthenes.

This study found that the inhibitory potential of xanthone derivatives against IRE1 depends on the number and location of prenylated side chains on the xanthone backbone. To broaden the structure- activity relationship (SAR), different xanthone derivatives with varying degrees of prenylation were synthesized using a divergent strategy (Fig. 1). Inhibition of IRE1 was then assessed in a cell-based assay, and a molecular docking study was performed.

Keywords: antifungal, xanthone, total synthesis, IRE1 inhibitors

References

- Charpentier, T., Viault, G., Le Ray Richomme, A.-M., Bataillé-Simoneau, N., Helesbeux, J.-J., Blon, N., Bastide, F., Marchi, M., Aligon, S., Bruguère, A., Dinh, C. P., Benbelkacem, Z., Dallery, J.-F., Simoneau, P., Richomme, P., Guillemette, T., 2023. Natural products targeting the fungal unfolded protein response as an alternative crop protection strategy. *J. Agric. Food Chem.* 71, 13706-13716.
- Fisher, M. C., Alastruey-Izquierdo, A., Berman, J., Bicanic, T., Bignell, E. M., Bowyer, P., Bromley, M., Brüggemann, R., Garber, G., Cornely, O. A., Gurr, S. J., Harrison, T. S., Kuijper, E., Rhodes, J., Sheppard, D. C., Warris, A., White, P. L., Xu, J., Zwaan, B., Verweij, P. E., 2022. Tackling the emerging threat of antifungal resistance to human health. *Nature Reviews. Microbiol.* 20, 557-571.
- Joubert, A., Simoneau, P., Campion, C., Bataillé-Simoneau, N., Iacomini-Vasilescu, B., Poupard, P., François, J.M., Georgeault, S., Sellier, E., Guillemette, T., 2011. Impact of the unfolded protein response on the pathogenicity of the necrotrophic fungus *Alternaria brassicicola*. *Mol. Microbiol.* 79, 1305-1324.

* Presenting author. E-mail address: guillaume.viault@univ-angers.fr (Guillaume Viault)

S3.P137 Cytotoxic effects of xanthone derivatives towards leukemia cells

Xiaocong Fu^{a,b,*}, Siau Hui Mah^b

^a Shanghai Novopathway Biotechnology Co. Ltd, Shanghai 201203, China

^b School of Bioscience, Taylor's University, Lakeside Campus, Subang Jaya, Selangor, Malaysia

Xanthones derivatives are a group of natural products with interesting pharmaceutical properties including antioxidant, anti-inflammatory, anti-bacterial and anti-cancer, activities (Gunter et al. 2023; Wong et al. 2020). Many previous studies have been focused on the anticancer effects but limited on their anti-leukemia effects with detailed mechanisms of action. Thus, this study aimed to evaluate the cytotoxic effects of synthesized xanthone derivatives towards leukemia by using A20, HL20 and Jurkat cells through the MTT assay. A total nineteen xanthone derivatives were evaluated at a concentration of 10 μ M and four of the derivatives were found to be show strong cytotoxic effects. These xanthone derivatives exhibited at least 90% of cytotoxicity towards all the cell lines. Thus, a range of concentrations of these derivatives were evaluated to obtain their IC₅₀ values. The results have shown that their IC₅₀ values are less than 3 μ M, indicating their potent cytotoxicity. The structure-activity relationship revealed that the halides are nitrogen containing substituent groups are the main contributors to the cytotoxic effects of xanthones. In summary, xanthone derivatives are potential lead compounds for leukemia cancer drugs. Thus, studies on the molecular mechanism pathways for the cytotoxic xanthone derivatives are under progress.

Keywords: xanthones derivatives, anticancer, cytotoxicity, MTT assay, structure-activity relationship

References

- Gunter, N. V., Teh, S.S., Jantan, I., Cespedes-Acuña, C.L., Mah, S H. 2023. The mechanisms of action of prenylated xanthones against breast, colon, and lung cancers, and their potential application against drug resistance. *Phytochem. Rev.* 22, 467-503.
- Wong, K.W., Teh, S.S., Law, K.P., Ismail, I.S., Sato, K., Mase, N., Mah, S.H. 2023. Synthesis of benzylated amine- substituted xanthone derivatives and their antioxidant and anti-inflammatory activities. *Arch. Pharm.* 356, e2200418.

* Presenting author. E-mail address: fuxiaocong2022@126.com (Xiaocong Fu)

S3.P138 Bimatoprost potentiates hair growth via Akt/Wnt/ β -catenin signaling by targeting the K_{ATP} channels

Jung-Il Kang^{a,b}, Youn Kyung Choi^{a,b}, Da Hee Yang^{a,b}, Eun-Sook Yoo^{a,b}, Hee-Kyoung Kang^{a,b,*}

^a Department of Medicine, College of Medicine, Jeju National University, Jeju 63243, South Korea

^b Jeju Research Center for Natural Medicine, Jeju National University, Jeju 63243, South Korea

Various prostaglandins (PGs) and PG synthases are involved in hair growth or hair loss. Male pattern hair loss is caused by increased PGD2 (Garza, 2012), but the mechanism by which PGE2 or PGF2 α promotes hair growth is unclear. Here, we investigated the effect and mechanisms of bimatoprost, an ethyl amide derivative of 17-phenyl trinor PGF2 α , on the hair growth. When C57BL/6 mice were applied topically with bimatoprost, not only rapidly progressive skin color changes but also large- sized hair structures were observed. To elucidate the mechanism by which bimatoprost promotes hair growth, we evaluated whether bimatoprost activate the Akt and Wnt/ β -catenin pathways. Bimatoprost induced cell proliferation through activation of the Akt and wnt/ β -catenin pathways in rat dermal papilla cells (rDPC), which are hair follicle cells, as well as in NIH3T3 cells, and similar results were observed in the minoxidil-treated group. Moreover, all effects of bimatoprost were completely inhibited in rDPC and NIH3T3 cells using glibenclamide, a selective K_{ATP} channel blocker. Bimatoprost significantly increased the level of cPGES, which is a PGE2 synthesis enzyme, in both 2D rDPC and 3D spheroids. In conclusion, these results indicate that bimatoprost induces the proliferation of rDPC by the activation of the K_{ATP} channel/Akt/Wnt/ β -catenin signaling cascade and increases cPGES level, which ultimately exerts a superior hair growth effect.

Acknowledgments: This research was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(RS-2023-00270936) and by the Korea government(MSIT) (2023R1A2C1007234)].

Keywords: bimatoprost, hair growth, wnt/ β -catenin, cPGES, K_{ATP} channel

References

Garza, L. A., Liu, Y., Yang, Z., Alagesan, B., Lawson, J. A., Norberg, S. M., Loy, D. E., Zhao, T., Blatt, H. B., Stanton,

D. C., Carrasco, L., Ahluwalia, G., Fischer, S. M., FitzGerald, G. A., Cotsarelis, G., 2012. Prostaglandin D2 inhibits hair growth and is elevated in bald scalp of men with androgenetic alopecia. *Sci. Transl. Med.* 4, 126ra34.

* Presenting author. E-mail address: pharmkhk@jejunu.ac.kr (Hee-Kyoung Kang)

S3.P139 Large scale synthesis, anticonvulsant effect and toxicity assessment of cannabigerol and cannabinalol

Evangelos Dadiotis^a, Lida Evmorfia Vagiaki^{b,c}, Vangelis Mitsis^d, Eleni Melliou^a, Kyriaki Sidiropoulou^{b,c}, Prokopios Magiatis^{a,*}

^a Laboratory of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Greece

^b Department of Biology, University of Crete, Greece

^c Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology, Greece

^d Ekati Alchemy Lab SL, Barcelona, Spain

Cannabigerol (CBG) and cannabinalol (CBN) are prominent cannabinoids with diverse molecular targets. This study presents a novel, environmentally friendly approach to the total and large-scale synthesis of CBG and an improved method for the scaled-up synthesis of CBN. For CBG, we utilized Amberlyst 15 resin in the condensation of olivetol with geraniol. This method offered a more manageable and less toxic alternative to the BF₃·OEt₂ approach cited in literature (Nguyen et al., 2022) with 55% yield. Regarding CBN, we optimized the iodine-promoted process (Pollastro et al., 2018) by replacing toluene with heptane and by applying an air stream inside a chemical reactor, removing effectively hydrogen iodide, achieving 97% conversion of cannabidiol (CBD) to CBN in a Kg scale reaction. Our pharmacological investigation revealed that both CBG and CBN exhibit potent anti-seizure effects. Following induction of seizure activity in hippocampal mouse brain slices using the *ex vivo* high potassium model, we recorded decreased spontaneous activity (>30%) in a dose- dependent manner (1-40 uM). Their anticonvulsant effect was similar to the one of CBD, an FDA approved anti-epileptic drug branded as Epidiolex. Furthermore, their safety profiles were assessed through acute oral toxicity experiments in mice, with doses up to 2000 mg/kg being nontoxic. Our findings highlight the anticonvulsant properties of CBG and CBN and suggest further *in vivo* investigation to assess their potential as candidates for new or supplementary oral anti-epileptic treatments. Their scalable synthesis, anticonvulsant potential, and favorable toxicity profiles highlight their significance in the ongoing research for effective epilepsy therapies.

Keywords: cannabinoids, anticonvulsants, cannabinalol, cannabigerol, cannabidiol

References

- Nguyen, G.-N., Jordan, E.N., Kayser, O., 2022. Synthetic strategies for rare cannabinoids derived from *Cannabis sativa*. J. Nat. Prod. 85, 1555–1568.
- Pollastro, F., Caprioglio, D., Marotta, P., Moriello, A.S., De Petrocellis, L., Tagliatella-Scafati, O., Appendino, G., 2018. Iodine-promoted aromatization of *p*-menthane-type phytocannabinoids. J. Nat. Prod. 81, 630–633.

* Presenting author. E-mail address: magiatis@pharm.uoa.gr (Prokopios Magiatis)

S3.P140 Thymol-based heterocyclic derivatives: synthesis and SARS-CoV-2 M_{pro} molecular docking studies

Geneive E. Henry*, Eden L. Parks

Department of Chemistry, Susquehanna University, Selinsgrove, PA 17870, USA

SARS-CoV-2 main protease (SARS-CoV-2 M^{pro}) enzyme is involved in the replication of the SARS-CoV-2 virus, and is therefore an important target in the development of drug treatments for COVID-19 disease. Plant based natural products of the monoterpene class have been explored as inhibitors of the enzyme, including thymol, a major component of thyme essential oil. In this study, twelve heterocyclic derivatives of thymol containing various heterocyclic moieties were synthesized. Molecular docking studies were performed with SARS-CoV-2 M^{pro} (PDB ID; 6lu7) to determine the influence of the heterocyclic unit on docking affinity. All of the derivatives showed interaction with His-41 and Cys-145, which are important in the catalytic mechanism of the SARS-CoV-2 M^{pro} enzyme. Docking scores ranged from -6.4 kcal to -8.0 kcal/mol, with evidence of structure-docking affinity correlations among the heterocyclic units. These scores are in a similar range to that of ML-188 (-7.5 kcal/mol), a known SARS-CoV-2 M^{pro} inhibitor.

Keywords: thymol, heterocycles, SARS-CoV-2 M^{pro} docking affinity

* Presenting author. E-mail address: henry@susqu.edu (Geneive E. Henry)

S3.P141 Expanding chemical space around structures from *Strychnos variabilis*: late-stage functionalization strategies using photochemistry

Ledoux Allison^a, Schmitz Astrid^a, Lesenfants Cindy^a, Leka Kristi^a, Frédéricich Michel^a, Garigliany Mutien-Marie^b, Francotte Pierre^{a,*}

^a University of Liège Faculty of Medicine, CIRM, Building B36 Quartier Hôpital 4000, Liège, Belgium

^b University of Liège Faculty of veterinary Medicine, FARA, Building B42 Quartier Vallée 2 Avenue de Cureghem 6 4000, Liège, Belgium

Strychnos variabilis De Wild. is a small deciduous tree native to the regions in and around Brazzaville and Kinshasa, situated on both banks of the Congo River in Africa. Despite its geographical prevalence, *S. variabilis* leaves were found to contain only trace amounts of indoline alkaloids, particularly from the retuline series. However, our laboratory, over the past three decades, isolated a plethora of rare flavonoids, including flavonol glycosides, from these leaves (Brasseur and Angenot, 1987). Flavonoids, renowned for their antiviral properties, particularly in glycosidic form, have garnered interest due to their enhanced solubility and efficacy. Therefore, our study sought to explore the potential antiviral activity of *S. variabilis* leaves against SARS-CoV-2. Initial findings indicated significant antiviral activity in the total extract, with an IC₅₀ range of 12.5 to 6.25 µg/mL. Encouragingly, our investigation into two specific flavonoids, variabilosides C and D, revealed even greater potency, with an IC₅₀ range of 6.25 to 3.12 µg/mL. Building upon these promising results, we adopted late-stage functionalization (LSF) strategy through photochemistry (Pitre and Overman, 2022) to chemodiversify *S. variabilis* extracts. LSF may alter functions already present in the structure, but more importantly it is able to target the C–H bonds of drug leads as points of diversification for generating new analogs. Thus, chemodiversification was achieved and preliminary insights were obtained. The antiviral activity of these modified extracts against SARS-CoV-2, were correlated with these modifications. In conclusion, our study highlights the efficacy of photochemistry in expanding the chemical space of natural products, offering innovative solutions for drug discovery and development.




Fig. 1. Variabilosides C and D isolated from *S. variabilis*.

Keywords: *Strychnos variabilis*, SARS-CoV-2, chemodiversification, late-stage functionalisation

References

- Brasseur, T., Angenot, L., 1987. Four acylated flavonol glycosides from leaves of *Strychnos variabilis*. *Phytochemistry* 26, 3331-3334.
- Pitre, S.P., Overman, L.E., 2022. Strategic use of visible-light photoredox catalysis in natural product synthesis. *Chem. Rev.* 122, 1717-1751.

* Presenting author. E-mail address: pierre.francotte@uliege.be (Pierre Francotte)

S3.P142 Effects of metabolic modulators on the accumulation of specialized metabolites in *Hypogymnia physodes* (L.) Nyl.

Izabela Baczevska^{a,*}, Sławomir Dresler^{a,b}, Agnieszka Hanaka^b

^a Department of Analytical Chemistry, Medical University of Lublin, Chodźki 4a, 20-093 Lublin, Poland

^b Department of Plant Physiology and Biophysics, Institute of Biological Science, Maria Curie- Skłodowska University, Akademicka 19, 20-033 Lublin, Poland

Lichens are symbiotic organisms consisting of a fungal component (mycobiont) and a plant component (photobiont). These organisms are a source of many biologically active substances, among which the most significant group are the so-called lichen acids. This group of compounds includes about 1,000 identified metabolites specific only to lichenized fungi. Secondary metabolites of lichens may account for up to 25% of their dry weight. They are characterized by antioxidant, antimicrobial, anti-inflammatory, photoprotective, and anticancer properties. Six common lichen species were used in this study: *Hypogymnia physodes* (L.) Nyl, *Xanthoria parietina* (L.) Th. Fr., *Evernia prunastri* (L.) Ach., *Cladonia uncialis* (L.) Weber ex F.H. Wigg., *Physcia adscendens* (Fr.) H. Olivier., and *Pseudevernia furfuracea* (L.) Opf. The lichen thallus was subjected to different metabolic modulators (methyl jasmonate, salicylic acid, nitrate oxide, selenite, zinc, chitosan lactate). After 48-hour incubation in stress conditions, the accumulation of lichen acids and allantoin was examined. The antioxidant potential exhibited by the prepared extracts was also studied. Although the results obtained did not provide sufficient evidence for a common pattern of action of metabolic modulators in all the lichen species studied, alterations in the metabolite content related to the metabolic modulator-induced lichen response were observed. The data showed significantly increased accumulation of specific compounds in some lichen species, especially after the exposure to salicylic acid, methyl jasmonate, NO, and chitosan.

Keywords: allantoin, antioxidant potential, elicitation, lichens, lichen acids, secondary metabolites

References

- Pichler, G., Muggia, L., Carniel, F. C., Grube, M., Kranner, I., 2023. How to build a lichen: from metabolite release to symbiotic interplay. *New Phytol.* 238, 1362-1378.
- Stasińska-Jakubas, M., Hawrylak-Nowak, B., Dresler, S., Wójciak, M., Rubinowska, K., 2023. Application of chitosan lactate, selenite, and salicylic acid as an approach to induce biological responses and enhance secondary metabolism in *Melissa officinalis* L. *Ind. Crops Prod.* 205, 117571.
- Thakur, M., Kapoor, B., Kapoor, D., & Sharma, N. R., 2023. Lichens: A promising source of anti-cancerous activity and their molecular mechanisms. *S. Afr. J. Bot.* 159, 155-163.

* Presenting author. E-mail address: izabela.baczevska98@gmail.com (Izabela Baczevska)

S3.P143 Synthesis of polymethoxyflavone - zapotin (5,6,2',6'-tetramethoxyflavone) and its new structural analogues

Monika Tomczykowa^{a,*}, Damian Pawelski^a, Jakub Władysław Strawa^b

^a Department of Organic Chemistry, Faculty of Pharmacy with the Division of Laboratory Medicine, Medical University of Białystok, ul. Mickiewicza 2A, 15-222 Białystok, Poland

^b Department of Pharmacognosy, Faculty of Pharmacy with the Division of Laboratory Medicine, Medical University of Białystok, ul. Mickiewicza 2A, 15-230 Białystok, Poland

The synthesis of 5,6-substituted flavonoid derivatives is intricate. Synthesis methods used so far are multi-step, long-lasting, time-consuming with a low total product yield, and therefore uneconomical (Maiti et al., 2007). Polymethoxyflavones (PMF) are of particular interest due to their broad spectrum of biological activities, including anti-inflammatory and anticancer properties. According to the literature data, zapotin, is an extremely interesting example of PMF (Strawa et al., 2021) with broad biological potential, especially as an anti-tumor and chemo-preventive factor (Changhong et al., 2022). Difficulties related to the synthesis of 5,6-substituted PMF using known synthetic methods are the reason why they have been obtained mainly through isolation from plant material so far. The amount of zapotin isolated from plant raw materials is scarce, and the process of obtaining it is very troublesome and inefficient. The synthesis of zapotin and its analogues was carried out to optimize synthesis conditions. Optimization of the conditions aimed at reducing the number of synthesis steps which led to the shortening of total synthesis time. Changing the synthesis strategy from linear to more branched allowed to easily obtain zapotin **1** and its analogues **2-3** using more selective and easiest to perform transformations. The synthesis we propose does not require using completely anhydrous conditions as well as halogenated solvents, at any stage of the synthesis. Intermediate products can be easily purified by crystallization leading to the final products - compounds **1-3** - in a crystalline, stable form, with relatively good yield (46%-48%) (Fig. 1).




Fig. 1. Structure of zapotin **1** and its synthesized analogues **2-3**

Keywords: zapotin, polymethoxyflavones, synthesis

References

- Changhong, L., Taotao, D., Jun, Ch., Mingshun, Ch., Ruihong, L., Chengmei, L., Liqing, D., McClements, D.J., 2023. Modification of flavonoids: methods and influences on biological activities. *Crit. Rev. Food Sci. Nutr.* 63, 10637-10658.
- Maiti, A., Cuendet, M., Kondratyuk, T., Croy, V. L., Pezzuto, J. M., Cushman, M., 2007. Synthesis and cancer chemopreventive activity of zapotin, a natural product from *Casimiroa edulis*. *J. Med. Chem.* 50, 350-355.
- Strawa, J.W., Jakimiuk, K., Tomczyk, M., 2021. Zapotin, a polymethoxyflavone, with potential therapeutic attributes. *Int. J. Mol. Sci.* 22, 13227.

* Presenting author. E-mail address: monika.tomczyk@umb.edu.pl (Monika Tomczykowa)

S3.P144 Anti-leishmanial activity of synthesized and characterized semisynthetic derivatives of betulin

Aanchal Loshali^{a,*}, Vidhu Aeri^a, Sandhya Bawa^b

^a Department of Pharmacognosy and Phytochemistry, School of Pharmaceutical Education and Research, Jamia Hamdard, 110062 New Delhi, India

^b Department of Pharmaceutical Chemistry, School of Pharmaceutical Education and Research, Jamia Hamdard, 110062 New Delhi, India

Betulin is a naturally occurring pentacyclic triterpene that is isolated from the bark of *Betula utilis* D. Don, a member of the Betulaceae family (Verma et al., 2014). It belongs to the lupane series and is known by the systematic name 3, 28-dihydroxy-20(29) lupen or Lup-20(29)-en-3, 28-diol. Betulin has three available sites for simple chemical modification, namely the secondary hydroxyl group at position C-3, the primary hydroxyl group at position C-28 and the isopropenyl side chain at position C-19. A broad range of biological activities, including antibacterial, antiviral, anticancer, and hepatoprotective qualities, are exhibited by betulin and its derivatives (Amiri et al., 2020). The aim of the present study was to anti-leishmanial activity of synthesized and characterized semisynthetic derivatives of betulin. The synthesized derivatives were characterized using various spectroscopic techniques, including NMR, IR, and MS. Anti-leishmanial activity was done by *in vitro* promastigote assay, *ex vivo* cytotoxicity assay and amastigote-macrophages assay (Sharma et al., 2023). The synthesized betulin derivatives was attached on C3 hydroxyl group and it was characterized by spectroscopic techniques. The synthesized derivatives was tested against *L. donovani* and confirmed its antileishmanial activity against the standard drug amphotericin B by *in vitro* studies.

Keywords: *Betula utilis*, betulin, synthesis, characterization, antileishmanial

References

- Verma, D., Ajoankar, S., Sahu, S., Rane, M., Teli, N., 2014. Pharmacological and phytochemical properties of *Betula utilis*: An overview. Res. J. Pharm. Biol. Chem. Sci. 5, 284–288.
- Amiri, S., Dastghaib, S., Ahmadi, M., Mehrbod, P., Khadem, F., Behrouj, H., Aghanoori, M.R., Machaj, F., Ghamsari, M., Rosik, J., Hudecki, A., 2020. Betulin and its derivatives as novel compounds with different pharmacological effects. Biotechnol. Adv. 38, 107409.
- Sharma, S., Anjaneyulu Yakkala, P., Beg, M.A., Tanwar, S., Latief, I., Khan, A., Sharma, K., ur Rehman, S., Selvapandiyani, A., Shafi, S., 2023. 5-Arylidene-2, 4-thiazolidinediones as cysteine protease inhibitors against *Leishmania donovani*. Chemistry Select. 8, 202302415.

* Presenting author. E-mail address: aanchaloshali_sch@jamaiahamdard.ac.in (Aanchal Loshali)

S3.P145 Exploring novel amino acid synthesis for plant-derived active cyclopeptide acquisition

Zheng-Ming Liang*, Hong-Jie Zhang

School of Chinese Medicine, Hong Kong Baptist University, 7 Baptist University Road, Kowloon Tong, Hong Kong SAR, People's Republic of China

MVA, a novel plant cyclic heptapeptide containing dimethylcyclopropyl-glycine (DMCPA) unique amino acids, exhibits remarkable anticancer activity with IC_{50} values in the nanomolar range against various cancer cell lines (Zhang, 2014). The need for the total synthesis of MVA is underscored by its rare occurrence in nature for further bioactivity studies and elucidation of its mechanism of action. In our efforts to create analogs of MVA, we substituted some common amino acids like cyclopropyl-glycine for DMCPA. However, the anticancer activities of the synthesized analogs significantly decreased, highlighting the crucial role of the unique amino acids in maintaining the anticancer potency of MVA. As DMCPA is not commercially available, the critical step in acquiring MVA is the synthesis of DMCPA to advance towards our ultimate goal of preparing MVA. The current research outlines our total synthetic approach for DMCPA.




Fig. 1. Total synthesis of DMCPA

Given that DMCPA and glutamic acid share the same backbone length, the chiral centre of the resulting amino acid does not necessitate construction due to the predetermined configuration of the starting material (Fig. 1). The synthesis of DMCPA involves crucial stages like forming cyclopropane ring and converting the carboxylic acid into a methyl group. Consequently, it was determined to use it as the primary ingredient for DMCPA preparation. Through a series of subsequent reactions, including Corey-Chaykovsky and the Appel reactions, the final DMCPA can be produced.

The study was financially supported by the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No. HKBU12142016 and HKBU12103021).

Keywords: cyclopeptide, plant compound, cytotoxicity, unique amino acid, synthesis

References

Zhang, H.J., 2014. Anticancer and anti-obesity cyclic peptide agents. U.S. Patent 2014/0107018 A1, April 17, 2014.

* Presenting author. E-mail address: zhengming@hkbu.edu.hk (Zheng-Ming Liang)

S3.P146 Structure modification of plant-derived miliusanes compounds to explore their therapeutic potential as anticancer agents

Wen-Jian Xie^{a,*}, Yi-Xuan Xia^a, Stephen Deyrup^{a,b}, Hong-Jie Zhang^a

^a School of Chinese Medicine, Hong Kong Baptist University, 7 Baptist University Road, Kowloon Tong, Hong Kong SAR, People's Republic of China

^b Department of Chemistry and Biochemistry, Siena College, 515 Loudon Road, 12211 NY, USA

Geranyl meroterpenoids, a class of naturally occurring compounds produced through a hybrid biogenetic pathway, have recently captured significant attention due to their multifaceted bioactivities. Miliusanes represent a striking example discovered in plants of the *Miliusa* genus, characterized by an 18-carbon skeleton formed from the fusion of homogentisic acid and a geranyl monoterpene. In our screening program involving a wide range of plant extracts, *Miliusa sinensis* emerged as a promising anticancer plant lead (Zhang et al., 2016). Subsequent bioassay-guided fractionation of two *Miliusa* species led to the discovery of 43 novel miliusanes (Zhang et al., 2006; Xu et al., 2019; Xie et al., 2023). Some of them exhibited potent *in vitro* and *in vivo* antitumor activities, with their mechanisms of action partially attributed to their ability to modulate the p21- dependent cellular senescence pathway. The current study presents our ongoing endeavors aimed at obtaining novel miliusanes through the structural modification of the miliusol prototype. Consequently, we have successfully synthesized over 100 miliusane derivatives (Fig. 1), some of which demonstrated 5-10 times higher potency than miliusol *in vitro*. Further *in vivo* assessments revealed that two compounds featuring Michael adducts exhibited a more balanced activity profile concerning anticancer efficacy and toxicity.




Fig. 1. Structure modification strategy based on the miliusol lead molecule.

These findings have provided us with valuable structure-activity relationship insights, which will facilitate the advancement of miliusanes as potential anticancer drug candidates.

This project was supported by Hong Kong Baptist University, Research Committee, Initiation Grant- Faculty Niche Research Areas (IG-FNRA) 2021/22, Hong Kong (RC-IG-FNRA/21-22/SCM/01).

Keywords: *Miliusa sinensis*, meroterpenoids, miliusanes, antitumor

References

- Zhang, H.J., Li, W.F., Fong, H.H.S., Soejarto, D.D., 2016. Discovery of bioactive compounds by UIC-ICBG drug discovery program in the 18 years since 1998. *Molecules* 21, 1448-1464.
- Zhang, H.J., Ma, C., Hung, N.V., Cuong, N.M., Tan, G.T., Santarsiero, B.D., Mesecar, A.D., Soejarto, D.D., Pezzuto, J.M., Fong, H.H.S., 2006. Miliusanes, a class of cytotoxic agents from *Miliusa sinensis*. *J. Med. Chem.* 49, 693-708.
- Xu, X.Y., Tsang, S.W., Guan, Y.F., Liu, K.L., Pan, W.H., Lam, C.S., Lee, K.M., Xia, Y.X., Xie, W.J., Wong, W.Y., Lee, M.M.L., Tai, W.C.S., Zhang, H.J., 2019. *In vitro* and *in vivo* antitumor effects of plant-derived miliusanes and their induction of cellular senescence. *J. Med. Chem.* 62, 1541-1561.
- Xie, W.J., Xu, X.Y., Xia, Y.X., Lam, C.S., Chen, T., Zhi, Y.S., Lee, K.M., Xun, S., Deyrup, S.T., He, Z.D., Zhang, H.J., 2023. Identification and bioactivity evaluation of miliusanes from *Miliusa sinensis*. *Bioorg. Chem.* 140, 106797-106809.

* Presenting author. E-mail address: xiewenjian@hkbu.edu.hk (Wenjian Xie)

Genetic engineering of food crops and medicinal plants

S3.P147 Seed propagation, tissue culture and *in-vitro* conservation studies in *Aristolochia tagala* Cham: an endangered medicinal plant of western ghats of India

Raviraja Gavali Shetty^{a,*}, Rajasekharan^b

^a Zonal Agricultural and Horticultural Research Station, Brahmavara, India (KSNUAHS, Shivamogga, India

^b ICAR- Indian Institute of Horticultural Research, Hesaraghatta, Bangalore, India

Aristolochia tagala Cham. is a threatened medicinal plant of Aristolochiaceae family. It is mainly used to treat snake bites and other poisonous bites. Apart from this, roots are also used to treat bone fracture, malaria, and various dermatological conditions. Due to indiscriminate use of its roots and deforestation, its population is reducing in its natural habitat. Seed germination is low due to the presence of insufficient endosperm, which leads to low seed viability. Hence the conservation of this species is crucial. Hence, this study was undertaken. Fresh seeds of *A. tagala* were collected from the Field Gene Bank of RET medicinal plants. Among different treatments, the water soaking treatment responded well with the maximum in all growth parameters like rate of germination, seedling height and seedling vigor index followed by treatment with thiourea 2.0 per cent. In tissue culture, nodal segments of seedlings as an explant and cultured on MS medium supplemented with various combinations of growth regulators like BAP, kinetin and NAA. Among different combinations, minimum days taken for shoot initiation (6.06 ± 0.04), the maximum number of shoots (3.20 ± 0.04) and the maximum number of the leaves (3.18 ± 0.02) were obtained in treatment combination of BAP (2.0 mg/L) and NAA (0.5 mg/L). The maximum shoot length (1.98 ± 0.02) was obtained in BAP (2 mg/L). The satisfactory results after three months of *in vitro* conservation were obtained by using the combination of BAP (2.0 mg/l) + NAA (0.50 mg/L) in terms of high survival per cent (100), maximum number of shoots per explant (2.40 ± 0.04) and the maximum number of leaves (2.73 ± 0.02). Thus, MS media supplemented with BAP (2.0 mg/l) + NAA (0.50 mg/L) can be used for both *in vitro* propagation and also conservation of *Aristolochia tagala* Cham. under reduced culture condition and low light intensity ($2.97 \mu\text{m}^{-2} \text{s}^{-1}$) and low temperature (10°C).

Keywords: propagation, conservation, endangered, seeds, treatment, germination

References

- Rajasekharan P.E., Ambika, S., Ganeshan, S., 2009. *In vitro* conservation of *Tylophora indica*: A threatened medicinal plant. J. Genet. Evol. 2, 1-11.
- Senapathi, S.K., Aparajitha, S., Rout, G.R., 2013. Micropropagation and assessment of genetic stability in *Celastrus paniculatus*: an endangered medicinal plant. Biologia (Springer) 68, 627-632.
- Tiwari, R.K.S., Chandra, K.K., Dubey, S., 2018. Techniques for breaking seed dormancy and its efficacy on seed germination of six important medicinal plant species. Int. J. Environ. Agric. Biotechnol. 11, 293-301
- Tiwari, S., Singh, K., Shah, P., 2007. *In vitro* propagation of *Oroxylum indicum* - an endangered medicinal tree. J. Biotechnol. 6, 299-301.

* Presenting author. E-mail address: rrshetty2059@gmail.com (Raviraja Gavali Shetty)

S3.P148 Efficient genetic transformation of *Nigella damascena* callus

Magdalena Klimek-Chodacka*, Dominik Huber, Magdalena Pieczara, Rafał Barański

University of Agriculture in Kraków, al. Mickiewicza 21, 31-120 Kraków, Poland

Nigella damascena L., commonly known as love-in-a-mist, originates from the Mediterranean basin and Southwest Asia. Its seeds are abundant in essential oils, rich in β -elemene and damascenine, which demonstrate antimicrobial and molluscicidal properties. Recent investigations have also explored the pharmacological effects of damascenine, unveiling its potential as an analgesic, antipyretic, anti-inflammatory, and antiedematous agent. As conventional methods of producing these compounds can be inefficient, various biotechnological approaches are increasingly being used to improve their production efficiency. The use of medicinal plants in *in vitro* cultures provides a controlled environment for the production of therapeutic compounds, thereby improving access to natural remedies. Regulation of the production levels of target compounds can be achieved through a variety of approaches, including elicitation, manipulation of growth parameters, and modification at the molecular level. Despite the promising potential of *N. damascena*, research on this species remains limited, with only a few scientific studies available, none of which involve genetic transformation, highlighting the need for further research in this area. In our study, we developed an efficient protocol for the genetic transformation of *N. damascena* callus cells using *Agrobacterium tumefaciens* (*Rhizobium tumefaciens*). We achieved a remarkably high transformation efficiency, which opens the door for further research and applications. The introduction of reporter genes has facilitated the evaluation of transformation efficiency, simplifying the process. As a result, it is now possible to insert different genes or use the clustered regularly interspaced short palindromic repeats (CRISPR) – Cas9 system for targeted mutagenesis, paving the way for genetic manipulation in *N. damascena* and deepening our understanding of its medicinal properties.

Keywords: *Nigella damascena*, genetic transformation, *in vitro*, callus, *Agrobacterium tumefaciens*

* Presenting author. E-mail address: m.chodacka@urk.edu.pl (M. Klimek-Chodacka)

Natural cosmetics, nutraceuticals, and food supplements

S3.P149 Identifying antioxidant compounds within a mixture through dereplication and online DPPH strategies: A study on the CO₂ byproduct of the Makwaen Pepper

Jérémy Perret^{a,b,c,*}, Shahad Ragheed Georges^b, Julien Cordonnier^b, Simon Remy^b, Florent Allais^c, Jérôme Masson^a, Jean-Hugues Renault^b

^a Cosmo International Ingredients, 855 av. Maurice Donat, 06250 Mougins, France

^b University of Reims Champagne-Ardenne, ICMR 7312, 51687 Reims, France

^c URD Agro-Biotechnologies Industrielles (ABI), CEBB, AgroParisTech, 51110 Pomacle, France

Natural antioxidants are widely used in cosmetics, nutraceuticals, and food supplements as preservatives or to protect against reactive oxygen species (Brewer, 2011). Exploring byproducts from various industries - often considered waste - presents a substantial and promising approach for developing natural and sustainable antioxidant ingredients in a circular economy perspective.

This study aimed to develop a robust and sensitive online screening workflow to identify compounds with anti-free radical activity in chemically complex natural extracts. Compared to previously reported results (Irina et al., 2000), the response's sensitivity and resolution were significantly improved by introducing centrifugal partition chromatography as a first separation dimension before the online workflow HPLC-DPPH-UV-HRMS². The dereplication step was then performed by combining NMR and MS (Cordonnier et al., 2023) data to improve the annotation confidence level and finally merged with the free-radical activity information.

As a proof of concept, the developed original and efficient workflow was performed on the CO₂ byproducts of the Makwaen pepper (*Zanthoxylum myriacanthum*), known for its medicinal properties and culinary seasoning. While CO₂ extract of pepper fruits has gained commercial attention, particularly in the fragrance industry for their sulphury fruity olfactive profile (Sriwichai et al. 2019), non-volatile compounds remain insufficiently explored.

The results were visualized as a multi-informative molecular network by matching annotated compounds with a reliability level score and anti-free radical activity activities. For instance, hydroquinone, isoquercetin, hyperoside, dicaffeoylquinic acids, chlorogenic acids and lignan glucosides were annotated and identified to be responsible for the activity of CPC fractions.

Keywords: antioxidant, dereplication, molecular networking, centrifugal partition chromatography, *Zanthoxylum sp*

References

- Brewer M.S., 2011. Natural Antioxidants: Sources, Compounds, Mechanisms of Action, and Potential Applications. *Compr. Rev. Food Sci. Food Saf.*, 10:221–247.
- Irina I. Koleva, Harm A. G. Niederländer, and Teris A. van Beek, 2000. An Online HPLC Method for Detection of Radical Scavenging Compounds in Complex Mixtures. *Analytical Chemistry*. 72 (10), 2323-2328
- Cordonnier J., Remy S., Renault J., Nuzillard. 2023. Versa DB: Assisting ¹³C NMR and MS/MS Joint Data Annotation Through On-Demand Databases. *Chem. Methods*, 4, e202300020.
- Sriwichai, T.; Junmahasathien, T.; Sookwong, P.; Potapohn, N.; Sommano, S.R., 2019. Evaluation of the Optimum Harvesting Maturity of Makhwaen Fruit for the Perfumery Industry. *Agriculture*. 9, 78.

* Presenting author. E-mail address: jeremy.perret@univ-reims.fr (Jérémy Perret)

S3.P150 Establishing human intestinal slice cultures as an ex vivo model of inflammatory bowel diseases for the pharmacological characterization of herbal extracts

Julia Werner^{a,b,*}, Laura Schiller^{a,b}, Kerstin Hill^c, Sonja Kallendrusch^d, Cica Vissiennon^{a,b}

^a Institute of Medical Physics and Biophysics, Medical Faculty, Leipzig University, Härtelstraße 16- 18, 04107 Leipzig, Germany

^b Repha GmbH Biologische Arzneimittel, Alt-Godshorn 87, 30855 Langenhagen, Germany

^c Rudolph-Boehm-Institute for Pharmacology and Toxicology, Medical Faculty, Leipzig University, Härtelstraße 16-18, 04107 Leipzig, Germany

^d Institute of Clinical Research and System Medicine, Medical Faculty, Health and Medical University Potsdam, Schiffbauergasse 14, 14471 Potsdam, Germany

Inflammatory bowel diseases (IBD) are characterized by recurrent or progressive inflammatory alterations of the intestine. Besides standard therapy, complementary treatments, especially phytomedicines, are frequently used. The traditional herbal combination of myrrh, coffee charcoal and chamomile flower extract is included in the German S3-guideline for 'Colitis ulcerosa' based on clinical investigations (Kucharzik et al. 2020). Previous cell culture studies demonstrated anti-inflammatory, spasmolytic and barrier-stabilizing effects of these herbal extracts (Vissiennon et al. 2017; Weber et al. 2020).

This project aims to capture the complex human pathophysiology in an ex vivo model by establishing precision cut intestinal slices (PCIS) from colon biopsies of IBD patients. Influences of the herbal test substances (reference compound: budesonide) on inflammation were monitored. Initial results from hematoxylin-eosin staining confirm the preservation of intestinal morphology in PCIS ex vivo for 24 h (88,4% of samples) and compared to healthy tissue, IBD-typical morphological changes were observed in PCIS of IBD patients. Using immunohistochemistry, distinctions in specific immune cell populations could be detected, revealing comparable percentages of CD68⁺ macrophages (IBD: 1,57%±0,80%; healthy: 0,96%±0,27%) but increased neutrophils (IBD: 3,96%±1,02%; healthy: 0,82%±0,44%). The release of 24 mediators (such as TNFα, IL8, IL6) into the culture media was quantified using multiplex-ELISA. Compared to untreated controls, budesonide (1μM) treatment showed less inflammation, an enhanced epithelial morphology and a significantly altered the mediator pattern.

Combining results from histology and mediator release, this model allows investigation of the effects of herbal extracts from myrrh, coffee charcoal and chamomile flower within the context of IBD's complex pathophysiology.

Keywords: inflammatory bowel diseases, ex vivo, PCIS, herbal extracts

References

- Kucharzik, T.; Dignass, A.U.; Atreya, R.; Bokemeyer, B.; Esters, P.; Herrlinger, K. et al., 2020. Aktualisierte S3-Leitlinie Colitis ulcerosa – Living Guideline. *Z. Gastroenterol.* 58 (12), e241-e326.
- Vissiennon, C.; Goos, K.-H.; Arnhold, J.; Nieber, K., 2017. Spasmolytische und antiinflammatorische Wirkmechanismen eines pflanzlichen Arzneimittels bestehend aus Myrrhe, Kamillenblüten und Kaffeekehl. *Wien Med Wochenschr.* 167 (7-8), 169–176.
- Weber, L.; Kuck, K.; Jürgenliemk, G.; Heilmann, J.; Lipowicz, B.; Vissiennon, C., 2020. Anti-Inflammatory and Barrier- Stabilising Effects of Myrrh, Coffee Charcoal and Chamomile Flower Extract in a Co-Culture Cell Model of the Intestinal Mucosa. *Biomolecules.* 10 (7), 1033.

* Presenting author. E-mail address: julia.werner@medizin.uni-leipzig.de (Julia Werner)

S3.P151 Comparative investigation of frankincense nutraceuticals: pharmacokinetic/pharmacodynamic study

Michael Schmiech^a, Emaad Abdel-Kahaar^a, Judith Ulrich^a, Maximilian Pfeiffer^a, Amira Duweb^{a,b}, Oliver Zolk^c, Tatiana Syrovets^{a,*}, Thomas Simmet^a

^a Institute of Experimental and Clinical Pharmacology, Toxicology, and Pharmacology of Natural Products, Ulm University, Ulm, Germany

^b Institute of Clinical Pharmacology, Brandenburg Medical School, Immanuel Klinik Rüdersdorf, Rüdersdorf, Germany

Frankincense extracts have been widely used in traditional medicine for centuries and more recently in complementary medicine. Therefore, frankincense components such as boswellic and lupeolic acids are of great therapeutic interest (Sterk et al. 2004, Syrovets et al. 2005). Sixteen prescription-free commercial frankincense nutraceuticals marketed in Europe were characterized by high-performance liquid chromatography and tandem mass spectrometry (Schmiech et al. 2019). The frankincense nutraceuticals showed large differences regarding composition and total contents of boswellic and lupeolic acids. Frankincense nutraceuticals significantly inhibited the release of proinflammatory cytokines by LPS-stimulated whole human blood and isolated peripheral blood mononuclear cells. Interestingly, boswellic and lupeolic acid contents in the frankincense nutraceuticals correlated with their ability to inhibit release of proinflammatory cytokines. Among eight different boswellic and lupeolic acids tested, acetyl- β -boswellic acid exhibited the most profound effect on cytokine release.




Fig. 1. Production of frankincense nutraceuticals. *Boswellia* tree grown in Somalia (a), harvesting frankincense oleogum resin through bark cuttings (b), commercial frankincense nutraceuticals (c) (Schmiech et al. 2019).

In a single-dose cross-over clinical trial, pharmacokinetics/pharmacodynamics of two frankincense nutraceuticals, micellar and native, were compared. Administration of the micellar extract increased statistically significant C_{max} , AUC_{t-48} and shortened T_{max} for all boswellic and lupeolic acids compared to the native extract. This resulted in increased relative bioavailability. Administration of both preparations reduced the release of proinflammatory cytokines by LPS-stimulated whole blood. This study demonstrates the mechanisms of the antiinflammatory properties of frankincense nutraceuticals.

Keywords: boswellia, boswellic acid, lupeolic acid, cytokine, triterpenoid

References

- Schmiech, M., Lang, S.J., Ulrich, J., Werner, K., Rashan, L.J., Syrovets, T. and Simmet, Th., 2019. Comparative investigation of frankincense nutraceuticals: Correlation of boswellic and lupeolic acid contents with cytokine release inhibition and toxicity against triple-negative breast cancer cells. *Nutrients* 11: 2341.
- Sterk, V., Büchele, B. and Simmet, Th., 2004. Effect of food intake on the bioavailability of boswellic acids from a herbal preparation in healthy volunteers. *Planta Med* 70: 1155-1160.
- Syrovets, T., Büchele, B., Krauss, C., Laumonnier, Y. and Simmet, Th., 2005. Acetyl-boswellic acids inhibit lipopolysaccharide-mediated TNF- α induction in monocytes by direct interaction with I κ B kinases. *J Immunol* 174: 498- 506.

* Presenting author. E-mail address: tatiana.syrovets@uni-ulm.de (Tatiana Syrovets)

S3.P152 Bioactive potential of deglycosylated flavonoids extracted from immature oranges in skin health

Anne Mandeau*, Lila Zebiri, Quentin Subrenat, Romain Duroux, Julie Leignadier, Joan Attia

R&D department, Lucas Meyer Cosmetics, Toulouse, France

Citrus species are rich in flavonoids, especially the bitter orange *Citrus aurantium* L. In the early stages of maturation, the albedo constitutes a significant portion of the fruit's weight and contains most of the flavonoids naringin and neohesperidin (>80% of the fruit content) (Castillo, 1992).

This study explores the use of immature fruits, which fall during maturation and are discarded as waste, like a valuable resource for producing flavonoid-enriched fruit extracts.

After hydro-alcoholic extraction of these fruits and acidic deglycosylation, the extract containing up to 70% of both naringenin and hesperetin has been obtained, showing comparatively better results than non-hydrolyzed one on *in vitro* anti-inflammatory activity in human dermal fibroblasts.

Knowing the numerous biological activities of the bitter compound hesperetin, the potential involvement of bitter-taste receptors activation was examined. These receptors are known to be expressed beyond the oral cavity, playing a role in immune defense (Patel, 2018). They are also considered in the skin as a warning system to detect toxic compounds, including bacterial toxins.

After identifying the bitter taste receptor T2R14 in the skin, our results indicate that the hydrolyzed orange extract acted as an agonist. By using a T2R14 antagonist, we also demonstrated that this receptor was implicated in mediating the anti-inflammatory response in skin cells stimulated with a bacterial endotoxin (LPS).

Conclusively, our findings reveal that this unique hydrolyzed extracts from immature *Citrus aurantium* L. fruit can activate cutaneous T2R14 receptors, contributing to skin defense mechanisms and attenuating inflammation, thereby highlighting the potential of repurposing agricultural byproducts in cosmetic applications.

Keywords: *Citrus aurantium*, Flavonoids, bitter taste receptors, T2R14

References

- Castillo, J., Benavente, O., del Rio, J.A., 1992. Naringin and neohesperidin levels during development of leaves, flower buds, and fruits of *Citrus aurantium*. *Plant Physiol.* 99, 67-73.
- Patel, N.N., Workman, A.D., Cohen, N.A., 2018. Role of taste receptors as sentinels of innate immunity in the upper airway. *Journal of Pathogens*, Article ID 9541987, 8 pages, <https://doi.org/10.1155/2018/9541987>

* Presenting author. E-mail address: anne.mandeau@lucasmeyercosmetics.com (Anne Mandeau)

S3.P153 Protective effect on H₂O₂-induced oxidative stress in L6 myotubes of *Mitragyna diversifolia* leaf extract

Il-ham Hama^a, Rinnapat Boonsut^b, Supattra Limsuwanchote^b, Juraithip Wungsintaweeikul^{a,*}

^a Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand

^b Department of Pharmacology, Division of Health and Applied Sciences, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand

Mitragyna diversifolia (Wall. ex G. Don) Havil. (MD), also known as kratum [Thai], belongs to the Rubiaceae family and is widely distributed in southern Thailand (Smitinand 2014). This study aimed to evaluate the antioxidative constituents of MD leaf extract. The phytochemical constituents, including total phenolic, total condensed tannin, and total flavonoid contents, were analyzed. Antioxidant activities were assessed using techniques such as the 2,2-diphenyl-1-picrylhydrazyl radical scavenging assay (DPPH), ferric reducing antioxidant power (FRAP), and lipid peroxidation assays. Methanolic crude extracts of leaves, branches, stem barks, and flowers of MD were prepared and screened.

The results indicated that the leaf crude extract (LCE) had the highest total phenolic, total tannin, and total flavonoid contents. The antioxidative capacity of LCE exhibited IC₅₀ values of 459.19 ± 9.08 µg/mL (DPPH) and 150.93 ± 7.59 µg/mL (lipid peroxidation), with an FRAP value of 279.88 ± 2.54 µM gallic acid equivalent/g extract. After fractionating the LCE through the Toyopearl HW-40 column, fractions (Fr) 1-5 were collected. Antioxidant evaluations revealed that Fr 5 demonstrated about a two-fold enhancement of antioxidation. Analyses of Fr 5 using LC-MS/MS and TLC fingerprinting revealed epicatechin as the putative antioxidative agent.

All fractions were further evaluated for their protective effect on H₂O₂-induced oxidative stress. After treating rat L6 myotubes with Fr 1-5 for 24 h, the protective effect found in cells treated with 25 µg/mL of Fr 5. The present study concludes that MD extract has antioxidative potential and reduces oxidative stress in the experimental cell line.

Keywords: *Mitragyna diversifolia*, kratum, antioxidant, oxidative stress, L6 myotubes

References

Smitinand, T., 2014. Thai plant names (revised edition), Forest Herbarium, Royal Forest Department, Bangkok.

* Presenting author. E-mail address: juraithip.w@psu.ac.th (Juraithip Wungsintaweeikul)

S3.P154 *Diospyros lotus* leaf extract and its main component myricitrin inhibit both histamine- dependent and histamine-independent itching

Seon Il Jang^{a,*}, Jae Young Shin^b, Byong Ok Cho^b, Ji Hyeon Park^b, Eun Seo Kang^b, Young-Soo Kim^c

^a Department of Health Management, Jeonju University, 303, Cheonjam-ro, Wansan-gu, Jeonju-si, Jeollabuk-do, 55069, Republic of Korea

^b Institute of Health & Science, Jeonju University, 303, Cheonjam-ro, Wansan-gu, Jeonju-si, Jeollabuk-do, 55069, Republic of Korea

^c Department of Food Science and Technology, Jeonbuk National University, 567, Baekjedae-ro, Deokjin-gu, Jeonju-si, Jeollabuk-do, 54896, Republic of Korea

The *Diospyros lotus* leaf extract (DLE) contains various polyphenols, including myricitrin (MC). Previous studies have demonstrated the alleviating effects of DLE on acute and chronic itching, such as atopic dermatitis. However, there has been a lack of research on the impact of DLE on the central nervous system, including the spinal cord, concerning itching. Therefore, this study aimed to investigate whether DLE and its main component, myricitrin, inhibit the activity and expression of molecules related to itching in the central nervous system and elucidate the underlying mechanisms. In this study, the effects of DLE and MC on histamine-dependent and histamine-independent itching in ICR mice injected with compound 48/80 or chloroquine were investigated through ELISA, immunohistochemistry, immunofluorescence staining, and Western blot analysis. According to the results, both DLE and MC effectively inhibited itching induced by two types of pruritogenic substances. Furthermore, DLE and MC suppressed the infiltration of mast cells and reduced the serum levels of histamine and IL-31. Additionally, this study provided the first *in vivo* evidence that DLE and MC inhibited the expression of itching-related receptors, GRPR and IL31-RA, in the spinal cord and suppressed the activation of STAT3, demonstrating their anti-pruritic effects in the central nervous system. In conclusion, our findings suggest that DLE, along with its main component MC, holds promise as a therapeutic agent for managing itching by targeting both histamine-dependent and histamine-independent pathways.

Keywords: *Diospyros lotus*, myricitrin, itching, CNS, IL-31

References

- Huang, C. C., Kim, Y. S., Olson, W. P., Li, F., Guo, C., Luo, W., et al. 2016. A histamine-independent itch pathway is required for allergic ocular itch. *J. Allergy Clin. Immunol.* 137(4), 1267-1270.
- Yosipovitch, G., Rosen, J. D., & Hashimoto, T., 2018. Itch: From mechanism to (novel) therapeutic approaches. *J. Allergy Clin. Immunol.* 142(5), 1375-1390.
- Ikoma, A., Steinhoff, M., Ständer, S., Yosipovitch, G., & Schmelz, M., 2006. The neurobiology of itch. *Nat. Rev. Neurosci.* 7(7), 535-547.

* Presenting author. E-mail address: sonjjang@jj.ac.kr (Seol Il Jang)

S3.P155 Beneficial effects of *Gryllus bimaculatus* extract on muscle atrophy in dexamethasone- stimulated C2C12 Cells

Seon Il Jang^{a,*}, Jae Young Shin^b, Eun Seo Kang^b, Ji Hyeon Park^b, Ji Hwan Heo^b, Se Yeon Jeong^b, Byong Ok Cho^b

^a Department of Health Management, Jeonju University, 303, Cheonjam-ro, Wansan-gu, Jeonju-si, Jeollabuk-do, 55069, Republic of Korea

^b Institute of Health & Science, Jeonju University, 303, Cheonjam-ro, Wansan-gu, Jeonju-si, Jeollabuk-do, 55069, Republic of Korea

Muscle atrophy, characterized by a progressive decrease in muscle mass due to an elevated ratio of protein degradation to protein synthesis, predominantly affects the elderly population. While various pharmaceuticals have been developed for its treatment, some drugs may induce undesirable side effects. Therefore, the exploration of natural substances with minimal side effects for improving muscle atrophy is imperative. In this study, we investigated the muscle atrophy-improving effects of *Gryllus bimaculatus* extract (BGE) using dexamethasone (DEX)-stimulated C2C12 cells through cell viability, morphological observations, assays, Western blot analysis, and real-time PCR. Results revealed that treatment of differentiated C2C12 cells with GBE promoted recovery from DEX- induced cell death and increased the thickness of muscle cells. Additionally, GBE treatment enhanced the expression of cleaved caspase proteins, inhibitors of cell death, and increased the bax/bcl2 ratio, indicating suppression of apoptosis. Furthermore, GBE treatment upregulated the expression of muscle synthesis factors, including mef2, myogenin, IGF-1, MYF5, and MYF6. Importantly, GBE treatment decreased the expression of muscle atrophy-related proteins, such as myostatin, MAFbx, and FoxO3a. In conclusion, GBE demonstrated promising effects in mitigating DEX-induced muscle atrophy in C2C12 cells by promoting cell survival, inhibiting apoptosis, and modulating muscle synthesis and atrophy-related factors. Further studies are warranted to explore its potential therapeutic applications in addressing muscle wasting conditions.

Keywords: *Gryllus bimaculatus*, muscle atrophy, muscle synthesis factor, muscle breakdown factor, apoptosis

References

- Sani, A., Hasegawa, K., Yamaguchi, Y., Panichayupakaranant, P., & Pengjam, Y., 2021. Inhibitory effects of curcuminoids on dexamethasone-induced muscle atrophy in differentiation of C2C12 cells. *Phytomedicine Plus*. 1(1), 100012.
- Kim, N. H., Lee, J. Y., & Kim, C. Y., 2023. Protective Role of Ethanol Extract of Cibotium barometz (Cibotium Rhizome) against Dexamethasone-Induced Muscle Atrophy in C2C12 Myotubes. *Int. J.Mol. Sci.* 24(19), 14798.
- Lee, Y., Kim, H. M., Kim, J. H., Lee, J. H., Zhang, K. X., Gao, E. M., et al., 2023. Chemical constituents of the *Ajuga multiflora* Bunge and their protective effects on dexamethasone-induced muscle atrophy in C2C12 myotubes. *Nat. Prod. Res.* 37(12), 1978-1985.

* Presenting author. E-mail address: sonjjang@jj.ac.kr (Seol Il Jang)

S3.P156 Anti-inflammatory effect of hyuganin from *Peucedanum japonicum* Thunberg leaves in lipopolysaccharide-stimulated RAW264.7 cells

Byoung Ok Cho^{a,*}, Ji Hyeon Park^a, Eun Seo Kang^b, Ji Hwan Heo^a, Se Yeon Jeong^b, Jae Young Shin^a, Seon Il Jang^a, Jang Hoon Kim^c

^a Institute of Health Science, Jeonju University, Jeonju-si, Jeollabuk-do 55069, Korea

^b Department of Environmental Science and Biotechnology, Jeonju University, Jeonju-si, Jeollabuk-do 55069, Korea

^c Department of Herbal Crop Research, National Institute of Horticultural & Herbal Science, RDA, Eumsung 27709, Korea

Peucedanum japonicum Thunberg has traditionally been used as a medicine to treat cold, pain, stroke, inflammation, and vascular disease in Korea, Japan, and Philippines (Hwang et al., 2022). A hyuganin (Fig. 1) is a natural khellactone coumarin of *P. japonicum* Thunberg (Heo et al., 2020).




Fig. 1. Structure of hyuganin

The goal of the present study was to investigate anti-inflammatory effects and underlying molecular mechanisms of hyuganin isolated from *P. japonicum* Thunberg leaves in lipopolysaccharide (LPS)-stimulated RAW264.7 cells. Anti-inflammatory effects of hyuganin were evaluated using nitric oxide (NO) assay, western blot analysis, real-time PCR, enzyme-linked immunosorbent assay (ELISA), and immunofluorescence assay. The hyuganin treatment reduced the production of NO and prostaglandin E₂ in LPS-stimulated RAW264.7 cells. Hyuganin also suppressed the protein expression of inducible nitric oxide synthase and cyclooxygenase-2 in LPS-stimulated RAW264.7 cells. Hyuganin decreased the secretion of the pro-inflammatory cytokines tumor necrosis factor- α and interleukin-6 in LPS-stimulated RAW264.7 cells. Additionally, hyuganin effectively inhibited NF- κ B activation and mitogen-activated protein kinases (MAPKs) phosphorylation in LPS-stimulated RAW264.7 cells. Furthermore, hyuganin increased the expression of anti-inflammatory enzyme heme oxygenase-1 through activation of nuclear factor erythroid 2-related factor 2 in LPS-stimulated RAW264.7 cells. These results suggest that hyuganin isolated from *P. japonicum* has potential as a candidate for the treatment of inflammatory diseases.

Keywords: *Peucedanum japonicum* Thunberg, hyuganin, anti-inflammation, NF- κ B, MAPKs

References

- Hwang, D., Ryu, H.W., Park, J.W., Kim, J.H., Kim, D.Y., Oh, J.H., Kwon, O.K., Han, S.B., Ahn, K.S., 2022. Effects of 3'-isovaleryl-4'-seneciylkhellactone from *Peucedanum japonicum* Thunberg on PMA-Stimulated Inflammatory Response in A549 Human Lung Epithelial Cells. *J. Microbiol. Biotechnol.* 32, 81-90.
- Heo, J.H., Eom, B.H., Ryu, H.W., Kang, M.G., Park, J.E., Kim, D.Y., Kim, J.H., Park, D., Oh, S.R., Kim, H., 2020. Acetylcholinesterase and butyrylcholinesterase inhibitory activities of khellactone coumarin derivatives isolated from *Peucedanum japonicum* Thunberg. *Sci. Rep.* 10, 21695.

* Presenting author. E-mail address: enzyme21@jj.ac.kr (Byoung Ok Cho)

S3.P157 Triterpene glycosides from the roots of *Deutzia x hybrida* “Strawberry Fields” (Hydrangeaceae) as sweet taste modulators

Efstathia Karachaliou^{a,*}, Antoine Bruguière^a, Marie-José Penouilh^b, Michel Picquet^b, Christine Belloir^a, Loïc Briand^a, Anne Claire Mitaine-Offer^a

^a Centre des Sciences du Goût et de l'Alimentation, CNRS, INRAE, Institut Agro, Université de Bourgogne, 21000 Dijon Cedex, France

^b Plateforme d'Analyse Chimique et de Synthèse Moléculaire de l'Université de Bourgogne (PACSMUB), Université de Bourgogne, BP 47870, 21078 Dijon Cedex, France

Deutzia Thunb. is a genus of the Hydrangeaceae family, among the seven families belonging to the Cornales order (Stevens et al., 2001). The 60 species of this genus are widespread in warm temperate climate, from Southeast Asia to the Philippines, West, and South of North America, also Central America. The *Deutzia* species hold an economic significance as ornamental plants, with hybrids such as *Deutzia x hybrida* or *Deutzia x rosea* (Huang et al., 2001). In garden centers, many cultivars can be found, especially the “Strawberry Fields”, a deciduous shrub with pink flowers bordered with white.

From a phytochemistry point of view, echinocystic acid glycosides, Deutzicoside A and B, have been previously isolated from *Deutzia corymbosa* (Malaviya et al., 1991). Echinocystic acid was found in literature, to possess antiviral, anti-inflammatory and antioxidation activities (Yu et al., 2019). However, information about their influence on the sweet taste receptor TAS1R2/TAS1R3 is currently lacking.

As the main research of our laboratory is the isolation and structural analysis of saponins, and the evaluation of their modulation of the receptors of sweet taste, the perspective of the isolation of echinocystic acid glycosides represents an interesting challenge. So, we chose to study the *Deutzia x hybrida* “Strawberry Fields” cultivar, with a microwave extraction of the air-dried roots. This poster presentation thoroughly describes the isolation protocol using various chromatographic techniques, such as VLC, flash chromatography and MPLC, and the structural analysis of the pure compounds. The evaluation of the activation of the TAS1R2/TAS1R3 receptor will be presented.




Fig. 1. Structure of echinocystic acid derivatives

Keywords: *Deutzia* (x) *hybrida* “Strawberry Fields”, Hydrangeaceae, echinocystic acid, glycosides, sweet-taste receptor

References

- Stevens P. F., 2001 onwards. Angiosperm Phylogeny Website. Available online at: <http://www.mobot.org/MOBOT/Research/APweb/welcome.html> (accessed on 31 January 2024).
- Huang S, Ohba H, Akiyama S, Wu Z.Y., Ravan P.H. (Eds.) 2001. *Deutzia*, Flora of China, Volume 8. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 379–395
- Malaviya N, Pal R, Khanna NM., 1991. Saponins from *Deutzia corymbosa*. *Phytochemistry*, 30, 2798–800.
- Yu H, Li W, Cao X, Wang X, Zhao Y, Song L, Chen J, Wang S, Chen B, Xu Y., 2019. Echinocystic acid, a natural plant extract, alleviates cerebral ischemia/reperfusion injury via inhibiting the JNK signaling pathway. *Eur J Pharmacol.*, 861, 172610.

* Presenting author. E-mail address: efikarahaliou@gmail.com (Efstathia Karachaliou)

S3.P158 Simultaneous analysis of bergapten and schinifoline in *Zanthoxylum schinifolium* Siebold & Zucc. seeds using HPLC and UPLC-MS/MS systems

ChangSeob Seo*

KM Science Research Division, Korea Institute of Oriental Medicine, Daejeon, South Korea

Zanthoxylum schinifolium Siebold & Zucc. belongs to the Rutaceae family and has been widely used as a spice (typically seeds and pericarp) in East Asian countries such as Korea, China, and Japan (Oh and Chung, 2014). The present study focused on developing and validating a simultaneous analytical method for marker substances (bergapten and schinifoline; Fig. 1) in *Z. schinifolium* seeds. This was achieved using high-performance liquid chromatography (HPLC) with a photo-diode array detector (DAD) and ultra-performance liquid chromatography (UPLC) with tandem mass spectrometry systems.

In the regression equation, all markers showed a coefficient of determination of ≥ 0.9990 . Marker recovery was 96.90–105.16% (relative standard deviation (RSD) ≤ 2.23), and the intra- and interday precision was RSD < 3.00 . Bergapten and schinifoline were detected in the seeds at 1.70–2.85 mg/g and 0.19–0.94 mg/g, respectively. This analytical method will improve quality control of *Z. schinifolium* seeds. Additionally, this assay will provide basic data and quality assurance for future biological activity experiments or clinical applications.




Fig. 1. Chemical structures of the two markers selected of *Z. schinifolium* seeds.

Keywords: simultaneous analysis, *zanthoxylum schinifolium*, bergapten, schinifoline

References

Oh, M., Chung, M.S., 2014. Effects of oils and essential oils from seeds of *Zanthoxylum schinifolium* against foodborne viral surrogates. Evid. Based Complement. Alternat. Med. 2014, 135797.

* Presenting author. E-mail address: csseo0914@kiom.re.kr (ChangSeob Seo)

S3.P159 Unveiling the mystery of Australian *Apis mellifera* propolis

Trong D. Tran^{a,b,*}

^a Centre for Bioinnovation, University of the Sunshine Coast, Queensland, Australia

^b School of Science, Technology and Engineering, University of the Sunshine Coast, Queensland, Australia

Bees incorporate plant resins into beeswax to produce propolis (bee glue) for sealing their hives, preventing intruders, and maintaining an antiseptic environment for the colony and larvae (Marcucci, 1995). Propolis has been identified as one of the good sources of bioactive compounds to improve human immunity and prevent a wide spectrum of human diseases such as microbial infections, inflammation, cancer, diabetes, heart, Alzheimer, Parkinson, early aging, and atherosclerosis (Braakhuis, 2019). Over the last three decades, the interest in propolis chemistry and its pharmacological properties has noticeably grown in a world (Tran et al., 2020). However, Australian propolis has remained understudied with paucity in the knowledge of natural diversity and biological activities.

Australia possesses a well-renowned pristine environment owing to the vast stretches of unique and diverse native flora. As a result, Australian propolis is expected to be unique and highly diverse. Here the first study of the quality and chemical diversity of Australian propolis produced by honey bees *Apis mellifera* is presented. The chemical composition, bioactivities, and botanical origin of the premium propolis type which is unique to Queensland state of Australia (Fig. 1) are then introduced.



Fig. 1. Premium Queensland propolis produced by *A. mellifera*

Keywords: *Apis mellifera*, Queensland propolis, chemical diversity, bioactivities, botanical origin

References

- Marcucci, M. C. Propolis: Chemical composition, biological properties and therapeutic activity. *Apidologie*, 1995, 26, 83–99.
- Braakhuis, A. Evidence on the health benefits of supplemental propolis. *Nutrients*, 2019, 11, 2705.
- Tran, T.D.; Ogbourne, S.M.; Brooks, P.R.; Sánchez-Cruz, N.; Medina-Franco, J.L.; Quinn, R.J. Lessons from Exploring Chemical Space and Chemical Diversity of Propolis Components. *Int. J. Mol. Sci.* 2020, 21, 4988.

* Presenting author. E-mail address: ttran1@usc.edu.au (Trong D. Tran)

S3.P160 Chemical profile and antioxidant capacity of Chilean traditional fermented beverages of algarrobo and chañar

Christina Mitsi*, Javier Echeverría

Laboratory of Bioactive Natural Products, Department of Environmental Sciences, Faculty of Chemistry and Biology, University of Santiago of Chile, Chile

Despite being an integral part of the popular nutritional and medicinal culture of Chile since prehispanic time, scientific literature provides little evidence regarding the chemical composition and bioactivity of *chichas*, the traditional Chilean, low-alcohol fermented beverages. Therefore, the present study aimed to investigate the effect of fermentation with *Saccharomyces cerevisiae* on the chemical profile and bioactivity of extracts of fruits of chañar (*Geoffroea decorticans* (Gill. ex Hook. et Arn.) Burkart) and algarrobo (*Neltuma chilensis* (Molina) C.E.Hughes & G.P.Lewis).

Alcoholic fermentation led to a decrease in the concentration of sugars, the generation of ethanol and glycerol and an increase of organic acids content. Moreover, fermentation increased the total phenolic (TPC) and total flavonoid (TFC) contents of the algarrobo and chañar extracts, while chromatographic analyses revealed that fermentation induced both qualitative and quantitative changes at the chemical profile of both plant extracts.

Regarding their antioxidant potential and despite the variability observed between fruits and among tests, the fermentative process increased the *in vitro* antioxidant capacity of the plant extracts as evaluated applying the DPPH[•], FRAP, ABTS^{•+} and CUPRAC tests. This increase of the antioxidant capacity was also observed while evaluating the intracellular ROS levels in Vero and HepG2 cell lines.

Finally, *in vitro* studies suggest that both the fermented and non-fermented extracts of algarrobo and chañar do not have a cytotoxic effect ($EC_{50} > 200 \mu\text{g/mL}$) on the Vero, HUVEC, HepG2 and Caco-2 cell lines.

Keywords: *Geoffroea decorticans*, *Neltuma chilensis*, fermented beverages, antioxidant capacity, cytotoxicity

* Presenting author. E-mail address: christina.mitsi@usach.cl (Christina Mitsi)

S3.P161 Anti-glycation activity of corn silk water extract and establishment of methods for evaluating its quality

Ryuichiro Suzuki*, Aiko Sano

Faculty of Pharmacy and Pharmaceutical Sciences, Josai University, 1-1 Keyakidai, Sakado, Saitama, Japan

This presentation covers the anti-glycation activity of corn silk water extract and the structural elucidation of its active components. The evaluation methods for the quality control of corn silk water extracts will be included using chemometric techniques.

Corn silk is the stigma of *Zea mays* (Poaceae), and dried corn silk is used in traditional medicine as a diuretic. Furthermore, corn silk is also known to prevent diabetes complications, such as nephropathy [Suzuki et al., 2005].

Reducing sugars can non-enzymatically react with amino groups in proteins, forming Schiff bases and Amadori products to produce advanced glycation end products (AGEs). This process is known as glycation. The formation of AGEs induces diabetes complications [Singh et al., 2001].

We previously reported that the water extract of corn silk exhibited stronger anti-glycation activity, and the active compounds in the water extract were lignin-carbohydrate complexes [Sano et al., 2022]. Corn silk's demonstrated inhibitory activity suggests its effectiveness as a valuable material for developing into functional foods.

In brief we aimed to develop prediction methods for the anti-glycation activity of water corn silk extract using spectroscopy, such as ¹H-NMR and FTIR, combined with multivariate statistical analysis. As a result, each corn silk water extract was classified according to its inhibitory activity. Regression models were built for the discriminant analysis. Best-fitting regression models were obtained for the corn silk water extracts. These results indicate that ¹H-NMR and FTIR combined with multivariate chemometrics can be utilized as useful techniques for the detection of active corn silk extracts.

Keywords: *Zea mays*, anti-glycation activity, lignin, chemometrics

References

- Hasanudin, K., Hashim, P., Mustafa, S., 2012. Corn Silk (*Stigma Maydis*) in Healthcare: A Phytochemical and Pharmacological Review. *Molecules* 17, 9697-9715.
- Suzuki, R., Okada, Y., Okuyama, T., 2005. The Favorable Effect of Style of *Zea mays* L. on Streptozotocin Induced Diabetic Nephropathy. *Biol. Pharm. Bull.* 28, 919-920.
- Sano, A., Inoue, Y., Suzuki, R., 2022. Lignin, an active component in the corn silk water extract, inhibits glycation. *Sci. Rep.* 12, 17764.
- Singh, R., Barden, A., Mori, T., Beilin, L., 2001. Advanced glycation end-products: a review. *Diabetologia* 44, 129-146.

* Presenting author. E-mail address: ryu_suzu@josai.ac.jp (Ryuichiro Suzuki)

S3.P162 Bioactivity-Based Molecular Networking for the comparison of *Calophyllum inophyllum* (tamanu) oil obtained via three ecofriendly extraction processes

Émilie Stierlin^a, Raimana Ho^a, Stéphane Greff^b, Elnur Garayev^b, Gaëtan Herbette^c, Sandrine Perino^d, Farid Chemat^d, Rémy Le Guével^e, Phila Raharivelomanana^{a,*}

^a SECOPOL, UMR 241, UPF, BP 6570, 98702 Faa'a, French Polynesia

^b IMBE, UMR 7263 CNRS, AMU-AU, rue de la Batterie des Lions, 13007 Marseille, France

^c AMU, CNRS, Centrale Marseille, FSCM, Spectropole, 13013 Marseille, France

^d Avignon University, INRAE, UMR 408, GREEN Extraction Team, 84000 Avignon, France

^e ImPACcell-BIOSIT, Université de Rennes 1, Campus de Villejean, 35043 Rennes, France

Tamanu oil, obtained from the seeds of *Calophyllum inophyllum* L. (Calophyllaceae), was traditionally used to cure various skin problems and ailments in French Polynesia (Pétard P, 1986). Nuts and especially oil are also used for skin care (Raharivelomanana et al., 2018). It was reported to treat different kinds of skin affections as well as conjunctivitis and used as natural cosmetic ingredient. Aiming to produce high quality tamanu oil from an eco-friendly process, ecological extraction of this oil from nuts had been carried out using different methods such as ultrasound- assisted and microwave-assisted extractions as well Soxhlet extraction using agro-solvents. Bioactivity-Based Molecular Networking was performed in order to compare their eco-extracts chemical composition and cytotoxic properties. Ultrasound assisted extraction seemed to be an innovative and interesting process to extract tamanu oil yielding a rich extract within bioactive compounds being responsible of its cytotoxic activity shown on different cancer cell lines (IC₅₀ < 0.1%V/V): HuH7 (liver), CaCo-2 (colon), MDA-MB-231 (breast), MDA-MB-468 (breast), HCT116 (colon), PC3 (prostate), MCF-7 (breast). The obtained results indicated that the tamanu eco-extracts can be used as an interesting source of natural ingredient for potential nutraceutical and cosmetic applications.

Keywords: *Calophyllum inophyllum*, Eco-extraction, Molecular networking, Cosmetic application

References

Pétard P., 1986. Plantes utiles de Polynésie et rāau tahiti. Ed. Revue et augmentée HaerePo No Tahiti. 224-225. Raharivelomanana P., Ansel J.-L., Lupo E., Mijouin L., Guillot S., Butaud J.-F., Ho R., Lecellier G. and Pichon C., 2018. Tamanu oil and skin active properties: from traditional to modern cosmetic uses. OCL. 25(5).

* Presenting author. E-mail address: phila.raharivelomanana@upf.pf (Phila Raharivelomanana)

S3.P163 Regenerative farming of medicinal crops as a source of robust, sustainable and novel supply chains for the cosmetic industry (BIOSURE)

Simon J. Jackson^{a,*}, Paolo Goi^a, Dario Fornara^b, Enrico Martani^b, Gabriele Costantino^c

^a Davines Group - Botanical Research Department, Via Don Angelo Calzolari 55/A 43126 Parma, Italy

^b Davines Group – EROC (European Regenerative Organic Centre), Via Don Angelo Calzolari 55/A 43126 Parma, Italy

^c University of Parma, Department of Food and Drugs, Parco Area delle Scienze, 27/A, 43124 Parma PR, Italy

The European Regenerative Organic Center, established in Parma, Italy in 2021, focuses on regenerative organic agriculture (ROA) in partnership with the Rodale Institute. This study aims to assess the qualitative and quantitative benefits of growing medicinal crops, particularly in terms of carbon sequestration and secondary metabolite production, compared to commercial farming practices. ROA methods, including minimal soil disturbance, crop rotation, absence of harmful chemicals, cover crops for nitrogen fixation, and increased biodiversity, are examined for their potential to enhance yield and quality of medicinal plants. In 2023, six species (Yarrow, Marigold, Red clover, Chamomile, and Lemon balm) were introduced, and this study investigates the impact of growing conditions on secondary metabolites. Initial findings indicate a significant increase in microbial count and secondary metabolite yield. Parameters evaluated include carbon sequestration, organic matter, nitrogen content, microorganism colony counts, and secondary metabolite production. This study introduces the concept of medicinally interesting Bio-actives, Sustainably sourced and Regeneratively farmed (BIOSURE) using best practice guidelines. Collaborating with the University of Parma, the study compares regenerative practices with commercial agriculture, aiming to develop open-access best practices for laboratory guidance documentation in phytochemical research.



Fig. 1. The 56 test beds set up incorporating medicinal herbs in EROC, Parma, Italy

Keywords: Regenerative farming, Medicinal crops, EROC, Yarrow, ROA, BIOSURE

References

- Fornara D.A. et al 2008, Plant functional composition influences rates of soil carbon and nitrogen accumulation *Journal of Ecology*, 96, 314–322
- De Deyn G.B. et al 2011 Additional carbon sequestration benefits of grassland diversity restoration *Journal of Applied Ecology*, 48, 600–608
- Li Q et al 2016, Grass–legume ratio can change soil carbon and nitrogen storage in a temperate steppe grassland *Soil & Tillage Research* 157 23–31
- Heinrich M et al. 2022, Best Practice in the chemical characterisation of extracts used in pharmacological and toxicological research—The ConPhyMP—Guidelines. *Frontiers in Pharmacology*.13:953205.

* Presenting author. E-mail address: s.jackson@davines.it (Simon James Jackson)

S3.P164 Immunomodulatory activities of aqueous extracts of *Dendrobium officinale* and *Dendrobium nobile* in peripheral blood mononuclear cells of tongue cancer patients

Grace Gar-Lee Yue^a, Si Gao^a, Leanne Lee Leung^b, Jian Xu^c, Hongwen Li^c, Jason Ying-Kuen Chan^b, Clara Bik-San Lau^{a,d,*}

^a Institute of Chinese Medicine and State Key Laboratory of Research on Bioactivities and Clinical Applications of Medicinal Plants, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^b Department of Otorhinolaryngology, Head and Neck Surgery, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

^c Department of Dentistry, Longgang E.N.T. Hospital & Shenzhen Key Laboratory of E.N.T, Institute of E.N.T, Shenzhen, and Shenzhen Longgang Institute of Stomatology, Shenzhen, Guangdong, China

^d Department of Pharmacology and Pharmacy & School of Chinese Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong SAR, China

Dendrobium officinale Kimura et Migo. and *Dendrobium nobile* Lindl. (Orchidaceae family), have long been used as traditional Chinese herbal medicines. The anti-tumor and immunological properties of their polysaccharides have been previously reported (Liu et al., 2011; Wang et al., 2017; Yang et al., 2023). The present study aimed to investigate the immunomodulatory activities of the whole aqueous extracts of *D. officinale* (DOW) and *D. nobile* (DNW) in peripheral blood mononuclear cells (PBMCs) of tongue cancer patients.

Aqueous extracts of the raw herbs of *D. officinale* and *D. nobile* were individually prepared. Blood samples were collected from consented tongue cancer patients, in which PBMCs were isolated and incubated with various concentrations of DOW or DNW for 24 hours. The culture supernatants were then collected for cytokines measurement using ELISA. The proportions of CD3⁺CD4⁺ and CD3⁺CD8⁺ cells in cultured PBMCs were determined by flow cytometry.

Results showed that DOW and DNW (0.8 and 1.6 mg/mL) could modulate the productions of IL-2, IL-10, TNF- α and IFN- γ in PBMCs in concentration-dependent manner. Both DOW and DNW (1.6 mg/mL) significantly increased the populations of cytotoxic T cells (CD3⁺CD8⁺) and helper T cells (CD3⁺CD4⁺). The stimulatory activity of DNW on helper T cells was higher than that of DOW at 0.8 mg/mL.

This is the first report on the immunomodulatory activities of aqueous extracts of *D. officinale* and *D. nobile* in PBMCs of tongue cancer patients, which provide insight for further development of these extracts as adjuvant for tongue cancer patients.

Keywords: *Dendrobium officinale*, *Dendrobium nobile*, tongue cancer, peripheral blood mononuclear cells, cytokines

References

Liu, X.F., Zhu, J., Ge, S.Y., Xia, L.J., Yang, H.Y., Qian, Y.T., Ren, F.Z., 2011. Orally administered *Dendrobium officinale* and its polysaccharides enhance immune functions in BALB/c mice. *Nat Prod Commun.* 6(6), 867-70.

Wang, J.H., Zuo, S.R., Luo, J.P., 2017. Structural analysis and immuno-stimulating activity of an acidic polysaccharide from the stems of *Dendrobium nobile* Lindl. *Molecules.* 22(4), 611.

Yang, W., Chen, D., Ji, Q., Zheng, J., Ma, Y., Sun, H., Zhang, Q., Zhang, J., He, Y., Song, T., 2023. Molecular mechanisms underlying the anticancer property of *Dendrobium* in various systems of the human body: A review. *Biomed Pharmacother.* 165, 115223.

* Presenting author. E-mail address: cbslau@hku.hk (Clara Bik-San Lau)

S3.P165 An innovative bioactive ingredient preparation method without column chromatography and gel filtration: High-purity 3S,3'S-astaxanthin from bioresources

Wei-Cheng Yuan^a, Tung-Ying Wu^b, Pei-Yi Chu^c, Fang-Rong Chang^a, Yang-Chang Wu^{c,*}

^a Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung, Taiwan

^b Department of Biological Science & Technology, Meiho University, Pingtung, Taiwan

^c Graduate Institute of Integrated Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

3S,3'S-Astaxanthin (3S,3'S-AST), an outstanding natural antioxidant that may be beneficial to individuals with degenerative diseases, diabetes, cancer, and cardiovascular disease prevention (Liu et al., 2023; Nishida et al., 2023). However, the preparation methods of 3S,3'S-AST from bioresources require large volumes of solvents, are time-consuming, and have low extraction efficiencies. Therefore, the development of a simple yet systematic and complete preparation for extraction, isolation, and purification of high-purity 3S,3'S-AST is necessary.

In this study, a highly efficient methodology for 3S,3'S-AST preparation from natural microalgae (*Haematococcus pluvialis*) and genetically modified yeast (*Kluyveromyces marxianus*) with a combination of enzyme-assisted extraction and salt-assisted liquid-liquid extraction (SALLE) was achieved. The highest yield of 3S,3'S-AST indicated that FoodPro® CBL for yeast cell walls hydrolysis could significantly enhance extraction and obtain, with the help of SALLE procedure, quantified 3S,3'S-AST 80% in purity from microalgae and 99% in purity from yeast through cation chelation, respectively.

In oxygen radical antioxidant capacity (ORAC) assay, the antioxidant capacity of high-purity 3S,3'S-AST product was 18.3 times higher than that of the original raw material extract. This new 3S,3'S-AST preparation may replace previous methods and has the potential to be scaled up in the manufacture of high-purity 3S,3'S-AST from low-value bioresources of raw materials to high-value products in the food and/or drug industries with lower cost and simple equipment.

Keywords: 3S,3'S-Astaxanthin, *Kluyveromyces marxianus*, SALLE

References

- Liu, X., Xie, J., Zhou, L., Zhang, J., Chen, Z., Xiao, J., Cao Y., Xiao, H. Recent advances in health benefits and bioavailability of dietary astaxanthin and its isomers. *Food Chemistry*, **2023**, 404, 134605.
- Nishida, Y., Berg, P. C., Shakersain, B., Hecht, K., Takikawa, A., Tao, R., Kakuta Y., Uragami C., Hashimoto, H., Misawa N., Maoka, T. Astaxanthin: Past, Present, and Future. *Marine Drugs*, **2023**, 21, 514.

* Presenting author. E-mail address: yachwu@mail.cmu.edu.tw (Yang-Chang Wu)

S3.P166 Bioguided isolation and feature-based identification of bioactive molecules from *Fagraea berteriana* targeting dermal papilla cells and the Wnt pathway

Kristelle Hughes^a, Raimana Ho^b, Stéphane Greff^c, Gaëtan Herbette^d, Edith Filaire^e, E. Ranouille^e, Jean-Yves Berthon^e, Phila Raharivelomanana^{c,*}

^a Pharmacognosy Research Group, Louvain Drug Research Institute, Université Catholique de Louvain, Avenue E. Mounier 72, B1.72.03, B-1200 Brussels, Belgium

^b SECOPOL, UMR 241, University of French Polynesia, BP 6570, 98702 Faa'a, French Polynesia

^c IMBE, UMR 7263 CNRS, IRD, Aix Marseille, Avignon Université, rue de la Batterie des Lions, 13007 Marseille, France

^d Aix Marseille Université, CNRS, Centrale Marseille, FSCM, Spectropole, Service 511, Campus Saint-Jérôme, 13397 Marseille, France

^e Greentech SA, Biopôle Clermont-Limagne, 63360 Saint-Beauzire, France

Fagraea berteriana (Gentianaceae) is a small tree, whose scented flowers are used in French Polynesia as fragrance in the preparation of monoi (Girardi et al., 2015). More unconventionally, its ground fruits were used on cadavers to prevent hair loss (Handy, 1923).

Aiming to understand the effect of the fruits of *F. berteriana* on the hair growth cycle and identify bioactive metabolites, we focused our study on this plant part extract.

Bioguided fractionation of the EtOAc extract of the fruits of *F. berteriana* (FEAE) by Combiflash led to the selection of sub-fraction, FF1, that showed significant hair follicle dermal papilla cells (HFDPCs) proliferation after both 24 and 48h of treatment.

Further investigation of molecular mechanisms revealed a 17% and 50% increase (p-value < 0.001) in β -catenin protein expression and *CCND1* gene expression, respectively, compared to vehicle, of FF1. These findings suggested an upregulation of the Wnt pathway in dermal papilla cells (Hughes et al., 2021).

Several classes of compounds including mainly subclasses of terpenoids were tentatively identified by LC-MS/MS dereplication such as iridoids and secoiridoids (dihydroactinidiolide, boonein, loganic acid or swertiamarin), triterpenoids (*cis/trans-p*-coumaroyloxy maslinic acid and *cis/trans-p*- coumaroyloxy corosolic acid) and other compounds such as flavonoids and coumarins were isolated from FF1.

Statistical analyses highlighted that C-glycosylated flavonoids, iridoids and secoiridoids from the fruits of *F. berteriana*, were positively correlated to this bioactivity, indicating these amongst others, as potential bioactive molecules involved in the proliferation of HFDPCs. These obtained results will be discussed.

Keywords: *Fagraea berteriana*, β -catenin, dermal papilla cells, metabolomics, hair growth

References

Girardi, C., Butaud, J.F., Ollier, C., Ingert, N., Weniger, B., Raharivelomanana, P., Moretti, C., 2015. Herbal medicine in the Marquesas Islands. *Journal of Ethnopharmacology* 161, 200–213.

Handy, E.S.C., 1923. *The Native Culture in the Marquesas*. Bernice P. Bishop Museum, Honolulu, Hawaii.

Hughes, K., Ho, R., Chazaud, C., Hermitte, S., Greff, S., Butaud, J.-F., Filaire, E., Ranouille, E., Berthon, J.-Y., Raharivelomanana, P., 2021. In Vitro Hair Dermal Papilla Cells Induction by *Fagraea berteriana*, a Tree of the Marquesan Cosmetopoeia (French Polynesia). *Cosmetics* 8, 13.

* Presenting author. E-mail address: phila.raharivelomanana@upf.pf (Phila Raharivelomanana)

S3.P167 GABA prevents sarcopenia through regulation of protein turnover and inflammaging in 21- 25-month-old C57BL/6J mice

Boo-Yong Lee*, Hyun-Ji Oh, Gunju Song

Department of Food Science and Biotechnology, College of Life Science, CHA University, Seongnam, Kyeonggi, 13488, Republic of Korea

Sarcopenia is a progressive decline of skeletal muscle mass, strength, and functions in elderly people (Larsson et al., 2019). Sarcopenia, which is age-related muscle atrophy not only decreases the mobility and the exercise capacity but also increases the risks of complications, falls, and mortality, consequently deteriorating the quality of the individual's life and health. Muscle mass and muscle strength reach their maximum around the age of 30, begin to decrease by more than 1% every year after the age of 40, decrease twice as fast after the age of 70 (Pereira et al., 2013).

Here, Gamma-aminobutyric acid (GABA) (**Fig. 1**) is a naturally occurring neurotransmitter synthesized from glutamate by the enzyme glutamic acid decarboxylase (Dhakal et al., 2012). In this study, we investigated the effect of GABA on improving sarcopenia by suppressing muscle protein degradation. GABA (10 or 30 mg/kg/day) was administrated orally daily to young (3-5 months) and old (21-25 months) C57BL/6 mice for 7 weeks.




Fig. 1. Structure of Gamma-aminobutyric acid

The administration of GABA increased gastrocnemius and quadriceps muscle strength, mass and muscle fiber size in old mice. In addition, we found that GABA inhibits sarcopenia by improving muscle protein turnover, which was imbalanced due to aging, through activation of Akt/mTOR/FoxO3a signaling pathways. GABA also regulated the inflammaging that is hallmarks of age-related muscle atrophy, such as the imbalance of M1/M2 macrophage ratio and pro-inflammatory cytokine levels. Thus, GABA can be used in the development of a functional health food customized for the elderly to improve sarcopenia.

Keywords: Sarcopenia, Gamma-aminobutyric acid, Akt/mTOR/FoxO3a signaling pathways

References

- Larsson, L., Degens, H., Li, M., Salviati, L., Lee, Y. I., Thompson, W., ... & Sandri, M. (2019). Sarcopenia: aging-related loss of muscle mass and function. *Physiological reviews*, 99(1), 427-511.
- Pereira, A. D. F., Silva, A. J., Costa, A. M., Monteiro, A. M., Bastos, E., & Marques, M. C. (2013). Muscle tissue changes with aging. *Acta medica portuguesa*, 26, 51-55.
- Dhakal, R., Bajpai, V. K., & Baek, K. H. (2012). Production of GABA (γ-aminobutyric acid) by microorganisms: a review. *Brazilian Journal of Microbiology*, 43, 1230-1241.

* Presenting author. E-mail address: bylee@cha.ac.kr (Boo-Yong Lee)

S3.P168 The influence of disturbed gravity on the content of bioactive phytochemicals in selected sprouts of edible plants from the Brassicaceae family

Marta Grudzińska^{a,b}, Agnieszka Galanty^c, Agata Kołodziejczyk^d, Paweł Żmudzki^e, Ewelina Prochownik^b, Paweł Paśko^b

^a Doctoral School of Medical and Health Sciences, Jagiellonian University Medical College, 16 Łazarza Str., 31-530, Cracow, Poland

^b Department of Food Chemistry and Nutrition, Faculty of Pharmacy, Jagiellonian University Medical College, Medyczna 9, 30-688 Cracow, Poland

^c Department of Pharmacognosy, Faculty of Pharmacy, Jagiellonian University Medical College, Medyczna 9, 30-688 Cracow, Poland

^d Space Technology Centre, AGH University of Technology, 36 Czarnowiejska St, 30-054 Cracow, Poland

^e Department of Medicinal Chemistry, Faculty of Pharmacy, Jagiellonian University Medical College, Medyczna 9, 30-688 Cracow, Poland

One of the studies described that microgravity can increase the antioxidant activity and content of phenolic compounds in mung bean seedlings (Nakajima et al., 2019). This inspired us to examine the influence of disturbed gravity conditions provided by the Random Positioning Maschine on the content of bioactive compounds in the sprouts from four representatives of Brassicaceae family: broccoli, kale, brussels sprouts and kohlrabi. In the first part of the experiment, the process of growing sprouts under microgravity and normal gravity conditions was carried out. Moreover, different lighting conditions (day and night) and harvesting periods (5, 6, 7 days) were tested. Sprouts were extracted with methanol and analyzed quantitatively by HPLC and UPLC-MS/MS (Paśko et al., 2021). The PAL and cytochrome P450, which are responsible for the synthesis of sulfur and phenolic compounds and stress factor - abscisic acid level in the sprouts was determined by ELISA assay.

Qualitative analysis showed the presence of glucosinolates, isothiocyanates and phenolic compounds, which could be responsible for the biological activity of sprouts (Grudzińska et al., 2023). In our study, especially interesting results were obtained for kohlrabi and kale sprouts, in which disturbed gravity as a stress factor caused a significant increase in the content of compounds like: glucoerucin, progoitrin, sinigrin and sulforaphane in comparison to the normal gravity condition.

Because of the insufficient amount of research describing this issue, we wanted to investigate that further and contribute to the discovery of new and innovative methods of harvesting functional foods, such as Brassica sprouts.

Keywords: disturbed gravity, Brassicaceae, sprouts, bioactive compounds, phytochemicals

References

- Grudzińska, M., Galanty, A., Paśko, P., 2023. Can edible sprouts be the element of effective chemopreventive strategy? - A systematic review of in vitro and in vivo study. *Trends Food Sci. Technol.* 139, 104130.
- Nakajima, S., Ogawa, Y., Suzuki, T., Kondo, N., 2019. Enhanced Antioxidant Activity in Mung Bean Seedlings Grown under Slow Clinorotation. *Microgravity Sci. Technol.* 31, 395–401.
- Paśko, P., Galanty, A., Tyszcza-Czochara, M., Żmudzki, P., Zagrodzki, P., Gdula-Argasińska, J., Prochownik, E., Gorinstein, S., 2021. Health Promoting vs Anti-nutritive Aspects of Kohlrabi Sprouts, a Promising Candidate for Novel Functional Food. *Plant Foods Hum Nutr* 76, 76–82.

* Presenting author. E-mail address: marta.grudzinska@doctoral.uj.edu.pl (Marta Grudzińska)

S3.P169 Equivalency of DNA sequencing, HPTLC chromatographic analysis, botanical microscopy, for botanical identity/authentication: A statistical evaluation and quantitation of identification uncertainty

Sidney Sudberg^{a,*}, Khanh Tran^a, Dinah Yu^a, Anthony Fontana^a, Robert LaBudde (Deceased)^b, Zhengfei Lu^c, Yanjun Zhang^c, Adam Faller^c

^a Alkemist Labs, 12661 Hoover Street, Garden Grove, CA 92841, USA

^b Least Cost Formulations, 824 Timberlake Dr., Virginia Beach, VA, 23464, USA

^c Herbalife, 950 190th St., Torrance, CA, 90502, USA

A small-scale single laboratory study was carried out to illustrate methodology for establishing equivalence between different qualitative analytical methods used for identifying botanicals. Samples of different lots from various sources of known inclusion *Mentha x piperita* (known positive) and exclusion *Mentha spicata* (known negative) species were obtained, and measurements made in replicate by DNA NGS sequencing, high performance thin-layer chromatography (HPTLC) and microscopy on each sample. Special statistical methodology for proving the claim of equivalency among methods is illustrated and a predictive model for the evaluation of botanical identification and its associated uncertainty will be presented.

Keywords: *Mentha x piperita*, botanical identification, method equivalency, identification uncertainty, positive predictive value

References

- Do, T.K. et al. 2022. Complementary developing solvents for simpler and more powerful routine analysis by high- performance thin-layer chromatography, JPC – Journal of Planar Chromatography – Modern TLC, 35(3), 299–311.
- Ellison, S. L., & Gregory, S. 1998. Quantifying uncertainty in qualitative analysis. *The Analyst*, 123(5), 1155–1161.
- Gobert, V. et al. 2006. Heterogeneity of three molecular data partition phylogenies of mints related to *M. x piperita* (Mentha; Lamiaceae), *Plant Biology*, 8(4), pp. 470–485.
- Hibbert, B., et. al., 2023. Assessment of performance and uncertainty in qualitative chemical analysis. *Chemistry International*, 45(1), 46–50.
- Monaghan, T. F., et. al., 2021. Foundational statistical principles in medical research: sensitivity, specificity, positive predictive value, and negative predictive value, *Medicina*, 57, 503.

* Presenting author. E-mail address: sidney@alkemist.com (Sidney Sudberg)

S3.P170 Bioassay-guided characterization of the Bergamot (*Citrus aurantium* var. *bergamia*) leaf- extract

Marika Allodi^{a,*}, Beatrice Conversazioni^a, Chiara Spaggiari^a, Gabriele Costantino^a, Sonia Vaccaro^b, Beatrice De Carne^b, Paolo Goi^b

^a Food and Drug Department, University of Parma, Parco Area delle Scienze, 27/A 43124, Parma, Italy

^b Davines Group - Botanical Research Department, Via Don Angelo Calzolari 55/A 43126 Parma, Italy

Nowadays, the use of plant-derived ingredients and sustainable processing methods is becoming increasingly important in the cosmetics industry. In this context, the close collaboration between industry and academia aims to improve the competitiveness of ingredients also in terms of performance, making the characterization of bioactive compounds a key-player in research. The chronic inflammation is one the major responsible of skin disorders, making more and more efforts needed to identify ingredients that counteract the inflammatory process.

The leaves of Bergamot (*Citrus aurantium* var. *bergamia*) are a rich source of polyphenols with potential anti-inflammatory and antioxidant properties (Baron et al., 2021). For this reason, they can be exploited to obtain interesting leaf-extracts for use in cosmetic formulations to alleviate skin disorders.

Our bergamot water leaf-extract, obtained with a rapid solid-liquid dynamic extraction (RSLDE) through Naviglio[®] extractor (Naviglio,2003), once optimized showed a relevant content of flavanones, such as hesperidin, naringenin and neoeriocitrin. Bergamot leaf-extract was analyzed in terms of antioxidant activity by means of ABTS assay, in which it demonstrated distinct radical scavenging properties.

Based on these premises, our investigation now aims to study the anti-inflammatory potential on keratinocytes with *in vitro* assays aimed at reproducing the skin inflammatory state. In particular, HaCat cells will be treated with LPS, capable to induce the phlogosis process, modulate the gene expression of inflammatory mediators, such as IL-1 α , IL-1 β , IL-6, IL-22, TNF- α and their quantitative levels.

Keywords: extract, cosmetics, *Citrus aurantium* var. *bergamia*, inflammation, keratinocytes

References

- Baron, G., Altomare, A., Mol, M., Garcia, J. L., Correa, C., Raucci, A., Mancinelli, L., Mazzotta, S., Fumagalli, L., Trunfio, G., Tucci, L., Lombardo, E., Malara, D., Janda, E., Mollace, V., Carini, M., Bombardelli, E., & Aldini, G. (2021). Analytical Profile and Antioxidant and Anti-Inflammatory Activities of the Enriched Polyphenol Fractions Isolated from Bergamot Fruit and Leave. *Antioxidants* (Basel, Switzerland), 10(2), 141.
- Daniele Naviglio (2003) Naviglio's Principle and Presentation of an Innovative Solid-Liquid Extraction Technology: Extractor Naviglio[®], *Analytical Letters*, 36:8, 1647-1659

* Presenting author. E-mail address: marika.allodi@unipr.it (Marika Allodi)

S3.P171 From nature to hair: using lignin from *Acacia* wood to produce hair conditioners

Carla Varela^{a,b,c,*}, Catarina Fernandes^{a,d}, Solange Magalhães^a, Bruno Medronho^{d,e}, Luís Alves^a, M. Graça Rasteiro^a, Daniela Cabaça

^a CERES, Department of Chemical Engineering, University of Coimbra, 3030-790 Coimbra, Portugal

^b iCBR/CACC/FMUC, University of Coimbra, 3000-548 Coimbra, Portugal

^c CIBB, University of Coimbra, Portugal

^d MED – Mediterranean Institute for Agriculture, Environment and Development & CHANGE – Global Change and Sustainability Institute, University of Algarve, 8005-139 Faro, Portugal

^e FSCN, Surface and Colloid Engineering, Mid Sweden University, Sundsvall, Sweden

Conditioning agents are compounds often used in hair care formulations for physical and aesthetical features improve. Some of the conventional conditioners are produced from non-renewable resources, and can be toxic to aquatic organisms with poor biodegradability. These concerns and the preference for natural products with low environmental impact have prompted the research of renewable feedstocks for the development of cosmetic formulations.

Lignin is a natural aromatic polymer with attractive properties, such as biocompatibility, biodegradability, UV protection, and others. This justifies the increased focus on lignin being widely used for many industrial applications, especially as an additive for cosmetic formulations (Fernandes et al. 2023).

Lignin's hydrophobic nature can help to restore the hydrophobic barrier characteristic of hair, and the existence of functional groups allow for chemical modifications to enhance its interaction with hair, namely by introducing cationic moieties. Hence, our research aims to use lignin as a resource to prepare natural conditioning agents using a natural and abundant source, like wood from *Acacia*, which is an invasive plant in Portugal.

In this work, lignin extracted from *Acacia* wood was chemically modified to prepare cationic derivatives (Figure 1) that can act as hair conditioning agents. Their characterization and inclusion in formulations for further studies of interaction with biomimetic models for hair surface are also presented. It is our belief that new lignin-based conditioning agents will efficiently repair damaged hair without compromising the environment or the human health.




Fig. 1. Overall structure of cationic lignin derivatives.

Keywords: *Acacia* wood, lignin-based derivatives, natural-based hair care conditioners

References

Fernandes, C., Medronho, B., Alves, L., Rasteiro, M.G., 2023. On hair care physicochemistry: From structure and degradation to novel biobased conditioning agents. *Polymers* 15, 608, 1-23.

Acknowledgments

This work was financially supported by the Portuguese Foundation for Science and Technology (FCT) via the Research and Technological Development Project (2022.06810.PTDC), the research PhD grant (2021.05991.BD) and the Strategic Research Centre Project (UIDB/00102/2020).

* Presenting author. E-mail address: carlalvarela@gmail.com (Carla Varela)

S3.P172 Phytochemical exploration and biological evaluation of four *Thymus* L. species from Türkiye

Pierre-Eric Campos^{a,*} Audrey Le Cabec^a, Ebru Özdemir Nath^{b,c}, Gül İpek Gündoğan^d, Emilie Destandau^a, Sefa Sönmez^{b,c}, Gül İpek^d

^a Institute of Organic and analytical chemistry (ICOA), UMR 7311, University of Orléans, France

^b Department of Pharmaceutical Botany, Faculty of Pharmacy, Altınbaş University, İstanbul 34147, Türkiye

^c Natural Products Research and Development Center (DÜAGEM), Altınbaş University, İstanbul 34147, Türkiye

^d Department of Histology and Embryology, Faculty of Medicine, İstanbul Yeni Yüzyıl University, İstanbul 34010, Türkiye

Türkiye is rich in term of plant diversity with a high rate of endemism. Indeed, the country is shared between three biodiversity hotspots: Caucasus, Iran-Anatolia region and the Mediterranean basin (Noroozi et al. 2019). However, even if this richness translates into a high traditional use of plants for medicinal or cosmetic purposes, numerous of these plants remain unexplored in term of phytochemistry.

Ethnobotanical studies revealed that *Thymus* L. species are used for cosmetic purposes. *Thymus brachychilus* Jalas, *Thymus leucostomus* Hausskn. & Velen., *Thymus migricus* Klokov & Des.-Shost., *Thymus sipyleus* Boiss. were obtained from different localities, turned into herbarium material and plant species were determined. To better understand their uses, metabolic activities of ethanolic extracts of these 4 species have been evaluated on Keratinocyte cells (HaCat) and Endothelial cells (HuVeC) while cytotoxicity have been evaluated on L929 fibroblast cells. The extracts didn't show cytotoxicity and were ineffective on HaCat but demonstrated activation of HuVeC which can be promising for wound healing treatments.

A phytochemical study of these extracts has been made by UHPLC-HRMS/MS. Data were represented using molecular networking and annotation tools as SIRIUS4 (Dührkop et al., 2019) and Tima-R (Rutz et al., 2019) were employed to increasing annotation confidence and to refine the compound identification. This methodology allowed us to identify main compounds of these extracts as triterpenoids, such as Uvaol, known for improving the functioning of endothelial cells (Carmo et al., 2020) and flavonoids, including polymethoxylated flavonoids, known for modulating endothelial cell functions (Lee et al., 2014).

Keywords: *Thymus* genus, metabolomics, mass spectrometry, molecular network, bioactivity

References

- Noroozi, J., Zare, G., Sherafati, M., et al. 2019. Patterns of Endemism in Turkey, the Meeting Point of Three Global Biodiversity Hotspots, Based on Three Diverse Families of Vascular Plants. *Front. Ecol. Evol.* 7.
- Dührkop, K., Fleischauer, M., Ludwig, M., et al. 2019. SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure information. *Nat Methods* 16, 299–302.
- Rutz, A., Dounoue-Kubo, M., Ollivier, S., et al. 2019. Taxonomically Informed Scoring Enhances Confidence in Natural Products Annotation. *Front Plant Sci* 10, 1–15.
- Carmo, J., Cavalcante-Araújo, P., Silva, J., et al. 2020. Uvaol Improves the Functioning of Fibroblasts and Endothelial Cells and Accelerates the Healing of Cutaneous Wounds in Mice. *Molecules* 25 (21), 4982.
- Lee, T. H., Jung, H., Park, K. H., et al. 2014. Jaceosidin, a Natural Flavone, Promotes Angiogenesis via Activation of VEGFR2/FAK/PI3K/AKT/NF-KB Signaling Pathways in Endothelial Cells. *Exp Biol Med (Maywood)*, 239 (10), 1325– 1334.

* Presenting author. E-mail address: pierre-eric.campos@univ-orleans.fr (Pierre-Eric Campos)

S3.P173 Isoflavone-rich clover sprouts and their effect on thyroid cells in vitro

Agnieszka Galanty^a, Wojciech Paździora^{a,b,*}, Monika Niepsuj^a, Irma Podolak^a, Marta Grudzińska^{b,c}, Paweł Paśko^c

^a Department of Pharmacognosy, Faculty of Pharmacy, Jagiellonian University Medical College, Medyczna 9, 30-688 Cracow, Poland

^b Doctoral School of Medical and Health Sciences, Jagiellonian University Medical College, 16 Łazarza Str., 31-530, Cracow, Poland

^c Department of Food Chemistry and Nutrition, Faculty of Pharmacy, Jagiellonian University Medical College, Medyczna 9, 30-688 Cracow, Poland

In our previous study we described the effect of red (*Trifolium pratense* L.), white (*T. repens* L.), crimson (*T. incarnatum* L.), Persian (*T. resupinatum* L.) clover sprouts, on different breast and prostate cancer cells. We proved that estrogen-dependent MCF7 breast cancer cells, and androgen- dependent LNCap cancer cells were most susceptible to the tested sprouts, when compared to hormone-independent cell lines. Interestingly, the observed cytotoxic effect was not related to the isoflavone content in the sprouts (Galanty et al., 2022).

As a follow-up, in this study we focused on the effect of the clover sprouts on non-neoplastic (Nthy- ori 3-1) and cancerous (FTC133, TPC1, 8505C) thyroid cells in vitro, as well as determining anti- inflammatory properties. Sprouts were cultured under standard conditions for 3, 5, 7, and 10 days after seeding, and extracted with methanol. The isoflavone content was determined by HPLC-UV- VIS. Cell viability was determined using the MTT assay. NO release was measured by the Griess reaction.

Red clover sprouts were richest in isoflavones (up to 426.2 mg/100 g dw), with the lowest content noted for crimson clover. Formononetin and ononin were dominant isoflavones, with genistein and daidzin in smaller amounts. The sprouts of different clover species showed varied effects on thyroid cancer cells, with papillary carcinoma TPC1 being the most sensitive, and undifferentiated 8505 thyroid cancer cells the most resistant. The strongest cytotoxic effect was mainly observed for the sprouts harvested in 7 and 10 day, irrespective of clover species. Furthermore, at sub-cytotoxic doses the tested sprouts showed high anti-inflammatory potential.

Keywords: *Trifolium* sp., clover sprouts, isoflavones, thyroid cells, cancer

References

Galanty, A., Niepsuj, M., Grudzińska, M., Zagrodzki, P., Podolak, I., Paśko, P., 2022. In the Search for Novel, Isoflavone- Rich Functional Foods - Comparative Studies of Four Clover Species Sprouts and Their Chemopreventive Potential for Breast and Prostate Cancer. *Pharmaceuticals*. 15(7), 806. <https://doi.org/10.3390/ph15070806>

* Presenting author. E-mail address: wojciech.pazdziora@doctoral.uj.edu.pl (Wojciech Paździora)

S3.P174 Anti-histaminic and anti-inflammatory activities of *Ribes nigrum* L. leaf extract CPSP-01

Sophie Holowacz^{a,*}, Anne Leblanc^a, Adrien Brulefert^b, Estelle Wanecz^c, Gérald Chene^c, Claude Blondeau^a, Lucile Berthomier^a, César Cotte C.^a, Valérie Bardot V.^a

^a PiLeJe, 31-35 rue de la Fédération, 75015 Paris, France

^b Qima, 1^{bis} rue des plantes, 86160 Gençay, France

^c Ambiotis, Canal Biotech 2, 3 rue des satellites, 31400 Toulouse, France

The prevalence of allergies is rising dramatically worldwide. Due to the side effects of existing drugs, the search for new solutions remains a challenge, hence the increased interest in exploring the potential of plants. Leaf extracts of *Ribes nigrum* L., with their high proanthocyanidin content, are possible candidates. In our study, we evaluated the *in vitro* potential of a blackcurrant leaf extract, CPSP-01 (aqueous extract, DER 3:1, LC-MS characterized), in various human cell models. Histamine release was assessed using mast cells isolated from plastic surgery and activated by compound 48/80. Pro-inflammatory cytokine production was studied in monocyte-derived M1 macrophages primed with LPS/IFN- γ , as well as in peripheral blood mononuclear cells (PBMC) stimulated with PMA/A23187. Cells were pretreated with 3 non-cytotoxic doses (5.6, 16.7 and 50.0 μ g/ml) of CPSP-01 extract before stimulation with the inducing agent. Cromoglycate and dexamethasone were used as positive controls.

Interestingly, CPSP-01 extract significantly inhibited compound 48/80-induced histamine release in human mast cells in a dose-dependent manner (23% to 90% inhibition). The extract dose-dependently decreased the production of pro-inflammatory cytokines (e.g. IL-6 and TNF- α) following stimulation in both macrophage and PBMC models. CPSP-01 extract also inhibited CD80 marker expression and increased CD163 marker expression, suggesting a shift from M1 polarized macrophages to an M2 phenotype.

Overall, our results suggest that the blackcurrant leaf extract CPSP-01 tested could be of interest in the management of allergy-related symptoms by limiting the release of histamine and pro-inflammatory mediators. Further investigations will be needed to determine underlying mechanisms.

Keywords: *Ribes nigrum*, allergy, histamine, inflammation

* Presenting author. E-mail address: s.holowacz@pileje.com (Sophie Holowacz)

S3.P175 In vitro physiological effects and phytochemical exploration of four *Sideritis* L. species from Türkiye

Ebru Özdemir Nath^{a,b,*}, Pierre-Eric Campos^c, Gül İpek Gündoğan^d

^a Department of Pharmaceutical Botany, Faculty of Pharmacy, Altınbaş University, İstanbul 34147, Türkiye

^b Natural Products Research and Development Center (DÜAGEM), Altınbaş University, İstanbul 34147, Türkiye

^c Institute of Organic and analytical chemistry (ICOA), UMR 7311, University of Orléans, France

^d Department of Histology and Embryology, Faculty of Medicine, İstanbul Yeni Yüzyıl University, İstanbul 34010, Türkiye

Sideritis L. is a member of the Lamiaceae family. There are approximately 218 taxa in the world and 60 taxa in Türkiye belonging to the *Sideritis* genus. Among these 60 taxa, 40 are endemic to Türkiye (Duman, 2012; WFO, 2024). According to ethnobotanical studies, *Sideritis* species have been used in the treatment of skin disease, cold, cough, wound, diarrhoea (Tuzlacı, 2016). The *Sideritis* genus has a wide range of chemical components, including lignans, sterols, iridoids, terpenoids and flavonoids (González-Burgos, 2011).

Four different *Sideritis* taxa were collected for this study from different localities and after identification stored at Herbarium of Altınbaş University, Faculty of Pharmacy (HERA). The ethanolic extracts of these four species were tested for their cytotoxicity on L929 fibroblast cells and their metabolic activities on keratinocyte cells (HaCat) and endothelial cells (HuVeC) in order to better understand their uses. *Sideritis congesta* P.H.Davis & Hub.-Mor. and *Sideritis montana* L. have proven ineffective on keratinocytes but demonstrated activation of dermal cells. Intriguingly, *Sideritis trojana* Bornm. selectively promote the division of HuVeC cells, while *Sideritis germanicopolitana* subsp. *germanicopolitana* Bornm. stimulate a more significant proliferation of fibroblast cells compared to the negative control group.

Extracts have been analysed by UHPLC-DAD-ELSD and are mainly composed by flavonoids and terpenoids. Analysis have shown a large difference in terpenoids composition. UHPLC-HRMS/MS analysis are in progress to refine the compound identification and better understand the differences of physiological effects of these extracts.

Keywords: *Sideritis*, keratinocyte cells, endothelial cells, fibroblast cells, phytochemistry

References

- Duman, H. (2012). *Sideritis*. <http://www.bizimbitkiler.org.tr> (Accessed on 11.02.2024).
- González-Burgos, E.; Carretero, M.E.; Gómez-Serranillos, M.P. 2011. *Sideritis* Spp.: Uses, Chemical Composition and Pharmacological Activities—A Review. *J Ethnopharmacol*, 135, 209–225, doi:10.1016/j.jep.2011.03.014.
- Tuzlacı, E. 2016. Türkiye bitkileri geleneksel ilaç rehberi. İstanbul Tıp Kitabevleri, İstanbul. World Flora Online, *Sideritis*: <https://wfoplantlist.org/taxon> (Accessed on 11.02.2024).

* Presenting author. E-mail address: pharmebru@gmail.com (Ebru Özdemir Nath)

S3.P176 Inhibitory effects of dietary Schisandra supplements on CYP2C19 and CYP2J2 in human liver microsomes

Do-Hyeon Gwon^{a,*}, Zhuoning Liang^a, Bae-Gon Kang^a, So-Young Park^{a,b}, Taeho Lee^a, Song Kyung-Sik^a, Kwang-Hyeon Liu^{a,b}

^a BK21 FOUR Community-Based Intelligent Novel Drug Discovery Education Unit, College of Pharmacy and Research Institute of Pharmaceutical Science, Kyungpook National University, Daegu 41566, Republic of Korea

^b Mass Spectrometry Based Convergence Research Institute, Kyungpook National University, Daegu 41566, Republic of Korea, Republic of Korea

This study investigated the inhibitory effects of five Schisandra supplements on CYP2C19 and CYP2J2 in human liver microsomes. Swanson, Thompson, and Nature's Way supplements displayed notable inhibitory effects on CYP2C19, with IC₅₀ values of 0.011, 0.035, and 0.025 mg/mL, respectively. The contents of dibenzocyclooctadiene lignans in these supplements were higher than those in other two supplements (Planetary Herbals and Only Natural). Dibenzocyclooctadiene lignans have been known to inhibit CYP2C19 enzymes. In addition, only Thompson's product inhibited CYP2J2 activity in human liver microsomes with an IC₅₀ value of 0.010 mg/mL among five Schisandra supplements. The contents of dibenzocyclooctadiene lignans in Thompson were similar to those in Swanson. Therefore, dibenzocyclooctadiene lignans might not be associated with the inhibition of CYP2J2 enzyme. Thompson supplements consist of the extracts of milk thistle, Schisandra, globe artichoke, and dandelion, while other four supplements consist of Schisandra extracts. We are currently confirming the CYP2J2 inhibitory potential of the main components of milk thistle, globe artichoke, and dandelion to identify major components involved in the CYP2J2 inhibition of Thompson supplements.




Fig. 1. Chemical structures of the seven schisandra lignans: (a) deoxyschisandrins; (b) gomisin A; (c) gomisin B; (d) gomisin C; (e) gomisin N; (f) schisandrins; and (g) wuweizisus C.

Keywords: *Schisandra chinensis*, DDI, CYP2C19, CYP2J2, LC/MS-MS

References

- Seo, H.-J.; Ji, S.-B.; Kim, S.-E.; Lee, G.-M.; Park, S.-Y.; Wu, Z.; Jang, D.S.; Liu, K.-H. Inhibitory Effects of Schisandra Lignans on Cytochrome P450s and Uridine 5'-Diphospho-Glucuronosyl Transferases in Human Liver Microsomes. *Pharmaceutics* 2021, 13, 371.
- Kang, Bae-Gon, et al. "Inhibitory Effects of Dietary Schisandra Supplements on CYP3A Activity in Human Liver Microsomes." *Mass Spectrometry Letters* 13.4 (2022): 152-157.

* Presenting author. E-mail address: dohun365@gmail.com (Do-Hyeon Gwon)

S3.P177 Inhibitory effects of dietary *Schisandra* supplements on cytochrome P450 3A in human liver microsomes

Bae-Gon Kang^{a,*}, Do-Hyeon Gwon^a, Kyung-Sik Song^a, Kwang-Hyeon Liu^{a,b}

¹ BK21 FOUR Community-Based Intelligent Novel Drug Discovery Education Unit, College of Pharmacy and Research Institute of Pharmaceutical Sciences, Kyungpook National University, Daegu 41566, Republic of Korea

² Mass Spectrometry Based Convergence Research Institute, Kyungpook National University, Daegu 41566, Republic of Korea

We investigated the contents of dibenzocyclooctadiene lignans (Fig. 1.) in dietary Schisandra supplements and the CYP3A inhibitory potential of dietary Schisandra supplements in human liver microsomes (HLMs). *Schisandra chinensis* and its fruits have been used as a traditional herbal medicine to treat liver dysfunction, fatigue, and chronic coughs. Dibenzocyclooctadiene lignans in four dietary Schisandra supplements were analyzed using liquid chromatography-tandem mass spectrometry (LC-MS/MS). The mean correlation coefficient (r^2) of the respective weighted calibration curves was over 0.992. The limit of quantification (LOQ) for deoxyschisandrin, gomisin A, gomisin B, gomisin C, gomisin N, wuweizisu C, and schisandrin were 5, 30, 15, 5, 5, 5, and 40 ng/ml, respectively. Contents of gomisin B and gomisin C followed in order: Nature's Way > Swanson > Planetary Herbals > Only Natural. In addition, we evaluated the CYP3A inhibitory potential of four dietary Schisandra supplements in human liver microsomes. Midazolam, a representative CYP3A probe substrate, was incubated in HLMs in the presence or absence of dietary Schisandra supplements. After incubation, 1'-hydroxymidazolam (CYP3A-specific metabolite of midazolam) in the samples were analysed by liquid chromatography and triple quadrupole mass spectrometry. At a concentration of 0.05 mg/ml, Schisandra supplements from Nature's Way, Swanson, Planetary Herbals, and Only Natural inhibited CYP3A activity by 93.9, 70.8, 33.6, and 24.8%, respectively. Nature's Way, which exhibited the strongest inhibition against CYP3A, had the highest contents of gomisin B and gomisin C, which potently inhibit CYP3A activity. The in vivo pharmacokinetics of this product should be examined to determine whether the clinical relevance of inhibiting CYP3A activity by dietary Schisandra supplementation.




Fig. 1. Structures of dibenzocyclooctadiene lignans: (a) deoxyschisandrin; (b) gomisin A; (c) gomisin B; (d) gomisin C; (e) gomisin N; (f) schisandrin; and (g) wuweizisu C

Keywords: CYP3A, dietary Schisandra supplements, inhibition, lignan

References

- Seo., H.-J, Ji., S.-B, Kim., S.-E, Lee., G.-M, Park., S.-Y, Wu., Z, Jang., D.S, Liu., K.-H. 2021. Inhibitory Effects of Schisandra Lignans on Cytochrome P450s and Uridine 5'-Diphospho-Glucuronosyl Transferases in Human Liver Microsomes. *Pharmaceutics* 13, 371.
- Szopa., A, Dziurka., M, Warzecha., A, Kubica., P, Klimek-Szczykutowicz., M, Ekiert. 2018. Targeted Lignan Profiling and Anti-Inflammatory Properties of Schisandra rubriflora and Schisandra chinensis Extracts. *H. Molecules* 23, 3103.

* Presenting author. E-mail address: baegon0709@naver.com (Bae-Gon Kang)

S3.P178 Direct extraction and formulation of *Bixa orellana* L. using a green biphasic solvent

Marine Chambaud^{a,b,*}, Rémi Guyomarch^b, Emilie Destandau^a

^a Institut de Chimie Organique et Analytique, University of Orléans, Orléans, France

^b Terre de Couleur, 6 rue de Châtenay, Rochecorbon, France

Achiote (*Bixa orellana* L., Bixaceae) is the source of annatto, one of the main pigments used worldwide. Annatto is obtained from the seeds of achiote and gets its characteristic orange/red colour from apocarotenoids. However, the seeds of achiote also contain volatile compounds and some polyphenols, specifically chalcones, tannins and phenolic acids (Hirko et al., 2022). All of those, and apocarotenoids, have shown activities that can be used in cosmetics such as antioxidant, anti-UV, anti-ageing or antimicrobial (Hoang et al., 2021).

An ultrasound-assisted extraction using a biphasic oil/water solvent allowed the simultaneous extraction of all those compounds. Ultrasounds generate cavitation bubbles that disrupt cell membranes and the oil-water interface, thus facilitating the exchanges between the plant and both phases. The duration of extraction, frequency and plant/solvent ratio were optimised to get the brightest colour in the oil phase and the highest total phenolic content in the aqueous phase. Multiple oils with different compositions (sunflower, coconut, grapeseed, virgin olive, and jojoba oil) were compared to assess their influence on the colour (UV-Vis spectrometry), composition, determined by UHPLC-HMRS completed by a statistical data treatment, and antioxidant activity (DPPH, ABTS) of the extracts. The volatile fraction was characterised by SPME-GC-MS and the aqueous phase by UHPLC-HRMS/MS.

The biphasic solvent allowed the direct formulation of the final extract in an O/W emulsion. A one- pot process combining the extraction and the formulation was then performed. This study shows the possibility of formulating plant bioactive compounds in only one step to obtain multi-functional green products.

Keywords: Green extraction, *Bixa orellana*, antioxidants, HRMS, cosmetics

References

Hirko, B., Getu, A., 2022. *Bixa orellana* (Annatto Bixa): A Review on Use, Structure, Extraction Methods and Analysis.

J. Agron. Technol. Eng. Manag. 5, 687-696.

Hoang, H.T., Moon, J.-Y., Lee, Y.-C., 2021. Natural Antioxidants from Plant Extracts in Skincare Cosmetics: Recent Applications, Challenges and Perspectives. *Cosmetics* 8, 106.

* Presenting author. E-mail address: marine.chambaud@univ-orleans.fr (Marine Chambaud)


S3.P179 Phenylpropanoids and lignans in hemp seeds (*Cannabis sativa L. semen*): identification and quantification

Ilaria Maria Crimi*, Pietro Allegrini^a, Stefania Bosisio, Eric de Combarieu, Nicola Sardone, Francesco Sicolo

Indena SpA, Settala (Milan), Italy

Cannabis sativa L. has been named the plant of “the thousand and one molecules” (Christelle et al., 2016), but a recent database actually claimed the presence of > 6000 identified compounds in commercial Cannabis (Wishart et al., 2024). The phytochemical (secondary metabolites) profile of the plant is dominated by phytocannabinoids in leaves and flowers, and by phenylpropanoids in the seeds, whose high content of essential nutrients (polyunsaturated fatty acids, proteins, vitamins, minerals) make them valuable from a nutritional standpoint (Callaway, 2004). Hemp seeds contain monomeric (phenolic amides) and dimeric (lignanamide) phenylpropanoids (Chaymae et al., 2022), both of nutraceutical interest for their pleiotropic profile of bioactivity (anti-inflammatory, antioxidant, antiviral, anticancer, antidiabetic, antiobesity) (Christelle et al., 2016).

We present a validated HPLC-UV method for the identification and quantification of mono- and dimeric phenylpropanoids in hemp flour, bran and derivatives thereof (Fig.1). This reverse-phase HPLC-UV analytical method can separate, identify and quantify 12 compounds within the two classes of compounds (Fig. 1).



S3.P180 A refined extract of Rosehip modulates skin genes: New evidence for a nutricosmetic

Cynthia G. Suarez^{a,*}, Cecilia Bender^b, Björn Feistel^a, René Roth-Ehrang^a

^a Finzelberg GmbH & Co. KG, Koblenzer Str. 48-56, 56626 Andernach, Germany

^b Istituto Kurz Italia s.r.l., Via Golfo dei Poeti 1/A, 43126 - Parma, Italy

Rose hips are pseudo-fruits from the plants of the *Rosa* genus (Rosaceae) which have been used both for alimentation and for medicinal purposes thanks to their high level content of bioactive compounds (Marmol et al., 2017).

Previous studies have shown an anti-inflammatory effect of a refined aqueous extract from rosehip peels (*Rosa canina* L.) (DEV 4:1), through cytokine release reduction (Walbroel et al., 2009) and TNF alpha inhibition (Feistel et al. 2012).

The purpose of this study was to explore the potential of a standardized Rose hip extract as a “beauty from within” product by examining the changes in gene expression it induces in human keratinocytes. To achieve this, HaCaT cells were subjected to treatments with standardized Rose hip extract or vehicle control. Subsequently, total RNA was extracted and sequenced using RNAseq, a differential gene expression analysis was applied.

The results revealed that the Rose hip extract treatment led to significant ($p < 0.05$) changes in the expression of 5650 genes compared to the control group. Further analysis showed that the Rose hip extract modulates the expression of many genes functionally categorized into pathways related to skin health, including anti-inflammatory function, synthesis of structural proteins of the epidermis, integrity and function of the skin barrier, and antioxidant defense.

These findings provide new exciting evidence for the potential of refined Rose hip extract to promote skin health and beauty through modulation of gene expression in skin cells. Further studies are warranted to validate these findings in vivo and elucidate the underlying mechanisms of action.

Keywords: Rosehip extract, skin health, cosmeceutical, nutraceutical, differential gene expression analysis

References

- Mármol, I., Sánchez-de-Diego, C., Jiménez-Moreno, N., Ancín-Azpilicueta, C., Rodríguez-Yoldi, M.J. 2017. Therapeutic Applications of Rose Hips from Different *Rosa* Species. *Int. J. Mol. Sci.* 18, 1137.
- Walbroel, B & Feistel, B. 2009. A new refined Rose hip extract and its anti-arthritis potential. *Zeitschrift für Phytotherapie*. 2009, 30.
- Feistel B. & Walbroel B. 2012: A purified Rose hip extract – effective in a new joint health concept. Vienna 2012 – congress Phytotherapeutika.; *Zeitschrift für Phytotherapie* 2012; 33, S15

* Presenting author. E-mail address: Cynthia.suarez-rizzo@finzelberg.com (Cynthia G. Suarez)

S3.P181 Complementary LC-MS/MS analyses of alkaloid-rich fractions of the seeds of *Plukenetia volubilis* L. by LTQ Orbitrap XL and ZenoTOF 7600

Kristelle Hughes, Marie-France Hérent, Joëlle Leclercq

Pharmacognosy Research Group, Louvain Drug Research Institute, Université Catholique de Louvain, Avenue E. Mounier 72, B1.72.03, B-1200 Brussels, Belgium

Plukenetia volubilis L. (Euphorbiaceae), better known as Sacha inchi (SI), is an oleaginous vine that produces seeds consumed in several countries in South America. However, while the seeds possess nutritional benefits (Goyal et al., 2022), an assessment of potentially toxic metabolites such as their alkaloid content has only been done by colorimetric assays or IR spectroscopy (Kumar et al., 2020; Srichamnong et al., 2018). The lack of in-depth identification of the type of alkaloids present in the seeds of SI has prevented their acceptance in the European Food Market (Authority (EFSA), 2020). Our aim was thus to investigate the chemical composition of an alkaloid-rich fraction of the defatted seeds, raw (SIRA) and after roasting (SIRO). The two fractions were analyzed by HPLC-HRMS/MS using a LTQ Orbitrap XL and UPLC-HRMS/MS with a ZenoTOF 7600. The main families identified by dereplication and in silico methods with both instruments were cyclopeptides mainly in SIRA, and primary fatty acid amides, more abundant in SIRA than SIRO. Differences in the detection and identification of sphingosine derivatives, true alkaloids and lysophosphatidylcholines between the Orbitrap and ZenoTOF analyses will be further discussed in the poster.

The combined results of the two instruments show a variety of nitrogen-containing compounds. Moreover, our analyses suggest that roasting induces degradation of these compounds, although further quantification is necessary. Mapping the complete alkaloid profile of the seeds will help to better assess its potential toxicity during human consumption.

Keywords: Sacha inchi, alkaloids, ZenoTOF 7600, fatty acid amides

References

- Authority (EFSA), E. F. S. (2020). Technical Report on the notification of roasted seeds from *Plukenetia volubilis* L. as a traditional food from a third country pursuant to Article 14 of Regulation (EU) 2015/2283. *EFSA Supporting Publications*, 17(3), 1817E.
- Goyal, A., Tanwar, B., Kumar Sihag, M., & Sharma, V. (2022). Sacha inchi (*Plukenetia volubilis* L.) : An emerging source of nutrients, omega-3 fatty acid and phytochemicals. *Food Chemistry*, 373, 131459.
- Kumar, B., Smita, K., Debut, A., & Cumbal, L. (2020). Andean Sacha Inchi (*Plukenetia Volubilis* L.) Leaf-Mediated Synthesis of Cu₂O Nanoparticles : A Low-Cost Approach. *Bioengineering*, 7(2), 54.
- Srichamnong, W., Ting, P., Pitchakarn, P., Nuchuchua, O., & Temviriyankul, P. (2018). Safety assessment of *Plukenetia volubilis* (Inca peanut) seeds, leaves, and their products. *Food Science & Nutrition*, 6(4), 962-969.

* Presenting author. E-mail address: kristelle.hughes@uclouvain.be (Kristelle Hughes)

S3.P182 Compounds of edible fruits vs. bacterial nucleases – *in silico* study

Monika E. Czerwińska^{a,*}, Łukasz Szeleszczuk^b

^a Department of Biochemistry and Pharmacogenomics, Medical University of Warsaw, Banacha 1, 02-097 Warsaw, Poland

^b Department of Organic and Physical Chemistry, Medical University of Warsaw, Banacha 1, 02-097 Warsaw, Poland

The extracts from fruits of *Hippophaë rhamnoides* L. (HRE), *Chaenomeles japonica* (Thunb.) Lindl. ex Spach (CJE), and *Cornus mas* L. (CME) are inhibitors of hydrolases, such as pancreatic lipase and α -amylase, engaged in the digestion of fats and polysaccharides (Siegien et al., 2021). To continue our studies we raise the question of whether the constituents of these plant materials influence bacterial hydrolases participating in the replication of nucleic acids.

The study aimed to evaluate the binding mode of the most abundant constituents of HRE, CJE, and CME with the bacterial nucleases *in silico*. The molecular docking study of isorhamnetin glycosides, procyanidins B1 and B2, catechin/epicatechin (Siegien et al. 2021), as well as loganic acid and cornuside (Olędzka et al., 2022) with bacterial enzymes was performed. Endonuclease 1 (EC 3.1.21.1), colicin E9, and ribonuclease H (EC 3.1.26.4) from *Escherichia coli* were taken into consideration.

The results of molecular docking for selected compounds with endonuclease 1 from *E. coli* are shown in Fig. 1. All three ligands (Fig. 1) bind in the same region of the protein and form interactions with common residues.




Fig. 1. Molecular docking results for isorhamnetin-3-O- β -D-glucosyl-7-O- α -L-rhamnoside (A), procyanidin B1 (B), and loganic acid (C) with endonuclease 1 from *E. coli*.

The most relevant flexibility of binding mode with *E. coli* endonuclease 1 was observed in the case of procyanidin B1. Six different poses for this ligand have been obtained, with significantly various orientations of the molecule in the binding site. The obtained complexes have been then subjected to molecular dynamics and MM/GBSA calculations.

Keywords: *Hippophaë rhamnoides*, *Chaenomeles japonica*, *Cornus mas*, gut microbiota

References

- Siegien, J., et al. 2021 Pancreatic lipase and α -amylase inhibitory activity of extracts from selected plant materials after gastrointestinal digestion *in vitro*. Food Chem. 355, 129414
- Olędzka, A., et al. 2022. Potentially bio-accessible metabolites from an extract of *Cornus mas* fruit after gastrointestinal digestion in vitro and gut microbiota ex vivo treatment. Nutrients. 14, 2287

Acknowledgments

The study was financially supported by the Medical University of Warsaw (grant no. WF1A/1/F/GW/N/23).

* Presenting author. E-mail address: monika.czerwinska@wum.edu.pl (Monika Czerwińska)


S3.P183 Iridoids from *Stenaria nigricans* and their insect repellency assessment

Fazila Zulfiqar^{a,*}, Abbas Ali^a, Zulfiqar Ali^a, Ikhlas A. Khan^{a,b}

^a National Center for Natural Products Research, School of Pharmacy, University of Mississippi, University, MS, USA

^b Division of Pharmacognosy, Department of BioMolecular Sciences School of Pharmacy, University of Mississippi, University, MS, USA

The methanolic extract of the aerial parts of *Stenaria nigricans* (Lam.) was tested for repellency against mosquitoes and imported hybrid fire ants, which showed biting deterrence similar to DEET against *Aedes aegypti* L. Following the bioassay-guided approach, seven iridoid glycosides were isolated from the active subfractions of the methanolic extract. Their structures were elucidated based on the 1D and 2D NMR and mass techniques. The isolates were tested for their biting deterrent activity and some of them showed the potential to be used as repellents against mosquitoes based on BDI values with biting deterrence similar to DEET. In an in vitro digging bioassay, none of these compounds showed repellency against hybrid imported fire ants.



Keywords: *Stenaria nigricans*, Rubiaceae, mosquitoes, iridoid glycosides, biting deterrent

Presenting author. E-mail address: fazila@olemiss.edu (Fazila Zulfiqar)

S3.P184 Natural deep eutectic solvents: a sustainable alternative for the extraction and stabilization of carotenoids and chlorophylls from spinach leaves

Ramla Sahli*, Trishiyaa Ragit Jeevakumar, Iron Mike Ardeza, Clément Wahl, Leslie Boudesocque-Delaye^a

EA 7502 Synthèse et Isolement de Molécules BioActives (SIMBA), Université de Tours, 31 Avenue Monge, 37200 Tours, France

The trends in green extraction focus on cost-effective and eco-friendly solvents. Natural Deep Eutectic Solvents (NaDES) are among the most promising options. They have proven to be useful in the sustainable production of bioactive plant-derived natural extracts. Spinach leaf extracts (*Spinacia oleracea*) are widely recognized as valuable sources of bioactive compounds, particularly chlorophylls and carotenoids (Hayes et al., 2020). Bajkacz et al. (2020) reported a green method for chlorophyll extraction using NaDES. The aim of this study is to improve the extraction efficiency of chlorophylls and carotenoids from spinach leaves using NADES in combination with ultrasound- assisted extraction (UAE). We investigated hydrophobic and hydrophilic NaDES to evaluate their selectivity and stabilization ability towards these fragile pigments. In order to maximize the concentration of the target compounds, biomass preparation and UAE parameters were optimized. The global contents of chlorophylls and carotenoids were estimated using high-throughput 96-well plate spectrophotometric assays. Additionally, the NaDES extract was characterized using liquid chromatography-diode array detector (UHPLC-DAD). The results indicated that hydrophobic NaDES provided extracts rich in carotenoids, while hydrophilic NaDES were more effective in stabilizing chlorophylls. To improve extraction selectivity, a one-step solid/liquid/liquid extraction was implemented. This work aims to produce enriched spinach extracts that can be used as natural colorant or additive ingredients, or as ready-to-use extracts.

Keywords: *Spinacia oleraceae*, NaDES, ultrasound assisted extraction, chlorophylls, carotenoids

References

- Hayes M., Pottorff M., Kay C., Van Deynze A., Osorio-Marin J., Lila M.A., Iorizzo M., Ferruzzi M.G., 2020. In Vitro Bioaccessibility of Carotenoids and Chlorophylls in a Diverse Collection of Spinach Accessions and Commercial Cultivars. *J Agric. Food Chem.* 68 (11), 3495-3505.
- Bajkacz, S., Rusin, K., Wolny, A., Adamek, J., Erfurt, K., Chrobok, A., 2020. Highly efficient extraction procedures based on Natural Deep Eutectic Solvents or Ionic Liquids for determination of 20-hydroxyecdysone in spinach. *Molecules.* 25:20, 4736.

* Presenting author. E-mail address: ramla.sahli@univ-tours.fr (Ramla Sahli)

S3.P185 Baking a difference: elevating bread with *Portulaca oleracea* L. seeds

Viana Castañeda-Loaiza^a, Maria João Rodrigues^{a,*}, Zoltán Cziáky^b, Luísa Custódio^a

^a Centre of Marine Sciences, University of Algarve, Faculty of Sciences and Technology, Ed. 7, Campus of Gambelas, 8005-139 Faro, Portugal

^b Agricultural and Molecular Research and Service Institute, University of Nyíregyháza, Nyíregyháza, Hungary

Portulaca oleracea L. seeds, rich in fatty acids like linoleic, palmitic, α -linolenic, and oleic acids, have been explored for their potential to enhance food products (Mousavi et al., 2017). Incorporating these seeds in powdered or whole form into bread has been studied to boost nutritional value (Sirvam et al., 2010). Experiments involved adding seed powder and whole seeds to wheat flour for bread-making at varying concentrations (0, 5, 10, 15, and 30%), resulting in thirteen bread variants. These were analyzed for mass loss upon baking, moisture, firmness, and consumer preference through sensory tests. The most preferred formulations contained low seed concentrations (5% seed powder, 5% and 10% whole seeds). Further analyses of these breads showed they were rich in nutrients, antioxidants, and specific metabolites like gentisic acid-O-glucoside, gentisic, ferulic, and isoferulic acids, which were not found in the control bread. Low seed concentrations improved the bread's lipid and fibre content, and antioxidant capacity, and maintained low levels of anti-nutritional factors such as condensed tannins, α -amylase, and trypsin inhibitors. It is concluded that adding small amounts of *P. oleracea* L. seeds to bread can enhance its nutritional and functional qualities while minimizing antinutritional compounds.

Keywords: functional food, bread, *Portulaca oleracea* L., nutritional composition, sensory evaluation

References

- Mousavi, S. R. J., & Niazmand, R., 2017. Fatty acids composition and oxidation kinetic parameters of purslane *Portulaca oleracea* seed oil. *Agricultural Research*, 6, 421-426.
- Sivam, A. S., Sun-Waterhouse, D., Quek, S., & Perera, C. O., 2010. Properties of bread dough with added fiber polysaccharides and phenolic antioxidants: A review. *Journal of Food Science*, 75(8), R163-R174.

Acknowledgments

This research was funded by Portuguese national funds from the FCT - Foundation for Science and Technology, through projects UIDB/04326/2020, UIDP/04326/2020, LA/P/0101/2020, and the GreenValue (PTDC/BAAAGR/1391/2020). VCL received a PhD grant from FCT (2020.04541.BD), MJR benefited from an FCT program contract (UIDP/04326/2020), and LC was supported by the FCT Scientific Employment Stimulus (CEECIND/00425/2017).

* Presenting author. E-mail address: mjrodrigues@ualg.pt (Maria João Rodrigues)

S3.P186 Chemical and sensory profile of different oregano individuals originated from Vertisco mountain

Aggeliki Ainalidou^a, Giannis Mourtzinou^b, Konstantinos Klimantakis^b, Petroula Tsitlakidou^b, Eirini Vallianatou^c, Katerina Karamanoli^{a*}

^a Department of Field Crops and Ecology, School of Agriculture, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece

^b Department of Food Science and Technology, School of Agriculture, Aristotle University of Thessaloniki, Thessaloniki, Greece

^c Symbeeosis, 19002, Paiania, Greece

Current study aims to evaluate botanical, chemical and sensory characteristics of different oregano individuals cultivated in the same field and originated from a natural population of Vertisco mountain in Greece. Fourteen oregano individuals (coded: I1-I14) that differ in their morphological characteristics, taxonomical classified as *Origanum vulgare* subsp. *hirtum* (Link) Ietswaart, also known as “Greek oregano”, were selected for further investigation. The *intended outcome* was to select the most prominent individuals in order to use them for propagation and mass production of high value planting material. Essential oil composition and its antioxidant capacity of all oregano individuals were explored. Aroma properties of selected individuals were further determined by a trained sensory panel using quantitative descriptive analysis (QDA). According to obtained results, essential oil yield ranged from 1.63 % to 5.98% d.w. in individuals I7 and I13, respectively. The main constituent of all essential oils was carvacrol with the highest value in I5 (81.217%). Further, evaluation of antioxidant capacity revealed I4 as the most active individual. High level of phenolic compounds was recorded in I13, I4, and I8, whilst I8 and I13 biotypes had also the higher levels of flavonoids. The examination of aroma attributes by the the sensory panel, indicated that I5 and I13 exhibited the most prominent odour properties. Overall, our results indicated that I5 and I13 are the oregano individuals with the superior combination of characteristics, and possible interest for food industry, thus these individuals were selected for mass production of better observed quality planting material.

Keywords: Greek oregano, carvacrol, antioxidant capacity, odor properties, sensory

Acknowledgments

This research has been conducted in the frame of the Regional Operational Programme Central Macedonia 2014-2020 (“Oregano Standardization and Added Value elaboration - O-SAVE”, action code: KMP6-0284665) that was co-financed by Greek national funds and the European Union (European Regional Development Fund).

Presenting author. E-mail address: katkar@agro.auth.gr (Karamanoli Katerina)

S3.P187 Valuing the medicinal flora of Côa valley (Portugal) through the development of new plant- based cosmetics

Mário Pedro Marques^{a,b,*}, Euclides Landim^{a,b}, Carla Varela^{a,b,c}, Joana Marques^d, Ricardo M. F. da Costa^d, Luís A. E. Batista de Carvalho^d, Ana Silva^{b,e,f}, Maria Teresa Cruz^{b,e,f}, Maria Inês Dias^{g,h}, Lillian Barros^{g,h}, Aida Carvalho^{i,j,k}, Paulo J. Oliveira^{b,e}, Célia Cabral^{a,b,l,*}

^a University of Coimbra, iCBR/CACC, FMUC, 3000-548 Coimbra, Portugal

^b University of Coimbra, CIBB/UC, 3000-548 Coimbra, Portugal

^c University of Coimbra, CERES, Department of Chemical Engineering, Coimbra, Portugal

^d University of Coimbra, QFM-UC, Rua Larga, 3004-535 Coimbra, Portugal

^e University of Coimbra, CNC-UC 3004-504 Coimbra, Portugal

^f University of Coimbra, FFUC, 3000-548 Coimbra, Portugal

^g Polytechnic Institute of Bragança (IPB), CIMO, 5300-253 Bragança, Portugal

^h IPB, SusTEC, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

ⁱ IPB, Campus de Santa Apolónia, 5300-253 Bragança, Portugal

^j CiTUR, Pólo Guarda, Av. Dr. Francisco Sá Carneiro 50, 6300-559 Guarda, Portugal

^k Fundação Côa Parque, Rua do Museu, 5150-620 Vila Nova de Foz Côa, Portugal

^l University of Coimbra, CFE/UC, Calçada Martim de Freitas, 3000-456 Coimbra, Portugal

From the ethnobotanical survey carried out in the Côa Valley Archeologic Park during our project CôaMedPlants, we recorded over 500 species with promising medicinal properties. From those, we selected the 9 species more representative in traditional medicine for the treatment of skin diseases. Considering these traditional uses, in this work we evaluated their potential as natural ingredients for cosmetics. Hence, hydroalcoholic extracts (EtOH 80%) were prepared, chemically characterized through FTIR-ATR and HPLC-ESI-MSⁿ, and the antioxidant activity was assessed through cell-free techniques. The cytotoxicity of extracts was determined *in vitro* using Normal Human Dermal Fibroblasts (NHDF). The majority of these nine plants' extracts did not present toxicity up to 0.2 and 0.4 mg/mL. These non-toxic concentrations were evaluated as cytoprotective for NHDF against the oxidative toxicity induced by 0.5 mM *tert*-butyl hydroperoxide and 1.5 mM hydrogen peroxide. From these assays, a very significant cytoprotective potential was found with *Cistus ladanifer* L. subsp. *ladanifer*, *Cistus salviifolius* L., and *Pistacia terebinthus* L. that showed a protection of about 20- 30% in metabolic activity. Noteworthy, the most cytoprotective ones were then evaluated concerning skin irritation by using the SkinEthic[™] Reconstructed Human Epidermis (RHE) model, in compliance with the OECD Test Guideline No. 439, revealing that none of these extracts presented irritant activity. In conclusion, five out of the nine studied plant species (*P. terebinthus*, *C. salviifolius*, *C. ladanifer*, *Thymus mastichina* (L.) L. subsp. *mastichina* and *Verbascum pulverulentum* Vill.) showed outstanding antioxidant and cytoprotective potential, thus guiding for further development into plant-based cosmetics.

Keywords: Côa Valley (Portugal), antioxidant, cytotoxicity, cytoprotection, skin irritation

Acknowledgments: This project is financed by FCT, Portugal, through the following projects: UIDB/04539/2020 - DOI: 10.54499/UIDB/04539/2020, UIDP/04539/2020 - DOI: 10.54499/UIDP/04539/2020, LA/P/0058/2020 - DOI: 10.54499/UIDP/04539/2020 and CôaMedPlants (COA/BRB/0019/2019 - DOI: 10.54499/COA/BRB/0019/2019). MPM is supported by a Foundation for Science and Technology (FCT) PhD grant (Reference: PRT/BD/153391/2021). EL is supported by a Foundation for Science and Technology (FCT) PhD grant (Reference: PRT/BD/154364/2022). These PhD grants are financed by national budget and co-financed through the European Social Fund (ESF) and the Regional Operational Por_Centro.

*Presenting author. E-mail address: mpedromarques1995@gmail.com (Mário Pedro Marques)

S3.P188 Identification of anti-inflammatory compounds in *Parasenecio adenostyloides* using bioactivity-based molecular networking

Yoo Kyong Han^{a,*}, Yoon-Young Sung^a, Heung Joo Yu^a, Chang-Hwa Song^b, Dong-Seon Kim^a

^a KM Science Research Division, Korea Institute of Oriental Medicine, 1672 Yuseongdae-ro, Yuseong-gu, Daejeon 34054, Republic of Korea

^b Department of microbiology, College of Medicine, Chungnam National University, 266 Munhwa-ro, Jung-gu, Daejeon 35015, Republic of Korea

In Korea, the genus *Parasenecio* consists of about 10 species. Among them, *Parasenecio adenostyloides* (PA) is a perennial herb of the Asteraceae family and is a rare plant with a narrow global distribution area, and its young leaves are edible (Nam, 2006). But there have been no previous research papers on the constituents and bioactivities of this plant. In this study, we obtained the chemical profile of the PA extract by component analysis using LC-QTOF-MS/MS. And, it was confirmed that the PA extract had a significant inhibitory activity in a concentration-dependent manner on the expression of RANTES, an important cytokine that mediates eosinophil chemotaxis, in TNF- α -induced human bronchial epithelial cells (BEAS-2B). Based on these results, we applied the bioactivity-based molecular networking concept to identify candidate active molecules directly from fractionated bioactive extracts (Nothias, 2018). The ten fractions (PAF 1-10) obtained by performing column chromatography on the extract and the inhibitory activity of each fraction were converted into scores, which were mapped to the molecular networking results. Through this, three compounds (PAC 1-3) were identified, including not only the main peak of the PAF 4, which showed the highest activity, but also the structurally similar constituents contained in the cluster. And these identified compounds showed significant inhibitory activity on RANTES expression in BEAS-2B cells. Our results suggested that bioactivity-based molecular networking could be useful for effectively discovering active constituents in unknown ingredients and that PA could be a useful functional ingredient for respiratory diseases such as asthma.

Keywords: *Parasenecio adenostyloides*, Asteraceae, molecular networking, RANTES, BEAS-2B cell

References

- Nam, G.H., Park, M.S., Jeong, H.J., Chung, G.Y., 2006. Taxonomic study on the genus *Parasenecio* (Compositae) of Korea by the morphology and somatic chromosome numbers. Korean J. Plant Res. 19, 323-330.
- Nothias, L.F., Nothias-Esposito, M., Silva, R.D., Wang, M., Protsyuk, I., Zhang, Z., Sarvepalli, A., Leyssen, P., Touboul, D., Costa, J., Paolini, J., Alexandrov, T., Litaudon, M., Dorrestein, P.C., 2018. Bioactivity-based molecular networking for the discovery of drug leads in natural product bioassay-guided fractionation. J. Nat. Prod. 18, 758-767.

* Presenting author. E-mail address: yookyong05@kiom.re.kr (Yoo Kyong Han)

S3.P189 Metabolic response of *Panax ginseng* to root-rot: a molecular networking approach

Kyu Won Kim^{a,b,*}, Kyung Ho Lee^{b,c}, Yena Kim^{b,c}, Misnik Choi^{a,b}, Geonha Park^b, Young Pyo Jang^{a-c}

^a Department of Biomedical and Pharmaceutical Sciences, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

^b Division of Pharmacognosy, College of Pharmacy, Kyung Hee University, Seoul 02447, Republic of Korea

^c Department of Integrated Drug Development and Natural Products, Graduate School, Kyung Hee University, Seoul 02447, Republic of Korea

The quality of *Panax ginseng* C.A. Meyer root is profoundly influenced by soil microorganisms. It is known that the diversity of microbial species decreases frequently as the ginseng matures. This change leads to the presence of more pathogenic fungal species like *Ilyonectria destructans*(syn. *Cylindrocarpon destructans*) and changes in soil condition (Nguyen et al., 2016) (Farh et al., 2018), causing persistent root rot and significantly reducing ginseng's value and economic viability (Yeon et al.,2007). Despite ongoing efforts to address root rot, there remains a notable research gap regarding changes in ginseng root metabolites caused by the disease. This study was designed to employ molecular networking to analyze the correlation between root rot and ginseng root metabolites. Extracts from normal and diseased ginseng root were subjected to analysis using High Performance LC-ESI-Tandem Mass Spectrometry, and molecular networking was performed with GNPS. The resulting molecular network displayed the formation of clusters and nodes. The ratio of nodes between groups was evaluated, and annotation and identification were conducted using in silico tools such as SIRIUS and NAP, particularly for nodes exhibiting an increasing trend after strain inoculation. This study presented metabolites that were significantly increased due to the root rot. It is suggested as a dammarane-based derivative (C₃₈H₆₂O₈) and an oligopeptide-based material (C₁₉H₃₈N₈O₅, C₂₃H₃₈N₈O₄). However, neither substance has been reported in *P. ginseng*. Further experiments are needed to identify these substances and the structure. Significance of this study lies in providing a metabolomics-based foundation for further research on this disease.

Keywords: *Panax ginseng* C.A. Meyer, root rot, metabolomics, molecular networking, ESI-MS/MS

References

- Nguyen, N. L., Kim, Y. J., Hoang, V. A., Subramaniam, S., Kang, J. P., Kang, C. H., & Yang, D. C. (2016). Bacterial diversity and community structure in Korean ginseng field soil are shifted by cultivation time. *PloS one*, 11(5), e0155055.
- Farh, M. E. A., Kim, Y. J., Kim, Y. J., & Yang, D. C. (2018). *Cylindrocarpon destructans*/*Ilyonectria radicola*-species complex: Causative agent of ginseng root-rot disease and rusty symptoms. *Journal of ginseng research*, 42(1), 9-15.
- Yeon, B. Y., Hyeon, G. S., Bae, Y. S., Lee, S. W., Seong, N. S., & Kang, S. W. (2007). Changes of soil chemical properties and root injury ratio by progress years of post-harvest in continuous cropping soils of ginseng. *Korean Journal of Medicinal Crop Science*, 15(3), 157-161.

* Presenting author. E-mail address: rbdnjs1227@naver.com (Kyu Won Kim)

S3.P190 Collagenase and tyrosinase inhibitory compounds from fish gut bacteria *Ruegeria atlantica* and *Pseudoalteromonas neustonica*

Jonghwan Kim^{a,*}, Su Jung Hwang^b, Gyu Sung Lee^a, Ju Ryeong Lee^a, Hye In An^b, Hong Sik Im^c, Minji Kim^a, Sang-Seob Lee^c, Hyo-Jong Lee^b, Chung Sub Kim^{a,b}

^a Department of Biopharmaceutical Convergence, Sungkyunkwan University, Suwon 16419, Republic of Korea

^b School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea

^c College of Biotechnology and Bioengineering, Sungkyunkwan University, Suwon 16419, Republic of Korea

Ruegeria atlantica and *Pseudoalteromonas neustonica* are fish gut bacteria isolated from the gut of *Pagrus major* (Yoshihito et al., 1998) and *Acanthopagrus schlegelii* (Hwang et al., 2016), respectively. A total of 22 compounds (**1–22**) were isolated from these two bacteria; 16 compounds (**1–16**) from *R. atlantica* and 6 compounds (**17–22**) from *P. neustonica*. Their chemical structures were elucidated by spectroscopic and spectrometric data analysis and chemical synthesis. Compounds **11** and **13** showed strong collagenase inhibitory activities of 31.91 and 36.43% at 20 μ M, respectively, comparable to or surpassing that of the positive control, epigallocatechin gallate (EGCG, 34.66%). Compounds **11** and **14** also showed mild tyrosinase inhibitory activity with 6.73 and 13.68%, respectively. All the tested compounds displayed no significant antibacterial activity against *Escherichia coli* and *Bacillus subtilis* up to 100 μ M. The collagenase- and tyrosinase-inhibitor compound **11**, cyclo(L-Pro-D-Leu), was found to be stable under heat (50°C) and UV light (254 and 365 nm) for up to 6 days. These results indicate that compound **11** could be developed into a cosmeceutical with anti-aging properties.




Fig. 1. The anti-aging compounds isolated from fish gut bacterium, *Ruegeria atlantica*.

Keywords: anti-collagenase, anti-tyrosinase, metabolites, *Ruegeria atlantica*, *Pseudoalteromonas neustonica*

References

- Hwang, C. Y., Lee, I., Hwang, Y. J., Yoon, S. J., Lee, W. S., Cho, B. C., 2016. *Pseudoalteromonas neustonica* sp. nov., Isolated from the Sea Surface Microlayer of the Ross Sea (Antarctica), and Emended Description of the Genus *Pseudoalteromonas*. Int. J. Syst. Evol. Microbiol. 66 (9), 3377-3382.
- Yoshihito, U., Aiko, H., Akira, Y., Junta, S., 1998. Reclassification of marine Agrobacterium species: Proposals of *Stappia stellulata* gen. nov., comb. nov., *Stappia aggregata* sp. nov., nom. rev., *Ruegeria atlantica* gen. nov., comb. nov., *Ruegeria gelatinovora* comb. nov., *Ruegeria algicola* comb. nov., and *Ahrensia kieliense* gen. nov., sp. nov., nom. rev. J. Gen. Appl. Microbiol. 44 (3), 201-210.

* Presenting author. E-mail address: doidboy@skku.edu (Jonghwan Kim)

S3.P191 Phytochemical study of five Polynesian plant extracts for skin care application

Marion Chambon^{a,*}, Raimana Ho^a, Beatrice Baghdikian^b, Gaëtan Herbette^c, Sok-Siya Bun-Llopet^b, Elnur Garayev^b, Émilie Thiébault^d, Laurent Duca^d, Remy Le Guével^e, Phila Raharivelomanana^a

^a UMR 241 SECOPOL, Université de Polynésie française, BP 6570, Faaa, Tahiti, French Polynesia

^b IMBE, Aix Marseille Univ, Avignon Univ, CNRS, IRD, Service of Pharmacognosie, Faculty of Pharmacy, Marseille, France

^c Aix Marseille Université, CNRS, Centrale Méditerranée, FSCM, Spectropole, Service 511, Campus Saint-Jérôme, Marseille, France

^d CNRS UMR 7369, Matrice Extracellulaire et Dynamique Cellulaire (MEDyC), Université de Reims Champagne Ardenne, Reims, France

^e Univ Rennes, CNRS, INSERM, BIOSIT UAR 3480, US_S 018, Impacell, Rennes, France

Five Polynesian plants, among the most traditionally used for skin care (pimples, wounds, burns, dermatoses) were studied and presented herein: leaves and nuts of *Calophyllum inophyllum*, flowers of *Gardenia taitensis*, rhizomes of *Curcuma longa*, leaves of *Cordia subcordata* and aerial roots of *Ficus prolixa* (Pétard, 2019). Eco-extraction was performed using ultrasound-assisted extraction (UAE) using ethanol:water (70:30). This method allowed reduced extraction time and lower energy consumption, within increased yields.

Plant extracts had already shown antioxidant properties evaluated by total phenolic content (TPC), free radical scavenging DPPH, and Ferric Reducing Antioxidant Power activity (FRAP) assays. Online HPLC-DPPH allowed the identification of phenolic bioactive compounds. *F. prolixa* extract was the most active and showed antioxidant intracellular activity on keratinocytes by AntiOxidant Power 1 (AOP1) assay (Chambon *et al.*, 2023).

Inflammatory properties were evaluated using NFκB transcription factor assay on THP1 cells and LOX assay on keratinocytes and fibroblasts. The first results indicated an anti-inflammatory activity of *C. subcordata* extract and a pro-inflammatory activity of *F. prolixa* extract.

The metabolomic analytical approach allowed the characterization of the chemical composition by using LC-MS² and molecular networking. Some compound classes were specific to a plant such as neoflavonoids, pyranocoumarins, and chromanones found in *C. inophyllum* or curcuminoids found in *C. longa*. Compounds annotated for *F. prolixa* and *C. subcordata* were described for the first time in these two indigenous Polynesian tree extracts.

These results highlight the potential of Polynesian plants from cosmetopeia and pharmacopeia, as a source of bioactive compounds for skin care applications.

Keywords: French Polynesia, cosmetopoeia, antioxidant, inflammatory, metabolomic

References

Chambon, M., Ho, R., Baghdikian, B., Herbette, G., Bun-Llopet, S. S., Garayev, E., Raharivelomanana, P., 2023. Identification of Antioxidant Metabolites from Five Plants (*Calophyllum inophyllum*, *Gardenia taitensis*, *Curcuma longa*, *Cordia subcordata*, *Ficus prolixa*) of the Polynesian Pharmacopoeia and Cosmetopoeia for Skin Care. Antioxidants. 12, 1870.

Petard, P., 2019. Quelques plantes utiles de Polynésie française : et ra'au tahiti. Editions Haere Po.

* Presenting author. E-mail address: marion.chambon@doctorant.upf.pf (Marion Chambon)

S3.P192 Eco-extraction and characterization by HPLC-HRMS/MS of *in vitro* culture of the carnivorous plant *Drosera*

Rym Attia^a, Alexander Castro Grijalba^a, David Da Silva^a, Laurine Garros^a, Christophe Hano^a, Arnaud Lanoue^b, Nathalie Giglioli-Guivarch^b, Emilie Destandau^{a,*}

^a UMR7311, Institut de Chimie Organique et Analytique, Université d'Orléans, CNRS, Orléans, France

^b EA2106 Biomolécules et Biotechnologies Végétales, Université de Tours, Tours, France

Drosera rotundifolia (round-leaved sundew) is an insectivorous plant from the *Droseraceae* family which is used since medieval times as a remedy for coughs and pulmonary diseases. *Drosera* extracts present antimicrobial, antifungal, and antitumor properties which are mainly attributed to two groups of phytochemical compounds characteristic of this genus: 1,4-naphthoquinones and flavonoids. Due to the substantial decline of *Drosera* habitat, the cultivation of this plant by *in vitro* propagation has been developed in order to enhance the amount of available plant resource. To be able to develop a new ingredient for cosmetic application from *in vitro* *Drosera*, green extraction methods and solvents were evaluated using microwave assisted extractions. Solvents generally recognized as safe, such as ethanol-water and propylene glycol-water mixtures and NaDES composed of lactic acid-glucose- water (LGH) were used to perform hybrid *Drosera* extracts (model). Extraction conditions were optimised using design of experiment and solvation calculation with CosmoRS approach were performed. CosmoRs, approach predicted a higher solubility of phenolic compounds in ethanol-water mixtures compared to the other tested solvents. Extraction optimization led to the similar conclusion obtaining a higher extraction yield using ethanol-water mixture with optimal conditions defined as EtOH % 62 %, extraction time 3 cycles of 15s and microwave power 130 W. Phytochemical compositions of obtained extracts were characterized using UHPLC-ESI-Q-TOF showing some slight differences in extraction selectivity, however the main compounds detected were myricetin hexose and quercetin hexose, dimethylelagic acid hexose and methyljuglone hexose. Other flavonoids and naphthoquinones derivatives were also identified in *Drosera* extracts.

Keywords: *in vitro* cultures, phytochemical study, eco-extraction optimization

Acknowledgments

The authors thank the Cosmetosciences program and “la Région Centre-Val de Loire” for founding (project INNOCOSM).

* Presenting author. E-mail address: emilie.destandau@univ-orleans.fr (Emilie Destandau)

S3.P193 Antioxidant potential and phytochemical analysis of an infusion obtained from the leaves of *Talisia esculenta*

Katia Castanho Scortecci^{a,b,c,*}, Maria Lúcia da Silva Cordeiro^{b,c}, Leticia Barbosa Santos^a, Verônica Giuliani de Queiroz Aquino Martins^{b,c}, Ariana Pereira da Silva^{b,c}, Wesley de Souza Paiva^d, Hasan Karimi^a, Ana Carolina Luchiari^e, Hugo Alexandre Oliveira Rocha^{c,e}, Fabio Boylan^a

^a School of Pharmacy and Pharmaceutical Sciences, Trinity Biomedical Sciences Institute, Trinity College Dublin, Dublin, Ireland

^b Laboratório de Transformação de Plantas e Análise em Microscopia (LTPAM), Departamento de Biologia Celular e Genética, Universidade Federal do Rio Grande do Norte (UFRN), Natal, RN, Brazil

^c Programa de Pós-graduação em Bioquímica e Biologia Molecular, Centro de Biociências, UFRN, Natal, RN, Brazil

^d Laboratório de Biotecnologia de Polímeros Naturais (BIOPOL), Departamento de Bioquímica, Universidade Federal do Rio Grande do Norte (UFRN), Natal, RN, Brazil

^e Departamento de Fisiologia e Comportamento, Universidade Federal do Rio Grande do Norte (UFRN), Natal, RN 59078-970, Brasil

Talisia esculenta (Sapindaceae) is a native plant in northeast of Brazil. Its fruits are consumed fresh and used in pastry making. Additionally, in folk medicine, the infusion of its leaves is utilised for treating high blood pressure, hip pain, and rheumatism, although scientific and toxicological information regarding its efficacy still needs to be provided. Previous research (Cordeiro et al., 2023) demonstrated that the infusion extract could protect 3T3 cell line against oxidative stress. This study evaluated the antioxidative potential of leaf infusion using zebrafish as an animal model. Zebrafish embryos were treated with 100 µg/mL of leaf infusion, and no mutagenic effects on embryo development were observed. Subsequently, these embryos were exposed to leaf infusion and an oxidative stressor to evaluate antioxidant protection. CuSO₄ was used as an oxidative stressor. Results indicated that *T. esculenta* infusion conferred protection against the harmful effects of oxidative stressors in this animal model. Phytochemical analysis of the leaves infusion revealed the presence of two major flavonoids. *T. esculenta* infusion was extracted with ethyl acetate and *n*-butanol and further submitted to separate its compounds using high-speed countercurrent chromatography (HSCCC). Quercitrin (quercetin- 3-O-rhamnose) was isolated in one run of the HSCCC analysis using Ethyl acetate:Butanol:Water (4.5:0.5:5). Rutin was further purified using Sephadex LH-20. These two flavonoids, which had their structures confirmed by Nuclear Magnetic Resonance (NMR), were the major flavonoids in the water infusion of *Talisia esculenta*. Further biological assays are necessary to determine whether the antioxidant potential is solely attributed to these two major flavonoids or if other unidentified compounds contribute to this effect.

Keywords: Pitomba, *Danio rerio*, cytotoxicity, HSCCC, NMR

References

Cordeiro, M.L.S., Aquino-Martins, V.G.Q., da Silva, A.P., Naliato, G.F.S., Silveira, E.R., Theodoro, R.C., da Santos, D.Y.A.C., Rocha, H.A.O., Scortecci, K.C., 2023. Exploring the Antioxidant Potential of *Talisia esculenta* using In Vitro and In Vivo Approaches. *Nutrients*. 2023 Sep 4;15(17):3855. doi: 10.3390/nu15173855.

Acknowledgments

Supported by: CNPq, CAPES PRINT.

* Presenting author. E-mail address: kacscort@yahoo.com (Kátia Scortecci)

S3.P194 Prospecting the phytochemical and antioxidant effects in vivo and in vitro of the ethanolic and aqueous extracts of the seeds of *Geoffroea decorticans* burkart (Fabaceae) on the animal model *Tenebrio molitor*

Ariana Pereira da Silva^{a,b}, Maria Lúcia da Silva Cordeiro^{a,b}, Verônica Giuliani de Queiroz Aquino Martins^{a,b}, Georggia Fatima Silva Naliato^{b,c}, Raquel Cordeiro Theodoro^{b,c}, Carlos Henrique Salvino Gadelha Meneses^d, Hugo Alexandre Oliveira Rocha^{b,e}, Kátia Castanho Scortecchi^{a,b,*}

^a Laboratório de Transformação de Plantas e Análise em Microscopia (LTPAM), Departamento de Biologia Celular e Genética, Universidade Federal do Rio Grande do Norte (UFRN), Natal, RN, Brazil

^b Programa de Pós-graduação em Bioquímica e Biologia Molecular, Centro de Biociências, UFRN, Natal, RN, Brazil

^c Instituto de Medicina Tropical, Universidade Federal do Rio Grande do Norte (UFRN), Natal 59077-080, Rio Grande do Norte, Brazil

^d Laboratório de Biotecnologia Vegetal (LBV), Departamento de Biologia, Centro de Ciências Biológicas e da Saúde, Universidade Estadual da Paraíba (UEPB) Campina Grande 58429500, Paraíba, Brazil

^e Laboratório de Biotecnologia de Polímeros Naturais (BIOPOL), Departamento de Bioquímica, Universidade Federal do Rio Grande do Norte (UFRN), Natal, RN, Brazil

Geoffroea decorticans, commonly known as Chañar, is a plant indigenous to Chile (Fabaceae). Various parts of the plant, including its bark, flowers, fruit, and leaves, are utilized in folk medicine for their expectorant activity, particularly in treating bronchopulmonary disorders and pain relief. Additionally, they exhibit antioxidant and antinociceptive activities (Costamagna, 2013, Isla et al., 2021;). However, there are no studies reporting such activities with Chañar seed extracts, which are used as food. This study aimed to evaluate the ethanolic (EE) and aqueous (EA) extracts of Chañar seeds. Phytochemical analysis reveals the presence of phenolic compounds and flavonoids in both extracts. High-Performance Liquid Chromatography (HPLC) coupled with Gas Chromatography-Mass Spectrometry/Mass Spectrometry (GC-MS/MS) identified key components such as phytol, alpha-tocopherol, vitexin, and rutin. Biochemical assays assessing total antioxidant capacity (TAC), reducing power, 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging, and copper and iron chelation demonstrated excellent antioxidant potential for both extracts. *Tenebrio molitor* larvae were utilized as an animal model to evaluate the toxicity of the EE extract and its antioxidant potential against the oxidative stressor agent CuSO₄. No toxicity was observed for the EE extract at concentration of 100 µg/mL. Moreover, the EE exhibited the ability to protect the larvae from the oxidative stress as evidenced by survival frequency over fifteen days and the melanisation rate, indicative of oxidative stress and inflammation. These findings suggest that the EE and EA extracts of Chañar seeds could serve as a potential source of bioactive compounds with antioxidant properties.

Keywords: chañar, extracts, HPLC, copper sulphate, *Tenebrio molitor*

References

- Costamagna, M. S.; Ordonez, R.M.; Zampini, I.C.; Sayago, J.E. & Isla, M.I. Nutritional and antioxidant properties of *Geoffroea decorticans*, an Argentinean fruit, and derived products (flour, arrope, decoction and hydroalcoholic beverage), Food Research International, 2013.
- Isla, M.I.; Ezquer, M.E.; Leal, M.; Moreno, M.A.; Zampini, I.C. Flower beverages of native medicinal plants from Argentina (*Acacia caven*, *Geoffroea decorticans* and *Larrea divaricata*) as antioxidant and anti-inflammatory. J Ethnopharmacol., v.281, p.114490, 2021.

Acknowledgments

Supported by CNPq, CAPES.

* Presenting author. E-mail address: kacscort@yahoo.com (Kátia Scortecchi)

S3.P195 Standardisation of secondary metabolite extraction and characterisation of *Citrus aurantium* var. *bergamia* leaves using green extraction technology

Beatrice Conversazioni^{a,*}, Chiara Spaggiari^a, Marika Allodi^a, Gabriele Costantino^a, Sonia Vaccaro^b, Beatrice De Carne^b, Paolo Goi^b

^a University of Parma, Department of Food and Drugs, Parco Area delle Scienze 27/A, 43124 Parma, Italy

^b Davines Group - Botanical Research Department, Via Don Angelo Calzolari 55/A, 43126 Parma, Italy

This study is aimed at optimizing a natural extract of *Citrus aurantium* var. *bergamia* leaves, with regard to the phytochemicals concentration in parallel with the stability of its macroscopic features, considering its final use in a range of cosmetic formulations.

The cosmetic industry increasingly seeks active compounds of natural origin, favoring more and more a sustainable production cycle. Plants rich in secondary metabolites such as *Citrus aurantium* var. *bergamia* (bergamot) offer promising biological activities in cosmetics. Specifically, bergamot leaves, considered a byproduct in the business of bergamot essential oil, have been proved to be a noteworthy source of flavonoids, including neoeriocitrin and naringenin (Baron et al., 2021), compounds associated with anti-inflammatory and anti-aging activities (Denaro et al., 2021; Lim and Kim, 2018). In our work we focused on a green extraction technology, specifically the Naviglio extractor*, which utilizes a rapid solid-liquid dynamic extraction process (Naviglio, 2003), allowing to maximize active compounds yield using environmentally friendly solvents under low temperature conditions.

The research assessed various parameters of the Naviglio extractor* (e.g. extraction time and number of cycles) and the composition of the extractive mixture composition to optimize the extract in terms of stability of qualitative and quantitative characteristics. The solvents mixture resulting as most performing consists of water/glycerin/butylene glycol in relative concentrations of 70/10/20 (%w/w) added to dried biomass in a ratio of 1/20 (biomass/solvents w/w). This optimized extraction method has shown to provide a stable extract for potential use in skin care cosmetics.

Keywords: natural extract, green extraction, *Citrus x bergamia*, cosmetic ingredients, phytochemicals

References

- Baron, G., Altomare, A., Mol, M., Garcia, J.L., Correa, C., Raucci, A., Mancinelli, L., Mazzotta, S., Fumagalli, L., Trunfio, G., et al., 2021. Analytical Profile and Antioxidant and Anti-Inflammatory Activities of the Enriched Polyphenol Fractions Isolated from Bergamot Fruit and Leave. *Antioxidants*, 10, 141.
- Denaro, M., Smeriglio, A., Trombetta, D., 2021. Antioxidant and Anti-Inflammatory Activity of Citrus Flavanones Mix and Its Stability after In Vitro Simulated Digestion. *Antioxidants*, 10, 140.
- Lim, K.H., Kim, G.R., 2018. Inhibitory effect of naringenin on LPS-induced skin senescence by SIRT1 regulation in HDFs. *Biomedical Dermatology*, 2, 26.
- Naviglio, D., 2003. Naviglio's Principle and Presentation of an Innovative Solid-Liquid Extraction Technology: Extractor Naviglio*. *Analytical Letters*, 36:8, 1647-1659.

* Presenting author. E-mail address: beatrice.conversazioni@unipr.it (Beatrice Conversazioni)

S3.P196 Sustainable isolation and improved structural characterization of mycosporin-like amino acids: natural UV-filters from red algae

Michel Thomas^{a,*}, Rieusset Laura^b, Simmler Charlotte^c, Stéphane Greff^c, Romain Bouhlo^a, Kévin Hardouin^b, Nathalie Bourgougnon^b, Grégoire Audo^a, Gilles Bedoux^b

^a Gilson Purification SAS, 22 rue Bourseul, 56890 Saint Avé, France

^b Laboratoire de Biotechnologie et Chimie Marine, Université Bretagne Sud, CNRS - EMR 6076, France

^c Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, UMR 7263, CNRS, IRD, Aix Marseille Université, Avignon Université, Station Marine d'Endoume, France

Mycosporin-like amino acids (MAAs) are low-molecular-weight, colorless and water-soluble compounds produced by a wide variety of marine organisms including corals, fungi, algae and cyanobacteria (Geraldès and Pinto, 2021). They are recognized as a potential source of natural sunscreens due to their capacity to absorb a very large proportion of UV rays. MAAs are also characterized by other biological activities such as antioxidant, anti-aging and anti-inflammatory properties (Rosic, 2021) (Fonsesca et al., 2023). MAAs are thus of growing interest for their capacity to replace controversial photoprotective agents in sunscreen products. Nevertheless, their enrichment or isolation remain a challenge in cosmetic industry.

This presentation will cover the development of a sustainable purification strategy of MAAs that can be easily transposed to an industrial scale. In this context, centrifugal partition chromatography (CPC) using aqueous two-phase system was particularly adapted to isolate MAAs (Michel et al. 2023). The developed method allowed to obtain pure compounds and enriched fraction from a *Porphyra* sp. crude extract. Structure elucidation employing extensive NMR spectroscopy and HRMS allowed the identification of MAAs such as Porphyra-334 (**1**, Fig. 1).




Fig. 1. Structure of porphyra-334 **1**

This work provides a new approach, easily scalable, to produce enriched cosmetical ingredients or to purify MAAs that can be used in diverse formulation to enhance the efficacy of natural sunscreens.

Keywords: *Porphyra* sp, mycosporin-like amino acids, isolation, CPC, UV-filters

References

- Geraldès, V., Pinto, E., 2021. Mycosporine-like amino acids (MAAs): Biology, chemistry and identification features. *Pharmaceuticals*, 14, 63.
- Rosic, N.N., 2021. Recent advances in the discovery of novel marine natural products and mycosporine-like amino acid UV-absorbing compounds. *Appl Microbiol Biotechnol.* 105, 7053–7067.
- Fonseca, S., Amaral, M.N., Reis, C.P., Custódio, L., 2023. Marine natural products as innovative cosmetic ingredients. *Mar. Drugs*. 21, 170.
- Michel, T., Loyet C., Boulho, R., Eveno, M., Audo, G., 2023. Investigation of alternative two-phase solvent systems for purification of natural products by centrifugal partition chromatography. *Phytochem. Anal.* (Online ahead of print).

* Presenting author. E-mail address: tmichel@gilson.com (Thomas Michel)

S3.P197 Cosmeceutical stability assessment of bergamot: behavioural studies on its potential as an active ingredient

Sonia Vaccaro^a, Beatrice De Carne^a, Paolo Goi^a, Beatrice Conversazioni^{b*}, Chiara Spaggiari^b, Gabriele Costantino^b

^a Davines Group - Botanical Research Department, Via Don Angelo Calzolari 55/A 43126 Parma, Italy

^b University of Parma, Department of Food and Drugs, Parco Area delle Scienze, 27/A, 43124 Parma, Italy

This study displays the stability investigation concerning a natural extract from *Citrus aurantium* var. *bergamia* (bergamot) leaves, focusing on assessing cosmeceutical properties, particularly their stability post-heating.

Our research is oriented towards the development of innovative cosmeceutical ingredients sourced from botanical origins, which include bioactive compounds promoting cosmetic functionalities. Notably, studies on flavonoid fractions from bergamot reveal significantly higher anti-inflammatory activity of flavanones mix compared to single molecules, highlighting a synergistic mechanism of this bergamot fraction (Denaro et al., 2021). Implementing natural extracts as active ingredients also requires consideration of the cosmetic formulation process, such as heating treatments. Given the heat sensitivity of flavonoids, they can be subject to degradation, resulting in a change of the final properties (Chaaban et al., 2017).

Therefore, a stability study of bergamot extract was conducted, monitoring phytochemical concentrations post-heating at 35°C, 40°C and 45°C. The compounds analysed were flavanones, i.e. neoeriocitrin, naringenin and hesperidin, and a flavonol, i.e. quercetin. They have been quantified at time 0, post-thermal process, and after 1 and 2 months of storage at room temperature and 45°C. Quercetin resulted as the most heat-sensitive molecule, showing since time 0 the most significant decline, with 19 % content loss when the extract is heated at 45°C. The degradation of this flavonol got worse over time, especially when stored at 45°C, reaching a 47% reduction by the second month. Based on these findings, it was proposed an acceptability threshold of 40°C for heating the extract during cosmetic formulation.

Keywords: bergamot, flavonoids, thermal stability, cosmeceutical ingredient, quercetin

References

- Chaaban, H., Ioannou, I., Chebil, L., et al., 2017. Effect of heat processing on thermal stability and antioxidant activity of six flavonoids. *J Food Process Preserv.* 41:e13203.
- Denaro, M., Smeriglio, A., Trombetta, D., 2021. Antioxidant and Anti-Inflammatory Activity of Citrus Flavanones Mix and Its Stability after In Vitro Simulated Digestion. *Antioxidants*, 10, 140.

* Presenting author. E-mail address: beatrice.conversazioni@unipr.it (Beatrice Conversazioni)

S3.P198 *Erythrina* spp. as natural anti-inflammatory ingredients for cosmeceuticals: an *in-silico* study

Natalia Blicharska^{a,*}, Dino Rotondo^a, Simon Jackson^b, Paolo Goi^b, Veronique Seidel^a

^a Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, 161 Cathedral Street, Glasgow UK

^b Davines Group - Botanical Research Department, Via Don Angelo Calzolari 55/A, 43126 Parma, Italy

Many inflammatory skin conditions depend upon activation of the nuclear factor NF- κ B pathway which triggers the transcription of genes coding for pro-inflammatory mediators (Bell *et al.*, 2003). Considering the rising demand for natural cosmeceuticals, developing products containing anti-inflammatory botanical ingredients capable of modulating the NF- κ B pathway is worth investigating. *Erythrina* spp. are used traditionally to treat skin ailments and alleviate inflammation. Their extracts and phytochemicals have demonstrated anti-inflammatory activity (Jiménez-Cabrera *et al.*, 2020; Khumalo *et al.*, 2021; Susilawati *et al.*, 2023). These anti-inflammatory phytochemicals include flavonoids, a class of compounds already known to target the NF- κ B pathway (Kadioglu *et al.*, 2015). In this study, an *in-silico* molecular docking approach using AutoDock Vina was employed to predict the binding affinities and ligand efficiencies of 18 flavonoid derivatives, from *Erythrina* spp. with anti-inflammatory activity, against three key proteins of the NF- κ B pathway. These included the NF- κ B p50/p65 heterodimer, the I κ B kinase subunit β (IKK β) and the NEMO/IKK association domain protein.

The isoflavonoids alpinumisoflavone (**1**) and wighteone (**2**) showed the best ligand efficiencies towards the NF- κ B p50/p65 complex, while the pterocarpans erybraedin A (**3**) and phaseollidin (**4**) showed the strongest ligand efficiencies towards IKK β and NEMO/IKK, respectively. To the best of our knowledge, this is the first report on the interactions of *Erythrina* flavonoids with proteins of the NF- κ B pathway.

Further *in-vitro* studies are warranted to validate the effects of these compounds on the NF- κ B pathway and confirm the potential of *Erythrina* spp. as natural anti-inflammatory ingredients for cosmeceuticals.

Keywords: *Erythrina*, anti-inflammatory, NF- κ B, *in-silico*, molecular docking

References

- Bell, S., Degitz, K., Quirling, M., Jilg, N., Page, S., Brand, K., 2003. Involvement of NF-kappaB signalling in skin physiology and disease. *Cell Signal.* 15, 1-7
- Kadioglu, O., Nass, J., Saeed, M.E., Schuler, B., Efferth, T., 2015. Kaempferol is an anti-inflammatory compound with activity towards NF- κ B pathway proteins. *Anticancer Res.* 35, 2645-2650
- Jiménez-Cabrera, T., Bautista, M., Velázquez-González, C., *et al.*, 2020. Promising antioxidant activity of *Erythrina* genus: An alternative treatment for inflammatory pain? *Int. J. Mol Sci.* 22, 1-19
- Khumalo, G.P., Sadgrove, N.J., Van Vuuren, S.F., Van Wyk, B.E., 2021. South Africa's best bark medicines prescribed at the Johannesburg Muthi markets for skin, gut and lung infections: MIC's and brine shrimp lethality. *Antibiotics.* 10, 1-27
- Susilawati, E., Levita, J., Susilawati, Y., Sumiwi, S.A., 2023. Pharmacology activity, toxicity and clinical trials of *Erythrina* genus plants (Fabaceae): an evidence-based review. *Front. Pharmacol.* 14, 1-9

* Presenting author. E-mail address: natalia.blicharska@strath.ac.uk (Natalia Blicharska)

S3.P199 Residual solvents in botanical extracts used in food supplements: a regulatory conundrum for scientific justification

Deepak Mundkinajeddu*, Rojison Koshy, Nithin J., Aboli Girme, Amit Agarwal

Research and Development Division, Natural Remedies Pvt Ltd, Bangalore, India

Botanical extracts are widely used in food supplements across the world for a variety of health benefits. Commercially the extracts are prepared typically using water and ethanol as solvents. Since the supplements fall under the food law, use of other solvents like methanol are generally not allowed and encouraged as extraction solvents. Accordingly, stringent limits (less than 10 ppm as per EU legislation) are set to ensure absence of solvents like methanol in botanical extracts used in food supplements.

On the other hand, we analyzed using headspace gas chromatography (GC) and observed the presence of many solvents like methanol and acetone in botanical extracts, even when these solvents are not used for extraction. Many crude herb powders (before extraction) as such were found to contain solvents like methanol and acetone. Particularly content of methanol was found to be higher and ranged from 25 to 1000 ppm. In addition, we prepared aqueous extracts of the plants under controlled condition in the laboratory and again found the presence of methanol and acetone in the extracts. Results of our findings will be covered in the presentation.

These results reflect the commercial challenges for regulatory compliance of botanical extracts under food supplements. Scientific explanations for the presence of solvents include naturally occurring solvents, artefacts generated during extraction and processing and as part of sample preparation in headspace for GC analysis (Deepak 2014, Uematsu 2002). These aspects will be discussed during the presentation.

Keywords: food supplements, botanical extracts, residual solvents, methanol, acetone

References

- Mundkinajeddu D., Agarwal A., 2014. Residual methanol in botanical dietary ingredients – perspectives of a manufacturer. *Herbalgram*. 11, 1-4.
- Uematsu Y., Suzuki K., Iida K., Hirata K., Ueta T., Kamata K., 2002. Determination of low levels of methanol and ethanol in licorice extract by large volume injection head-space GC. *J. Food Hyg. Soc. Japan*. 43, 295-300.

* Presenting author. E-mail address: deepak@naturalremedy.com (Deepak Mundkinajeddu)

S3.P200 *Grateloupia lanceolata* promotes hair growth by inhibiting 5 α -reductase activity and opening KATP channels

Da Hee Yang^{a,b}, Jung-Il Kang^{a,b}, Youn Kyung Choi^{a,b}, Eun-Sook Yoo^{a,b}, Hee-Kyoung Kang^{a,b,*}

^a Department of Medicine, College of Medicine, Jeju National University, Jeju 63243, South Korea

^b Jeju Research Center for Natural Medicine, Jeju National University, Jeju 63243, South Korea

The marine environment is rich in biologically active natural products, and the diverse biological activities of marine algae are known. This study was conducted to evaluate the effects of *Grateloupia lanceolata*, one of red algae, on hair growth. When examined the activity of 5 α -reductase, which converts testosterone to dihydrotestosterone, a main cause of androgenetic alopecia (Kaufman, 1998). *Grateloupia lanceolata* extract significantly inhibited the activity of 5 α -reductase. Minoxidil, a hair- growing drug, promotes the proliferation of NIH3T3 fibroblast via the opening of KATP channels (Sanders, 1996). The *G. lanceolata* extract significantly increased the proliferation of NIH3T3 fibroblasts. When isolated rat vibrissa hair follicles were treated with the *G. lanceolata* extract for 21 days, the *G. lanceolata* extract significantly increased the hair fiber length of vibrissa hair follicles. These results suggest that *G. lanceolata* promotes hair growth through the inhibition of 5 α -reductase activity and the opening of KATP channels.

Keywords: *Grateloupia lanceolata*, hair growth, 5 α -reductase, KATP channels

References

- Kaufman, K. D., Olsen, E. A., Whiting, D., Savin, R., DeVillez, R., Bergfeld, W., Price, V. H., Van Neste, D., Roberts, J. L., Hordinsky, M., Shapiro, J., Binkowitz, B., & Gormley, G. J. (1998). Finasteride in the treatment of men with androgenetic alopecia. Finasteride Male Pattern Hair Loss Study Group. *Journal of the American Academy of Dermatology*, 39(4 Pt 1), 578–589.
- Sanders, D. A., Fiddes, I., Thompson, D. M., Philpott, M. P., Westgate, G. E., & Kealey, T. (1996). In the absence of streptomycin, minoxidil potentiates the mitogenic effects of fetal calf serum, insulin-like growth factor 1, and platelet- derived growth factor on NIH 3T3 fibroblasts in a K⁺ channel-dependent fashion. *The Journal of investigative dermatology*, 107(2), 229–234.

Acknowledgments

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (RS-2023-00270936) and by the Korea government (MSIT) (2023R1A2C1007234)]

* Presenting author. E-mail address: pharmkhk@jejunu.ac.kr (Hee-Kyoung Kang)

S3.P201 Anti-inflammatory effect of *Lysimachia mauritiana* extract on particulate matter 10 Plus Diesel Exhaust Particles (PM10D)-induced respiratory disease mouse model and standardization using UPLC-qTof/MS analysis

Yoon-Young Sung^{a,*}, Heung Joo Yuk^a, Seung-Hyung Kim^b, Chang-Hwa Song^c, Dong-Seon Kim^a

^a KM Research Science Division, Korea Institute of Oriental Medicine, Daejeon 34054, Republic of Korea

^b Institute of Traditional Medicine and Bioscience, Daejeon University, Daejeon 34520, Republic of Korea

^c Department of microbiology College of Medicine Chungnam National University, 266 Munhwa-ro, Jung-gu, Daejeon 35015, Republic of Korea

Exposure to particulate matter (PM) causes considerable respiration-related health risks. *Lysimachia mauritiana* (LM) (Primulaceae) is known as a medicinal plant with anti-tumor, anti-viral, and anti-inflammatory effects. It is native to eastern Asia, where it is widespread in coastal areas. The effect of LM extract on airway inflammation was investigated in mice exposed to a fine dust mixture (PM10D) of PM10 (PM diameter < 10 μ m) and diesel exhaust particles (DEP). LM suppressed the number of immune cells and secretion of cytokines in bronchoalveolar lavage fluid (BALF) and lung of PM10D mice. Airway inflammation and collagen fibrosis in the lung after PM10D exposure were investigated by histopathological analysis, and LM ameliorated these symptoms. LM decreased the mRNA expression of MUC5AC, CXCL1, TNF- α , MIP-2, TRPA1, and TRPV1 in lung. The main candidate components from LM that exhibited anti-inflammatory effects were investigated by ultra-performance liquid chromatography quadrupole time of flight mass spectrometry (UPLC-qTof/MS). The seven phytochemicals were tentatively identified by MS analysis. Compound 3 among the flavonoids, which are easy to detect even under UV, is identified into biorobin (keampferol 3-O-robinobioside) and was confirmed as candidate of standardization. These findings suggest that LM may be a potential nutraceutical for the treatment of respiratory diseases in human.

Keywords: airway, fine dust, Lung, *Lysimachia mauritiana*

References

Lactobacillus paracasei ATG-E1 improves particulate matter 10 plus diesel exhaust particles (PM10D)-induced airway inflammation by regulating immune responses. Lee YS, et al. Front Microbiol. 2023;14:1145546.

Acknowledgments

This study was supported by the project (KSN23214313) from the KIOM of Korea.

*Presenting author. E-mail address: yysung@kiom.re.kr (Yoon-Young Sung)

S3.P202 Analysis of dietary supplements containing adaptogenic herb *Astragalus trimestris*

Kateřina Knížková*, Martin Kuchař, Petra Cihlářová

Forensic Laboratory of Biologically Active Substances, Department of Chemistry of Natural Compounds, University of Chemistry and Technology, Technická 5, 160 00 Prague, Czech Republic

Astragalus trimestris L. is an adaptogenic herb deeply rooted in traditional Chinese medicine. According to the presence of many bioactive substances, including compounds like saponins, flavonoids and alkaloids, the herb has significant beneficial effects on human health. *A. trimestris* is used for its anxiolytic, anti-inflammatory, antiviral, antimicrobial and immunomodulatory effects and also in the treatment of various diseases. The key bioactive compound unique to *A. trimestris*, cycloastragenol, has demonstrated the ability to enhance telomerase activity, which may contribute to promoting longevity. On the market is *A. trimestris* available for tea preparations, offered in the form of chopped or ground roots, as tinctures or encapsulated formulations.

This study aimed to develop novel analytical methods for quantification of selected bioactive compounds in dietary supplements containing *A. trimestris* using high-performance liquid chromatography coupled with triple quadrupole spectrometry (HPLC-MS/MS). For this purpose, a wide spectrum of dietary supplements was purchased on the Czech market. Regarding the optimisation experiments for each group of bioactive substances (polyphenols, saponins and steroids), different extraction solvents were selected. Various columns and mobile phase compositions were investigated, and ion source parameters were optimised.

Using the developed methods, selected bioactive compounds in dietary supplements containing *A. trimestris* were successfully quantified by HPLC-MS/MS analysis. Apart from the characteristic glucosides of cycloastragenol, supplements contained notable amounts of phytosteroids (β -sitosterol), isoflavones (calycosin-7-O- β -D-glucoside) or phenolic acids (salicylic acid). Due to the dimensionality of the data, multivariate analysis was essential for interpretation. Factor analysis revealed distinct clusters of different product types in the score plot.

Keywords: *Astragalus trimestris*, HPLC-MS/MS, cycloastragenol, adaptogen, longevity

* Presenting author. E-mail address: knizkovk@vscht.cz (Kateřina Knížková)

S3.P203 The GreenValue project: exploring salt tolerant plants as sources of innovative food additives

Célia Quintas^a, Isabel Ratão^a, Catarina G. Pereira^b, Maria J. Rodrigues^b, Luísa Custódio^{b,*}

^a Department of Food Engineering of the Algarve University (DFE), Mediterranean Institute for Agriculture, Environment and Development (MED), scientific hub of the University of Algarve

^b Centre of Marine Sciences (CCMAR), University of Algarve, Campus de Gambelas, 8005 139 Faro, Portugal

Food additives help extend the shelf-life and distribution of food products, but they are linked to health issues like allergies and cancer, causing growing consumer skepticism. There's a trend towards healthier options, like functional foods that offer health benefits beyond basic nutrition. This presentation will cover some results obtained during the GreenValue R&D project, that explored biomass from selected species of salt tolerant plants (halophytes) as sources of innovative food additives in bakery and dairy food products. For example, while natural extracts from fruit peels from the invasive species *Carpobrotus edulis* were used to functionalize yogurts, biomass from the aromatic halophytes *Chrithmum maritimum* and *Helychrisum picardii* were used to fortify soft goat cheese. In another approach, biomass from cultivated *Arthrocnemum macrostachyum* was used to replace salt during the preparation of soft goat cheese. Greenvalue results open new avenues on the use of salt tolerant plants, recognized for their ecological role in the rehabilitation of saline soils and as sources of nutrients and bioactive products, as innovative and sustainable food ingredients.

Funding: Work funded by Foundation for Science and Technology (FCT) through project UIDB/04326/2020, UIDP/04326/2020, LA/P/0101/2020 and PTDC/BAA-AGR/1391/2020, and the operational programs CRESC Algarve 2020 and COMPETE 2020 through project EMBRC.PTALG- 01-0145-FEDER-022121. FCT supported L.C. by the FCT Scientific Employment Stimulus (CEECIND/00425/2017) and M.J.R. by the FCT program contract (UIDP/04326/2020).

Keywords: food additives, functional ingredients, halophytes

* Presenting author. E-mail address: lcustodio@ualg.pt (Luísa Custódio)

S3.P204 Seasonal variation of the indole alkaloid profile of *Vinca minor* determined by GC-MS

Dominik Skalka, Jandirk Sendker, Verena Spiegler*

Institute of Pharmaceutical Biology and Phytochemistry, University of Münster, Corrensstraße 48, 48149 Münster, Germany

Seasonal variations in content and composition have frequently been reported for plant secondary metabolites, including monoterpene indole alkaloids (MIA) (Osman et al., 2019). This also applies to the content of vincamine (Girre et al., 1978), the major MIA of *Vinca minor* L., a subshrub from the family of Apocynaceae. However, less is known about variations of other MIA from this species. In this study, we employed GC-MS to investigate the alkaloid profile of *V. minor* harvested at different time points during the growing season.

Six replicates per sample were prepared by acid-base extraction of the aerial parts of the plant to obtain fractions enriched in alkaloids. Although one limitation of GC is the requirement for volatile substances, the MIAs passed the GC column showing chromatographic separation and acceptable peak shapes. Moreover, their characteristic EI fragmentation patterns facilitated compound identification. To assess the variation in the alkaloid profile, the relative proportion of the peak areas was determined for the major alkaloids.

Generally, the MIA composition varied significantly across the samples from spring to fall. However, major changes were observed for few compounds only, whereas the ratio of most alkaloids remained unaffected by the season. With respect to the analytical method, GC-MS performed similarly well as LC-MS in the detection of MIA from *V. minor* within the current study.

Keywords: *Vinca minor*, Apocynaceae, indole alkaloids, GC-MS

References

- Girre, L., Cillard, P., Cillard, J., Amoros, M., Durand, D., 1978. Changes in the vincamine content of the leaves of lesser periwinkle (*Vinca minor*) during growth. *Plant. Med. Phytother.* 12, 5–23.
- Osman, A.G., Haider, S., Chittiboyina, A.G., Khan, I.A., 2019. Utility of alkaloids as chemical and biomarkers for quality, efficacy, and safety assessment of botanical ingredients. *Phytomedicine* 54, 347–356.

* Presenting author. E-mail address: verena.spiegler@uni-muenster.de (Verena Spiegler)

S3.P205 Review of honeybee-sourced, ingested natural products and their health uses

Barbara C. Sorkin^{a,*}, Margaret A. Nagai-Singer^a, Johanna Dwyer^a, Alessandra Durazzo^b, Massimo Lucarini^b, Michael Heinrich^{c,d}, Gisela Butera^e

^a Office of Dietary Supplements, U.S. National Institutes of Health, 6705 Rockledge Drive, MSC 7991, Bethesda, MD, 20892, USA

^b CREA-Research Centre for Food and Nutrition, Via Ardeatina 546, 00178 Rome, Italy

^c UCL School of Pharmacy, Pharmacognosy and Phytotherapy, London, UK

^d Department of Pharmaceutical Sciences and Chinese Medicine Resources, Chinese Medicine Research Centre, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^e Division of Library Services, U.S. National Institutes of Health, Bethesda, MD, 20892-1150, USA

Honey bee-sourced products (HBP) such as bee pollen, honey, propolis and royal jelly are chemically complex natural products, containing varied combinations of probiotics, proteins, saccharides, polyphenols, lipids, minerals, and vitamins. The composition of HBP varies with region and season of harvest, among other variables. For over 10,000 years humans have used honey for flavor and HBP for diverse, purported health benefits including improving energy, enhancing fertility, sharpening cognition, alleviating depression, and for anti-cancer and anti-viral effects. To assess the range of HBP and associated clinical outcomes reported we used the Covidence screening software in retrieving over 1000 published clinical trials of ingested HBP. from PubMed and EMBASE. Propolis and royal jelly were each tested for more than a dozen different outcomes, and honey mostly to improve growth or alleviate inflammation. In addition to those above, outcomes tested for propolis and royal jelly included alleviation of non-alcoholic fatty liver disease and improvement of glucose homeostasis. Since most retrieved publications reported positive associations between specific HBP and health outcomes new research on the specific chemistry and targets involved may be helpful. However, most of the published trials had small sample sizes and lacked other indicators of research rigor. Researchers should assess the quality of publications reporting putative HBP efficacy before investing substantial effort in generating the mechanistic data required for meaningful product standardization and optimal trial design (Sorkin, et al., 2020; Heinrich, et al., 2022).

Keywords: honeybee product constituents, biological activities

References

- Sorkin, B.C.,Kuszak, A.J., Bloss, G., Fukagawa N.K., Hoffman F.A., Jafari, M., Barrett, B., Brown, P.L., et al. 2020. Improving natural product research translation: From source to clinical trial. *FASEB J.* 34, 41-65..
- Heinrich, M.,et al. 2022. Best Practice in the chemical characterisation of extracts used in pharmacological and toxicological research – The ConPhyMP Guidelines 12. *Front. Pharmacol.* 13.

* Presenting author. E-mail address: sorkinb@nih.gov (Barbara C. Sorkin)

S3.P206 Evaluating the analytical profile of ten hydrophilic *Cucurbita pepo* seed extracts

Adina-Elena Grasu^{a,*}, Roman Senn^b, Christiane Halbsguth^b, Georg Boonen^b, Veronika Butterweck^b, Alexander Schenk^b, Anca Miron^a

^a Grigore T. Popa University of Medicine and Pharmacy Iasi, Universitatii Str. 16, 700115, Iasi, Romania

^b Max Zeller & Söhne AG, Seeblickstrasse 4, 8590 Romanshorn, Switzerland

The *Cucurbita pepo* L. species exhibits a remarkable phenotypic diversity (Christenhusz et al., 2016) with various fruit morphologies, ranging from spherical to elongated and flattened shapes (Schaffer et al., 2016). The characterization of lipophilic constituents in seeds of *C. pepo* was comprehensively explored (Dotto et al., 2020), while the understanding of the hydrophilic metabolites remains comparatively underexplored. Therefore, this study generated ten hydrophilic extracts from a total of ten varieties of *C. pepo* seeds all sourced from Europe, ensuring that the collected specimens reflect the observed diversity within the fruits. Oil free, hydroethanolic extracts were obtained from hulled, hull-less, and zucchini seeds. Identification and quantification of free amino acids, sugars and lignans within hydrophilic extracts was achieved aiming to compare their composition of secondary metabolites using UHPLC-DAD-MS or HPLC-RI methods. Notably, the amount of free amino acid varied across the species. The Snow-White variety contained the highest content of free amino acids (~6.45%) and highest glutamine amount (~2.8%), while zucchini varieties were rich in cucurbitine, a genus specific non-proteinogenic amino acid (Mihranian et al., 1968). The presence of saccharides like glucose, fructose and the raffinose family of oligosaccharides was also determined, identifying sucrose in all varieties as the predominant sugar in the range of 19.3% - 28.7%. Lignan identification showed a high diversity in seeds with husks, while their absence in extracts of hullless seeds was observed. Our data of the evaluation of hydrophilic extracts underline the compositional variability of *C. pepo* seeds obtained from different sources in Europe.

Keywords: *Cucurbita pepo*, amino acids, cucurbitine, sucrose, lignans

References

- Christenhusz, M. J. M., Byng, J. W. (2016). The number of known plants species in the world and its annual increase. *Phytotaxa*. 261 (3): 201–217.
- Dotto, J.M., Chacha, J.S., 2020. The potential of pumpkin seeds as a functional food ingredient: A review. *Scientific African*, Volume 10, article e00575.
- Mihranian, V.H., & Abou-Chaar, C.I. (1968). Extraction, detection, and estimation of cucurbitin in *Cucurbita* seeds. Schaffer, A.A., Paris, H.S., 2016. Cucurbitaceae Melons, Squash, Cucumber. *Encyclopedia of Applied Plant Sciences*. Volume 2, 209-217.

* Presenting author. E-mail address: adina-elena.grasu@d.umfiasi.ro (Adina-Elena Grasu)

S3.P207 Uncovering the potential of cyanobacterial pheophorbide derivatives for metabolic disorders

Natália Gonçalves Silva^{a,b,*}, Vitor Vasconcelos^{a,b}, Siegfried Ussar^{c,d}, Ralph Urbatzka^a

^a Interdisciplinary Centre of Marine and Environmental Research (CIIMAR/CIMAR), University of Porto, Avenida General Norton de Matos, s/n, 4450-208 Matosinhos, Portugal

^b FCUP, Faculty of Science, Department of Biology, University of Porto, Rua do Campo Alegre, 4169-007 Porto, Portugal

^c Research Unit Adipocytes & Metabolism, Helmholtz Diabetes Center, Helmholtz Munich, 85764 Neuherberg, Germany

^d German Center for Diabetes Research (DZD), 85764 Neuherberg, Germany

The research on natural products is driven to discover new therapeutic solutions for diseases. Here, we aimed to discover novel compounds with lipid lowering activity. Bioactivity screening led to the identification of *Cyanobium* sp. LEGE 06097 as a promising strain (Costa et al., 2019), and bioassay-guided fractionation led to the isolation of 15^l-hydroxy-lactone pheophorbide a (**1**). The structure was confirmed based on 1D- and 2D-NMR spectroscopy, together with LC-HR-ESI-MS/MS analysis (Kamarulzaman et al., 2011). For the first time, (**1**) was shown to reduce neutral lipids in zebrafish larvae with an EC₅₀ value of 1.9 µM (1.2 µg/mL) after 48 h exposure, while the structurally related pheophorbide a (**2**) had an EC₅₀ value of 2.7 µM (1.6 µg/mL). Preliminary studies revealed insights into the modes of action for both compounds. Isoproterenol-induced lipolysis in primary subcutaneous adipocytes isolated from C57BL/6J mice was assessed together with the expression of the transcriptional factors EGR1, FOXO1 and SP1 by semi-quantitative real-time qPCR. Compound (**2**) significantly inhibited lipolysis by 61.7% and 81.6% and down-regulated the expression of all three genes at 2.7 µM and 27 µM (16 µg/mL), respectively. Mitochondrial respiration was measured in HepG2 and differentiated brown adipocytes for both compounds using a Seahorse Extracellular Flux Analyzer. Both compounds reduced FCCP induced maximal respiration in brown adipocytes at both tested concentrations, and at the highest concentration in HepG2 cells. In conclusion, pheophorbide derivatives have promising effects on lipid metabolism, which may be the first step to develop them into nutraceuticals in the future.

Keywords: pheophorbide derivatives, lipid metabolism, zebrafish larvae, lipolysis, mitochondrial respiration

References

- Costa, M., Rosa, F., Ribeiro, T., Hernandez-Bautista, R., Bonaldo, M., Gonçalves Silva, N., Eiriksson, F., Margrét Thorsteinsdóttir, M., Siegfried Ussar, S., Urbatzka, R., 2019. Identification of cyanobacterial strains with potential for the treatment of obesity-related co-morbidities by bioactivity, toxicity evaluation and metabolite profiling. *Mar. Drugs* 17, 280.
- Kamarulzaman, A.F., Shaari, K., Hock Ho, A.S., Lajis, N.H., Teo, S.H., Lee, H.B., 2011. Derivatives of Pheophorbide-a and Pheophorbide-b from Photocytotoxic *Piper penangense* Extract. *Chem. Biodivers.* 8, 494-502.

* Presenting author. E-mail address: nsilva@ciimar.up.pt (Natália Silva)

S3.P208 Phyllobilins derived from chlorophyll degradation are overlooked, bioactive phytochemicals in plant-based food

Christian Elvert^a, Fabian Hammerle^a, Christian Doppler^b, David Bernhard^b, Simone Moser^a, Cornelia A. Karg^{a,*}

^a Department of Pharmacognosy, Institute of Pharmacy, University of Innsbruck, Innrain 80-82, 6020 Innsbruck, Austria

^b Division of Pathophysiology, Institute of Physiology and Pathophysiology, Medical Faculty, Johannes-Kepler-University Linz, Krankenhausstrasse 7a, 4020 Linz, Austria

The breakdown of the green plant pigment chlorophyll in colorful fall foliage is a spectacular natural phenomenon and one of the most visible signs of life on Earth. It is estimated that approximately 10 billion tons of chlorophyll are degraded annually on Earth (Brown et al., 1991). However, the degradation of chlorophyll isn't limited to deciduous trees and shrubs; it also takes place during ripening or post-harvest storage in fruits and vegetables like apples and savoy cabbage (Müller et al., 2007; Karg et al., 2019). The breakdown products, known as phyllobilins, are therefore components of our diet, but were long considered mere by-products of the detoxification process of chlorophyll. Recent research has initiated a paradigm shift by revealing potent bioactivities of phyllobilins, such as antioxidative and anti-inflammatory properties (Wang et al., 2021). Nevertheless, their presence in plant-based food has been overlooked until now. Our research focuses on the comprehensive analysis of the 'phyllobilin-fingerprint' in various plant-based products purchased from the supermarket after post-harvest storage. Furthermore, we study the pharmacokinetic characteristics of phyllobilins with a specific emphasis on their distribution and metabolism in both animal models and human with the ultimate aim of spotlighting phyllobilins as essential and bioactive constituents within our everyday diet thereby considering especially stored produce as potential nutraceuticals. Given the widespread presence of phyllobilins in various plant species, these investigations also hold promise for their potential as ingredients of phytopharmaceuticals.

Keywords: chlorophyll breakdown, phyllobilins, bioactivities, pharmacokinetics, plant-based food

References

- Brown, S.B., Houghton, J.D., Hendry, G.A.F. 1991. In: Chlorophylls. Scheer, H., editor. CRP Press; Boca Raton, USA. pp. 465–489.
- Karg, C.A., Schilling, C.M., Allmendinger, L., Moser, S. 2019. Isolation, characterization, and antioxidative activity of a dioxobilin-type phyloxanthobilin from savoy cabbage. *Journal of Porphyrins and Phthalocyanines*, 23, 881-888.
- Müller, T., Ulrich, M., Ongania, K. H., and Kräutler, B. 2007. Colorless tetrapyrrolic chlorophyll catabolites found in ripening fruit are effective antioxidants. *Angewandte Chemie International Edition* 46, 8699-8702.
- Wang, P., Karg, A.C., Frey, N., Frädrich, J., Vollmar, A.M., Moser, S. 2021. Phyllobilins as a challenging diverse natural product class: Exploration of pharmacological activities. *Archiv der Pharmazie*. 354, 2100061.

* Presenting author. E-mail address: cornelia.karg@uibk.ac.at (Cornelia Karg)

S3.P209 Development of a simple liquid chromatographic method for the determination of sitosterol in dietary supplements

Mădălina-Georgiana Buț^a, Ruxandra Ștefănescu^{b,*}, Amelia Tero-Vescan^a

^a Medical Chemistry and Biochemistry Department, Faculty of Medicine in English, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Târgu Mureș, 540142 Târgu Mureș, Romania

^b Pharmacognosy and Phytotherapy Department, Faculty of Pharmacy, George Emil Palade University of Medicine, Pharmacy, Science, and Technology of Târgu Mureș, 540142 Târgu Mureș, Romania

β-Sitosterol (SIT) is a plant-derived compound classified as a phytosterol that is often found in various plant foods and is used as a supplement for its health advantages including potential effects on cholesterol levels, inflammation, and symptoms associated with benign prostatic hyperplasia, such as lower urinary tract symptoms in men. This prompts inquiries into the efficacy of supplementation, which may be related to the quality of readily available dietary supplements (DS).

The present study aimed to develop and validate a routine, rapid, simple, and efficient method for determining the level of SIT to verify the authenticity of selected DS promoted and sold to consumers. An HPLC/DAD method was developed and validated for the quantitative determination of the SIT. The separation was performed using a reversed-phase C18 column with a mixture of 97% methanol and 3% water in isocratic elution at a flow rate of 1 ml/min. The validated method was specific and linear in the concentration range of 15-400 µg/ml and was precise and accurate at all concentration levels according to the official guidelines for validating analytical methods. The sample pretreatment procedure was optimized to be less time-consuming than any other published method, especially because derivatization was not required.

In conclusion, the developed method provides a reliable means for the rapid and accurate determination of SIT levels in DS, facilitating quality assurance and regulatory compliance in the industry. The method was successfully applied to evaluate the SIT content present in the DS from Romanian market.

Keywords: sitosterol, phytosterols, dietary supplements, liquid chromatography

References

- Buț, M. G., Jitcă, G., Imre, S., Vari, C. E., Ösz, B. E., Jitcă, C. M., & Tero-Vescan, A. (2023). The Lack of Standardization and Pharmacological Effect Limits the Potential Clinical Usefulness of Phytosterols in Benign Prostatic Hyperplasia. *Plants*, 12(8), 1722.
- Salehi, B., Quispe, C., Sharifi-Rad, J., Cruz-Martins, N., Nigam, M., Mishra, A. P., Konovalov, D. A., Orobinskaya, V., Abu-Reidah, I. M., Zam, W., Sharopov, F., Venneri, T., Capasso, R., Kukula-Koch, W., Wawruszak, A., & Koch, W. (2021). Phytosterols: From Preclinical Evidence to Potential Clinical Applications. *Frontiers in pharmacology*, 11, 599959.

* Presenting author. E-mail address: ruxandra.stefanescu@umfst.ro (Ruxandra Ștefănescu)

S3.P210 Effects of Crude leaf extract of *Ficus* sp. on the proliferation and differentiation of C2C12 muscle cell line

Teresa, Ma. Theresa B.^{*}, Dela Rosa, Angelo Mc Court S., Dychiao, Garret Royce K., Ipulan-Colet, Lerrrie Ann^a

Institute of Biology, National Science Complex, UP Diliman, Quezon City, Philippines

Philippine biodiversity is known as source of potential bioactive compounds that can be used as food supplements or medicine. Among these prospective species is *Ficus* sp. which reported of its potential medical use. This study will investigate the effects of Crude leaf extracts (CLE) on myoblast count, proliferation, myotube formation, MyoD expression, antioxidant properties through DPPH assay and effect on glucose consumption in a dose dependent manner. For myoblast count and KI 67 expression, C2C12 myoblast will be observed after 6, 8, 24 and 48 hrs. of CLE treatments. For induction of differentiation assay, myotube area, stained with May-Grünwald Giemsa, will be examined at 1, 3, and 7 day post treatment (dpt) and images will be obtained and analyzed to determine myotube area using ImageJ. In the proliferation assay, 0.09765-3.125 µg/mL concentrations moderately increased cell counts compared to the negative control and cells cultured with 12.5 µg/mL extract yielded significantly lower cell counts versus controls. Similarly, groups treated with this concentration produced significantly lower myotube areas in the induction assay. At 0.7812 µg/mL of 3 dpt, results showed that there is a significant increase in myoblast differentiation. To determine the possible mechanism of action, the antioxidant activity and glucose consumption of cells will be tested using DPPH assay and glucose reading from cell culture media using glucometer. Results will be correlated with phytochemical analysis for the presence of specific secondary metabolites which may contribute to the effects observed in this study.

Keywords: cell culture, crude, extract, myoblast, C2C12

References

- Dugdale HF, Hughes DC, Allan R, Deane CS, Coxon CR, Morton JP, Stewart CE, Sharples AP. The role of resveratrol on skeletal muscle cell differentiation and myotube hypertrophy during glucose restriction. *Mol. Cell. Biochem.* 2018; 444(1–2): 109–123
- Ogawa M, Mizofuchi H, Kobayashi Y, Tsuzuki G, Yamamoto M, Wada S, Kamemura K. Terminal differentiation program of skeletal myogenesis is negatively regulated by O-GlcNAc glycosylation. *Biochim. Biophys Acta Gen. Subj.* 2012; 1820(1): 24–32.
- Recuenco MC, de Luna JRP, Magallon NG, Salamaney, KC. Phytochemical screening, total phenolics, and antioxidant and antibacterial activities of selected Philippine indigenous fruits. *Philipp J. Sci.* 2020; 149(3a): 697–710.
- Zammit PS. Function of the myogenic regulatory factors Myf5, MyoD, Myogenin and MRF4 in skeletal muscle, satellite cells and regenerative myogenesis. *Semin Cell Dev Biol.* 2017;72:19–32. pmid:29127046

^{*} Presenting author. E-mail address: mbteresa@up.edu.ph (Ma. Theresa B. Teresa)

S3.P211 *In vitro* analysis of antioxidant and antidiabetic activity of four plant extracts from the Algarve region

Isabel S. Carvalho^a, Lejsa Jakupović^b, Marijana Zovko Končić^{b,*}

^a MED - Mediterranean Institute for Agriculture, Environment and Development, CHANGE – Associated Laboratory, FCT - Faculty of Science and Technology, Food Science Laboratory, UALG - University of Algarve, Campus de Gambelas, Ed. 8, 8005-139, Faro, Portugal

^b University of Zagreb Faculty of Pharmacy and Biochemistry, A. Kovačića 1, 10000 Zagreb, Croatia

Diabetes mellitus (DM) is a metabolic disease that result from changes in glucose's metabolism and/or deficient insulin production/action. The main goal of present work is to study *in vitro* the potential antioxidant capacity and antidiabetic of aqueous extracts of four plants species of the Algarve region (*Aristolochia baetica* L., *Chelidonium majus* L., *Dittrichia viscosa* L. and *Lavandula virilis* L.). The methods used in the analysis of antioxidant activity were: Total phenolic (TPC) and flavonoid (TFC) content, total antioxidant activity (TAA), DPPH antiradical activity as well as ferric-reducing antioxidant (FRAP) and reducing (RP) power. To analyze the anti-diabetic activity, the enzymatic inhibition assays of α -glucosidase and α -amylase were used. Additionally, the ability of extracts to alleviate consequences of diabetes by inhibiting the enzymes elastase and collagenase was investigated. According to the results, the aqueous extracts of *D. viscosa* showed the best antioxidant activity (1132.99 ± 19.54 mg TE/g dw in FRAP, 623.35 ± 24.02 mg TE/g dw in the RP method and an IC_{50} value of 25.85 ± 0.75 μ g/mL in the DPPH method) as well as the highest content values in total phenolic compounds obtained (477.10 ± 22.29 mg GAE/g dw) and the highest in total flavonoids (22.87 ± 1.73 mg QE/g dw). Aqueous extracts from this plant were also found to show the best inhibitory activity in the α -glucosidase inhibition assay (IC_{50} of 0.61 ± 0.06 mg/mL), which was shown to be more potent than the positive control acarbose, which showed an IC_{50} of 35.17 ± 3.25 mg/mL. *D. viscosa* extracts demonstrated notable anti-elastase and anti-collagenase activity thereby confirming its' status as a promising candidate for development of new antidiabetics.

Keywords: Antioxidants, Algarve, α -glucosidase, *Dittrichia viscosa*

* Presenting author. E-mail address: mzovko@pharma.hr (Marijana Zovko Končić)

S3.P212 Innovative approach towards the qualitative analysis of herbal products using UHPLC-PDA- HRMS/MS: case study on species discrimination within the Lamiaceae family

Loïc Loffredo^{a,b}, Francis Hadji-Minaglou^a, Pierre Champy^b

^a BotaniCERT, Grasse Biotech, Grasse (06), France

^b Paris-Saclay University, CNRS UMR 8076 BioCIS, Orsay (91), France

The use of natural products, such as herbal products, in cosmetics, dietary supplements and pharmaceuticals has grown significantly in recent years, requiring robust measures to guarantee the authenticity of botanical material (Gafner et al., 2023). Existing quality control methods, mainly based on pharmacopoeia standards, are unfortunately inadequate due to the impossibility of assessing the overall chemical composition of ingredients.

In response to this challenge, an untargeted methodology is here developed to enhance the qualitative analysis of plant-derived material. This methodology, based on UHPLC-PDA-HRMS/MS, aims at conducting exhaustive compositional analyses, using qualitative and semi-quantitative comparisons with profiles obtained from authenticated plant material, for precise discrimination between different species, as well as detection of unauthorized substances, of product dilutions, and of undeclared mixtures. Our work focuses mainly on the qualitative aspects of this methodology, applied to the Lamiaceae family because of the wide variety of chemical families found in all genera (Fig. 1).




Fig. 1. Few families found in Lamiaceae. From left to right, Harpagide (Iridoid), Salvanolic acid (phenolic acid), Carnosic acid (diterpene), Luteolin-7-O-glucuronide (flavone) and Menthol (monoterpene)

The development of this analytical model involves untargeted statistical methods combined with molecular networks analysis and annotation tools (Nothias-Scaglia et al., 2015). Our primary objective is to establish a robust discrimination process based on PLS regression within the Lamiaceae family, which will subsequently be extended to the analysis of other plant families.

Keywords: Lamiaceae, HRMS, untargeted, authenticity, dietary supplements

References

- Gafner, S., Blumenthal, M., Foster, S., Cardellina, J.H., Khan, I.A., Upton R., 2023. Botanical Ingredient Forensics: Detection of Attempts to Deceive Commonly Used Analytical Methods for Authenticating Herbal Dietary and Food Ingredients and Supplements. *Journal of Natural Products*. 86, 460-472.
- Nothias-Scaglia, L.F., Esposito, M., Costa, J. Paolini, J., Touboul, D., Litaudon, M., 2015. Les réseaux moléculaires, une approche bio-informatique globale pour interpréter les données de spectrométrie de masse tandem. *Spectra Analyse*. 307, 65-70.

* Presenting author. E-mail address: loic.loffredo@botanicert.com (Loïc Loffredo)

S3.P213 Phenolic composition, antioxidant, and enzyme inhibitory activities of *Saxifraga paniculata* Mill.

Stevan Samardžić^{a,*}, Ivona Veličković^b, Dušan Ušjak^c, Marija Đorđić Crnogorac^d, Miloš Zbiljić^e, Marina Milenković^f, Tatjana Stanojković^d, Branislava Lakušić^e, Zoran Maksimović^a

^a Department of Pharmacognosy, University of Belgrade – Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

^b Institute of Botany and Botanical Garden “Jevremovac”, Department for Morphology and Plant Systematics, University of Belgrade – Faculty of Biology, Studentski trg 16, 11000 Belgrade, Serbia

^c Laboratory for Molecular Biology, Institute of Molecular Genetics and Genetic Engineering, University of Belgrade, Vojvode Stepe 444a, 11221 Belgrade, Serbia

^d Institute of Oncology and Radiology of Serbia, Pasterova 14, 11000 Belgrade, Serbia

^e Department of Botany, University of Belgrade – Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

^f Department of Microbiology and Immunology, University of Belgrade – Faculty of Pharmacy, Vojvode Stepe 450, 11221 Belgrade, Serbia

Alpine saxifrage (*Saxifraga paniculata* Mill.), a rosette-forming subshrub, is distributed in North America, Europe, and Asia. This study aimed to investigate the chemical composition and *in vitro* biological activities of the methanolic extract obtained from the aerial parts of the plant. The phenolic profile and content of the four compounds were determined by LC-DAD-ESI-MS. Total phenolic content (TPC), total flavonoid content (TFC), antioxidant potential as well as enzyme inhibitory capacity were evaluated spectrophotometrically. The antimicrobial activity was tested by the broth- microdilution method, while the cytotoxic effect was examined by the MTT assay. The analysis revealed the presence of 23 compounds, including quercetin, six quercetin glycosides, seven kaempferol glycosides, two caffeoylquinic acids, three galloylhexosides, two digalloylhexosides, galloylquinic acid, and (epi)gallocatechin gallate. The most abundant constituent was quercitrin (1.62%), followed by chlorogenic acid (0.28%), kaempferol deoxyhexoside (0.12%), and rutin (0.07%). TPC and TFC were 26.71% and 2.12%, respectively. The demonstrated antioxidant activity was significant (DPPH test, IC₅₀=9.30 µg/mL; FRAP test, FRAP value=2.89 mmol Fe²⁺/g; ABTS test, IC₅₀=34.22 µg/mL). The extract potently inhibited α-glucosidase (IC₅₀=1.86 µg/mL) and, to a lesser extent, 15-lipoxygenase (IC₅₀=49.60 µg/mL) and α-amylase (IC₅₀=9.95 mg/mL). The minimum inhibitory concentration (MIC) against *Acinetobacter baumannii* was 1000 µg/mL; MIC was higher against the other six tested strains of microorganisms. No effect on cancer cell lines (HeLa, HT-29, HCT-116, LS174) and healthy cells (MRC-5) was observed. The findings expanded the scarce literature data on *S. paniculata* and highlighted its potential, warranting further investigation.

Keywords: *Saxifraga paniculata*, flavonoids, gallic acid derivatives, *in vitro* biological activities

* Presenting author. E-mail address: stevan.samardzic@pharmacy.bg.ac.rs (Stevan Samardžić)

S3.P214 What does the percentage of TBA% say about *Boswellia serrata* extract?

Péter Püski^a, Tímea Körmöczi^b, Róbert Berkecz^b, Anita Barta^a, Ákos Bajtel^a, Tivadar Kiss^{a,*}

^a University of Szeged, Faculty of Pharmacy, Institute of Pharmacognosy Eötvös u 6, 6720 Szeged, Hungary

^b University of Szeged, Faculty of Pharmacy, Institute of Pharmaceutical Analysis Somogyi u 4, 6720 Szeged, Hungary

Boswellia serrata extract (BSE) is used in the management of osteoarthritis. Despite promising pharmacological activities, BSEs are poorly characterized chemically, in food supplements and even in some study drugs applied in clinical trials. Generally, three parameters are used to describe the frankincense extract content: total boswellic acid content (TBA%) mainly for commercial bulk extracts; while 11-keto- β -boswellic acid (KBA) and acetyl-11-keto-boswellic acid (AKBA) contents. TBA% is determined by acid-base titration, while KBA and AKBA content are quantified by HPLC; however, there are no clear minimal values defined for BSE in human application.

In our work, we have analyzed fourteen commercial bulk BSEs used for food supplement production. Our goal was to analyze KBA, AKBA and TBA%, respectively. According to the certificate of analysis of the purchased samples the TBA% was in range of 65% to 90%, and no information about KBA and AKBA contents was provided. The samples were subjected to acid-base titration and HPLC analysis. Targeted UPLC-MS and ¹H-NMR were applied to determine possible adulterants. All samples were compared with the authentic USP reference BSE.

According to our results the KBA and AKBA content of only two products were comparable to the authentic sample. Although the TBA% content of the other ten extracts complied with the expected amount, these samples contained citric acid (6%–11%). Our results suggest undeclared addition of citric acid to comply with declared contents of TBA when using titration methods. Incorporation of citric acid to industrial samples – in order to mimic boswellic acids in titration analysis, – was demonstrated for the first time.

Keywords: *Boswellia serrata*, frankincense, KBA, AKBA, TBA

* Presenting author. E-mail address: kiss.tivadar@szte.hu (Tivadar Kiss)

S3.P215 Pharmacokinetic study of cycloastragenol administered in a liposomal formulation to rats

Petra Cihlarova^{a,*}, Katerina Knizkova^a, Martin Kuchar^{a,b}

^a Forensic Laboratory of Biologically Active Substances, Department of Chemistry of Natural Compounds, University of Chemistry and Technology Prague, Czech Republic

^b National Institute of Mental Health, Klecany, Czech Republic

Cycloastragenol is a triterpenoid saponin found in representatives of the genus *Astragalus* (Fabaceae), namely in *A. trimestris* L., widely used in Chinese traditional medicine. Its most significant proven effect on the organism is its ability to activate the enzyme telomerase, leading to an extension of telomeres and consequently of cellular lifespan. Cycloastragenol is often the active substance of various food supplements, either in its pure form or as a part of *A. trimestris* plant or extracts. The limitation of the effectiveness of these products is the low bioavailability of cycloastragenol which is around 25% after oral administration (He et al., 2022). One potential approach to improve bioavailability is the administration of cycloastragenol as a part of a liposome. In this study, the bioavailability of cycloastragenol administered to rats in the form of liposomes or simple solution is evaluated after HPLC-MS analysis of serum. Consequently a curve describing the pharmacokinetic profile is constructed.

Before the HPLC-MS analysis, the serum samples were prepared using protein precipitation with acetonitrile. Ruscogenin was used as an internal standard. For the chromatographic separation, a C₁₈ column and methanol and 5 mM solution of ammonium formate in water were used. The QTRAP mass spectrometer was operating in ESI+ mode. The reached limit of quantification was 0.7 ng/mL of serum.

The results indicate a promising improvement in the bioavailability of cycloastragenol after incorporation into a liposome compared to administration in the form of a solution, which may lead to better utilization of the positive effects of cycloastragenol.

Keywords: cycloastragenol, *Astragalus trimestris*, telomerase, HPLC-MS, liposome

References

He, M., Wang, K., Che, H., Wang, H., Yang, K., Zhang, G., Yao, J., Wang, U., 2022. A comprehensive review of cycloastragenol: Biological activity, mechanism of action and structural modifications. Eur. J. Med. Chem. Rep. 5.

*Presenting author. E-mail address: petra.cihlarova@vscht.cz (Petra Cihlarova)

S3.P216 An extract of *Boehmeria nivea* induces apoptosis by disrupting oncogenic signaling in human pancreatic cancer cells

Sang-Chul Han^a, Ji Su Han^{a,b}, Seon-A Yoon^c, Hyejin Hyeon^c, Young-Min Ham^c, Yong-Hwan Jung^c, Hye-Jin Boo^{a,b,*}

^a Department of Histology, College of Medicine, Jeju National University, Jeju, Republic of Korea

^b Jeju Research Center for Natural Medicine, Jeju National University, Jeju, Republic of Korea

^c Biodiversity Research Institute, Jeju Technopark, Jeju, Republic of Korea

Pancreatic cancer is one of the most lethal solid tumors due to poor prognosis. Studying pancreatic cancer is essential to addressing the challenges associated with its high mortality rate, limited treatment options, and the difficulty of early detection. *Boehmeria nivea* demonstrates anti-inflammatory and antioxidant properties effective against a variety of diseases. However, little is known about its anti-cancer effects in human pancreatic cancer cells. In the present study, we aimed to explore the potential anti-cancer properties of *Boehmeria nivea* extract against human pancreatic cancer cells and elucidate its molecular mechanism of action. We evaluated the anti-cancer effects of *Boehmeria nivea* water extract through a comprehensive approach, including WST-1 assay, colony formation assay, western blot analysis, Annexin-V/PI staining, migration and invasion assay, as well as animal experiments utilizing a xenograft mouse model. The results revealed a significant inhibition of the growth and colony formation of pancreatic cancer cells upon treatment with *Boehmeria nivea* extract. Furthermore, the extract induced apoptosis in pancreatic cancer cells by enhancing the expression levels of cleaved PARP and cleaved caspase-3, while concurrently inhibiting oncogenic signaling pathways such as MAPK and PI3K/AKT pathways. In vivo experiments using a mouse xenograft model demonstrated that *Boehmeria nivea* extract not only suppressed tumor growth, but also induced apoptosis in human pancreatic cancer cells. These findings offer compelling evidence supporting the notion that *Boehmeria nivea* extract may be a potential therapeutic agent for pancreatic cancer.

Keywords: natural products, *Boehmeria nivea*, apoptosis, pancreatic cancer, carcinogenesis

References

Halbrook, C.J., Lyssiotis C.A., Magliano, M.P., Maitra, A., 2023. Pancreatic cancer: Advances and challenges. *Cell*. 186, 1729-1754.

Hu, Z.I., O'Reilly, E.M., 2024. Therapeutic developments in pancreatic cancer. *Nat. Rev. Gastroenterol. Hepatol.* 21, 7- 24.

Naeem, A., Hu, P., Yang, M., Zhang, J., Liu, Y., Zhu, W., Zheng, Q., 2022. Natural Products as Anticancer Agents: Current Status and Future Perspectives. *Molecules* 27, 1-64.

Yao, P., Liang, S., Liu, Z., Xu, C., 2024. A review of natural products targeting tumor immune microenvironments for the treatment of lung cancer. *Front. Immunol.* 15, 1-26.

Acknowledgments

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2022R1I1A1A01073350, RS-2023-00270936) and was supported by “Regional Innovation Strategy (RIS)” through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (MOE) (2023RIS-009).

* Presenting author. E-mail address: hjboo@jejunu.ac.kr (Hye-Jin Boo)

S3.P217 Chemical screening and biological activity of various species of wood-decay fungi of the genus *Ganoderma*

Petr Maršík^{a,*}, Jan Šťastný^a, Lucie Bašniarová^a, Ivan Jablonský^b, Jaroslav Matějka^a, Anna Mascellani-Bergo^a, Jan Tauchen^a, Andrea Bau Bernardos^c, Ángela Morellá Aucejo^c, Araceli Lérída Viso^c, María Dolores Marcos Martínez^c, Ramón Martínez-Mánez^c, Jakub Tremel^d, Petra Herczogová^d, Pavel Klouček^a

^a Department of Food Science, Faculty of Agrobiological Sciences, Food and Natural Resources, The Czech University of Life Sciences Prague, Kamýcká 129, 165 21 Prague 6 – Suchbátka, Czech Republic

^b Department of Horticulture, Faculty of Agrobiological Sciences, Food and Natural Resources, The Czech University of Life Sciences Prague, Kamýcká 129, 165 21 Prague 6 – Suchbátka, Czech Republic

^c Instituto Interuniversitario de Investigación de Reconocimiento Molecular y Desarrollo Tecnológico (IDM), Universitat Politècnica de València-Universitat de València, Camino de Vera, s/n, Valencia 46022, Spain

^d Department of Molecular Pharmacy, Faculty of Pharmacy, Masaryk University, Palackého Třída 1946/1, 612 00 Brno, Czech Republic

Saprophytic wood-decay fungi of the genus *Ganoderma* are widely used in traditional Asian medicine. The most famous species *Ganoderma lucidum*, (known also as Reishi or Lingzhi), is used as nutraceutical for a supportive medication in the treatment of cancer and to combat side effects during chemotherapy due to its immunomodulatory abilities (Shiao, 2003).

In this work, the differences among chemical profiles of eleven strains belonging to 7 species of *Ganoderma* genus cultivated in culture were evaluated. Methanolic extracts of fruiting bodies were analyzed using UHPLC/MS HRAM (Q-TOF) with subsequent evaluation using multivariate statistical methods. The metabolomic analysis revealed, that *Ganoderma pfeifferi* species had the most distinct metabolic profile compared to all other strains. It showed significantly higher content of several compounds tentatively identified as triterpene derivatives considered to be the principal anticancer constituents of the genus (Ahmad, 2020). Main triterpenoid compounds distinguishing *G. pfeifferi* from the other compared strains were then identified by NMR spectroscopy and MS/MS as applanoxidic acid A, ganoellipsic acid A, ganoderone A and ganoderone B. An anticancer (HeLa cell assay) and anti-inflammatory (COX-2 and NF-κB/AP-1 assays) effects of the MeOH extracts were tested. The extract of *G. pfeifferi* had the strongest anticancer effect in the HeLa anticancer assay. The highest anti-inflammatory potential of NF-κB/AP-1 inhibition was found in several strains of *Ganoderma lucidum* and *G. applanatum*, which were comparable to the positive control prednisolone. In contrast, inhibition of COX-2, the key enzyme of inflammatory processes, was weak or negligible for all extracts.

Keywords: *Ganoderma*, fungi, metabolomics, triterpenes, anticancer assay

References

- Shiao, M.-S., 2003. Natural Products of the Medicinal Fungus *Ganoderma lucidum*: Occurrence, Biological Activities, and Pharmacological Functions. *Chem. Rec.* 3, 172–80.
- Ahmad, M.F., 2020. *Ganoderma lucidum*: A rational pharmacological approach to surmount cancer. *J. Ethnopharmacol.* 260, 113047.

* Presenting author. E-mail address: marsik@af.czu.cz (Petr Maršík)

S3.P218 Identification of a marine cyanobacterium *Rivularia* sp. LEGE 07159 with promising bioactivity on appetite regulation

Ana C. Fonseca^{a,b,*}, Javier Sanz Moxó^a, Marta Moutinho^b, Leonor Ferreira^{a,b}, Vitor Vasconcelos^{a,b}, Mariana Alves Reis^a, Ralph Urbatzka^a

^a CIIMAR - Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Terminal de Cruzeiros Porto de Leixões, Avenida General Norton de Matos S/N, 4450-208 Matosinhos, Porto, Portugal

^b Faculty of Sciences, University of Porto, Rua do Campo Alegre S/N, 4169-007 Porto, Portugal

Obesity is a global health concern with a significant increase in developed countries. Appetite plays a crucial role in food intake representing a target for anti-obesity drugs. Cyanobacteria are known to produce a wide range of bioactive compounds with pharmaceutical relevance. Forty-six cyanobacterial strains belonging to the LEGE – CC of CIIMAR were screened for their ability to reduce the food intake in zebrafish larvae. The screening revealed that out of the 138 analysed fractions, obtained by increasing polarity fractionation (hexane, ethyl acetate and methanol), 14 fractions were able to decrease the labelled *Paramecia bursaria* intake in at least 50 % and 4 different fractions in at least 70%. The screening pinpointed the marine cyanobacterium *Rivularia* sp. LEGE 07159 as a promising strain to produce appetite suppressant compounds, showing 72% of appetite inhibiting activity in the methanolic fraction. The LC-HRESIMS/MS data of this fraction was compared with inactive fractions with the same polarity. The metabolomics network constructed using the GNPS platform revealed 217 unique masses in the bioactive fraction. Sixty-five nodes (30%) were characterized, according to the ClassyFire superclass, as lipids and lipid-like molecules, benzenoids, organic acids and derivatives, phenylpropanoids and polyketides and organoheterocyclic compounds. Some nodes corresponded to compounds with previously described effects on appetite inhibition. Molecular analyses by qPCR revealed an increase of mRNA expression of orexigenic and anorexigenic genes in zebrafish larvae, namely of *cartpt*, *npv*, and *ghrl*. Future work will focus on the isolation of the bioactive compounds and further characterization of the molecular mechanism.

Keywords: Obesity, appetite suppression, bioactive compounds, cyanobacteria

* Presenting author. E-mail address: ana.fonseca@ciimar.up.pt (Ana Catarina Fonseca)

S3.P219 Potent xanthine oxidase inhibitory activity of acacetin from *Agastache rugosa* (Fisch. & C.A.Mey.) Kuntze *in vitro* and molecular docking studies

Heung Joo Yuk, Yoon-Young Sung, Dong-Seon Kim

KM Research Science Division, Korea Institute of Oriental Medicine, Daejeon 34054, Republic of Korea

The main causes of hyperuricemia are the over-production of uric acid, insufficient excretion in the urine, or a combination of both. Xanthine oxidase (XO) inhibitors that block uric acid synthesis in the body can play an important role in preventing hyperuricemia. The aerial parts of *Agastache rugosa* are used as traditional medicine in Asia. A 50% ethanol extract has shown potent XO inhibitory activity. To investigate the major constituents responsible for this effect, bioassay-guided purification led to the isolation of five compounds identified as acacetin glycosides (1–4) and acacetin (5). Acacetin exhibited the most potent inhibitory activity with a half-maximal inhibitory concentration (IC₅₀) value of 0.5 µM, lower than that of allopurinol, which is commonly used as a XO inhibitor. In addition, it showed the strongest effect compared to other flavonoids, such as apigenin and quercetin etc., which are known to have a strong inhibition on XO. Kinetic analyses found that acacetin was a reversible mixed XO inhibitor. Additionally, a molecular docking study using MOE™ tool was carried out to establish the binding mode of the most active flavones with the enzyme, showing important interactions with its catalytic residues. Furthermore, as a result of the hydrolysis of the extract, it was almost converted to acacetin, which displayed the highest efficacy; the content increased by more than five times that before treatment as determined by UPLC-qTof MS. [Acknowledgment : This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (NRF-2022R1C1C1009298).]

Keywords: xanthine oxidase, acacetin, *Agastache rugosa*, gout, hyperuricemia

References

- Santi, M.D., Paulino Zunini, M., Vera, B., Bouzidi, C., Dumontet, V., Abin-Carriquiry, A., Grougnet, R., Ortega, M.G., 2018. Xanthine oxidase inhibitory activity of natural and hemisynthetic flavonoids from *Gardenia oudiepe* (Rubiaceae) in vitro and molecular docking studies. *Eur. J. Med. Chem.* 143, 577-582.
- Yuk, H.J., Kim, J.W., Sung, Y.Y., Kim, D.S., 2021. Phloroglucinol Derivatives from *Dryopteris crassirhizoma* as Potent Xanthine Oxidase Inhibitors. *Molecules* 26, 122.
- Yuk, H.J., R, H.W., Kim, D.S., 2023. Potent Xanthine Oxidase Inhibitory Activity of Constituents of *Agastache rugosa* (Fisch. and C.A.Mey.) Kuntze. *Foods* 12, 573.

* Presenting author. E-mail address: yukhj@kiom.re.kr (Heung Joo Yuk)

S3.P220 Analysis and biological evaluation of Chios Mastic Gum towards its anti-ageing properties

Eleni V. Mikropoulou^a, Aikaterini Argyropoulou^b, Marianna Vanioti^b, Maria Chalkiadaki^a, Aikaterini Basdeki^a, Stéphane Poigny^c, Leandros Skaltsounis^a, Maria Halabalaki^{a,*}

^a Division of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis, Zografou, 15771, Athens, Greece

^b PharmaGnose S.A., 57th km Athens-Lamia National Road, Oinofyta, 32011, Greece

^c Mibelle Group Biochemistry, Mibelle AG, 5033 Buchs, Switzerland

Chios Mastic Gum (CMG) obtained from *Pistacia lentiscus* var. *chia* (Anacardiaceae) is of high pharmacological significance due to its countless assigned properties (Ottria et al., 2023). It finds numerous applications and its use in the area of cosmetics and hygiene is widespread (Soulaïdopoulos et al., 2022). CMG is included in many cosmetic products offering skin care and anti-ageing protection being recommended for the care of photoaged skin and moisturization while it is beneficial for skin types prone to acne and black spots. CMG is a remarkably complex natural resin with an abundance of approximately 120 chemical compounds, belonging to natural polymers, triterpenes, volatile metabolites (monoterpenes, sesquiterpenes etc.) and phenolic compounds. The major triterpenic acids of CMG are masticadienonic (MNA) and iso masticadienonic acids (IMNA) (Pachi et al., 2021, Svingou et al., 2023). In the current work an analytical platform for the quality control of CMG and related cosmetic products was established while CMG fractions (mastic gum without polymer, acidic and neutral fractions and a mixture of MNA and IMNA) were evaluated for their anti-ageing properties. A reliable, robust and simple HPTLC methodology for the determination of MNA and IMNA, was developed and validated. The method was proven to be repeatable and selective for the specific constituents. The method was applied for the analysis of crude resin authentic CMG samples as well as commercial cosmetic products. The biological assays performed showed the promising inhibition activity of the materials in various enzymatic assays and specifically against collagenase, elastase and tyrosinase.

Keywords: Chios Mastic Gum, cosmetics, analysis, biological evaluation, anti-ageing

References

- Ottria, R., Xynomilakis, O., Casati, S., Abbiati, E., Maconi, G., Ciuffreda, P. 2023. Chios Mastic Gum: Chemical Profile and Pharmacological Properties in Inflammatory Bowel Disease: From the Past to the Future. *Int. J. Mol. Sci.* 24, 12038. Pachi, V.K., Mikropoulou, E.V., Gkiouvetidis, P., Siafakas, K., Argyropoulou, A., Angelis, A., Mitakou, S., Halabalaki, M. 2021. Traditional uses, phytochemistry and pharmacology of Chios mastic gum (*Pistacia lentiscus* var. *Chia*, Anacardiaceae): A review. *J. Ethnopharmacol.* 273, 113961.
- Soulaïdopoulos, S., Tsiogka, A., Chrysohoou, C., Lazarou, E., Aznaouridis, K., Doundoulakis, I., Tyrovola, D., Tousoulis, D., Tsioufis, K., Vlachopoulos, C., Lazaros, G. 2022. Overview of Chios Mastic Gum (*Pistacia lentiscus*) Effects on Human Health. *Nutrients* 14, 590.
- Svingou, D., Mikropoulou, E.V., Pachi, V.K., Smyrnioudis, I., Halabalaki, M. 2023. Chios mastic gum: A validated method towards authentication. *J. Food Compos. Anal.* 115, 104997

* Presenting author. E-mail address: mariahal@pharm.uoa.gr (Maria Halabalaki)

S3.P221 Metabolomics of *Withania somnifera* L. extracts by a combined LC-MS and NMR approach

Luciana Maria Polcaro*, Sonia Piacente, Milena Masullo

Department of Pharmacy, University of Salerno, Via Giovanni Paolo II 132, 84084 Fisciano (SA), Italy

Withania somnifera L. (Solanaceae), for over 3000 years, has been considered an essential herb in Ayurvedic Medicine, and the part of the plant of greatest interest is represented by the roots. The roots contain several secondary metabolites, mainly belonging to steroidal lactones called withanolides, which possess various pharmacological activities, including neuroprotective activity. Currently, the demand for *W. somnifera* extracts is increasing (Manjiri 2024), and it is necessary to find an ecologically friendly extraction strategy that allows a higher extraction yield using a lower amount of solvents and energy. Therefore, *W. somnifera* roots were submitted to different extraction protocols like macerations, decoctions, ultrasound-assisted extraction and solid-liquid dynamic extraction (SLDE-Naviglio) using EtOH and EtOH/H₂O (50:50, 60:40, 75:25, 100:0) as bio-solvents. The extracts were analyzed using a combined approach based on LCESI/QExactive/MS/MS and NMR analysis. The LC-MS and NMR data were analyzed using Principal Component Analysis to highlight how the extraction method can affect the chemical profile of the extracts (Fig.1). The extracts obtained using EtOH 100:0 were the best in terms of withanolides.




Fig.1. PCA score scatter plot of *W. somnifera* extracts of LC-MS data.

Moreover, based on the use of this plant in the treatment of neurological disorders and the evidence that in Parkinson's disease neurons that contain the dark-brown cytoplasmic pigment neuromelanin are particularly susceptible to neurodegeneration (Carballo-Carbajal et al., 2019), the tyrosinase inhibitory activity of the extracts was herein tested by a spectrophotometric assay. The obtained results showed how all *W. somnifera* extracts inhibited the tyrosinase enzyme in a range from 32.86 and 85.36 µg/ml in terms of IC₅₀ values.

Keywords: *Withania somnifera*, extractions, LC-MS analysis, NMR analysis

References

Manjiri, K., 2024. Withania Somnifera Extract Market Report. Pharma & Healthcare.

Carballo-Carbajal, I., Laguna, A., Romero-Giménez, J., Cuadros, T., Bové, J., Martínez-Vicente, M., Parent, A., Gonzalez-Sepulveda, M., Peñuelas, N., Torra, A., Rodríguez-Galván, B., Ballabio, A., Hasegawa, T., Bortolozzi, A., Gelpi, E., & Vila, M., 2019. Brain tyrosinase overexpression implicates age-dependent neuromelanin production in Parkinson's disease pathogenesis. Nat. Commun. 10(1), 973.

* Presenting author. E-mail address: lpolcaro@unisa.it (Luciana Maria Polcaro)

S3.P222 A food supplement derived from high-phenolic Kalamon table olives reduced cholesterol in patients with mild dyslipidemia

Annia Tsolakou^a, Eleni Melliou^{a,*}, Martha S. Katsarou^b, Prokopios Magiatis^a, Nikolaos Drakoulis^b

^a Laboratory of Pharmacognosy and Natural Products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Greece

^b Research Group of Clinical Pharmacology and Pharmacogenomics, Department of Pharmacy, National and Kapodistrian University of Athens, Greece

This study investigated the effects of a nutritional supplement CHOLESTOLIVE[®] derived from high-phenolic table olives (*Olea europaea* L. var. Kalamon) on lipid parameters in patients with mild dyslipidemia. The supplement was produced employing solely natural and mechanical methods, through a process involving removal of olive oil, aqueous extraction of tyrosol, hydroxytyrosol, and lactic acid, removal of water, and removal of salt. The obtained extract was formulated to hard capsules for oral use and the phenolic content of two capsules was adjusted to be equivalent to the amount obtained after daily consumption of five table olives (hydroxytyrosol 28 mg, tyrosol 12 mg, lactic acid 81 mg).

In a 30-day clinical study, that was performed in three major hospitals in Greece, 48 volunteers (19 men, 29 women) with mild dyslipidemia (mean total cholesterol 232.54 ± 26.88 mg/dL) were enrolled and instructed to take two capsules daily during meals, while maintaining their lifestyle and diet. The study resulted in a significant reduction of total cholesterol (-4.16%, p.s.=5%, Z=-2.518, p=0.012) and LDL cholesterol (-5.67%) levels following supplementation. Triglyceride levels showed a modest reduction but did not reach statistical significance. HDL cholesterol was not affected throughout the study. These findings suggest that the nutritional supplement may have beneficial effects on reducing total cholesterol and LDL cholesterol levels, highlighting the potential health benefits of the bioactive compounds found in table olives, particularly hydroxytyrosol, tyrosol and lactic acid, related to cardiovascular well-being and metabolic health. Larger-scale research is needed to confirm these results and investigate underlying mechanisms.

Keywords: olives, cholesterol, food supplement, clinical trial, hydroxytyrosol

* Presenting author. E-mail address: emelliou@pharm.uoa.gr (Eleni Melliou)

S3.P223 The rising phenomenon of cannabis ‘tea’ drinking: a safe and healthy alternative?

Matěj Malý*, František Beneš, Zuzana Bínová, Jana Hajšlová

Department of Food Analysis and Nutrition, University of Chemistry and Technology, Technická 5, 166 28 Prague 6, Czech Republic

Drinking ‘tea’ prepared from *Cannabis sativa* L. leaves and inflorescence has gained popularity in recent years due to its claimed beneficial effects on consumers’ health. However, even hemp may contain, besides various phytocannabinoids and other bioactive secondary metabolites, a non-negligible amount of psychotropic delta-9-tetrahydrocannabinol (Δ^9 -THC) and its precursor, delta-9-tetrahydrocannabinol acid (Δ^9 -THCA). To evaluate the risk of exceeding the Acute Reference Dose (ARfD) for Δ^9 -THC (1 μ g/kg body weight, set by the European Food Safety Authority, EFSA), when drinking cannabis tea, we quantified, using UHPLC-HRMS/MS method, 42 phytocannabinoids and 12 flavonoids transferred from three different *Cannabis* cultivars into aqueous infusions (1g of dried *Cannabis* + 250 ml of water, boiled for 10 minutes), including less-known trans- Δ^9 -THC isomers, cis- Δ^9 -THC and exo-THC. The transfer of phytocannabinoid acids was, on average, 15 times higher than their neutral, more lipophilic analogues. Moreover, only 0.4 to 1.9% of Δ^9 -THC was transferred under conditions of infusion preparation. Considering a 70kg individual drinking 250 ml of cannabis tea daily, Δ^9 -THC equivalent intake would correspond to 15, 21, and 180% of ARfD, depending on *Cannabis* variety. The average transfer rate for targeted flavonoids was 53%. However, the addition of 20 g of cream with 10 % fat to the tea, as recommended by some producers, resulted in a significant increase in the content of phytocannabinoids in the infusion (26 - 246 times for Δ^9 -THC) due to their lipophilic nature. Therefore, the risk assessment the way of cannabis teas preparation must be carefully considered.

Keywords: cannabis tea, phytocannabinoid transfer, UHPLC-HRMS/MS, Δ^9 -THC, health risk assessment

* Presenting author. E-mail address: matej.maly@vscht.cz (Matej Malý)

S3.P224 HPLC-MS analysis and AChE activity of aerial parts and root extracts of *Taraxacum mirabile* Wagenitz, an endemic species of Turkey

Seçil Karahüseyin^{a*}, Nurten Özsoy^b, Aynur Sari^c

^a Department of Pharmacognosy, Faculty of Pharmacy, Cukurova University Balcali, Saricam- Adana, Turkey

^b Department of Biochemistry, Faculty of Pharmacy, Istanbul University Beyazit, Fatih-Istanbul, Turkey

^c Department of Pharmacognosy, Faculty of Pharmacy, Istanbul University Beyazit, Fatih-Istanbul, Turkey

This presentation will cover the phenolic compounds of *Taraxacum mirabile* Wagenitz aerial parts and root extracts determined by HPLC-MS method and acetylcholinesterase enzyme activity (AChE) results of petroleum ether, dichloromethane, ethyl acetate and butanol fractions of *T. mirabile* Wagenitz aerial parts and root extracts. The aerial parts and root dichloromethane fractions of *T. mirabile* Wagenitz were analyzed for phenolic compounds and HPLC-MS results were obtained as follows:

The substances found in the aerial parts dichloromethane fraction were chlorogenic acid, fumaric acid, verbascoside, chicoric acid, caffeic acid, luteolin-7-rutinoside, vanillic acid, luteolin-7- glycoside, apigenin-7-glycoside, salicylic acid, luteolin, apigenin, hispidulin and chrysin compounds. The substances found in the root dichloromethane fraction were chlorogenic acid, fumaric acid, verbascoside, chicoric acid, caffeic acid, luteolin-7-rutinoside, vanillic acid, luteolin-7-glycoside, salicylic acid, luteolin, apigenin, hispidulin and chrysin compounds. According to these results, the most abundant compound in the aerial parts dichloromethane fraction was luteolin, while the most abundant compound in the root dichloromethane fraction was vanillic acid.

AChE inhibitory activities of the ethyl acetate, butanol, dichloromethane and petroleum ether extracts from roots and aerial parts of *T. mirabile* Wagenitz were tested using Ellman's colorimetric method in 96-wells microplates (Ellman et al., 1961). All the extracts showed strong inhibitory effect against acetylcholinesterase up to 90%, and thus may be beneficial in the treatment of Alzheimer's disease. The highest inhibitory activity was obtained with the dichloromethane and petroleum ether extracts from roots, followed by dichloromethane and petroleum extracts from aerial parts, ethyl acetate and butanol extract from roots, whereas only moderate inhibition of the enzyme was observed for the ethyl acetate and butanol extracts from aerial parts.

Keywords: *Taraxacum mirabile* Wagenitz, Aerial parts, Root, HPLC-MS, AChE activity

References

Ellman, G. L., Courtney, K. D., Andres, V., & Featherstone, R. M., 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology*. 7(2), 88–95.

* Presenting author. E-mail address: skarahuseyin@cu.edu.tr (Seçil Karahüseyin)

S3.P225 Dietary soft electrophiles upregulate lipid pro-resolving molecules in a *Drosophila* model of Parkinson's disease

Swarnali Chatterjee^{a,*}, Bianca McCarty^a, Madison Bever^a, Qiaoli Liang^b, Lukasz Ciesla^a

^a Department of Biological Sciences, The University of Alabama, Tuscaloosa AL 35487

^b Department of Chemistry and Biochemistry, The University of Alabama, Tuscaloosa AL 35487

As humans age, there is a decline in cognitive abilities due to chronic systemic inflammation, known as inflammaging. This further leads to age-related morbidities and neurodegeneration (Maitra et al., 2019). Neurodegenerative diseases, like Parkinson's disease (PD), currently have no effective treatment options, emphasizing the need for improved management strategies. Epidemiological data suggest that a regular dietary intake of flavonoids and omega-3 fatty acids enhances cognitive capacity in both animal models and humans and delays the onset of age-related neurological diseases (Seigler et al., 2021). We propose that dietary phytochemicals like flavonoids may mimic the physiological effects of omega-3-derived pro-resolving and anti-inflammatory molecules. Therefore, they might be involved in the process of the resolution of neuroinflammation. Additionally, we propose that the anti-inflammatory and neuroprotective activity of these dietary molecules is directly related to their soft electrophilic properties. Through a variety of analytical approaches, we have identified these compounds *in vivo* in the context of a living organism (*Drosophila melanogaster*), focusing on their anti-inflammatory properties (Maitra et al., 2021). Our preliminary results suggest that dietary soft electrophiles, such as gardenin A, thymoquinone, and vitamin E vitamers, enhance the production of lipid pro-resolving molecules in a paraquat-induced *Drosophila* model of PD. We have recently shown that oral administration of a polymethoxyflavonoid, gardenin A, modifies the lipid profile in a mammalian model of PD (Hack et al., 2023). My future studies will include an insight into the mechanisms by which these phytochemicals might bring about pro-resolution effects in this *Drosophila* PD model by inhibiting cyclooxygenase (COX) activity. Thus, lipophilic soft electrophiles may have potential therapeutic applications in addressing neuroinflammation-related conditions.

Keywords: neurodegenerative, inflammation, cognitive, phytochemicals, pro-resolution

References

- Maitra, U., Scaglione, M. N., Chtarbanova, S. & O'Donnell, J. M. (2019) Innate immune responses to paraquat exposure in a *Drosophila* model of Parkinson's disease. *Scientific Reports* 9, 12714.
- Seigler, D. S., Friesen, J. B., Bisson, J., Graham, J. G., Bedran-Russo, A., McAlpine, J. B., & Pauli, G. F. (2021). Do certain flavonoid IMPS have a vital function?. *Frontiers in Nutrition*, 8, 762753.
- Maitra, U., Harding, T., Liang, Q., & Ciesla, L. (2021). GardeninA confers neuroprotection against environmental toxin in a *Drosophila* model of Parkinson's disease. *Communications Biology*, 4(1), 162.
- Hack, W., Gladen-Kolarsky, N., Chatterjee, S., Lang, Q., Maitra, U., Ciesla, L., & Gray, N. E. (2023). Gardenin A improves cognitive and motor function in A53T- α -syn mice. *bioRxiv*, 2023-10.

*Presenting author. E-mail address: schatterjee5@crimson.ua.edu (Swarnali Chatterjee)

S3.P226 Comprehensive analysis of commercial kratom products: alkaloids, polyphenols, heavy metals and microbial contaminations

Hana Walaská*, Petra Cihlářová, Martin Kuchař, Lenka Kohoutová, David Kačer

Forensic Laboratory of Biologically Active Substances, Department of Chemistry of Natural Compounds, UCT Prague, Technická 3, 166 28 Prague, Czech Republic

Kratom (*Mitragyna speciosa*) leaves have been traditionally used in Asia and Indochina for pain relief, fatigue reduction, or mood lifting. These effects are primarily attributed to its indole and oxindole bioactive alkaloids, among which mitragynine is the major one, accompanied by over 40 others, which interact with opioid, serotonin, and dopamine receptors (Brown et al., 2017). Kratom has recently become popular in Western society as well. Here, however, it is often sold in the form of finely ground leaves or concentrated extracts and drinks (Swogger et al., 2022). In many countries, these products are outside legislative regulations, and their production is therefore not subject to any quality and safety requirements (Prozialeck et al., 2019).

The aim of this study was to develop analytical methods for quantification of five kratom alkaloids (mitragynine, 7-hydroxymitragynine, speciogynine, speciociliatine, and paynanthein) and 33 polyphenols in kratom products, utilizing a Vanquish UHPLC system and Orbitrap Exploris 120 mass spectrometer. We successfully developed analytical methods for the quantification of alkaloids and polyphenols in kratom samples. We employed these methods to analyze a number of commercial samples sourced from the Czech market, subsequently conducting assessments for heavy metal content and microbial contamination.

Our comprehensive analysis and critical evaluation of the results significantly enhances the understanding of the diverse pharmacological effects exhibited by various types of kratom. Particularly, our investigation contributes substantially to the assessment of safety parameters associated with the commercial products available for consumption.

Keywords: Kratom alkaloids, *Mitragyna speciosa*, HPLC-MS, mitragynine, polyphenols

References

- Brown, P. N., Lund, J. A., Murch, S. J., 2017. A botanical, phytochemical and ethnomedicinal review of the genus *Mitragyna* Korth: Implications for products sold as kratom. *J. Ethnopharmacol.* 202, 302-325.
- Swogger, M. T., Smith, K. E., Albert, G. R., Grundmann, O., Veltri, C. A., Henningfield, J. E., Busch, L. Y., 2022. Understanding Kratom Use: A Guide for Healthcare Providers. *Front. Pharmacol.* 13.
- Prozialeck, W. C., Avery, B. A., Boyer, E. W., Grundmann, O., Henningfield, J. E., Kruegel, A. C., McMahon, L. R., McCurdy, C. R., Swogger, M. T., Veltri, C. A., Singh, D., 2019. Kratom policy: The challenge of balancing therapeutic potential with public safety. *Int. J. Drug Policy* 70, 70-77.

* Presenting author. *E-mail address:* walaskah@vscht.cz (Hana Walaska)

S3.P227 Diarylheptanoids from *Corylus avellana* cultivar “Tonda di Giffoni”: antioxidant effects through the activation of the Nrf2 pathway as assessed via molecular docking and biological evaluation

Antonietta Cerulli^{a,b,*}, Simona De Vita^a, Milena Masullo^a, Carmen Bove^a, Paloma Bermejo Bescós^c, Giuseppe Bifulco^a, Sonia Piacente^{a,b}

^a Department of Pharmacy, University of Salerno, Via Giovanni Paolo II 134, 84084 Fisciano, Italy

^b Agritech National Research Center, Corso Umberto 40, 80138 Naples, Italy

^c Departamento de Farmacología, Farmacognosia y Botánica, Universidad Complutense de Madrid, 28040 Madrid, Spagna

Hazelnut (*Corylus avellana* L., Betulaceae) represents one of the most consumed foods, with a high content of healthy lipids. Food and Drug Administration (FDA) has recognized hazelnut as a “heart- healthy” food due to its nutritional and nutraceutical properties related to a large range of bioactive and health-promoting compounds (Bottone, 2019).

Kernels and byproducts of the Protected Geographical Indication (PGI) hazelnut cultivar “Tonda di Giffoni” contain diarylheptanoids derivatives, natural compounds able to prevent oxidative damage of human plasma lipids (Bottone, 2019).

Based on the effects of diarylheptanoids on oxidative stress, the antioxidant effects of giffonins E (1), H (2), J (3), and carpinontriol B (4), isolated from “green” extract (EtOH: H₂O, 50:50) of green leafy covers and kernels, were assessed using molecular docking experiments and *in vitro* tests. The isolated compounds were tested for their ability to disrupt the interaction between Kelch-like ECH- associated protein 1 (Keap1) and nuclear erythroid 2-related factor 2 (Nrf2), triggering the activation of the Nrf2-mediated pathway and consequently promoting the transcription of antioxidant enzymes (Bello, 2017). The results suggested an antioxidant effect of giffonin H and carpinontriol B. Moreover, compounds 1-4 were tested for the antioxidant and cytoprotective properties in human neuroblastoma SH-SY5Y and human embryonic kidney cell lines HEK/tau. showing a significant protection against H₂O₂-induced cytotoxic damage in both cell models. Finally, tested compounds were able to reduce cellular oxidative stress by decreasing ROS production and levels of lipid peroxidation, increasing cell viability, and attenuating cell apoptosis (caspase-3 activity).

Keywords: diarylheptanoids, *C. avellana* “Nocciola di Giffoni” PGI, green leafy covers, molecular docking, antioxidant activity

References

- Bottone, A., Cerulli, A., D’Urso, G., Masullo, M., Montoro, P., Napolitano, A., Piacente, S, 2019. Plant Specialized Metabolites in Hazelnut (*Corylus avellana*) Kernel and Byproducts: An Update on Chemistry, Biological Activity, and Analytical Aspects. *Planta Med* 2019, 85, 840-855.
- Bello, M., Morales-González, J.A, 2017. Molecular recognition between potential natural inhibitors of the Keap1-Nrf2 complex. *International journal of biological macromolecules* 2017, 105, 981-992.

Acknowledgments

This study was carried out within the Agritech National Research Center and received funding from the European Union Next-GenerationEU (PIANO NAZIONALE DI RIPRESA E RESILIENZA (PNRR) – MISSIONE 4 COMPONENTE 2, INVESTIMENTO 1.4 – D.D. 1032 17/06/2022, CN00000022). This manuscript reflects only the authors’ views and opinions, neither the European Union nor the European Commission can be considered responsible for them.

* Presenting author. E-mail address: acerulli@unisa.it (Antonietta Cerulli)

S3.P228 Valorization of Almond Okara as a valuable source of polyphenols and proteins

Mariam Taha^{a,*}, Jennifer Samaillie^a, Sevser Sahpaz^a, Nicolas Blanchemain^b, Krasimir Dimitrov^a, Céline Rivière^a

^a Joint Research Unit 1158, BioEcoAgro, Univ. Lille, 59650 Villeneuve-d'Ascq, France

^b Inserm U1008 Controlled Drug Delivery Systems & Biomaterials, University of Lille, France

Almond milk is one of the most popular plant-based beverages. However, its production generates a solid residue byproduct called “okara”, which is underutilized despite its potential bioactive constituents. Our objective is to deepen our knowledge of almond okara, composition with a focus on the levels of polyphenols, fibers, proteins, lipids, and vitamin E.

Solid-liquid extractions were conducted, varying several parameters including extraction time, solvent concentration, solid/solvent ratio, and extraction temperature. The results showed that the highest extraction yield was obtained after 90 minutes of extraction, which could be explained by considering that the extended exposure allowed the solvent to infiltrate into the matrix, thereby facilitating the solubility process.

Moreover, the results highlighted that the polyphenol yield in almond okara was positively associated with solvent concentration and temperature (Fig.1). Specifically, we found that using 50% ethanol at 60°C resulted in the highest levels of polyphenols, tannins, flavonoids, and antioxidant activity. This can be explained by the fact that hydro-ethanolic solvent increases the permeability of plant tissues enabling better mass transfer by diffusion. Moreover, high temperatures decrease surface tension and solvent viscosity and in return enhance the degree of diffusion and solubility of analytes.




Fig.1: Total polyphenol (TP) content, in mg gallic acid equivalent/ 100 g dry okara, obtained in different ethanol concentrations and extraction temperatures.

In conclusion, our results demonstrated that through its composition, almond okara has health- promoting potential. This makes it a very promising byproduct to facilitate the formulation of innovative industrial approaches, including cosmetics and nutraceuticals.

Keywords: Almond okara, solid-liquid extraction, phenolic compounds, antioxidant activity

* Presenting author. E-mail address: mariam.taha@univ-lille.fr (Mariam Taha)

S3.P229 Application of annular centrifugal extraction (ACE) in honey - isolation and identification of honey bioactive compounds

Gabriela Belén Lemus Ringlele^{a,*}, Petros Gkiouvetidis^a, Dimitris Michailidis^b, Elodie Bossard^c, Lorène Gourguillon^c, Aikaterini Argyropoulou^{a,b}, Stavros Beteinakis^a, Alexios-Leandros Skaltsounis^a, Maria Halabalaki^a

^a Division of Pharmacognosy and Natural products Chemistry, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimioupoli, Zografou, 15771, Athens, Greece

^b PharmaGnose S.A., 57th km Athens-Lamia National Road, Oinofyta, 32011, Greece

^c LVMH recherche, Advances biobased and bioinspired ingredients, Saint-Jean-de-Braye, France

Honey's intricate chemical composition and viscous nature pose challenges in handling. In recent literature, different extraction techniques have been extensively discussed (Hanim Hassan et al., 2022). Nevertheless, these conventional methods present many limitations, i.e., low yields, so there is a need for an efficient, repeatable, and reproducible extraction method. Thus, the primary aim of this study was the development of an extraction protocol using annular centrifugal extraction (ACE) that would allow the recovery of an enriched in non-sugar compounds extract while simultaneously eliminating the sugar content of honey. This approach involves single-stage extractors based on the continuous counter-current liquid-liquid extraction principle with applications mostly in the industrial field (Hamamah et al., 2022). As for food commodities, ACE has only been applied by our group in the recovery of biophenols and lignans from olive and sesame oil, respectively, and is its first application in honey (Michailidis et al., 2019). Following extraction and characterization of the obtained extracts with liquid chromatography - high-resolution tandem mass spectrometry (LC- HRMS/MS), the second aim of this study was the isolation and identification of possible bioactive compounds. After fractionation with fast centrifugal partition chromatography (FCPC) and purification with other chromatographic techniques, more than 20 compounds were isolated and identified from Greek and French honeys. Among these compounds belonging to simple phenols, megastigmanes, and fatty acids, 1-(4-methoxyphenyl)-ethane-1,2-diol, indicated in our previous study as a biomarker for heather honey, was isolated and identified for the first time in honey.

Keywords: Honey, extraction, ACE, isolation, identification

References

- Hamamah, Z. A., & Grützner, T. (2022). Liquid-Liquid Centrifugal Extractors: Types and Recent Applications – a Review. In *ChemBioEng Reviews* (Vol. 9, Issue 3, pp. 286–318). John Wiley and Sons Inc.
- Hanim Hassan, N., Cacciola, F., Chong, N. S., Arena, K., Marriott, P. J., & Wong, Y. F. (2022). An updated review of extraction and liquid chromatography techniques for analysis of phenolic compounds in honey. In *Journal of Food Composition and Analysis* (Vol. 114). Academic Press Inc.
- Michailidis, D., Angelis, A., Aligiannis, N., Mitakou, S., & Skaltsounis, L. (2019). Recovery of sesamin, sesamol, and minor lignans from sesame oil using solid support-free liquid-liquid extraction and chromatography techniques and evaluation of their enzymatic inhibition properties. *Frontiers in Pharmacology*, 10(JUN).

Acknowledgements

The research work was supported by the Hellenic Foundation for Research and Innovation (HFRI) under the 4th Call for HFRI PhD Fellowships (Fellowship number: 9427).

* Presenting author. E-mail address: glemus@pharm.uoa.gr (Gabriela Belén Lemus Ringlele)

S3.P230 Ornamental varieties of *Achillea millefolium* L. as a source of skin-lightening ingredients for natural cosmetics

Katarzyna Gawel-Bęben^{a,*}, Paulina Lechwar^a, Magdalena Lasota^a, Karolina Czech^a, Wirginia Kukula-Koch^b

^a Department of Cosmetology, University of Information Technology and Management in Rzeszow, 2 Sucharskiego Str., 35-225 Rzeszów, Poland

^b Department of Pharmacognosy with Medicinal Plant Unit, Medical University of Lublin, 1 Chodzki Str., 20-093 Lublin, Poland

Achillea millefolium L. is one of the best known medicinal plants with long history of usage in both traditional and modern medicine. (Strzępek-Gomółka et al., 2021a) *Achillea Millefolium* Oil, Extract, Flower Extract, Flower Water, Flower/Leaf/Stem Extract and Flower/Leaf/Stem Juice are also listed in the Cosmetic Ingredients Database CosIng as anti-seborrheic, cleansing, refreshing, skin conditioning, soothing, antioxidant and fragrance ingredients of cosmetics. (CosIng) In addition to the common *A. millefolium* with white flowers, there are many ornamental varieties with colorful flowers. In this study the phytochemical content, antioxidant and tyrosinase-inhibitory properties of *A. millefolium* hydroethanolic extracts obtained from the flowers of ornamental cultivars Skysail White (W), Yellow (Y), Bright Pink (P) and Fire (W) were compared. The content of phytochemicals between varieties was significantly different – W extract showed the highest polyphenolic and flavonoid levels (160.16 and 62.12 µg/mL, respectively), whereas Y cultivar contained the least polyphenols and flavonoids (86.77 and 35.05 µg/mL, respectively). The W and Y extracts showed also the highest and lowest antioxidant activity, respectively. Interestingly, Y extract was the most potent inhibitor of tyrosinase (41% inhibition at 50 µg/mL). The observed activity may result from the specific content of phytocompounds previously described as tyrosinase inhibitors. (Strzępek- Gomółka et al., 2021b) The extracts from *A. millefolium* varieties were also analysed using EpiDerm Phototoxicity Test according to OECD TG 498 guidelines. None of the tested extracts were phototoxic following UVA exposure. The Y and P extracts slightly increased the viability of UVA- treated EpiDerm tissues, suggesting their photoprotective potential.

Keywords: *Achillea millefolium*, cosmetic, tyrosinase, LCMS, phototoxicity

References

- Strzępek-Gomółka M, Gawel-Bęben K, Kukula-Koch W. 2021. *Achillea* Species as Sources of Active Phytochemicals for Dermatological and Cosmetic Applications. *Oxid Med Cell Longev*. 2021a:6643827.
- Strzępek-Gomółka M, Gawel-Bęben K, Angelis A, Antosiewicz B, Sakipova Z, Kozhanova K, Głowniak K, Kukula- Koch W. 2021b. Identification of Mushroom and Murine Tyrosinase Inhibitors from *Achillea biebersteinii* Afan. Extract. *Molecules*. 26(4):964.
- CosIng Cosmetic Ingredient Database of the European Commission (Accessed on 16 February 2024). Available Online: <https://Ec.Europa.Eu/Growth/Tools-Databases/Cosing/>.

* Presenting author. E-mail address: kagawel@wsiz.edu.pl (Katarzyna Gawel-Bęben)

S3.P231 Impact of the drying process and incorporation of natural preservatives on the viability of lactic acid bacteria in yogurts

Jonata M. Ueda^{a,b,c}, Patricia Morales^c, Márcio Carrocho^{a,b}, Sandrina A. Heleno^{a,b,*}

^a Mountain Research Centre (CIMO), Polytechnic Institute of Bragança, Campus Santa Apolónia, 5300-253 Bragança, Portugal

^b Associate Laboratory for Sustainability and Technology (SusTEC), Polytechnic Institute of Bragança, Campus Santa Apolónia, 5300-253 Bragança, Portugal

^c Dept. of Nutrition and Bromatology II, Faculty of Pharmacy, Complutense University of Madrid, Madrid, Spain

Dairy products are among the most widely consumed foods globally, with yogurts particularly demanded for their flavour and nutritional benefits, including essential macronutrients, vitamins, minerals, and probiotics. To mitigate food waste, methods to prolong shelf life are emphasized, such as the incorporation of food preservatives or the creation of powdered foods (Ueda *et al.*, 2023). Therefore, the aim of this study was to assess the survival of total lactic acid bacteria in yogurts after the incorporation of natural extracts (rosemary, basil, and sage), as well as to evaluate their viability after the freeze-drying process. Yogurts were prepared in the laboratory, with lyophilized plant extracts (30 mg/kg) added. Total lactic acid bacteria were monitored for 14 days, alongside the assessment of lactic acid and lactose level using HPLC-DAD and HPLC-RI, respectively. Subsequently, the viability of *Streptococcus thermophilus* and *Lactobacillus bulgaricus* was evaluated in freeze-dried yogurts without the addition of the vegetable extracts (Kok and Hutkins, 2018; Terzaghi and Sandine, 1975). The results indicated a slight increase in lactic acid bacteria counts, as well as an increase in lactic acid and a reduction in the lactose content over the 14-day period, confirming the yogurt fermentation process. During the lyophilization, a slight decrease in the viability of both bacteria was observed, but it remained within the Codex Standards limits (minimum of 10^7 CFU/g). Therefore, the incorporation of plant extracts and the freeze-drying process did not significantly change the viability of lactic acid bacteria in yogurts, remaining a probiotic food.

Keywords: yogurts, lactic acid bacteria, powdered foods, freeze-drying, natural products

References

- Ueda, J. M., Morales, P., Fernández-Ruiz, V., Ferreira, A., Barros, L., Carrocho, M., Heleno, S. A., 2023. Powdered Foods: Structure, Processing, and Challenges: A Review. *Applied Sciences*, 13(22), 12496.
- Kok, C. R., Hutkins, R., 2018, Yogurt and other fermented foods as sources of health-promoting bacteria. *Nutrition Reviews*, 76, 4-15.
- Terzaghi, B. E., Sandine, W. E., 1975. Improved Medium for Lactic Streptococci and Their Bacteriophages. *Applied Microbiology*, 29, 6, 807-813.

* Presenting author. E-mail address: sheleno@ipb.pt (Sandrina A. Heleno)

S3.P232 NMR metabolomics and chemometrics of *Malus domestica* (Suckow) Borkh. varieties from Molise region of Italy

Vadym Samukha^{a,*}, Francesca Fantasma^a, Gilda D'Urso^b, Vincenzo De Felice^a, Gabriella Saviano^a, Gianluigi Lauro^b, Agostino Casapullo^b, Giuseppe Bifulco^b, Maria Giovanna Chini^a, Maria Iorizzi^a

^a Department of Biosciences and Territory, University of Molise, Pesche (IS), Italy

^b Department of Pharmacy, University of Salerno, Fisciano (SA), Italy

Natural products identified in plants are an important resource for both food and drug discovery field. This is because they offer not only primary metabolites that satisfy our food requirement but also a large library of secondary metabolites which can show different bioactivities and can find a use in different areas such as nutraceuticals, cosmeceuticals and for the drug discovery.

This project is based on the valorization of the Molise region of Italy, exploiting several autochthonous varieties belonging to *Malus domestica* (Suckow) Borkh. (Fig. 1): the metabolic profile, with primary and secondary metabolites, can vary in the same species qualitatively and quantitatively if it grows in different environmental conditions, including the type of soil (Sampaio et al., 2016). Apples are rich of nutritional properties and represent an important source of polyphenols which are responsible for their antioxidant and anti-inflammatory properties and possess a role in the prevention of degenerative diseases (Geană et al., 2021).

The phytoconstituents of the varieties of *Malus domestica* from Molise region of Italy will be identified through 1D and 2D-NMR analysis of the extracts obtained by a specific extraction protocol. The qualitative analysis will be followed by a quantitative approach of the most important metabolites identified and by a chemometric analysis with the PCA and PLS-DA techniques in order to observe differences among the varieties. Moreover, the total phenolic, flavonoids, and condensed tannins content will be assessed, and in vitro antioxidant activity will be evaluated by using DPPH scavenging activity, the ABTS scavenging assay, and ferric-reducing antioxidant power (FRAP).




Fig. 1. Selected *Malus domestica* varieties of Molise region in Italy (a: Annurca; b: Gelata; c: Limoncella; d: Zitella)

Keywords: *Malus domestica*, NMR, metabolomics, chemometrics, metabolic profile

References

- Geană, E.I. "Profiling of phenolic compounds and triterpene acids of twelve apple (*Malus domestica* Borkh.) cultivars." *Foods* 10.2 (2021): 267.
- Sampaio, B.L. "Effect of the environment on the secondary metabolic profile of *Tithonia diversifolia*: a model for environmental metabolomics of plants." *Scientific reports* 6.1 (2016): 1-11.

Acknowledgments

The research activity was supported by: University of Molise (Start-Up 2022): PROJECT_20232024_BIOACTIVE_START_UP_CHINI - Isolation, characterisation, activity and evaluation of natural and synthetic bioactive compounds. MUR (PRIN 2022 PNRR): P2022MWY3P - Old but Gold! Identification of molecular platforms for age-associated diseases to promote healthy and active aging.

* Presenting author. E-mail address: v.samukha@studenti.unimol.it (Vadym Samukha)

S3.P233 Exploring *Genista tridentata* leaves extract: novel insights for functional food in diabetes management

Inês Martins Laranjeira^{a,b,c,d,*}, Marlene Silva^e, Diana Fonseca-Rodrigues, Gonçalves, C.^f, Dias, A. C. P.^{c,d}, Pinto-Ribeiro, F.^{a,b}

^a Life and Health Sciences Research Institute (ICVS), School of Medicine, Campus de Gualtar, University of Minho, 4710-057 Braga, Portugal

^b ICVS/3B's - PT Government Associate Laboratory, Braga/Guimarães, Portugal

^c CITAB - Centre for the Research and Technology of Agro-Environmental and Biological Sciences, University of Trás-os-montes e Alto Douro, Portugal

^d Centre of Molecular and Environmental Biology (CBMA), Department of Biology, University of Minho, Braga, Portugal

^e Polytechnic Institute of Viana do Castelo, Viana do Castelo, Portugal

^f Department of Biology, University of Aveiro, 3810-193 Aveiro, Portugal

Genista tridentata (L.) Willk., a Leguminosae (Fabaceae) species native to the Iberian Peninsula, Morocco, Algeria, and Tunisia, boasts versatility in both culinary and traditional medicine (Laranjeira et al., 2023). Recognized for its properties, this plant is used in folk medicine and by populations for the management of several pathologies like, diabetes, hypertension, respiratory and digestive disorders (Coelho et al., 2011; Balanč et al., 2016). This study aimed to evaluate the in vivo antinociceptive effects of *G. tridentata* leaves (GtL) methanolic extracts in rats with experimental Type 2 Diabetes mellitus (T2DM). Using the Streptozotocin-Nicotinamide model (Masiello et al., 1998), male Wistar rats were induced with T2DM, divided into Sham (n = 14) and Diab (n = 38) groups. Diabetic rats were further divided into three experimental groups: (i) Diab treated with PBS (n = 14), (ii) GtL_100 treated with 100 mg/kg (n = 11), and (iii) GtL_300 treated with 300 mg/kg of GtL extract (n = 13), administered daily via gavage, for four weeks. Sham animals received PBS (n = 14). Weekly nociceptive assessments included mechanical and thermal hyperalgesia and allodynia. Post-euthanasia, internal organs underwent histopathological examination. T2DM induced thermal allodynia and thermal and mechanical hyperalgesia, which was decreased after *G. tridentata* treatments. Concomitantly, *G. tridentata* treatment ameliorated the histopathological changes in T2DM animals' internal organs. These findings suggest *G. tridentata* leaves may offer beneficial effects in addressing T2DM-related complications, potentially due to its antioxidant and anti-inflammatory properties, supporting its potential in developing nutraceutical products.

Keywords: *Genista tridentata*, *Pterospartum tridentatum*, Type 2 Diabetes, Medicinal Plants, nociception, Functional foods

References

- Laranjeira, I. M., Dias, A. C. P., & Pinto-Ribeiro, F. L. 2023. *Genista tridentata* Phytochemical Characterization and Biological Activities: A Systematic Review. *Biology*, 12 (11), 1387.
- Coelho, M.T., Gonçalves, J.C., Alves, V., Moldão-Martins, M. 2011. Antioxidant activity and phenolic content of extracts from different *Pterospartum tridentatum* populations growing in Portugal. *Procedia Food Sci.*, 1, 1454–1458.
- Balanč, B., Kalušević, A., Drvenica, I., Coelho, M.T., Djordjević, V., Alves, V.D., Sousa, I., Moldão-Martins, M., Rakic, V., Nedovic, V., et al. 2016. Calcium–alginate–inulin microbeads as carriers for aqueous carqueja extract. *J. Food Sci.*, 81, E65–E75.
- Masiello, P., Broca, C., Gross, R., Roye, M., Manteghetti, M., Hillaire-Buys, D., Novelli, M. & Ribes, G. 1998. Experimental NIDDM: development of a new model in adult rats administered streptozotocin and nicotinamide. *Diabetes*, 47 (2), 224–229.

* Presenting author. E-mail address: inesmartinslaranjeira@gmail.com (Inês Martins Laranjeira)

S3.P234 A formulation containing palmitoylethanolamide and phenolic compounds counteracts diabetes and related hepatic damage in mice: in vivo and in vitro evidence

Stefania Melini^{a,*}, Claudio Pirozzi^a, Adriano Lama^a, Federica Comella^a, Nicola Opallo^a, Filomena Del Piano^b, Giuseppina Mattace Raso^a, Rosaria Meli^a

^a Department of Pharmacy, School of Medicine, University of Naples Federico II, 80131 Naples, Italy

^b Department of Veterinary Medicine and Animal Production, University of Naples Federico II, 80137 Naples, Italy

Glucose- and lipotoxicity play a key role in the development of diabetes, namely diabetes-related-to- obesity (Michaelidou et al., 2023). The N-acyl ethanolamine palmitoylethanolamide (PEA) and some phenolic compounds, such as rutin and hydroxytyrosol (HT), have shown different metabolic, anti- inflammatory and antioxidant effects (Annunziata et al., 2020, Pirozzi et al., 2016, Muvhulawa et al., 2022). Here, we evaluated the beneficial activity of a formulation containing PEA co-micronized with rutin and associated with HT, named NORM3, in counteracting insulin resistance (IR) and hepatic metabolic alterations in high-fat diet (HFD)-fed mice.

Male C57Bl/6J mice received standard chow diet or HFD for 19 weeks; a HFD group administered NORM3 (PEA 10 mg/kg/die-Rutin 2 mg/kg/die, HT 0,5 mg/kg/die *per os*) from week 12 up to week 19.

First, NORM3 limited body weight of obese mice, after 4 weeks of treatment, and fat mass. HFD+NORM3 animals showed an improved insulin sensitivity, as shown by oral glucose and insulin tolerance tests and the reduced HOMA-IR index. In liver, morphological analysis showed that NORM3 reduced macro- and microvacuolar steatosis caused by HFD. Contextually, NORM3 treatment restored hepatic insulin signaling, activating AKT/PI3K pathway, and reduced the mRNAs of crucial enzymes of gluconeogenesis. Moreover, NORM3 counteracted hepatic lipid dysmetabolism caused by HFD, and associated inflammation and oxidative stress. In HepG2 cells, we proved the synergistic effect of the single components of NORM3 in reducing ROS production and inflammation and increasing antioxidant defense.

Our findings identify NORM3 as a potential nutritional approach in counteracting IR and related hepatic dysmetabolism due to diabetes.

Keywords: N-acyl ethanolamines, polyphenols, obesity, diabetes, glucose metabolism, oxidative stress, MAFLD

References

- Annunziata C, Lama A, Pirozzi C, Cavaliere G, Trinchese G, Di Guida F, Nitrato Izzo A, Cimmino F, Paciello O, De Biase D, Murru E, Banni S, Calignano A, Mollica MP, Mattace Raso G, Meli R. Palmitoylethanolamide counteracts hepatic metabolic inflexibility modulating mitochondrial function and efficiency in diet-induced obese mice. *FASEB J.* 2020 ;34(1):350-364.
- Michaelidou M, Pappachan JM, Jeeyavudeen MS. Management of diabetes: Current concepts. *World J Diabetes.* 2023;14(4):396-411.
- Muvhulawa N, Dlodla PV, Ziqubu K, Mthembu SXH, Mthiyane F, Nkambule BB, Mazibuko-Mbeje SE. Rutin ameliorates inflammation and improves metabolic function: A comprehensive analysis of scientific literature. *Pharmacol Res.* 2022;178:106163.
- Pirozzi C, Lama A, Simeoli R, Paciello O, Pagano TB, Mollica MP, Di Guida F, Russo R, Magliocca S, Canani RB, Raso GM, Calignano A, Meli R. Hydroxytyrosol prevents metabolic impairment reducing hepatic inflammation and restoring duodenal integrity in a rat model of NAFLD. *J Nutr Biochem.* 2016;30:108-15.

* Presenting author. E-mail address: stefania.melini@unina.it (Stefania Melini)

S3.P235 The *Vitis vinifera* cvs. callus *in vitro* culture extracts as a potential cosmetic antioxidant

Marta Sharafan^a, Magdalena Malinowska^{a,*}, Arnaud, Lanoue^c, Manon Ferrier^c, Florent Magot^c, Nathalie Giglioli-Guivarc'h^c, Halina Ekiert^b, Agnieszka Szopa^b, Elżbieta Sikora^a

^a Department of Organic Chemistry and Technology, Faculty of Chemical Engineering and Technology, Cracow University of Technology, 24 Warszawska st., 31-155 Kraków, Poland

^b Chair and Department of Pharmaceutical Botany, Medical College, Jagiellonian University, 9 Medyczna st., 30-688 Kraków, Poland

^c Université de Tours, EA 2106, Biomolécules et Biotechnologies Vegetales, UFR des 8 Sciences Pharmaceutiques, 31 Monge av., F37200 Tours, France

The innovative active ingredients of natural origin, especially the new antioxidants, are constantly in demand in the production of cosmetics (Oresajo et al., 2012).

Grapevine (*Vitis vinifera* L.) is one of the major fruit crop over the world. Several studies on *Vitis vinifera* extracts showed their strong antioxidant potential for which phenolic compounds are most likely responsible (Sharafan et al., 2023).

The aim of the study was to evaluate the antioxidant potential of *in vitro* callus extracts obtained from different grapevine cultivars (3 red cultivars: Marechal Foch, Regent, Rondo and 3 white cultivars: Chardonnay, Hibernat, Seyval Blanc). The cultures were maintained over 30-day growth cycles on two media: Murashige and Skoog (MS) and Schenk and Hildebrandt (SH) with various concentration of plant growth regulators: 6-benzylaminopurine (BA) and indole-3-butyric acid (IBA) or 1- naphthaleneacetic acid (NAA). The extracts were prepared using ultrasound mediated extraction, in 96% ethanol. The ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) and FRAP (ferric reducing antioxidant power) assays were used to determine the antioxidant potential of extracts.

The results of ABTS assays showed that all the extracts possessed relatively high antioxidant activity. The ABTS inhibition reached 51.38%. In the case of the FRAP assay, the ferric ion reducing activity reached up to 10.4%. The best results in ABTS and FRAP assays were obtained for cv. Hibernat maintained on medium 'W2' (MS, 1.5 ml/l BA and 0.2 ml/l NAA).

The obtained results indicate that *V. vinifera* cv. callus *in vitro* cultures could be proposed as valuable alternative of natural cosmetic ingredient that is custom-produced in a controlled environment.

Keywords: *Vitis vinifera*, vine grape, *in vitro* cultures, plant callus culture, antioxidant capacity

References

- Oresajo, C., Pillai, S., Manco, M., Yatskayer, M., McDaniel, D., 2012. Antioxidants and the skin: understanding formulation and efficacy. *Dermatol Ther.* 25(3):252-259.
- Sharafan, M., Malinowska, M.A., Kubicz, M., Kubica P., Gémin, M-P., Abdallah, C., Ferrier, M., Giglioli-Guivarc'h, N., Hano, C., Sikora, E., Lanoue, A., Szopa, A., 2023. Shoot cultures of *Vitis vinifera* (vine grape) Different Cultivars as a Promising Innovative Cosmetic Raw Material – The Phytochemical Profiling, Antioxidant Potential and Whitening Activity. *Molecules.* 28(19), 6868.
- Sharafan, M., Malinowska, M.A., Ekiert, H., Kwaśniak, B., Sikora, E., Szopa, A., 2023. *Vitis vinifera* (Vine Grape) as a Valuable Cosmetic Raw Material. *Pharmaceutics.* 15, 1372.

Acknowledgments

This research was funded by PHC Polonium 2022 joint research project between France and Poland (entitled 'Cosmetic potential of viticulture byproducts as novel functional ingredients for skin barrier recover) grant no. BPN/BFR/2022/1/00011 and project no. N42/DBS/000273.

* Presenting author. E-mail address: magdalena.malinowska@poczta.fm (Magdalena Malinowska)

S3.P236 Phytochemical analysis of *Echinacea* preparations from the pharmacies and herbal markets in Türkiye

Ayşe Esra Karadağ^a, Rengin Baydar^{a,b,*}

^a Department of Pharmacognosy, School of Pharmacy, Istanbul Medipol University, 34810 Istanbul, Türkiye

^b Institute of Health Sciences, Yeditepe University, 34755 Istanbul, Türkiye

Echinacea species are Asteraceae family plants containing mainly chlorogenic acid, cichoric acid, caffeic acid, echinacoside. Aerial parts, roots, and especially flowering tops of *Echinacea* sp. were used for treatment of recurrent respiratory tract infections and wounds, ethnobotanically. This study aims to evaluate the phytochemical quality (according to European Pharmacopoeia) of 5 different *Echinacea* containing commercial products (ECCP) available on pharmacies/herbal markets in Türkiye. Detailed information about ECCP planned to be analyzed is given in Table 1.

Table 1. Detailed information about 5 different *Echinacea* preparations used in this study

<i>Echinacea</i> preparations	Dosage form	Ingredients/Dosage (if available)
Sample-1	Capsule	265 mg <i>E. purpurea</i> (aerial parts) powder and 65 mg <i>Echinacea</i> root extract
Sample-2	Capsule	100 mg <i>E. purpurea</i> , zinc, vitamin C, β -glukan, <i>Rosa canina</i> L. / 2 capsules (daily recommended dosage)
Sample-3	Capsule	150 mg <i>E. purpurea</i> , β -glukan, vitamin C, zinc
Sample-4	Liquid	300 mg <i>Echinacea</i> extract, β -glukan, propolis, vitamin C, zinc
Sample-5	Powder	<i>Echinacea</i> spp. (roots) powder

Sample 1, 2, 3 and 4 was directly macerated with ethanol for 3 days. Sample-5 was purchased as dried root parts of the plant. Then, it was grinded and macerated with ethanol for 3 days. After maceration, extracts were filtered through Whatman No:1 filter paper and concentrated by using rotary-evaporator.

Standard compounds (cichoric acid, chlorogenic acid, caffeic acid, and echinacoside) and ECCP extracts were dissolved in ethanol (10 mg/mL) and filtered via 0.22 μ m membrane filters prior to HPLC analysis. An Agilent 1100 HPLC-DAD system was used to conduct HPLC studies. C18 column (100 x 4.6mm, 10 μ m) and A: water:formic acid (100:0.1, v/v), B: acetonitrile were used as stationary and mobile phases, respectively. The gradient elution in HPLC was applied as 10-78% B (0-18 min), 90-10% B (18-21 min). Flow rate was 1.2 mL/min and injection volume was 5 μ L with a temperature of 26°C. The detection was performed at 330 nm. The reference standards were prepared and analyzed at the concentrations of 10, 20, 30, 40 and 50 μ g/mL. The quantitative analysis of ECCP extracts was calculated by using the calibration curve of reference standards. Results of the quantitative analysis of ECCP extracts are given in Table 2. Chlorogenic acid was the only standard detected in Sample-1 as 0.1% (w/w). Sample-2 and Sample-4 were lack of any standard compounds. Cichoric acid, chlorogenic acid and caffeic acid were detected in Sample-3 at the concentration of 0.4%, 0.6% and 0.2% (w/w), respectively. Sample-5 revealed that it contains cichoric acid, chlorogenic acid, echinacoside and caffeic acid at the concentration of 0.1%, 0.6%, 0.2% and 0.3% (w/w), respectively. All results were evaluated according to European Pharmacopoeia. To the best of our knowledge, in this study, pharmacy and herbal market's *Echinacea* supplements were tested together for their phytochemical content for the first time.

Table 2. Phytochemical contents of the samples evaluated by HPLC method (w/w)

	Cichoric acid	Chlorogenic acid	Echinacoside	Caffeic acid
Sample-1	Not detected	Not detected	Not detected	0.1%
Sample-2	Not detected	Not detected	Not detected	Not detected
Sample-3	0.4%	0.6%	Not detected	0.2%
Sample-4	Not detected	Not detected	Not detected	Not detected
Sample-5	0.1%	0.6%	0.2%	0.3%

Keywords: *Echinacea*, herbal supplement, HPLC, pharmacy

* Presenting author. E-mail address: eczrenginbaydar@gmail.com (Rengin Baydar)

S3.P237 Untargeted metabolomics analyses to identify a new sweet compound released during post- fermentation maceration of wine

Marie Le Scanff^{a,*}, Laurence Marcourt^{b,c}, Adriano Rutz^{b,c}, Warren Albertin^a, Jean-Luc Wolfender^{b,c}, Axel Marchal^a

^a Univ. Bordeaux, Bordeaux INP, INRAE, BSA, OENO, UMR 1366, ISVV, F-33140 Villenave d'Ornon, France

^b School of Pharmaceutical Sciences, University of Geneva, Centre Médical Universitaire (CMU), Geneva, Switzerland

^c Institute of Pharmaceutical Sciences of Western Switzerland (ISPSO), University of Geneva, Centre Médical Universitaire (CMU), Geneva, Switzerland

The sensory quality of a wine is mainly based on its aroma and its flavor. In particular, gustatory balance of red wines relies on sourness, bitterness and sweetness. Even if certain compounds were already identified as contributing to sweetness, certain taste modifications remain largely unexplained (Cretin et al., 2019, Fayad et al., 2021). Some empirical observations combined with sensory analyses have shown that an increase of wine sweetness occurs during post-fermentation maceration. This step is a key stage of red winemaking during which the juice is left in contact with the marc, that contains the solid parts of the grape (seeds, skins and sometimes stems). The present work aimed to identify a new taste-active compound that contributes to this gain of sweetness.

Recent developments have highlighted the interest of untargeted metabolomic analysis for natural product research (Hubert et al., 2017, Wolfender et al., 2019). Using similar tools, an original approach using UHPLC-HRMS/MS has been developed to discover new sweet molecules released during post-fermentation maceration. Among different markers highlighted, one compound was selected and considered for a targeted purification by various separative techniques (SPE, CPC and HPLC-preparative). It was unambiguously identified by NMR as N6-succinyladenosine and was reported for the first time in wine. It was quantitated at an average concentration of 3.16 mg/L in 85 red wines. Furthermore, sensory analysis revealed its pronounced sweetness. In addition to discovering a new sweet compound in wine, this study offers new tools for studying taste-active compounds in natural matrices.

Keywords: Untargeted metabolomics analysis, UHPLC-HRMS/MS, wine, sweetness, taste

References

- Cretin, B., Waffo-Teguo, P., Dubourdieu, D., Marchal, A., 2019. Taste-guided isolation of sweet-tasting compounds from grape seeds, structural elucidation and identification in wines. *Food Chemistry* 272, 388–395.
- Fayad, S., Le Scanff, M., Waffo-Teguo, P., Marchal, A., 2021. Understanding sweetness of dry wines: First evidence of astilbin isomers in red wines and quantitation in a one-century range of vintages. *Food Chemistry* 352, 129293.
- Hubert, J., Nuzillard, J.-M., Renault, J.-H., 2017. Dereplication strategies in natural product research: How many tools and methodologies behind the same concept? *Phytochem Rev.* 16, 55–95.
- Wolfender, J.-L., Nuzillard, J.-M., van der Hooft, J.J.J., Renault, J.-H., Bertrand, S., 2019. Accelerating Metabolite Identification in Natural Product Research: Toward an Ideal Combination of Liquid Chromatography–High-Resolution Tandem Mass Spectrometry and NMR Profiling, in Silico Databases, and Chemometrics. *Analytical Chemistry*. 91, 704– 742.

* Presenting author. E-mail address: marie.le-scanff@u-bordeaux.fr (Marie Le Scanff)

S3.P238 Do you know what's really hiding in popular herbal supplements? – evaluation of products containing *Cistus* leaves and extracts

Justyna Baranowska*, Marta Karolina Komoń, Małgorzata Kołtun-Jasion, Anna Karolina Kiss

Department of Pharmaceutical Biology, Faculty of Pharmacy, Medical University of Warsaw, 1 Banacha Str., 02-097 Warsaw, Poland

Cistus plants, also known as rockroses, are Mediterranean-native shrubs belonging to the Cistaceae family. Their flowers, leaves, and aerial parts have been used in traditional medicine to treat many conditions, including the common cold, gastric ulcers, wounds, and many inflammatory diseases (Barrajón-Catalán et al., 2016). Nowadays, products containing those species are becoming increasingly popular due to their somehow proven efficacy in reducing the average duration and severity of upper respiratory tract infections (Kalus et al., 2009; Guzelmeric et al., 2023). Manufacturers supply and market a wide range of *Cistus* dietary supplements, including infusion teas, capsules, and tablets. In order to evaluate their quality, we tested sixteen of those products that are available on the Polish market, using HPTLC methods to compare their fingerprint profiles to authentic standards. Additionally, we used HPLC-DAD-MS/MS method to in deep analyze their qualitative profile. Five out of sixteen supplements were found to be adulterated. The majority of the counterfeit preparations were in the form of tablet and capsules. Instead of *Cistus* extracts, products probably contained buckwheat, green tea, or oak extracts. In the other products, we identified typical *Cistus* phytochemicals, including characteristic myricetin derivatives, tiliroside, punicalagins, and dimeric prodelphinidin. The investigation showed that, despite what the producers claim, several items on the market do not contain *Cistus*. This study also pointed out the significance of controlling the raw herbal material used in the manufactures.

Keywords: *Cistus*, Cistaceae, supplements, HPTLC, LC-MS

References

- Barrajón-Catalán, E., Tomás-Menor, L., Morales-Soto, A., Bruñá, N. M., López, D. S., Segura-Carretero, A., Micol, V. 2016. Rockroses (*Cistus* sp.) Oils. Essential Oils in Food Preservation, Flavor and Safety, 649–658.
- Guzelmeric, E., Reis, R., Sen, N.B., Celik, C., Özhan, Y., Petrikaite, V., Sipahi, H., Aydin, A., Yesilada, E. 2023 Insights into the Anti-inflammatory, Analgesic, and Anticancer Potentials of the Standardized Extracts From Three *Cistus* L. Species. Journal of Herbal Medicine 41.
- Kalus, U., Grigorov, A., Kadecki, O., Jansen, J.-P., Kiesewetter, H., Radtke, H. 2009 *Cistus incanus* (CYSTUS052) for treating patients with infection of the upper respiratory tract. Antiviral Research, 84(3), 267–271.

* Presenting author. E-mail address: justyna.baranowska222@gmail.com (Justyna Baranowska)

S3.P239 A smartphone-based methodology for analyte quantification and antioxidant activity determination directly in emulsions

Vassiliki Gardiki^a, Georgia Eleni Tsotsou^{a,b}, Anastasia Kyriaki Tsara^a, Georgia Bakara^a, Spyros Papageorgiou^{a,b}, Athanasia Varvaresou^{a,b*}

^a Department of Biomedical Sciences, University of West Attica, Egaleo 122 43, Greece

^b Laboratory of Chemistry, Biochemistry and Cosmetic Science, Department of Biomedical Sciences, University of West Attica, Egaleo 122 43, Greece

This work investigated the width of applicability of a methodology for the quantification of analytes directly into emulsions with oil internal phase and aqueous external phase (O/W). The methodology relied on a specific reaction (enzyme-catalysed, or not) between the analyte of interest and a suitable chromogen towards a coloured product. Colour intensity was captured by a mobile phone camera and analysed by the ImageJ software (Schneider et al., 2012) into its R, G and B components (Fig. 1). A linear relationship was observed between at least one of the R, G or B components or their average, and analyte concentration in O/W emulsions for the investigated analytes (glucose (Tsotsou and Tsara, 2023), urea (Tsotsou, 2024), and total iron). Satisfactory coefficients of correlation (R^2) (≥ 0.984), and relative error values ($\leq \pm 15.9\%$), as well as wide linear dynamic ranges were determined. Interferences were also reported.



Fig. 1. Setup for the proposed extraction-free quantification methodology

The robustness of the methodology was demonstrated in terms of camera specifications, emulsion properties (pH, viscosity, conductivity, transparency), ambient temperature and light conditions. Preliminary results suggested its applicability to antioxidant capacity measurement by the FRAP assay, directly into an O/W emulsion containing Unisoorth EG-28[®] (R^2 of calibration curve: 0.989). Unisoorth EG-28[®] named according to INCI system as water, propyl gallate, gallyl glucoside, epigallocatechin gallatyl glucoside, is an antioxidant cosmetic active ingredient complex that is composed of gallic acid glucoside, of epigallocatechin gallate glucoside, and propyl gallate. This methodology is proposed as an extraction-free, green procedure, suitable even for minimally equipped settings, to serve quality control needs of emulsions. It may also prove useful for quantifying antioxidant capacity of natural extracts into emulsified matrices.

Keywords: emulsion, smartphone-based analysis, digital image colourimetry, urea/glucose/phenolics/total iron analysis, antioxidant, cosmetic ingredients

References

- Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis, Nat. Methods 9, 671-675.
- Tsotsou, G.E., Tsara, A.K., 2023. Extraction-free, in situ Analysis of Glucose in Cosmetic Formulations based on Digital Image Colourimetry by Smartphone. J. Cosm. Sci. 74, 289-300.
- Tsotsou, G.E., 2024. Extraction-free analysis in cosmetics by digital image colorimetry, illustrated by the quantification of urea. Heliyon 10, e25503.

* Presenting author. E-mail address: avarvares@uniwa.gr (Athanasia Varvaresou)

S3.P240 Phytochemical analysis of *Laurus nobilis* L. leaves essential oil: rationalization of antioxidant and anti-inflammatory activities by combined *in vitro* and *in silico* approach

Francesca Fantasma^{a,*}, Vadym Samukha^a, Michela Aliberti^b, Ester Colarusso^b, Gabriella Saviano^a, Vincenzo De Felice^a, Gianluigi Lauro^b, Agostino Casapullo^b, Maria Giovanna Chini^a, Giuseppe Bifulco^b, Maria Iorizzi^a

^a Department of Biosciences and Territory, University of Molise, C.da Fonte Lappone snc, 86090 Isernia, Italy

^b Department of Pharmacy, University of Salerno, via Giovanni Paolo II 132, 84084 Salerno, Italy

This research focuses on natural product chemistry, explicitly targeting the structural characterization of secondary metabolites derived from plant matrices. These secondary metabolites have been assessed as prospective “new drugs” due to their intriguing pharmacological properties demonstrated in both *in vitro* and *in vivo* studies (Newman and Cragg, 2020). These properties include anti-inflammatory, antioxidant, antitumor, antiviral, and antibacterial activities.

The evergreen Mediterranean shrub *Laurus nobilis* L., commonly known as bay laurel, has a long history of traditional use in cuisines and folk medicine due to its health-promoting properties. These properties are now being scientifically elucidated through various biological activities.

Samples of *L. nobilis* were collected from three different regions in central-southern Italy. The leaves were subjected to hydro-distillation, and essential oils (EOs) were analyzed using gas chromatography (GC/FID) for quantification of volatile compounds and gas chromatography/mass spectroscopy (GC/MS) for identification. The multivariate analysis of the main components establishes the correlations of the different chemical compounds with the corresponding antioxidant and anti-inflammatory activities.

In vitro, antioxidant activity was then evaluated by different methods (DPPH, ABTS, FRAP assay) and anti-inflammatory activity through inhibition of selected enzymes involved in eicosanoid biosynthesis through the three metabolic routes: cyclooxygenase (COX), 5-lipoxygenase (5-LO), cytochrome P450 (CYP450) pathways (Sun et al., 2021). In more detail, here, the simultaneous targeting of different, i.e., COXs and Soluble Epoxide Hydrolase (sEH) of the EOs was evaluated. Furthermore, the *in silico* studies were applied to rationalize the biological activities in inflammatory events observed for the main secondary metabolites of the different varieties of *L. nobilis* EOs at the molecular level.

Keywords: *Laurus nobilis* L., phytochemical analysis, essential oil, metabolomics, natural product

References

- Newman, D.J., Cragg, G.M., 2020. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J. Nat. Prod.* 83, 3, 770–803.
- Sun, c., Zhang, X., Morisseau, C., Hwang, S.H., Zhang, Z., Hammock, B.D., 2021. Discovery of Soluble Epoxide Hydrolase Inhibitors from Chemical Synthesis and Natural Products. *J. Med. Chem.* 64, 1, 184–215.

Acknowledgments

The research activity was supported by Ministero dell'Università e della Ricerca (MUR)—(PRIN 2022 PNRR project), grant number “P2022MWY3P—Old but Gold! Identification of molecular platforms for age-associated diseases to promote healthy and active aging”.

* Presenting author. E-mail address: fantasma@unimol.it (Francesca Fantasma)

S3.P241 A study on the biological activities of loquat leaf extract using various extraction methods

Jung-Wook Kang^{a,*}, Suk-Min Son^b, In-Chul Lee^c

^a College of Fusion and Convergence, Seowon University, Cheongju, Republic of Korea

^b Department of Food Engineering, Seowon University, Cheongju, Republic of Korea

^c Department of Bio-Cosmetic Science, Seowon university, Cheongju, Republic of Korea

Loquat (*Eriobotrya japonica*) is a member of the Rosaceae family and the Pomoideas subfamily, cultivated in East Asia. The aim of this study was to examine the physicochemical properties of extracts obtained through different extraction methods of *Eriobotrya japonica* leaf extract. We used 70% ethanol extraction (EE), infrared-assisted extraction (IAE), and ultrasound-assisted extraction (UAE). To determine the antioxidant and anti-aging effects of *Eriobotrya japonica* leaf extract, we investigated various biological activities induced by different extracts. Extraction using IAE resulted in the highest chlorogenic acid content compared to other methods, as analyzed by high-performance liquid chromatography. It was observed that the IAE-associated extract exhibited higher antioxidant activities than the others, as determined by DPPH and ABTS assays. Significant inhibitory effects on elastase and collagenase activities were found using the IAE method, which might be attributed to its anti-wrinkle effects. These results suggest that selecting the extraction method is crucial in obtaining loquat leaf extract with specific bioactive properties, which can be useful in functional materials.

Keywords: loquat, *Eriobotrya japonica*, antioxidant, anti-wrinkle, cosmetic

References

Zhu X et al., Traditional uses, phytochemistry, pharmacology, and toxicity of *Eriobotrya japonica* leaves: a summary, *Journal of Ethnopharmacology*, 298:115566, 2022.

Altemimi A, et al., Phytochemicals: extraction, isolation, and identification of bioactive compounds from plant extracts. *Plants*, 6:42, 2017.

* Presenting author. E-mail address: jwkkang@gmail.com (Jung-Wook Kang)

S3.P242 Study on modulation of inflammation expression by *Brassica oleracea* L. leaves extract

Jung-Wook Kang^{a,*}, Dan-Hee Yoo^a, Hyo-Min Kim^b, Ji-Won Hur^b, Kyeong-A Kwon^c, Kyung-Eun Lee^c, Min-Yeong Kim^c, Yong-Jun Kwon^c, Su-Bin Yu^c, Ju-Hyun Park^c, In-Chul Lee^d

^a College of Fusion and Convergence, Seowon University, Cheongju, Republic of Korea

^b College of Pharmacy, Kyungpook National University, Daegu, Republic of Korea

^c Department of Cosmetic Engineering, Seowon University, Cheongju, Chungbuk, Korea

^d Department of Bio-Cosmetic Science, Seowon university, Cheongju, Republic of Korea

Brassica oleracea L., commonly known as cabbage, belongs to the plant family Brassicaceae. It holds significant nutritional and medicinal importance due to its rich biochemical and phytochemical composition. In this study, the antioxidant and anti-inflammatory activities were investigated to confirm the potential of *Brassica oleracea* L leaves extract. as a natural cosmetic material. The total polyphenol content of hot water extract (KRD) and 70% ethanol extract (KRE) was measured, resulting in 124 mg TAE/100 g and 144 mg TAE/100 g, respectively. To affect the antioxidant activity of various extracts, we confirmed their ability to donate electron and scavenge ABTS radical. It was observed that the activity of KRD and KRE increased in a dose-dependent manner. The inhibitory effects on inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) were assessed through western blot and reverse transcription-PCR. Treatment with KRD and KRE confirmed the concentration-dependent suppression of iNOS and COX-2 protein and mRNA expression. These findings suggest that kailan could serve as a promising natural cosmetic material.

Keywords: *Brassica oleracea*, antioxidant, anti-inflammation, nitric oxide, cosmetic

References

Folin O, Dennis W. On phosphotungstic-phosphomolybdic compounds as color reagent. J. Biol Chem., 12, pp.239-243, (1912).

Blois MS. Antioxidant determination by the use of a stable free radical. Nature., 26, pp.1199-1120, (1985).

Pellegrin N, Roberta R, Min Y, Catherine RE. Screening of dietary carotenoids and carotenoid-rich fruit extracts for antioxidant activities applying 2,2'-azinobis(3-ethylbenzo-thiazoline-6-sulfonic acid) radical cation decolorization assay. Method Enzymol., 299, pp.379-389, (1998).

Carmichael J, DeGraff WG, Gazdar AF, Minna JD, Mitchell JB. Evaluation of a tetrazolium based semiautomated colorimetric assay: assessment of chemosensitivity testing. Cancer Res., 47, pp.936-942, (1987).

Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrate, nitrite, and [15N] nitrate in biological fluids. Analytical Biochem., 126, pp.131-138, (1982).

* Presenting author. E-mail address: jwkkang@gmail.com (Jung-Wook Kang)

S3.P243 Glucosinolates in *Brassica incana* subsp. *raimondoi* (Brassicaceae) leaf extract: composition, content, and anti-inflammatory activity

Giuseppe Antonio Malfa^{a,b,*}, Claudia Di Giacomo^{a,b}, Simone Ronsisvalle^{a,b}, Simone Bianchi^{a,b}, Federica Sipala^a, Rosaria Acquaviva^{a,b}

^a Department of Drug and Health Science, University of Catania, Catania, Italy

^b Research Centre on Nutraceuticals and Health Products (CERNUT), University of Catania, Catania, Italy

Brassica incana subsp. *raimondoi* (Brassicaceae) is a taxon endemic to Sicily (Italy), restricted to the specific geographic area of Castelmola (Messina). It grows at an altitude of about 500 m a.s.l. on abrupt limestone cliffs overlooking the Ionian Sea and is typified by various distinctive traits. *B. raimondoi* belongs to the *B. oleracea* cytodeme, a group of species characterized by the same diploid genome (2n=18) and which attracts great attention for the genetic improvement of the horticultural varieties of *B. oleracea* in terms of resistance to biotic and abiotic stresses and phytochemical compounds such as the glucosinolates (Malfa et al., 2020). Initially described at a specific level, it was not long ago assigned to the rank of subspecies (Malfa et al., 2020; 2022). As part of our ongoing research on Sicilian Brassicaceae taxa, in this study the methanolic leaf extract of *B. raimondoi*, analyzed by UPLC-MS/MS, evidenced the presence of one aromatic, four indolic, and eleven aliphatic glucosinolates. Among these, gluconapin, glucoraphanin, glucoraphasatin, and sinigrin represented more than 50% of the total glucosinolate content.



Fig. 1. *Brassica incana* subsp. *raimondoi*

Since we found that the phytochemicals in the methanolic extract had no cytotoxic effects on HDF cells, we tested the extract's anti-inflammatory effects on this cell line by inducing inflammation with IL-1 β exposure. Quantitative analysis of ROS and nitrite/nitrate levels, which increase as hallmarks of inflammation in cells, showed that the extract can counteract the harmful effects of the pro-inflammatory cytokine in a dose-dependent manner starting from a concentration of 50 μ g/mL.

Keywords: *Brassica*, glucosinolates, chemotaxonomy, oxidative stress, inflammation

References

- Malfa G.A., Acquaviva R., Bucchini A.A.E., Ragusa S., Raimondo F.M., Spadaro V., 2020. The Sicilian wild cabbages as biological resources: Taxonomic update and a review on chemical constituents and biological activities. *Flora Mediterr.* 30, 245–260.
- Malfa, G. A., Pappalardo, F., Miceli, N., Taviano, M. F., Ronsisvalle, S., Tomasello, B., Bianchi, S., Davi, F., Spadaro, V., Acquaviva, R., 2023. Chemical, Antioxidant and Biological Studies of *Brassica incana* subsp. *raimondoi* (Brassicaceae) Leaf Extract. *Molecules*, 28, 1254.

* Presenting author. E-mail address: giuseppe.malfa@unict.it (Giuseppe Antonio Malfa)

S3.P244 *Ex vivo* and clinical evaluation of efficacy and mode of action of an *Anogeissus leiocarpus* and retinol-containing eye cream

Rocky Graziose*, Martin Keh, Kristine Schmalenberg, Nadine Pernodet, Claude Saliou

The Estee Lauder Companies

In an *ex vivo* study, the phytochemistry and bioactivity of *Anogeissus leiocarpus* bark, alone and when combined with retinol were investigated. Additionally, the clinical impact of a novel anti-aging eye cream with these ingredients was assessed. High-Performance Thin Layer Chromatography (HPTLC) Phytochemical scouting analysis of an ethanolic extract of *A. leiocarpus* bark revealed the presence of 10-12 unidentified flavonoids, several unidentified saponins and two compounds tentatively identified as Caffeic and Ellagic acid. Previously excised human skin (*ex vivo*) showed a marked increase in extracellular matrix components (Collagen, Elastin and Fibrillin) when individually treated with *A. leiocarpus* extract and Retinol. When the two ingredients were combined in a treatment, extracellular matrix components showed a significant improvement when compared to the effect of either ingredient alone. Separately, the effects of a novel anti-aging eye cream containing proprietary levels of retinol and *A. leiocarpus* were assessed via clinical trial involving 29 female subjects (aged 35 to 64, average age 57) with mild-to-moderate age-related eye conditions. After 12 weeks of usage, the cream significantly reduced wrinkles, fine lines, puffiness, dark circles, as assessed by expert grading. The study underscores the effectiveness of the cream in addressing age-related eye concerns, which can be linked to the action of its key ingredients.

Keywords: *Anogeissus leiocarpus*, phytochemical scouting analysis, retinol, clinical test, anti-aging

* Presenting author. E-mail address: rgrazios@estee.com (Rocky Graziose)

S3.P245 Targeted Isolation of natural products that modify sweet taste perception through metabolomics, molecular networking and sensory evaluation of *Symplocos* spp. (Symplocaceae)

Qingxi Su, Lynn Ni, Bonnie Bao, Feng Qiu, Yanpeng Hou, Jing Li, Jeanmarie Carr, Diana Klaser Cheng*

International Flavors and Fragrances, R&D Union Beach, New Jersey 07735 USA

As health consciousness rises, sugar reduction becomes a trend. Taste is paramount and achieving the right balance between health and taste is a challenge. Modifying the taste profile of high intensity sweeteners to mimic that of sugar is one solution and taste modulators have been identified from natural sources. Leaves of *Symplocos* spp. have sweet taste and were partly attributed to the presence of dihydrochalcone glucosides (Tanaka et al., 1980; Kinghorn et al. 2010). These dihydrochalcones were approved by Flavor & Extracts Manufacturers Association (FEMA) for use as flavoring substances (Cohen et al., 2022; Smith et al., 2011).

In our research, we found that *Symplocos* spp. contain additional compounds reported to have sweetness perception modifying function. Six extracts from *Symplocos* species were evaluated by a sensory panel, and one species showed more sweetness intensity. To identify molecules responsible for the variation of sensory quality, we performed molecular network and metabolomics analysis. With molecular network analysis, we investigated three classes of compounds associated with sweet modulating properties: dihydrochalcones, megastigmanes and flavonoids. Through differential analysis we identified a fourth class of compounds present at higher levels in the extract with a higher rating of sweetness. These compounds were selected for isolation to verify their taste function.




Fig. 1. PCA loadings plot of LCMS ions in negative mode. Dots in blue represent compounds unique to our target species.

Keywords: *Symplocos*, Symplocaceae, sweet, dihydrochalcones, megastigmanes

References

- Cohen, S.M., et al. 2022 GRAS Flavoring Substances 30.
- Kinghorn, A.D., et al., 2010. Natural Products as Sweeteners and Sweetness Modifiers. In: Comprehensive Natural Products II Chemistry and Biology., eds Mander, L. and Liu, H.W. Elsevier, London publishers. 3, 269–315.
- Smith, R.L., et al. 2011. GRAS Flavoring Substances 25. Food Technology, 65, 44-75.
- Tanaka, T., et al. 1980. Dihydrochalcone-glucosides as Sweet Principles in *Symplocos* ssp. Planta Medica. 40, 81-83.

* Presenting author. E-mail address: Diana.KlaserCheng@iff.com (Diana Klaser Cheng)

S3.P247 Investigation of the Greek microbial diversity for the discovery of novel antiaging products

Konstantinos Gaitanis^{a,*}, Nikolaos Tsafantakis^a, Eirini Gkogkou^b, Despoina D. Gianniou^b, Stavroula Kaili^c, Paris Laskaris^c, Dimitris Hatzinikolaou^c, Ioannis P. Trougkos^b, Amalia D. Karagouni^c, Nikolas Fokialakis^a

^a Section of Pharmacognosy & Chemistry of Natural Products, Department of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou 15771, Athens, Greece

^b Section of Cell Biology and Biophysics, Department of Biology, Panepistimiopolis Zografou 15784, Athens, Greece

^c Section of Botany, Department of Biology, Panepistimiopolis Zografou 15784, Athens, Greece

Greek ecosystems present a rich yet under-explored area for investigating Actinobacteria, well-known for their production of bioactive compounds. To this purpose, the “AntiAging” project sought to uncover novel, promising natural compounds with anti-aging properties, which could be incorporated into cosmeceutical/nutraceutical formulations. In total, 1000 isolates originating from the Greek natural environment and belonging to the ATHUBA collection (Athens University Bacterial & Archaea Culture Collection)—some originating from unique environments (caverns, volcanoes, thermal springs, etc.)—were investigated. A unique library of ~2000 extracts (EtOAc and MeOH/H₂O) was generated, and the tyrosinase and elastase inhibition activity was evaluated via enzymatic assays. The top 3 microbial strains, belonging to *Amycolatopsis* sp. and *Streptomyces* sp. families, combining a high production of secondary metabolites and bioactivity on both cell-free and cell-based assays, were grown using four different cultivation media and subjected to enzymatic assays. LCMS based metabolomics was then applied to detect significant differences among strains and fermentation conditions. One selected strain was then subjected to a scale up process (12L) and its EtOAc extract was fractionated. A customized bio-guided workflow was then applied, combining a high-throughput dereplication and molecular networking method (UPLC-HRMS). Purification of the targeted components led to the isolation of several compounds and structure elucidation was achieved using 1D and 2D NMR combined to HRESIMS. The latter results and the plethora of identified molecules show a hidden yet powerful potential of harnessing Greek microbial wealth in the context of the science of anti-aging.

Keywords: *Actinobacteria*, extract library, LCMS, metabolomics, NMR

* Presenting author. E-mail address: gaitanisk@pharm.uoa.gr (Konstantinos Gaitanis)

S3.P248 EthnoHERBS: Conservation of european biodiversity through exploitation of traditional herbal knowledge for the development of innovative products

Evanthia Dina^{a,*}, Nikolaos Fokialakis^a, Vladimir Dimitrov^b, Zora Dajic^c, Anna Rita Bilia^d, Katarina Šavikin^e, Alberto Dias^{f,l}, Andreas Afantitis^g, Aikaterini Termentzi^h, Juan Ramón Muñoz Montañóⁱ, Paolo Allegrini^j, Konstantinos Gardikis^k, Mirko Oluški^m, Hristina Hristovaⁿ, Nektarios Aligiannis^a

^a Department of Pharmacognosy and Natural Products Chemistry, Faculty of Pharmacy, National and Kapodistrian University of Athens, Greece

^b Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Sofia, Bulgaria

^c Faculty of Agriculture, University of Belgrade, Serbia

^d Department of Chemistry "Ugo Schiff", University of Florence, Italy

^e Institute for Medicinal Plants Research "Dr Josif Pančić", Belgrade, Serbia

^f Department of Biology and CITAB-UM research Center, University of Minho, Portugal

^g NOVAM, NTE TYRAS Cyprus

^h Department of Pesticides Control and Phytopharmacy, Benaki Phytopathological Institute, Athens, Greece

ⁱ Zurko Research SL, Madrid, Spain

^j Indena SpA, Milan, Italy

^k Apivita, Industrial park of Markopoulo, Athens, Greece

^l TecMinho, Campus de Azurém, University of Minho, Guimaraes, Portugal

^m MACVAL, Novi Sad, Serbia

ⁿ VenusRoses Labsolutions, Sofia, Bulgaria

EthnoHERBS project aims to record and evaluate information on Southeast European traditional knowledge, explore in a high-throughput manner the biodiversity of Balkan Peninsula flora and elaborate cutting-edge technologies in Natural Products chemistry to discover and develop innovative cosmeceutical products against skin disorders. Based on ethnobotanical/ethnopharmacological studies and ancient-medieval manuscripts, 1289 plant species have been identified for their traditional use against skin ailments (Tsioutsiou et al., 2022). The evaluation of the aforementioned data resulted in the selection of 240 plant species, that were collected according to good botanical practices and laws. Moreover, the selected species investigated through in silico procedure which was used in combination with biodiversity data to select the most promising herbs for further phytochemical analysis. Furthermore, the selected plant material was extracted in lab-scale with Ultrasound Assisted Extraction using dichloromethane, methanol and methanol:water 50:50, successively and the generated extracts (*ca.* 720) were forwarded for the investigation of their phytochemical profile was performed using chromatometric (TFC, TPC) and analytical HPTLC, HPLC-DAD, and NMR methodologies and cell- free bioassays for evaluation of their biological properties (antioxidant, enzymes inhibition). Based on the chemical profiles and preliminary biological results *ca.* 100 extracts have been selected for further investigation for the isolation of bioactive compounds. The phytochemical characterization of *Momordica charantia*, *Cistus creticus* subsp. *eriocephalus* (Viv.) Greuter et Burdet and *Juglans regia* extracts, coupled with their biological evaluation on skin enzymes and on skin cell lines, prospects for the discovery of bioactive natural compounds for the treatment of skin diseases.

Keywords: ethnoherbs, skin diseases, enzyme inhibitors

References

Tsioutsiou, E.E., Amountzias, V., Vontzalidou, A., Dina, E., Stevanović, Z.D., Cheilari, A., Aligiannis, N. 2022. Medicinal Plants Used Traditionally for Skin Related Problems in the South Balkan and East Mediterranean Region—A Review. *Front. Pharmacol.* 13:936047. doi: 10.3389/fphar.2022.936047.

Acknowledgments

This research has been funded under the European H2020-MSCA-RISE-2018 (ID 823973) project "EthnoHERBS - Conservation of European Biodiversity through Exploitation of Traditional Herbal Knowledge for the Development of Innovative Products".

* Presenting author. E-mail address: euantina@pharm.uoa.gr (Evanthia Dina)

S3.P249 *Artocarpus thailandicus* heartwood extract extends lifespan and promotes healthy aging behaviour in *Caenorhabditis elegans*

Chamaiphorn Wongwan^{a,d}, Panthakarn Rangsinth^{b,c}, Nilubon Sornkaew^{a,d}, Wudtipong Vongthip^{a,d}, Sunita Nilkhet^{a,b}, Chanin Sillapachaiyaporn^{a,d}, Anchalee Prasansuklab^{d,e}, Siriporn Chuchawankul^b, Tewin Tencomnao^{a,d,*}

^a Department of Clinical Chemistry, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^b Department of Transfusion Medicine and Clinical Microbiology, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

^c Department of Pharmacology and Pharmacy, The University of Hong Kong, Pokfulam, Hong Kong SAR, China

^d Natural Products for Neuroprotection and Anti-ageing Research Unit, Chulalongkorn University, Bangkok 10330, Thailand

^e College of Public Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand

Artocarpus thailandicus (AT) and its primary bioactive compound (oxyresveratrol) were believed to possess a high antioxidant capacity. However, the beneficial effect and mechanism of AT and oxyresveratrol on healthy aging phenotype and oxidative stress *in vivo* studies have never been investigated. We aimed to study the effects and mechanisms of AT and oxyresveratrol on healthy aging phenotype and oxidative stress resistance. The total phenolic and flavonoid contents were determined. Then, the ROS scavenging activity of these extracts was tested by DPPH and ABTS assays. To study healthy aging phenotype, survival rate under oxidative stress, lifespan assay, egg laying, and pharyngeal pumping rate were performed. The molecular mechanism affected-healthy aging phenotype was investigated by measuring the expression of HSP16.2, *sod-3*, *gst-4*, and nuclear translocation of DAF-16 and SKN-1 has been studied. Additionally, the oxyresveratrol level was quantified by HPLC, and NMR was employed to identify the phytochemicals in AT. Its ethanol crude extract (ATE) possesses a strong scavenging activity against DPPH and ABTS free radicals. ATE and oxyresveratrol improved the pharyngeal pumping rate including attenuating ROS accumulation of juglone-induced oxidative stress. Moreover, HSP16.2, an oxidative stress resistance biomarker, was decreased. In addition, DAF-16 and SKN-1 nuclear translocation was obvious, stimulating *sod-3* and *gst-4* gene expression. Moreover, the major phytochemical constituent of ATE can exhibit powerful antioxidant activity. In conclusion, our study exhibited a novel mechanism of ATE and oxyresveratrol against oxidative stress. Both ATE and oxyresveratrol have the potential to serve as dietary supplements to promote healthy aging.

Keywords: *Artocarpus thailandicus*, oxyresveratrol, *C. elegans*, oxidative stress

* Presenting author. E-mail address: tewin.t@chula.ac.th (Tewin Tencomnao)

S3.P250 Utilization of ethnobotanical data to discover potent inhibitors of skin related enzymes through the application of in silico based methodology

Antigoni Cheilari^{a,*}, Panagiotis Lagarias^b, Antonios Drakopoulos^b, George Lambrinidis^c, Andreas Afantitis^b, Nektarios Aligiannis^a

^a Department of Pharmacognosy and Natural Products Chemistry, Faculty of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou 17551, Greece

^b NovaMechanics Ltd, Nte tyras 16 Karantoki Building, Nicosia 1065, Cyprus

^c Department of Pharmaceutical Chemistry, Faculty of Pharmacy, National and Kapodistrian University of Athens, Panepistimiopolis Zografou 17551, Greece

EthnoHERBS aims to evaluate information on Southeastern European traditional knowledge for the treatment of skin diseases and elaborate cutting-edge technologies in Natural Products chemistry. After the systematic reviewing of published ethnobotanical surveys in Balkan countries, 1289 taxa were recorded for their skin beneficial effects. The recorded information was assessed based on current scientific data concerning the biological properties of the identified plants resulting in the collection of 240 species of Balkan flora and the preparation of 720 extracts. Complementary to enzyme inhibition bioactivity evaluation, in silico studies were introduced in the assortment process of the most promising ones.

Protein and ligand structure preparation were performed by means of the NovaMechanics Asclepios KNIME (Fig.1) pipeline (Papadopoulou et al., 2021). The metabolites of species were searched in Reaxys and their structures were converted to SMILES strings. The ligand structure preparation involved addition of missing hydrogen atoms, conversion of the 2D structure to 3D by performing energy minimization to obtain a low energy conformation. After removal of duplicates and exclusion of ligands <175 Daltons, the ligands were reduced from *ca.* 210,000 to 20,000. The latter were docked within the experimental binding cavity of selected enzymes such as elastase, tyrosinase, collagenase, xanthine oxidase, cyclooxygenase, and lipoxygenase. The receptors were prepared, protonated and hydrogen-treated with PDBFixer node and docking calculations were performed with the Vina-GPU software (Eastman et al., 2017). Among the most promising species that were identified were *Momordica charantia*, *Juglans regia* and *Cistus creticus* subsp. *eriocephalus*.




Fig. 1. Enalos Asclepios KNIME pipeline for enzyme inhibitors discovery automation

Keywords: in silico, Balka flora, enzyme inhibitors, skin disorders

References

- Eastman, P., Swails, J., Chodera, J.D., McGibbon, R.T., Zhao, Y., Beauchamp, K.A., et al. 2017. OpenMM 7: Rapid development of high performance algorithms for molecular dynamics. PLoS Comput Biol 13(7): e1005659.
- Papadopoulou, D., Drakopoulos, A., Lagarias, P., Melagraki, G., Kollias, G., Afantitis, A. 2021. In silico identification and evaluation of natural products as potential tumor necrosis factor function inhibitors using advanced Enalos Asclepios KNIME Nodes. Int J Mol Sci. 22(19):10220.

* Presenting author. E-mail address: cheilarianti@pharm.uoa.gr (Antigoni Cheilari)

S3.P251 Chemical characterization of a *Schisandra sphenanthera* fruit extract, an adaptogen from a sustainable panda-friendly conservation program

Alison D. Pawlus^{a,*}, Paul Kiehne^a, Nevena Karaman-Jurukovska,^b Kate Sammons^a, Jeanna Zguris^a

^a Hair Innovation and Technology, Aveda R&D, Blaine MN, USA

^b Bioscience, The Estée Lauder Companies R&D, Melville NY, USA

Schisandra sphenanthera Rehd. et Wils (Schisandraceae), commonly known as Southern Schisandra, *nan wu wei zi*, is used interchangeably with the adaptogen, *S. chinensis*, in traditional Chinese medicine (TCM). It is endemic to southern China, where it is used in indigenous medicines of ethnic minorities of Sichuan, including Qiang, Tibetan, and Yi, where it grows in forests of Minshan and Qinling mountains. Its habitat overlaps with the habitat of the Giant Panda (Brinckmann, 2018). This ingredient meets FairWild, Giant Panda Friendly Products, and Organic Wild-crop Harvesting Practice Standards.

Schisandra species contain unique dibenzocyclooctadiene lignans (DBCL) and nortriterpenoids with a wide range of biological activities. As part of our interest in finding sustainable, high functioning ingredients, we chemically characterized a *S. sphenanthera* extract obtained from a hot water (<90°C) high pressure reiterative process using UPLC-MS. Using LC-MS to obtain a chemical profile of this important botanical, we tentatively identified several schindilactone nortriterpenoids from *S. sphenanthera* fruit for the first time. This extract was shown to activate nrf2, a major transcription factor that regulates the cellular defense through the expression of genes involved in oxidative stress response and drug detoxification, measured by NQO1 activity in normal human epidermal keratinocytes.

Keywords: *Schisandra sphenanthera*, Schisandraceae, lignans, sustainable sourcing

References

Brinckmann, J.A., Luo, W., Xu, Q., He, X., Wu, J., Cunningham A.B., 2018. Sustainable harvest, people and pandas: Assessing a decade of managed wild harvest and trade in *Schisandra sphenanthera*. J. Ethnopharmacol. 224, 522-534.

* Presenting author. E-mail address: apawlus@aveda.com (Alison D. Pawlus)

S3.P252 LC-MS/MS analysis of polyphenolic and triterpene components and antioxidant activity of cold-pressed rose oils

Marta Olech

Department of Pharmaceutical Botany, Medical University of Lublin, 1 Chodźki Str., 20-093 Lublin, Poland

Cold-pressed oils are preferred over refined edible oils and are gaining growing consumer interest since they are characterized by higher levels of bioactive compounds (Rabiej-Kozioł et al., 2023). Among vegetable oils, rose oil is one of the most expensive and most highly valued, especially for its cosmetic use (Ayati et al., 2018).

The aim of the study was to compare chemical composition and antioxidant activity of rose oils available on the Polish market. Special attention was paid to the liquid chromatography-mass spectrometry (LC-MS/MS) analysis of polyphenolic and triterpene components as potential markers for oil authentication. Another goal was to determine to what extent polar and non-polar components are responsible for oil's antioxidant activity. Antioxidant assays (the ABTS^{•+} radical scavenging, H-ORAC, and L-ORAC) were adjusted to enable fast analysis of oil samples on microplates.

It was demonstrated that the polyphenolic profile, colour and carotenoid content (1.22-79.53 mg/mL) of oils from different producers differ significantly. The dissimilarities in the content of quercetin, protocatechuic, syringic, salicylic and caffeoylquinic acids were particularly noticeable. Interestingly, analysis of triterpene content did not reveal significant qualitative differences. The TEAC values of the oils varied from 349.06 to 907.18 µg Trolox/mL, and most of this activity was found to be caused by the non-polar components of the oily fraction (515.01 - 849.17 µg Trolox/mL).

The observed differences suggest large discrepancies in rose oil production (e.g. botanical origin, quality and preparation of plant material) and the need to establish quality standards and methods for verifying oil's authenticity.

Keywords: cold-pressed oil, LC-MS, polyphenols, hydrophilic oxygen radical absorbance capacity, lipophilic oxygen radical absorbance capacity

References

- Ayati, Z., Amiri, M.S., Ramezani, M., Delshad, E., Sahebkar A., Emami, S.A., 2018. Phytochemistry, traditional uses and pharmacological profile of rose hip: a review. *Curr. Pharm. Des.* 24, 4101-4124.
- Rabiej-Kozioł, D., Momot-Ruppert, M., Stawicka, B., Szydłowska-Czerniak, A., 2023. Health benefits, antioxidant activity, and sensory attributes of selected cold-pressed oils. *Molecules.* 28, 5484.

* Presenting author. *E-mail address:* marta.olech@umlub.pl (Marta Olech)

S3.P253 Chemical insights into a bioactive polyphenolic extract from *Camellia japonica* L. flower

Humaira Khan*, Lara Comune, Simona Piccolella, Severina Pacifico

Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania 'Luigi Vanvitelli', Via Vivaldi 43, 81100 Caserta, Italy

Camellia is a genus of flowering plants, which belongs to Theaceae family. Although some *Camellia* spp. are broadly exploited for their essential oils and tea for their leaves, a few species are grown as garden ornamentals (Garcia-Jares et al., 2017). Among these latter, *C. japonica* L. varieties are distinguishable based on their flower anatomical features (Pereira et al., 2022), whereas those with flowers double full in shape and red in colour are the most fascinating also in a nutraceutical and/or cosmeceutical perspective.

In this context, *C. japonica* L. flowers, collected from the botanical garden of the University of Campania (Caserta, Italy), underwent ultrasound assisted maceration in formic acid acidified ethanol, and extract obtained was chromatographed on Amberlite XAD-4 resin to achieve an anthocyanin rich fraction (CjA).

CjA fraction was first chemically investigated using spectroscopic (ATR FT-IR, and UV-Vis) techniques, and then profiled by means of Ultra-High-Performance Liquid Chromatography-High-Resolution Mass Spectrometry (UHPLC-HRMS). The fraction, which consisted of anthocyanins (mostly acylated), procyanidins and flavonols, exhibited a marked antiradical activity. In fact, it massively scavenged DPPH radical and ABTS radical cation, with calculated ID₅₀ values equal to

14.2 and 2.1 µg/mL, respectively. Furthermore, when cytotoxicity was assessed by employing MTT assay on human cell lines, CjA fraction appeared to be cytotoxic, also at the lowest tested doses towards tested cancer cells, while cytotoxic effects were not observed on HaCaT keratinocytes. These preliminary data suggest further investigation needs to be carried out, for better exploiting these high-value flowers.

Keywords: *Camellia japonica*, polyphenolic extract, cytotoxicity, antioxidant activity

References

- Pereira, A.G., Garcia-Perez, P., Cassani, L., Chamorro, F., Cao, H., Barba, F.J., Simal-Gandara, J., Prieto, M.A. 2022. *Camellia japonica*: A phytochemical perspective and current applications facing its industrial exploitation. Food Chem.:X, 13, 100258.
- Garcia-Jares, C., Sanchez-Nande, M., Lamas, J. P., Lores, M. 2017. Profiling the fatty acids content of ornamental camellia seeds cultivated in Galicia by an optimized matrix solid-phase dispersion extraction. Bioeng., 4, 87.

* Presenting author. E-mail address: humaira.khan@unicampania.it (Humaira Khan)

S3.P254 Complementary LC-MS/MS analyses of alkaloid-rich fractions of the seeds of *Plukenetia volubilis* L. by LTQ Orbitrap XL and ZenoTOF 7600

Kristelle Hughes*, Marie-France Hérent, Joëlle Leclercq

Pharmacognosy Research Group, Louvain Drug Research Institute, Université Catholique de Louvain, Avenue E. Mounier 72, B1.72.03, B-1200 Brussels, Belgium

Plukenetia volubilis L. (Euphorbiaceae), better known as Sacha inchi (SI), is an oleaginous vine that produces seeds consumed in several countries in South America. However, while the seeds possess nutritional benefits (Goyal et al., 2022), an assessment of potentially toxic metabolites such as their alkaloid content has only been done by colorimetric assays or IR spectroscopy (Kumar et al., 2020; Srichamnong et al., 2018). The lack of in-depth identification of the type of alkaloids present in the seeds of SI has prevented their acceptance in the European Food Market (Authority (EFSA), 2020). Our aim was thus to investigate the chemical composition of an alkaloid-rich fraction of the defatted seeds, raw (SIRA) and after roasting (SIRO). The two fractions were analyzed by HPLC-HRMS/MS using a LTQ Orbitrap XL and UPLC-HRMS/MS with a ZenoTOF 7600. The main families identified by dereplication and in silico methods with both instruments were cyclopeptides mainly in SIRA, and primary fatty acid amides, more abundant in SIRA than SIRO. Differences in the detection and identification of sphingosine derivatives, true alkaloids and lysophosphatidylcholines between the Orbitrap and ZenoTOF analyses will be further discussed in the poster.

The combined results of the two instruments show a variety of nitrogen-containing compounds. Moreover, our analyses suggest that roasting induces degradation of these compounds, although further quantification is necessary. Mapping the complete alkaloid profile of the seeds will help to better assess its potential toxicity during human consumption.

Keywords: Sacha inchi, alkaloids, ZenoTOF 7600, fatty acid amides

References

- Authority (EFSA), E. F. S. (2020). Technical Report on the notification of roasted seeds from *Plukenetia volubilis* L. as a traditional food from a third country pursuant to Article 14 of Regulation (EU) 2015/2283. *EFSA Supporting Publications*, 17(3), 1817E. <https://doi.org/10.2903/sp.efsa.2020.EN-1817>
- Goyal, A., Tanwar, B., Kumar Sihag, M., & Sharma, V. (2022). Sacha inchi (*Plukenetia volubilis* L.) : An emerging source of nutrients, omega-3 fatty acid and phytochemicals. *Food Chemistry*, 373, 131459. <https://doi.org/10.1016/j.foodchem.2021.131459>
- Kumar, B., Smita, K., Debut, A., & Cumbal, L. (2020). Andean Sacha Inchi (*Plukenetia Volubilis* L.) Leaf-Mediated Synthesis of Cu₂O Nanoparticles : A Low-Cost Approach. *Bioengineering*, 7(2), 54. <https://doi.org/10.3390/bioengineering7020054>
- Srichamnong, W., Ting, P., Pitchakarn, P., Nuchuchua, O., & Temviriyankul, P. (2018). Safety assessment of *Plukenetia volubilis* (Inca peanut) seeds, leaves, and their products. *Food Science & Nutrition*, 6(4), 962-969. <https://doi.org/10.1002/fsn3.633>

* Presenting author. E-mail address: kristelle.hughes@uclouvain.be (Kristelle Hughes)

S3.P255 Effect of different extraction methods on determination of *Drosera capensis* active ingredients

Onur Sinan Türkmen*, Melike Kucuk

Çanakkale Onsekiz Mart University Department of Field Crops Çanakkale, Turkey Margeht Biotechnology Ltd. Çanakkale, Turkey

Drosera capensis cape sundew is a carnivorous plant native to South Africa (Jobson and Conn, 2012). *Drosera capensis* is also known contained the 1,4-naphthoquinones and 7-methyljuglone chemicals are used for antimicrobial and antifungal purposes and prevent tumor formation (Ziaratnia et al., 2009). In this study, *Drosera capensis* was propagated by tissue culture method extracted by ultrasound, soxhlet and microwave extraction methods and its 1,4-naphthoquinones content determined by HPLC method. It was concluded that different extraction methods had a significant impact on the determination of the active ingredient content in *D. capensis*.

Keywords: cape sundew, 1,4-naphthoquinones, ultrasound, Soxhlet, microwave

References

- Jobson R. W. and Conn B. J., 2012. *Drosera capensis* (Droseraceae), a new naturalised record for Australia. *Journal of Plant Systematics* 14: 89–92.
- Ziaratnia S.M., Kunert K.J., Lall N., 2009. Elicitation of 7-methyljuglone in *Drosera capensis*. *South Africa Journal of Botany* 75:97-103.

Acknowledgments

This work was supported by Çanakkale Onsekiz Mart University The Scientific Research Coordination Unit, Project number: 2023-FBA-4619

* Presenting author. E-mail address: onurturkmen@comu.edu.tr (Onur Sinan Türkmen)

S3.P256 Valorization of *Olea europaea* L. cv. Caiazzana pruning waste: a step towards a sustainable approach for resource functional recovery

Hamid Mushtaq*, Sarmad Ahmad Qamar, Simona Piccolella, Severina Pacifico

Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania "Luigi Vanvitelli", Via Vivaldi 43, 81100 Caserta, Italy

Olive leaves by the pruning and harvesting of olive trees (*Olea europaea* L.) represent one of the olive oil industry by-products, whose valorization needs to be pursued also in an innovative food & nutraceutical scenario. In this context, also taking into account cultivar biodiversity, leaves from *Olea europaea* L. cv. Caiazzana, an autochthonous cultivar of Southern Italy, collected after pruning in an orchard near Caiazzo (Caserta, Italy), and freeze-dried, underwent ultrasound-assisted extraction (UAE) (Pacifico et al., 2022) using ethanol. The alcoholic extract was further fractionated to achieve a polyphenol fraction, which was quali-quantitatively profiled by UHPLC-ESI-QqTOF-MS/MS techniques, and screened for its antiradical capability, Fe(III) reducing power, and antibacterial activity. The cytotoxicity was also carried out towards normal-like and cancer cells. Nevertheless, the exhausted leaves were explored for recovering pure cellulose, characterized chemically by ATR FT-IR tools, and morphologically by scanning electron microscopy (SEM). Thus, cellulose was properly derivatized and cellulose-based films entrapping olive leaf polyphenol fraction were fabricated. FTIR spectroscopy and SEM suggested the occurrence of hydrogen-bonding among the film components. The films maintained the antioxidant efficacy of the bioactive extract and advantageously release the polyphenols when allowed them to adhere to a depletable product, such as meat (storage time considered 10 days), preserving it from oxidation, and functionally enriching it in bioactive compounds. Thus, the recovery of olive leaves opens up to their favourable use for the creation of active films capable of enriching foods in our diet with functional constituents.

Keywords: *Olea europaea* L. leaves, polyphenols, cellulose, nutraceutical & food sectors

References

Pacifico, S., Bláha, P., Faramarzi, S., Fede, F., Michaličková, K., Piccolella, S., Ricciardi, V., Manti, L., 2022. Differential Radiomodulating Action of *Olea europaea* L. cv. Caiazzana Leaf Extract on Human Normal and Cancer Cells: A Joint Chemical and Radiobiological Approach. *Antioxidants*, 2022, 11, 1603.

* Presenting author. E-mail address: hamid.mushtaq@unicampania.it (Hamid Mushtaq)

S3.P257 Plants from Lublin region as potential sources of ingredients with cosmetic value

Kamila Kusio-Targońska^a, Katarzyna Gaweł-Beben^b, Krzysztof Kamil Wojtanowski^a, Wirginia Kukuła-Koch^a

^a Department of Pharmacognosy with Medicinal Plant Unit, Medical University of Lublin, 1 Chodzki Str. 20-093 Lublin, Poland

^b Department of Cosmetology, The University of Information Technology and Management in Rzeszow, 2 Sucharskiego Str, 35-225, Rzeszów, Poland

Nowadays, ingredients used in cosmetics are expected to perform multiple functions and protect the skin against harmful effects of environmental factors. Due to ongoing environmental changes, there is a great demand for cosmetic ingredients that have both antioxidant and skin brightening properties. This project aimed to develop analytical conditions that would allow the identification of extracts with antioxidant and tyrosinase-inhibitory potential among the medicinal plants occurring in the Lublin region. From the tested ethanol, water and 50% ethanol extracts of 10 plant species four were found to be the most active, namely *Matricaria chamomilla*, *Sambucus nigra*, *Urtica dioica* and *Melissa officinalis*. Their percentage inhibition of the DPPH radical was calculated as 38,561%, 35,791%, 16,711% and 50,536% respectively. The tyrosinase inhibition was calculated as 81,16% for the water: ethanol extract (50:50 v/v), 20,29% for the ethanol extract, 28,14% for the ethanol extract and 16,91% for the water extract respectively, whereas the total phenolic content was determined using the Folin Ciocalteu reagent. Gallic Acid Equivalent (GAE) was calculated as 13,35 for the water: ethanol extract (50:50 v/v), 16,69 for the ethanol extract 13,98 for the ethanol extract, 16,68 for the water extract respectively.

The HPLC-ESI-QTOF-MS/MS fingerprinting was performed for the analyzed extracts. *Melissae* water extract was found to be the most rich. Among the major components caffeic, chlorogenic, caftaric, rosmarinic, syringic, genistic, p-coumaric and ferulic acid, hydroxy jasmonic acid glucoside, rutine, apigenin, were identified.

Also, the most active extract was subjected to an on-line assay using an HPLC-MS instrument coupled with a bioreactor to determine single constituents with anti-tyrosinase properties in the extract.

Keywords: *Melissa officinalis*, antioxidant, anti-tyrosinase, cosmetics, HPLC-ESI-QTOF-MS/MS

Presenting author. E-mail address: kamilakusio@poczta.fm (Kamila Kusio-Targońska)

S3.P258 Anti-neuroinflammatory effects of *Elsholtzia blanda* Benth on lipopolysaccharide-stimulated BV-2 microglial cells

Yeo Dae Yoon^c, Krishna K. Shrestha^b, Seung-Hwa Baek^{a,*}

^a Zebrafish Translational Medical Research Center, Korea University College of Medicine, Seoul, Republic of Korea

^b Ethnobotanical Society of Nepal, 107 Guchha Marg, New Road, MJ Mart, Kathmandu, Nepal

^c Bioevaluation Center, Korea Research Institute of Bioscience and Biotechnology, Ochang, Cheongwon, Chungbuk, Korea

Elsholtzia blanda (Benth.) Benth. (EBB) is a medicinal plant in Nepal with traditional use. Several studies have shown that EBB has bioactive properties, including antimicrobial (Nguyen et al., 2022; Ishwori et al., 2015) and anti-inflammatory effects (Duong et al., 2023). In this study, the methanol extract of EBB was tested to evaluate its anti-inflammatory effect against the lipopolysaccharide (LPS)-induced microglial macrophage cells (BV2). In LPS-activated BV2 microglial cells, EBB significantly reduced the induced proinflammatory mediators, e.g. TNF- α and PEG2, as well as nitric oxide (NO) accumulation. EBB also attenuated LPS-induced ROS in BV2 cells. In addition, both protein and mRNA expression levels of iNOS and COX-2, which were upregulated by LPS, were significantly decreased by EBB extract. Also, EBB extract effectively reduced the transcriptional expression and further suppressed the increased production of inflammatory cytokines by LPS stimulation. The results obtained in this study showed that EBB extract effectively suppressed the inflammatory response induced by LPS in BV-2 cells. We suggest that EBB extract holds promise as a preventive agent against diseases triggered by microglial inflammatory responses.

Keywords: *Elsholtzia blanda* (Benth.) Benth., microglial macrophage (BV2), neuroinflammation, lipopolysaccharide, anti-inflammatory effect

References

- Nguyen V. N. and Hoang T. B., 2022. The antimicrobial activity and chemical composition of *Elsholtzia blanda* (Benth.) Benth. essential oils in Lam Dong Province, Viet. *Nam. J. Innov. Sustain. Dev.* 14. 72-77.
- Ishwori L., Talukdar A. D., Singh P. K., Choudhury D. M., Nath D., 2015. Antibacterial activity of some selected plants traditionally used as medicine in Manipur. *African J. Biotech.* 13, 1491-1495.
- Duong T. D., Do T. T., Pham H. Y., Nguyen H. H., Bui H. T., Phan V. K., 2023. Elsholblanosides A-D, four New oleuropeic acid derivatives isolated from *Elsholtzia blanda* and their inhibition of NO production in LPS-activated RAW264.7 Cells. *Chem Biodivers.* 20. e202300785.

* Presenting author. E-mail address: micro340@gmail.com (Seung-Hwa Baek)

S3.P259 Metabolite profiles and biological activities of *Capsicum chinense* cell cultures

Amir Akhgari^{a,*}, Maria Sulli^b, Anna Ramata Stunda^c, Suvi T. Häkkinen^a, Liisa Nohynek^a, Aleksander Salwinski^d, Olivia Costantina Demurtas^b, Mārtiņš Boroduškis^c, Maria Pajumo^a, Richard Twyman^e, Kirsi-Marja Oksman-Caldentey^a, Gianfranco Diretto^b, Heiko Rischer^a

^a VTT Technical Research Centre of Finland Ltd., Tekniikantie 21, FIN-02150 Espoo, Finland

^b Italian National Agency for New Technologies, Energy, and Sustainable Development (ENEA), Casaccia Research Center, 00123 Roma, Italy

^c Alternative Plants Ltd., 2 Podraga Str., LV-1007 Riga, Latvia

^d Plant Advanced Technologies, 19 Avenue de la Forêt de Haye, 54500 Vandoeuvre-lès-Nancy, France

^e TRM Ltd, Scarborough, UK

Research and industry pursue natural alternatives to synthetic antioxidants and antiaging agents for safer use in food and cosmetics. Chili fruits harbor capsaicinoids, polyphenols, carotenoids and fatty acids, offering anti-inflammatory, antioxidant, and antitumoral effects (Alonso-Villegas et al., 2023). Consistent agricultural production of these constituents is challenging due to yield variability, highlighting sustainable sourcing importance for cosmetic and pharmaceutical industries. Alternatively, plant cell culture offers a renewable, eco-friendly production system.

In the frame of the EU-InnCoCells project, we established *Capsicum chinense* var. Pimento cell suspension cultures in dark and photoperiodic growth conditions. Subsequently, hydrophilic and hydrophobic compounds were extracted by hydroethanol and ethyl acetate, respectively. Extracts were then subjected to comprehensive metabolite profiling. Chemical analyses revealed that the hydrophilic extracts contain capsaicinoids, sesquiterpenes, phenolic acids, flavonoids and phenolamides, whilst the hydrophobic extracts included carotenoids and fatty acids. The most abundant components were phenolamides. Phenolamides have been reported to possess skin depigmentative and antioxidative activities (Roumani et al., 2020).

Finally, antioxidant, antimicrobial and cytotoxicity of cell culture hydrophilic extracts were investigated. Extracts exhibited potent antioxidant properties. Hydrophilic extracts effectively inhibited the growth of Gram-positive *Staphylococcus aureus* and *S. aureus* MRSA strains.

The hydrophilic extracts of all cell cultures were not cytotoxic at concentrations of 2.5 and 1.25 mg/mL, respectively, against the BALB/c 3T3 and HaCaT cell lines. Furthermore, cell extracts did not exhibit phototoxicity in the BALB/c 3T3 cell line at a concentration below 2.5 mg/mL.

Keywords: Chili, Plant cell culture, Antimicrobial and antioxidant activity, Cytotoxicity, Cosmetics

References

Alonso-Villegas, R., González-Amaro, R.M., Figueroa-Hernández, C.Y., Rodríguez-Buenfil, I.M., 2023. The Genus *Capsicum*: A review of bioactive properties of its polyphenolic and capsaicinoid composition. *Molecules* 28, 1–23.

Roumani, M., Duval, R.E., Ropars, A., Risler, A., Robin, C., Larbat, R., 2020. Phenolamides: Plant specialized metabolites with a wide range of promising pharmacological and health-promoting interests. *Biomed. Pharmacother.* 131, 110762.

* Presenting author. E-mail address: amir.akhgari@vtt.fi (Amir Akhgari)

S3.P260 Liposome encapsulation of *Lavandula austroapennina* N.G. Passal., Tundis & Upson leaf and stem extracts for cosmeceutical innovation

Claudia Gravina^a, Simona Piccolella^a, Cristina Alonso^b, Meritxell Martí^b, Marialuisa Formato^a, Severina Pacifico^a, Luisa Coderch^b, Assunta Esposito^a

^a Department of Environmental, Biological and Pharmaceutical Sciences and Technologies, University of Campania 'Luigi Vanvitelli', Via Vivaldi 43, 81100 Caserta, Italy

^b Surfactants and Nanobiotechnology Department, Institute of Advanced Chemical of Catalonia of CSIC, (IQAC-CSIC), Jordi Girona 18-26, 08034 Barcelona

Plant-derived specialized metabolites are highly researched for their efficacy against disease and for their use in cosmetics and dietary supplements (Elshafie et al., 2023). However, their poor water solubility and stability pose a challenge, impacting their efficacy and bioavailability (Bhalani et al., 2022). Liposomes are extensively applied as lipid-based carriers for topical drug delivery, improving the penetration of bioactive compounds through skin layers (Chaves et al., 2023). In this context, extracts from the endemic species *Lavandula austroapennina* N.G. Passal. Tundis & Upson, were explored for their cosmeceutical potential.

L. austroapennina plants from Mt. Cervati were fractionated into its different organs and extracted using ultrasound-assisted maceration in *n*-hexane and methanol. Phytochemical analysis through UHPLC-HR-MS/MS identified fatty acids and olean- and ursane-type triterpenes in lipophilic extracts, and hydroxycinnamoyl and 8-hydroxyphenylpropanoid acid derivatives in alcoholic ones. Previous studies highlighted the superior bioactivity of stem lipophilic and leaf alcoholic extracts (Chianese et al. 2023; Gravina et al. 2023), selected for encapsulation in liposome-based nanocarriers. Accordingly, extracts-containing liposomes, preliminarily characterized by ATR-FTIR spectroscopy, underwent cytotoxic and healing capability assessment towards keratinocytes cell models. The extracts-enriched liposomes showed a healing capacity higher than the parental extracts they were derived from. A joint approach employing UHPLC-HR-MS/MS and *in vitro* skin permeation studies revealed that encapsulation in liposomes improved bioactive delivery through/into a specific skin layer. These findings suggest the effectiveness of liposome-based nanoparticles in improving the bioavailability and biological activity of specialized metabolites, thereby reinforcing the cosmeceutical potential of *L. austroapennina* extracts.

Keywords: cosmeceuticals, encapsulation, percutaneous absorption, UHPLC-QqTOF-HR-MS, ATR- FTIR

References

- Bhalani, D.V., Nutan, B., Kumar, A., Singh Chandel, A.K., 2022. Bioavailability Enhancement Techniques for Poorly Aqueous Soluble Drugs and Therapeutics. *Biomedicines* 10(9),2055.
- Chaves, M.A., Ferreira, L.S., Baldino, L., Pinho, S.C., Reverchon, E., 2023. Current applications of liposomes for the delivery of vitamins: a systematic review. *Nanomaterials*, 13, 1557.
- Elshafie, H.S., Camele, I., Mohamed, A.A., 2023. A Comprehensive Review on the Biological, Agricultural and Pharmaceutical Properties of Secondary Metabolites Based-Plant Origin. *Int. J. Mol. Sci.* 24(4), 3266.
- Chianese, A., Gravina, C., Morone, M.V., et al. 2023. *Lavandula austroapennina*: Assessment of the Antiviral Activity of Lipophilic Extracts from Its Organs. *Viruses* 15(8),1648.
- Gravina, C., Formato, M., Piccolella, S., Fiorentino, M., Stinca, A., Pacifico, S., Esposito, A., 2023. *Lavandula austroapennina* (Lamiaceae): Getting Insights into Bioactive Polyphenols of a Rare Italian Endemic Vascular Plant. *Int. J. Mol. Sci.* 24(9),8038.

* Presenting author. E-mail address: claudia.gravina@unicampania.it (Claudia Gravina)

S3.P261 Effect of combining proteolytic enzymes papain and chymotrypsin with Alexandrite laser hair removal on the degree of facial hirsutism

Aikaterini Liatsopoulou^a, Athanasia Varvaresou^{a,b,*}, Evangelia Protopapa^a, Foteini Mellou^{a,b}

^a Department of Biomedical Sciences, University of West Attica, Egaleo 122 43, Greece

^b Laboratory of Chemistry, Biochemistry and Cosmetic Science, Department of Biomedical Sciences, University of West Attica, Egaleo 122 43, Greece

Papain (**1**) is a proteolytic enzyme derived from the latex of *Carica papaya*, which is used for depilatory preparations (Mellou et al., 2020). Chymotrypsin (**2**) is a pancreatic serine protease (Fu et al., 2021). Iontophoresis of **1** and **2** to skin of experimental animals has shown long-lasting depilatory effects (Protopapa et al., 1999). Photo-epilation is the treatment of choice for hirsutism. However, it does not offer complete and persistent results (Mikiel et al., 2020). Therefore, we opted to combine photo-epilation with application of these enzymes, as a more natural approach, to enhance treatment efficacy.

In this randomized controlled clinical study, 59 adult Caucasian women with facial hirsutism were divided into two groups: Group I, treated with combination of laser Alexandrite 755nm and iontophoresis of aqueous solutions **1** (0.48%) and **2** (0.29%) and Group II, treated with laser Alexandrite 755nm alone. The degree of facial hirsutism (0-4) was assessed by a dermatologist and the participants, based on the modified Ferriman-Gallwey scoring system. After 10 sessions, a comparison between the two groups of the degree of hirsutism was performed.

Participants of Group I had a statistically significantly lower overall degree of hirsutism compared to Group II, according to both the dermatologist's (p=0.014) and the participants' assessment (p=0.009). Iontophoresis of **1** and **2** provides a more natural adjuvant treatment, with a potential to enhance laser hair removal efficacy in hirsute patients.

Keywords: papain, chymotrypsin, photo-epilation, facial hirsutism, efficacy

References

- Fu, Z., Akula, S., Thorpe, M., Hellman, L., 2021. Marked difference in efficiency of the digestive enzymes pepsin, trypsin, chymotrypsin, and pancreatic elastase to cleave tightly folded proteins. *Biological Chemistry*. 402(7), 861-867.
- Mellou, F., Varvaresou, A., Papageorgiou, S., Protopapa, E., Liatsopoulou, E., 2020. Papain: An Enzyme of Multiple Applications. *Rev. Clin. Pharmacol. Pharmacokinet. Int.* 34, 107-111.
- Mikiel, D., Olszewska, B., Polanska, A., Adamski, Z., Zaba, R., Danczak-Pazdrowska, A., 2020. Principles of management of women with hirsutism – a dermatologist's perspective. *Dermatol Rev/Przegl Dermatol.* 107, 424-40.
- Protopapa, E.E., Gaissert, H., Xenakis, A., Avramiotis, S., Stavrianeas, N., Sekeris, C.E., Schenkelf, J., Alonso, A., 1999. The effect of proteolytic enzymes on hair follicles of transgenic mice expressing the lac Z-protein in cells of the bulge region. *J Eur Acad Dermatology Venereol.* 13(1), 28-35.

* Presenting author. E-mail address: avarvares@uniwa.gr (Athanasia Varvaresou)

S3.P262 Simultaneous analysis of bergapten and schinifoline in *Zanthoxylum schinifolium* Siebold & Zucc. seeds using HPLC and UPLC-MS/MS systems

ChangSeob Seo*

KM Science Research Division, Korea Institute of Oriental Medicine, Daejeon, South Korea

Zanthoxylum schinifolium Siebold & Zucc. belongs to the Rutaceae family and has been widely used as a spice (typically seeds and pericarp) in East Asian countries such as Korea, China, and Japan (Oh and Chung, 2014). The present study focused on developing and validating a simultaneous analytical method for marker substances (bergapten and schinifoline; Fig. 1) in *Z. schinifolium* seeds. This was achieved using high-performance liquid chromatography (HPLC) with a photo-diode array detector (DAD) and ultra-performance liquid chromatography (UPLC) with tandem mass spectrometry systems.

In the regression equation, all markers showed a coefficient of determination of ≥ 0.9990 . Marker recovery was 96.90–105.16% (relative standard deviation (RSD) ≤ 2.23), and the intra- and interday precision was RSD < 3.00 . Bergapten and schinifoline were detected in the seeds at 1.70–2.85 mg/g and 0.19–0.94 mg/g, respectively. This analytical method will improve quality control of *Z. schinifolium* seeds. Additionally, this assay will provide basic data and quality assurance for future biological activity experiments or clinical applications.




Fig. 1. Chemical structures of the two markers selected of *Z. schinifolium* seeds.

Keywords: simultaneous analysis, *Zanthoxylum schinifolium*, bergapten, schinifoline

References

Oh, M., Chung, M.S., 2014. Effects of oils and essential oils from seeds of *Zanthoxylum schinifolium* against foodborne viral surrogates. Evid. Based Complement. Alternat. Med. 2014, 135797.

* Presenting author. E-mail address: csseo0914@kiom.re.kr (ChangSeob Seo)

S3.P263 Health-promoting effects of the hydroalcoholic extract from the *Allium sativum* L. ecotype “Sulmona red garlic” in the colon

Claudio Ferrante^{a*}, Fabrizio Masciulli^{b,c}, Cinzia Ingallina^{b,c}, Luisa Mannina^{b,c}, Maria Loreta Libero^a, Simonetta Cristina Di Simone^a, Alessandra Acquaviva^a, Nilofar Lucia Recinella^a, Sheila Leone^a, Luigi Brunetti^a, Giustino Orlando^{a,*}, Gokhan Zengin^d, Luigi Menghini^{a,*}, Annalisa Chiavaroli^a

^a Department of Pharmacy, G. d'Annunzio University of Chieti-Pescara, Chieti, Italy

^b Food Chemistry Lab, Department of Chemistry and Technology of Drugs, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185 Rome, Italy

^c NMR Lab, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185 Rome, Italy ^dPhysiology and Biochemistry Laboratory, Department of Biology, Science Faculty, Selcuk University, 42130 Konya, Türkiye

In the present study, we investigated the chemical composition and biological properties of the hydroalcoholic extract from the aerial bulbils of the *Allium sativum* L. ecotype “Sulmona red garlic”, a traditional ecotype of middle Italy cultivated in the Abruzzo region. The study aimed at exploring the potential application of a plant material, namely aerial bulbils, that is considered a waste material in the “Sulmona red garlic” supply chain. The aerial bulbils extract was tested in different biological models, including an ecotoxicological model constituted by the crustacean *Daphnia magna* and cell two different cultures (C2C12 and HCT116 cell lines), with the aim to unravel the limits of its biocompatibility and protective effects. Additionally, a deep phytochemical analysis was conducted for determining the content of specialized metabolites. NMR analyses demonstrated the content of alliin and allicin, in the aerial bulbils where different phenolic compounds were identified and quantified, among which the prominent were catechin, chlorogenic acid, and gallic acid. The extract was biocompatible in the *Daphnia magna* model (up to a concentration of 10 mg/mL) and in the non-tumoral C2C12 cell line, in the concentration range of 10-1000 µg/mL. By contrast, a concentration-dependent cytotoxic effect was observed in the human colon cancer HCT116 cell line. This effect was related, albeit partially, to the inhibition of the transient receptor potential (TRP) M8 (TRPM8), whose upregulation has been implicated in inflammation and carcinogenesis. Additionally, the extract inhibited the gene expression of TNF-α (tumor necrosis factor), HIF1-α (hypoxia-inducible factor), and VEGFA (vascular endothelial growth factor), thus further demonstrating promising health-promoting effects, especially against inflammatory bowel diseases.

Keywords: garlic, organosulphur compounds, phenolic compounds, cytotoxicity, colon, TRPM8

References

Lasalvia, A., Cairone, F., Cesa, S., Maccelli, A., Crestoni, M. E., Menghini, L., Carradori, S., Marinacci, B., Gallorini, M., & Elsallabi, O. 2022. Characterization and Valorization of ‘Sulmona Red Garlic’ Peels and Small Bulbs. *Antioxidants*, 11 (11), 2088.

*Presenting author: claudio.ferrante@unich.it (Claudio Ferrante)

S3.P264 Anti-inflammatory effect of *Lysimachia mauritiana* extract on particulate matter 10 plus diesel exhaust particles (PM10D)-induced respiratory disease mouse model and standardization using UPLC-qTof/MS analysis

Yoon-Young Sung^{a,*}, Heung Joo Yuk^a, Seung-Hyung Kim^b, Chang-Hwa Song^c, Dong-Seon Kim^a

^a KM Research Science Division, Korea Institute of Oriental Medicine, Daejeon 34054, Republic of Korea

^b Institute of Traditional Medicine and Bioscience, Daejeon University, Daejeon 34520, Republic of Korea

^c Department of microbiology College of Medicine Chungnam National University, 266 Munhwa-ro, Jung-gu, Daejeon 35015, Republic of Korea

Exposure to particulate matter (PM) causes considerable respiration-related health risks. The effect of *Lysimachia mauritiana* (LM) extract on airway inflammation was investigated in mice exposed to a fine dust mixture (PM10D) of PM10 (PM diameter < 10 µm) and diesel exhaust particles (DEP). LM suppressed the number of immune cells and secretion of cytokines in bronchoalveolar lavage fluid (BALF) and lung of PM10D mice. Airway inflammation and collagen fibrosis in the lung after PM10D exposure were investigated by histopathological analysis, and LM ameliorated these symptoms. LM decreased the mRNA expression of MUC5AC, CXCL1, TNF-α, MIP-2, TRPA1, and TRPV1 in lung. The main candidate components from LM that exhibited anti-inflammatory effects were investigated by ultra-performance liquid chromatography quadrupole time of flight mass spectrometry (UPLC-qTof/MS). The seven phytochemicals were tentatively identified by MS analysis. Compound 3 among the flavonoids, which are easy to detect even under UV, is identified into biorobin (keampferol 3-O-robinobioside) and was confirmed as candidate of standardization. These findings suggest that LM may be a potential therapeutic agent for the treatment of respiratory diseases in human.

Keywords: airway inflammation, fine dust mixture, *Lysimachia mauritiana*, lung

References

- Kyung, S.Y., Jeong, S.H., 2020. Particulate-Matter related respiratory diseases. *Tuberc. Respir. Dis.* 83, 116–121. Tangavel, P., Park, D., Lee, Y.C., 2022. Recent insights into Particulate Matter (PM2.5)-mediated toxicity in humans: An overview. *Int. J. Environ. Res. Public Health.* 19, 7511.
- Lee, Y.S., Yang, W.K., Park, Y.R., Park, Y.C., Park, I.J., Lee, G.J., Kang, H.S., Kim, B.K., Kim, S.H., 2022. *Opuntia ficus-indica* alleviates particulate matter 10 plus diesel exhaust Particles (PM10D)-induced airway inflammation by suppressing the expression of inflammatory cytokines and chemokines. *Plants.* 11, 520.

Acknowledgment

This study was supported by the project (KSN23214313) and (NSN2213010) from the KIOM of Korea.

* Presenting author. E-mail address: yysung@kiom.re.kr (Yoon-Young Sung)

S3.P265 Optimization, estimation, and improvement of lutein content from marigold flowers using *Lactobacillus* species by solid-state fermentation

Zuha Rahman*, Vidhu Aeri, Bibhu Prasad Panda

Department of Pharmacognosy and Phytochemistry, School of Pharmaceutical Education and Research, Jamia Hamdard, New Delhi-110062, India

Lutein is primarily sourced from marigold petals which are used in the cosmetic, nutraceutical, and pharmaceutical industries (Anna et al., 2020). This contributes to overall eye health and brain diseases (Mitra et al., 2021; Yagi et al., 2021). In this study, we investigated the improvement in lutein content in marigold flowers (*Calendula officinalis* Linn) via solid-state fermentation using three different *Lactobacillus* species such as *L. rhamnosus*, *L. casei*, and *L. plantarum*. Microorganisms were isolated from the National Dairy Research Institute (India) and fermented for 10 days in an incubator with fresh and dried marigold petals. Fermented and non-fermented petals were extracted with hexane and ethanol, and the presence of lutein was confirmed by HPLC at 446 nm. Using hexane as a solvent, the experiment revealed variations in the lutein content of different types of petals. Fresh petals (1 kg) fermented with *L. plantarum* contained 4.0% lutein, whereas dried petals (1 kg) fermented with *L. rhamnosus* contained 0.99% lutein. Non-fermented fresh petals contained 1.1% lutein, whereas non-fermented dried petals contained 0.4% lutein. Furthermore, the use of ethanol also yielded satisfactory results. When fresh marigold flowers (1 kg) were fermented by *L. plantarum*, they contained 0.43% lutein and dried petals (1 kg) contained 0.52% lutein when fermented by *L. rhamnosus*. Based on these findings, *L. plantarum* enhanced lutein content by 3-fold in hexane. In addition, ethanol is a useful green replacement for conventional hazardous solvents, and *Lactobacillus* species are preferred to enhance the extraction of lutein from marigold petals in research laboratories and industries.

Keywords: Lutein, *Lactobacillus* species, marigold, solid-state fermentation

References

- Anna, B., Cristina, C., Maria, C. T., Nicolae, C., 2020. Determination of carotenoids in extracts from species of *Tagetes* and *Calendula*. The Moldovan Medical Journal, 63(4), 23-26.
- Mitra, S., Rauf, A., Tareq, A. M., Jahan, S., Emran, T. B., Shahriar, T. G., Rengasamy, K. R., 2021. Potential health benefits of carotenoid lutein: An updated review. Food and Chemical Toxicology, 154, 112328.
- Yagi, A., Nouchi, R., Butler, L., Kawashima, R., 2021. Lutein has a positive impact on brain health in healthy older adults: a systematic review of randomized controlled trials and cohort studies. Nutrients, 13(6), 1746.

* Presenting author. E-mail address: zuharahman_sch@jamiahamdard.ac.in (Zuha Rahman)

S3.P266 Metabolite Profiling and Molecular Network Shows Kinkeloids as promoting of collagen synthesis from *Combretum micranthum*

Souhila Messaili^{1,*}, Doha Haggouch¹, Mikaela Bignard¹, Pierre-Eric Campos², Emilie Destandau², Isabelle Thuillier¹, José Ginestar¹, Eldra Delannay^{1,*}

¹ Sisley Paris, 32, Avenue des Béthunes - CS30003 - SAINT-OUEN LAUMÔNE 95067 Cergy Pontoise Cedex, France

² Institut de Chimie Organique et Analytique (ICOA), UMR 7311, Université d'Orléans, France

This presentation will investigate the molecular composition and a new cosmetically relevant biological activity of *Combretum micranthum* extract and enriched fractions to begin to establish a structure-activity relationship. *Combretum micranthum*, a plant native to Africa, has a well-documented traditional use in the treatment of various ailments such as fever, diabetes and malaria (Yoro, Tine et al., 2024). Its pharmaceutical benefits include nephroprotective, anti-inflammatory, antioxidant and antimicrobial properties have been demonstrated (Olajide, Olumayokun et al., 2003). In addition, its potential for cosmetic applications is being explored due to its depigmenting, anti-inflammatory and UV damage repairing properties (胡舒婷 et al., 2020). First, an extract of *Combretum micranthum* was prepared and selected for its overall biological response, then fractionated to obtain simplified molecular fractions. One fraction was particularly enriched in kinkeloids, a family of compounds specific to this species. All fractions and the crude extract were then tested on biological targets to evaluate and compare their cosmetic activities. Molecular networks were constructed from the UHPLC-MS/HRMS data to better characterize the extract and fractions and to highlight structure-activity relationships. This study highlights the metabolic profiling of a butylene glycol extract of *Combretum micranthum*, showing its main chemical families and revealing that the kinkeloids identified by HRMS and NMR promote an increase in collagen I synthesis, an interesting cosmetic activity not previously described for these compounds or for any *Combretum micranthum* extract.

Keywords: preparative chromatography fractionation, UHPLC-HRMS/MS, molecular network, NMR, collagen I synthesis

References

- Yoro, Tine, Madiaye Sene, Cheikhouna Gaye, Alioune Diallo, Benjamin Ndiaye, Idrissa Ndoye, and Alassane Wele. "Combretum Micranthum G. Don (Combretaceae): A Review on Traditional Uses, Phytochemistry, Pharmacology and Toxicology." *Chemistry & Biodiversity*, February 14, 2024, e202301606.
- Olajide, Olumayokun A., J. Modupe Makinde, and David T. Okpako. "Evaluation of the Anti-Inflammatory Property of the Extract of Combretum Micranthum G. Don (Combretaceae)." *InflammoPharmacology* 11, no. 3 (September 2003): 293–98.
- 胡舒婷, 西蒙J.E., 吴忆箴, 王明福, and 吴庆立. "Use of combretum micranthum extract in cosmetics. World Intellectual Property Organization" WO2020052571A1, filed September 10, 2019, and issued March 19, 2020.

* Presenting author. E-mail address: souhila.messaili@sisley.fr (Souhila Messaili)

S3.P267 Determination of the anti-inflammatory activity of fruits of some varieties of *Rubus idaeus* and *Rubus occidentalis* (Rosaceae)

Mirosława Krauze-Baranowska^{a,*}, Magdalena Majdan^b, Aleksandra Radwańska^c, Antoni Nasal^c, Barbara Sparzak-Stefanowska^a

^a Department of Pharmacognosy with Medicinal Plant Garden, Faculty of Pharmacy, Medical University of Gdańsk, Gen. J. Hallera 107, 80-416 Gdańsk, Poland

^b Department of Toxicology and Bromatology, Faculty of Pharmacy, Medical University of Warsaw, Banacha 1, Warsaw, 02-097, Poland

^c Department of Biopharmacy and Pharmacodynamics, Faculty of Pharmacy, Medical University of Gdańsk, Gen. J. Hallera 107, 80-416 Gdańsk, Poland

Red raspberry fruits (*Rubus idaeus* L., Rosaceae) are a remedy known in folk medicine for treating colds. In Poland, the species is cultivated in many varieties (Szymanowska et al., 2018). However, another raspberry species originating from North America - black raspberry (*Rubus occidentalis* L.) occurs in Poland in only two varieties, including the 'Litacz' variety (Kaume et al., 2012). Raspberries are a source of anthocyanins - cyanidin glycosides and ellagitannins with proven antioxidant and anti-inflammatory effects (Gomathi et al., 2024). The aim of the study was to evaluate the *in vivo* anti-inflammatory effect of fruits from selected varieties of red raspberry and black raspberry. The analysis of the biologically active compounds in the tested fruits was carried out using the HPLC–DAD–ESI/MS. In the pharmacological studies, a plethysmometric method was used to measure carrageenan-induced rat paw edema compared to indomethacin (reference compound). *R. occidentalis* 'Litacz' fruit extract at a dose of 500 mg/kg (p.o.) revealed the strongest anti-inflammatory effect (inhibition of rat paw swelling was 54.88% at 1 hour after administration of carrageenan, compared to indomethacin - 39.56%) ($p < 0.05$). The anti-inflammatory effect was dose-dependent except for the dose of 1000 mg/kg (23.68%). However, at this dose, at 5 hour of the experiment there was a statistically significant increase in the inhibition of edema (55%) ($p < 0.05$). The richest source of cyanidin glycosides, especially its triglycosides, turned out to be the fruits of the *R. occidentalis* 'Litacz'. Their slow hydrolysis to free aglycone - cyanidin, probably determines the long-lasting anti-inflammatory effect.

Keywords: raspberry fruits, cultivar varieties, anti-inflammatory activity

References

- Szymanowska, U., Baraniak, B., Bogucka-Kocka, A., 2018. Antioxidant, Anti-Inflammatory, and Postulated Cytotoxic Activity of Phenolic and Anthocyanin-Rich Fractions from Polana Raspberry (*Rubus idaeus* L.) Fruit and Juice-*In Vitro* Study. *Molecules*. 21;23(7):1812. doi: 10.3390/molecules23071812
- Kaume, L., Howard, L. R., Devareddy, L., 2012. The blackberry fruit: a review on its composition and chemistry, metabolism and bioavailability, and health benefits Review. *J Agric Food Chem*. 13;60(23):5716-27. doi: 10.1021/jf203318p
- Gomathi, R., Umamaheswari T. N., Prethipa, R., 2024. Evaluation of Antioxidant, Anti-inflammatory, and Antimicrobial Activities of Raspberry Fruit Extract: An *In Vitro* Study. *Cureus*, 12;16(2):e54045. doi: 10.7759/cureus.54045.

* Presenting author. E-mail address: krauze@gumed.edu.pl (Mirosława Krauze-Baranowska)

S3.P268 Unveiling Thailand's ethnopharmacological heritage to find new natural sweeteners

Antoine Bruguère^{a,*}, Efsthia Karachaliou^a, David Pertuit^a, Sorawit Watakajaturaphon^b, Pimonnart Promyos^b, Natnaree Yeerong^b, Pensak Jantrawut^b, Christine Belloir^a, Loïc Briand^a, Anne Claire Mitaine-Offer^a

^a Centre des Sciences du Goût et de l'Alimentation CSGA, CNRS, INRAE, Institut Agro, Université de Bourgogne, 21000 Dijon Cedex, France

^b Department of Pharmaceutical Sciences, Faculty of Pharmacy, Chiang Mai University, Chiang Mai 50200, Thailand

Thailand's biodiverse landscape and traditional practices in parts of southwest Asia offer a unique opportunity to explore the rich array of medicinal plants and their cultural significance. In this context, three species were sourced from local Thai markets to investigate their potential for yielding novel glycosides, such as sweet-tasting saponins.

First, the Monk fruit *Siraitia grosvenorii* (Swingle) C. Jeffrey ex A.M. Lu & Zhi Y. Zhang (Cucurbitaceae) (MPNS Portal, 2024) is a natural sweetener used for over 300 years to treat pharyngitis and pharyngeal pain (Gong et al., 2019). *Albizia myriophylla* Benth. (Fabaceae), is also prevalent in Thai and Vietnamese traditional medicine. Its stems are utilized as a substitute for licorice due to their sweet taste and it is highly valued for treating diseases of oral cavity like dental caries and aphthous ulcers (Joycharat et al., 2016). *Sapindus rarak* DC. (Sapindaceae) pericarp powder and extracts are widely used for their sweet taste and in hair and skin care, due to their rich concentration in saponins and thus natural foaming abilities. The pericarps also find applications in laundry detergents and in Ayurveda and folklore medicine for treating asthma, skin diseases and psychiatric disorders (Kora, 2023).

This poster presentation provides a comprehensive overview of the isolation protocol used to obtain crude saponin mixtures from each plant, beginning with microwave-assisted extraction and incorporating various chromatographic techniques including VLC, flash chromatography, and MPLC. Furthermore, it assesses their capacity to activate the sweet taste receptor TAS1R2/TAS1R3, offering insights into their potential effects.

Keywords: Glycosides, *Siraitia grosvenorii*, *Sapindus rarak*, *Albizia myriophylla*, TAS1R2/TAS1R3 receptor

References

Medicinal Plant Names Services Portal, Royal Botanic Gardens, Kew, Version 13, accessed on 10/04/2024.

Gong X., Chen N., Ren K., Jia J., Wei K., Zhang L., Lv Y., Wang J., Li M. 2019. The Fruits of *Siraitia grosvenorii*: A Review of a Chinese Food-Medicine. *Front Pharmacol.* 10, 1400.

Joycharat N., Boonma C., Thammavong S., Yingyongnarongkul B.E., Limsuwan S., Voravuthikunchai S.P., 2016. Chemical constituents and biological activities of *Albizia myriophylla* wood. *Pharm Biol.* 54, 62-73.

Kora A. J., Inamuddin, Adetunji C.O. (Eds). 2023. Applications of Next Generation Biosurfactants in the Food Sector, Chapter 22, 459-477. Academic Press

* Presenting author. E-mail address: antoine.bruguere@u-bourgogne.fr (Antoine Bruguère)

S3.P269 Recovery effect of phloroglucinol 1-O-β-D-glucopyranoside from the brown alga *Agarum clathratum* subsp. *yakishiriense* on neomycin-induced otic hair cell damage in zebrafish

Yukyoungh Jeon^a, Youn Hee Nam^b, Tong Ho Kang^b, Jong Hwan Kwak^{a,*}

^a School of Pharmacy, Sungkyunkwan University, Suwon 16419, Korea

^b Department of Oriental Medicine Biotechnology, College of Life Sciences, Kyung Hee University, Gyeonggi-do 17104, Korea

Marine plants are popular and abundant food ingredients mainly in Asian countries, and also known to have many beneficial biological activities. *Agarum clathratum* subsp. *yakishiriense* Yamada ex

G.H. Boo & P. C. Silva is a perennial brown alga in the family Costariaceae and is widely distributed in the coastal area of the East Sea in Korea (Boo et al., 2012). Previous studies on *A. clathratum* have reported neuroprotective effect (Kim et al., 2014) and anticancer effect on epithelial ovarian cancer (Kim et al., 2023).

As part of our continuing search to find potent anti-ototoxic compound, we investigated the recovery effect of hearing loss for EtOH and water extracts of *A. clathratum* subsp. *yakishiriense* on a zebrafish model. The EtOH extract of *A. clathratum* subsp. *yakishiriense* which showed ameliorative effect in a zebrafish ototoxicity model, was suspended in water and then partitioned consecutively with organic solvents to give CH₂Cl₂, EtOAc, *n*-BuOH, and H₂O fractions. Among these fractions, the H₂O fraction revealed the strongest protection against the otic hair cell damage in zebrafish models. The water fraction was subjected to column chromatographic separation. Phloroglucinol 1-O-β-D-glucopyranoside along with five compounds were isolated from the water fraction of *A. clathratum* subsp. *yakishiriense*. The isolated compounds were investigated the ameliorative effect on neomycin-induced ototoxicity in the zebrafish. Phloroglucinol 1-O-β-D-glucopyranoside showed significant recovery on otic hair cell damage at concentration of 1 μg/mL. In addition, its mechanism of action was studied. In conclusion, phloroglucinol 1-O-β-D-glucopyranoside revealed potent recovery effect for the hearing loss.

Keywords: *Agarum clathratum* subsp. *yakishiriense*, Costariaceae, phloroglucinol 1-O-β-D-glucopyranoside, neomycin-induced ototoxicity, zebrafish

References

- Boo, S.M., Ko, Y.D., 2012. Marine Plants from Korea. Seoul, Junghaeng-Sa, 114.
- Kim, I.H., Yoo, K.-Y., Park, J.H., Yan, B.C., Ahn, J.H., Lee, J.-C., Kwon, H.M., Kim, J.-D., Kim, Y.-M., You, S.G., Kang, I.-J., Won, M.-H., 2014. Comparison of neuroprotective effects of extract and fractions from *Agarum clathratum* against experimentally induced transient cerebral ischemic damage. *Pharm. Biol.* 52, 335-343.
- Kim, L., Hong, T., Ham, J., Lim, W., 2023. Effects of *Agarum clathratum* extract on cell death and calcium ion levels of ovarian cancer cell. *Mol. Cell. Toxicol.* 19, 303-310.

Acknowledgements

This work was supported by the National Research Foundation of Republic of Korea (NRF), and the grant was funded by the Korea government (2021R1A2C1014595).

* Presenting author. E-mail address: jhkwak@skku.edu (Jong Hwan Kwak)

S3.P270 Giuseppe Raddi and the plants: Taxonomy, Uses and Conservation in the 21st century

Juliana de Paula-Souza^a, Rafael Raddi^{b,*}

^a Botany Department, Federal University of Santa Catarina, Rua Eng. Agrônomo Andrei Cristian Ferreira 216, 88040-535, Florianópolis, SC, Brazil

^b Comitato Giuseppe Raddi - Ludwig-Maximilians-Universität München, Geschwister-Scholl-Platz 1 80539 München, Germany

Giuseppe Raddi was an Italian naturalist who surveyed the flora of Rio de Janeiro in 1817, producing a rich collection of around 4000 plant specimens. The difficulties in studying historical collections were greatly attenuated by the digitization of biological collections and libraries, but the volume of materials produced by Raddi combined with particularities of his life history poses a greater challenge in the study of his work.

The present work aims a brief assessment of the relevance and contribution of Raddi in plant taxonomy, and, given the impossibility of tabulating for analysis all the specimens collected by him, we will use the names he described to bring his work to the 21st century reality, in relation to the uses made of the species reported by him and their confirmation by science, and their state of conservation, almost 200 years after his death.

We used the main online botanical databases to search for names containing “Raddi”, resulting in 368 names which were checked as to their priority and legitimacy, generating a list of 273 names that correspond to the universe of taxa dealt by Raddi, 236 of them occurring in Brazil.

Evidence of Raddi’s brilliance is the fact that he navigated with equal dexterity every group of land plants known to science, and that his name has been celebrated in 40 names of taxa, during his lifetime and after. His dedication to non-flowering plants, however, might explain the lower number of medicinal plants described in his works. From the 236 Brazilian taxa surveyed, only 20 had their conservation status evaluated, evidencing the size of Brazil’s natural heritage and our responsibility towards future generations.

Keywords: biodiversity, conservation, historical botany, Giuseppe Raddi, medicinal plants

*Presenting author. E-mail address: raddi@giusepperaddi.org (Rafael Raddi)

Index of Authors

Index of Authors

Symbole

, Gertie Janneke Oostingh 681
' ztürk, Meltem 806

A

Aas, Gregor 595
Abbasi, Anham 800, 802
Abbee, Pieter Van den 104
Abdallah, Rola 662
AbdelKader, Maged 815
Abdel-Kahaar, Emaad 923
Abe, Ikuro 53
Aberkane, Fatima 743
Abreu, Lucas Silva 397
Abril, Sarabia Dominguez Jazmin 613
Abubakar, Mustapha Ngaski 62
Acar, Ayten 462
Acero, Nuria 760, 761
Achard-Baccati, Clémentine 783
Acharya, Deepa D. 481
Acquaviva, Alessandra 1034
Acquaviva, Rosaria 1015
Adaikpoh, Barbara I. 798
Adamska-Szewczyk, Aldona 474
Adamu, Ramadhani 393
Adebisi, Oluwaseun 762
Adegoke, Abdullahi Adeyemi 882, 883
Adekomi, Adedayo 762
Adeleke, Oluwakemi 513
Adelsberger, Sigrid 525
Adenmosun, Adeola 762
Adepiti, Awodayo O. 826
Adewale, Omowumi 762, 763
Adeyem, Abiodun 358, 656
Adhami, Hamid-Reza 108, 531
Adinehpour, Fatemeh 835
Adnan, Noorazlina 373, 374, 446, 447
Adomou, Aristide 48
Aeri, Vidhu 916
Afantitis, Andreas 1019, 1021
Afonso, Carlos A.M. 33
Afoullouss, Sam 250, 325
Agarwal, Amit 971

Agbaje, Maryam Titilope 883
Agbankpé, Jérrold 48
Aguiñiga-Sánchez, Itzen 563
Agusti, Susana 285, 286, 288, 292
Ah Kim, Jeong 564
Ahmad, Sayeed 759
Ahmadu, Augustine A. 559
Ahmaduc, Augustine A. 558
Ahmat, Norizan 373, 374, 446
Ahmed, Mohammed M. A. 674
Ahmed, Shopno R. 493
Ahn, Hyojin 70
Ahn, Mi-Jeong 390
Ainalidou, Aggeliki 958
Ajibade, Adefisoye Martins 438
Ak, Güneş 609
Akhgari, Amir 130
Akinboye, Ayodele O. 753
Akinlade, Mary Ololade 883
Akissi, Zachee L. E. 748
Akrap, Karla 704
Aktar, Jesmin 63
Alamro, Abir 738
Alan-Albayrak, Elif 318, 319
Al-Awadhi, Fatma H. 346
Al-Awadh, Saeed 727
Albarico, Genesis 340
Albayrak, Gokay 319, 324
Albertin, Warren 1009
Albrecht, Larissa 846
Alderotti, Francesca 889
Aldini, Giancarlo 197, 212
AlDosari, Abdullah 815
Alehaideb, Zeyad 738
Alencar, Severino de 663
Alexander, Kelsey L. 260
Alexander, N. Reed 68
Algaffar, Shereen Abd 436, 78
Algaffar, Shereen O. Abd 410
Al-Hemedawi, Ryath 778
Al-Hindy, Momen 781
Al-Hmadi, Hekmat B. 37, 625
Ali, Abbas 955
Aliberti, Michela 1012
Aligiannis, Nektarios 329, 873, 1019, 1021
Ali, Hazhmat 339
Alilou, Mostafa 299, 304
Ali, Zulfiqar 415, 955
Alla, Chaimae 748
Allais, Florent 921
Allard, Pierre-Marie 816, 131, 901
Allegrini, Paolo 1019

Allegrini, Pietro	951	An, Joon Soo	400
Allison, Ledoux	377, 913	Annamneedi, Venkata Prakash	615
Allodi, Marika	942, 967	Annang, Frederick Boye	77
Al-Maddboly, Lamiaa	458	Antoine, Marie-Hélène	229
Al-Madhi, Faisal A.	727	Antoniadi, Ioanna	172
Almaliti, Jehad	781	Antoniadi, Lemonia	571
Almeida, Joana	33	Antonopoulos, Romanos K.	329
Almogren, Aljowhara	738	Anukanon, Shisanupong	880
Alonso, Cristina	1031	Anwar, Zeeshan	404
Alqahtani, Omaish	456	Anywar, Godwin	47, 719
Alruzaihi, Marwah	738	Apel, Cécile	258, 389, 376
Alsharidah, Amani	738	Arabi, Shiva Pouramin	835
Alsharif, Bashaer	578, 579	Arampatzis, Athanasios	829
Alshehri, Osama H.	727	Ara, Tabassum	367
Al-Suliman, Bader A.	727	Araya-Cloutier, Carla	206, 237, 383
Alsup, Tyler	469	Arbelaez, Natalia	545, 797
Altomare, Claudio Fabio	303	Ardeza, Iron Mike	160
Altucci, Lucia	302	Areesanan, Alexander	593
Aluta, Uzeme P.	557	Argyropoulou, Aikaterini	230, 992, 1001
Alvanoudi, Panagiota	731	Arias-Duran, Luis	614
Álvarez-Rodríguez, Javier	45	Armagan, Guliz	640
Alves, Luís	111	Armengol, Eva Sanchez	84
Al-Yafei, Thoraya	347	Army, Monika Kerry	351
Amakorová, Petra	878	Arnaudin, Ingrid	189
Amand, Séverine	85	Arroo, Randolph	133
Amanlou, Massoud	108	Arsenijević, Jelena	395
Ambe, Gael Noel Neh Neba	654	Arturo-Schaan, M.	751
Ambrosio, Sergio Ricardo	785	Arundhathi, Dev J.R.	538
Amerikanou, Charalampia	142	Arundhathi, Dev JR	537
Amin, Sahar	560	Asali, Mazen Mounir Abu	347
Ammar, Ramy M.	635	Asekunowo, Anita Kelechi	460
Amon, Denick	417	Ashmawy, Naglaa	436
Amos-Tautua, Bamidele Martin	368	Asiimwe, Savina	719
Anacak, Gunay Yetik	319	Asi, Ranelle Janine	886
Anagnostou, Maria	868	Aslan-Erdem, Sinem	768
An, Chae-Yeong	566	Assimopoulou, Andreana	827, 829
Andersen, Aaron J.C.	386	Assimopoulou, Andreana N.	261, 731
Andersen, Aaron John Christian	365	Astrid, Schmitz	913
Andersen, Jeanette Hammer	341	Ataputta, Nirasha	205
Andersen, Raymond J.	278, 860	Atmani, Djebbar	512
Andersson, Håkan S.	295, 414	Attia, Joan	924
Andolfi, Anna	210, 405	Attia, Rym	451, 964
Andrade-Cetto, Adolfo	666, 667	Aucejo, Ángela Morellá	989
Andrén, Per E.	707	Aucher, Willy	416
André, Rebeca	819	Audo, Gregoire	700
Andrés, M. Fe.	255	Audo, Grégoire	885, 968
Angelis, Apostolis	571	Audoin, Coralie	277
Angelov, Andrea	431	Augustinović, Mario	141
Angenot, Luc	54	Augustynowicz-Kopeć, Ewa	408
Angielczyk, Mirosław	399	Augustyn, Wilma	840
An, Hye In	962	Aumer, Carola	568
An, Hyo-Jin	220	Autzen, Sabrina	747



Bedran-Russo, Ana K.	893	Bifulco, Giuseppe	623, 999, 1004, 1012
Beemelmaans, Christine	323	Bigliati, Mauro	771
Beerhues, Ludger	87, 88	Bignard, Mikaela	1037
Beest, Ellen ter	383	Bildstein, Augustin	74
Behsaz, Bahar	775	Bilela, Lada Lukić	182
Belen, Nazlican	319	Bilgi, Çiğdem	581, 612
Bellanger, Jean-Michel	65	Bilgi, Eyüp	265
Belloir, Christine	929, 1039	Bilia, Anna Rita	333, 871, 875, 889, 890, 137
Belofsky, Gil	70	Binev, Biser	599
Belouzard, Sandrine	74	Binev, Biser K.	598, 600, 603
Benarbiab, Amine	363	Bing, Jade	481
Benarbia, Mohammed El Amine	726, 743	Bínová, Zuzana	735, 995
Bendel, Pamela	510	Birim, Dervis	640
Bender, Andreas	790	Bisson, Jean-François	584
Bender, Cecilia	952	Blaheta, Roman A.	846
Benedetti, Giada	241	Blainski, Andressa	230, 728, 730
Beneš, František	735, 995	Blair, Taylor	744
Beniddir, Mehdi A.	806, 818, 824, 56, 65, 72	Blanchard, Patricia	55
Béni, Zoltán	213	Blanchemain, Nicolas	1000
Benjamin, Palmaerts	377	Blanchet, Marine	73
Berenice, Aguilar Guadarrama Angelica	613	Blanchfield, Joanne	221
Berger, Leonie	323	Blanco-Penedo, Isabel	45
Berger, Tim	316, 450, 872	Blasi-Romero, Anna	424
Bergfeld, Arne	253	Blažeković, Biljana	331, 332
Bergonzi, Maria Camilla	871, 137	Blee, James	406, 749
Berjanski, Mark	888	Blicharska, Natalia	741, 970
Berkecz, Róbert	986	Blodgett, Joshua A.V.	193
Berlinck, Roberto G. S.	486	Blondeau, Claude	946
Berlioz-Barbier, Alexandra	903	Blon, Nadège	258, 700
Bernadat, Guillaume	824	Blumenthal, Mark	94
Bernardos, Andrea Bau	989	Boakye, Yaw Duah	417
Bernhard, David	980	Boatey, Susana Dufie	417
Bernieri, Anna	546	Bocquet, Laetitia	74
Berry, Olivier	263	Bodó, Enikő	213
Berscheid, Anne	79	Boesger, Hannah	560
Bertaina, Lorenza	836	Bogalho, Hugo	242
Berthomier, Lucile	946	Bogdanov, Alexander	185
Berthon, Jean-Yves	240, 938	Boisard, Séverine	55
Bertolotti, Luigi	750	Bojarová, Pavla	82
Bertoni, Francesco	359	Bojkova, Denisa	72
Bertrand, Samuel	231, 257, 263, 830	Boleti, Haralabia	841, 166
Bescós, Paloma Bermejo	999	Bollaro, Flavia	192, 240
Beteinakis, Stavros	845, 847, 1001	Bollen, Melissa	520
Bever, Madison	997	Bolletta, Valentina	703
Bhambra, Avninder	654	Bonaterrea, Gabriel A.	449, 652
Bharathi, Avula	415	Boneva, Blagovesta D.	599
Bialas, Elena	494	Bonnassieux, Justine	414
Białas, Marcel	622	Bonnet, Olivier	196, 54
Bianchi, Simone	1015	Bonnet, Susan L.	812
Biasiotti, Matilde	579	Bonnet, Sylvestre	76
Biccigo, Mayra	510	Bonn, Guenther K.	51
Biccigo, Mayra Gabriella	518	Bonn, Günther K.	100

Boo, Hye-Jin	752, 988	Briand, Loïc	929
Boonen, Georg	97, 98, 162	Bringmann, Gerhard	78
Boonsut, Rinnapat	542, 925	Brissoit, Alice	192
Boppré, Michael	122	Brito, Ângela	501
Boratyński, Filip	219, 83	Brodsky, Katerina	82
Borges, Ana	632, 852	Bronze, Maria do Rosário	491
Borges, Warley de Souza	200, 904	Brötz-Oesterhelt, Heike	79
Borie, Nicolas	38, 164	Brown, Paula N.	57
Borkova, Lucie	74	Brożyna, Malwina	394
Bornikoel, Jan	79	Brück, Thomas	795
Boroduškis, Mārtiņš	1030	Brüelhart, Edouard	816
Borromeo, Rod Vincent	886	Bruguière, Antoine	55, 929
Bory, Alexandre	254	Bruijn, Wouter de	206
Bosch-Morell, Francisco	761	Brulefert, Adrien	946
Bosch-Navarrete, Cristina	77	Brunel, Jean-Michel	73, 384
Bösch, Nina M.	28, 854	Brunelle, Alain	216
Bosisio, Stefania	951	Brunetti, Cecilia	889
Bossard, Elodie	1001	Brunetti, Luigi	1034
Bot, Maxime Le	726	Bruni, Natascia	750
Bouderbala, Salima	160	Bruyère, A.	751
Boudesocque-Delaye, Leslie	207, 233, 688, 814, 815, 160	Brzóska, Małgorzata M.	642
Boudjelal, Mohamed	738	Bucar, Franz	407, 441
Boudreau, Paul D.	674	Bucquial, Leopoldo	444
Bouguellid, Ghania	512	Budai, Anthony	164
Bouhlo, Romain	968	Budnik, Kornelia	867
Boujedaini, Naoual	584	Budzy, Elea	688
Bouloumpasi, Elisavet	714	Bueno, Paula Carolina Pires	360
Bourgougnon, Nathalie	968	Bufaino, Eliana Martino	728, 730
Bourguet-Kondracki, Marie-Lise	73, 384	Bugayong-Janagap, Adrienne Marrie	444
Bourguignon, Anna	820	Bui-Hoang, Minh	36
Bousarghin, L.	244	Bui, Nguyen-Bien-Thuy	307
Boustie, Joël	192	Buisson, Didier	85
Boutet, Stéphanie	389	Bujor, Alexandra	655
Bove, Carmen	999	Bülöw, Fabian	487
Boylan, Fabio	578, 579, 641, 820, 965	Bülöw, Fabian E.	543
Bozinovikj, Elizabeta Krstevska	780	Bültemeier, Arne	478
Bozkurt, Buket	577	Bungar, Jeffrey Clement	886
Braca, Alessandra	241, 392, 675	Bun-Llopet, Sok-Siya	782, 963
Braga-Henriques, Andreia	182	Bunnell, Amelia	828
Bratiichuk, Dmytro	480	Buommino, Elisabetta	411, 422
Brauner, Annelie	424	Burch, Jessica E.	838
Braun, Theodor	100	Burdet, Frederic	901
Bréard, Dimitri	55	Burdette, Joanna E.	535, 659
Breud, Célia	782	Burkart, Martin	121
Breinlinger, Steffen	190	Burroughs, Megan	115
Bremer, Swen	97	Bushman, Timothy J.	291
Brendel, Fabian L.	478	Bussmann, Hendrik	97
Bren, Urban	442, 443, 445	Bustos-Brito, Celia	777
Breynaert, Annelies	84	Butaud, Jean-François	682
Briand, Enora	263	Butdapheng, Khunkhang	471
Briand, Loic	1039	Butera, Gisela	977
		Buț, Mădălina-Georgiana	981

Butterweck, Veronika	97, 98, 162	Carlucci, Vittorio	862
Bu, Youngmin	565, 586	Carne, Beatrice De	942, 967, 969
Bwaila, King	848	Carneiro, Ariadne Magalães	695
Bye, Robert	165	Carneiro, Ariadne Magalhães	694
Bystrom, Laura	696	Carocho, Marcio	490
Bytešníková, Zuzana	465	Carocho, Márcio	491, 1003
Byun, Woong Sub	551	Caro, Yanis	226
		Carranza-Aranda, Ahtziri Socorro	544
		Carreira, Erick	901
		Carr, Jeanmarie	1017
		Carvalho, Aida	959
		Carvalho, Aline	820
		Carvalho, Isabel S.	983
		Carvalho, João E.	510
		Carvalho, João Ernesto de	518
		Carvalho, Ludmilla Tonani	296
		Carvalho, Luís A. E. Batista de	959
		Carvalho, Maria F.	182, 191
		Carvalho, Paulo Roberto	510
		Casadellà-Xifra, Josep	45
		Casapullo, Agostino	1004, 1012
		Castañeda-Loaiza, Viana	227, 957
		Castillo, Francisco	668
		Castro-Falcón, Gabriel	79
		Cathey, Lucas	828
		Čechová, Jana	465
		Cécile, Apel	380
		Cedergreen, Nina	113
		Celep, Engin	572
		Cerulli, Antonietta	999
		Cespi, Marco	703
		Chae, Hee-Sung	161
		Chaichamnong, Nattiya	604
		Chaiwangrach, Nathareen	900
		Cha, Jin Wook	249
		Chaker, El Kalamouni	380
		Chalkiadaki, Maria	992
		Chambaud, Marine	683, 950
		Chambon, Marion	164
		Cha, Min-Ho	402, 403
		Champavier, Yves	189
		Champy, Pierre	196, 984
		Chana, Ju-I	671
		Chandra, Amit	94
		Chandrasekaran, Sriram	837
		Chandra, Sona	790
		Chang, Chia-Chuan	195, 473, 521
		Chang, Chi-Ting	773
		Chang, Fang-Rong	425, 788, 789, 937
		Chang, Meng-Ting	773
		Chang, Pi-Kai	831
		Chang, Tsu-Yuan	195, 521
Cabaça, Daniela	943		
Cabalfin, Zeb Jose Joshua	444		
Cabec, Audrey Le	944		
Cabral, Célia	809, 959		
Cabrera-Pardo, Jaime Roberto	336		
Cadelis, Melissa M.	73, 384		
Cadena-Iñiguez, Jorge	544, 563		
Cadoret, Justine	885		
Caffrey, Conor	296, 297		
Cahlíková, Lucie	151		
Cailloux, Olivier	806, 818		
Çakılcıoğlu, Uğur	609		
Cala, Antonio	171		
Calcul, Laurent	250		
Caldwell, Kim A.	291		
Calhelha, Ricardo	242		
Calhelha, Ricardo C.	491, 632, 852		
Caljon, Guy	47		
Calvo, María Luisa Martín	632, 852		
Camangi, Fabiano	675		
Camara, Aboubacar Kadiatou	379		
Camara, Mohamed Kerfalla	379		
Câmara, Paulo Eduardo Aguiar Saraiva	709		
Campos, Pierre-Eric	692, 816, 944, 947, 1037		
Camuzet, Charline	74		
Canaan, Stéphane	782		
Caneschi, Alice	769		
Caniça, Manuela	327, 328		
Cankaya, I. Irem Tatli	624		
Canoyra, Antonio	760		
Cantrell, Charles L.	161		
Cao, Shugeng	193		
Cappellacci, Loredana	152		
Capucchio, Maria Teresa	44, 771, 750, 123		
Carafa, Vincenzo	302		
Cárdenas, Denny	545		
Cárdenas, Pablo D.	113		
Cárdenas, Paco	274		
Cardoso, Weverson Cavalcante	200		
Carla, Hamann	736		
Carlson, Skylar	264		

C

Chang, Yuan-Shiun	800	Chikowe, Ibrahim	848
Chan, Jason Ying-Kuen	936	Chik, Wai-I	895
Chanut, Stéphanie	583, 584	Chini, Maria Giovanna	134, 1012
Chao, Chih-Hua	307	Chinou, Ioanna	873
Chao, Chi-Hsein	521	Chin, Young-Won	566
Chaput, Dale	250, 325	Chiocchio, Ilaria	136
Charline, Leroy	736	Chłopecka, Magdalena	627
Charlotte, Simmler	968	Cho, Byong Ok	926, 927
Charoensup, Rawiwan	278, 858, 860, 866	Cho, Byoung Ok	928
Charpentier, Thomas	258, 908	Choi, Hee-Jeong	402
Chassagne, François	523, 682	Choi, Hyuk Joon	833
Chathuranga, Kiramage	426, 427	Choi, In-Wook	547
Chatterjee, Swarnali	997	Choi, Jaewoo	98
Chatzopoulou, Paschalina	714	Choi, Jin Won	313
Chaudhary, Mridul kant	519	Choi, Minsik	725, 175
Chaudhary, Mridul Kant	677	Choi, Misnik	961
Chauvin, Isabel M.	260	Choi, Sang Yoon	536, 547
Chavez, Benjamin G.	478	Choi, Young H.	298
Che, Chun-Tao	876	Choi, Young Hae	176
Cheenpracha, Sarot	860	Choi, Youn Kyung	697, 910
Cheilari, Antigoni	329, 873, 1021	Choi, Yu Kyung	249
Chemat, Farid	934	Cholke, Kaushavi	413
Chemin, Guillaume	189	Cholke, Kaushavi Anil	629
Chen, Bai-Sen	895	Chollet-Krugler, Marylène	192
Chen, Chi	505	Chollet, Sébastien	38
Chen, Ching-Yeu	307	Chollet, Sylvie	74
Chen, Chun-Lin	180, 186, 779	Choonong, Rattanathorn	471
Chene, Gérald	946	Choquesillo-Lasarte, Duane	77
Cheng, Diana Klaser	1017	Cho, Sanghyun	141
Cheng, Jing-Jy	179	Cho, Seung Hye	272
Cheng, Kun-Ching	516	Cho, Won-Kyung	403, 457, 458
Cheng, Ming-Che	195, 521	Cho, Yeo Rang	271, 272
Cheng, Ming-Chu	773	Christochowitz, Sandra	533
Cheng, Yuan-Bin	773	Christofilos, Dimitrios	827, 829
Chen, Hsi-Mei	195	Chuang, Ling	478
Chen, Hui-Ming	671	Chuang, Wu-Chang	800
Chen, Jian	685	Chuchawankul, Siriporn	757, , 167
Chen, Lo-Yun	29	Chu, Man-Hsiu	793
Chen, Mei-Chuan	307	Chung, Seungtae	721
Chen, Meng-Chin	195	Chun, Simin	551
Chen, Pai-Hsin	517	Chu, Pei-Yi	937
Chen, Shao-Nong	893, 894	Čičak, Marijo	849
Chen, Tai-Yang	671	Cicek, Serhat Sezai	694
Chen, You-Ying	180	Cielecka-Piontek, Judyta	238, 867, 869
Chen, Yu	88	Ciesla, Lukasz	291, 997
Chen, Yu-Hsin	773	Cieśla, Joanna	763
Chen, Yu-Li	425	Cihlarova, Petra	987
Cheseto, Xavier	796	Cihlářová, Petra	974, 998
Chevalley, Alia	207, 233, 679, 814	Cinatl Jr., Jindrich	72
Chianese, Giuseppina	37, 422, 625	Cindy, Lesenfans	913
Chiavaroli, Annalisa	1034	Cioni, Emily	675
Chibli, Lucas Apolinário	694, 695	Cisseb, Sekhou	363

de Freitas, Ailane S.	510	Dias, A. C. P.	1005
Degotte, Gilles	54	Dias, Alberto	1019
Dékány, Miklós	213	Dias, Alberto C. P.	633
De LaMare, Jo Anne	840	Dias, Edelson	349
Delannay, Eldra	1037	Dias, Maria Inês	959
Dela Rosa, Angelo Mc Court S.	982	Dias, Sara Leite	478
De Leo, Marinella	392, 675	Diaz, Caridad	77
Delgadillo, David A.	838	Díaz, Caridad	668
Delgado-Tiburcio, Eugenia Elisa	563	Díaz, Carmen E.	255
Delgoda, Rupika	120	Díaz-Franco, Cristina	114
Deli, Aristeia	284	Dickens, Ava	223
de Lima, Bruna Ribeiro	694, 695	Dickmeis, Thomas	785
DellaGreca, Marina	210, 405	Didara, Mislav	766
Delnavazi, Mohammad-Reza	108, 531	Dietl, Andreas	494
Delporte, Cedric	229	Dilnashin, Hagera	661
Demaye, Anaïs	216	Dimitrov, Krasimir	1000
de Melo Junqueira, Marcelo	785	Dimitrov, Vladimir	1019
Demiray, Hatice	581	Dimovski, Aleksandar	780
Demir, Gulcan	318	Dina, Evanthia	1019
Demirkiran, Ozlem	337	Dinala, Mmabatho Monica	840
Demiroz, Tugce	319	Ding, Ling	365, 706
Demir, Serdar	319	Dinis, Pedro	499
de Moraes, Lygia Silva	838	Dinteren, Sarah van	383
Demougeot, Céline	604, 655	Diretto, Gianfranco	1030
Demurtas, Olivia Costantina	1030	Dirks, René	237
Deneka, Maxym	874	Dirsch, Verena M.	813, 834, 836
Deng, Biao	865	Di Stasi, Mauro	392
de Oliveira, Katia Mara	785	Divoux, Laura	160
de O. Sousa, Ilza Maria	510	Djurič, Marija	445
de Paz- Silava, Sheriah Laine	886	Dobhal, Yogita	890
de Pedro, Nuria	668	Dobrzynski, Maciej	31, 554
Derbaj, Ghada	240	Dobrzyński, Maciej	555
Derbre, Séverine	397	Do, Hai Nhat	188
Derbré, Séverine	702, 55	Doležal, Karel	125
Dervishi, Malboro	113	Doll, Ursula	710
Desmarets, Lowiese	74	Dolzhko, Diana	489
De Souza, Letícia Teixeira Marcos	785	Dołomisiewicz, Wioleta	643
de Souza Silva, Jefferson	709	Domingo-Fernández, Daniel	396
Despres, Alexandra	160	Domínguez-Martín, Eva María	809
Desrat, Sandy	376, 389	Donadioa, Giuliana	135
Desrini, Sufi	416	Donadio, Giuliana	423, 701
Destandau, Emilie	683, 692, 816, 944, 950, 964, 1037	Donato, Rosa	333
De Tommasi, Nunziatina	701, 835	Donny-Clark, Aaron	546
Detti, Cassandra	889	Donoso, Kathleen Mae	444
Dévéhat, Françoise Lohezic-Le	192	Doorandishan, Mina	344
Deyaert, Stef	104	Doppler, Christian	980
Deyrup, Stephen	918	Doskocil, Ivo	340
Deyrup, Stephen T.	234	dos Santos, Raquel Alves	786
Diallo, Mamadou Saliou Telly	379	Drakopoulos, Antonios	1021
Diamantakos, Panagiotis	739, 844	Drakoulis, Nikolaos	994
Diana, A. Barrera A.	776	Dramis, Martina	197, 225
		Drebka, Alexandra	646, 647

Dresler, Sławomir	914	Edrad-Ebel, RuAngelie	367
Dreux-Zhiga, Assia	240	Edwardsen, Bente	386
Drewe, Jürgen	97, 162	Efferth, Thomas	809
Driscoll, Daniel	678	Eglseer, Karin Zitterl	770
Drobac, Milica	395, 605, 606	Eichenauer, Elisabeth	509
Duangyod, Thidarat	278, 860	Ekaterina-Michaela, Tomou	333
Duan, Yao-Tao	60	Ekejiuba, Chioma M.	638
Duanyong, Thidarat	866	Ekiert, Halina	306, 338, 1007
Duarte, Carlos M.	285, 286, 288, 292	Ekuadzi, Edmund	417
Duarte, Rodrigo dos Santos Pinto	821	Elahi, Ava	670
DuBore, Ericka	70	Elansary, Hosam O.	308
Dubuisson, Jean	74	El-Awaad, Islam	87
Duca, Laurent	963	Elevado, Danielle	444
Dugay, Annabelle	397	ElGamal, Ali	815
Duggan, Brendan	64	Elhabiri, Mourad	74
Du, Guohao	865	El-Morsi, Rasha	458
Dumonceaux, Tim	317	El-Salam, Mohamed Abd	458
Duncan, Katherine R.	67	El-Sayed, El-Sayed R.	83
D'Unienville, Amandine	699	El-Sayed, R.	219
DunkiĆ, Niko	704	Elsayed, Somayah	499
DunkiĆ, Valerija	607, 698, 704	Elvert, Christian	826, 980
Dunman, Paul M.	418	Emami, S. Noushin	414
Durán, Alexandra G.	322	Embry, Michelle	863
Durazzo, Alessandra	977	Emeline, Houël	801
Duroux, Romain	924	Emilie, Pinault	232
Dürr, Lara	31, 554, 555	Emokpae, Osagie	291
D'Urso, Gilda	1004	Engler, Robin	901
Dutra, Jessyca Aparecida Paes	200	Enslin, Gill	39
Duvaud, Lauriane	534	Eparvier, Véronique	110, 378
Duweb, Amira	923	Erazo, Gabriela	610
Dworak-Schultz, Inken	47	Erdogan, Gizem	768
Dwyer, Johanna	977	Erguc, Ali	324
Dyar, M. Darby	713	Ergün, Sebahattin	127
Dychiao, Garret Royce K.	982	Eric, Hallot	377
Dyer, Lee A.	115	Eriksson, Camilla	433
Dymarska, Monika	364, 894	Ernst, Lukas	87, 88
Dziągwa-Becker, Magdalena	394	Erol, Özlem	298
Dziurka, Michał	308	Escandón-Rivera, Sonia	666

E

Ebada, Sherif Elsayed	454
Ebadi, Mohammad-Taghi	514, 658
Ebel, Rainer	67
Ebiziem, Chukwuma Joseph	372
Echevarría-Becerra, José-Luis	732
Echeverría, Javier	932
Eckert, Franziska	664
Eckert, Johannes A.	28, 854
Eckert, Nikolas	494
Edmonson, William	141

F

Fabien, Rakotondrabe Tojofaniry	887

Fàbregab, Emma	45	Filazi, Ayhan	768
Faith, Siriya	763	Filho, Edilson Beserra De Alencar	451
Fajitova, Pavla	781	Filho, Valdir Cechinel	734
Falcao, B. P.	275	Filipek, Agnieszka	276
Faleiros, Renata	786	Filipe, Marcia	819
Faller, Adam	941	Filippini, Raffaella	690
Famuyide, Ibukun M.	753	Fill, Taicia P.	486
Fandohan, Adandé Berlamain	691	Fiorentino, Antonio	300, 302, 411
Fan, Minxia	887	Fiorito, Filomena	405
Fanny, Roussi	380	Firooznia, Fardad	44
Fantasma, Francesca	1004, 1012	Flade, Jens	657
Farag, Mohamed A.	133	Fladerer, Johannes-Paul	441, 864
Faramarzi, Mohammad Ali	531	Fleury, Fabrice	231
Faraone, Immacolata	862	Flores-Bocanegra, Laura	165
Faruk, Mohamed Abbas Umare	401	Flores-Vallejo, Rosario	201
Faruque, Mohammad Omar	369	Flori, Lorenzo	241
Farzamarzi, Mohammad Ali	108	Foglio, Mary Ann	510, 518
Farzaneh, Avishan	108	Foitová, Ivona	125
Faske, Trevor M.	115	Fokialakis, Nikolaos	1019
Fauconnier, Marie-Laure	74	Fokialakis, Nikolas	329, 330, 500, 1018
Faustino, Rita	491	Fon, Mathias	386
Fausto, Rui	497	Fonseca, Ana C.	294, 295, 168
Favrod, Sina	856	Fonseca-Rodrigues, Diana	1005
Feineis, Doris	502	Fontaine, Jean-Xavier	74
Feistel, Björn	657, 952	Fontaine, Véronique	824
Fekner, Zygmunt	463	Fontana, Anthony	941
Felegyi-Tóth, Csenge Anna	213	Formato, Marialuisa	1031
Felice, Vincenzo De	1004, 1012	Formisano, Carmen	623
Feng, Yunjiang	107, 439	Fornara, Dario	897, 935
Fenical, William	246	Forte, Claudio	123
Fermišek, Miha	445	Foster, Adam	66
Fernandes, Catarina	111	Foster, Kimberley	120
Fernandes, Isabel P.	242	Foubert, Kenn	84
Fernandes, Yan	349	Frana, Sergio Alexandre	124
Fernández-Godino, Rosario	668	Franc, Ales	44
Fernández, Naiara	491	François,Louvet	232
Ferrante, Claudio	1034	Francotte, Pierre	54
Ferrati, Marta	152	Franke, Jakob	478
Ferraz, Natalia	424	Franken, Sebastian	97
Ferreira, Antonio G.	486	Franssen, Maurice	206
Ferreira, Leonor	294, 990	Franzblau, Scott G.	141
Ferrier, Manon	1007	Fraternal, Daniele	701
Ferrini, Francesco	889	Fratini, Filippo	392
Ferron, S.	244	Frederich, Michel	440
Fester, Karin	502	Frédérich, Michel	196, 861, 54
Fetter, Artem	67	Freitas, Ailane Souza de	518
Fiala, Johannes	76	Freitas, Anibal	297
Fiebich, Bernd L.	211	Frešer, Franjo	442
Figueiredo, Laisla Rodrigues	786	Freymann, Elodie	47
Figueiró, Harley Cardoso	694	Freytag, Virginie	97
Figueroa, Mario	246	Fricke, Anna	360
Filaire, Edith	938	Froment, Luke	688

Fuchs, James R.	535, 659
Fuglsang, Anja	113
Fuhrmann, Gregor	825
Fulco, Umberto Laino	75
Fumuso, Fernanda	453
Funari, Cristiano Soleo	694, 695
Funez, Luiz Adriano	734
Funmilola, Akomolafe Seun	459
Fürnkranz, Ursula	299, 304
Fu, Xiaocong	909
Fu, Yanan	865
Fyhrqvist, Pia	355, 464, 466

G

Gabriela, Ávila-Villarreal	613, 732
Gadar, Kavita	404
Gadiya, Yojana	396
Gahounga, Fabrice Jean Eudes Crédo	691
Gai, Francesco	44
Gaitanis, Konstantinos	1018
Galanty, Agnieszka	214, 218, 940, 945
Galica, T.	275
Gallah, Umar Shehu	339
Gallard, Jean-François	824
Galperin, Ryan	70
Gałazyn-Sidorczuk, Małgorzata	642
Gamage, Nuwan	428
Ganesan, A.	357
Gantert, Thomas	549
Gao, Si	936
Garayev, Elnur	782, 783, 934, 963
Garbe, Leif-Alexander	47, 121
Garbiec, Ewa	867
Garcés, Any C.	157
García-Contreras, Rodolfo	246
Gardan-Salmon, D.	751
Gardikis, Konstantinos	1019
Gardiki, Vassiliki	1011
Gargiulo, Ernesto	422
Garg, Munish	774
Garg, Neha	148
Garigliany, Mutien-Marie	412
Garner, Amanda L.	775
Garó, Eliane	31, 99, 554, 555
Garrett, Rafael	820
Garros, Laurine	964
Gaslonde, Thomas	85
Gaspar, Lorena Rigo	296
Gattringer, Jasmin	669
Gaudry, Arnaud	901, 131

Gautam, Vasuk	888
Gauthier, Leah	317, 66
Gauthier, Nathalie	743
Gaweł-Bęben, Katarzyna	1002, 1028
Gawlik, Urszula	644
Gbaguidi, Fernand Ahokanou	899
Gbemu, Makafui Adzo	417
Geetha, Rajeswari Gopal	113
Gelani, Chona D.	539
Genilloud, Olga	315, 316, 77
Genta-Jouve, Gregory	231
Genta-Jouve, Grégory	277
Gentil, Emmanuel	231
Genzel, Franziska	689
Georges, Shahad Ragheed	921
Georgiev, Milen I.590, 598, 599, 600, 601, 603, 173	
Georgoula, Eleni	500
Georgiyants, Victoriya	842
Gerber, Luc	764
Germanò, Maria Paola	675, 835
Gershenson, Jonathan	88
Gertsch, Jürg	413, 628, 629
Gerwick, Lena	251, 781
Gerwick, William H.	251, 781
Geshkovski, Nikola	630, 851
Gevrenova, Reneta	532
Gezer, Emre	265
Gębalski, Jakub	202, 745
Ghani, Nurunajah Ab.	447
Ghasemi, Ghader	658
Gheza, Gabriele	136
Ghotbi, Marjan	68
Giacomo, Claudia Di	1015
Gianniou, Despoina D.	1018
Gibbons, S.	357
Giddings, Lesley-Ann	712, 713
Giglioli-Guivarc'h, Nathalie	1007
Giglioli-Guivarch, Nathalie	964
Gigopulu, Olga	630, 851
Gilles, Ducret	232
Gil, Tae-Young	220
Gindro, Katia	254
Ginestar, José	1037
Ginigini, Joape	67
Giovannetti, Rita	703
Girão, Mariana	191
Girard, Corine	604, 655
Girardot, Marion	416
Giraud, Florence Thomas	257
Girme, Aboli	971
Girst, Gábor	817
Gisèle, E. Loe	377

Gjorgievska, Veronika Stoilkovska	780	Graczyk, Filip	548
Gjorgievska, Jovana	630	Graczyk, Piotr	639, 649
Gkiouvetidis, Petros	1001	Grafakou, Maria-Eleni	86
Gkogkou, Eirini	1018	Graf-Schiller, Sandra	764
Glasl, Sabine	509	Graikou, Konstantia	873
Glavač, Nina Kočevar	676	Granica, Sebastian	102, 309, 432, 433, 472, 479, 489, 492, 585, 153
Gloaguen, Yoann	790	Grard, Roxane	160
Glomb, Marcus A.	747	Grasu, Adina-Elena	978
Glover, Adrian	67	Gravina, Claudia	169
Glukhov, Evgenia	781	Gray, Nora	98
Godlewska, Sylwia	356	Gray, Nora E.	520
Goel, Bharat	888	Graziose, Rocky	1016
Goh, Asly Poh-Tze	870	Greaves, David R.	877
Goi, Paolo	741, 897, 935, 942, 967, 969, 970	Greff, Stéphane	217, 934, 938, 968
Gokie, Elizabeth	291	Grienke, Ulrike	525, 813, 72
Goll, Anne K. C.	253	Griffin, Elisha	150
Golubova, Daria	784	Grifoni, Lucia	333, 871, 137
Gołaszewska, Agata	650	Grigoriou, Galateia	827
Gomes, Liv Nunes Almeida	124	Grijalba, Alexander Castro	964
Gonçalves, Ana Cristina	809	Griñán-Lisón, Carmen	668
Gonçalves, Beatriz	633	Grisel, Clément	389
Gonçalves, C.	1005	Grivard, Antoine	258
Goncalves, Frédéric	583	Grohs, Lisa	47
Gonçalves, Miguel	242	Grollitsch, Selina	441
Gonçalves, Rita de Cássia Ribeiro	200	Gromek, Kamila	358, 656
Gonou-Zagou, Zacharoula	329, 500	Grondin, Antonio	468, 900, 131, 901
González, Benita	710	Grougnet, Raphael	451
González-Coloma, Azucena	255	Grougnet, Raphaël	391
González, Eduardo Villicaña	84	Grovel, Olivier	231, 263
Gonzalez, Gabriel	292	Gruba, Marcin	463
González, Ignacio	315	Gruber, Christian W.	669
Gonzalez, Luis Alberto	807	Gruber, Thomas Olaf	595
González- Menéndez, Victor	77	Grudzińska, Marta	940, 945
González-Menéndez, Víctor	668	Gründemann, Carsten	593
González-Pacanowska, Dolores	77	Grundler, Florian M. W.	689
Goodrich, Laurie	744	Grzechnik, Eryk	763
Goossens, Alain	32, 130, 155	Grzegorzczak, Agnieszka	366
Goracci, Giulia	889	Grzelczyk, Joanna	219
Göransson, U.	580, 708	Grzelka, Kajetan	684
Göransson, Ulf	274, 295, 296, 414, 433, 434, 61, 619, 143	Grześkowiak, Łukasz	585
Gori, Antonella	889	Gucwa, Magdalena	639
Gorr, Gilbert	293	Guerin, Aurélie	700
Gorzolka, Karin	382	Guerrero-Garzón, Jaime Felipe	184
Gostner, Johanna M.	826	Guerrini, Alessandro	746, 767
Götz, Sinja	409	Gu, Eun-Young	720
Goulart, Yara	242	Guevara, Elizabeth	245
Gourguillon, Lorène	1001	Guével, Remy Le	963
Gouveia, Goncalo	839	Guével, Rémy Le	934
Górska, Renata	362	Guilet, David	726, 795
Grabowska, Karolina	214, 218	Guillemette, Thomas	908
Gracz-Bernaciak, Joanna	504	Guillemette, Thomas	258

Heleno, Sandrina A.	242, 490, 491, 632, 852, 1003	Holm, Meredith	287
Helen, Skaltsa	333	Holowacz, Sophie	946
Helf, Maximilian J.	839	Hölzl, Claudia	681
Hellhammer, Juliane	664	Hong, Junhwa	551
Hellinger, Roland	669	Hong, Suckchang	551
Hell, Tanja	31, 554, 555	Ho, Raimana	682, 934, 938, 164
Henckel, Melissa M.	582	Horgan, Mark J.	117
Hendra, Rudi	350, 351	Hornyák, Marta	252
Henneck, Alina	587	Horton, Tammy	67
Henquet, Maurice G.L.	857	Hosseini, Faezeh Sadat	108
Henriques, C. Filipe	168	Hosseini, Seyed Fakhreddin	835
Henry, Geneive E.	912	Hošťálková, Anna	591
Hensel, Andreas	104	Hostnik, Gregor	442, 443, 445
Heo, Chang-Su	156	Houari, Inès	277
Heo, Ji Hwan	927, 928	Houdkova, Marketa	340
Herath, H. M. T. Bandara	685	Houehanou, Thierry Dèhouégnon	691
Herbette, Gaëtan	934, 938, 164	Hounzangbe-Adoté, Sylvie Mawule	899
Herczogová, Petra	989	Hou, Yanpeng	1017
Herde, Marco	478	Hovtvian, Dariia	484
Hérent, Marie-France	953, 1025	Hozain, David	85
Herida, Nove Kris	444	Hristova, Hristina	1019
Hering, Anna	639, 649	Hrouzek, P.	275
Herl, Nicole	576	Hryć, Benita	902
Hermann, Lukas	790	Hsiao, Pei-Wen	671
Hermans, Nina	84	Hsin-Yi, Tang	789
Hernández, Aarón Vázquez	529, 530	Hsu, Yu-Chi	521
Hernández-Granillo, Carlos D.	246	Huang, Hui-Chi	307
Hernández, Leyre	105	Huang, Tianqi	352
Herrera-Durigneux, Tomas	378	Huber, Dominik	920
Herrera-Rodríguez, Sara Elisa	544	Huber, Florian	806
Herrmann, Fabian	382	Huber, Lukas A.	100
Herve, Fabienne	263	Hubert, Alexandra	74
Hesami, Pourya	353	Huber, Tiffany	552
Hettiarachchi, Chamari	274	Hubrich, Florian	28, 854
Heymann, Thomas	747	Hu, Dandan	101
Hilbert, Jean-Louis	748	Hughes, Chambers C.	79
Hilger, Daniel	872	Hughes, Kristelle	899, 938, 953, 1025
Hille, Katharina	36	Hu, Hao-Chun	788, 789
Hill, Kerstin	922	Hunyadi, Attila	46, 413, 788, 817
Hirsbrunner, Gaby	758, 764	Huo, Chen	267, 268
Hlengwa, Sbonelo S.	448, 452	Hur, Jae-Seon	313
Hoang, Minh Bui	117	Hur, Ji-Won	1014
Hobaiter, Catherine	47	Hur, Yegang	400
Höck, Anja	76	Husiev, Yurii	76
Hoffie, Iris	478	Hussain, Shah	100
Hoffie, Robert E.	478	Hussein, Ahmed A.	819
Hofmann, Lena	36	Husson, Cécile	229
Högger, Petra	133	Huwlyer, Felix	628
Hohmann, Judit	339, 345	Huwlyer, Jörg	820
Holandino, Carla	820, 821, 846	Hu, Yifei	193
Holding, Matthew	287	Huymann, Lesley	76
Holinger, Mirjam	764	Huynh, Kim-Dung	99

Hwang, Bang Yeon	312
Hwang, Hyung Yeon	725
Hwang, Jinhee	729
Hwang, Su Jung	962
Hwang, Tsong-Long	29, 425
Hyeon, Hyejin	752, 988
Hymete, Ariaya	506
Hyun, Jaehae	721
Hyun, Jin Won	697

I

Iannotti, Fabio Arturo	37, 625
Icenhour, Daniel	90
Idborg, Helena	619
Ignea, Codruta	60
Igoli, John Ogbaji	460
Iker, Tabitha	47
Ilić, Milan	605
Imbert, Christine	416
Im, Hong Sik	962
Inçi, Nejla	534
Ingalalli, Vijay	790
Ingallina, Cinzia	1034
Ingebrigtsen, Richard A.	386
Ingkaninan, Kornkanok	604
Inngjerdingen, Marit	431
Inoue, Hirofumi	541
Ioannou, Efstathia	285, 286, 288, 292, 293, 60
Iorizzi, Maria	1004, 1012
Ioset, Jean-Robert	315
Iovane, Valentina	405
Ióca, Laura P.	486
İpek, Gül	944
Ippolito, Dorotea	44, 123
Ipulan-Colet, Lerrie Ann	982
Irakli, Maria	714
Irianti, Marina Ika	383
Irvine, William	120
Isaksson, Johan Mattias	341
Isca, Vera M. S.	33, 819
Iseki, Masanori	541
Ishihara, Katsuhiko	541
Iskandar, Benni	791
Islam, Nurul	327, 328
Islam, Zohorul	327, 328
Ivanova, Stanislava	590
Ivanov, Kalin	590

J

Jabal, Khadijah A.	618
Jablonsky, Ivan	259
Jablonský, Ivan	989
Jackson, Simon	741, 970
Jackson, Simon J.	897, 935
Jacob, Souopgui	377
Jacobsson, Erik	61, 295, 143
Jacquemin, Johan	233, 814
Jacquin, Justine	74
Jadhao, Mahendra	425
Jafernik, Karolina	308, 309
Jagatap, Vilas	375
Jagora, Adrien	65
Jaidee, Wuttichai	858, 866
Jaisi, Amit	342, 485
Jakimiuk, Katarzyna	472
Jakimov, Dimitar	605
Jakobsson, Per-Johan	619, 672
Jakschitz, Thomas	100
Jakubowska, Klaudia	540
Jakupović, Lejsa	705, 983
Jalil, Banaz	526, 121
Janeczko, Tomasz	364, 894
Jang, Dae Sik	203, 204
Jang, Seon Il	926, 927, 928
Jang, Yoon Seo	266
Jang, Young Pyo	723, 724, 725, 175
Jansen, Olivia	861, 377
Jantan, Ibrahim	906
Jantas, Danuta	540
Jantrawut, Pensak	1039
Jarmusch, Scott A.	365
Jarratt, Abigail	713
Jarret, Robert L.	161
Jaspars, Marcel	67
Jassbi, Amir Reza	344
Jaśpińska, Joanna	684
Jeevakumar, Trishiyaa Ragit	956
Jefford, Hannah	511
Jeffrey, Christopher S.	115
Jelić, Dubravko	331
Jena, Sudipta	507
Jennifer, Nambooze	812
Jennings, Laurence K.	359
Jenny, Urs	758
Jensen, Paul R.	185
Jeong, Eunah	822
Jeong, Se Yeon	927, 928
Jeong, Se Yun	270, 280, 281
Jeong, Woo-Hyun	475

Jeong, Woo-Jin	390	Kamatou, Guy	39
Jeon, Yukyoung	1040	Kampasakali, Elli	827
Jessen, Henning J.	381	Kampranis, Sotirios C.	60
Ji, Jianfeng	865	Kang, Bae-Gon	722, 948, 949
Jiménez, Marisol Cristina	391	Kang, Dong-Min	390
Jiménez, Vicente Montes	463	Kang, Eun Seo	926, 927, 928
Jing-Ru, Weng	186	Kang, Hee-Kyoung	697, 910
Jing, Shu-Xi	893	Kang, Jong Soon	156
Johari, Muhammad Sulaiman Mohd	373	Kang, Jung-Il	697, 910
Johnson, Holly	94	Kang, Jung-Wook	1013, 1014
Jones, Dean P.	488	Kang, Kyo Bin	247, 248, 249, 806, 822
Jongsma, Maarten A.	857	Kang, Tae Jin	615
Joo, Hyejin	565, 586	Kang, Tong Ho	1040
Jordão, Ana Carolina	296	Kapepula, Paulin Mutwale	440
Joshi, Darshana	502	Kapp, Karmen	813
Jui-Hsin, Su	186	Kaptein, Suzanne	420
Juliano, Haddad	380	Karachaliou, Efstathia	929, 1039
Jullian, Valérie	523	Karadağ, Ayşe Esra	609, 1008
Jullien, Kateline	861	Karady, Michal	711
Jung, Jeyong	222	Karagouni, Amalia D.	1018
Jung, Yong-Hwan	752, 988	Karahüseyin, Seçil	996
Junior, Raimundo Goncalves De Oliveira	397	Karaman-Jurukovska, Nevena	1022
Junior, Raimundo Gonçalves de Oliveira	451	Karamanoli, Katerina	958
Júnior, Valdemar Lacerda	200	Karampasi, Christina	868
Junka, Adam	394	Karanfilova, Ivana Cvetkovikj	780
Jun, Mingoo	204	Karapandzova, Marija	780
Jürgenliemk, Guido	568, 595	Karapandzova, Marja	630
Jürgens, Franziska M.	157	Karavergou, Sofia	299
Jurk, Kerstin	655	Karg, Cornelia	826
Justus, Nicolas de Aguiar	205	Karg, Cornelia A.	980
		Karimi, Hasan	965
		Karioti, Anastasia	875
		Karlsberger, Lea	482
		Karpierz, K.	708
		Karpouchtsi, Maria	31, 554
		Karreman, Hubert J.	42
		Karzis, Joanne	753
		Kasap, Melis	640
		Kashyap, Akanksha	537, 538
		Kaškelienė, Erika	524
		Katanaev, Vladimir L.	900
		Kato, Massuo Jorge	397
		Katsaros, Fotis	105
		Katsarou, Martha S.	994
		Kaur, Amninder	409
		Kaweesa, Elizabeth N.	535, 659
		Kaya, Gulen Irem	631
		Kayser, Leonie	595
		Kazemzadeh, Houman	108
		Keasling, Adam	223
		Keel, William	250
		Keh, Martin	1016
Kabamba, Nadege Ngombe	440		
Kaboudari, Ata	392		
Kačer, David	998		
Kachor, Anna	874		
Kahankova, Zdenka	259		
Kahn, Shahebraj	141		
Kaili, Stavroula	1018		
Kaiser, Marcel	183, 321, 455		
Kaixin, Gao	672		
Takeya, Hideaki	147		
Kakudidi, Esezah	719		
Kalamouni, Chaker El	389		
Kalema-Zikusoka, Gladys	47		
Kalemba, Danuta	644		
Kaliora, Andriana C.	142		
Kallendrusch, Sonja	922		
Kalli, Sylvia	398		
Kamarozaman, Aisyah Salihah	373, 374, 446, 447		

K

Kelber, Olaf	104, 449, 450, 863, 635, 646, 647, 652, 653, 657, 86	Kim, Jonghwan	962
Keller, Amy C.	582	Kim, Jongwhan	721
Keller, Lena	409	Kim, Kang Sub	204
Keller, Morris	593	Kim, Ki Hyun	266, 267, 268, 269, 270, 271, 272, 273, 280, 281
Kellner, Harald	454	Kim, Kyung Ah	273
Kennelly, Edward J.	215, 140	Kim, Kyu Won	961
Kenny, Owen	406, 749	Kim, Minji	962
Kerr, Tyler A.	181	Kim, Min Ju	723, 740
Keshri, Priyanka Kumari	661	Kim, Min-Yeong	1014
Ketterer, Paul	481	Kim, Seung Bum	400
Khalid, Sami	436	Kim, Seung-Hyung	973, 1035
Khalid, Sami A.	410	Kim, Soyeon	636
Khalid, Sami Ahmed	78	Kim, So Yeon	561
Khalighian, Najmeh	344	Kim, Yena	723, 961
Khalil, Atif Ali Khan	390	Kim, Yeongseo	249
Khan, Ajmal	388	Kim, Yong Bum	720
Khana, Shabana I.	161	Kim, You Ah	833
Khan, Hafizurrahman	519	Kim, Young-Soo	926
Khan, Humaira	1024	Kindl, Marija	332
Khan, Ikhlas A.	415, 685, 161	Kinghorn, A. Douglas	535, 659
Khayyal, Mohamed T.	660	Kinscherf, Ralf	652
Khederlarian, Lila	790	Kinscherf, Ralf	449
Khoo, Mary Gaik Hong	163	Kiraga, Łukasz	627
Khumalo, Gugulethu P.	439	Kirby, Christopher W.	317, 66
Khuniad, Chuanchom	170	Kirchhoffer, Olivier	629, 901
Kicel, Agnieszka	320	Kirchhoffer, Olivier A.	468, 131
Kiehne, Paul	1022	Kirchweiger, Benjamin	813, 72
Kielich, Natalia	504	Kirkegaard, Rasmus	184
Kieninger, Hildegard	385	Kis, Peter	437
Kientz, Gaëtan	702	Kiss, Anna	276, 527
Kiermaier, Josef	576	Kiss, Anna K.	540, 573
Kikionis, Stefanos	288, 621	Kiss, Anna Karolina	596, 1010
Kikuchi, Takashi	228	Kiss, Tivadar	986
Kilani, Dina	512	Kiss, Tobias	299
Kiledal, Anders	287	Kitagawa, Rodrigo Rezende	200
Kim, Brandon J.	291	Kirci, Damla	609
Kim, Chae Kyung	723	Klanert, Gerald	522
Kim, Chung Sub	235, 249, 177	Klein, Joshua D.	126
Kim, Dae Geun	220	Klein Junior, Luiz Carlos	734
Kim, Do-Hyeong	833	Klein, Thomas	864
Kim, Dong-Seon	561, 960, 973, 991, 1035	Klimantakis, Konstantinos	958
Kimel, Katarzyna	356	Klimek-Chodacka, Magdalena	920
Kim, Eun-Sook	312	Klimiené, Asta	524
Kim, Hye Jin	564	Kloucek, Pavel	259
Kim, Hyo-Min	1014	Klouček, Pavel	989
Kim, Hyungwoo	636	Knizkova, Katerina	987
Kim, Hyun-Ji	722	Knižková, Kateřina	974
Kimishima, Atsushi	71	Koch, Lukas	208
Kim, Jang Hoon	928	Koch, Mirijam	553
Kim, Jang-Hoon	565, 586	Koeberle, Andreas	36, 792, 826
Kim, Jeong Ah	686, 687	Koeberle, Solveigh	792

Koeberle, Solveigh C.	36	Koudoro, Yaya	48
Koech, Samson Cheruiyot	342	Koufali, Christina	330, 500
Ko, Eun Hye	729	Koutsaviti, Aikaterini	285, 60, 286, 292
Köhler, Thilo	254	Kovačević, Nada	395, 605, 606
Kohoutová, Lenka	998	Koval, Alexey	900
Kojić, Vesna	605	Kowalski, Rafał	463
Kokkini, Stella	261, 827	Koycheva, Ivanka K.	599, 602
Kokoska, Ladislav	340	Kozioł, Łukasz	361
Kokotkiewicz, Adam	353	Kozłowska, Weronika	394, 487, 532, 622
Kolarik, Miroslav	323	Kraithong, Witsarut	470
Kolarsky, Noah Gladen	520	Krajewska, Agnieszka	644
Koller, James T.	888	Kramer, Nynke	206
Köllner, Tobias G.	88	Krause, Rui Werner Maçedo	460
Kolodziejczyk-Czepas, Joanna	528	Krauze-Baranowska, Mirosława	354, 356, 1038
Kołodziejczyk, Agata	940	Krawczyk-Łebek, Agnieszka	364, 894
Kołodziejczyk-Czepas, Joanna	527	Kremenovica, Valentina	717
Kołtun-Jasion, Małgorzata	276, 527, 1010	Kremer, Dario	607, 698, 704
Komendo, Aleksandra	472	Kress, Márcia Regina von Zeska	296
Komoń, Marta Karolina	1010	Krettler, Christoph A.	396
Kompek, Albert	864	Kries, von Jens-Peter von	31
Končić, Marijana Zovko	705, 983	Krishnan, Sandhya	628
Kongkaew, Chuenjid	733	Krishnan, Sandhya R.	413
Konieczka, Paweł	41	Kristi, Leka	913
König, Beatrix	550	Kristofova, Radoslava	44
Konkol, Marcin	463	Kritsanida, Marina	391
Koo, Man Hyung	187	Krittanaï, Supaluk	471
Kopf, Adrian H.	237	Kroning, Kayla	560
Korábečný, Jan	151	Křoustková, Jana	151
Korablova, Olha	262	Król, Agata	353
Korczak, Maciej	489, 153	Krstic, Sanja	569
Korhonen, Petrus	787	Krstić, Gordana	339
Korinek, Michal	401, 425	Krüerke, Daniel	552
Kořínek, Michal	591	Krüger, David	680
Körmöczy, Tímea	986	Kruk, Aleksandra	309, 472, 489, 585
Koroleva, Arina	832	Krull, Nyssa K.	141
Korona-Główniak, Izabela	308	Krupova, Monika	341
Korostil, Isaiah I.	234	Krüsselmann, Philip	430, 432, 434
Korotkova, Marina	672	Krysa, Mikołaj	394
Korzeniowska, Weronika	487, 532	Krzykwa, Julie	863
Ko, Seok-Jae	565, 586	Kubacka, Monika	218
Kose, Yavuz Bulent	314	Kubica, Paweł	306, 308, 309, 338
Köse, Yavuz Bülent	289, 290	Kubiszewska, Aleksandra	492
Kosgahakumbura, Lakmini	274	Kuchar, Martin	987
Koshovyi, Oleh	484	Kuchař, Martin	974, 998
Koshy, Rojison	971	Kucuk, Melike	1026
Kostenko, Yuliia	430, 432, 434	Kuhn, Stefan	146
Kostner, Claudia	792	Kukula-Koch, Wirginia	571, 579, 634
Kostrzewa-Susłow, Edyta	364, 894	Kukuła-Koch, Wirginia	1002, 1028
Kosturko, Stefan	425	Kula-Maximenko, Monika	252
Kosturko, Štefan	591	Kulawik, Anna	869
Kotoku, Naoyuki	71	Kulevanova, Svetlana	780
Kouam, Simeon Fogue	256	Kulić, Žarko	121

Kulinowski, Łukasz	103, 308	Lameiras, Pedro	164
Kumar, Bhanu	661	Landegren, Ulf	433
Kumar, Jyotsna	435, 853	Landim, Euclides	959
Kumboonma, Pakit	278	Landrum, Leslie	411
Kumlehn, Jochen	478	Langat, Linda	678
Künstle, Gerald	556, 567, 574, 575, 617	Langat, Moses	511, 678
Kuo, Ping-Chung	516	Langbecker, Konrad	121
Kurach, Łukasz	628	Langeder, Julia	553
Kuruldak, Ebru	616	Lang, Rebecca M.	493
Kurylenko, Olena	874	Lankin, David C.	787
Ku, Sejin	724	Lanoue, Arnaud	964
Kushkevych, Ivan	458	Lanoue, Arnaud,	1007
Kusio-Targońska, Kamila	1028	Laphookhieo, Surat	278, 860
Kuszel, Urszula	472	Laranjeira, Inês Martins	633, 1005
Kvasnicová, Marie	292, 648	Lardicci, Claudio	241
Kwak, Jong Hwan	1040	Larlhimi, Abdelhalim	257
Kwon, Chaesun	313	Larsen, Thomas O.	386
Kwon, Hak Cheol	203	Laskaris, Paris	1018
Kwon, Jaeyoung	203	Lasnel, Lucas	699
Kwon, Kyeung-A	1014	Lasota, Magdalena	1002
Kwon, Mi Jeong	564	Latek, Urszula	754, 755, 756
Kwon, Oran	94	Latta, Olaf	381
Kwon, Yong-Jun	1014	Lau, Clara Bik-San	101, 198, 199, 936
Kyung-Sik, Song	948	Lau, David Tai-Wai	101

L

Labanowsky, Jérôme	416	Laura, Rieusset	968
Laborde, Quentin	414	Laurent, Patrick	899
Laborde, Quentine	295	Lauro, Gianluigi	623, 1004, 1012
LaBudde, Robert	941	Lawal, Bilqis Abiola	559
Lacandalo, Lina S.	539	Law, Kung Pui	906
Lackner, Michaela	76	Leach, David	221
Lacoue-Nègre, Marion	903	Leão, Pedro N.	182, 191, 501
Ladirat, Stephanie	482	Lebelo, Sogolo	588, 589
Laezza, Carmen	209	Leblanc, Anne	946
Laffleur, Flavia	84	Leblanc, Karine	258
Lagarias, Panagiotis	1021	Le Cabec, Audrey	692, 816
Lage, Miguel Angel	491	Lechwar, Paulina	1002
Lager, Estelle	584	Leclercq, Joëlle	953, 1025
Lahngong, Methodius	54	Ledoux, Allison	196, 440, 861, 54
Laiche, Mouna Hind	808	Lee,	427
Lai, Guan-Hua	773	Lee, Ae Sin	536
Lai, Kuei-Hung	29	Lee, Beom-Joon	565, 586
Lakušić, Branislava	594, 985	Lee, Boo-Yong	939
Lakušić, Dmitar	594	Lee, Brian L.	888
Lallemand, Marie-Christine	85	Lee, Chia-Lin	800
Lama, Adriano	1006	Lee, Ching-Kuo	179, 517, 793, 898, 899
Lamas, Marcelo Einicker	821	Lee, Choon M.	488
Lambrinidis, George	1021	Lee, Chun	516
Lam, Chu Shing	896	Lee, Da Eun	271
		Lee, Dahyeon	312
		Lee, Gyu Sung	235, 249, 962
		Lee, Ha Yeon	220
		Lee, Hung Kay	101

Lee, Hwa-Sun	156	Le, Ngoc-Thao-Hien	420
Lee, Hyo-Jong	962	Lengo, Mambu	232
Lee, Hyunyoung	722	Lenz, Myong-Su	382
Lee, In-Chul	1013, 1014	Leo, Marinella De	241
Lee, Jae-Cheol	271	Leonard, Carmen	626
Lee, Jin A.	740	Leonardo, Jemma Dawn	444
Lee, Jin Woo	222, 312, 313	Leone, Sheila	1034
Lee, Jong-Soo	426, 428, 429	Leonhard, Bettina	820
Lee, Juho	475	León-González, Antonio-José	760
Lee, Julia Kin-Ming	198	Le Pogam, Pierre	806
Lee, Jung-Ho	141	Le Ray, Anne-Marie	700
Lee, Ju Ryeong	962	Le Ray Richomme, Anne-Marie	908
Lee, Kyeong-Ryeol	475	Leroy, Charline	196
Lee, Kyung-Eun	1014	Le, Seo Yoon	280
Lee, Kyungha	247, 806	Le, Thi Thanh	687
Lee, Kyung Ho	724, 175	Leuchtenmüller, Niklas	790
Lee, Mina	247	Leung, George Pak-Heng	167
Lee, Min Ah	156	Leung, Leanne Lee	936
Lee, Myong-Min	403	Leupin, Marianne	717
Lee, Na Young	833	Le, Van-Tuyen	231
Lee, Peter A.	713	Levchuk, Iryna	262
Lee, Roun	725	Lev, Katherine L.	287, 837
Lee, Sang Kook	551	Lewis, Jared	560
Lee, Sang-Seob	962	Le-Xuan, Loc	36
Lee, Seogyeong	565, 586	Liang, Qiaoli	997
Lee, Seong Mi	729	Liang, Zheng-Ming	917
Lee, Seon-Kyeong	475	Liang, Zhuoning	948
Lee, Seo Yoon	268, 269, 270	Liao, Hsing-Yu	773
Lee, Seulah	187	Liasi, Georgia	829
Lee, Shih-Yu	831	Liatsopoulou, Aikaterini	1032
Lee, Shoei-Sheng	473	Liaw, Chia-Ching	307
Lee, Sullim	204, 273	Liberato, Walleska	893
Lee, Taeho	722, 948	Libero, Maria Loreta	1034
Lee, Wai-Fung	198	Li, Chunshun	193
Lee, Wookyu	721	Li, Danny	461
Lee, Yun Mi	561	Lien, Yi-Yang	773
Lee, Yun-Sil	174	Li, Hongwen	936
Lee, Zhi Xuan	163	Li, Jing	1017
Lefèvre, Gabriel	74	Lima, Clarissa Silva	641
Legay, Sylvain	522	Lima, Débora Machado de	334, 335
Legeay, Marc	795	Li, Min	371
Legssyer, Abdelkhaleq	748	Limsuwanchote, Supattra	542, 925
Lehner, Martin D.	533, 549, 550	Lim, Yijae	551
Lehtivarjo, Juuso	787	Lim, Young Woon	248
Leibold, Thomas	293	Lin, Brendan	418
Leignadier, Julie	924	Linda, Tetty Marta	350
Leka, Kristi	412	Linder, Thomas	834, 836
Lela, Ludovica	862	Ling, Zhengyi	476
Le Masle, Agnès	903	Linh, Phan	380
Lembo, Francesca	422	Lin, Hsiu-Mei	791
Lemieszek, Marta Kinga	472	Linnington, Roger G.	79, 888, 140
Lemoine, Sandrine	584	Lin, Jun-Wei	671

[illegible]

Maake, Caroline	211, 439
Maas, Elke	420
Mabou, F.D.	244
Machado Jr., Rogerio Jose	518
Macherla, Venkat	879
Ma, Chi Thanh	352
Macías, Francisco A.	114, 117, 322, 171
Maciejewska-Turska, Magdalena	399, 620
Mackenzie, Thomas A.	77
Macquigneau, Wadi	699
Mada, Olubunmi Olaitan	883
Madduri, Krishna	481
Madelou, Nikoleta Anna	739, 843, 844
Maeschli, Ariane	764
Magalhães, Mariana	809
Magalhães, Solange	111
Maggi, Filippo	152
Magiatis, Prokopios	284, 329, 330, 841, 843, 844, 845, 166, 911
Magiera, Anna	527, 528
Magnutzki, Alexander	100
Magot, Florent	1007
Magozwi, Kemboi	840
Maharjan, Sailesh	341
Mahmoud, Abdelhalim Babiker	321
Ma, Hongwu	87
Mah, Siau Hui	906, 909
Maier, Claudia S.	98

Maier, Jakob	820	Markowitsch, Sascha D.	846
Mainka, Marta	611	Marney, Luke C.	98
Majdan, Magdalena	1038	Maron, Etienne	105
Ma, Jin Yeul	403, 457, 458, 850	Maróstica, Mário Roberto	663
Makhloufi, Hind	189	Marquesa, Thaywane Azevedo	335
Makhubu, Fikile N.	753	Marques, Joana	959
Makinde, Emmanuel	107	Marques, Mário Pedro	809, 959
Makino, Toshiaki	93	Marquez, Lewis	47
Makitrynsky, Roman	409	Marshall, Erin M.	828
Makreski, Petre	630, 851	Marsik, Petr	259
Maksimović, Zoran	395, 985	Maršík, Petr	989
Malarz, Janusz	540	Martani, Enrico	935
Maldonado, Karen Cristina Comin	124	Martí, Meritxell	1031
Malee, Kulawadee	858	Martín-Cordero, Carmen	760
Malee, Kulwadee	866	Martinez, Agathe	164
Malfa, Giuseppe Antonio	1015	Martínez-Ferrer, Magaly	610
Malicka, Maria	487, 532	Martínez, Isabel Ramón	376
Malik, Muhammad Irfan	750, 123	Martínez-Máñez, Ramón	989
Ma, Linlin	107, 139	Martínez, María Dolores Marcos	989
Malinowska, Magdalena	1007	Martínez-Rodríguez, Oswaldo P.	246
Malinowska, Magdalena Anna	309	Martínez Yerena, J.A.	275
Mallema, Yassine	363	Martín, Jesús	77, 315
Mallem, Yassine	743	Martins, Carlos Henrique Gomes	335
Malmir, Maryam	327, 328, 358	Martins, Giovanna Gomes	821
Malones, Angel Mae	444	Martins, Teresa P.	501
Malý, Matěj	995	Martins, Verônica Giuliani de Queiroz Aquino	965, 966
Mambu, Lengo	189	Martzen, Mark R.	546
Mamede, Lucia	54	Ma, Rui	141
Mandau, Anne	924	Marylène, Viana	232
Mandova, Tsvetelina	75	Marzorati, Stefania	746, 767
Mandrone, Manuela	136	Mascellani-Bergo, Anna	989
Maneerat, Tharakorn	278, 860, 866	Masciulli, Fabrizio	1034
Manfredi, Kirk P.	245	Mäser, Pascal	183, 455
Mangieri, Claudia	862	Masi, Marco	210
Manniello, Matthew	546	Massironi, Alessio	746, 767
Mannina, Luisa	1034	Masson, Jérôme	921
Mante, Priscilla Kolibea	417	Masugarut, Pisitchai	471
Mantzouridou, Fani Th.	731	Masuku, Nelisiwe	588, 589
Manzanares, Paloma	490	Masullo, Milena	623, 993, 999
Maphanga, Veronica	39	Mata, Rachel	165
Maquoi, Erik	861	Matasyoh, Josphat	454
Marceau, Levasseur	801	Matejka, Jaroslav	259
Marčetić, Mirjana	594, 605, 606	Matějka, Jaroslav	989
Marchal, Axel	1009	Mateo, Marta	105
Marchal, Luc	38	Mateos, Katherine	713
Marc, Litaudon	380	Matevska-Geskovska, Nadica	780
Marcotullio, Maria Carla	176	Mathiron, David	702
Marcourt, Laurence	468, 900, 901, 131	Matías, Mariana Sanchez	391
Mareš, J.	275	Matkowski, Adam	394, 487, 532, 622, 684
Mareš, Jan	190	Matos, Ana C.	33
Marijan, Marijan	705	Matsuda, Kenichi	476
Markovics, Alex	126		

Matsuzaki, Michihisa	71	Menghini, Luigi	1034
Mattarelli, Paola	769	Menicagli, Virginia	241
Mattar, Nariman	126	Mennet von Eiff, Mónica	574
Matuszewski, Maciej	484	Mennet-von Eiff, Mónica	556, 575
Mauduit, Morgane	365, 366	Menon, Arya	775
Mayer, Philipp	764	Menoud, Valérie	764
May, Tom	76	Merckx, Anaïs	451
Mazahery, Ahmad Reza	886	Mert, Ozan	318
Mazen, Ahmed M.A.	311	Mertzaniadis, Dimitrios	261, 827
Mazucato, Vitor de Souza	62	Mervić, Mateja	332
Mazur, Oliwia	504	Meslet-Cladière, Laurence	231
Mbaka, G.O.	859	Mesmar, Joelle E.	662
Mbanu, Augustine	669	Messaili, Souhila	1037
McAlpine, James B.	893	Metsä-Ketelä, Mikko	495, 832, 89
McCadden, Caitlin	469	Meunier, Manon	55
McCarthy, Ronan	404	Meyer, Nadin	664
McCarty, Bianca	997	Meyer, Susan	210
McEachern, Emily	291	Mgbeahuruike, Eunice Ego	355, 464, 466
McFadden, Jeremy	481	Michailidis, Dimitris	230, 1001
McGaw, Lyndy J.	753	Michaliski, Lamonielli F.	486
McKay, Scott	888	Michel, Frédéric	377, 913
McKenna, Lorraine	755	Michel, Martin	550
McLeman, Amy	243	Michels, Franck	702
McLennan, Matthew	47	Michel, Thomas	700
McPhail, Kerry L.	68	Michrafy, Ines	336
Medina-Díaz, Irma Martha	613	Migliorini, Ivan Lucas de Freitas	283
Medronho, Bruno	111	Miguel, Sónia P.	632, 852
Meents, Tomke	87	Mihaylova, Liliya V.	600, 601, 603, 173
Meesomperm, Peerapong	733	Mihaylova, Nikolina M.	599
Mehiou, Afaf	748	Mihovilovic, Marko D.	834, 836
Mehjabin, Jakia Jerin	476	Mikropoulou, Eleni V.	142
Mehl, Florence	901	Mikros, Emmanuel	847
Mei, Liyan	619	Mikucki, Jill A.	713
Mejri, Yassine M.	806, 818	Milella, Luigi	862
Melcher, Silas F.	533	Milenković, Marina	985
Melini, Stefania	1006	Miles, Christopher O.	386
Meli, Rosaria	1006	Milic, Nikola	330
Mellick, George D.	107	Miller, Kate	742
Melliou, Eleni	284, 285, 329, 841, 843, 844, 845, 166, 911	Milosevic, Marija	100
Mellou, Foteini	1032	Min, Byung Sun	564, 686, 687, 720, 850
Melnyk, Natalia	608	Minceva, Mirjana	145
Melnyk, Sofia	480, 874	Minorics, Renáta	817
Melo, Matheus Reis Santos de	334	Min, Seung Eui	720
Melo, Michelle Nonato de Oliveira	821	Mioletti, Silvia	750
Melzig, Matthias F.	382	Mirabile, Salvatore	835
Mendel, Marta	754, 755, 756, 765	Miron, Anca	655, 978
Mendonça, Iatã	349	Misra, Ankita	519, 661, 677
Meneses, Carlos Henrique Salvino Gadelha	966	Miszczak, R.	708
Meneses, Vanya	165	Miszczak, Rebecca	707
Menett-von Eiff, Mónica	617	Miszczuk, Edyta	627
Menezes, Gabriela de Lima	75	Misztal, Tomasz	650
		Mitaine-Offer, Anne Claire	929, 1039

N

O'Connor, Roberta	453	Oostingh, Gertie Janneke	597
O'Donoghue, Anthony J.	781	Opallo, Nicola	1006
Odou, Solène	207, 160	Opletal, Lubomír	239
Odumosu, Patricia	559	Opoku, Melvin Osei	469
Ogbole, Omonike O.	882	Orčić, Dejan	772
Ogechukwu, Omeje Edwin	651	Orčić, Dejan Z.	576
Oger, Elise	908	Orhan, Ilkay Erdogan	132
Ogoamaka, Osadebe Patience	651	Orhanović, Stjepan	698
Ogunniyi, Queeneth Abiola	882, 883	Orjala, Jimmy	260
O'Hara, Peter	406, 749	Orlandini, Serena	871
Oh, Dong-Chan	59	Orlando, Giustino	1034
Oh, Hyun-Ji	939	Orozco, Jahir	797
Oh, Kyung-Eon	833	Oroz, Tamara	105
Oh, Min Hyuk	740	Ortega, Teresa	761
Oh, Seh Wook	729	Ortiz, Adriana Aburto	529
Oh, Won Keun	508	Ortiz, Sergio	899
Oiry, Catherine	596	Osano, Anne	796
Ojewumi, Anthony Wale	883	Osiagwu, D.	859
Okafor, Fabian	372	Osmaniye, Derya	314
Okamura, Yu	88	Osores, Esteban	710
Okino, Tatsufumi	279, 476	Ospina, Victoria	545
Oklestkova, Jana	878	Ostash, Bohdan	480
Okoro, Ijeoma Solomon	460	Ostash, Iryna	874
Oksman-Caldentey, Kirsi-Marja	130	Ősz, Bianca Eugenia	421
Oktavia, Listiana	877	Otogo, Elvis N'Nang	824
Okten, Suzan	337	Otsuki, Kouharu	228
Oladele, Johnson	762	Otto, Daniel Jordan	365
Oladipo, Adewale	588, 589	Ottolina, Gianluca	197, 225
Olajide, Olumayokun A.	557	Ouachinou, Jérôme	48
Olatunji, J. O.	498	Oueslati, Saoussen	376
Olech, Marta	643, 644, 645, 1023	Ouoba, Kampadilemba	737
Olędzka, Agata J.	484	Owczarek-Januszkiewicz, Aleksandra	320
Oliveira, Adriana Passos	821, 846	Owens, Skyler L.	493
Oliveira, Alberto de	335	Oyaseiye, Precious Ezougha	368
Oliveira, Ana Ligia Leandrini de	283	Oyejide, J.O.	859
Oliveira Junior, Raimundo Gonçalves de	85	Oyelola, Roseline	762
Oliveira, Paula	491	Oyeyode, John Oluwafemi	883
Oliveira, Paulo J.	959	Ozbek, Emine Nur	319
Olivier, Bonnet	736	Özenver, Nadire	290
Olivon, Florent	675	Oziamara, Okoro Nkwachukwu	651
Olounlade, Pascal Abiodoun	691	Ozkurt, Muhammet Cagri	624
Olounladé, Pascal Abiodoun	899	Özsoy, Nurten	996
Olšanský, Milan	125	Öztürk, Meltem	818
Olszewska, Monika A.	528	Ozturk, Sura Baykan	319
Olszewska, Monika Anna	527		
Olubi, O.O.	859		
Oluški, Mirko	1019		
Olusola, Olajuyigbe Olufunmiso	438		
Omarsdottir, Sesselja	141		
Omosa, Leonidah Kerubo	183		
O'Neill, Sarah	811		
Onurdag, Fatma Kaynak	337		

Ó

Ólafsdóttir, Elín S. 282

P

Pabich, Martyna	528	Park, Yun Seo	222
Pablo, María José Cubero	43	Parlar, Zeliha	624
Pabst, Jean-Yves	737	Parzonko, Andrzej	573
Paciencia, Mateus Luís Barradas	124	Pastore, Arianna	37, 625
Pacifico, Severina	112, 1027, 169	Paśko, Paweł	940, 945
Paczkowska-Walendowska, Magdalena	238	Patel, Harun	375
Paek, Ockjin	721	Patiwettham, Saharat	471
Paetz, Christian	87, 88	Patron, Nicola	784
Pagni, Marco	901	Pattarachotananant, Nattaporn	167
Paguet, Anne-Sophie	74	Patyra, Andrzej	596
Paini, Jacopo	225	Paula-Souza, Juliana de	1041
Paiva, Wesley de Souza	965	Paulc, Valerie J.	346
Pajot, Emmanuel	258	Pauli, Guido F.	893, 52, 146
Pajumo, Maria	1030	Paul, Valerie J.	109
Pakora, Gilles	397	Pavanello, Larissa	518
Pakulski, Zbigniew	648	Pavličević, Aleksandra	772
Palanisamy, Uma Devi	870	Pawelski, Damian	915
Palaric, Cécile	74	Pawlak, Bernard	463
Paliunytė, Jurgita	637	Pawlus, Alison D.	1022
Palm, Marlon	449, 652	Pawłowska, Karolina A.	432
Palomino, Olga	761	Paździora, Wojciech	945
Palo-Nieto, Carlos	424	Pecio, Łukasz	532
Paloukopoulou, Charikleia	875	Pecio, Solomiya	532
Panagiotidou, Christina	714	Pecková, Radka	125
Panagou, Anastasia	621	Pedrosa, Mariana C.	490
Panda, Bibhu Prasad	1036	Peintner, Ursula	76
Panda, Pratap Chandra	507	Pel, Pisey	566
Panda, Sujogya Kumar	388	Pelzer, Florian	592
Panieri, Giuliana	828	Peña-Valdivia, Cecilia Beatriz	563
Pan, Li	535, 659	Pěňčík, Aleš	172
Papachristodoulou, Anastasia	847	Peng, Bo-Rong	29
Papadopoulou, Kyriaki	286	Peng, Cheng-Huan	804
Papageorgiou, Spyros	1011	Peng, Jiangnan	796
Papaioannou, Varvara	845	Penouilh, Marie-José	929
Papakonstantinou, Aikaterini	841, 166	Pepe, Jessica L.	234
Papantonaki, Anastasia-Ioanna	868	Pereira, Catarina G.	975
Papassotiropoulos, Andreas	99	Pereira, Lucas Natã	734
Parchman, Thomas L.	115	Pérez-Bonilla, Mercedes	315
Parisi, Valentina	423, 835, 135	Pérez-Juanchi, Diana	777
Park, Byoungjun	833	Pérez-Moreno, Guiomar	77
Park, Geonha	723, 724, 725, 961	Pérez-Thierry	217
Park, Ji Hyeon	926, 927, 928	Pérez, Valeria Reyes	165
Park, Ji-Young	729	Pérez-Victoria, Ignacio	77
Park, Ju-Hyun	1014	Perhal, Alexander F.	813, 834, 836
Park, Jun Woo	615	Perino, Sandrine	934
Park, Min-Ha	187	Perković, Milana	849
Park, Sangjun	636	Pernodet, Nadine	1016
Parks, Eden L.	912	Perret, Jérémy	921
Park, So-Young	722, 948	Perry, Matthew	742
Park, Thomas J.	876	Perse, Xavier	160
		Pertuit, David	1039
		Pertz, Olivier	31, 554, 555

Pervaiz, Javeria	670	Piqué-Borràs, Maria-Riera	556, 567, 574, 575, 587, 617
Perz, Martyna	364	Pires, Loren Monielly	786
Petek, Anja	443	Pirhadi, Somayeh	344
Peter, Samuel	717	Pirozzi, Claudio	1006
Peters, Harrison	888	Pitakbut, Thanet	825
Peters, Justin	245	Pithon, Shamsia	226
Petit, Bastien	336	Piwowarski, Jakub P.	430, 432, 434, 435, 479, 489, 492, 153
Petit, Thomas	226	Piwowarski, Jakub Patryk	608
Petraitytė, Evelina	637	Placha, Iveta	44, 770
Petránková, Barbora	82	Platon, Estelle	700
Petras, Daniel	872	Pliakou, Christiana	829
Petrásková, Lucie	82	Pluskal, Tomáš	849
Petrelli, Riccardo	152	Pobłocka-Olech, Loretta	356
Petrović, Jelena	772	Podolak, Irma	214, 218, 945
Petrov, Kamen	790	Poetzsch, Michael	856
Petzer, Inge-Marie	753	Pogam, Pierre Le	824, 65
Petz, Petra	345	Poigny, Stéphane	992
Pezelj, Ivana	698	Polak, Joanna	754, 755, 756
Pfeiffer, Maximilian	923	Polcaro, Luciana Maria	993
Pferschy-Wenzig, Eva-Maria	311, 312, 86, 371	Poli, Ferruccio	136
Pham, Huong T.	248, 822	Polyzois, Alexandros	839
Phan, Chin-Soon	476	Pommerening, Tanja	450
Phelippe, Myriam	207	Ponce-Zea, Jorge-Eduardo	508
Phi, Kim-Hoa	187	Ponticelli, Maria	862
Philbin, Casey S.	115	Pont, Thibaut Robiou du	231
Phillips, Tabitha J.	190	Popowski, Dominik	479, 489, 492
Phukhatmuen, Piyaporn	860	Pos, Klaas Martinus	383
Phuneerub, Pravaree	866	Potterat, Olivier	534, 593
Piacente, Sonia	623, 993, 999	Poulain, Clément	257
Piano, Filomena Del	1006	Poupon, Erwan	806
Piaz, Fabrizio Dal	135	Poynton, Ella F.	888
Piccolella, Simona	112, 1027, 169	Pradel, Charlotte	160
Picot, Laurent	85	Prasad, Chandra Prakash	537, 538
Picquet, Michel	929	Prasansuklab, Anchalee	167
Pidaný, Filip	151	Preet, Gagan	67
Pieczara, Magdalena	920	Premprasert, Charoenwong	542
Piel, Jörn	28, 854	Preto, António José	396
Pierce, Savannah	264	Prieto-Davó, Alejandra	246
Piergentili, Alessia	152	Prochownik, Ewelina	940
Pierre, Francotte	736, 913	Promnart, Phunrawie	278
Pieters, Luc	47, 393, 420, 479	Promyos, Pimonnart	1039
Pietruszka, Jörg	689	Protopapa, Evangelia	1032
Piffet, Caroline	105	Pryce, Christopher R.	549
Pigott, Maria	570	Przyborowski, Jerzy Andrzej	320
Ping-Jyun, Sung	186	Ptach, Wiesław	754
Pinilla-Rosas, José M.	750	Ptaszyńska, Aneta A.	745
Pin, Matthew	888	Puglisi-Weening, Melany	264
Pinon, Aline	189	Pupo, Mônica Tallarico	283
Pintado, Manuela	490	Purdy, Trevor N.	79
Pinto-Ribeiro, F.	1005	Purger, Dragica	345
Pinto-Ribeiro, Filipa	633		
Piovan, Anna	690		

Püski, Péter	986
Putalun, Waraporn	470, 471
Pylkkö, Tuomas	301

Q

Qader, Mallique	141
Qamar, Sarmad Ahmad	112
Qiu, Feng	1017
Qi, Yunci	193
Quattrini, Andrea	325
Quave, Cassandra	47
Quave, Cassandra L.	418, 419, 488
Queiroz, Emerson F.	468
Queiroz, Emerson Ferreira	900, 131
Queiroz, Nubia C.A.	510
Quetin-Leclercq, Joëlle	899, 54
Quiceno, Eyson	545
Quijano, Leovigildo	777
Quinchia, Jennifer	807
Quinn, Ronald J.	139
Quintas, Célia	975
Quiros-Guerrero, Luis	468, 131
Quiros-Guerrero, Luis-Manuel	900, 901

R

Rabelo, Luan da S. D.	510
Rabenau, Holger F.	72
Rabini, Sabine	635
Rabone, Muriel	67
Racziewicz, Imelda	74
Raddi, Rafael	1041
Radetzki, Silke	31, 554
Radwańska, Aleksandra	1038
Raeber, Justine	855, 856
Raganathan, Dinesh	367
Raharivelomanana, Phila	682, 934, 938, 164
Rahman, Zuha	1036
Raimondo, Francesco Maria	835
Rajasekharan	919
Rakotondraibe, Harianantenaina L.	98
Rallis, Michael	621
Rallis, Michail Ch.	868
Ramada, Marcelo Henrique Soller	124
Raman, Babu	481
Rämä, Teppo	341
Rambaud, Caroline	74
Ramírez-Apan, Teresa	777
Ramos, Alba Maria	750

Ramos, Maria C.	77
Ramtanon, Ian	903
Ramtoola, Zeibun	811
Rangsinth, Panthakarn	167
Ranouille, E.	938
Rarová, Lucie	292
Rárová, Lucie	648, 878
Rasgado-Bonilla, Fátima Azucena	544
Raskopf, Esther	646, 647
Raso, Giuseppina Mattace	1006
Rasol, Nurulfazlina Edayah	446
Rasteiro, M. Graça	111
Ratajac, Radomir	772
Ratão, Isabel	975
Rath, Santosh Kumar	310
Ratnayake, Ranjala	346
Räuber, Gesine Karas	770
Raucci, Carlo	303
Raviraj, Thiyagarajan	425
Rawat, Poonam	519, 661
Ray, Asit	507
Ray, G. Joseph	787
Raymond, Louis	702
Ray Richomme, Anne-Marie Le	258
Rayya, Noura	146
Reazo, Renzo Luis D.	539
Rebbani, Essadia	871
Rebets, Yuriy	480, 874
Rédei, Dóra	339
Redl, Martina	30, 153
Reeves, Desiree T.	582
Rego, Adriana	501
Reguengo, Livia	663
Rehan, Mohammad Zaker	750, 771
Reher, Raphael	316, 450, 872
Rehman, Najeebb	815
Rehrl, Johanna	681
Reif, Marcus	592
Reimer, Daniela	79
Reinhard, Steffen	224
Reinhardt, Jakob K.	555
Reis, Mariana	294, 168
Reis, Mariana Alves	990
Remy, Simon	903, , 164
Renault, Jean-Hugues	38, 903, , 164
Resende, P. de	357
Restrepo, Adriana	797
Reusch, Jane E.-B.	582
Reuter, Margaret M.	837
Reyes, Fernando	315, 191, 77, 129
Rial, Carlos	114, 171
Ribeiro, Arthur Barcelos	785

Sahri, Nihad	662	Sasagawa, Masa	546
Saidene, Naima	512	Satyral, Prabodh	436, 437
Saidu, Muhammad Bello	339	Saubion, Frédéric	55
Saivish, Marielena Vogel	75	Saukel, Johannes	509
Sajed, Tanvir	888	Saunier, Clémentine	782
Salgaço, Mateus	663	Sauvager, A.	244
Saliba, Ana Sofia	663	Saviano, Gabriella	1004, 1012
Saliba, Jennifer	189	Šavikin, Katarina	1019
Salih, Enass	464, 466	Savova, Martina S.	590, 599, 601, 603, 173
Salih, Enass Y.A.	355	Sawicki, Rafał	408
Saliou, Claude	1016	Sayed, Hesham MB	87
Saliu, Adenike Mariam	883	Scalabrino, Gaia	404, 570
Salmon, Melissa	784	Scandar, Samir	176
Salomon, Christine	453	Scanff, Marie Le	1009
Salomon, Christine E.	70	Scarafile, Donatella	769
Saltan, Nagehan	314	Schäfer, Lizanne	455
Salvador, Manuel	128	Schärer, Hans-Jakob	678
Salvatore, Maria Michela	210, 405	Schempp, Florence M.	79
Salwinski, Aleksander	1030	Schenk, Alexander	978
Samaillie, Jennifer	702, 74	Scheuber, Victoria	764
Samardžić, Stevan	605, 985	Schiller, Laura	922
Sambayi, Godfrey	393	Schinkovitz, Andreas	55
Samborskyy, Markiyan	874	Schleker, Sylvia	689
Samdal, Ingunn A.	386	Schlemmer, Markus	552
Šamec, Dunja	305, 326	Schlittenlacher, Theresa	765
Samimi, Aria	857	Schmalenberg, Kristine	1016
Sammons, Kate	1022	Schmelzer, Sebastian	587
Sampietro, Diego	391	Schmidli, Anina	856
Samples, Robert	712, 713	Schmidtke, Michaela	553
Samuel, Monday Dadi	372	Schmidt, Thomas J.	157, 183, 455, 456
Samukha, Vadym	1004, 1012	Schmiech, Michael	923
Sánchez-Hidalgo, Marina	315	Schmietendorf, Hannes	553
Sánchez-Vázquez, María	610	Schnabel, Thomas	597
Sandargo, Birthe	323	Schnee, Sylvain	254
Sandasi, Maxleene	39, 805	Schneider, Yannik K.	301
Sandy, Desrat	380	Schnürch, Michael	834, 836
Šangut, Iva Jurčević	305, 326	Schoenhardt, Katherine	879
Sangwan, Payare L.	310	Scholl, Annika J.	646, 647
Sano, Aiko	933	Schoss, Katja	676
Santiago-Rosario, Mitzy	610	Schrey, Hedda	454
Santopietro, Federica	862	Schroeder, Frank C.	839
Santos, Ana Carolina Costa	334	Schüler, Vincent	224
Santos, Anna Livia Oliveira	335	Schultz, Fabien	47, 121
Santos, Gustavo	297	Schultz, Pamela J.	775
Santos, Leticia Barbosa	965	Schultz, Trevor	70
Santos, Mario Ferreira Conceição	786	Schulze, Christian	385
Santos-Martinez, María Jose	578	Schulzke, Esther	323
Saraiva, Lucília	33	Schuster, Anja	597, 681
Sardone, Nicola	951	Schuster, Sabine	776
Sari, Aynur	616, 996	Schwaiger, Stefan	36, 84, 117, 114
Sarker, Satyajit D.	170	Schwark, Markus	190
Sarotti, Ariel M.	193	Schwarzinger, Jacqueline	525

Schwarz, Patrik F.	813, 834	Sherman, David	461
Schwikkard, Sianne	150	Sherman, David H.	287, 288, 837, 58
Schwindl, Sebastian	568	Shetty, Raviraja Gavali	919
Scobie, Darren	67	Shetye, Gauri	141
Scognamiglio, Monica	300	Sheu, Jyh-Horng	505
Scortecci, Katia Castanho	641, 965	Shiels, Katie	897
Scortecci, Kátia Castanho	966	Shimada, Yusuke	71
Scotti, Francesca	211	Shim, Sang Hee	222, 313
Scozzoli, Maurizio	769	Shin, Hee Jae	156
Sebree, Joshua	245	Shin, Jae Young	926, 927, 928
Secundo, Francesco	197, 212, 225	Shin, Min-kyoo	313
Seeberger, Peter H.	50	Shin, Mi Rae	740
Seegers, Stephan	410	Shin, Soyoung	721
Seglab, Fatiha	347	Shinyuy, Lahngong Methodius	377
Segone, Ramoagi	805	Shrestha, Krishna K.	1029
Seidel, Veronique	741, 970	Shuaib, Nurah Eniola	348
Seligmann, Benedikt	478	Shum, David	315
Selo, Mohammed A.	298, 404	Shyur, Lie-Fen	773
Semdé, Rasmané	737	Siah, Ali	74
Sendker, Jandirk	404, 976	Sicolo, Francesco	951
Senevirathne, Amal	429	Sideratou, Zili	105
Senn, Roman	978	Sidiropoulou, Kyriaki	166, 911
Sens-Albert, Carla	550	Sieniawska, Elwira	399, 408, 620, 902
Seo, Chang Seob	930	Siewert, Bianka	76
Seo, Chang-Seob	1033	Sigrist, Hannes	549
Seo, Jaeuk	721	Sikora, Elżbieta	1007
Serino, Elena	37, 623, 625	Sillapachaiyaporn, Chanin	757, 1020
Séron, Karin	74	Silva Almeida, Jackson Roberto Guedes da	85
Serra, Ana Teresa	491	Silva, Ana	959
Serra, Arianna	76	Silva, Ariana Pereira da	965, 966
Serrano-Jara, Daniel	43	Silva, David Da	964
Serrano, Rita	358, 656	Silva, Dulce	349
Serra, Stefano	83	Silva, Givaldo S. da	349
Seršen, Sara	445	Silva, Jefferson de Souza	124
Sesink, David	681	Silva, Karolinne Beloti	786
Setiawan, Andi	71	Silva, Marlene	1005
Setzer, William N.	410	Silva, Natália Gonçalves	979
Sevin, Gulnur	318, 319	Silva, Olga	327, 328, 358, 656
Seyfarth, Lena	291	Silva, Rita de Cássia Melo Vilhena de Andrade	
Shaaban, Khaled A.	89	Fonseca da	734
Shadmani, Leila	68	Silva, William Cezar de Lima	904
Shaffer, Jacob M.C.	713	Silver, Jack	810
Shah-Hosseini, Kija	646, 647	Sim, Ha-Yeon	220
Shaito, Abdullah A.	347, 662	Simin, Nataša	772
Shanaida, Mariia	262	Simmet, Thomas	923
Sharafan, Marta	309, 1007	Simmler, Charlotte	217
Sharma, Hitender	774	Simões-Wuest, Ana Paula	534
Sharma, Nisha	146	Simoneau, Philippe	908
Shaw, Pang-Chiu	101	Simone, Mancini	392
Shcherbakova, Anastasiia	635	Simone, Simonetta Cristina Di	1034
Shen, Jincheng	113	Simonet, Ana M.	322
Sheridan, Helen	298, 404, 570	Simpson, Matthew	250

Simpson, Pelle	879	Sönmez, Sefa	944
Sinaeve, Sébastien	229	Son, Ninh The	194
Singer, Grant	488	Sonntag, Alexander	382
Singh, Dharendra	519	Son, So-Ri	203, 204
Singh, Garima	282	Son, Suk-Min	1013
Singh, Shekhar	661	Son, Yoorae	636
Singh, Shoor Vir	759	Sorbo, Luca Del	405
Singh, Surya Pratap	661	Sorkin, Barbara C.	977
Singkhram, Sukritta	470	Sornkaew, Nilubon	1020
Sipala, Federica	1015	Šoštarić, Ivan	395
Sipari, Nina	464, 466	Soto-Hernández, Marcos	544, 563
Sirak, Aleksandra	484	Soto, Javier A.	545
Sisto, Francesca	395	Soumah, Aboubacar	379
Sita, Dan Dragoş	421	Soumayanath, Amala	98
Sivieri, Katia	663	Soumyanath, Amala	520
Siwe-Noundou, Xavier	840	Souopgui, Jacob	54
SiweNoundou, Xavier	460	Souquet, Florence	397
Skala, Tomas	259	Sousa, Raquel Maria Ferreira	334, 335
Skalicka-Wozniak, Krystyna	628	Souza, Priscila de	734
Skalicka-Woźniak, Krystyna	39, 102, 103, 308	Spaggiari, Chiara	942, 967, 969
Skalický, Vladimír	172	Sparzak-Stefanowska, Barbara	354, 1038
Skalka, Dominik	976	Spatafora, Joseph W.	68
Skalli, Souad	803	Šperanda, Marcela	766
Skaltsa, Helen	823, 868	Spiegel, P. Clint	70
Skaltsounis, Alexios Leandros	571	Spiegler, Verena	694, 976
Skaltsounis, Alexios-Leandros	38, 1001	Spinozzi, Eleonora	152
Skaltsounis, Leandros	992	Spórna-Kucab, Aneta	366
Skaltsounis, Leandros A.	845, 149	Spriano, Filippo	359
Sklute, Eli C.	713	Spyropoulos, Georgios	292
Skowrońska, Weronika	608, 611	Spyros, Tziakas	333
Slavkovska, Violeta	606	Squarisi, Iara Silva	785
Slazak, B.	708	Srivastava, Sharad	519, 661, 677
Slazak, Blazej	61	Srivastava, Varsha	759
Slippers, Bernard	323	Srivastava, Vikas	519
Słomka, A.	708	Sroka-Bartnicka, Anna	394
Słomka, Aneta	707	Stadler, Marc	323, 454, 881
Smakosz, Aleksander	532	Stadler, Nina	689
Šmejkal, Karel	467	Stajich, Jason E.	68
Smyth, Thomas J.	406, 749	Stamataki, Christina	330
Snodgrass, Harrison	560	Stambouli, Belkes N.	215
Sobeh, Mansour	662	Standley, Ashlyn	223
Soeberdt, Michael	100	Stanković, Jelena S. Katanić	569
Soldati, Thierry	468, 901	Stanojković, Tatjana	985
Solhaug, Anita	386	Šťastný, Jan	259, 989
Solís, Isabel Martínez	761	Statha, Dimitra	868
Solis, Pablo	99, 555	Stavropoulou, Maria-Ioanna	873
Somer, Nehir Unver	640	Ştefănescu, Ruxandra	421, 981
Somoza, V.	567	Štefan, Maja Bival	331, 332
Song, Chang-Hwa	960, 973, 1035	Stefano, Gandolfi	225
Song, Gunju	939	Stefanowicz-Hajduk, Justyna	639, 649
Song, Kyung-Sik	722, 949	Stefkov, Gjoshе	630, 780, 851
Song, Ying-Chyi	665	Steinbeck, Christoph	146

Tanaka, Takashi	541	Thomas, Michel	885, 968
Tang, Shen	343	Thomas, Olivier P.	359
Tang, Yi	181, 838	Thomassin, Jean-Michel	105
Tan, Jin Yi	141	Thorel, Audrey	583
Tanniou, Simon	263	Thuerig, Barbara	678
Tan, Sheri-Ann	163	Thuillier, Isabelle	1037
Tansley, Connor	784	Tiefenbacher, Stefanie	371
Taoubi, Khalil	583	Tietjen, Ian	379
Tarabasz, Dominik	571, 634	Tirkkonen, Heli	832, 89
Tarfa, Florence	558, 559	Titton, Doriane	682
Taştan, Pelin	462	Todorova, Monika N.	590, 598, 601, 603, 173
Tate, Rothwell	367	Todorova, Velislava	590
Tauchen, Jan	340, 989	Tolossa, Ketema	506
Tavares, Denise Crispim	334, 785	Tomasi, S.	244
Tayeb, Bizhar A.	817	Tomasi, Sophie	192, 240
Tchetan, Esaïe	691, 899	Tomczyk, Michał	472, 642, 650
Tchinda, Alembert	440	Tomczykowa, Monika	915
Tchinda, Alembert Tiabou	196	Tomita, Ryoko	541
Tchoumtchoua, Job	105	Tommasi, Nunziatina De	135
Tedesco, Doriana Eurosia Angela	746, 767	Tomou, Ekaterina-Michaela	823
Teerapongpisan, Passakorn	278, 860	Tong, Man-Ho	101
Tehan, Richard M.	68	Tong, Xiaolin	371
Teh, Soek Sin	906	Tormo, Jose R.	668
Teixeira, Thaiz Rodrigues	296, 297	Totoson, Perle	604, 655
Tekieli, Anna	366	Touboul, David	110, 378
Tekielska, Dorota	465	Tragust, Simon	208
Tel, Banu Cahide	624	Trajkovska, Ana	780
Telukunta, Kiran K.	502	Tranchimand, Sylvain	240
Tembeni, B.	498	Trancoso, Daniel Stoffle	904
Tembu, Jacqueline	840	Tran, Khanh	941
Temiz, Burak	290	Tran-Thi, Van Anh	36
Tencomnao, Tewin	757, , 167	Tran, Trong D.	931
Tenderly, Vidya Febrasca	425	Trautman, Ian A.	68
Teng, Yu-Ning	515	Treml, Jakub	467, 989
Teodoro, Amanda Leal da Silva	709	Tremmel, Sabine	533
Teresa, Ma. Theresa B.	982	Trempat, Pascal	584
Termentzi, Aikaterini	1019	Trentin, Riccardo	699
Tero-Vescan, Amelia	421, 981	Tretter, Zachary M.	888
Teruna, Hilwan Yuda	351	Trifan, Adriana	102
Tesárek, Adam	467	Tripathi, Ashootosh	775
Testai, Lara	241	Tröger, Wilfried	592
Teufel, Robin	31, 99, 100, 381, 555	Trougakos, Ioannis P.	1018
Tews, Friederike	574	Tsafantakis, Nikolaos	330, 500, 1018
Thaisaeng, Wachirasak	880	Tsai, Wei-Cheng	831
Thamnarak, Wanlaya	880	Tsalikis, Lazaros	829
Thankappan, Babumon M.	120	Tsara, Anastasia Kyriaki	1011
Thasana, Nopporn	880	Tsitlakidou, Petroula	958
Theodoridis, Konstantinos	829	Tsolakou, Annia	994
Theodoro, Raquel Cordeiro	966	Tsotsou, Georgia Eleni	1011
Thiébauld, Émilie	963	Tsujikawa, Kazutake	71
Thies, Jennifer L.	291	Tsukamoto, Sachiko	91
Thomas, Gaslonde	380	Tsypik, Olga	409

Viault, Guillaume	258, 700, 908	Wang, Shen-Chieh	671
Vicente, Francisca	77	Wang, Shih-Wei	505
Vicuna, Elena Diaz	123	Wang, Wenjing	560
Vieira, Paulo Cezar	62	Wang, Xiao	64
Viljoen, Alvaro	39, 626, 805	Wang, Xinhui	386
Vincken, Jean-Paul	206, 383	Wang, Yan Alexander	860
Viso, Araceli L�rida	989	Warpeha, Katherine M.	876, 116
Vissiennon, Cica	922	Washio, Kenji	476
Vitale, Rosa Maria	37, 625	Wasilewicz, Andreas	72
Vita, Simona De	999	Watakajaturaphon, Sorawit	1039
Vitiello, Maria	241	Watts, Darrell	696
Vitsos, Andreas	621, 868	Wdowiak, Kamil	869
Vladimir-Kne�zevi�, Sanda	331, 332	Weber, M�rton	213
Vlad Luca, Simon	102	Weeraratne, Nirodha	63
Vlasova, Inna	430, 432, 434	Weghuber, Julian	482, 522
Vogel, Hermine	710	Wei, Ching-Peng	793
Vogt, Dominic	664	Weinberg, Jaclyn	488
Voltolini, Adrielli Tenfen	734	Weisenburger, Sabrina	533, 549, 550
Vongthip, Wudtipong	1020	Weishaupt, Ramon	664
Vongwairat, Sapondanai	471	Weiss, Alexander	792
von Kries, von Jens-Peter	554	Weiss, Alexander K.H.	117
Vorderegger, Sabrina	681	Wei, Xiuting	469
Vorgersb, Tha�s	363	Wei, Yanjie	825
Voss, James De	221	Welti, St�phane	229
Vo, Thanh-Hoa	307	Welzel, Janika	672
Vrabl, Pamela	76	Weng, Aleksander	487
Vrca, Ivana	607, 698, 704	Weng, Alexander	543, 382
Vroomen, Sophie	412	Weng, Jing-Ru	180, 779
Vuji�, Lovorka	705	Wenner, Cynthia A.	546
Vuli�, Berislav	766	Werner, Julia	922

W

Wagner, Brian	317, 66	Weston, Leslie A.	63
Wahl, Cl�ment	956	Weston, Paul A.	63
Wakimoto, Toshiyuki	81	White, John Kerr	424
Walask�, Hana	998	Whyms, Sophie	570
Wald, Florian	47	Wiater, Adrian	472
Walig�rski, Piotr	308	Wiejak, Rafa�	463
Walkenhorst, Michael	758, 764, 765	Wiese-Klinkenberg, Anika	689
Walker, Donald M.	68	Wilcox, Dominique	70
Walker, Heather	657	Wilde, Susan B.	190
Waloszczyk, Magdalena	477	Wilhelm, Anke	812
Walsh, John J.	618	Williams, Jessica	453
Wanecz, Estelle	946	Wils, Laura	679
Wang, Caroline	838	Wilson, Lavonne	120
Wangchuk, Phurpa	742	Wilson, Tyler A.	535, 659
Wa-ngensteen, Helle	431	Wi�nska, Patrycja	763
Wang, Hui-Chun	401	Wishart, David S.	888
Wang, Pinqi	877	Witte, Claus-Peter	478
Wang, Rongbin	89	Wittman, Caroline	98
		Wojtanowski, Krzysztof Kamil	1028
		Wolfender, Jean-Luc	468, 900, 901, 110, 902, 131
		Wolff, Carsten	236
		Wolf, Johanna	478

Wolfram, Evelyn	102, 717	Yang, Huihai	101
Wolfrum, Christian	901	Yang, Hyun Ok	352
Wong, Chun Kwok	199	Yangın, Sevcen	768
Wong, Kwan-Ho	101	Yang, Jinkyu	636
Wongwan, Chamaiphorn	1020	Yang, Liping	98
Woo, Sunmin	47, 418	Yang, Wonjin	636
Wójciak, Magdalena	202	Yao, Haoyi	828
Wójcik, Sara	238	Yeerong, Natnaree	1039
Wright, J.	357	Yen, Hung-Rong	665, 800, 802
Wróbel-Biedrawa, Dagmara	214, 218	Yeniçeri, Esmâ	581
Wu, Chia-Hung	516	Yen, I-Chuan	831
Wu, Chunyan	901	Yeomans, Tim	897
Wu, Lin	181, 838	Yerena, José A. Martínez	190
Wungsintaweekul, Juraithip	542, 925	Yevide, Sèdami Igor Armand	691
Wüst, Ana Paula Simões	552	Yiga, Nicholas	716
Wu, Tian-Shung	516	Yim, Nam Hui	402, 403
Wu, Tung-Ying	937	Yim, Nam-Hui	850
Wu, Xiaohua	193	Yilmaz, Fatıma Nur	616
Wu, Yang-Chang	937	Yilmaz, Sevdan	127
Wybraniec, Sławomir	361, 362	Yoldaş, Fatma Gül	572
Wziątek, Anita	463	Yoo, Dan-Hee	1014

X

Xavier, Cachet	380	Yoon, Seon-A	752, 988
Xavier, Karina B.	437	Yoon, Yeo Dae	1029
Xavier, Virginie	491	Young, Louise	367
Xiao, Hui-Hui	810	Young, Ryan M.	236
Xiao, Jianbo	133	Youn, Ui Joung	187
Xiao, Tiqiao	865	You-Ying, Chen	186
Xiao, Wei-Lie	199	Yuan, Jui-Hung	790
Xiao, Xiansha	499	Yuan, Wei-Cheng	937
Xiao, Youli	485	Yu, Dinah	941
Xia, Yi-Xuan	918	Yue, Grace Gar-Lee	101, 198, 199, 936
Xie, Honglan	865	Yu, Heung Joo	960
Xie, Wen-Jian	918	Yu, Jae Sik	352
Xi, Wenhui	825	Yuk, Heung Joo	973, 991, 1035
Xuebo, Hu	369	Yurdakok-Dikmen, Begum	768
Xue, Yanling	865	Yurduseven, Semanur	768
Xu, Jian	936	Yusakul, Gorawit	470
Xu, Maonian	282	Yu, Su-Bin	1014
		Yusuf, Abayomi	811
		Yu, Szu-Yin	788, 789
		Yzebe, Olivier	692

Y

Yagmur, Mervé	233, 814
Yalcin, Funda N.	94
Yalçın, Funda Nuray	289, 290
Yamada, Keith	832
Yáñez-Ibarra, Guadalupe	732
Yang, Da Hee	697, 910

Z

Zabiegała, Bożena	353
Zabost, Anna	408
Zadarko, Alicja	434
Zaghini, Anna	769
Zajac, Jan	763

Zakaria, Nurul Shahira	446
Zalar, Matja	442
Zalewski, Przemysław	867, 869
Załoski, Daniel	202, 548, 745
Zamudio, Sergio	777
Zannotti, Marco	703
Zapata, Maxwell	70
Zbiljić, Miloš	594, 985
Zblewska, Daria	487, 532
Zebiri, Lila	924
Zehl, Martin	184
Zehuai, Wen	672
Zelei, Ágoston	213
Zengin, Gökhan	609, 620, 1034
Zentek, Jurgen	585
Zentek, Jürgen	430, 432, 434
Zeybek, Gulhan	631
Zgórká, Grażyna	474
Zguris, Jeanna	1022
Zhang, Haolin	133
Zhang, Hong-Jie	895, 896, 917, 918
Zhang, Mingshu	433, 619, 672
Zhang, Yalong	181
Zhang, Yanjun	941
Zhang, Yi	112
Zhao, Hongxing	433
Zhao, Yi	140
Zhao, Yong	60
Zheng, Tao	101, 198
Zhigang, Rao	826
Zhihua, Yang	672
Zhou, Yanyan	371
Zhuang, Yihao	775
Zhu, Chenxi	828
Zhu, Jin-Lin	860
Zhu, Yiyang	199
Zidorn, Christian	503
Zielińska, Sylwia	394, 487, 622
Zimmermann-Klemd, Amy Marisa	593
Zimowska, Beata	673
Zitterl-Eglseer, Karin	44
Ziyyat, Abderrahim	748
Zlatković, Bojan	606
Zölch, Sabrina	35, 568
Zolett, Gabrieli	734
Zolk, Oliver	923
Zonno, Maria Chiara	303
Zonoubi, Niloofar	531
Zorrilla, Jesús G.	171
Zotchev, Sergey B.	184
Zulfiqar, Fazila	415, 955
Zupkó, István	339, 817

Zurfluh, Leonie

534



Żarowska, Barbara

894

Żmudzki, Paweł

940

